




2013-2014 Influenza Report Week 03

January 12 – January 18, 2014

About our flu activity reporting

MSDH relies upon selected sentinel health practitioners across the state to report the percentage of total patient visits consistent with an influenza-like illness (ILI: fever of 100°F or higher AND cough and/or sore throat). Also, providers are supplied with specimen collection kits. Samples are submitted to the Mississippi Public Health Laboratory for influenza PCR testing. Reports are used to estimate the state's ILI rate and the magnitude of the state's influenza activity. Reports represent only the distribution of flu in the state, not an actual count of all flu cases statewide. **Information is provisional only and may change depending on additional reporting from sentinel providers.**

State ILI Surveillance



The state ILI rate had been slowly, but steadily increasing since week 41. However, during week 01 and continuing through week **03** (01/12-18/2014), the overall state ILI


rate (**5.6%**) **decreased** from last week (**6.3%**), but was comparable to this time last year (**5.9%**). |

Figure 1


Total number of patients treated by sentinel providers in the last three weeks. | **Table 1**

2013-2014 Influenza Season				
CDC Week	Week Ending	Total patients	ILI symptoms	ILI Rate (%)
03	Jan. 18	17163	958	5.6%
02	Jan. 11	14834	939	6.3%
01	Jan. 04	16231	1260	7.8%

During week **03**, **one** district (3) had an increase in ILI activity, while **four** districts (2, 7, 8, and 9) had a decrease. **Four** districts (1, 4, 5, and 6) remained about the same. *Information is provisional only and may change depending on additional reporting from sentinel providers.* | **Table 2**



MSDH District ILI Rates (%) 2013-2014		
District	Week 02	Week 03
State	6.3	5.6
I	3.5	3.1
II	16.2	7.8
III	6.8	12.9
IV	4.7	4.8
V	2.2	2.0
VI	14.9	14.9
VII	6.5	4.8
VIII	6.6	4.4
IX	7.2	6.7




Since week 35, the percentage of reported ILI cases has been highest among those in the **5-24 years** of age group. During week **03**, however,


the percentage of reported ILI was highest among individuals in the **5-24 years** and **25-49 years** of age groups. | [Figure 2](#)

The 2013-14 state ILI rate was **above** the national, Region 4, and the state baselines for week **03**. |

Figure 3




Mississippi ILI Rates 2010-2014 | **Figure 4**



Flu Testing Reports



From week **40** (week ending October 5th) through week **03** (week ending January 18th), **116** positive influenza samples were identified by MSDH. One hundred one (101) of the samples were identified as influenza A (2009 H1N1). Fourteen samples were identified as influenza A (subtype not performed) and one was identified as influenza B.

The influenza cases were identified from the following counties: Adams (1), Alcorn (1), Benton (1), Chickasaw (1), Clay (1), Coahoma (5), Copiah (1), Covington (1), Forrest (4), George (2), Grenada (2), Hancock (1), Harrison (17), Hinds (2), Itawamba (1), Jackson (16), Jones (2), Lamar (1), Lawrence (3), Leake (1), Lee (15), Lincoln (1), Marion (1), Marshall (5), Monroe (7), Neshoba (1), Oktibbeha (5), Pearl River (1), Pike (1), Prentiss (1), Stone (1), Tallahatchie (1), Tate (3), Tunica (1), Washington (3), and Yalobusha (1). The counties of four influenza cases were unknown. | [Figure 5](#)



National and Mississippi Pediatric Mortality Surveillance

Nationally, **eight** influenza-associated pediatric deaths were reported to CDC during week **03**. Five deaths were associated with a 2009 H1N1 virus and occurred during weeks 01 (week ending January 4th), 02 (week ending January 11th), and 03 (week ending January 18th). Three were associated with an influenza A virus for which no subtyping was performed and occurred during week 02 (week ending January 11th).| [Figure 6](#)




Twenty-eight influenza-associated pediatric deaths have been reported nationally during the 2013-2014 season. Of the 28 deaths, 26 (93%) have been attributed to influenza A viruses, one (4%) to an influenza A/B virus, and one (4%) to an influenza virus whose type was not determined.

influenza A viruses, one (4%) to an influenza A/B virus, and one (4%) to an influenza virus whose type was not determined.| [Figure 7](#)

Mississippi has had **one** influenza-associated pediatric death reported during this influenza season.

National ILI Surveillance


For week **03**, the MS ILI rate (5.6%) continued to remain **above** the national ILI rate (3.4%), but continued to follow national trends. | [Figure 8](#)



Mississippi reported **“Regional”** flu activity for week **03**. | [Table 3](#)

Level of Flu Activity	Definition
No Activity	Overall clinical activity remains low and there are no lab confirmed cases.
Sporadic	Isolated cases of lab confirmed influenza in the state; ILI activity is not increased <u>OR</u> A lab-confirmed outbreak in a single institution in the state; ILI activity is not increased.
Local	Increased ILI within a single region AND recent (within the past 3 weeks) laboratory evidence of influenza in that region. ILI activity in other regions is not increased <u>OR</u> two of more institutional outbreaks (ILI or lab confirmed) within a single region AND recent (within the past 3 weeks) lab confirmed influenza in that region. Other regions do not have increased ILI and virus activity is no greater than sporadic in those regions
Regional	Increased ILI in at least 2 regions but fewer than half of the regions AND recent (within the past 3 weeks) lab confirmed influenza in the affected regions <u>OR</u> Institutional outbreaks (ILI or lab confirmed) in at least 2 regions but fewer than half of the regions AND recent lab confirmed influenza in the affected regions.
Widespread	Increased ILI and/or institutional outbreaks (ILI or lab confirmed) in at least half of the regions AND recent (within the past 3 weeks) lab confirmed influenza in the state.

During week **03**, influenza activity **remained high** in the United States.¹ | [Figure 9](#)



¹For up-to-date information on flu activity nationwide, please refer to the CDC’s website:
<http://www.cdc.gov/flu/weekly/fluactivitiesurv.htm>.

Additional information:

Centers for Disease Control and Prevention	http://cdc.gov/flu/
Centers for Disease Control and Prevention FluView	http://www.cdc.gov/flu/weekly/
Flu.gov	http://www.flu.gov/
MSDH Flu and Pneumonia	http://msdh.ms.gov/msdhsite/ static/14,0,199.html
Google Flu Trends	http://www.google.org/flutrends/
World Health Organization FluNet	http://www.who.int/influenza/gisrs_laboratory/flunet/en/

Page intentionally blank

Appendix

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6





Figure 7

**Percentage of Influenza-Associated Pediatric Deaths
by Influenza Type and Subtype, Nationwide,
September 30, 2013 - January 18, 2014
(CDC Week 40, 2013 - Week 03, 2014)**


■ Flu A ■ Flu B ■ Flu A/B Not Distinguished ■ Flu A/B



N=28

Figure 8

Comparison of Nationwide ILI Rates to Mississippi ILI Rates Weeks 40-03 | 2013-2014 Flu Season



US ILI rates from the Centers for Disease Control and Prevention: <http://www.cdc.gov/flu/weekly/>

CDC Week

Figure 9

