




2011-2012 Influenza Report Week 17 *April 22 – April 28, 2012*

About our flu activity reporting

MSDH relies upon selected sentinel health practitioners across the state to report the percentage of non-trauma visits consistent with an influenza-like illness (ILI: fever > 100°F AND cough and/or sore throat). Also, providers are supplied with specimen collection kits: Samples are submitted to the Mississippi Public Health Laboratory for influenza PCR testing. Reports are used to estimate the state’s ILI rate and the magnitude of the state’s influenza activity. Reports represent only the distribution of flu in the state, not an actual count of all flu cases statewide. **Information is provisional only and may change depending on additional reporting from sentinel providers.**

State ILI Surveillance



For week 17 (04/22/2012-04/28/2012), the overall state ILI rate (**3.8%**) was **comparable** to last week (**3.5%**) and to this time last year (**3.6%**). For the past six weeks, the state ILI has remained stable. | **Figure 1**

Total number of non-trauma patients treated by sentinel providers in the last three weeks. | **Table 1**


2011-2012 Influenza Season				
CDC Week	Week Ending	Non-trauma patients	ILI symptoms	ILI Rate (%)
17	Apr. 28	12127	457	3.8%
16	Apr. 21	12966	451	3.5%
15	Apr. 14	13512	522	3.9%

During week 17, **two** districts (2 and 6) had an increase in ILI activity, while **one** district (3) had a decrease. **Six** districts (1, 4, 5, 7, 8, and 9) remained about the same.


Information is provisional only and may change depending on additional reporting from sentinel providers. /

Table 2

MS Public Health Districts | **Map**



MSDH District ILI Rates (%) 2011-2012		
District	Week 16	Week 17
<i>State</i>	3.5	3.8
I	2.4	2.0
II	0.9	5.1
III	6.0	3.4
IV	5.1	5.3
V	0.8	1.0
VI	4.2	6.1
VII	2.3	2.7
VIII	6.0	6.0
IX	3.6	4.1




The number of reported ILI cases has been highest among people in the 0 to 4 years of age group, followed by those in the 5 to 24 years of age group.

During Week


17, the number of reported ILI cases was highest among individuals in the **0-4 years** of age group. |

Figure 2

The 2011-12 state ILI rate was **below** baseline for week 17. | **Figure 3**




Mississippi ILI Rates 2009-2012 | **Figure 4**



Flu testing Reports

From week **35** (week ending September 3rd) through week **17** (week ending April 21st), fifty-seven (**57**) positive influenza samples were identified by MSDH. Fifty-three were influenza A and four were influenza B. Thirty were subtyped as influenza A (H3) and nine were subtyped as influenza A (H1). Fourteen samples did not have subtyping performed.

The influenza cases were identified from the following counties: Benton, Calhoun (4), Chickasaw, Coahoma, Copiah (2), Covington, DeSoto (2), Forrest, Harrison (3), Hinds (6), Lawrence, Leflore, Lincoln, Marshall (9), Monroe (2), Neshoba, Oktibbeha, Panola, Pike, Prentiss, Rankin (3), Stone, Tate (2), and Yazoo (10). | **Figure 5**



National ILI Surveillance

Nationally, there were two influenza-associated pediatric deaths reported to CDC during week **17**. One was associated with a 2009 H1N1 virus and the other was associated with a seasonal influenza A (H3) virus. The reported deaths occurred during week 7 (week ending February 18th) and week 14 (week ending April 7th). The total number of influenza-associated pediatric deaths reported during the 2011-2012 season is **20**. Additional data can be found at:


<http://gis.cdc.gov/GRASP/Fluview/PedFluDeath.html>.

Mississippi has had no reported influenza-associated pediatric deaths during this influenza season.


Mississippi reported “**Sporadic**” activity for week 17. | **Table 4**

Level of Flu Activity	Definition
No Activity	Overall clinical activity remains low and there are no lab confirmed cases.
Sporadic	Isolated cases of lab confirmed influenza in the state; ILI activity is not increased <u>OR</u> A lab-confirmed outbreak in a single institution in the state; ILI activity is not increased.
Local	Increased ILI within a single region AND recent (within the past 3 weeks) laboratory evidence of influenza in that region. ILI activity in other regions is not increased <u>OR</u> two of more institutional outbreaks (ILI or lab confirmed) within a single region AND recent (within the past 3 weeks) lab confirmed influenza in that region. Other regions do not have increased ILI and virus activity is no greater than sporadic in those regions
Regional	Increased ILI in at least 2 regions but fewer than half of the regions AND recent (within the past 3 weeks) lab confirmed influenza in the affected regions <u>OR</u> Institutional outbreaks (ILI or lab confirmed) in at least 2 regions but fewer than half of the regions AND recent lab confirmed influenza in the affected regions.
Widespread	Increased ILI and/or institutional outbreaks (ILI or lab confirmed) in at least half of the regions AND recent (within the past 3 weeks) lab confirmed influenza in the state.

For week 17, the MS ILI rate continued to remain **above** the national rate. | **Figure 6**



During week **17**, influenza activity declined nationally and in most regions, but remained elevated in some areas of the United States.¹ | **Figure 7**



¹For up-to-date information on flu activity nationwide, please refer to the CDC’s website:
<http://www.cdc.gov/flu/weekly/fluactivity.htm>

Additional information:

Centers for Disease Control and Prevention	http://cdc.gov/flu/
Flu.gov	http://www.flu.gov/
MSDH Flu and Pneumonia	http://msdh.ms.gov/msdhsite/ static/14,0,199.html
Google Flu Trends	http://www.google.org/flutrends/
World Health Organization FluNet	http://www.who.int/csr/disease/influenza/influenznetwork/flunet/en/

Page intentionally blank

Appendix

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7

