



2017-2018 Influenza Surveillance Report

Week 19

May 6 – May 12, 2018


About our flu activity reporting

MSDH relies upon selected sentinel health practitioners across the state to report the percentage of total patient visits consistent with an influenza-like illness (ILI: fever of 100°F or higher AND cough and/or sore throat). Also, providers are supplied with specimen collection kits. Samples are submitted to the Mississippi Public Health Laboratory for influenza PCR testing. Reports are used to estimate the state's ILI rate and the magnitude of the state's influenza activity. Reports represent only the distribution of flu in the state, not an actual count of all flu cases statewide. **Information is provisional only and may change depending on additional reporting from sentinel providers.**

Content

- ❖ [State ILI Surveillance](#)
- ❖ [Influenza Outbreaks](#)
- ❖ [Flu Testing Reports](#)
- ❖ [National and Mississippi Pediatric Mortality Surveillance](#)
- ❖ [National ILI Surveillance](#)
- ❖ [Appendix](#)
 - [Figure 1](#) (Statewide ILI Data, Current Season compared to Previous Season)
 - [Figure 2](#) (Percentage of ILI Cases by Age Group, Mississippi)
 - [Figure 3](#) (State ILI Rates 2014-2018 (YTD))
 - [Figure 4](#) (Number of Reported Influenza Outbreaks by Type and Subtype, Mississippi)
 - [Figure 5](#) (Comparison of Statewide ILI Rate to Positive Influenza Isolates by Type and Subtype, Mississippi)
 - [Figure 6](#) (Number of Influenza-Associated Pediatric Deaths by Week of Death and Influenza Type, Nationwide)
 - [Figure 7](#) (Percentage of Influenza-Associated Pediatric Deaths by Influenza Type, Nationwide)
 - [Figure 8](#) (Comparison of Nationwide and Region 4 ILI Rates to Mississippi ILI Rates)
 - [Figure 9](#) (CDC FluView Weekly Influenza Activity Map)

State ILI Surveillance




During week **19** (05/06/18-05/12/18), the overall state ILI rate (**1.8%**) was **comparable** to the previous week (**2.1%**) and to this time last year (**2.2%**). | [Figure 1](#)


Total number of patients treated by sentinel providers in the last three weeks. | **Table 1**

2017-2018 Influenza Season					
CDC Week	Week Ending	Number of reports received from Sentinel Providers	Total patients	ILI symptoms	ILI Rate (%)
19	May 12	127	17226	315	1.8
18	May 5	135	18111	386	2.1
17	Apr. 28	135	18241	405	2.2

During week **19**, one district (3) had an increase in ILI activity, while one district (6) had a decrease. The remaining seven districts (1, 2, 4, 5, 7, 8, and 9) remained about the same. *Information is provisional only and may change depending on additional reporting from sentinel providers.* | **Table 2**




MSDH District ILI Rates (%) 2017-2018		
District	Week 18	Week 19
State	2.1	1.8
I	2.4	2.0
II	2.8	2.5
III	0.7	1.8
IV	2.0	1.6
V	1.8	1.9
VI	6.8	3.3
VII	2.1	2.2
VIII	1.3	0.9
IX	1.9	1.8



Overall, the percentage of reported ILI cases has been highest among those in the **0-4** and **5-24 years** of age groups. During week **19**, the percentage of ILI cases increased in the

5-24 years of age group. | [Figure 2](#)


Mississippi ILI Rates 2014-2018 | [Figure 3](#)



Influenza Outbreaks

Outbreaks are reportable in Mississippi as a Class 1A event and must be reported by telephone within **24 hours** of first knowledge or suspicion to the Mississippi State Department of Health. For more information on reportable diseases and conditions, please refer to the [MSDH List of Reportable Diseases and Conditions](#).

Between week 40 (week ending October 7th) and week **19** (week ending May 12th), 162 influenza outbreaks were reported to MSDH, with the last outbreak reported to MSDH on March 12, 2018. MSDH investigates all reported outbreaks, and of the 162 reported outbreaks, complete information was available for all of them. Twenty-four (15%) of the outbreaks were attributed to influenza A (H1), 62 (38%) were due to influenza A (H3), 14 (8%) were due to influenza A, unknown subtype, 41 (25%) were due to influenza B (Yamagata), seven (4%) were due to influenza B, unknown lineage, and 17 (10%) were due to an unknown influenza type. (Note: Three outbreaks were associated with two different influenza strains.) | [Figure 4](#)




The influenza outbreaks have occurred in the following counties: Adams (5), Alcorn (4), Amite (1), Attala (2), Benton (1), Bolivar (2), Calhoun (2), Choctaw (1), Clarke (1), Clay (1), Coahoma (1), Covington (3), DeSoto (6), Forrest (5), Greene (1), Grenada (2), Hancock (1), Harrison (6), Hinds (13), Holmes (1), Itawamba (1), Jackson (3), Jasper (1), Jones (6), Lafayette (2), Lamar (3), Lauderdale (4), Lee (3), Lincoln (2), Lowndes (3), Madison (2), Marshall (1), Monroe (5), Neshoba (7), Newton (2), Noxubee (1), Oktibbeha (1), Panola (1), Pearl River (2), Pike (3), Pontotoc (3), Prentiss (3), Quitman (2), Rankin (8), Scott (2), Simpson (3), Smith (3), Stone (2), Sunflower (1), Tallahatchie (1), Tate (1), Tippah (2), Tunica (1), Union (3), Walthall (3), Warren (2), Washington (4), Wilkinson (1), Winston (1), Yalobusha (1), and Yazoo (3).

For additional information on infection control measures in health care facilities and managing influenza outbreaks in long-term care facilities, please refer to the CDC’s webpages:

<https://www.cdc.gov/flu/professionals/infectioncontrol/index.htm> and <https://www.cdc.gov/flu/professionals/infectioncontrol/ltc-facility-guidance.htm>, respectively.

Flu Testing Reports



Since week 40 (week ending October 7th), **280** laboratory confirmed influenza samples have been identified. Seventy-eight (28%) were identified as influenza A (H1), 127 (45%) were identified as influenza A (H3), two (1%) were identified as influenza A, unknown subtype and 73 (26%) were identified as influenza B. The last positive influenza sample identified by MSDH occurred in week 9 (week ending March 3, 2018). | [Figure 5](#)



The influenza cases were identified from the following counties: Adams (7), Alcorn (6), Attala (2), Bolivar (3), Calhoun (3), Choctaw (1), Clarke (1), Clay (3), Coahoma (6), Covington (11), DeSoto (6), Forrest (11), Greene (3), Grenada (3), Harrison (14), Hinds (15), Itawamba (1), Jackson (6), Jasper (2), Jones (10), Kemper (1), Lafayette (2), Lamar (3), Lauderdale (8), Leake (1), Lee (5), Leflore (3), Lincoln (1), Lowndes (10), Madison (4), Marion (2), Marshall (13), Monroe (5), Neshoba (10), Newton (2), Noxubee (1), Oktibbeha (7), Panola (1), Pearl River (3), Pike (5), Pontotoc (5), Prentiss (3), Quitman (3), Rankin (8), Scott (4), Sharkey (2), Simpson (3), Smith (2), Stone (4), Sunflower (1), Tallahatchie (1), Tate (3), Tippah (7), Tunica (1), Union (4), Walthall (2), Warren (4), Washington (4), Wilkinson (3), Winston (5), Yalobusha (1), and Yazoo (6). The counties for three of the cases were unknown.

National and Mississippi Pediatric Mortality Surveillance

Nationally, **three** influenza-associated pediatric deaths were reported to CDC during week **19**. One death was associated with an influenza A(H1N1)pdm09 virus and occurred during week 16 (week ending April 21, 2018). One death was associated with an influenza B virus and occurred during week 19 (week ending May 12, 2018). One death was associated with an influenza virus for which type was not determined and occurred during week 14 (week ending April 7, 2018). **One hundred sixty-eight** influenza-associated pediatric deaths have been reported to CDC for the 2017-2018 season. | [Figure 6](#)




Of the **168** influenza-associated pediatric deaths reported nationally during the 2017-2018 season, 102 (61%) have been attributed to influenza A viruses, 62 (37%) to influenza B viruses, three (2%) to an influenza virus for which type was not determined, and one (1%) to an influenza virus co-infection.

for which type was not determined, and one (1%) to an influenza virus co-infection. | [Figure 7](#)

2017 – 2018 Influenza Season | Week 19 Influenza Surveillance Report| May 06 – May 12, 2018
 Mississippi has had **three** influenza-associated pediatric deaths reported during this influenza season.
 For additional information on influenza-associated pediatric deaths, please refer to the [CDC's FluView](https://www.cdc.gov/fluview/).


National ILI Surveillance

During week **19**, the MS (1.8%), national (1.3%) and Region 4 (1.0%) ILI rates remained about the same as the previous week. | [Figure 8](#)



US and Region 4 ILI rates from the Centers for Disease Control and Prevention: <http://www.cdc.gov/flu/weekly/>.

During week **19**, influenza activity **continued to decrease** in the United States.¹ | [Figure 9](#)



¹For up-to-date information on flu activity nationwide, please refer to the CDC’s website: <http://www.cdc.gov/flu/weekly/fluactivitysurv.htm>.

Mississippi reported “**No Activity**” for the influenza activity during week **19**. | [Table 3](#)

Level of Flu Activity	Definition
No Activity	Overall clinical activity remains low and there are no lab confirmed cases.
Sporadic	Isolated cases of lab confirmed influenza in the state; ILI activity is not increased <u>OR</u> A lab-confirmed outbreak in a single institution in the state; ILI activity is not increased.
Local	Increased ILI within a single region AND recent (within the past 3 weeks) laboratory evidence of influenza in that region. ILI activity in other regions is not increased <u>OR</u> two of more institutional outbreaks (ILI or lab confirmed) within a single region AND recent (within the past 3 weeks) lab confirmed influenza in that region. Other regions do not have increased ILI and virus activity is no greater than sporadic in those regions
Regional	Increased ILI in at least 2 regions but fewer than half of the regions AND recent (within the past 3 weeks) lab confirmed influenza in the affected regions <u>OR</u> Institutional outbreaks (ILI or lab confirmed) in at least 2 regions but fewer than half of the regions AND recent lab confirmed influenza in the affected regions.
Widespread	Increased ILI and/or institutional outbreaks (ILI or lab confirmed) in at least half of the regions AND recent (within the past 3 weeks) lab confirmed influenza in the state.

Additional influenza information:

Centers for Disease Control and Prevention	http://cdc.gov/flu/
Centers for Disease Control and Prevention FluView	http://www.cdc.gov/flu/weekly/
MSDH Flu and Pneumonia	http://msdh.ms.gov/msdhsite/_static/14,0,199.html
World Health Organization FluNet	http://www.who.int/influenza/gisrs_laboratory/flunet/en/

Appendix

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

**Number of Influenza-Associated Pediatric Deaths by Week of Death
and Influenza Type, Nationwide,
October 1, 2017 - May 19, 2018
(CDC Week 40, 2017 - Week 20, 2018)**




Figure 7

**Percentage of Influenza-Associated Pediatric Deaths
by Influenza Type, Nationwide,
October 1, 2017 - May 19, 2018
(CDC Week 40, 2017 - Week 20, 2018)
N = 168**

■ Flu A ■ Flu B ■ Flu A/B Not Distinguished ■ Flu A/B




Figure 8




Figure 9

