

MISSISSIPPI STATE DEPARTMENT OF HEALTH

# PRAMS

Pregnancy  
Risk  
Assessment  
Monitoring  
System

2011

Surveillance Report Brief



## **Acknowledgments**

The 2011 Mississippi Pregnancy Risk Assessment Monitoring System (PRAMS) Surveillance Report was made possible through the support and effort of many individuals working to improve the health and well-being of mothers and their infants. Sincere appreciation is extended to mothers who participated in the survey and contributed directly to the quality of the surveillance report.

Thanks also go to the Centers for Disease Control and Prevention (CDC), the Mississippi PRAMS staff, Mississippi State Department of Health, Office of Health Data and Research, and Office of Vital Statistics for their contributions to the project.

## **Contact information**

Lei Zhang, PhD, MSc, MBA  
Director  
Office of Health Data and Research  
Mississippi State Department of Health  
PO Box 1700  
570 East Woodrow Wilson  
Jackson, MS 39215-1700  
Email: [lei.zhang@msdh.ms.gov](mailto:lei.zhang@msdh.ms.gov)  
Phone: (601) 576-8165

## **Suggested citation**

Office of Health Data and Research, Mississippi State Department of Health. Mississippi PRAMS Surveillance Report, Year 2011 Births, Jackson, MS: Mississippi Department of Health, 2015.

## **Sources for more information:**

CDC PRAMS provides more information about PRAMS on its website, <http://www.cdc.gov/prams>

# Introduction

The Pregnancy Risk Assessment Monitoring System (PRAMS) is led by the Centers for Disease Control and Prevention (CDC) in collaboration with state health departments. The goal of this project is to improve the health outcomes of mothers and infants. State-specific findings describe maternal experiences and behaviors during the time before pregnancy through the months just after delivery. Reports are used to plan and evaluate health programs, to inform policymakers and the general public. Mississippi (MS) PRAMS addresses several Title V Maternal and Child Health priorities that include factors related to low birth weight, infant mortality, access to care, teen births, and unhealthy behaviors during pregnancy (e.g. cigarette smoking).

In this report, responses to questions are provided for all women with a recent birth. Methods are explained in the appendix, including guidelines for interpreting the report. The topics covered include:

1. Alcohol and drug use
2. Low birth weight, preterm birth, and preconception care
3. Nutrition and physical activity
4. Sexually transmitted disease
5. Pregnancy and birth rate
6. Violence
7. Infant feeding and care
8. Delivery or pregnancy history
9. Mental health
10. Maternal morbidity
11. Demographics/SES
12. Miscellaneous

Healthy People 2020 objectives related to the questions are given whenever possible (U.S. Department of Health and Human Services. Office of Disease Prevention and Health Promotion. Healthy People 2020. Washington, DC. Available at <http://www.health.gov/healthypeople>. Accessed February 26<sup>th</sup>, 2014.).

## Interpretation of Text and Tables

The PRAMS survey requires special weighted calculations to account for the survey design and present population level estimates. "Percent of women" refers to the proportion of Mississippi female residents with an in-state live birth minus those women who adopt infants and other exclusions. This report does not include mothers who are excluded according to CDC protocol guidelines: out of state births to state residents, in state births to nonresidents, infants whose birth certificates lack the mother's last name, birth certificates that are processed too late after the birth occurred (i.e. more than six months), multiple gestation infants (higher than triplets), adopted infants, and surrogate births.

Because of the survey design, we report a margin of error and the 95% confidence interval (CI). Categories of maternal characteristics that have non-overlapping 95% CI are designated as statistically significantly different from each other. This conservative test of significance can be used to describe findings for each PRAMS questionnaire item. Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.

## Phase 6 Survey Guide

Alcohol and drug use		Page
Question #	Alcohol use	
21c	Discussion with health care workers (during any prenatal care visits): Drinking alcohol during pregnancy affecting baby	8
31	Any alcohol consumption in the past 2 years	11
32a	Alcohol use per week, 3 months before pregnancy	12
32b	Binge drinking 3 months before pregnancy	12
33a	Alcohol use per week, last 3 months of pregnancy	12
33b	Binge drinking, last 3 months of pregnancy	13
55f	Discussion with health care workers (12 months before pregnancy): Quitting drinking alcohol	20
Question #	Tobacco use	
21a	Discussion with health care workers (during any prenatal care visits): Quitting smoking during prenatal care	8
26	Any smoking in past 2 years	9
27	Number of cigarettes per day, 3 months before pregnancy	10
28	Number of cigarettes per day, last 3 months of pregnancy	10
29	Last 3 months of pregnancy, smoking now	11
55e	Discussion with health care workers (12 months before pregnancy): Quitting smoking	19
63i	Discussion with health care workers (before discharge from hospital): Smoking around baby	23
Question #	Drug use	
21e	Discussion with health care workers (during any prenatal care visits): Using safe medicine during pregnancy	8
21f	Discussion with health care workers (during any prenatal care visits): Illegal drug use affecting baby	8

<b>Low birth weight, preterm birth and preconception care</b>		<b>Page</b>
<b>Question #</b>	<b>Infant morbidity</b>	
37	Baby's due date	14
39	Baby's birth date	14
42	Baby in intensive care after birth	15
43	Baby's hospital stay after birth	15
44	Is the baby alive now?	16
<b>Question #</b>	<b>Pregnancy recognition</b>	
16	Weeks or months when sure pregnant	6
<b>Question #</b>	<b>Prenatal care</b>	
21	Discussions with health care workers during prenatal care visits	8
55	Discussions with health care workers, 12 months before pregnancy	20
56	Test for birth defects during prenatal care	20
57	Reasons for not having birth defect tests	20
58	Problems with health care workers during prenatal care	21
63	Health care workers instructions to mother before discharge	23
<b>Question #</b>	<b>Prenatal care initiation</b>	
16	Weeks or months when sure pregnant	6
17	Weeks or months of first prenatal care visit	6
18	Prenatal care as early as wanted	7
19	Barriers to prenatal care	7

<b>Nutrition and physical activity</b>		<b>Page</b>
<b>Question #</b>	<b>Nutrition and physical activity</b>	
1a	Dieting 12 months before pregnancy to lose weight	1
1b	Exercising 3 or more days per week	1
55b	Discussion with health care workers (12 months before pregnancy): Healthy weight	20
55d	Discussion with health care workers (12 months before pregnancy): Healthy eating and water intake	20
<b>Question #</b>	<b>Screening risk of chronic disease</b>	
1	Activities 12 months before pregnancy	1
4	Mother's weight before pregnancy	2
5	Mother's height	2
21a	Discussion with health care workers (during any prenatal care visits): How smoking during pregnancy affects baby	8
21c	Discussion with health care workers (during any prenatal care visits): How drinking alcohol during pregnancy affects baby	8
24	Gestational diabetes this pregnancy	9
25e	High blood pressure or hypertension during pregnancy	9
26	Any smoking in past 2 years	9
27	Number of cigarettes per day, 3 months before pregnancy	10
28	Number of cigarettes per day, last 3 months of pregnancy	10
29	Last 3 months of pregnancy, smoking now	11
30	Smoking rules inside home	11
31	Any alcohol consumption in the past 2 years	11
32a	Alcohol per week, 3 months before pregnancy	12
32b	Binge drinking 3 months before pregnancy	12
33a	Alcohol per week, last 3 months of pregnancy	12
33b	Binge drinking, last 3 months of pregnancy	13
55b	Discussion with health care workers (12 months before pregnancy): Healthy weight	20
55c	Discussion with health care workers (12 months before pregnancy): Getting annual checkup	20
64a	Close family members (blood relatives) had diabetes	23
64b	Close family members (blood relatives) had heart problems	23
64c	Close family members (blood relatives) had hypertension	23
<b>Question #</b>	<b>Multivitamin use</b>	
3	Multivitamin, prenatal vitamin, or a folic acid vitamin 12 months before pregnancy	2

55a	Discussion with health care workers (12 months before pregnancy): Taking a multivitamin with folic acid	20
-----	---	----

<b>Nutrition and physical activity (continued)</b>		Page
Question #	Obesity	
4	Women's weight before pregnancy	2
5	Women's height	2
55b	Discussion with health care workers (12 months before pregnancy): Healthy weight	20
64	Immediate family members (blood relatives) who have chronic diagnostic conditions	23
Question #	Women, Infants and Children (WIC) Program	
23	On WIC during pregnancy	8

<b>Sexually transmitted disease / HIV</b>		Page
Question #	Human Immunodeficiency Virus (HIV) test	
21j	Health care workers talk about getting testing for HIV during prenatal care visits	8
22	HIV test during pregnancy or delivery	8
Question #	Sexually Transmitted Diseases (STDs), Urinary Tract Infection (UTI), vaginal infection	
21h	Discussion with health care workers (during any prenatal care visits): Signs and symptoms of preterm labor	8
25g	Preterm or early labor	9
25h	Premature rupture of membranes	9
59	UTI, STDs, vaginal infection (include <i>Bacterial Vaginosis</i> or group B Streptococcus) during most recent pregnancy	20
60	What type of vaginal infection	21

<b>Pregnancy intention and contraception</b>		<b>Page</b>
<b>Question #</b>	<b>Contraception</b>	
12	Feelings about being pregnant before pregnancy	4
13	Trying to get pregnant before pregnancy	5
14	Doing anything to prevent pregnancy before pregnancy	5
15	Reasons for not doing anything to prevent pregnancy before pregnancy	5
52	Doing anything now to prevent pregnancy before pregnancy	18
53	Reasons for not doing anything now to prevent pregnancy before pregnancy	18
55g	Discussion with health care workers (12 months before pregnancy): Birth control	20
<b>Question #</b>	<b>Pregnancy intention</b>	
12	Feelings about being pregnant before pregnancy	4
13	Trying to get pregnant before pregnancy	5
14	Doing anything to prevent pregnancy before pregnancy	5
34h	Husband or partner did not want pregnancy 12 months before delivery	13
55g	Discussion with health care workers (12 months before pregnancy): Birth control	20

<b>Violence</b>		<b>Page</b>
<b>Question #</b>	<b>Physical abuse</b>	
211	Discussion with health care workers (during any prenatal care visits): Physical abuse by husband or partner	8
35	Did husband or partner physically abuse 12 months before pregnancy	14
36	Did husband or partner physically abuse during pregnancy	14



<b>Infant feeding and care</b>		<b>Page</b>
<b>Question #</b>	<b>Breastfeeding</b>	
21b	Discussion with health care workers (during any prenatal care visits): Breastfeeding	8
46	Ever breastfed or pumped breast milk	16
47	Currently breastfeeding or pumping breast milk	16
48	Duration of breastfeeding or pumping breast milk	16
<b>Question #</b>	<b>Infant health care</b>	
51	Did baby receive one week check-up	18
63j	Discussion with health care workers (before discharge from hospital): Taking baby to doctor	23
63k	Discussion with health care workers (before discharge from hospital): Signs of baby's illness	23
63l	Discussion with health care workers (before discharge from hospital): Baby's care at home	23
<b>Question #</b>	<b>Sleep behaviors</b>	
50	Baby's sleeping position	17
63c	Discussion with health care workers (before discharge from hospital): How or where to lay baby down to sleep	23
63d	Discussion with health care workers (before discharge from hospital): How to prepare baby's bed for sleeping	23
<b>Question #</b>	<b>Feeding</b>	
49a	Baby's age when consumed liquid other than breast milk	17
49b	Baby's age when consumed food	17
63f	Discussion with health care workers (before discharge from hospital): Baby's position for feeding	23
63g	Discussion with health care workers (before discharge from hospital): Frequency for feeding baby	23

<b>Delivery or pregnancy history</b>		<b>Page</b>
<b>Question #</b>	<b>Hospital length of stay</b>	
38	Hospital entry date for delivery	14
39	Baby's birth date	14
40	Women's stay in the hospital after baby born	14
<b>Question #</b>	<b>Pregnancy history</b>	
8	Previous live birth	3
9	Previous low birth weight birth	3
10	Previous premature infant	4
11	Baby spacing	4
65	Previous miscarriage or baby death	23
66	Reasons for previous baby death	24

<b>Mental health</b>		<b>Page</b>
<b>Question #</b>	<b>Mental health</b>	
1f	Visited health care workers for depression treatment or assessment 12 months before pregnancy	1
21k	Discussion with health care workers (during any prenatal care visits): Depression during pregnancy or after delivery	8
34	Stressful events 12 months before delivery	13
54	Depression symptoms after pregnancy	19
64d-f	Immediate family members (blood relatives) who have depression, postpartum depression, or anxiety	23

<b>Maternal morbidity</b>		<b>Page</b>
<b>Question #</b>	<b>Maternal morbidity</b>	
1c	Regularly taking prescription medicines other than birth control 12 months before pregnancy	1
7	Health care workers told mothers they had Type 1 or Type 2 diabetes before pregnancy	3
24	Health care workers told mothers they had gestational diabetes during pregnancy	9
25	Problems during pregnancy	9
54	Depression symptoms after pregnancy	19
59	<i>UTI, STDs</i> , vaginal infection (include <i>Bacterial vaginosis</i> or group B <i>Streptococcus</i> ) during most recent pregnancy	21
60	What type of vaginal infection	21
67	Health care workers ever told mothers they had asthma	24

<b>Demographics/Socioeconomic status</b>		<b>Page</b>
<b>Question #</b>	<b>Demographics</b>	
2	Insurance coverage the month before pregnancy	1
6	Mom's date of birth	2
20	Health insurance for prenatal care	7
23	WIC during pregnancy	8
41	Health insurance for delivery	15
68	12 months before delivery, yearly total household income before taxes	24
69	12 months before delivery, number of people who depended on household income	25
<b>Question #</b>	<b>Delivery payment</b>	
41	Health insurance coverage for delivery	15
<b>Question #</b>	<b>Insurance coverage</b>	
2	Insurance coverage the month before pregnancy	1
40	Discharge date after delivery	14
41	Health insurance coverage for delivery	15
<b>Question #</b>	<b>Medicaid</b>	
2c	Insurance coverage the month before pregnancy	1
40	Discharge date after delivery	14
41c	Health insurance to cover delivery of new baby	15
<b>Question #</b>	<b>Socioeconomic status/Services</b>	
2	Insurance coverage the month before pregnancy	1
40	Discharge date after delivery	14
68	12 months before delivery, yearly total household income before taxes	24
69	12 months before delivery, number of people who depended on household income	25
<b>Question #</b>	<b>WIC Program</b>	
23	WIC during pregnancy	8

<b>Miscellaneous</b>		<b>Page</b>
<b>Question #</b>	<b>Oral health</b>	
1h	Had teeth cleaned by a dentist or dental hygienist 12 months before pregnancy	1
62	Care of teeth during pregnancy	22
<b>Question #</b>	<b>Support during pregnancy</b>	
61	Things that would help women during pregnancy	22
<b>Question #</b>	<b>Seat belt use</b>	
21d	Discussion with health care workers (during any prenatal care visits): Seat belt use during pregnancy	8

## Before pregnancy

### 1. At any time during the 12 months before you got pregnant with your new baby, did you do any of the following things?

#### Women's behaviors 12 months before pregnancy

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. I was dieting (changing my eating habits) to lose weight	33.9	30.0	37.8
b. I was exercising 3 or more days of the week	40.9	36.9	44.8
c. I was regularly taking prescription medicines other than birth control	18.2	15.1	21.4
d. I visited a health care worker to be checked or treated for diabetes	12.8	10.1	15.5
e. I visited a health care worker to be checked or treated for high blood pressure	15.5	12.6	18.5
f. I visited a health care worker to be checked or treated for depression or anxiety	10.0	7.6	12.3
g. I talked to a health care worker about my family medical history	23.4	19.9	26.8
h. I had my teeth cleaned by a dentist or dental hygienist	46.2	42.2	50.2

### 2. During the month before you got pregnant with your new baby, were you covered by any of these health insurance plans?

#### Health insurance coverage during the month before pregnancy

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. Health insurance from your job or the job of your husband, partner, or parents	46.2	42.2	50.2
b. Health insurance that you or someone else paid for (not from a job)	5.4	3.5	7.2
c. Medicaid	22.4	19.3	25.6
d. TRICARE or other military health care	2.9	1.4	4.2
e. Indian Health Service	0.3	0.0	0.7
f. SCHIP	0.7	0.1	1.2
g. Other source(s)	1.4	0.4	2.3
h. I did not have any health insurance before I got pregnant	24.3	21.0	27.8

Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.

**3. During the *month before* you got pregnant with your new baby, how many times a week did you take a multivitamin, a prenatal vitamin, or a folic acid vitamin?**

Vitamin intake before pregnancy

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. I didn't take a multivitamin, prenatal vitamin, or folic acid vitamin at all	57.3	53.3	61.3
b. 1 to 3 times a week	10.3	7.8	12.7
c. 4 to 6 times a week	7.3	5.1	9.5
d. Every day of the week	25.0	21.4	28.5

**4. Just before you got pregnant with your new baby, how much did you weigh?**

Women's weight from this question is used to calculate Body Mass Index (BMI) before pregnancy.

**5. How tall are you without shoes?**

Women's height from this question is used to calculate Body Mass Index (BMI) before pregnancy.

Women's Body Mass Index (BMI) before pregnancy

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Underweight, BMI <18.5 kg/m <sup>2</sup>	10.7	8.3	13.1
Normal weight, BMI 18.5 kg/m <sup>2</sup> - 24.9 kg/m <sup>2</sup>	48.6	44.6	52.6
Overweight, BMI 25 kg/m <sup>2</sup> -29.9 kg/m <sup>2</sup>	17.0	14.0	20.1
Obese, BMI 30+ kg/m <sup>2</sup>	23.7	20.2	27.1

**6. What is *your* date of birth?**

Women's age group (years)

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
19 and younger	8.1	7.2	9.0
20-24	34.9	31.1	38.7
25-34	47.4	43.5	51.3
35 and older	9.4	6.9	11.8

**7. Before you got pregnant with your new baby, were you ever told by a doctor, nurse, or other health care worker that you had Type 1 or Type 2 diabetes?** This is not the same as gestational diabetes or diabetes that starts during pregnancy.

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Women who had Type 1 or Type 2 diabetes	5.3	3.4	7.1

**8. Before you got pregnant with your new baby, did you ever have any other babies who were born alive?**

*Note: Women who say “No” on this question, go to question 12*

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Women who had previous live birth	51.0	47.1	54.9

Previous live birth grouped

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
0	41.9	38.1	45.6
1	32.1	28.4	35.9
2	14.3	11.4	17.2
3-5	11.1	8.5	13.7
≥6	0.6	0.0	1.2

*Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.*

**9. Did the baby born just before your new one weigh more than 5 pounds, 8 ounces (2.5 kilos) at birth?**

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Women who had a baby with birth-weight >2.5 kilograms in previous pregnancy	91.2	89.3	93.0

**10. Was the baby *just before* your new one born *more than 3 weeks* before his or her due date?**

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Women who had previous preterm birth	16.1	12.0	20.2
<b>Women who had previous live births, low birth weight (LBW) or preterm births (PTB)</b>			
	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
No previous live births	49.2	45.3	53.1
Normal previous live births (not LBW or PTB)	40.3	36.3	44.2
Low birth weight births	2.3	1.1	3.5
Preterm births	5.5	3.6	7.3
Low birth weight & preterm births	2.7	1.5	3.9

**11. When your new baby was born, how old was the child born *just before* your new baby?**

Age of previous child

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. 0 to 12 months	2.8	1.1	4.5
b. 13 to 18 months	7.4	4.3	10.4
c. 19 to 24 months	15.5	11.2	19.7
d. More than 2 years, less than 3 years	24.7	19.5	29.8
e. 3 to 5 years	31.3	25.8	36.8
f. More than 5 years	18.3	13.8	22.9

*Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.*

**12. Thinking back to *just before* you got pregnant with your *new* baby, how did you feel about becoming pregnant?**

Women's feelings about becoming pregnant

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
I wanted to be pregnant sooner	11.1	8.5	13.6
I wanted to be pregnant later	37.7	33.9	41.4
I wanted to be pregnant then	33.9	30.0	37.7
I did not want to be pregnant	17.4	14.3	20.4

**13. When you got pregnant with your new baby, were you trying to get pregnant?**

*Note: Women who say “No” on this question, go to question 14*

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Women trying to get pregnant*	40.2	36.2	44.1

\*Go to question 16

**14. When you got pregnant with your new baby, were you or your husband or partner doing anything to keep from getting pregnant?** (Some things people do to keep from getting pregnant include not having sex at certain times [natural family planning or rhythm] or withdrawal, and using birth control methods such as the pill, condoms, vaginal ring, IUD, having their tubes tied, or their partner having a vasectomy.)

*Note: Women who say “No” on this question, go to question 15*

Birth control use among women with unintentional pregnancies

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Women or their husband or partner using birth control*	45.4	40.4	50.5

\*Go to question 16

**15. What were your reasons or your husband’s or partner’s reasons for not doing anything to keep from getting pregnant?**

Reasons for not doing anything to keep from getting pregnant

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. I didn’t mind if I got pregnant	31.0	24.6	37.5
b. I thought I could not get pregnant at that time	32.9	26.5	39.2
c. I had side effects from the birth control method I was using	7.9	4.5	11.2
d. I had problems with getting birth control when I needed it	5.1	2.2	7.9
e. I thought my husband or partner or I was sterile (could not get pregnant at all)	8.3	4.5	12.1
f. My husband or partner didn’t want to use anything	22.9	17.4	28.4
g. Other(s)	10.0	5.8	14.1



## During pregnancy

The next questions are about the prenatal care you received during your most recent pregnancy. Prenatal care includes visits to a doctor, nurse, or other health care worker before your baby was born to get checkups and advice about pregnancy.

**16. How many weeks or months pregnant were you when you were *sure* you were pregnant?** (For example, you had a pregnancy test or a doctor or nurse said you were pregnant.)

Women sure of pregnancy

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
First trimester	82.1	79.0	85.1
Second trimester	17.9	14.8	20.9
Third trimester	0.0	0.0	0.0

**17. How many weeks or months pregnant were you when you had your first visit for prenatal care?**

These results do not count women's visits that were only for a pregnancy test or only for WIC (the Special Supplemental Nutrition Program for Women, Infants, and Children).

First prenatal care visit

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
First trimester prenatal care	75.9	72.5	79.2
Second trimester prenatal care	22.6	19.3	25.8
Third trimester prenatal care	1.6	0.5	2.5

*Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.*

Number of prenatal care visits

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
≤ 8 visits	11.5	9.1	13.8
9 – 11 visits	18.7	15.7	21.8
≥12 visits	69.8	66.3	73.3

**18. Did you get prenatal care as early in your pregnancy as you wanted?**

*Note: Women who say “No” on this question, go to question 19*

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Women who got prenatal care as early as they wanted*	84.0	81.2	86.8

\*Go to question 20

**19. Did any of these things keep you from getting prenatal care at all or as early as you wanted?**

*Note: Women who did not go for prenatal care, continue to number 22*

Barriers to prenatal care

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. I couldn't get an appointment when I wanted one	29.2	20.6	37.8
b. I didn't have enough money or insurance to pay for my visits	34.0	25.1	43.0
c. I had no transportation to get to the clinic or doctor's office	14.4	7.7	21.1
d. The doctor or my health plan did not start care as early as I wanted	14.2	7.5	20.8
e. I had too many other things going on	17.4	10.4	24.5
f. I couldn't take time off from work or school	14.2	7.7	20.8
g. I didn't have my Medicaid card	37.1	28.0	46.2
h. I had no one to take care of my children	8.7	3.1	14.2
i. I didn't know that I was pregnant	39.0	30.3	47.8
j. I didn't want anyone else to know I was pregnant	19.6	12.7	26.6
k. I didn't want prenatal care	3.3	0.4	6.2

*Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.*

**20. Did any of these health insurance plans help you pay for your prenatal care?**

Health insurance coverage for prenatal care

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. Health insurance from your job or the job of your husband, partner, or parents	39.9	35.9	43.9
b. Health insurance that you or someone else paid for (not from a job)	1.8	0.6	2.8
c. Medicaid	59.2	55.2	63.2
d. TRICARE or other military health care	2.8	1.3	4.1
e. Indian Health Service	0.3	0.0	0.7
f. SCHIP	0.2	0.0	0.4
g. Other source(s)	0.8	0.1	1.5
h. I did not have health insurance to help pay for my prenatal care	2.4	1.1	3.6

*Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.*

**21. During any of your prenatal care visits, did a doctor, nurse, or other health care worker talk with you about any of the things listed below?**

Women's discussion with doctor, nurse or health workers

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. How smoking during pregnancy could affect my baby	80.1	76.7	83.4
b. Breastfeeding my baby	90.0	87.5	92.5
c. How drinking alcohol during pregnancy could affect my baby	81.6	78.3	84.8
d. Using a seat belt during my pregnancy	64.9	61.0	68.7
e. Medicines that are safe to take during my pregnancy	95.0	93.4	96.7
f. How using illegal drugs could affect my baby	78.2	74.8	81.6
g. Doing tests to screen for birth defects or diseases that run in my family	89.0	86.5	91.5
h. The signs and symptoms of preterm labor (labor more than 3 weeks before the baby is due)	86.4	83.7	89.2
i. What to do if my labor starts early	88.4	85.9	90.9
j. Getting tested for HIV (the virus that causes AIDS)	75.4	71.9	78.9
k. What to do if I feel depressed during my pregnancy or after my baby is born	72.7	69.1	76.3
l. Physical abuse to women by their husbands or partners	55.1	51.0	59.1

**22. At any time during your most recent pregnancy or delivery, did you have a test for HIV (the virus that causes AIDS)?**

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Women who had an HIV test during pregnancy	77.9	74.3	81.6

**23. During your most recent pregnancy, were you on WIC (the Special Supplemental Nutrition Program for Women, Infants, and Children)?**

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Women on WIC program during pregnancy	65.9	62.1	69.8

**24. During your most recent pregnancy, were you told by a doctor, nurse, or other health care worker that you had gestational diabetes (diabetes that started during this pregnancy)?**

	Percent	95% CI	
		Lower	Upper
Women who had gestational diabetes during this pregnancy	13.3	10.6	16.1

**25. Did you have any of the following problems during your most recent pregnancy?**

Problems during the most recent pregnancy

	Percent	95% CI	
		Lower	Upper
a. Vaginal bleeding	18.4	15.3	21.5
b. Kidney or bladder (urinary tract) infection	34.6	30.8	38.4
c. Severe nausea, vomiting, or dehydration	39.9	36.0	43.8
d. Cervix had to be sewn shut (cerclage for incompetent cervix)	1.0	0.5	1.5
e. High blood pressure, hypertension (including pregnancy-induced hypertension [PIH]), preeclampsia, or toxemia	14.0	11.3	16.7
f. Problems with the placenta (such as abruptio placenta or placenta previa)	3.5	2.1	4.8
g. Labor pains more than 3 weeks before my baby was due (preterm or early labor)	19.2	16.2	22.3
h. Water broke more than 3 weeks before my baby was due (premature rupture of membranes [PROM])	4.6	3.3	5.9
i. I had to have a blood transfusion	1.5	0.5	2.4
j. I was hurt in a car accident	1.1	0.3	1.8

*Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.*

**The next questions are about smoking cigarettes around the time of pregnancy (before, during, and after).**

**26. Have you smoked any cigarettes in the past 2 years?**

*Note: Women who say "No" on this question, go to question 30*

	Percent	95% CI	
		Lower	Upper
Women who smoked any cigarettes in the past 2 years*	22.4	19.2	25.7

\* Go to question 27

**27. In the 3 months before you got pregnant, how many cigarettes did you smoke on an average day?**  
(A pack has 20 cigarettes.)

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Women who smoked 3 months before pregnancy	19.5	16.4	22.5

Amount of cigarettes smoked on an average day in the 3 months before pregnancy

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. 41 cigarettes or more	4.5	0.8	8.0
b. 21 to 40 cigarettes	6.4	2.3	10.5
c. 11 to 20 cigarettes	31.7	24.0	39.5
d. 6 to 10 cigarettes	22.6	15.7	29.4
e. 1 to 5 cigarettes	15.4	9.9	20.8
f. Less than 1 cigarette	6.4	2.3	10.5
g. I didn't smoke then	13.0	7.4	18.5

*Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.*

**28. In the last 3 months of your pregnancy, how many cigarettes did you smoke on an average day?**  
(A pack has 20 cigarettes)

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Women who smoked in last 3 months of pregnancy	11.1	8.7	13.6

Amount of cigarettes smoked on an average day in the last 3 months of pregnancy

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. 41 cigarettes or more	4.8	0.9	8.7
b. 21 to 40 cigarettes	1.4	0.0	3.2
c. 11 to 20 cigarettes	7.6	3.5	11.8
d. 6 to 10 cigarettes	12.3	7.0	17.6
e. 1 to 5 cigarettes	18.6	12.1	25.1
f. Less than 1 cigarette	4.8	1.5	8.1
g. I didn't smoke then	50.3	42.1	58.5

*Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.*

**29. How many cigarettes do you smoke on an average day *now*? (A pack has 20 cigarettes.)**

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Women who smoke now	16.6	13.7	19.5

**Amount of cigarettes mothers smoke on an average day now**

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. 41 cigarettes or more	3.6	0.5	6.8
b. 21 to 40 cigarettes	4.0	1.3	6.8
c. 11 to 20 cigarettes	25.1	17.7	32.4
d. 6 to 10 cigarettes	20.7	14.0	27.4
e. 1 to 5 cigarettes	15.0	9.5	20.4
f. Less than 1 cigarette	5.6	1.9	9.3
g. I didn't smoke then	26.0	18.7	33.2

**30. Which of the following statements best describes the rules about smoking *inside* your home *now*?**

**Rules about smoking inside the home**

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. No one is allowed to smoke anywhere inside my home	93.7	91.9	95.6
b. Smoking is allowed in some rooms or at some times	5.1	3.5	6.7
c. Smoking is permitted anywhere inside my home	1.1	0.3	1.9

*Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.*

**The next questions are about drinking alcohol around the time of pregnancy (before, during, and after).**

**31. Have you had any alcoholic drinks in the *past 2 years*? A drink is 1 glass of wine, wine cooler, can or bottle of beer, shot of liquor, or mixed drink.**

*Note: Women who say "No" on this question, go to question 34*

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Women who drank alcohol in the past 2 years*	45.7	41.7	49.7

*\*Go to question 32a*

**32a. During the 3 months before you got pregnant, how many alcoholic drinks did you have in an average week?**

Amount of drinks women had in an average week before pregnancy

	Percent	95% CI	
		Lower	Upper
a. 14 drinks or more a week*	2.0	0.2	3.7
b. 7 to 13 drinks a week*	2.4	0.6	4.2
c. 4 to 6 drinks a week*	7.7	4.4	11.0
d. 1 to 3 drinks a week*	25.3	20.1	30.6
e. Less than 1 drink a week*	41.4	35.4	47.4
f. I didn't drink then**	21.1	16.2	25.9

Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.

\*Go to question 32b

\*\*Go to question 33a

**32b. During the 3 months before you got pregnant, how many times did you drink 4 alcoholic drinks or more in one sitting? A sitting is a two hour time span.**

Frequency of having 4 or more alcoholic drinks in one sitting before pregnancy

	Percent	95% CI	
		Lower	Upper
a. 6 or more times	3.5	0.9	6.0
b. 4 to 5 times	4.6	1.7	7.5
c. 2 to 3 times	18.4	13.0	23.8
d. 1 time	15.9	10.9	20.9
e. I didn't have 4 drinks or more in 1 sitting	57.5	50.7	64.3

Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.

**33a. During the last 3 months of your pregnancy, how many alcoholic drinks did you have in an average week?**

Amount of alcoholic drinks women had in an average week during the last 3 months of pregnancy

	Percent	95% CI	
		Lower	Upper
a. 14 drinks or more a week*	0.2	0.0	0.7
b. 7 to 13 drinks a week*	6.6	0.0	15.1
c. 4 to 6 drinks a week*	1.1	0.0	2.2
d. 1 to 3 drinks a week*	13.4	3.0	23.9
e. Less than 1 drink a week*	9.6	6.0	13.2
f. I didn't drink then**	85.2	81.0	89.5

Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.

\*Go to question 33b

\*\*Go to question 34

**33b. During the last 3 months of your pregnancy, how many times did you drink 4 alcoholic drinks or more in one sitting? A sitting is a two hour time span.**

Frequency of having 4 or more alcoholic drinks in one sitting during the last 3 months of pregnancy

	Percent	95% CI	
		Lower	Upper
a. 6 or more times	0.2	0.0	0.7
b. 4 to 5 times	6.6	0.0	15.1
c. 2 to 3 times	1.1	0.0	2.2
d. 1 time	13.4	3.0	23.9
e. I didn't have 4 drinks or more in 1 sitting	78.5	65.6	91.3

*Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.*

**Pregnancy can be a difficult time for some women. The following questions are about events that may have happened before and during the most recent pregnancy.**

**34. This question is about events that may have happened during the 12 months before your new baby was born.**

Stressful events during the 12 months before new baby was born

	Percent	95% CI	
		Lower	Upper
a. A close family member was very sick and had to go into the hospital	25.2	21.7	28.6
b. I got separated or divorced from my husband or partner	7.5	5.4	9.5
c. I moved to a new address	29.8	26.1	33.5
d. I was homeless	2.1	1.0	3.3
e. My husband or partner lost his job	11.9	9.3	14.5
f. I lost my job even though I wanted to go on working	9.7	7.4	12.1
g. I argued with my husband or partner more than usual	21.8	18.6	25.0
h. My husband or partner said he didn't want me to be pregnant	7.0	5.1	8.9
i. I had a lot of bills I couldn't pay	19.5	16.3	22.8
j. I was in a physical fight	3.6	2.2	5.0
k. My husband or partner or I went to jail	2.7	1.5	3.7
l. Someone very close to me had a problem with drinking or drugs	9.0	6.8	11.2
m. Someone very close to me died	18.4	15.3	21.5



**35. During the 12 months before you got pregnant with your new baby, did your husband or partner push, hit, slap, kick, choke, or physically hurt you in any other way?**

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Women who had physical abuse from husband or partner before pregnancy	4.8	3.2	6.5

**36. During your most recent pregnancy, did your husband or partner push, hit, slap, kick, choke, or physically hurt you in any other way?**

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Women who had physical abuse from husband or partner during pregnancy	3.2	1.9	4.5

**The next questions are about your labor and delivery.**

**37. When was your baby due?**

**38. When did you go into the hospital to have your baby?**

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Women who delivered baby in hospital	99.5	98.9	100.0

**39. When was your baby born?**

**40. When were you discharged from the hospital after your baby was born?**

Women's length of stay in the hospital

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
1-3 nights	81.6	78.5	84.6
4-7 nights	15.6	12.8	18.4
≥8 nights	1.4	0.5	2.4

**41. Did any of these health insurance plans help you pay for the *delivery* of your new baby?**

Health insurance coverage for delivery

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. Health insurance from your job or the job of your husband, partner, or parents	37.4	33.5	41.3
b. Health insurance that you or someone else paid for (not from a job)	1.7	0.6	2.7
c. Medicaid	63.8	59.9	67.7
d. TRICARE or other military health care	2.7	1.3	4.1
e. Indian Health Service	0.1	0.0	0.2
f. SCHIP	0.1	0.0	0.3
g. Other source(s)	0.1	0.0	0.2
h. I did not have health insurance to help pay for my delivery	1.8	0.7	2.8

*Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.*

## After pregnancy

**42. After your baby was born, was he or she put in an intensive care unit?**

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
The baby was in ICU after birth	11.7	9.4	13.9

**43. After your baby was born, how long did he or she stay in the hospital?**

Baby's length of stay in the hospital

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. Less than 24 hours (<1 day)*	1.2	0.3	2.1
b. 24 to 48 hours (1-2 days)*	44.4	40.4	48.4
c. 3 to 5 days*	44.9	40.9	48.9
d. 6 to 14 days*	5.7	4.0	7.3
e. More than 14 days*	2.5	1.8	3.2
f. My baby was not born in a hospital*	0.6	0.0	1.3
g. My baby is still in the hospital**	0.7	0.0	1.2

*Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.*

\* Go to question 44

\*\* Go to question 46

**44. Is your baby alive now?***Note: Women who say "No" on this question, go to question 52*

	Percent	95% CI	
		Lower	Upper
Baby is alive now*	99.2	98.8	99.7

*\*Go to question 45***45. Is your baby living with you now?***Note: Women who say "No" on this question, go to question 52*

	Percent	95% CI	
		Lower	Upper
Baby lives with mother now*	99.5	98.9	100.0

*\*Go to question 46***46. Did you ever breastfeed or pump breast milk to feed your new baby after delivery, even for a short period of time?***Note: Women who say "No" on this question, go to question 49b*

	Percent	95% CI	
		Lower	Upper
Women who ever breastfed or pumped breast milk to feed baby*	51.1	47.1	55.1

*\* Go to question 47***47. Are you currently breastfeeding or feeding pumped milk to your new baby?***Note: Women who say "No" on this question, go to question 48*

	Percent	95% CI	
		Lower	Upper
Women who are currently breastfeeding or feeding pumped milk to baby*	32.9	27.4	38.5

*\*Go to question 49a***48. How many weeks or months did you breastfeed or pump milk to feed your baby?  
(Percentages represented are among women who initiated breastfeeding)**

Duration of breastfeeding or pumping milk

	Percent	95% CI	
		Lower	Upper
<1 week	13.9	9.1	18.5
1-3 weeks	55.7	48.7	62.6
4-7 weeks	25.9	19.8	32.0
≥8 weeks	4.5	1.5	7.4

*\*note: This is NOT exclusive breastfeeding.*

**49a. How old was your new baby the first time he or she drank liquids other than breast milk (such as formula, water, juice, tea, or cow's milk)?**  
**(Percentages represented are among women who initiated breastfeeding)**

Baby's first time drinking liquids other than breast milk

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
<1 week	35.8	29.7	41.8
1-3 weeks	40.0	33.9	46.1
4-7 weeks	20.4	15.3	25.3
≥8 weeks	3.8	1.4	6.0

**49b. How old was your new baby the first time he or she ate food (such as baby cereal, baby food, or any other food)?**  
**(Percentages represented are among women who initiated breastfeeding)**

Baby's first time ate food

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
<1 week	1.3	0.1	2.4
1-3 weeks	57.2	52.2	62.1
4-7 weeks	30.5	25.9	35.0
≥8 weeks	10.9	7.7	14.2

**50. In which *one* position do you *most often* lay your baby down to sleep now?**

Baby's sleeping position

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. On his or her side	17.3	14.2	20.3
b. On his or her back	62.5	58.6	66.4
c. On his or her stomach	16.2	13.2	19.2
d. On his or her side/back	3.0	1.7	4.3
e. On his or her side/stomach	0.6	0.0	1.2
f. On his or her back/stomach	0.1	0.0	0.2
g. All 3 positions	0.2	0.0	0.7

*Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.*

**51. Was your new baby seen by a doctor, nurse, or other health care worker for a *one week check-up* after he or she was born?**

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
The baby had check-up one week after delivery	86.2	83.4	89.0

**52. Are you or your husband or partner doing anything *now* to keep from getting pregnant?** (Some things people do to keep from getting pregnant include not having sex at certain times [natural family planning or rhythm] or withdrawal, and using birth control methods such as the pill, condoms, vaginal ring, IUD, having their tubes tied, or their partner having a vasectomy.)

*Note: Women who say “No” on this question, go to question 53*

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Women or their husband or partner were using birth control*	90.0	87.6	92.5

*\*Go to question 54*

**53. What are your reasons or your husband’s or partner’s reasons for not doing anything to keep from getting pregnant *now*?**

Reasons of women or their husband or partner for not using birth control now

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. I am not having sex	24.5	14.0	35.0
b. I want to get pregnant	13.1	4.5	21.7
c. I don’t want to use birth control	29.1	17.1	41.1
d. My husband or partner doesn’t want to use anything	11.1	3.2	19.0
e. I don’t think I can get pregnant (sterile)	6.4	0.0	13.1
f. I can’t pay for birth control	10.1	2.3	17.9
g. I am pregnant now	6.3	0.4	12.1
h. Other(s)	15.7	6.4	25.0

*Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.*

54. Below is a list of feelings and experiences that women sometimes have after childbirth. Read each item to determine how well it describes your feelings and experiences. Then, write on the line the number of the choice that best describes how often you have felt or experienced things this way *since your new baby was born*. Use the scale when answering:

1	2	3	4	5
Never	Rarely	Sometimes	Often	Always

Women's feelings after childbirth

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. I felt down, depressed, or sad			
Never	43.5	39.5	47.5
Rarely	19.9	16.7	23.1
Sometimes	23.1	19.7	26.4
Often	9.7	7.4	12.1
Always	3.8	2.4	5.2
b. I felt hopeless			
Never	68.2	64.5	71.9
Rarely	15.8	12.9	18.8
Sometimes	10.2	7.8	12.6
Often	3.5	2.1	4.9
Always	2.3	1.2	3.4
c. I felt slowed down			
Never	31.9	28.1	35.7
Rarely	22.4	19.0	25.8
Sometimes	23.5	20.1	26.9
Often	14.2	11.4	16.9
Always	8.0	5.9	10.1

## Other experiences

### 55. During the 12 months before you got pregnant with your new baby, did a doctor, nurse, or other health care worker talk with you about any of the things listed below?

Discussion with a doctor, nurse, or other health care workers during 12 months before pregnancy

	Percent	95% CI	
		Lower	Upper
a. Taking a multivitamin with folic acid	50.6	46.6	54.6
b. Maintaining or being a healthy weight	58.4	54.4	62.4
c. Getting a yearly check-up	72.3	68.7	75.9
d. Eating healthy and drinking water everyday	69.0	65.3	72.7
e. Quitting smoking	45.0	41.1	49.0
f. Quitting drinking alcohol	41.2	37.3	45.1
g. Using birth control	65.7	61.9	69.6

### 56. During any of your prenatal care visits, did you have any tests for birth defects?

*Note: Women who say "No" on this question, go to question 57*

	Percent	95% CI	
		Lower	Upper
Women who had any tests for birth defects during prenatal care*	65.5	61.3	69.7

\*Go to question 58

### 57. What were your reasons for not having tests for birth defects during your most recent pregnancy?

Reasons for not having birth defect test during pregnancy

	Percent	95% CI	
		Lower	Upper
a. I was not told about any tests for birth defects	21.5	15.6	27.5
b. I did not understand the reason for the test	1.9	0.0	3.8
c. I do not like having my blood drawn	4.7	1.5	8.0
d. I heard the tests were unreliable	14.9	9.6	20.3
e. I did not want to know if my baby had birth defects	50.6	43.2	58.1
f. The cost of the blood test was too high	7.1	3.2	11.0
g. Other(s)	18.0	12.2	23.8

*Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.*

**58. During your prenatal care visits, did you have any of the following problems with a doctor, nurse or other health care worker?**

Problems with a doctor, nurse or other health care workers during prenatal care

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. My doctor or nurse did not understand my concerns	2.8	1.6	4.0
b. My doctor or nurse did not answer my questions	3.6	2.1	5.1
c. I did not understand what my doctor or nurse was telling me	4.6	3.0	6.3
d. My doctor or nurse seemed not to care about me or my pregnancy	4.1	2.5	5.7

**59. During your most recent pregnancy, did a doctor, nurse, or other health care worker tell you that you had a urinary tract infection (UTI), a sexually transmitted disease (STD), or any vaginal infection, including Bacterial vaginosis or Group B Strep (Beta Strep)?**

*Note: Women who say "No" on this question, go to question 61*

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Women who had an infection (UTI, STD, <i>Bacterial vaginosis</i> or Group B Strep)*	40.4	36.5	44.4

\*Go to question 60

**60. What infection or disease were you told that you had?**

Type of infection/disease told by health care workers

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. Genital warts (HPV)	2.6	0.6	4.6
b. Herpes	2.4	0.4	4.4
c. Chlamydia	6.0	3.3	8.6
d. Gonorrhea	1.5	0.0	2.9
e. Pelvic inflammatory disease (PID)	0.9	0.0	1.9
f. Syphilis	1.0	0.0	2.4
g. Group B Strep (Beta Strep)	17.1	12.2	22.0
h. <i>Bacterial vaginosis</i>	11.8	7.8	15.8
i. <i>Trichomoniasis</i> (Trich)	3.3	1.3	5.4
j. Yeast infections	32.7	26.8	38.6
k. Urinary tract infection (UTI)	62.5	56.5	68.6
l. Other(s)	1.6	0.1	3.0

*Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.*



**61. During your most recent pregnancy, which one of the following things do you feel would have helped you during your pregnancy?**

Things that will help women during pregnancy

	Percent	95% CI	
		Lower	Upper
a. A pregnancy support group	14.0	11.3	16.7
b. A 1-800 Toll Free Help Line	7.1	5.1	9.1
c. Assistance with finding a daycare or child care provider	16.6	13.8	19.5
d. Better health insurance	14.9	12.0	17.8
e. A more supportive husband or partner	19.0	15.9	22.1
f. Home visits from a social worker or other health care worker to give me advice	5.9	4.1	7.6
g. Other	3.2	1.9	4.6
h. None of these things would have helped me	47.1	43.1	51.1

**62. This question is about the care of your teeth during your most recent pregnancy.**

Women's care of teeth during pregnancy

	Percent	95% CI	
		Lower	Upper
a. I went to a dentist or dental clinic for a checkup	29.5	25.8	33.2
b. I went to a dentist or dental clinic to have my teeth cleaned	25.6	22.0	29.1
c. I had painful, red, or swollen gums	12.8	10.0	15.5
d. I had to have a tooth pulled	5.0	3.2	6.7
e. I had cavities that needed to be filled	14.4	11.6	17.2
f. I needed to see a dentist for an abscess (infection surrounded by inflamed tissue)	4.4	2.8	6.0
g. A dental or other health care provider talked with me about how to care for my teeth and gums	24.0	20.5	27.5

**63. Before you were discharged from the hospital after having your new baby, did a doctor, nurse, social worker, or other health care worker talk with you about any of the following?**

Discussion before discharged from hospital after delivery

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. Having a car seat ready for my new baby	95.6	93.9	97.3
b. Having a crib ready for my new baby	73.0	69.3	76.6
c. How or where to lay my baby down to sleep	92.4	90.2	94.6
d. How to prepare my baby's bed for sleeping	76.2	72.7	79.6
e. How to dress my baby in cold or hot weather	71.0	67.3	74.7
f. How to position my baby for feeding	88.5	85.9	91.1
g. How often and how much to feed my baby	95.0	93.2	96.8
h. How to get help if I need it when breastfeeding the baby	85.3	82.4	88.1
i. How smoking near my baby could affect him or her	84.4	81.3	87.4
j. How often to take my baby to the doctor for shots or checkups	90.1	87.6	92.5
k. How to watch my baby for signs of possible illness	83.9	80.9	87.0
l. How to take care of my baby while at home	89.2	86.7	91.8

**64. Have any of your close family members who are related to you by blood (mother, father, sisters, or brothers) had any of the conditions listed below?**

Problems by close family members (blood relatives)

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. Diabetes	48.9	44.5	53.3
b. Heart problems	46.2	41.7	50.6
c. High blood pressure (Hypertension)	74.5	70.7	78.2
d. Depression	37.7	33.2	42.2
e. Postpartum depression	14.7	11.4	18.0
f. Anxiety	35.1	30.7	39.5

**65. Did you ever have a miscarriage or baby that died?**

*Note: Women who say "No" on this question, go to question 67*

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Women who had a miscarriage or baby that died*	17.5	14.4	20.6

\*Go to question 66

**66. Please check the primary reason for death.**

Reasons for baby's death

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. SIDS or Sudden Infant Death Syndrome (Crib Death)	1.6	0.0	3.9
b. Birth defect	4.1	0.2	8.1
c. Premature birth (born too early)	7.2	2.6	11.8
d. Miscarriage	75.3	66.9	83.7
e. Birth weight was too low	2.7	0.0	5.9
f. Pregnancy complications	6.0	1.4	10.5
g. Accident	2.4	0.0	5.6
h. Other	5.0	1.0	9.1
i. I don't know the cause of death	3.8	0.0	7.8

*Note: Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution.*

**67. Have you ever been told by a doctor, nurse, or other health care worker that you have asthma?**

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
Women who had asthma	10.1	7.8	12.5

**The following questions are about the 12 months before your new baby was born.**

**68. During the 12 months before your new baby was born, what was your yearly total household income before taxes? Include your income, your husband's or partner's income, and any other income you may have received.**

Annual total household income before taxes

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
a. Less than \$10,000	27.5	24.0	31.0
b. \$10,000 to \$14,999	14.0	11.1	16.8
c. \$15,000 to \$19,999	10.6	8.0	13.2
d. \$20,000 to \$24,999	9.0	6.6	11.3
e. \$25,000 to \$34,999	10.3	7.7	12.8
f. \$35,000 to \$49,999	10.2	7.6	12.8
g. \$50,000 or more	18.5	15.2	21.9

**69. During the 12 months before your new baby was born, how many people, including yourself, depended on this income?**

Number of people who depended on the household income

	Percent	95% CI	
		<i>Lower</i>	<i>Upper</i>
1 person	11.4	9.0	13.9
2 people	32.6	28.8	36.4
3 people	31.0	27.2	34.8
4 people	16.4	13.4	19.3
5 people	4.6	3.0	6.2
6 or more people	5.6	3.9	7.3

# Appendix

## Methods

### Purpose and History

In 1987, the Centers for Disease Control and Prevention (CDC) initiated the PRAMS project in collaboration with state health departments. The purpose was to gather data about maternal experiences and behaviors relevant to maternal and infant health, especially low birth weight and infant mortality. Thirty-seven states and New York City currently participate in this project.

Findings from PRAMS are used to enhance understanding of maternal behaviors and their relationship with adverse pregnancy outcomes. Data can be used at the state level to plan, implement, monitor and evaluate programs and to inform policymakers and the general public. PRAMS helps identify at risk populations and monitors progress over time. At the national level, PRAMS represents 78% of all U.S. births. For year 2011 births, the Phase VI questionnaire was used.

### Sampling

This description is based on information from the CDC PRAMS website <http://www.cdc.gov/prams/methodology.htm>. The PRAMS sample of women who have had a recent live birth was drawn from the state's birth certificate file. For birth year 2011, Mississippi PRAMS sampled 1,186 mothers (generalizing to a PRAMS-eligible population of 37,124). The sample was stratified by birth weight: women with very low birth weight (less than 1,500 g) or low birth weight (1,500 g to 2,499 g) infants were sampled at a higher rate. Records were excluded from the sample if the mother was a non-resident, if the mother gave birth out of state, if there was missing data from birth certificates, or if there was delayed processing of the birth certificates.

### Data Collection

Procedures and instruments are standardized to allow comparisons between states. Selected women were first contacted by mail. If there was no response, attempts were made to interview women by telephone.

### Weighting and Analysis

CDC PRAMS weighted the dataset, providing a final analysis weight that was the product of the weights for sampling, non-response, and non-coverage. Estimates for the PRAMS-eligible population were obtained by using the analysis weight and survey design variables. SAS 9.3 software was used for survey analysis.

### Variable Definitions

#### Birth Certificate Variables

The mother provided information for demographics (maternal age, race, educational level, and marital status). Age and educational levels were reported as completed years.

#### Survey Variables

##### Q4, Q5

Question 4 asks about women's weight before pregnancy. Question 5 asks about women's height before pregnancy. Both were used to measure Body Mass Index (BMI). According to CDC, BMI is calculated as weight in kilograms divided by the square of height in meters ( $\text{kg}/\text{m}^2$ ). For all adults 20 years and older, it was categorized as underweight if less than 18.5; normal if 18.5 to 24.9; overweight if 25.0 to 29.9; and obese if 30.0 or more (Body Mass Index: Considerations for Practitioners, Department of Health and Human Services Centers for Disease Controls and Prevention. Available at <http://www.cdc.gov/obesity/downloads/BMIforPractitioners.pdf>, accessed February 26<sup>th</sup>, 2014).

### ***Q6***

Question 6 asks about the women's birth date. The response was used to categorize women's ages into groups.

### ***Q9***

Question 9 asks about women's previous low birth weight births. Low birth weight is defined as birth weight less than 2,500 grams (United Nations Fund and World Health Organization. Low birth weight: country, regional, and global estimates. New York: UNICEF, 2004. Available at [http://www.childinfo.org/files/low\\_birthweight\\_from\\_EY.pdf](http://www.childinfo.org/files/low_birthweight_from_EY.pdf), accessed February 26<sup>th</sup>, 2014).

### ***Q11***

Question 11 asks about women's previous baby's age. The result is categorized into baby's age group and this question may be used to interpret birth spacing.

### ***Q26, Q27, Q28, Q29***

These questions ask about smoking habits among women. Responses were dichotomized as 'yes' or 'no'. For women who smoke, follow-up questions are asked about smoking habit questions before, during and after pregnancy.

### ***Q38, Q40***

Question 38 asks about women's admission to hospital. Question number 40 asks about women's discharge from the hospital after hospitalization. From these questions, the length of maternal hospitalization for delivery was calculated.

## **Statistical Methods**

This brief surveillance report was produced using the SAS version 9.3 with complex sample for weighted data. Descriptive statistics were used to create summary percentages from each question and the 95% confidence interval (CI). Percentages of maternal characteristics with non-overlapping 95% confidence intervals were considered statistically significant. This conservative test of significance was used for all PRAMS questionnaire items presented in this report. Trend tests were not computed. The approach for presenting point estimates and their variability is adapted from National Center for Health Statistics (NCHS). Details of NCHS guidelines are available at <http://www.cdc.gov/nchs/products/pubs/pubd/hestats/children2005/children2005.htm>. Accessed February 26<sup>th</sup>, 2014.

## **Strengths and Limitations**

### **Strengths:**

PRAMS is a population-based survey allowing generalization to all women with a live birth with relatively few exclusions. PRAMS data provides overall estimates of prevalence of maternal behaviors or health problems which are useful for monitoring trends and assessing a state's need for services (MS PRAMS Surveillance Protocol, February 19, 2011). This 2011 birth dataset represents a 63.6% response rate, slightly below the 65% response rate required by the CDC.

### **Limitations:**

Sample sizes for some subpopulations were too small for precise estimates. Responses with relative standard error above 30% are shaded in gray. Results should be interpreted with caution. PRAMS questionnaire data are self-reported and may be subject to inaccurate reporting. This may lead to recall bias, reporting bias and/or mode bias. Survey responses are also influenced by how respondents interpret the questions. If respondents interpret questions differently than expected, inaccurate data may result.

## **Survey Guide**

The survey guide was designed to provide direction for PRAMS data users on noteworthy topics in the PRAMS survey. A database of topics was developed from the Mississippi Maternal and Child Health (MCH) needs assessment and the CDC PRAMS web query categories. Each topic area is matched with questions from the survey and a brief summary of the question is provided.

**Sample and Population, year 2011 births**

Number of MS PRAMS respondents (unweighted number) and weighted estimates (numbers and percents of MS residents with an in-state live birth) by maternal characteristics used in tables.

<b>Maternal characteristics</b>	Unweighted counts	Number	<b>Weighted estimates</b>				
			95% CI <i>Lower</i>	95% CI <i>Upper</i>	Weighted percent	95% CI <i>Lower</i>	95% CI <i>Upper</i>
<b>Age (years)</b>							
All women	1,174	36,694	36,615	36,774	100.0	-	-
19 and younger	137	2,993	2,665	3,321	8.2	7.3	9.0
20-24	424	12,837	11,440	14,233	35.0	31.2	38.8
25-34	500	17,408	15,974	18,842	47.4	43.5	51.3
35+	113	3,457	2,560	4,353	9.4	7.0	11.9
<b>Race-ethnicity</b>							
All women	1,186	37,124	37,044	37,204	100.0	-	-
White	522	18,939	17,459	20,419	51.0	47.0	55.0
Black	644	17,384	15,908	18,859	46.8	42.9	50.8
Other	20	801	351	1,251	2.2	0.9	3.4
<b>Education (years completed)</b>							
All women	1,185	37,118	37,038	37,198	100.0	-	-
Less than high school	30	738	392	1,083	2.0	1.1	2.9
High school	223	6,162	5,172	7,152	16.6	13.9	19.3
Some college	389	10,874	9,551	12,196	29.3	25.7	32.9
College graduate	543	19,344	17,913	20,775	52.1	48.3	56.0
<b>Marital status</b>							
All women	1,186	37,124	37,044	37,204	100.0	-	-
Married	429	16,604	15,146	18,062	44.7	40.8	48.7
Not married	757	20,520	19,058	21,981	55.3	51.3	59.2
<b>Medicaid for PNC &amp;/or delivery</b>							
All women	1,186	37,124	37,044	37,204	100.0	-	-
No	325	13,122	11,684	14,559	35.3	31.5	39.2
Yes	861	24,002	22,563	25,442	64.7	60.8	68.5



**Your answers on this  
survey will help us improve  
the health of mothers and  
babies in Mississippi**



**MISSISSIPPI STATE DEPARTMENT OF HEALTH**

**[www.HealthyMS.com](http://www.HealthyMS.com)**

**1-866-HLTHY4U**

**(1-866-458-4948)**