

Selected Death Statistics, Residents of Adams County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 424 | 198 | 225 | 1,336.0 | 1,398.7 | 1,327.2 |
| 1 | Malignant Neoplasms | 93 | 41 | 52 | 293.0 | 289.6 | 306.7 |
| 2 | Heart Diseases | 91 | 40 | 51 | 286.7 | 282.6 | 300.8 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 29 | 22 | 7 | 91.4 | 155.4 | 41.3 |
| 4 | Diabetes Mellitus | 26 | 4 | 21 | 81.9 | 28.3 | 123.9 |
| 5 | Alzheimer's Disease | 20 | 13 | 7 | 63.0 | 91.8 | 41.3 |
| 5 | Accidents | 20 | 13 | 7 | 63.0 | 91.8 | 41.3 |
| 7 | Cerebrovascular Disease | 19 | 9 | 10 | 59.9 | 63.6 | 59.0 |
| 8 | Nephritis, Nephrotic Syndrome And Nephrosis | 17 | 4 | 13 | 53.6 | 28.3 | 76.7 |
| 9 | Septicemia | 16 | 8 | 8 | 50.4 | 56.5 | 47.2 |
| 9 | Influenza and Pneumonia | 16 | 8 | 8 | 50.4 | 56.5 | 47.2 |
| 11 | Hypertension/Hypertensive Renal Disease | 7 | 2 | 5 | 22.1 | 14.1 | 29.5 |
| 12 | Homicide | 5 | 0 | 5 | 15.8 | 0.0 | 29.5 |
| 13 | Suicide | 4 | 4 | 0 | 12.6 | 28.3 | 0.0 |
| 13 | Anemias | 4 | 0 | 4 | 12.6 | 0.0 | 23.6 |
| 15 | Chronic Liver Disease And Cirrhosis | 3 | 1 | 2 | 9.5 | 7.1 | 11.8 |
| 16 | Pneumonitis Due to Solids and Liquids | 2 | 2 | 0 | 6.3 | 14.1 | 0.0 |
| 17 | Complications Medical/Surgical Care | 1 | 0 | 1 | 3.2 | 0.0 | 5.9 |
| 17 | Parkinson's Disease | 1 | 1 | 0 | 3.2 | 7.1 | 0.0 |
| 17 | Tuberculosis | 1 | 0 | 1 | 3.2 | 0.0 | 5.9 |
| 17 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 2.5 | 0.0 | 3.7 |
| 17 | Maternal Causes** | 1 | 1 | 0 | 249.4 | 793.7 | 0.0 |
| 17 | In Situ, Benign and Unspecified Neoplasms | 1 | 0 | 1 | 3.2 | 0.0 | 5.9 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 9 | 4 | 5 | 28.4 | 28.3 | 29.5 |
| | All Other Causes | 37 | 21 | 16 | 116.6 | 148.3 | 94.4 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 424 | 198 | 225 | 209 | 102 | 107 | 215 | 96 | 118 |
| Under 1 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| (Neonatal) | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 3 | 1 | 2 | 2 | 0 | 2 | 1 | 1 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 6 | 1 | 5 | 5 | 1 | 4 | 1 | 0 | 1 |
| 25-34 | 13 | 5 | 8 | 9 | 4 | 5 | 4 | 1 | 3 |
| 35-44 | 10 | 2 | 8 | 6 | 1 | 5 | 4 | 1 | 3 |
| 45-54 | 38 | 11 | 26 | 22 | 7 | 15 | 16 | 4 | 11 |
| 55-64 | 61 | 25 | 36 | 30 | 15 | 15 | 31 | 10 | 21 |
| 65-74 | 84 | 41 | 43 | 47 | 24 | 23 | 37 | 17 | 20 |
| 75-84 | 106 | 52 | 54 | 51 | 26 | 25 | 55 | 26 | 29 |
| 85 and over | 102 | 60 | 42 | 36 | 24 | 12 | 66 | 36 | 30 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 31,737 |
| White | 14,156 |
| Black | 16,953 |
| Other | 628 |

Selected Death Statistics, Residents of Alcorn County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 482 | 436 | 45 | 1,289.5 | 1,360.5 | 961.9 |
| 1 | Heart Diseases | 134 | 123 | 11 | 358.5 | 383.8 | 235.1 |
| 2 | Malignant Neoplasms | 85 | 78 | 6 | 227.4 | 243.4 | 128.3 |
| 3 | Alzheimer's Disease | 28 | 28 | 0 | 74.9 | 87.4 | 0.0 |
| 4 | Emphysema and Other Chronic Lower Resp. Diseases | 24 | 24 | 0 | 64.2 | 74.9 | 0.0 |
| 4 | Accidents | 24 | 22 | 2 | 64.2 | 68.7 | 42.8 |
| 6 | Cerebrovascular Disease | 23 | 23 | 0 | 61.5 | 71.8 | 0.0 |
| 7 | Influenza and Pneumonia | 17 | 15 | 2 | 45.5 | 46.8 | 42.8 |
| 8 | Nephritis, Nephrotic Syndrome And Nephrosis | 15 | 11 | 4 | 40.1 | 34.3 | 85.5 |
| 9 | Septicemia | 12 | 10 | 2 | 32.1 | 31.2 | 42.8 |
| 10 | Chronic Liver Disease And Cirrhosis | 10 | 9 | 1 | 26.8 | 28.1 | 21.4 |
| 11 | Diabetes Mellitus | 7 | 3 | 4 | 18.7 | 9.4 | 85.5 |
| 12 | Homicide | 6 | 5 | 1 | 16.1 | 15.6 | 21.4 |
| 12 | Suicide | 6 | 6 | 0 | 16.1 | 18.7 | 0.0 |
| 14 | Pneumonitis Due to Solids and Liquids | 4 | 3 | 1 | 10.7 | 9.4 | 21.4 |
| 15 | Viral Hepatitis | 2 | 2 | 0 | 5.4 | 6.2 | 0.0 |
| 15 | Hypertension/Hypertensive Renal Disease | 2 | 2 | 0 | 5.4 | 6.2 | 0.0 |
| 17 | Complications Medical/Surgical Care | 1 | 0 | 1 | 2.7 | 0.0 | 21.4 |
| 17 | HIV Disease | 1 | 1 | 0 | 2.7 | 3.1 | 0.0 |
| 17 | Malnutrition/Nutritional Deficiency | 1 | 1 | 0 | 2.7 | 3.1 | 0.0 |
| 17 | Parkinson's Disease | 1 | 1 | 0 | 2.7 | 3.1 | 0.0 |
| 17 | Maternal Causes** | 1 | 1 | 0 | 241.0 | 291.5 | 0.0 |
| 17 | Congenital Malformations and Chrom. Abnormalities | 1 | 1 | 0 | 2.7 | 3.1 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 16 | 14 | 2 | 42.8 | 43.7 | 42.8 |
| | All Other Causes | 61 | 53 | 8 | 163.2 | 165.4 | 171.0 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 482 | 436 | 45 | 230 | 210 | 19 | 252 | 226 | 26 |
| Under 1 | 3 | 3 | 0 | 1 | 1 | 0 | 2 | 2 | 0 |
| (Neonatal) | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 |
| (Postneonatal) | 2 | 2 | 0 | 1 | 1 | 0 | 1 | 1 | 0 |
| 1-4 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| 15-24 | 5 | 5 | 0 | 4 | 4 | 0 | 1 | 1 | 0 |
| 25-34 | 6 | 5 | 1 | 4 | 4 | 0 | 2 | 1 | 1 |
| 35-44 | 19 | 17 | 2 | 12 | 12 | 0 | 7 | 5 | 2 |
| 45-54 | 34 | 27 | 7 | 18 | 15 | 3 | 16 | 12 | 4 |
| 55-64 | 71 | 65 | 6 | 46 | 42 | 4 | 25 | 23 | 2 |
| 65-74 | 104 | 91 | 13 | 54 | 46 | 8 | 50 | 45 | 5 |
| 75-84 | 116 | 107 | 8 | 54 | 50 | 3 | 62 | 57 | 5 |
| 85 and over | 122 | 116 | 6 | 36 | 36 | 0 | 86 | 80 | 6 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 37,380 |
| White | 32,046 |
| Black | 4,678 |
| Other | 656 |

Selected Death Statistics, Residents of Amite County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 154 | 104 | 50 | 1,219.4 | 1,421.2 | 968.2 |
| 1 | Heart Diseases | 37 | 28 | 9 | 293.0 | 382.6 | 174.3 |
| 2 | Malignant Neoplasms | 28 | 15 | 13 | 221.7 | 205.0 | 251.7 |
| 3 | Accidents | 12 | 9 | 3 | 95.0 | 123.0 | 58.1 |
| 4 | Emphysema and Other Chronic Lower Resp. Diseases | 10 | 9 | 1 | 79.2 | 123.0 | 19.4 |
| 5 | Septicemia | 7 | 6 | 1 | 55.4 | 82.0 | 19.4 |
| 5 | Diabetes Mellitus | 7 | 4 | 3 | 55.4 | 54.7 | 58.1 |
| 7 | Influenza and Pneumonia | 6 | 4 | 2 | 47.5 | 54.7 | 38.7 |
| 8 | Cerebrovascular Disease | 4 | 2 | 2 | 31.7 | 27.3 | 38.7 |
| 8 | Alzheimer's Disease | 4 | 3 | 1 | 31.7 | 41.0 | 19.4 |
| 10 | Homicide | 3 | 0 | 3 | 23.8 | 0.0 | 58.1 |
| 10 | Hypertension/Hypertensive Renal Disease | 3 | 3 | 0 | 23.8 | 41.0 | 0.0 |
| 12 | Congenital Malformations and Chrom. Abnormalities | 2 | 1 | 1 | 15.8 | 13.7 | 19.4 |
| 13 | Suicide | 1 | 1 | 0 | 7.9 | 13.7 | 0.0 |
| 13 | Chronic Liver Disease And Cirrhosis | 1 | 1 | 0 | 7.9 | 13.7 | 0.0 |
| 13 | Infections of Kidney | 1 | 1 | 0 | 7.9 | 13.7 | 0.0 |
| 13 | Pneumonitis Due to Solids and Liquids | 1 | 1 | 0 | 7.9 | 13.7 | 0.0 |
| 13 | Nephritis, Nephrotic Syndrome And Nephrosis | 1 | 1 | 0 | 7.9 | 13.7 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 1 | 1 | 0 | 7.9 | 13.7 | 0.0 |
| | All Other Causes | 25 | 14 | 11 | 198.0 | 191.3 | 213.0 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 154 | 104 | 50 | 82 | 56 | 26 | 72 | 48 | 24 |
| Under 1 | 3 | 1 | 2 | 1 | 1 | 0 | 2 | 0 | 2 |
| (Neonatal) | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 2 | 0 | 2 | 0 | 0 | 0 | 2 | 0 | 2 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 2 | 1 | 1 | 1 | 0 | 1 | 1 | 1 | 0 |
| 25-34 | 2 | 2 | 0 | 2 | 2 | 0 | 0 | 0 | 0 |
| 35-44 | 7 | 3 | 4 | 3 | 2 | 1 | 4 | 1 | 3 |
| 45-54 | 11 | 9 | 2 | 4 | 3 | 1 | 7 | 6 | 1 |
| 55-64 | 30 | 11 | 19 | 20 | 7 | 13 | 10 | 4 | 6 |
| 65-74 | 26 | 20 | 6 | 15 | 12 | 3 | 11 | 8 | 3 |
| 75-84 | 41 | 33 | 8 | 19 | 16 | 3 | 22 | 17 | 5 |
| 85 and over | 32 | 24 | 8 | 17 | 13 | 4 | 15 | 11 | 4 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 12,629 |
| White | 7,318 |
| Black | 5,164 |
| Other | 147 |

Selected Death Statistics, Residents of Attala County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 283 | 194 | 88 | 1,476.8 | 1,795.0 | 1,085.3 |
| 1 | Heart Diseases | 72 | 50 | 22 | 375.7 | 462.6 | 271.3 |
| 2 | Malignant Neoplasms | 57 | 38 | 19 | 297.4 | 351.6 | 234.3 |
| 3 | Alzheimer's Disease | 25 | 23 | 2 | 130.5 | 212.8 | 24.7 |
| 4 | Diabetes Mellitus | 14 | 6 | 8 | 73.1 | 55.5 | 98.7 |
| 5 | Emphysema and Other Chronic Lower Resp. Diseases | 12 | 7 | 5 | 62.6 | 64.8 | 61.7 |
| 6 | Cerebrovascular Disease | 11 | 8 | 3 | 57.4 | 74.0 | 37.0 |
| 7 | Accidents | 10 | 9 | 1 | 52.2 | 83.3 | 12.3 |
| 8 | Nephritis, Nephrotic Syndrome And Nephrosis | 8 | 2 | 6 | 41.7 | 18.5 | 74.0 |
| 9 | Influenza and Pneumonia | 6 | 4 | 2 | 31.3 | 37.0 | 24.7 |
| 9 | Septicemia | 6 | 3 | 3 | 31.3 | 27.8 | 37.0 |
| 9 | Pneumonitis Due to Solids and Liquids | 6 | 5 | 1 | 31.3 | 46.3 | 12.3 |
| 12 | Congenital Malformations and Chrom. Abnormalities | 2 | 1 | 1 | 10.4 | 9.3 | 12.3 |
| 12 | Chronic Liver Disease And Cirrhosis | 2 | 2 | 0 | 10.4 | 18.5 | 0.0 |
| 12 | Aortic Aneurysm and Dissection | 2 | 2 | 0 | 10.4 | 18.5 | 0.0 |
| 15 | Parkinson's Disease | 1 | 1 | 0 | 5.2 | 9.3 | 0.0 |
| 15 | Homicide | 1 | 1 | 0 | 5.2 | 9.3 | 0.0 |
| 15 | Conditions Originating in the Perinatal Period** | 1 | 0 | 0 | 4.2 | 0.0 | 0.0 |
| 15 | HIV Disease | 1 | 0 | 1 | 5.2 | 0.0 | 12.3 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 3 | 2 | 1 | 15.7 | 18.5 | 12.3 |
| | All Other Causes | 43 | 30 | 13 | 224.4 | 277.6 | 160.3 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total* ----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|--------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 283 | 194 | 88 | 153 | 111 | 42 | 130 | 83 | 46 |
| Under 1 | 3 | 1 | 1 | 1 | 1 | 0 | 2 | 0 | 1 |
| (Neonatal) | 3 | 1 | 1 | 1 | 1 | 0 | 2 | 0 | 1 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 25-34 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| 35-44 | 9 | 8 | 1 | 7 | 6 | 1 | 2 | 2 | 0 |
| 45-54 | 17 | 7 | 10 | 10 | 4 | 6 | 7 | 3 | 4 |
| 55-64 | 34 | 16 | 18 | 21 | 10 | 11 | 13 | 6 | 7 |
| 65-74 | 55 | 40 | 15 | 28 | 22 | 6 | 27 | 18 | 9 |
| 75-84 | 67 | 48 | 19 | 36 | 28 | 8 | 31 | 20 | 11 |
| 85 and over | 97 | 74 | 23 | 50 | 40 | 10 | 47 | 34 | 13 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 19,163 |
| White | 10,808 |
| Black | 8,108 |
| Other | 247 |

Selected Death Statistics, Residents of Benton County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 112 | 74 | 38 | 1,350.0 | 1,433.6 | 1,263.7 |
| 1 | Malignant Neoplasms | 30 | 19 | 11 | 361.6 | 368.1 | 365.8 |
| 2 | Heart Diseases | 29 | 21 | 8 | 349.6 | 406.8 | 266.0 |
| 3 | Accidents | 11 | 7 | 4 | 132.6 | 135.6 | 133.0 |
| 4 | Diabetes Mellitus | 6 | 0 | 6 | 72.3 | 0.0 | 199.5 |
| 4 | Alzheimer's Disease | 6 | 5 | 1 | 72.3 | 96.9 | 33.3 |
| 6 | Emphysema and Other Chronic Lower Resp. Diseases | 4 | 4 | 0 | 48.2 | 77.5 | 0.0 |
| 7 | Septicemia | 3 | 3 | 0 | 36.2 | 58.1 | 0.0 |
| 7 | Influenza and Pneumonia | 3 | 1 | 2 | 36.2 | 19.4 | 66.5 |
| 9 | Homicide | 1 | 0 | 1 | 12.1 | 0.0 | 33.3 |
| 9 | Tuberculosis | 1 | 1 | 0 | 12.1 | 19.4 | 0.0 |
| 9 | Pneumonitis Due to Solids and Liquids | 1 | 1 | 0 | 12.1 | 19.4 | 0.0 |
| 9 | Chronic Liver Disease And Cirrhosis | 1 | 0 | 1 | 12.1 | 0.0 | 33.3 |
| 9 | Suicide | 1 | 1 | 0 | 12.1 | 19.4 | 0.0 |
| 9 | Nephritis, Nephrotic Syndrome And Nephrosis | 1 | 1 | 0 | 12.1 | 19.4 | 0.0 |
| 9 | Cerebrovascular Disease | 1 | 1 | 0 | 12.1 | 19.4 | 0.0 |
| 9 | Parkinson's Disease | 1 | 1 | 0 | 12.1 | 19.4 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 0 | 0 | 0 | 0.0 | 0.0 | 0.0 |
| | All Other Causes | 12 | 8 | 4 | 144.6 | 155.0 | 133.0 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 112 | 74 | 38 | 57 | 38 | 19 | 55 | 36 | 19 |
| Under 1 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| (Neonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 25-34 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 35-44 | 5 | 4 | 1 | 3 | 2 | 1 | 2 | 2 | 0 |
| 45-54 | 6 | 4 | 2 | 5 | 3 | 2 | 1 | 1 | 0 |
| 55-64 | 17 | 11 | 6 | 7 | 5 | 2 | 10 | 6 | 4 |
| 65-74 | 21 | 15 | 6 | 11 | 9 | 2 | 10 | 6 | 4 |
| 75-84 | 33 | 21 | 12 | 16 | 10 | 6 | 17 | 11 | 6 |
| 85 and over | 27 | 17 | 10 | 12 | 7 | 5 | 15 | 10 | 5 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|-------|
| Total | 8,296 |
| White | 5,162 |
| Black | 3,007 |
| Other | 127 |

Selected Death Statistics, Residents of Bolivar County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 391 | 140 | 249 | 1,157.9 | 1,227.0 | 1,140.4 |
| 1 | Heart Diseases | 97 | 36 | 60 | 287.3 | 315.5 | 274.8 |
| 2 | Malignant Neoplasms | 76 | 26 | 50 | 225.1 | 227.9 | 229.0 |
| 3 | Hypertension/Hypertensive Renal Disease | 21 | 1 | 20 | 62.2 | 8.8 | 91.6 |
| 4 | Cerebrovascular Disease | 19 | 7 | 12 | 56.3 | 61.3 | 55.0 |
| 5 | Emphysema and Other Chronic Lower Resp. Diseases | 18 | 9 | 9 | 53.3 | 78.9 | 41.2 |
| 5 | Accidents | 18 | 8 | 10 | 53.3 | 70.1 | 45.8 |
| 7 | Diabetes Mellitus | 16 | 2 | 14 | 47.4 | 17.5 | 64.1 |
| 8 | Alzheimer's Disease | 14 | 9 | 5 | 41.5 | 78.9 | 22.9 |
| 9 | Nephritis, Nephrotic Syndrome And Nephrosis | 11 | 4 | 7 | 32.6 | 35.1 | 32.1 |
| 10 | Septicemia | 10 | 2 | 8 | 29.6 | 17.5 | 36.6 |
| 11 | Influenza and Pneumonia | 9 | 5 | 4 | 26.7 | 43.8 | 18.3 |
| 12 | Suicide | 6 | 4 | 2 | 17.8 | 35.1 | 9.2 |
| 13 | Homicide | 4 | 1 | 3 | 11.8 | 8.8 | 13.7 |
| 14 | Chronic Liver Disease And Cirrhosis | 3 | 0 | 2 | 8.9 | 0.0 | 9.2 |
| 14 | Anemias | 3 | 0 | 3 | 8.9 | 0.0 | 13.7 |
| 16 | Pneumonitis Due to Solids and Liquids | 2 | 0 | 2 | 5.9 | 0.0 | 9.2 |
| 16 | Aortic Aneurysm and Dissection | 2 | 0 | 2 | 5.9 | 0.0 | 9.2 |
| 16 | In Situ, Benign and Unspecified Neoplasms | 2 | 1 | 1 | 5.9 | 8.8 | 4.6 |
| 19 | Congenital Malformations and Chrom. Abnormalities | 1 | 0 | 1 | 3.0 | 0.0 | 4.6 |
| 19 | Diseases of Appendix | 1 | 1 | 0 | 3.0 | 8.8 | 0.0 |
| 19 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 2.1 | 0.0 | 2.9 |
| 19 | Parkinson's Disease | 1 | 0 | 1 | 3.0 | 0.0 | 4.6 |
| 19 | Malnutrition/Nutritional Deficiency | 1 | 1 | 0 | 3.0 | 8.8 | 0.0 |
| 19 | HIV Disease | 1 | 0 | 1 | 3.0 | 0.0 | 4.6 |
| 19 | Meningitis | 1 | 0 | 1 | 3.0 | 0.0 | 4.6 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 4 | 1 | 3 | 11.8 | 8.8 | 13.7 |
| | All Other Causes | 49 | 22 | 27 | 145.1 | 192.8 | 123.7 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 391 | 140 | 249 | 204 | 72 | 130 | 187 | 68 | 119 |
| Under 1 | 2 | 0 | 2 | 2 | 0 | 2 | 0 | 0 | 0 |
| (Neonatal) | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| (Postneonatal) | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 4 | 1 | 2 | 4 | 1 | 2 | 0 | 0 | 0 |
| 25-34 | 12 | 3 | 9 | 8 | 2 | 6 | 4 | 1 | 3 |
| 35-44 | 18 | 4 | 14 | 8 | 3 | 5 | 10 | 1 | 9 |
| 45-54 | 31 | 7 | 24 | 14 | 3 | 11 | 17 | 4 | 13 |
| 55-64 | 78 | 21 | 56 | 56 | 16 | 39 | 22 | 5 | 17 |
| 65-74 | 89 | 26 | 63 | 51 | 12 | 39 | 38 | 14 | 24 |
| 75-84 | 85 | 35 | 50 | 38 | 17 | 21 | 47 | 18 | 29 |
| 85 and over | 72 | 43 | 29 | 23 | 18 | 5 | 49 | 25 | 24 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 33,768 |
| White | 11,410 |
| Black | 21,834 |
| Other | 524 |

Selected Death Statistics, Residents of Calhoun County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 176 | 144 | 32 | 1,193.6 | 1,384.5 | 774.3 |
| 1 | Malignant Neoplasms | 51 | 41 | 10 | 345.9 | 394.2 | 242.0 |
| 2 | Heart Diseases | 48 | 41 | 7 | 325.5 | 394.2 | 169.4 |
| 3 | Alzheimer's Disease | 15 | 14 | 1 | 101.7 | 134.6 | 24.2 |
| 4 | Cerebrovascular Disease | 11 | 8 | 3 | 74.6 | 76.9 | 72.6 |
| 5 | Hypertension/Hypertensive Renal Disease | 7 | 6 | 1 | 47.5 | 57.7 | 24.2 |
| 6 | Accidents | 6 | 4 | 2 | 40.7 | 38.5 | 48.4 |
| 7 | Septicemia | 4 | 3 | 1 | 27.1 | 28.8 | 24.2 |
| 8 | Influenza and Pneumonia | 3 | 2 | 1 | 20.3 | 19.2 | 24.2 |
| 8 | Suicide | 3 | 1 | 2 | 20.3 | 9.6 | 48.4 |
| 8 | Nephritis, Nephrotic Syndrome And Nephrosis | 3 | 3 | 0 | 20.3 | 28.8 | 0.0 |
| 8 | Emphysema and Other Chronic Lower Resp. Diseases | 3 | 3 | 0 | 20.3 | 28.8 | 0.0 |
| 8 | Chronic Liver Disease And Cirrhosis | 3 | 1 | 2 | 20.3 | 9.6 | 48.4 |
| 13 | Diabetes Mellitus | 2 | 1 | 1 | 13.6 | 9.6 | 24.2 |
| 13 | In Situ, Benign and Unspecified Neoplasms | 2 | 2 | 0 | 13.6 | 19.2 | 0.0 |
| 15 | Viral Hepatitis | 1 | 1 | 0 | 6.8 | 9.6 | 0.0 |
| 15 | Pneumonitis Due to Solids and Liquids | 1 | 1 | 0 | 6.8 | 9.6 | 0.0 |
| 15 | Complications Medical/Surgical Care | 1 | 1 | 0 | 6.8 | 9.6 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 0 | 0 | 0 | 0.0 | 0.0 | 0.0 |
| | All Other Causes | 12 | 11 | 1 | 81.4 | 105.8 | 24.2 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 176 | 144 | 32 | 92 | 78 | 14 | 84 | 66 | 18 |
| Under 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Neonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 3 | 2 | 1 | 1 | 0 | 1 | 2 | 2 | 0 |
| 25-34 | 2 | 1 | 1 | 2 | 1 | 1 | 0 | 0 | 0 |
| 35-44 | 4 | 2 | 2 | 2 | 1 | 1 | 2 | 1 | 1 |
| 45-54 | 13 | 4 | 9 | 8 | 3 | 5 | 5 | 1 | 4 |
| 55-64 | 30 | 25 | 5 | 19 | 17 | 2 | 11 | 8 | 3 |
| 65-74 | 37 | 31 | 6 | 24 | 22 | 2 | 13 | 9 | 4 |
| 75-84 | 43 | 40 | 3 | 21 | 20 | 1 | 22 | 20 | 2 |
| 85 and over | 44 | 39 | 5 | 15 | 14 | 1 | 29 | 25 | 4 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 14,745 |
| White | 10,401 |
| Black | 4,133 |
| Other | 211 |

Selected Death Statistics, Residents of Carroll County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 128 | 102 | 25 | 1,248.3 | 1,536.8 | 717.6 |
| 1 | Heart Diseases | 40 | 27 | 12 | 390.1 | 406.8 | 344.4 |
| 2 | Malignant Neoplasms | 28 | 24 | 4 | 273.1 | 361.6 | 114.8 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 15 | 13 | 2 | 146.3 | 195.9 | 57.4 |
| 4 | Cerebrovascular Disease | 6 | 5 | 1 | 58.5 | 75.3 | 28.7 |
| 4 | Accidents | 6 | 4 | 2 | 58.5 | 60.3 | 57.4 |
| 6 | Influenza and Pneumonia | 3 | 2 | 1 | 29.3 | 30.1 | 28.7 |
| 7 | Hypertension/Hypertensive Renal Disease | 2 | 2 | 0 | 19.5 | 30.1 | 0.0 |
| 7 | Septicemia | 2 | 2 | 0 | 19.5 | 30.1 | 0.0 |
| 9 | Suicide | 1 | 0 | 1 | 9.8 | 0.0 | 28.7 |
| 9 | Complications Medical/Surgical Care | 1 | 0 | 1 | 9.8 | 0.0 | 28.7 |
| 9 | Peptic Ulcer | 1 | 1 | 0 | 9.8 | 15.1 | 0.0 |
| 9 | Congenital Malformations and Chrom. Abnormalities | 1 | 1 | 0 | 9.8 | 15.1 | 0.0 |
| 9 | Pneumonitis Due to Solids and Liquids | 1 | 1 | 0 | 9.8 | 15.1 | 0.0 |
| 9 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 10.0 | 0.0 | 28.6 |
| 9 | Chronic Liver Disease And Cirrhosis | 1 | 1 | 0 | 9.8 | 15.1 | 0.0 |
| 9 | Alzheimer's Disease | 1 | 1 | 0 | 9.8 | 15.1 | 0.0 |
| 9 | Diabetes Mellitus | 1 | 1 | 0 | 9.8 | 15.1 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 3 | 3 | 0 | 29.3 | 45.2 | 0.0 |
| | All Other Causes | 14 | 14 | 0 | 136.5 | 210.9 | 0.0 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 128 | 102 | 25 | 70 | 53 | 16 | 58 | 49 | 9 |
| Under 1 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| (Neonatal) | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 25-34 | 2 | 2 | 0 | 1 | 1 | 0 | 1 | 1 | 0 |
| 35-44 | 2 | 2 | 0 | 0 | 0 | 0 | 2 | 2 | 0 |
| 45-54 | 7 | 4 | 2 | 5 | 2 | 2 | 2 | 2 | 0 |
| 55-64 | 25 | 22 | 3 | 16 | 14 | 2 | 9 | 8 | 1 |
| 65-74 | 27 | 20 | 7 | 20 | 15 | 5 | 7 | 5 | 2 |
| 75-84 | 29 | 26 | 3 | 17 | 16 | 1 | 12 | 10 | 2 |
| 85 and over | 34 | 26 | 8 | 9 | 5 | 4 | 25 | 21 | 4 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 10,254 |
| White | 6,637 |
| Black | 3,484 |
| Other | 133 |

Selected Death Statistics, Residents of Chickasaw County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 182 | 125 | 56 | 1,051.2 | 1,330.9 | 736.5 |
| 1 | Heart Diseases | 45 | 36 | 9 | 259.9 | 383.3 | 118.4 |
| 2 | Malignant Neoplasms | 36 | 23 | 13 | 207.9 | 244.9 | 171.0 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 17 | 15 | 2 | 98.2 | 159.7 | 26.3 |
| 4 | Accidents | 14 | 10 | 4 | 80.9 | 106.5 | 52.6 |
| 5 | Cerebrovascular Disease | 11 | 6 | 5 | 63.5 | 63.9 | 65.8 |
| 6 | Alzheimer's Disease | 10 | 9 | 1 | 57.8 | 95.8 | 13.2 |
| 7 | Nephritis, Nephrotic Syndrome And Nephrosis | 9 | 3 | 6 | 52.0 | 31.9 | 78.9 |
| 8 | Pneumonitis Due to Solids and Liquids | 5 | 3 | 2 | 28.9 | 31.9 | 26.3 |
| 9 | Diabetes Mellitus | 4 | 2 | 2 | 23.1 | 21.3 | 26.3 |
| 9 | Septicemia | 4 | 3 | 1 | 23.1 | 31.9 | 13.2 |
| 11 | Influenza and Pneumonia | 3 | 2 | 1 | 17.3 | 21.3 | 13.2 |
| 12 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 3.8 | 0.0 | 7.4 |
| 12 | Aortic Aneurysm and Dissection | 1 | 1 | 0 | 5.8 | 10.6 | 0.0 |
| 12 | Chronic Liver Disease And Cirrhosis | 1 | 1 | 0 | 5.8 | 10.6 | 0.0 |
| 12 | Peptic Ulcer | 1 | 1 | 0 | 5.8 | 10.6 | 0.0 |
| 12 | Hypertension/Hypertensive Renal Disease | 1 | 1 | 0 | 5.8 | 10.6 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 4 | 2 | 2 | 23.1 | 21.3 | 26.3 |
| | All Other Causes | 15 | 7 | 7 | 86.6 | 74.5 | 92.1 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 182 | 125 | 56 | 86 | 56 | 29 | 96 | 69 | 27 |
| Under 1 | 2 | 0 | 2 | 1 | 0 | 1 | 1 | 0 | 1 |
| (Neonatal) | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| (Postneonatal) | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 25-34 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 35-44 | 5 | 4 | 1 | 5 | 4 | 1 | 0 | 0 | 0 |
| 45-54 | 11 | 6 | 5 | 7 | 4 | 3 | 4 | 2 | 2 |
| 55-64 | 22 | 12 | 9 | 12 | 5 | 6 | 10 | 7 | 3 |
| 65-74 | 36 | 23 | 13 | 18 | 10 | 8 | 18 | 13 | 5 |
| 75-84 | 50 | 39 | 11 | 22 | 18 | 4 | 28 | 21 | 7 |
| 85 and over | 55 | 40 | 15 | 20 | 14 | 6 | 35 | 26 | 9 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 17,313 |
| White | 9,392 |
| Black | 7,604 |
| Other | 317 |

Selected Death Statistics, Residents of Choctaw County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 89 | 65 | 23 | 1,073.1 | 1,142.8 | 939.9 |
| 1 | Heart Diseases | 24 | 17 | 7 | 289.4 | 298.9 | 286.1 |
| 2 | Malignant Neoplasms | 20 | 13 | 7 | 241.1 | 228.6 | 286.1 |
| 3 | Cerebrovascular Disease | 11 | 8 | 3 | 132.6 | 140.6 | 122.6 |
| 4 | Accidents | 10 | 9 | 0 | 120.6 | 158.2 | 0.0 |
| 5 | Alzheimer's Disease | 5 | 5 | 0 | 60.3 | 87.9 | 0.0 |
| 6 | Emphysema and Other Chronic Lower Resp. Diseases | 4 | 3 | 1 | 48.2 | 52.7 | 40.9 |
| 7 | Influenza and Pneumonia | 2 | 1 | 1 | 24.1 | 17.6 | 40.9 |
| 7 | Nephritis, Nephrotic Syndrome And Nephrosis | 2 | 1 | 1 | 24.1 | 17.6 | 40.9 |
| 9 | Chronic Liver Disease And Cirrhosis | 1 | 1 | 0 | 12.1 | 17.6 | 0.0 |
| 9 | Suicide | 1 | 1 | 0 | 12.1 | 17.6 | 0.0 |
| 9 | In Situ, Benign and Unspecified Neoplasms | 1 | 1 | 0 | 12.1 | 17.6 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 0 | 0 | 0 | 0.0 | 0.0 | 0.0 |
| | All Other Causes | 8 | 5 | 3 | 96.5 | 87.9 | 122.6 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 89 | 65 | 23 | 47 | 35 | 11 | 42 | 30 | 12 |
| Under 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Neonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 2 | 1 | 1 | 2 | 1 | 1 | 0 | 0 | 0 |
| 25-34 | 3 | 0 | 2 | 2 | 0 | 1 | 1 | 0 | 1 |
| 35-44 | 6 | 3 | 3 | 3 | 3 | 0 | 3 | 0 | 3 |
| 45-54 | 5 | 3 | 2 | 4 | 2 | 2 | 1 | 1 | 0 |
| 55-64 | 14 | 8 | 6 | 8 | 4 | 4 | 6 | 4 | 2 |
| 65-74 | 14 | 12 | 2 | 6 | 4 | 2 | 8 | 8 | 0 |
| 75-84 | 24 | 19 | 5 | 15 | 14 | 1 | 9 | 5 | 4 |
| 85 and over | 21 | 19 | 2 | 7 | 7 | 0 | 14 | 12 | 2 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|-------|
| Total | 8,294 |
| White | 5,688 |
| Black | 2,447 |
| Other | 159 |

Selected Death Statistics, Residents of Claiborne County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 126 | 31 | 95 | 1,387.7 | 2,392.0 | 1,244.1 |
| 1 | Heart Diseases | 48 | 14 | 34 | 528.6 | 1,080.2 | 445.3 |
| 2 | Malignant Neoplasms | 20 | 4 | 16 | 220.3 | 308.6 | 209.5 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 10 | 4 | 6 | 110.1 | 308.6 | 78.6 |
| 4 | Accidents | 6 | 0 | 6 | 66.1 | 0.0 | 78.6 |
| 4 | Cerebrovascular Disease | 6 | 1 | 5 | 66.1 | 77.2 | 65.5 |
| 6 | Nephritis, Nephrotic Syndrome And Nephrosis | 5 | 1 | 4 | 55.1 | 77.2 | 52.4 |
| 7 | Conditions Originating in the Perinatal Period** | 4 | 2 | 2 | 30.3 | 181.8 | 16.8 |
| 8 | Homicide | 3 | 0 | 3 | 33.0 | 0.0 | 39.3 |
| 8 | Alzheimer's Disease | 3 | 0 | 3 | 33.0 | 0.0 | 39.3 |
| 10 | Pneumonitis Due to Solids and Liquids | 2 | 0 | 2 | 22.0 | 0.0 | 26.2 |
| 10 | Diabetes Mellitus | 2 | 0 | 2 | 22.0 | 0.0 | 26.2 |
| 10 | Septicemia | 2 | 0 | 2 | 22.0 | 0.0 | 26.2 |
| 13 | Influenza and Pneumonia | 1 | 1 | 0 | 11.0 | 77.2 | 0.0 |
| 13 | Viral Hepatitis | 1 | 0 | 1 | 11.0 | 0.0 | 13.1 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 2 | 0 | 2 | 22.0 | 0.0 | 26.2 |
| | All Other Causes | 11 | 4 | 7 | 121.1 | 308.6 | 91.7 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births.

Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 126 | 31 | 95 | 66 | 13 | 53 | 60 | 18 | 42 |
| Under 1 | 4 | 2 | 2 | 1 | 0 | 1 | 3 | 2 | 1 |
| (Neonatal) | 4 | 2 | 2 | 1 | 0 | 1 | 3 | 2 | 1 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| 25-34 | 5 | 0 | 5 | 3 | 0 | 3 | 2 | 0 | 2 |
| 35-44 | 5 | 0 | 5 | 3 | 0 | 3 | 2 | 0 | 2 |
| 45-54 | 5 | 1 | 4 | 2 | 1 | 1 | 3 | 0 | 3 |
| 55-64 | 22 | 5 | 17 | 15 | 4 | 11 | 7 | 1 | 6 |
| 65-74 | 25 | 6 | 19 | 18 | 3 | 15 | 7 | 3 | 4 |
| 75-84 | 27 | 6 | 21 | 16 | 2 | 14 | 11 | 4 | 7 |
| 85 and over | 32 | 11 | 21 | 8 | 3 | 5 | 24 | 8 | 16 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|-------|
| Total | 9,080 |
| White | 1,296 |
| Black | 7,636 |
| Other | 148 |

Selected Death Statistics, Residents of Clarke County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 233 | 161 | 72 | 1,429.5 | 1,533.5 | 1,289.9 |
| 1 | Heart Diseases | 73 | 48 | 25 | 447.9 | 457.2 | 447.9 |
| 2 | Malignant Neoplasms | 42 | 30 | 12 | 257.7 | 285.7 | 215.0 |
| 3 | Accidents | 15 | 10 | 5 | 92.0 | 95.2 | 89.6 |
| 4 | Cerebrovascular Disease | 13 | 6 | 7 | 79.8 | 57.1 | 125.4 |
| 4 | Alzheimer's Disease | 13 | 9 | 4 | 79.8 | 85.7 | 71.7 |
| 6 | Emphysema and Other Chronic Lower Resp. Diseases | 11 | 10 | 1 | 67.5 | 95.2 | 17.9 |
| 7 | Septicemia | 8 | 6 | 2 | 49.1 | 57.1 | 35.8 |
| 8 | Influenza and Pneumonia | 7 | 5 | 2 | 42.9 | 47.6 | 35.8 |
| 9 | Diabetes Mellitus | 3 | 1 | 2 | 18.4 | 9.5 | 35.8 |
| 9 | Chronic Liver Disease And Cirrhosis | 3 | 2 | 1 | 18.4 | 19.0 | 17.9 |
| 11 | Nephritis, Nephrotic Syndrome And Nephrosis | 2 | 1 | 1 | 12.3 | 9.5 | 17.9 |
| 11 | Conditions Originating in the Perinatal Period** | 2 | 2 | 0 | 10.5 | 16.1 | 0.0 |
| 11 | Parkinson's Disease | 2 | 2 | 0 | 12.3 | 19.0 | 0.0 |
| 14 | Suicide | 1 | 1 | 0 | 6.1 | 9.5 | 0.0 |
| 14 | Hypertension/Hypertensive Renal Disease | 1 | 0 | 1 | 6.1 | 0.0 | 17.9 |
| 14 | HIV Disease | 1 | 0 | 1 | 6.1 | 0.0 | 17.9 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 5 | 3 | 2 | 30.7 | 28.6 | 35.8 |
| | All Other Causes | 31 | 25 | 6 | 190.2 | 238.1 | 107.5 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 233 | 161 | 72 | 111 | 78 | 33 | 122 | 83 | 39 |
| Under 1 | 3 | 2 | 1 | 2 | 1 | 1 | 1 | 1 | 0 |
| (Neonatal) | 2 | 2 | 0 | 1 | 1 | 0 | 1 | 1 | 0 |
| (Postneonatal) | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 1-4 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 4 | 1 | 3 | 2 | 0 | 2 | 2 | 1 | 1 |
| 25-34 | 4 | 1 | 3 | 2 | 1 | 1 | 2 | 0 | 2 |
| 35-44 | 4 | 2 | 2 | 1 | 0 | 1 | 3 | 2 | 1 |
| 45-54 | 18 | 11 | 7 | 13 | 8 | 5 | 5 | 3 | 2 |
| 55-64 | 34 | 22 | 12 | 20 | 14 | 6 | 14 | 8 | 6 |
| 65-74 | 42 | 27 | 15 | 20 | 15 | 5 | 22 | 12 | 10 |
| 75-84 | 64 | 52 | 12 | 32 | 24 | 8 | 32 | 28 | 4 |
| 85 and over | 59 | 43 | 16 | 18 | 15 | 3 | 41 | 28 | 13 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 16,299 |
| White | 10,499 |
| Black | 5,582 |
| Other | 218 |

Selected Death Statistics, Residents of Clay County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 252 | 128 | 124 | 1,246.0 | 1,573.1 | 1,045.6 |
| 1 | Heart Diseases | 62 | 34 | 28 | 306.6 | 417.8 | 236.1 |
| 2 | Malignant Neoplasms | 58 | 26 | 32 | 286.8 | 319.5 | 269.8 |
| 3 | Cerebrovascular Disease | 15 | 8 | 7 | 74.2 | 98.3 | 59.0 |
| 4 | Nephritis, Nephrotic Syndrome And Nephrosis | 11 | 7 | 4 | 54.4 | 86.0 | 33.7 |
| 5 | Accidents | 10 | 6 | 4 | 49.4 | 73.7 | 33.7 |
| 5 | Alzheimer's Disease | 10 | 5 | 5 | 49.4 | 61.4 | 42.2 |
| 7 | Emphysema and Other Chronic Lower Resp. Diseases | 9 | 8 | 1 | 44.5 | 98.3 | 8.4 |
| 8 | Diabetes Mellitus | 8 | 2 | 6 | 39.6 | 24.6 | 50.6 |
| 9 | Influenza and Pneumonia | 7 | 4 | 3 | 34.6 | 49.2 | 25.3 |
| 10 | Suicide | 5 | 5 | 0 | 24.7 | 61.4 | 0.0 |
| 10 | Septicemia | 5 | 0 | 5 | 24.7 | 0.0 | 42.2 |
| 12 | Conditions Originating in the Perinatal Period** | 3 | 0 | 3 | 14.9 | 0.0 | 22.1 |
| 12 | Pneumonitis Due to Solids and Liquids | 3 | 0 | 3 | 14.8 | 0.0 | 25.3 |
| 14 | Aortic Aneurysm and Dissection | 2 | 0 | 2 | 9.9 | 0.0 | 16.9 |
| 14 | Congenital Malformations and Chrom. Abnormalities | 2 | 0 | 2 | 9.9 | 0.0 | 16.9 |
| 14 | Hypertension/Hypertensive Renal Disease | 2 | 0 | 2 | 9.9 | 0.0 | 16.9 |
| 17 | Homicide | 1 | 0 | 1 | 4.9 | 0.0 | 8.4 |
| 17 | Peptic Ulcer | 1 | 0 | 1 | 4.9 | 0.0 | 8.4 |
| 17 | Parkinson's Disease | 1 | 1 | 0 | 4.9 | 12.3 | 0.0 |
| 17 | HIV Disease | 1 | 0 | 1 | 4.9 | 0.0 | 8.4 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 2 | 1 | 1 | 9.9 | 12.3 | 8.4 |
| | All Other Causes | 34 | 21 | 13 | 168.1 | 258.1 | 109.6 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births.
 Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 252 | 128 | 124 | 131 | 70 | 61 | 121 | 58 | 63 |
| Under 1 | 6 | 0 | 6 | 2 | 0 | 2 | 4 | 0 | 4 |
| (Neonatal) | 3 | 0 | 3 | 1 | 0 | 1 | 2 | 0 | 2 |
| (Postneonatal) | 3 | 0 | 3 | 1 | 0 | 1 | 2 | 0 | 2 |
| 1-4 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 5-9 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 10-14 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 15-24 | 5 | 4 | 1 | 3 | 3 | 0 | 2 | 1 | 1 |
| 25-34 | 6 | 1 | 5 | 5 | 0 | 5 | 1 | 1 | 0 |
| 35-44 | 4 | 3 | 1 | 4 | 3 | 1 | 0 | 0 | 0 |
| 45-54 | 18 | 8 | 10 | 8 | 4 | 4 | 10 | 4 | 6 |
| 55-64 | 37 | 14 | 23 | 21 | 9 | 12 | 16 | 5 | 11 |
| 65-74 | 41 | 23 | 18 | 31 | 17 | 14 | 10 | 6 | 4 |
| 75-84 | 59 | 34 | 25 | 36 | 22 | 14 | 23 | 12 | 11 |
| 85 and over | 73 | 38 | 35 | 18 | 9 | 9 | 55 | 29 | 26 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 20,225 |
| White | 8,137 |
| Black | 11,859 |
| Other | 229 |

Selected Death Statistics, Residents of Coahoma County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 297 | 94 | 201 | 1,197.2 | 1,610.1 | 1,082.0 |
| 1 | Malignant Neoplasms | 72 | 14 | 57 | 290.2 | 239.8 | 306.8 |
| 2 | Heart Diseases | 68 | 24 | 43 | 274.1 | 411.1 | 231.5 |
| 3 | Accidents | 22 | 10 | 12 | 88.7 | 171.3 | 64.6 |
| 4 | Diabetes Mellitus | 20 | 7 | 13 | 80.6 | 119.9 | 70.0 |
| 5 | Cerebrovascular Disease | 17 | 6 | 11 | 68.5 | 102.8 | 59.2 |
| 6 | Hypertension/Hypertensive Renal Disease | 12 | 1 | 11 | 48.4 | 17.1 | 59.2 |
| 7 | Emphysema and Other Chronic Lower Resp. Diseases | 9 | 4 | 5 | 36.3 | 68.5 | 26.9 |
| 8 | Homicide | 8 | 0 | 8 | 32.2 | 0.0 | 43.1 |
| 8 | Alzheimer's Disease | 8 | 7 | 1 | 32.2 | 119.9 | 5.4 |
| 10 | Influenza and Pneumonia | 7 | 4 | 3 | 28.2 | 68.5 | 16.1 |
| 11 | Pneumonitis Due to Solids and Liquids | 5 | 1 | 4 | 20.2 | 17.1 | 21.5 |
| 12 | Nephritis, Nephrotic Syndrome And Nephrosis | 4 | 1 | 3 | 16.1 | 17.1 | 16.1 |
| 13 | Suicide | 3 | 2 | 1 | 12.1 | 34.3 | 5.4 |
| 13 | Conditions Originating in the Perinatal Period** | 3 | 0 | 3 | 7.4 | 0.0 | 8.7 |
| 15 | Congenital Malformations and Chrom. Abnormalities | 2 | 1 | 1 | 8.1 | 17.1 | 5.4 |
| 15 | Septicemia | 2 | 1 | 1 | 8.1 | 17.1 | 5.4 |
| 17 | Chronic Liver Disease And Cirrhosis | 1 | 1 | 0 | 4.0 | 17.1 | 0.0 |
| 17 | In Situ, Benign and Unspecified Neoplasms | 1 | 0 | 1 | 4.0 | 0.0 | 5.4 |
| 17 | Anemias | 1 | 0 | 1 | 4.0 | 0.0 | 5.4 |
| 17 | HIV Disease | 1 | 0 | 1 | 4.0 | 0.0 | 5.4 |
| 17 | Viral Hepatitis | 1 | 1 | 0 | 4.0 | 17.1 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 1 | 0 | 1 | 4.0 | 0.0 | 5.4 |
| | All Other Causes | 29 | 9 | 20 | 116.9 | 154.2 | 107.7 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total* ----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|--------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 297 | 94 | 201 | 168 | 63 | 104 | 129 | 31 | 97 |
| Under 1 | 4 | 0 | 4 | 2 | 0 | 2 | 2 | 0 | 2 |
| (Neonatal) | 4 | 0 | 4 | 2 | 0 | 2 | 2 | 0 | 2 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 6 | 1 | 5 | 5 | 1 | 4 | 1 | 0 | 1 |
| 25-34 | 14 | 2 | 12 | 7 | 1 | 6 | 7 | 1 | 6 |
| 35-44 | 14 | 4 | 10 | 10 | 3 | 7 | 4 | 1 | 3 |
| 45-54 | 31 | 10 | 21 | 21 | 7 | 14 | 10 | 3 | 7 |
| 55-64 | 55 | 17 | 37 | 37 | 12 | 24 | 18 | 5 | 13 |
| 65-74 | 51 | 17 | 34 | 30 | 13 | 17 | 21 | 4 | 17 |
| 75-84 | 68 | 24 | 43 | 40 | 18 | 22 | 28 | 6 | 21 |
| 85 and over | 53 | 19 | 34 | 15 | 8 | 7 | 38 | 11 | 27 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 24,807 |
| White | 5,838 |
| Black | 18,576 |
| Other | 393 |

Selected Death Statistics, Residents of Copiah County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 323 | 196 | 127 | 1,121.6 | 1,445.3 | 858.5 |
| 1 | Heart Diseases | 86 | 49 | 37 | 298.6 | 361.3 | 250.1 |
| 2 | Malignant Neoplasms | 63 | 36 | 27 | 218.8 | 265.5 | 182.5 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 17 | 12 | 5 | 59.0 | 88.5 | 33.8 |
| 4 | Cerebrovascular Disease | 15 | 9 | 6 | 52.1 | 66.4 | 40.6 |
| 5 | Diabetes Mellitus | 13 | 7 | 6 | 45.1 | 51.6 | 40.6 |
| 6 | Accidents | 10 | 9 | 1 | 34.7 | 66.4 | 6.8 |
| 7 | Influenza and Pneumonia | 9 | 7 | 2 | 31.3 | 51.6 | 13.5 |
| 7 | Alzheimer's Disease | 9 | 8 | 1 | 31.3 | 59.0 | 6.8 |
| 9 | Septicemia | 6 | 4 | 2 | 20.8 | 29.5 | 13.5 |
| 10 | Nephritis, Nephrotic Syndrome And Nephrosis | 5 | 1 | 4 | 17.4 | 7.4 | 27.0 |
| 11 | Suicide | 4 | 3 | 1 | 13.9 | 22.1 | 6.8 |
| 11 | Chronic Liver Disease And Cirrhosis | 4 | 4 | 0 | 13.9 | 29.5 | 0.0 |
| 13 | Homicide | 3 | 2 | 1 | 10.4 | 14.7 | 6.8 |
| 14 | HIV Disease | 2 | 0 | 2 | 6.9 | 0.0 | 13.5 |
| 14 | Viral Hepatitis | 2 | 2 | 0 | 6.9 | 14.7 | 0.0 |
| 14 | Conditions Originating in the Perinatal Period** | 2 | 1 | 1 | 5.7 | 6.7 | 5.2 |
| 14 | Pneumonitis Due to Solids and Liquids | 2 | 2 | 0 | 6.9 | 14.7 | 0.0 |
| 14 | Parkinson's Disease | 2 | 2 | 0 | 6.9 | 14.7 | 0.0 |
| 19 | Complications Medical/Surgical Care | 1 | 0 | 1 | 3.5 | 0.0 | 6.8 |
| 19 | Peptic Ulcer | 1 | 0 | 1 | 3.5 | 0.0 | 6.8 |
| 19 | Aortic Aneurysm and Dissection | 1 | 1 | 0 | 3.5 | 7.4 | 0.0 |
| 19 | Hypertension/Hypertensive Renal Disease | 1 | 1 | 0 | 3.5 | 7.4 | 0.0 |
| 19 | Malnutrition/Nutritional Deficiency | 1 | 0 | 1 | 3.5 | 0.0 | 6.8 |
| 19 | Anemias | 1 | 1 | 0 | 3.5 | 7.4 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 2 | 2 | 0 | 6.9 | 14.7 | 0.0 |
| | All Other Causes | 61 | 33 | 28 | 211.8 | 243.3 | 189.3 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births.
 Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total* ----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|--------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 323 | 196 | 127 | 177 | 114 | 63 | 146 | 82 | 64 |
| Under 1 | 4 | 3 | 1 | 4 | 3 | 1 | 0 | 0 | 0 |
| (Neonatal) | 2 | 1 | 1 | 2 | 1 | 1 | 0 | 0 | 0 |
| (Postneonatal) | 2 | 2 | 0 | 2 | 2 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 3 | 3 | 0 | 3 | 3 | 0 | 0 | 0 | 0 |
| 25-34 | 4 | 1 | 3 | 2 | 1 | 1 | 2 | 0 | 2 |
| 35-44 | 15 | 7 | 8 | 8 | 5 | 3 | 7 | 2 | 5 |
| 45-54 | 35 | 18 | 17 | 24 | 15 | 9 | 11 | 3 | 8 |
| 55-64 | 56 | 27 | 29 | 38 | 18 | 20 | 18 | 9 | 9 |
| 65-74 | 59 | 35 | 24 | 29 | 19 | 10 | 30 | 16 | 14 |
| 75-84 | 68 | 44 | 24 | 40 | 27 | 13 | 28 | 17 | 11 |
| 85 and over | 79 | 58 | 21 | 29 | 23 | 6 | 50 | 35 | 15 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 28,797 |
| White | 13,561 |
| Black | 14,793 |
| Other | 443 |

Selected Death Statistics, Residents of Covington County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 241 | 178 | 62 | 1,239.6 | 1,462.1 | 889.3 |
| 1 | Heart Diseases | 58 | 47 | 11 | 298.3 | 386.1 | 157.8 |
| 2 | Malignant Neoplasms | 41 | 28 | 13 | 210.9 | 230.0 | 186.5 |
| 3 | Alzheimer's Disease | 20 | 16 | 4 | 102.9 | 131.4 | 57.4 |
| 3 | Cerebrovascular Disease | 20 | 13 | 7 | 102.9 | 106.8 | 100.4 |
| 5 | Emphysema and Other Chronic Lower Resp. Diseases | 18 | 14 | 3 | 92.6 | 115.0 | 43.0 |
| 6 | Accidents | 14 | 10 | 4 | 72.0 | 82.1 | 57.4 |
| 7 | Influenza and Pneumonia | 7 | 4 | 3 | 36.0 | 32.9 | 43.0 |
| 8 | Septicemia | 5 | 4 | 1 | 25.7 | 32.9 | 14.3 |
| 9 | Suicide | 4 | 4 | 0 | 20.6 | 32.9 | 0.0 |
| 9 | Pneumonitis Due to Solids and Liquids | 4 | 2 | 2 | 20.6 | 16.4 | 28.7 |
| 9 | Parkinson's Disease | 4 | 4 | 0 | 20.6 | 32.9 | 0.0 |
| 9 | Diabetes Mellitus | 4 | 1 | 3 | 20.6 | 8.2 | 43.0 |
| 13 | Homicide | 3 | 1 | 2 | 15.4 | 8.2 | 28.7 |
| 13 | Nephritis, Nephrotic Syndrome And Nephrosis | 3 | 2 | 1 | 15.4 | 16.4 | 14.3 |
| 13 | Aortic Aneurysm and Dissection | 3 | 2 | 1 | 15.4 | 16.4 | 14.3 |
| 16 | Cholelithiasis/Gallbladder Disorder | 1 | 1 | 0 | 5.1 | 8.2 | 0.0 |
| 16 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 3.8 | 0.0 | 9.6 |
| 16 | Anemias | 1 | 1 | 0 | 5.1 | 8.2 | 0.0 |
| 16 | In Situ, Benign and Unspecified Neoplasms | 1 | 0 | 1 | 5.1 | 0.0 | 14.3 |
| 16 | HIV Disease | 1 | 0 | 1 | 5.1 | 0.0 | 14.3 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 3 | 3 | 0 | 15.4 | 24.6 | 0.0 |
| | All Other Causes | 25 | 21 | 4 | 128.6 | 172.5 | 57.4 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births.
 Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 241 | 178 | 62 | 143 | 112 | 30 | 98 | 66 | 32 |
| Under 1 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| (Neonatal) | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| 15-24 | 3 | 2 | 1 | 2 | 2 | 0 | 1 | 0 | 1 |
| 25-34 | 5 | 1 | 4 | 4 | 1 | 3 | 1 | 0 | 1 |
| 35-44 | 6 | 3 | 3 | 5 | 3 | 2 | 1 | 0 | 1 |
| 45-54 | 12 | 9 | 3 | 6 | 4 | 2 | 6 | 5 | 1 |
| 55-64 | 26 | 11 | 15 | 14 | 8 | 6 | 12 | 3 | 9 |
| 65-74 | 50 | 38 | 11 | 29 | 22 | 6 | 21 | 16 | 5 |
| 75-84 | 56 | 43 | 13 | 37 | 32 | 5 | 19 | 11 | 8 |
| 85 and over | 81 | 71 | 10 | 45 | 40 | 5 | 36 | 31 | 5 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 19,442 |
| White | 12,174 |
| Black | 6,972 |
| Other | 296 |

Selected Death Statistics, Residents of Desoto County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|-------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 1,259 | 1,085 | 166 | 736.6 | 883.2 | 389.9 |
| 1 | Malignant Neoplasms | 343 | 296 | 44 | 200.7 | 241.0 | 103.4 |
| 2 | Heart Diseases | 293 | 266 | 27 | 171.4 | 216.5 | 63.4 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 86 | 82 | 4 | 50.3 | 66.8 | 9.4 |
| 4 | Diabetes Mellitus | 80 | 57 | 20 | 46.8 | 46.4 | 47.0 |
| 5 | Accidents | 66 | 62 | 4 | 38.6 | 50.5 | 9.4 |
| 6 | Alzheimer's Disease | 52 | 48 | 3 | 30.4 | 39.1 | 7.0 |
| 7 | Cerebrovascular Disease | 45 | 34 | 11 | 26.3 | 27.7 | 25.8 |
| 8 | Hypertension/Hypertensive Renal Disease | 25 | 23 | 2 | 14.6 | 18.7 | 4.7 |
| 9 | Suicide | 23 | 19 | 4 | 13.5 | 15.5 | 9.4 |
| 10 | Influenza and Pneumonia | 20 | 19 | 1 | 11.7 | 15.5 | 2.3 |
| 11 | Nephritis, Nephrotic Syndrome And Nephrosis | 18 | 10 | 8 | 10.5 | 8.1 | 18.8 |
| 11 | Chronic Liver Disease And Cirrhosis | 18 | 18 | 0 | 10.5 | 14.7 | 0.0 |
| 13 | Homicide | 11 | 8 | 3 | 6.4 | 6.5 | 7.0 |
| 13 | Septicemia | 11 | 9 | 2 | 6.4 | 7.3 | 4.7 |
| 15 | In Situ, Benign and Unspecified Neoplasms | 8 | 5 | 3 | 4.7 | 4.1 | 7.0 |
| 16 | Parkinson's Disease | 6 | 6 | 0 | 3.5 | 4.9 | 0.0 |
| 17 | Conditions Originating in the Perinatal Period** | 4 | 2 | 2 | 1.9 | 1.4 | 3.4 |
| 17 | Congenital Malformations and Chrom. Abnormalities | 4 | 1 | 3 | 2.3 | 0.8 | 7.0 |
| 17 | Pneumonitis Due to Solids and Liquids | 4 | 4 | 0 | 2.3 | 3.3 | 0.0 |
| 20 | Complications Medical/Surgical Care | 3 | 2 | 1 | 1.8 | 1.6 | 2.3 |
| 20 | Cholelithiasis/Gallbladder Disorder | 3 | 3 | 0 | 1.8 | 2.4 | 0.0 |
| 20 | Malnutrition/Nutritional Deficiency | 3 | 1 | 2 | 1.8 | 0.8 | 4.7 |
| 23 | Aortic Aneurysm and Dissection | 2 | 1 | 1 | 1.2 | 0.8 | 2.3 |
| 23 | Anemias | 2 | 2 | 0 | 1.2 | 1.6 | 0.0 |
| 23 | Viral Hepatitis | 2 | 2 | 0 | 1.2 | 1.6 | 0.0 |
| 26 | Legal Intervention | 1 | 1 | 0 | 0.6 | 0.8 | 0.0 |
| 26 | Maternal Causes** | 1 | 1 | 0 | 47.7 | 69.4 | 0.0 |
| 26 | Atherosclerosis | 1 | 1 | 0 | 0.6 | 0.8 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 10 | 10 | 0 | 5.9 | 8.1 | 0.0 |
| | All Other Causes | 114 | 92 | 21 | 66.7 | 74.9 | 49.3 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 1,259 | 1,085 | 166 | 668 | 583 | 81 | 591 | 502 | 85 |
| Under 1 | 6 | 3 | 3 | 5 | 2 | 3 | 1 | 1 | 0 |
| (Neonatal) | 6 | 3 | 3 | 5 | 2 | 3 | 1 | 1 | 0 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 4 | 4 | 0 | 4 | 4 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 2 | 2 | 0 | 1 | 1 | 0 | 1 | 1 | 0 |
| 15-24 | 17 | 13 | 4 | 14 | 11 | 3 | 3 | 2 | 1 |
| 25-34 | 23 | 20 | 3 | 19 | 18 | 1 | 4 | 2 | 2 |
| 35-44 | 42 | 31 | 11 | 22 | 18 | 4 | 20 | 13 | 7 |
| 45-54 | 104 | 86 | 18 | 62 | 55 | 7 | 42 | 31 | 11 |
| 55-64 | 204 | 167 | 37 | 126 | 105 | 21 | 78 | 62 | 16 |
| 65-74 | 271 | 231 | 37 | 149 | 124 | 24 | 122 | 107 | 13 |
| 75-84 | 327 | 293 | 29 | 171 | 156 | 12 | 156 | 137 | 17 |
| 85 and over | 259 | 235 | 24 | 95 | 89 | 6 | 164 | 146 | 18 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|---------|
| Total | 170,913 |
| White | 122,842 |
| Black | 42,571 |
| Other | 5,500 |

Selected Death Statistics, Residents of Forrest County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 722 | 482 | 238 | 945.9 | 1,052.2 | 842.1 |
| 1 | Heart Diseases | 147 | 97 | 49 | 192.6 | 211.7 | 173.4 |
| 2 | Malignant Neoplasms | 132 | 86 | 45 | 172.9 | 187.7 | 159.2 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 48 | 37 | 11 | 62.9 | 80.8 | 38.9 |
| 4 | Accidents | 38 | 28 | 10 | 49.8 | 61.1 | 35.4 |
| 5 | Cerebrovascular Disease | 34 | 20 | 14 | 44.5 | 43.7 | 49.5 |
| 5 | Diabetes Mellitus | 34 | 19 | 15 | 44.5 | 41.5 | 53.1 |
| 7 | Alzheimer's Disease | 33 | 25 | 8 | 43.2 | 54.6 | 28.3 |
| 8 | Septicemia | 24 | 13 | 11 | 31.4 | 28.4 | 38.9 |
| 9 | Nephritis, Nephrotic Syndrome And Nephrosis | 22 | 11 | 11 | 28.8 | 24.0 | 38.9 |
| 10 | Influenza and Pneumonia | 18 | 12 | 6 | 23.6 | 26.2 | 21.2 |
| 11 | Chronic Liver Disease And Cirrhosis | 13 | 11 | 2 | 17.0 | 24.0 | 7.1 |
| 12 | Suicide | 12 | 11 | 1 | 15.7 | 24.0 | 3.5 |
| 12 | Homicide | 12 | 3 | 9 | 15.7 | 6.5 | 31.8 |
| 14 | Hypertension/Hypertensive Renal Disease | 9 | 6 | 3 | 11.8 | 13.1 | 10.6 |
| 15 | Pneumonitis Due to Solids and Liquids | 8 | 5 | 3 | 10.5 | 10.9 | 10.6 |
| 16 | Conditions Originating in the Perinatal Period** | 4 | 2 | 2 | 3.9 | 3.6 | 4.4 |
| 16 | Parkinson's Disease | 4 | 4 | 0 | 5.2 | 8.7 | 0.0 |
| 16 | In Situ, Benign and Unspecified Neoplasms | 4 | 4 | 0 | 5.2 | 8.7 | 0.0 |
| 19 | Cholelithiasis/Gallbladder Disorder | 3 | 2 | 1 | 3.9 | 4.4 | 3.5 |
| 19 | Congenital Malformations and Chrom. Abnormalities | 3 | 2 | 1 | 3.9 | 4.4 | 3.5 |
| 19 | Anemias | 3 | 1 | 2 | 3.9 | 2.2 | 7.1 |
| 19 | Viral Hepatitis | 3 | 2 | 1 | 3.9 | 4.4 | 3.5 |
| 23 | Complications Medical/Surgical Care | 1 | 0 | 1 | 1.3 | 0.0 | 3.5 |
| 23 | Atherosclerosis | 1 | 1 | 0 | 1.3 | 2.2 | 0.0 |
| 23 | Aortic Aneurysm and Dissection | 1 | 0 | 1 | 1.3 | 0.0 | 3.5 |
| 23 | HIV Disease | 1 | 0 | 1 | 1.3 | 0.0 | 3.5 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 8 | 6 | 2 | 10.5 | 13.1 | 7.1 |
| | All Other Causes | 102 | 74 | 28 | 133.6 | 161.5 | 99.1 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 722 | 482 | 238 | 370 | 247 | 123 | 352 | 235 | 115 |
| Under 1 | 10 | 5 | 5 | 5 | 3 | 2 | 5 | 2 | 3 |
| (Neonatal) | 4 | 2 | 2 | 2 | 1 | 1 | 2 | 1 | 1 |
| (Postneonatal) | 6 | 3 | 3 | 3 | 2 | 1 | 3 | 1 | 2 |
| 1-4 | 2 | 0 | 2 | 1 | 0 | 1 | 1 | 0 | 1 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 2 | 1 | 1 | 2 | 1 | 1 | 0 | 0 | 0 |
| 15-24 | 7 | 2 | 5 | 4 | 1 | 3 | 3 | 1 | 2 |
| 25-34 | 14 | 7 | 7 | 8 | 5 | 3 | 6 | 2 | 4 |
| 35-44 | 26 | 9 | 17 | 14 | 5 | 9 | 12 | 4 | 8 |
| 45-54 | 74 | 44 | 30 | 44 | 27 | 17 | 30 | 17 | 13 |
| 55-64 | 122 | 66 | 56 | 82 | 42 | 40 | 40 | 24 | 16 |
| 65-74 | 144 | 101 | 43 | 70 | 52 | 18 | 74 | 49 | 25 |
| 75-84 | 157 | 119 | 36 | 89 | 70 | 19 | 68 | 49 | 17 |
| 85 and over | 164 | 128 | 36 | 51 | 41 | 10 | 113 | 87 | 26 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 76,330 |
| White | 45,809 |
| Black | 28,264 |
| Other | 2,257 |

Selected Death Statistics, Residents of Franklin County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 117 | 82 | 35 | 1,493.7 | 1,655.2 | 1,259.0 |
| 1 | Heart Diseases | 40 | 28 | 12 | 510.7 | 565.2 | 431.7 |
| 2 | Malignant Neoplasms | 26 | 19 | 7 | 331.9 | 383.5 | 251.8 |
| 3 | Accidents | 9 | 4 | 5 | 114.9 | 80.7 | 179.9 |
| 4 | Influenza and Pneumonia | 5 | 5 | 0 | 63.8 | 100.9 | 0.0 |
| 4 | Cerebrovascular Disease | 5 | 1 | 4 | 63.8 | 20.2 | 143.9 |
| 4 | Alzheimer's Disease | 5 | 5 | 0 | 63.8 | 100.9 | 0.0 |
| 7 | Suicide | 3 | 2 | 1 | 38.3 | 40.4 | 36.0 |
| 7 | Emphysema and Other Chronic Lower Resp. Diseases | 3 | 2 | 1 | 38.3 | 40.4 | 36.0 |
| 9 | Nephritis, Nephrotic Syndrome And Nephrosis | 2 | 1 | 1 | 25.5 | 20.2 | 36.0 |
| 9 | Diabetes Mellitus | 2 | 1 | 1 | 25.5 | 20.2 | 36.0 |
| 11 | Chronic Liver Disease And Cirrhosis | 1 | 1 | 0 | 12.8 | 20.2 | 0.0 |
| 11 | Parkinson's Disease | 1 | 1 | 0 | 12.8 | 20.2 | 0.0 |
| 11 | Septicemia | 1 | 1 | 0 | 12.8 | 20.2 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 2 | 2 | 0 | 25.5 | 40.4 | 0.0 |
| | All Other Causes | 12 | 9 | 3 | 153.2 | 181.7 | 107.9 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births.
 Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 117 | 82 | 35 | 53 | 39 | 14 | 64 | 43 | 21 |
| Under 1 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| (Neonatal) | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 25-34 | 4 | 1 | 3 | 1 | 1 | 0 | 3 | 0 | 3 |
| 35-44 | 4 | 2 | 2 | 3 | 1 | 2 | 1 | 1 | 0 |
| 45-54 | 3 | 2 | 1 | 1 | 0 | 1 | 2 | 2 | 0 |
| 55-64 | 12 | 10 | 2 | 7 | 7 | 0 | 5 | 3 | 2 |
| 65-74 | 22 | 14 | 8 | 14 | 7 | 7 | 8 | 7 | 1 |
| 75-84 | 31 | 23 | 8 | 13 | 11 | 2 | 18 | 12 | 6 |
| 85 and over | 39 | 30 | 9 | 13 | 12 | 1 | 26 | 18 | 8 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|-------|
| Total | 7,833 |
| White | 4,954 |
| Black | 2,780 |
| Other | 99 |

Selected Death Statistics, Residents of George County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 245 | 225 | 18 | 1,051.4 | 1,075.0 | 906.3 |
| 1 | Malignant Neoplasms | 51 | 49 | 2 | 218.9 | 234.1 | 100.7 |
| 2 | Heart Diseases | 49 | 42 | 6 | 210.3 | 200.7 | 302.1 |
| 3 | Accidents | 22 | 20 | 2 | 94.4 | 95.6 | 100.7 |
| 4 | Emphysema and Other Chronic Lower Resp. Diseases | 17 | 15 | 1 | 73.0 | 71.7 | 50.4 |
| 5 | Diabetes Mellitus | 15 | 14 | 1 | 64.4 | 66.9 | 50.4 |
| 6 | Alzheimer's Disease | 10 | 10 | 0 | 42.9 | 47.8 | 0.0 |
| 6 | Septicemia | 10 | 9 | 1 | 42.9 | 43.0 | 50.4 |
| 8 | Cerebrovascular Disease | 7 | 7 | 0 | 30.0 | 33.4 | 0.0 |
| 9 | Homicide | 4 | 4 | 0 | 17.2 | 19.1 | 0.0 |
| 10 | Influenza and Pneumonia | 3 | 3 | 0 | 12.9 | 14.3 | 0.0 |
| 10 | Nephritis, Nephrotic Syndrome And Nephrosis | 3 | 2 | 1 | 12.9 | 9.6 | 50.4 |
| 10 | Aortic Aneurysm and Dissection | 3 | 3 | 0 | 12.9 | 14.3 | 0.0 |
| 13 | Suicide | 2 | 2 | 0 | 8.6 | 9.6 | 0.0 |
| 13 | Pneumonitis Due to Solids and Liquids | 2 | 2 | 0 | 8.6 | 9.6 | 0.0 |
| 15 | Congenital Malformations and Chrom. Abnormalities | 1 | 1 | 0 | 4.3 | 4.8 | 0.0 |
| 15 | Complications Medical/Surgical Care | 1 | 1 | 0 | 4.3 | 4.8 | 0.0 |
| 15 | Cholelithiasis/Gallbladder Disorder | 1 | 1 | 0 | 4.3 | 4.8 | 0.0 |
| 15 | Pneumoconioses and Chemical Effects | 1 | 1 | 0 | 4.3 | 4.8 | 0.0 |
| 15 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 2.8 | 0.0 | 38.5 |
| 15 | Hernia | 1 | 1 | 0 | 4.3 | 4.8 | 0.0 |
| 15 | Parkinson's Disease | 1 | 1 | 0 | 4.3 | 4.8 | 0.0 |
| 15 | Anemias | 1 | 0 | 1 | 4.3 | 0.0 | 50.4 |
| 15 | In Situ, Benign and Unspecified Neoplasms | 1 | 1 | 0 | 4.3 | 4.8 | 0.0 |
| 15 | Hypertension/Hypertensive Renal Disease | 1 | 0 | 1 | 4.3 | 0.0 | 50.4 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 2 | 2 | 0 | 8.6 | 9.6 | 0.0 |
| | All Other Causes | 35 | 34 | 1 | 150.2 | 162.4 | 50.4 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total* ----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|--------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 245 | 225 | 18 | 117 | 108 | 8 | 128 | 117 | 10 |
| Under 1 | 2 | 0 | 2 | 1 | 0 | 1 | 1 | 0 | 1 |
| (Neonatal) | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| (Postneonatal) | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 6 | 5 | 1 | 4 | 3 | 1 | 2 | 2 | 0 |
| 25-34 | 8 | 7 | 1 | 4 | 4 | 0 | 4 | 3 | 1 |
| 35-44 | 8 | 7 | 1 | 3 | 3 | 0 | 5 | 4 | 1 |
| 45-54 | 21 | 18 | 3 | 15 | 14 | 1 | 6 | 4 | 2 |
| 55-64 | 36 | 34 | 1 | 21 | 20 | 1 | 15 | 14 | 0 |
| 65-74 | 52 | 48 | 3 | 26 | 23 | 2 | 26 | 25 | 1 |
| 75-84 | 56 | 55 | 1 | 23 | 23 | 0 | 33 | 32 | 1 |
| 85 and over | 56 | 51 | 5 | 20 | 18 | 2 | 36 | 33 | 3 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 23,303 |
| White | 20,931 |
| Black | 1,986 |
| Other | 386 |

Selected Death Statistics, Residents of Greene County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 126 | 106 | 20 | 879.5 | 1,035.6 | 513.1 |
| 1 | Heart Diseases | 38 | 31 | 7 | 265.3 | 302.9 | 179.6 |
| 2 | Malignant Neoplasms | 24 | 20 | 4 | 167.5 | 195.4 | 102.6 |
| 3 | Accidents | 9 | 8 | 1 | 62.8 | 78.2 | 25.7 |
| 3 | Cerebrovascular Disease | 9 | 7 | 2 | 62.8 | 68.4 | 51.3 |
| 5 | Alzheimer's Disease | 5 | 5 | 0 | 34.9 | 48.8 | 0.0 |
| 6 | Homicide | 4 | 2 | 2 | 27.9 | 19.5 | 51.3 |
| 6 | Diabetes Mellitus | 4 | 4 | 0 | 27.9 | 39.1 | 0.0 |
| 8 | Influenza and Pneumonia | 3 | 2 | 1 | 20.9 | 19.5 | 25.7 |
| 8 | Septicemia | 3 | 3 | 0 | 20.9 | 29.3 | 0.0 |
| 10 | Emphysema and Other Chronic Lower Resp. Diseases | 2 | 1 | 1 | 14.0 | 9.8 | 25.7 |
| 10 | Aortic Aneurysm and Dissection | 2 | 2 | 0 | 14.0 | 19.5 | 0.0 |
| 10 | HIV Disease | 2 | 1 | 1 | 14.0 | 9.8 | 25.7 |
| 13 | Suicide | 1 | 1 | 0 | 7.0 | 9.8 | 0.0 |
| 13 | Hernia | 1 | 1 | 0 | 7.0 | 9.8 | 0.0 |
| 13 | Pneumonitis Due to Solids and Liquids | 1 | 1 | 0 | 7.0 | 9.8 | 0.0 |
| 13 | Congenital Malformations and Chrom. Abnormalities | 1 | 1 | 0 | 7.0 | 9.8 | 0.0 |
| 13 | Chronic Liver Disease And Cirrhosis | 1 | 1 | 0 | 7.0 | 9.8 | 0.0 |
| 13 | Nephritis, Nephrotic Syndrome And Nephrosis | 1 | 1 | 0 | 7.0 | 9.8 | 0.0 |
| 13 | Parkinson's Disease | 1 | 1 | 0 | 7.0 | 9.8 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 2 | 2 | 0 | 14.0 | 19.5 | 0.0 |
| | All Other Causes | 12 | 11 | 1 | 83.8 | 107.5 | 25.7 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 126 | 106 | 20 | 71 | 58 | 13 | 55 | 48 | 7 |
| Under 1 | 2 | 2 | 0 | 2 | 2 | 0 | 0 | 0 | 0 |
| (Neonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 2 | 2 | 0 | 2 | 2 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 15-24 | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 |
| 25-34 | 3 | 3 | 0 | 3 | 3 | 0 | 0 | 0 | 0 |
| 35-44 | 6 | 4 | 2 | 3 | 1 | 2 | 3 | 3 | 0 |
| 45-54 | 12 | 6 | 6 | 8 | 4 | 4 | 4 | 2 | 2 |
| 55-64 | 14 | 13 | 1 | 8 | 8 | 0 | 6 | 5 | 1 |
| 65-74 | 28 | 27 | 1 | 22 | 21 | 1 | 6 | 6 | 0 |
| 75-84 | 35 | 30 | 5 | 17 | 14 | 3 | 18 | 16 | 2 |
| 85 and over | 24 | 19 | 5 | 7 | 4 | 3 | 17 | 15 | 2 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 14,326 |
| White | 10,236 |
| Black | 3,898 |
| Other | 192 |

Selected Death Statistics, Residents of Grenada County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 288 | 177 | 111 | 1,329.3 | 1,458.7 | 1,207.4 |
| 1 | Heart Diseases | 82 | 44 | 38 | 378.5 | 362.6 | 413.4 |
| 2 | Malignant Neoplasms | 52 | 28 | 24 | 240.0 | 230.8 | 261.1 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 19 | 16 | 3 | 87.7 | 131.9 | 32.6 |
| 4 | Cerebrovascular Disease | 16 | 9 | 7 | 73.8 | 74.2 | 76.1 |
| 5 | Alzheimer's Disease | 14 | 14 | 0 | 64.6 | 115.4 | 0.0 |
| 6 | Influenza and Pneumonia | 11 | 6 | 5 | 50.8 | 49.4 | 54.4 |
| 7 | Accidents | 10 | 8 | 2 | 46.2 | 65.9 | 21.8 |
| 8 | Nephritis, Nephrotic Syndrome And Nephrosis | 9 | 8 | 1 | 41.5 | 65.9 | 10.9 |
| 9 | Homicide | 7 | 2 | 5 | 32.3 | 16.5 | 54.4 |
| 10 | Hypertension/Hypertensive Renal Disease | 5 | 4 | 1 | 23.1 | 33.0 | 10.9 |
| 11 | Pneumonitis Due to Solids and Liquids | 4 | 2 | 2 | 18.5 | 16.5 | 21.8 |
| 11 | Chronic Liver Disease And Cirrhosis | 4 | 4 | 0 | 18.5 | 33.0 | 0.0 |
| 11 | Septicemia | 4 | 3 | 1 | 18.5 | 24.7 | 10.9 |
| 14 | Suicide | 3 | 3 | 0 | 13.8 | 24.7 | 0.0 |
| 14 | Conditions Originating in the Perinatal Period** | 3 | 0 | 3 | 11.0 | 0.0 | 21.3 |
| 16 | Aortic Aneurysm and Dissection | 2 | 1 | 1 | 9.2 | 8.2 | 10.9 |
| 16 | Diabetes Mellitus | 2 | 1 | 1 | 9.2 | 8.2 | 10.9 |
| 18 | Complications Medical/Surgical Care | 1 | 1 | 0 | 4.6 | 8.2 | 0.0 |
| 18 | Congenital Malformations and Chrom. Abnormalities | 1 | 1 | 0 | 4.6 | 8.2 | 0.0 |
| 18 | Parkinson's Disease | 1 | 1 | 0 | 4.6 | 8.2 | 0.0 |
| 18 | Malnutrition/Nutritional Deficiency | 1 | 1 | 0 | 4.6 | 8.2 | 0.0 |
| 18 | In Situ, Benign and Unspecified Neoplasms | 1 | 1 | 0 | 4.6 | 8.2 | 0.0 |
| 18 | HIV Disease | 1 | 0 | 1 | 4.6 | 0.0 | 10.9 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 0 | 0 | 0 | 0.0 | 0.0 | 0.0 |
| | All Other Causes | 35 | 19 | 16 | 161.5 | 156.6 | 174.0 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 288 | 177 | 111 | 148 | 93 | 55 | 140 | 84 | 56 |
| Under 1 | 4 | 1 | 3 | 1 | 0 | 1 | 3 | 1 | 2 |
| (Neonatal) | 3 | 0 | 3 | 1 | 0 | 1 | 2 | 0 | 2 |
| (Postneonatal) | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 |
| 1-4 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 6 | 3 | 3 | 6 | 3 | 3 | 0 | 0 | 0 |
| 25-34 | 10 | 3 | 7 | 4 | 2 | 2 | 6 | 1 | 5 |
| 35-44 | 9 | 2 | 7 | 5 | 2 | 3 | 4 | 0 | 4 |
| 45-54 | 25 | 10 | 15 | 15 | 7 | 8 | 10 | 3 | 7 |
| 55-64 | 44 | 17 | 27 | 30 | 14 | 16 | 14 | 3 | 11 |
| 65-74 | 55 | 39 | 16 | 37 | 27 | 10 | 18 | 12 | 6 |
| 75-84 | 66 | 52 | 14 | 28 | 22 | 6 | 38 | 30 | 8 |
| 85 and over | 68 | 50 | 18 | 21 | 16 | 5 | 47 | 34 | 13 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 21,666 |
| White | 12,134 |
| Black | 9,193 |
| Other | 339 |

Selected Death Statistics, Residents of Hancock County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 488 | 463 | 25 | 1,062.0 | 1,148.0 | 648.5 |
| 1 | Malignant Neoplasms | 109 | 106 | 3 | 237.2 | 262.8 | 77.8 |
| 2 | Heart Diseases | 92 | 85 | 7 | 200.2 | 210.8 | 181.6 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 46 | 46 | 0 | 100.1 | 114.1 | 0.0 |
| 4 | Accidents | 33 | 28 | 5 | 71.8 | 69.4 | 129.7 |
| 5 | Alzheimer's Disease | 23 | 22 | 1 | 50.1 | 54.5 | 25.9 |
| 6 | Cerebrovascular Disease | 17 | 15 | 2 | 37.0 | 37.2 | 51.9 |
| 7 | Nephritis, Nephrotic Syndrome And Nephrosis | 13 | 12 | 1 | 28.3 | 29.8 | 25.9 |
| 8 | Influenza and Pneumonia | 10 | 10 | 0 | 21.8 | 24.8 | 0.0 |
| 8 | Diabetes Mellitus | 10 | 10 | 0 | 21.8 | 24.8 | 0.0 |
| 10 | Suicide | 9 | 9 | 0 | 19.6 | 22.3 | 0.0 |
| 11 | Septicemia | 8 | 6 | 2 | 17.4 | 14.9 | 51.9 |
| 12 | Chronic Liver Disease And Cirrhosis | 7 | 7 | 0 | 15.2 | 17.4 | 0.0 |
| 12 | Parkinson's Disease | 7 | 7 | 0 | 15.2 | 17.4 | 0.0 |
| 14 | Complications Medical/Surgical Care | 6 | 6 | 0 | 13.1 | 14.9 | 0.0 |
| 15 | Hypertension/Hypertensive Renal Disease | 4 | 4 | 0 | 8.7 | 9.9 | 0.0 |
| 15 | Malnutrition/Nutritional Deficiency | 4 | 4 | 0 | 8.7 | 9.9 | 0.0 |
| 17 | Congenital Malformations and Chrom. Abnormalities | 3 | 3 | 0 | 6.5 | 7.4 | 0.0 |
| 18 | Conditions Originating in the Perinatal Period** | 2 | 2 | 0 | 3.8 | 4.3 | 0.0 |
| 19 | Maternal Causes** | 1 | 1 | 0 | 188.3 | 213.7 | 0.0 |
| 19 | Hernia | 1 | 1 | 0 | 2.2 | 2.5 | 0.0 |
| 19 | Pneumoconioses and Chemical Effects | 1 | 1 | 0 | 2.2 | 2.5 | 0.0 |
| 19 | Anemias | 1 | 1 | 0 | 2.2 | 2.5 | 0.0 |
| 19 | In Situ, Benign and Unspecified Neoplasms | 1 | 1 | 0 | 2.2 | 2.5 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 4 | 4 | 0 | 8.7 | 9.9 | 0.0 |
| | All Other Causes | 76 | 72 | 4 | 165.4 | 178.5 | 103.8 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 488 | 463 | 25 | 262 | 251 | 11 | 226 | 212 | 14 |
| Under 1 | 3 | 3 | 0 | 2 | 2 | 0 | 1 | 1 | 0 |
| (Neonatal) | 2 | 2 | 0 | 1 | 1 | 0 | 1 | 1 | 0 |
| (Postneonatal) | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 15-24 | 2 | 2 | 0 | 1 | 1 | 0 | 1 | 1 | 0 |
| 25-34 | 5 | 5 | 0 | 3 | 3 | 0 | 2 | 2 | 0 |
| 35-44 | 11 | 9 | 2 | 6 | 6 | 0 | 5 | 3 | 2 |
| 45-54 | 37 | 36 | 1 | 23 | 22 | 1 | 14 | 14 | 0 |
| 55-64 | 93 | 90 | 3 | 54 | 52 | 2 | 39 | 38 | 1 |
| 65-74 | 116 | 110 | 6 | 68 | 65 | 3 | 48 | 45 | 3 |
| 75-84 | 123 | 115 | 8 | 58 | 55 | 3 | 65 | 60 | 5 |
| 85 and over | 97 | 92 | 5 | 46 | 44 | 2 | 51 | 48 | 3 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 45,949 |
| White | 40,331 |
| Black | 3,855 |
| Other | 1,763 |

Selected Death Statistics, Residents of Harrison County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 1,948 | 1,575 | 346 | 978.6 | 1,133.2 | 724.0 |
| 1 | Heart Diseases | 502 | 407 | 92 | 252.2 | 292.8 | 192.5 |
| 2 | Malignant Neoplasms | 430 | 349 | 72 | 216.0 | 251.1 | 150.6 |
| 3 | Accidents | 123 | 102 | 19 | 61.8 | 73.4 | 39.8 |
| 4 | Emphysema and Other Chronic Lower Resp. Diseases | 104 | 99 | 4 | 52.2 | 71.2 | 8.4 |
| 5 | Cerebrovascular Disease | 100 | 72 | 25 | 50.2 | 51.8 | 52.3 |
| 6 | Nephritis, Nephrotic Syndrome And Nephrosis | 52 | 35 | 16 | 26.1 | 25.2 | 33.5 |
| 7 | Septicemia | 44 | 34 | 10 | 22.1 | 24.5 | 20.9 |
| 8 | Influenza and Pneumonia | 43 | 37 | 6 | 21.6 | 26.6 | 12.6 |
| 8 | Alzheimer's Disease | 43 | 35 | 7 | 21.6 | 25.2 | 14.6 |
| 10 | Suicide | 35 | 30 | 4 | 17.6 | 21.6 | 8.4 |
| 11 | Chronic Liver Disease And Cirrhosis | 34 | 27 | 7 | 17.1 | 19.4 | 14.6 |
| 12 | Parkinson's Disease | 15 | 13 | 2 | 7.5 | 9.4 | 4.2 |
| 13 | Homicide | 13 | 4 | 9 | 6.5 | 2.9 | 18.8 |
| 14 | Pneumonitis Due to Solids and Liquids | 12 | 10 | 2 | 6.0 | 7.2 | 4.2 |
| 15 | Conditions Originating in the Perinatal Period** | 10 | 4 | 5 | 3.5 | 2.1 | 6.4 |
| 15 | Diabetes Mellitus | 10 | 6 | 4 | 5.0 | 4.3 | 8.4 |
| 17 | Congenital Malformations and Chrom. Abnormalities | 9 | 7 | 2 | 4.5 | 5.0 | 4.2 |
| 17 | Hypertension/Hypertensive Renal Disease | 9 | 5 | 3 | 4.5 | 3.6 | 6.3 |
| 19 | HIV Disease | 8 | 2 | 6 | 4.0 | 1.4 | 12.6 |
| 19 | Viral Hepatitis | 8 | 8 | 0 | 4.0 | 5.8 | 0.0 |
| 21 | Complications Medical/Surgical Care | 6 | 5 | 1 | 3.0 | 3.6 | 2.1 |
| 21 | Anemias | 6 | 5 | 1 | 3.0 | 3.6 | 2.1 |
| 23 | Aortic Aneurysm and Dissection | 5 | 4 | 1 | 2.5 | 2.9 | 2.1 |
| 24 | Peptic Ulcer | 3 | 3 | 0 | 1.5 | 2.2 | 0.0 |
| 24 | Malnutrition/Nutritional Deficiency | 3 | 2 | 0 | 1.5 | 1.4 | 0.0 |
| 24 | In Situ, Benign and Unspecified Neoplasms | 3 | 3 | 0 | 1.5 | 2.2 | 0.0 |
| 27 | Cholelithiasis/Gallbladder Disorder | 2 | 1 | 1 | 1.0 | 0.7 | 2.1 |
| 28 | Legal Intervention | 1 | 0 | 1 | 0.5 | 0.0 | 2.1 |
| 28 | Maternal Causes** | 1 | 1 | 0 | 34.7 | 51.8 | 0.0 |
| 28 | Infections of Kidney | 1 | 1 | 0 | 0.5 | 0.7 | 0.0 |
| 28 | Diseases of Appendix | 1 | 1 | 0 | 0.5 | 0.7 | 0.0 |
| 28 | Pneumoconioses and Chemical Effects | 1 | 1 | 0 | 0.5 | 0.7 | 0.0 |
| 28 | Meningitis | 1 | 1 | 0 | 0.5 | 0.7 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 31 | 28 | 3 | 15.6 | 20.1 | 6.3 |
| | All Other Causes | 279 | 233 | 43 | 140.2 | 167.6 | 90.0 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 1,948 | 1,575 | 346 | 1,069 | 870 | 182 | 879 | 705 | 164 |
| Under 1 | 23 | 15 | 7 | 13 | 10 | 2 | 10 | 5 | 5 |
| (Neonatal) | 16 | 9 | 6 | 10 | 7 | 2 | 6 | 2 | 4 |
| (Postneonatal) | 7 | 6 | 1 | 3 | 3 | 0 | 4 | 3 | 1 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 2 | 0 | 2 | 0 | 0 | 0 | 2 | 0 | 2 |
| 10-14 | 3 | 3 | 0 | 1 | 1 | 0 | 2 | 2 | 0 |
| 15-24 | 27 | 18 | 9 | 19 | 12 | 7 | 8 | 6 | 2 |
| 25-34 | 42 | 31 | 11 | 34 | 25 | 9 | 8 | 6 | 2 |
| 35-44 | 60 | 42 | 15 | 36 | 26 | 7 | 24 | 16 | 8 |
| 45-54 | 186 | 140 | 41 | 112 | 84 | 24 | 74 | 56 | 17 |
| 55-64 | 327 | 239 | 80 | 205 | 156 | 46 | 122 | 83 | 34 |
| 65-74 | 374 | 307 | 64 | 224 | 181 | 41 | 150 | 126 | 23 |
| 75-84 | 458 | 393 | 61 | 240 | 207 | 31 | 218 | 186 | 30 |
| 85 and over | 446 | 387 | 56 | 185 | 168 | 15 | 261 | 219 | 41 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|---------|
| Total | 199,058 |
| White | 138,988 |
| Black | 47,793 |
| Other | 12,277 |

Selected Death Statistics, Residents of Hinds County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 2,152 | 836 | 1,308 | 882.9 | 1,258.1 | 757.9 |
| 1 | Heart Diseases | 567 | 194 | 370 | 232.6 | 292.0 | 214.4 |
| 2 | Malignant Neoplasms | 404 | 141 | 261 | 165.8 | 212.2 | 151.2 |
| 3 | Cerebrovascular Disease | 123 | 47 | 76 | 50.5 | 70.7 | 44.0 |
| 4 | Emphysema and Other Chronic Lower Resp. Diseases | 85 | 56 | 29 | 34.9 | 84.3 | 16.8 |
| 5 | Accidents | 73 | 29 | 44 | 30.0 | 43.6 | 25.5 |
| 6 | Alzheimer's Disease | 72 | 44 | 28 | 29.5 | 66.2 | 16.2 |
| 7 | Homicide | 69 | 5 | 64 | 28.3 | 7.5 | 37.1 |
| 8 | Diabetes Mellitus | 60 | 17 | 42 | 24.6 | 25.6 | 24.3 |
| 8 | Septicemia | 60 | 19 | 41 | 24.6 | 28.6 | 23.8 |
| 10 | Influenza and Pneumonia | 45 | 16 | 29 | 18.5 | 24.1 | 16.8 |
| 10 | Nephritis, Nephrotic Syndrome And Nephrosis | 45 | 12 | 33 | 18.5 | 18.1 | 19.1 |
| 12 | Hypertension/Hypertensive Renal Disease | 35 | 17 | 18 | 14.4 | 25.6 | 10.4 |
| 13 | HIV Disease | 34 | 4 | 30 | 13.9 | 6.0 | 17.4 |
| 14 | Chronic Liver Disease And Cirrhosis | 32 | 21 | 10 | 13.1 | 31.6 | 5.8 |
| 15 | Suicide | 22 | 12 | 9 | 9.0 | 18.1 | 5.2 |
| 16 | Parkinson's Disease | 19 | 15 | 4 | 7.8 | 22.6 | 2.3 |
| 17 | Pneumonitis Due to Solids and Liquids | 13 | 9 | 4 | 5.3 | 13.5 | 2.3 |
| 18 | Conditions Originating in the Perinatal Period** | 12 | 0 | 12 | 3.6 | 0.0 | 4.7 |
| 19 | Congenital Malformations and Chrom. Abnormalities | 11 | 3 | 8 | 4.5 | 4.5 | 4.6 |
| 20 | Viral Hepatitis | 6 | 4 | 2 | 2.5 | 6.0 | 1.2 |
| 21 | Atherosclerosis | 5 | 0 | 5 | 2.1 | 0.0 | 2.9 |
| 21 | In Situ, Benign and Unspecified Neoplasms | 5 | 3 | 2 | 2.1 | 4.5 | 1.2 |
| 23 | Anemias | 4 | 0 | 4 | 1.6 | 0.0 | 2.3 |
| 24 | Malnutrition/Nutritional Deficiency | 3 | 1 | 2 | 1.2 | 1.5 | 1.2 |
| 25 | Cholelithiasis/Gallbladder Disorder | 2 | 1 | 1 | 0.8 | 1.5 | 0.6 |
| 25 | Hernia | 2 | 2 | 0 | 0.8 | 3.0 | 0.0 |
| 25 | Peptic Ulcer | 2 | 1 | 1 | 0.8 | 1.5 | 0.6 |
| 25 | Aortic Aneurysm and Dissection | 2 | 0 | 2 | 0.8 | 0.0 | 1.2 |
| 29 | Diseases of Appendix | 1 | 1 | 0 | 0.4 | 1.5 | 0.0 |
| 29 | Meningitis | 1 | 0 | 1 | 0.4 | 0.0 | 0.6 |
| 29 | Syphilis | 1 | 0 | 1 | 0.4 | 0.0 | 0.6 |
| 29 | Tuberculosis | 1 | 0 | 1 | 0.4 | 0.0 | 0.6 |
| 29 | Complications Medical/Surgical Care | 1 | 0 | 1 | 0.4 | 0.0 | 0.6 |
| 29 | Female Pelvic Inflammatory Disease | 1 | 0 | 1 | 0.8 | 0.0 | 1.1 |
| 29 | Hyperplasia of Prostate | 1 | 0 | 1 | 0.9 | 0.0 | 1.3 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 24 | 9 | 15 | 9.8 | 13.5 | 8.7 |
| | All Other Causes | 309 | 153 | 156 | 126.8 | 230.3 | 90.4 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total* ----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|--------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 2,152 | 836 | 1,308 | 1,161 | 460 | 697 | 991 | 376 | 611 |
| Under 1 | 34 | 1 | 33 | 16 | 1 | 15 | 18 | 0 | 18 |
| (Neonatal) | 17 | 0 | 17 | 9 | 0 | 9 | 8 | 0 | 8 |
| (Postneonatal) | 17 | 1 | 16 | 7 | 1 | 6 | 10 | 0 | 10 |
| 1-4 | 7 | 0 | 7 | 4 | 0 | 4 | 3 | 0 | 3 |
| 5-9 | 2 | 2 | 0 | 2 | 2 | 0 | 0 | 0 | 0 |
| 10-14 | 2 | 0 | 2 | 2 | 0 | 2 | 0 | 0 | 0 |
| 15-24 | 51 | 8 | 43 | 41 | 5 | 36 | 10 | 3 | 7 |
| 25-34 | 63 | 16 | 46 | 47 | 13 | 34 | 16 | 3 | 12 |
| 35-44 | 101 | 20 | 81 | 59 | 13 | 46 | 42 | 7 | 35 |
| 45-54 | 192 | 45 | 143 | 116 | 29 | 84 | 76 | 16 | 59 |
| 55-64 | 360 | 100 | 258 | 244 | 74 | 169 | 116 | 26 | 89 |
| 65-74 | 391 | 148 | 242 | 247 | 97 | 150 | 144 | 51 | 92 |
| 75-84 | 445 | 212 | 233 | 203 | 97 | 106 | 242 | 115 | 127 |
| 85 and over | 502 | 283 | 219 | 178 | 128 | 50 | 324 | 155 | 169 |
| Unknown | 2 | 1 | 1 | 2 | 1 | 1 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|---------|
| Total | 243,729 |
| White | 66,449 |
| Black | 172,583 |
| Other | 4,697 |

Selected Death Statistics, Residents of Holmes County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 225 | 45 | 178 | 1,218.9 | 1,467.2 | 1,173.4 |
| 1 | Heart Diseases | 55 | 12 | 42 | 298.0 | 391.3 | 276.9 |
| 2 | Malignant Neoplasms | 41 | 7 | 34 | 222.1 | 228.2 | 224.1 |
| 3 | Accidents | 20 | 2 | 18 | 108.3 | 65.2 | 118.7 |
| 4 | Hypertension/Hypertensive Renal Disease | 18 | 4 | 14 | 97.5 | 130.4 | 92.3 |
| 5 | Cerebrovascular Disease | 14 | 7 | 7 | 75.8 | 228.2 | 46.1 |
| 5 | Diabetes Mellitus | 14 | 1 | 13 | 75.8 | 32.6 | 85.7 |
| 7 | Influenza and Pneumonia | 6 | 1 | 5 | 32.5 | 32.6 | 33.0 |
| 8 | Conditions Originating in the Perinatal Period** | 5 | 0 | 5 | 18.4 | 0.0 | 20.1 |
| 8 | Septicemia | 5 | 2 | 3 | 27.1 | 65.2 | 19.8 |
| 10 | Homicide | 4 | 0 | 4 | 21.7 | 0.0 | 26.4 |
| 10 | Alzheimer's Disease | 4 | 2 | 2 | 21.7 | 65.2 | 13.2 |
| 12 | Nephritis, Nephrotic Syndrome And Nephrosis | 3 | 0 | 3 | 16.3 | 0.0 | 19.8 |
| 12 | Emphysema and Other Chronic Lower Resp. Diseases | 3 | 1 | 2 | 16.3 | 32.6 | 13.2 |
| 14 | Suicide | 2 | 1 | 0 | 10.8 | 32.6 | 0.0 |
| 14 | Malnutrition/Nutritional Deficiency | 2 | 0 | 2 | 10.8 | 0.0 | 13.2 |
| 14 | HIV Disease | 2 | 0 | 2 | 10.8 | 0.0 | 13.2 |
| 17 | Infections of Kidney | 1 | 0 | 1 | 5.4 | 0.0 | 6.6 |
| 17 | Chronic Liver Disease And Cirrhosis | 1 | 0 | 1 | 5.4 | 0.0 | 6.6 |
| 17 | Pneumonitis Due to Solids and Liquids | 1 | 0 | 1 | 5.4 | 0.0 | 6.6 |
| 17 | In Situ, Benign and Unspecified Neoplasms | 1 | 0 | 1 | 5.4 | 0.0 | 6.6 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 2 | 0 | 2 | 10.8 | 0.0 | 13.2 |
| | All Other Causes | 21 | 5 | 16 | 113.8 | 163.0 | 105.5 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births.
 Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 225 | 45 | 178 | 117 | 23 | 93 | 108 | 22 | 85 |
| Under 1 | 5 | 0 | 5 | 3 | 0 | 3 | 2 | 0 | 2 |
| (Neonatal) | 5 | 0 | 5 | 3 | 0 | 3 | 2 | 0 | 2 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 25-34 | 13 | 0 | 13 | 7 | 0 | 7 | 6 | 0 | 6 |
| 35-44 | 10 | 0 | 10 | 8 | 0 | 8 | 2 | 0 | 2 |
| 45-54 | 22 | 3 | 19 | 11 | 1 | 10 | 11 | 2 | 9 |
| 55-64 | 37 | 5 | 31 | 24 | 4 | 19 | 13 | 1 | 12 |
| 65-74 | 43 | 9 | 34 | 25 | 6 | 19 | 18 | 3 | 15 |
| 75-84 | 40 | 11 | 29 | 20 | 6 | 14 | 20 | 5 | 15 |
| 85 and over | 53 | 17 | 35 | 17 | 6 | 11 | 36 | 11 | 24 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 18,459 |
| White | 3,067 |
| Black | 15,170 |
| Other | 222 |

Selected Death Statistics, Residents of Humphreys County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 93 | 31 | 62 | 1,064.0 | 1,508.5 | 946.3 |
| 1 | Heart Diseases | 31 | 12 | 19 | 354.7 | 583.9 | 290.0 |
| 2 | Malignant Neoplasms | 22 | 3 | 19 | 251.7 | 146.0 | 290.0 |
| 3 | Accidents | 6 | 3 | 3 | 68.6 | 146.0 | 45.8 |
| 3 | Diabetes Mellitus | 6 | 2 | 4 | 68.6 | 97.3 | 61.1 |
| 5 | Cerebrovascular Disease | 4 | 0 | 4 | 45.8 | 0.0 | 61.1 |
| 5 | Alzheimer's Disease | 4 | 4 | 0 | 45.8 | 194.6 | 0.0 |
| 7 | Nephritis, Nephrotic Syndrome And Nephrosis | 3 | 0 | 3 | 34.3 | 0.0 | 45.8 |
| 7 | Influenza and Pneumonia | 3 | 1 | 2 | 34.3 | 48.7 | 30.5 |
| 9 | Emphysema and Other Chronic Lower Resp. Diseases | 2 | 2 | 0 | 22.9 | 97.3 | 0.0 |
| 9 | Septicemia | 2 | 0 | 2 | 22.9 | 0.0 | 30.5 |
| 11 | Parkinson's Disease | 1 | 1 | 0 | 11.4 | 48.7 | 0.0 |
| 11 | HIV Disease | 1 | 0 | 1 | 11.4 | 0.0 | 15.3 |
| 11 | Viral Hepatitis | 1 | 0 | 1 | 11.4 | 0.0 | 15.3 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 0 | 0 | 0 | 0.0 | 0.0 | 0.0 |
| | All Other Causes | 7 | 3 | 4 | 80.1 | 146.0 | 61.1 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births.
 Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 93 | 31 | 62 | 45 | 13 | 32 | 48 | 18 | 30 |
| Under 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Neonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| 25-34 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| 35-44 | 7 | 1 | 6 | 4 | 1 | 3 | 3 | 0 | 3 |
| 45-54 | 10 | 1 | 9 | 7 | 0 | 7 | 3 | 1 | 2 |
| 55-64 | 13 | 2 | 11 | 8 | 1 | 7 | 5 | 1 | 4 |
| 65-74 | 18 | 4 | 14 | 9 | 1 | 8 | 9 | 3 | 6 |
| 75-84 | 22 | 12 | 10 | 10 | 5 | 5 | 12 | 7 | 5 |
| 85 and over | 21 | 11 | 10 | 7 | 5 | 2 | 14 | 6 | 8 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|-------|
| Total | 8,741 |
| White | 2,055 |
| Black | 6,552 |
| Other | 134 |

Selected Death Statistics, Residents of Issaquena County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|-------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 11 | 3 | 8 | 787.4 | 613.5 | 895.9 |
| 1 | Malignant Neoplasms | 4 | 1 | 3 | 286.3 | 204.5 | 335.9 |
| 2 | Heart Diseases | 3 | 2 | 1 | 214.7 | 409.0 | 112.0 |
| 3 | Nephritis, Nephrotic Syndrome And Nephrosis | 1 | 0 | 1 | 71.6 | 0.0 | 112.0 |
| 3 | Hypertension/Hypertensive Renal Disease | 1 | 0 | 1 | 71.6 | 0.0 | 112.0 |
| 3 | In Situ, Benign and Unspecified Neoplasms | 1 | 0 | 1 | 71.6 | 0.0 | 112.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 0 | 0 | 0 | 0.0 | 0.0 | 0.0 |
| | All Other Causes | 1 | 0 | 1 | 71.6 | 0.0 | 112.0 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 11 | 3 | 8 | 6 | 3 | 3 | 5 | 0 | 5 |
| Under 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Neonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 25-34 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 35-44 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 45-54 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 55-64 | 4 | 1 | 3 | 2 | 1 | 1 | 2 | 0 | 2 |
| 65-74 | 3 | 1 | 2 | 2 | 1 | 1 | 1 | 0 | 1 |
| 75-84 | 3 | 1 | 2 | 2 | 1 | 1 | 1 | 0 | 1 |
| 85 and over | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|-------|
| Total | 1,397 |
| White | 489 |
| Black | 893 |
| Other | 15 |

Selected Death Statistics, Residents of Itawamba County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 284 | 274 | 10 | 1,207.1 | 1,267.5 | 626.2 |
| 1 | Heart Diseases | 64 | 59 | 5 | 272.0 | 272.9 | 313.1 |
| 2 | Malignant Neoplasms | 57 | 56 | 1 | 242.3 | 259.0 | 62.6 |
| 3 | Alzheimer's Disease | 24 | 24 | 0 | 102.0 | 111.0 | 0.0 |
| 4 | Emphysema and Other Chronic Lower Resp. Diseases | 22 | 22 | 0 | 93.5 | 101.8 | 0.0 |
| 5 | Accidents | 17 | 17 | 0 | 72.3 | 78.6 | 0.0 |
| 6 | Cerebrovascular Disease | 14 | 12 | 2 | 59.5 | 55.5 | 125.2 |
| 7 | Septicemia | 9 | 8 | 1 | 38.3 | 37.0 | 62.6 |
| 8 | Nephritis, Nephrotic Syndrome And Nephrosis | 7 | 7 | 0 | 29.8 | 32.4 | 0.0 |
| 9 | Diabetes Mellitus | 6 | 6 | 0 | 25.5 | 27.8 | 0.0 |
| 10 | Pneumonitis Due to Solids and Liquids | 4 | 4 | 0 | 17.0 | 18.5 | 0.0 |
| 11 | Influenza and Pneumonia | 3 | 3 | 0 | 12.8 | 13.9 | 0.0 |
| 12 | Suicide | 2 | 2 | 0 | 8.5 | 9.3 | 0.0 |
| 12 | Aortic Aneurysm and Dissection | 2 | 2 | 0 | 8.5 | 9.3 | 0.0 |
| 12 | Hypertension/Hypertensive Renal Disease | 2 | 2 | 0 | 8.5 | 9.3 | 0.0 |
| 12 | Anemias | 2 | 1 | 1 | 8.5 | 4.6 | 62.6 |
| 16 | Complications Medical/Surgical Care | 1 | 1 | 0 | 4.3 | 4.6 | 0.0 |
| 16 | Congenital Malformations and Chrom. Abnormalities | 1 | 1 | 0 | 4.3 | 4.6 | 0.0 |
| 16 | Cholelithiasis/Gallbladder Disorder | 1 | 1 | 0 | 4.3 | 4.6 | 0.0 |
| 16 | Chronic Liver Disease And Cirrhosis | 1 | 1 | 0 | 4.3 | 4.6 | 0.0 |
| 16 | In Situ, Benign and Unspecified Neoplasms | 1 | 1 | 0 | 4.3 | 4.6 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 5 | 5 | 0 | 21.3 | 23.1 | 0.0 |
| | All Other Causes | 39 | 39 | 0 | 165.8 | 180.4 | 0.0 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births.
 Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 284 | 274 | 10 | 133 | 126 | 7 | 151 | 148 | 3 |
| Under 1 | 2 | 2 | 0 | 1 | 1 | 0 | 1 | 1 | 0 |
| (Neonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 2 | 2 | 0 | 1 | 1 | 0 | 1 | 1 | 0 |
| 1-4 | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 3 | 3 | 0 | 1 | 1 | 0 | 2 | 2 | 0 |
| 25-34 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 35-44 | 6 | 5 | 1 | 2 | 2 | 0 | 4 | 3 | 1 |
| 45-54 | 21 | 21 | 0 | 14 | 14 | 0 | 7 | 7 | 0 |
| 55-64 | 38 | 35 | 3 | 22 | 19 | 3 | 16 | 16 | 0 |
| 65-74 | 57 | 54 | 3 | 32 | 30 | 2 | 25 | 24 | 1 |
| 75-84 | 78 | 76 | 2 | 35 | 33 | 2 | 43 | 43 | 0 |
| 85 and over | 77 | 76 | 1 | 25 | 25 | 0 | 52 | 51 | 1 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 23,527 |
| White | 21,618 |
| Black | 1,597 |
| Other | 312 |

Selected Death Statistics, Residents of Jackson County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 1,318 | 1,084 | 218 | 933.8 | 1,048.2 | 700.3 |
| 1 | Malignant Neoplasms | 295 | 250 | 40 | 209.0 | 241.8 | 128.5 |
| 2 | Heart Diseases | 279 | 223 | 53 | 197.7 | 215.6 | 170.2 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 91 | 82 | 8 | 64.5 | 79.3 | 25.7 |
| 4 | Accidents | 84 | 67 | 16 | 59.5 | 64.8 | 51.4 |
| 5 | Cerebrovascular Disease | 72 | 55 | 15 | 51.0 | 53.2 | 48.2 |
| 6 | Alzheimer's Disease | 37 | 32 | 3 | 26.2 | 30.9 | 9.6 |
| 7 | Septicemia | 35 | 29 | 6 | 24.8 | 28.0 | 19.3 |
| 8 | Suicide | 31 | 26 | 5 | 22.0 | 25.1 | 16.1 |
| 9 | Chronic Liver Disease And Cirrhosis | 26 | 23 | 3 | 18.4 | 22.2 | 9.6 |
| 9 | Influenza and Pneumonia | 26 | 21 | 5 | 18.4 | 20.3 | 16.1 |
| 11 | Hypertension/Hypertensive Renal Disease | 23 | 17 | 6 | 16.3 | 16.4 | 19.3 |
| 12 | Diabetes Mellitus | 21 | 10 | 11 | 14.9 | 9.7 | 35.3 |
| 13 | Nephritis, Nephrotic Syndrome And Nephrosis | 20 | 13 | 7 | 14.2 | 12.6 | 22.5 |
| 14 | Parkinson's Disease | 13 | 12 | 0 | 9.2 | 11.6 | 0.0 |
| 15 | Pneumonitis Due to Solids and Liquids | 8 | 6 | 2 | 5.7 | 5.8 | 6.4 |
| 15 | Viral Hepatitis | 8 | 8 | 0 | 5.7 | 7.7 | 0.0 |
| 17 | Complications Medical/Surgical Care | 7 | 6 | 1 | 5.0 | 5.8 | 3.2 |
| 17 | Homicide | 7 | 5 | 2 | 5.0 | 4.8 | 6.4 |
| 17 | Congenital Malformations and Chrom. Abnormalities | 7 | 4 | 2 | 5.0 | 3.9 | 6.4 |
| 17 | Conditions Originating in the Perinatal Period** | 7 | 6 | 1 | 4.2 | 5.0 | 2.5 |
| 17 | Aortic Aneurysm and Dissection | 7 | 6 | 1 | 5.0 | 5.8 | 3.2 |
| 22 | In Situ, Benign and Unspecified Neoplasms | 4 | 4 | 0 | 2.8 | 3.9 | 0.0 |
| 23 | Hernia | 2 | 2 | 0 | 1.4 | 1.9 | 0.0 |
| 23 | Anemias | 2 | 0 | 2 | 1.4 | 0.0 | 6.4 |
| 23 | HIV Disease | 2 | 1 | 1 | 1.4 | 1.0 | 3.2 |
| 26 | Cholelithiasis/Gallbladder Disorder | 1 | 0 | 1 | 0.7 | 0.0 | 3.2 |
| 26 | Diseases of Appendix | 1 | 1 | 0 | 0.7 | 1.0 | 0.0 |
| 26 | Pneumoconioses and Chemical Effects | 1 | 1 | 0 | 0.7 | 1.0 | 0.0 |
| 26 | Acute Bronchitis and Bronchiolitis | 1 | 1 | 0 | 0.7 | 1.0 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 5 | 3 | 2 | 3.5 | 2.9 | 6.4 |
| | All Other Causes | 195 | 170 | 25 | 138.2 | 164.4 | 80.3 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 1,318 | 1,084 | 218 | 693 | 565 | 120 | 625 | 519 | 98 |
| Under 1 | 12 | 10 | 2 | 6 | 5 | 1 | 6 | 5 | 1 |
| (Neonatal) | 10 | 9 | 1 | 5 | 4 | 1 | 5 | 5 | 0 |
| (Postneonatal) | 2 | 1 | 1 | 1 | 1 | 0 | 1 | 0 | 1 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 2 | 1 | 1 | 2 | 1 | 1 | 0 | 0 | 0 |
| 10-14 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 15-24 | 17 | 15 | 2 | 9 | 9 | 0 | 8 | 6 | 2 |
| 25-34 | 36 | 25 | 10 | 28 | 20 | 7 | 8 | 5 | 3 |
| 35-44 | 53 | 40 | 11 | 30 | 22 | 7 | 23 | 18 | 4 |
| 45-54 | 100 | 76 | 23 | 61 | 47 | 14 | 39 | 29 | 9 |
| 55-64 | 219 | 170 | 48 | 126 | 93 | 32 | 93 | 77 | 16 |
| 65-74 | 259 | 214 | 39 | 139 | 115 | 20 | 120 | 99 | 19 |
| 75-84 | 369 | 315 | 51 | 204 | 176 | 27 | 165 | 139 | 24 |
| 85 and over | 250 | 218 | 30 | 87 | 77 | 10 | 163 | 141 | 20 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|---------|
| Total | 141,137 |
| White | 103,412 |
| Black | 31,131 |
| Other | 6,594 |

Selected Death Statistics, Residents of Jasper County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 206 | 125 | 81 | 1,240.9 | 1,623.6 | 931.4 |
| 1 | Heart Diseases | 58 | 35 | 23 | 349.4 | 454.6 | 264.5 |
| 2 | Malignant Neoplasms | 42 | 29 | 13 | 253.0 | 376.7 | 149.5 |
| 3 | Accidents | 18 | 13 | 5 | 108.4 | 168.9 | 57.5 |
| 3 | Cerebrovascular Disease | 18 | 10 | 8 | 108.4 | 129.9 | 92.0 |
| 5 | Emphysema and Other Chronic Lower Resp. Diseases | 8 | 5 | 3 | 48.2 | 64.9 | 34.5 |
| 6 | Influenza and Pneumonia | 7 | 4 | 3 | 42.2 | 52.0 | 34.5 |
| 7 | Septicemia | 6 | 3 | 3 | 36.1 | 39.0 | 34.5 |
| 8 | Diabetes Mellitus | 5 | 2 | 3 | 30.1 | 26.0 | 34.5 |
| 9 | Chronic Liver Disease And Cirrhosis | 3 | 1 | 2 | 18.1 | 13.0 | 23.0 |
| 9 | Alzheimer's Disease | 3 | 2 | 1 | 18.1 | 26.0 | 11.5 |
| 11 | Suicide | 2 | 2 | 0 | 12.0 | 26.0 | 0.0 |
| 11 | Congenital Malformations and Chrom. Abnormalities | 2 | 1 | 1 | 12.0 | 13.0 | 11.5 |
| 11 | Conditions Originating in the Perinatal Period** | 2 | 0 | 2 | 9.3 | 0.0 | 15.3 |
| 11 | Nephritis, Nephrotic Syndrome And Nephrosis | 2 | 0 | 2 | 12.0 | 0.0 | 23.0 |
| 11 | Hypertension/Hypertensive Renal Disease | 2 | 2 | 0 | 12.0 | 26.0 | 0.0 |
| 16 | Homicide | 1 | 0 | 1 | 6.0 | 0.0 | 11.5 |
| 16 | Pneumonitis Due to Solids and Liquids | 1 | 0 | 1 | 6.0 | 0.0 | 11.5 |
| 16 | Parkinson's Disease | 1 | 1 | 0 | 6.0 | 13.0 | 0.0 |
| 16 | Anemias | 1 | 0 | 1 | 6.0 | 0.0 | 11.5 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 2 | 1 | 1 | 12.0 | 13.0 | 11.5 |
| | All Other Causes | 22 | 14 | 8 | 132.5 | 181.8 | 92.0 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 206 | 125 | 81 | 98 | 58 | 40 | 108 | 67 | 41 |
| Under 1 | 4 | 1 | 3 | 2 | 0 | 2 | 2 | 1 | 1 |
| (Neonatal) | 4 | 1 | 3 | 2 | 0 | 2 | 2 | 1 | 1 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 3 | 2 | 1 | 3 | 2 | 1 | 0 | 0 | 0 |
| 25-34 | 3 | 2 | 1 | 3 | 2 | 1 | 0 | 0 | 0 |
| 35-44 | 7 | 5 | 2 | 3 | 2 | 1 | 4 | 3 | 1 |
| 45-54 | 22 | 10 | 12 | 12 | 3 | 9 | 10 | 7 | 3 |
| 55-64 | 35 | 20 | 15 | 20 | 11 | 9 | 15 | 9 | 6 |
| 65-74 | 46 | 30 | 16 | 25 | 16 | 9 | 21 | 14 | 7 |
| 75-84 | 52 | 36 | 16 | 20 | 14 | 6 | 32 | 22 | 10 |
| 85 and over | 34 | 19 | 15 | 10 | 8 | 2 | 24 | 11 | 13 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 16,601 |
| White | 7,699 |
| Black | 8,697 |
| Other | 205 |

Selected Death Statistics, Residents of Jefferson County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 99 | 19 | 80 | 1,302.8 | 1,769.1 | 1,236.7 |
| 1 | Heart Diseases | 36 | 9 | 27 | 473.7 | 838.0 | 417.4 |
| 2 | Malignant Neoplasms | 24 | 2 | 22 | 315.8 | 186.2 | 340.1 |
| 3 | Cerebrovascular Disease | 7 | 0 | 7 | 92.1 | 0.0 | 108.2 |
| 4 | Accidents | 5 | 0 | 5 | 65.8 | 0.0 | 77.3 |
| 4 | Nephritis, Nephrotic Syndrome And Nephrosis | 5 | 1 | 4 | 65.8 | 93.1 | 61.8 |
| 6 | Diabetes Mellitus | 3 | 0 | 3 | 39.5 | 0.0 | 46.4 |
| 6 | Septicemia | 3 | 1 | 2 | 39.5 | 93.1 | 30.9 |
| 8 | Influenza and Pneumonia | 2 | 1 | 1 | 26.3 | 93.1 | 15.5 |
| 8 | Hypertension/Hypertensive Renal Disease | 2 | 2 | 0 | 26.3 | 186.2 | 0.0 |
| 10 | Homicide | 1 | 0 | 1 | 13.2 | 0.0 | 15.5 |
| 10 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 9.7 | 0.0 | 11.2 |
| 10 | Cholelithiasis/Gallbladder Disorder | 1 | 1 | 0 | 13.2 | 93.1 | 0.0 |
| 10 | Chronic Liver Disease And Cirrhosis | 1 | 0 | 1 | 13.2 | 0.0 | 15.5 |
| 10 | Emphysema and Other Chronic Lower Resp. Diseases | 1 | 0 | 1 | 13.2 | 0.0 | 15.5 |
| 10 | Alzheimer's Disease | 1 | 1 | 0 | 13.2 | 93.1 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 0 | 0 | 0 | 0.0 | 0.0 | 0.0 |
| | All Other Causes | 6 | 1 | 5 | 79.0 | 93.1 | 77.3 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 99 | 19 | 80 | 56 | 12 | 44 | 43 | 7 | 36 |
| Under 1 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| (Neonatal) | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 15-24 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 25-34 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| 35-44 | 2 | 0 | 2 | 1 | 0 | 1 | 1 | 0 | 1 |
| 45-54 | 9 | 0 | 9 | 5 | 0 | 5 | 4 | 0 | 4 |
| 55-64 | 24 | 5 | 19 | 17 | 4 | 13 | 7 | 1 | 6 |
| 65-74 | 21 | 5 | 16 | 11 | 3 | 8 | 10 | 2 | 8 |
| 75-84 | 24 | 4 | 20 | 12 | 3 | 9 | 12 | 1 | 11 |
| 85 and over | 15 | 5 | 10 | 7 | 2 | 5 | 8 | 3 | 5 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|-------|
| Total | 7,599 |
| White | 1,074 |
| Black | 6,469 |
| Other | 56 |

Selected Death Statistics, Residents of Jeff Davis County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 153 | 68 | 85 | 1,294.2 | 1,486.3 | 1,199.9 |
| 1 | Heart Diseases | 35 | 14 | 21 | 296.1 | 306.0 | 296.4 |
| 2 | Malignant Neoplasms | 27 | 16 | 11 | 228.4 | 349.7 | 155.3 |
| 3 | Accidents | 15 | 4 | 11 | 126.9 | 87.4 | 155.3 |
| 4 | Influenza and Pneumonia | 8 | 5 | 3 | 67.7 | 109.3 | 42.3 |
| 5 | Suicide | 7 | 6 | 1 | 59.2 | 131.1 | 14.1 |
| 6 | Cerebrovascular Disease | 6 | 2 | 4 | 50.8 | 43.7 | 56.5 |
| 6 | Septicemia | 6 | 2 | 4 | 50.8 | 43.7 | 56.5 |
| 8 | Emphysema and Other Chronic Lower Resp. Diseases | 5 | 3 | 2 | 42.3 | 65.6 | 28.2 |
| 8 | Hypertension/Hypertensive Renal Disease | 5 | 3 | 2 | 42.3 | 65.6 | 28.2 |
| 10 | Homicide | 4 | 0 | 4 | 33.8 | 0.0 | 56.5 |
| 11 | Nephritis, Nephrotic Syndrome And Nephrosis | 3 | 0 | 3 | 25.4 | 0.0 | 42.3 |
| 11 | Alzheimer's Disease | 3 | 1 | 2 | 25.4 | 21.9 | 28.2 |
| 13 | Atherosclerosis | 2 | 2 | 0 | 16.9 | 43.7 | 0.0 |
| 13 | HIV Disease | 2 | 0 | 2 | 16.9 | 0.0 | 28.2 |
| 15 | Congenital Malformations and Chrom. Abnormalities | 1 | 0 | 1 | 8.5 | 0.0 | 14.1 |
| 15 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 8.3 | 0.0 | 14.3 |
| 15 | Pneumoconioses and Chemical Effects | 1 | 1 | 0 | 8.5 | 21.9 | 0.0 |
| 15 | Aortic Aneurysm and Dissection | 1 | 1 | 0 | 8.5 | 21.9 | 0.0 |
| 15 | Parkinson's Disease | 1 | 1 | 0 | 8.5 | 21.9 | 0.0 |
| 15 | Diabetes Mellitus | 1 | 1 | 0 | 8.5 | 21.9 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 1 | 0 | 1 | 8.5 | 0.0 | 14.1 |
| | All Other Causes | 18 | 6 | 12 | 152.3 | 131.1 | 169.4 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births.
 Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 153 | 68 | 85 | 84 | 36 | 48 | 69 | 32 | 37 |
| Under 1 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| (Neonatal) | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 2 | 1 | 1 | 1 | 0 | 1 | 1 | 1 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 7 | 4 | 3 | 5 | 2 | 3 | 2 | 2 | 0 |
| 25-34 | 4 | 1 | 3 | 1 | 0 | 1 | 3 | 1 | 2 |
| 35-44 | 6 | 1 | 5 | 5 | 0 | 5 | 1 | 1 | 0 |
| 45-54 | 18 | 6 | 12 | 14 | 6 | 8 | 4 | 0 | 4 |
| 55-64 | 31 | 11 | 20 | 20 | 6 | 14 | 11 | 5 | 6 |
| 65-74 | 28 | 15 | 13 | 16 | 9 | 7 | 12 | 6 | 6 |
| 75-84 | 25 | 15 | 10 | 17 | 11 | 6 | 8 | 4 | 4 |
| 85 and over | 31 | 14 | 17 | 5 | 2 | 3 | 26 | 12 | 14 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 11,822 |
| White | 4,575 |
| Black | 7,084 |
| Other | 163 |

Selected Death Statistics, Residents of Jones County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 750 | 576 | 173 | 1,098.3 | 1,227.4 | 866.0 |
| 1 | Heart Diseases | 198 | 145 | 52 | 289.9 | 309.0 | 260.3 |
| 2 | Malignant Neoplasms | 143 | 112 | 31 | 209.4 | 238.7 | 155.2 |
| 3 | Cerebrovascular Disease | 43 | 33 | 10 | 63.0 | 70.3 | 50.1 |
| 4 | Accidents | 39 | 25 | 14 | 57.1 | 53.3 | 70.1 |
| 5 | Emphysema and Other Chronic Lower Resp. Diseases | 31 | 28 | 3 | 45.4 | 59.7 | 15.0 |
| 6 | Diabetes Mellitus | 27 | 16 | 11 | 39.5 | 34.1 | 55.1 |
| 7 | Alzheimer's Disease | 26 | 24 | 2 | 38.1 | 51.1 | 10.0 |
| 8 | Influenza and Pneumonia | 23 | 20 | 3 | 33.7 | 42.6 | 15.0 |
| 9 | Septicemia | 18 | 15 | 3 | 26.4 | 32.0 | 15.0 |
| 10 | Nephritis, Nephrotic Syndrome And Nephrosis | 14 | 9 | 5 | 20.5 | 19.2 | 25.0 |
| 11 | Hypertension/Hypertensive Renal Disease | 12 | 6 | 6 | 17.6 | 12.8 | 30.0 |
| 12 | Pneumonitis Due to Solids and Liquids | 11 | 10 | 1 | 16.1 | 21.3 | 5.0 |
| 12 | Parkinson's Disease | 11 | 10 | 1 | 16.1 | 21.3 | 5.0 |
| 14 | Homicide | 8 | 3 | 5 | 11.7 | 6.4 | 25.0 |
| 14 | Suicide | 8 | 8 | 0 | 11.7 | 17.0 | 0.0 |
| 16 | Chronic Liver Disease And Cirrhosis | 7 | 6 | 1 | 10.3 | 12.8 | 5.0 |
| 17 | Conditions Originating in the Perinatal Period** | 5 | 2 | 3 | 5.2 | 3.4 | 8.4 |
| 18 | In Situ, Benign and Unspecified Neoplasms | 3 | 2 | 1 | 4.4 | 4.3 | 5.0 |
| 19 | Congenital Malformations and Chrom. Abnormalities | 2 | 2 | 0 | 2.9 | 4.3 | 0.0 |
| 19 | Aortic Aneurysm and Dissection | 2 | 2 | 0 | 2.9 | 4.3 | 0.0 |
| 19 | Anemias | 2 | 2 | 0 | 2.9 | 4.3 | 0.0 |
| 22 | Complications Medical/Surgical Care | 1 | 0 | 1 | 1.5 | 0.0 | 5.0 |
| 22 | Cholelithiasis/Gallbladder Disorder | 1 | 1 | 0 | 1.5 | 2.1 | 0.0 |
| 22 | Meningitis | 1 | 1 | 0 | 1.5 | 2.1 | 0.0 |
| 22 | Tuberculosis | 1 | 1 | 0 | 1.5 | 2.1 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 25 | 15 | 10 | 36.6 | 32.0 | 50.1 |
| | All Other Causes | 88 | 78 | 10 | 128.9 | 166.2 | 50.1 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 750 | 576 | 173 | 386 | 298 | 88 | 364 | 278 | 85 |
| Under 1 | 9 | 4 | 5 | 7 | 3 | 4 | 2 | 1 | 1 |
| (Neonatal) | 6 | 3 | 3 | 5 | 3 | 2 | 1 | 0 | 1 |
| (Postneonatal) | 3 | 1 | 2 | 2 | 0 | 2 | 1 | 1 | 0 |
| 1-4 | 4 | 3 | 1 | 0 | 0 | 0 | 4 | 3 | 1 |
| 5-9 | 2 | 0 | 2 | 1 | 0 | 1 | 1 | 0 | 1 |
| 10-14 | 2 | 1 | 1 | 0 | 0 | 0 | 2 | 1 | 1 |
| 15-24 | 10 | 5 | 5 | 7 | 4 | 3 | 3 | 1 | 2 |
| 25-34 | 15 | 6 | 8 | 4 | 3 | 1 | 11 | 3 | 7 |
| 35-44 | 27 | 20 | 7 | 17 | 12 | 5 | 10 | 8 | 2 |
| 45-54 | 47 | 26 | 21 | 26 | 16 | 10 | 21 | 10 | 11 |
| 55-64 | 113 | 83 | 30 | 77 | 54 | 23 | 36 | 29 | 7 |
| 65-74 | 145 | 117 | 28 | 82 | 68 | 14 | 63 | 49 | 14 |
| 75-84 | 188 | 158 | 30 | 97 | 81 | 16 | 91 | 77 | 14 |
| 85 and over | 188 | 153 | 35 | 68 | 57 | 11 | 120 | 96 | 24 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 68,290 |
| White | 46,927 |
| Black | 19,978 |
| Other | 1,385 |

Selected Death Statistics, Residents of Kemper County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 99 | 44 | 55 | 974.1 | 1,233.2 | 896.3 |
| 1 | Heart Diseases | 29 | 16 | 13 | 285.3 | 448.4 | 211.9 |
| 2 | Malignant Neoplasms | 25 | 12 | 13 | 246.0 | 336.3 | 211.9 |
| 3 | Cerebrovascular Disease | 6 | 1 | 5 | 59.0 | 28.0 | 81.5 |
| 4 | Accidents | 5 | 1 | 4 | 49.2 | 28.0 | 65.2 |
| 4 | Influenza and Pneumonia | 5 | 2 | 3 | 49.2 | 56.1 | 48.9 |
| 4 | Alzheimer's Disease | 5 | 3 | 2 | 49.2 | 84.1 | 32.6 |
| 7 | Nephritis, Nephrotic Syndrome And Nephrosis | 4 | 0 | 4 | 39.4 | 0.0 | 65.2 |
| 8 | Pneumonitis Due to Solids and Liquids | 3 | 1 | 2 | 29.5 | 28.0 | 32.6 |
| 8 | Emphysema and Other Chronic Lower Resp. Diseases | 3 | 2 | 1 | 29.5 | 56.1 | 16.3 |
| 8 | Diabetes Mellitus | 3 | 1 | 2 | 29.5 | 28.0 | 32.6 |
| 8 | Septicemia | 3 | 0 | 3 | 29.5 | 0.0 | 48.9 |
| 12 | Suicide | 1 | 1 | 0 | 9.8 | 28.0 | 0.0 |
| 12 | Homicide | 1 | 0 | 1 | 9.8 | 0.0 | 16.3 |
| 12 | Aortic Aneurysm and Dissection | 1 | 0 | 1 | 9.8 | 0.0 | 16.3 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 1 | 1 | 0 | 9.8 | 28.0 | 0.0 |
| | All Other Causes | 4 | 3 | 1 | 39.4 | 84.1 | 16.3 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 99 | 44 | 55 | 52 | 25 | 27 | 47 | 19 | 28 |
| Under 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Neonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 25-34 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 35-44 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 45-54 | 13 | 4 | 9 | 7 | 2 | 5 | 6 | 2 | 4 |
| 55-64 | 21 | 8 | 13 | 12 | 6 | 6 | 9 | 2 | 7 |
| 65-74 | 22 | 10 | 12 | 14 | 7 | 7 | 8 | 3 | 5 |
| 75-84 | 14 | 9 | 5 | 8 | 5 | 3 | 6 | 4 | 2 |
| 85 and over | 26 | 13 | 13 | 8 | 5 | 3 | 18 | 8 | 10 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 10,163 |
| White | 3,568 |
| Black | 6,136 |
| Other | 459 |

Selected Death Statistics, Residents of Lafayette County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|-------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 350 | 267 | 81 | 661.3 | 700.4 | 645.1 |
| 1 | Heart Diseases | 89 | 73 | 15 | 168.1 | 191.5 | 119.5 |
| 2 | Malignant Neoplasms | 71 | 46 | 25 | 134.1 | 120.7 | 199.1 |
| 3 | Cerebrovascular Disease | 29 | 24 | 5 | 54.8 | 63.0 | 39.8 |
| 4 | Emphysema and Other Chronic Lower Resp. Diseases | 17 | 14 | 3 | 32.1 | 36.7 | 23.9 |
| 5 | Accidents | 15 | 14 | 1 | 28.3 | 36.7 | 8.0 |
| 5 | Alzheimer's Disease | 15 | 10 | 5 | 28.3 | 26.2 | 39.8 |
| 7 | Septicemia | 11 | 8 | 3 | 20.8 | 21.0 | 23.9 |
| 8 | Influenza and Pneumonia | 10 | 8 | 2 | 18.9 | 21.0 | 15.9 |
| 9 | Nephritis, Nephrotic Syndrome And Nephrosis | 6 | 3 | 3 | 11.3 | 7.9 | 23.9 |
| 9 | Parkinson's Disease | 6 | 6 | 0 | 11.3 | 15.7 | 0.0 |
| 11 | Suicide | 4 | 2 | 1 | 7.6 | 5.2 | 8.0 |
| 12 | Chronic Liver Disease And Cirrhosis | 3 | 2 | 1 | 5.7 | 5.2 | 8.0 |
| 13 | Conditions Originating in the Perinatal Period** | 2 | 1 | 1 | 3.7 | 2.8 | 6.3 |
| 13 | Pneumonitis Due to Solids and Liquids | 2 | 2 | 0 | 3.8 | 5.2 | 0.0 |
| 13 | Aortic Aneurysm and Dissection | 2 | 2 | 0 | 3.8 | 5.2 | 0.0 |
| 13 | Diabetes Mellitus | 2 | 1 | 1 | 3.8 | 2.6 | 8.0 |
| 17 | Homicide | 1 | 0 | 1 | 1.9 | 0.0 | 8.0 |
| 17 | Congenital Malformations and Chrom. Abnormalities | 1 | 1 | 0 | 1.9 | 2.6 | 0.0 |
| 17 | Peptic Ulcer | 1 | 1 | 0 | 1.9 | 2.6 | 0.0 |
| 17 | In Situ, Benign and Unspecified Neoplasms | 1 | 1 | 0 | 1.9 | 2.6 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 1 | 1 | 0 | 1.9 | 2.6 | 0.0 |
| | All Other Causes | 61 | 47 | 14 | 115.2 | 123.3 | 111.5 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births.
 Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 350 | 267 | 81 | 198 | 159 | 37 | 152 | 108 | 44 |
| Under 1 | 3 | 2 | 1 | 2 | 2 | 0 | 1 | 0 | 1 |
| (Neonatal) | 2 | 1 | 1 | 1 | 1 | 0 | 1 | 0 | 1 |
| (Postneonatal) | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 15-24 | 2 | 0 | 2 | 0 | 0 | 0 | 2 | 0 | 2 |
| 25-34 | 3 | 2 | 0 | 3 | 2 | 0 | 0 | 0 | 0 |
| 35-44 | 9 | 6 | 3 | 4 | 2 | 2 | 5 | 4 | 1 |
| 45-54 | 33 | 17 | 15 | 21 | 11 | 9 | 12 | 6 | 6 |
| 55-64 | 41 | 25 | 16 | 27 | 15 | 12 | 14 | 10 | 4 |
| 65-74 | 67 | 53 | 14 | 42 | 36 | 6 | 25 | 17 | 8 |
| 75-84 | 77 | 61 | 16 | 41 | 36 | 5 | 36 | 25 | 11 |
| 85 and over | 114 | 100 | 14 | 57 | 54 | 3 | 57 | 46 | 11 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 52,930 |
| White | 38,123 |
| Black | 12,556 |
| Other | 2,251 |

Selected Death Statistics, Residents of Lamar County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|-------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 394 | 342 | 51 | 655.6 | 744.4 | 411.0 |
| 1 | Heart Diseases | 106 | 93 | 12 | 176.4 | 202.4 | 96.7 |
| 2 | Malignant Neoplasms | 77 | 61 | 16 | 128.1 | 132.8 | 128.9 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 34 | 33 | 1 | 56.6 | 71.8 | 8.1 |
| 4 | Accidents | 28 | 26 | 2 | 46.6 | 56.6 | 16.1 |
| 5 | Cerebrovascular Disease | 21 | 18 | 3 | 34.9 | 39.2 | 24.2 |
| 6 | Diabetes Mellitus | 15 | 14 | 1 | 25.0 | 30.5 | 8.1 |
| 7 | Alzheimer's Disease | 14 | 12 | 2 | 23.3 | 26.1 | 16.1 |
| 8 | Suicide | 6 | 5 | 1 | 10.0 | 10.9 | 8.1 |
| 8 | Nephritis, Nephrotic Syndrome And Nephrosis | 6 | 6 | 0 | 10.0 | 13.1 | 0.0 |
| 8 | Influenza and Pneumonia | 6 | 4 | 2 | 10.0 | 8.7 | 16.1 |
| 11 | Hypertension/Hypertensive Renal Disease | 5 | 4 | 1 | 8.3 | 8.7 | 8.1 |
| 11 | Parkinson's Disease | 5 | 5 | 0 | 8.3 | 10.9 | 0.0 |
| 13 | Homicide | 4 | 0 | 4 | 6.7 | 0.0 | 32.2 |
| 13 | Chronic Liver Disease And Cirrhosis | 4 | 4 | 0 | 6.7 | 8.7 | 0.0 |
| 13 | Septicemia | 4 | 4 | 0 | 6.7 | 8.7 | 0.0 |
| 16 | Congenital Malformations and Chrom. Abnormalities | 2 | 2 | 0 | 3.3 | 4.4 | 0.0 |
| 16 | Pneumonitis Due to Solids and Liquids | 2 | 2 | 0 | 3.3 | 4.4 | 0.0 |
| 16 | Aortic Aneurysm and Dissection | 2 | 2 | 0 | 3.3 | 4.4 | 0.0 |
| 16 | Anemias | 2 | 1 | 1 | 3.3 | 2.2 | 8.1 |
| 16 | In Situ, Benign and Unspecified Neoplasms | 2 | 2 | 0 | 3.3 | 4.4 | 0.0 |
| 21 | Complications Medical/Surgical Care | 1 | 1 | 0 | 1.7 | 2.2 | 0.0 |
| 21 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 1.3 | 0.0 | 6.3 |
| 21 | HIV Disease | 1 | 1 | 0 | 1.7 | 2.2 | 0.0 |
| 21 | Viral Hepatitis | 1 | 1 | 0 | 1.7 | 2.2 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 3 | 3 | 0 | 5.0 | 6.5 | 0.0 |
| | All Other Causes | 42 | 38 | 4 | 69.9 | 82.7 | 32.2 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births.
 Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 394 | 342 | 51 | 201 | 169 | 31 | 193 | 173 | 20 |
| Under 1 | 3 | 2 | 1 | 2 | 1 | 1 | 1 | 1 | 0 |
| (Neonatal) | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 2 | 1 | 1 | 1 | 0 | 1 | 1 | 1 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 15-24 | 14 | 11 | 3 | 9 | 7 | 2 | 5 | 4 | 1 |
| 25-34 | 10 | 8 | 2 | 5 | 3 | 2 | 5 | 5 | 0 |
| 35-44 | 6 | 4 | 2 | 5 | 3 | 2 | 1 | 1 | 0 |
| 45-54 | 34 | 29 | 5 | 17 | 14 | 3 | 17 | 15 | 2 |
| 55-64 | 60 | 45 | 14 | 34 | 25 | 8 | 26 | 20 | 6 |
| 65-74 | 80 | 70 | 10 | 50 | 43 | 7 | 30 | 27 | 3 |
| 75-84 | 96 | 89 | 7 | 43 | 40 | 3 | 53 | 49 | 4 |
| 85 and over | 90 | 84 | 6 | 35 | 33 | 2 | 55 | 51 | 4 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 60,099 |
| White | 45,942 |
| Black | 12,409 |
| Other | 1,748 |

Selected Death Statistics, Residents of Lauderdale County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 829 | 568 | 256 | 1,039.6 | 1,299.2 | 748.2 |
| 1 | Heart Diseases | 229 | 160 | 67 | 287.2 | 366.0 | 195.8 |
| 2 | Malignant Neoplasms | 160 | 110 | 49 | 200.7 | 251.6 | 143.2 |
| 3 | Accidents | 44 | 31 | 13 | 55.2 | 70.9 | 38.0 |
| 4 | Alzheimer's Disease | 43 | 37 | 6 | 53.9 | 84.6 | 17.5 |
| 5 | Cerebrovascular Disease | 41 | 28 | 13 | 51.4 | 64.0 | 38.0 |
| 6 | Emphysema and Other Chronic Lower Resp. Diseases | 40 | 34 | 6 | 50.2 | 77.8 | 17.5 |
| 7 | Nephritis, Nephrotic Syndrome And Nephrosis | 25 | 9 | 16 | 31.4 | 20.6 | 46.8 |
| 8 | Influenza and Pneumonia | 24 | 20 | 4 | 30.1 | 45.7 | 11.7 |
| 9 | Homicide | 15 | 3 | 12 | 18.8 | 6.9 | 35.1 |
| 9 | Septicemia | 15 | 9 | 6 | 18.8 | 20.6 | 17.5 |
| 11 | Suicide | 7 | 6 | 1 | 8.8 | 13.7 | 2.9 |
| 12 | Conditions Originating in the Perinatal Period** | 6 | 3 | 3 | 5.7 | 6.0 | 5.6 |
| 12 | Aortic Aneurysm and Dissection | 6 | 3 | 3 | 7.5 | 6.9 | 8.8 |
| 12 | Parkinson's Disease | 6 | 6 | 0 | 7.5 | 13.7 | 0.0 |
| 15 | Chronic Liver Disease And Cirrhosis | 4 | 3 | 1 | 5.0 | 6.9 | 2.9 |
| 16 | Pneumonitis Due to Solids and Liquids | 3 | 1 | 2 | 3.8 | 2.3 | 5.8 |
| 16 | In Situ, Benign and Unspecified Neoplasms | 3 | 2 | 1 | 3.8 | 4.6 | 2.9 |
| 18 | Complications Medical/Surgical Care | 2 | 1 | 1 | 2.5 | 2.3 | 2.9 |
| 18 | Diabetes Mellitus | 2 | 1 | 1 | 2.5 | 2.3 | 2.9 |
| 18 | Viral Hepatitis | 2 | 2 | 0 | 2.5 | 4.6 | 0.0 |
| 21 | Legal Intervention | 1 | 0 | 1 | 1.3 | 0.0 | 2.9 |
| 21 | Congenital Malformations and Chrom. Abnormalities | 1 | 0 | 1 | 1.3 | 0.0 | 2.9 |
| 21 | Hernia | 1 | 1 | 0 | 1.3 | 2.3 | 0.0 |
| 21 | Anemias | 1 | 0 | 1 | 1.3 | 0.0 | 2.9 |
| 21 | HIV Disease | 1 | 0 | 1 | 1.3 | 0.0 | 2.9 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 30 | 19 | 11 | 37.6 | 43.5 | 32.2 |
| | All Other Causes | 117 | 79 | 36 | 146.7 | 180.7 | 105.2 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 829 | 568 | 256 | 425 | 294 | 128 | 404 | 274 | 128 |
| Under 1 | 8 | 3 | 5 | 7 | 2 | 5 | 1 | 1 | 0 |
| (Neonatal) | 7 | 3 | 4 | 6 | 2 | 4 | 1 | 1 | 0 |
| (Postneonatal) | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 1-4 | 3 | 1 | 2 | 3 | 1 | 2 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 2 | 1 | 1 | 2 | 1 | 1 | 0 | 0 | 0 |
| 15-24 | 22 | 9 | 13 | 16 | 7 | 9 | 6 | 2 | 4 |
| 25-34 | 17 | 8 | 9 | 12 | 5 | 7 | 5 | 3 | 2 |
| 35-44 | 25 | 13 | 11 | 15 | 8 | 6 | 10 | 5 | 5 |
| 45-54 | 70 | 45 | 25 | 37 | 27 | 10 | 33 | 18 | 15 |
| 55-64 | 112 | 63 | 48 | 70 | 43 | 26 | 42 | 20 | 22 |
| 65-74 | 167 | 114 | 53 | 98 | 72 | 26 | 69 | 42 | 27 |
| 75-84 | 194 | 147 | 46 | 96 | 75 | 20 | 98 | 72 | 26 |
| 85 and over | 209 | 164 | 43 | 69 | 53 | 16 | 140 | 111 | 27 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 79,739 |
| White | 43,720 |
| Black | 34,214 |
| Other | 1,805 |

Selected Death Statistics, Residents of Lawrence County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 135 | 99 | 35 | 1,079.8 | 1,181.8 | 890.6 |
| 1 | Heart Diseases | 30 | 29 | 1 | 240.0 | 346.2 | 25.4 |
| 2 | Malignant Neoplasms | 27 | 19 | 8 | 216.0 | 226.8 | 203.6 |
| 3 | Accidents | 11 | 8 | 3 | 88.0 | 95.5 | 76.3 |
| 4 | Cerebrovascular Disease | 8 | 4 | 4 | 64.0 | 47.7 | 101.8 |
| 5 | Emphysema and Other Chronic Lower Resp. Diseases | 7 | 6 | 1 | 56.0 | 71.6 | 25.4 |
| 6 | Diabetes Mellitus | 5 | 1 | 4 | 40.0 | 11.9 | 101.8 |
| 7 | Septicemia | 4 | 2 | 2 | 32.0 | 23.9 | 50.9 |
| 8 | Conditions Originating in the Perinatal Period** | 2 | 1 | 1 | 12.1 | 9.3 | 17.9 |
| 8 | Nephritis, Nephrotic Syndrome And Nephrosis | 2 | 0 | 2 | 16.0 | 0.0 | 50.9 |
| 8 | Influenza and Pneumonia | 2 | 2 | 0 | 16.0 | 23.9 | 0.0 |
| 8 | Atherosclerosis | 2 | 1 | 1 | 16.0 | 11.9 | 25.4 |
| 8 | Hypertension/Hypertensive Renal Disease | 2 | 2 | 0 | 16.0 | 23.9 | 0.0 |
| 8 | Alzheimer's Disease | 2 | 2 | 0 | 16.0 | 23.9 | 0.0 |
| 14 | Homicide | 1 | 0 | 0 | 8.0 | 0.0 | 0.0 |
| 14 | Suicide | 1 | 1 | 0 | 8.0 | 11.9 | 0.0 |
| 14 | Chronic Liver Disease And Cirrhosis | 1 | 1 | 0 | 8.0 | 11.9 | 0.0 |
| 14 | HIV Disease | 1 | 0 | 1 | 8.0 | 0.0 | 25.4 |
| 14 | Viral Hepatitis | 1 | 1 | 0 | 8.0 | 11.9 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 2 | 2 | 0 | 16.0 | 23.9 | 0.0 |
| | All Other Causes | 24 | 17 | 7 | 192.0 | 202.9 | 178.1 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births.
 Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 135 | 99 | 35 | 66 | 48 | 17 | 69 | 51 | 18 |
| Under 1 | 3 | 1 | 2 | 1 | 0 | 1 | 2 | 1 | 1 |
| (Neonatal) | 2 | 1 | 1 | 1 | 0 | 1 | 1 | 1 | 0 |
| (Postneonatal) | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| 1-4 | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 25-34 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 35-44 | 9 | 5 | 4 | 3 | 1 | 2 | 6 | 4 | 2 |
| 45-54 | 14 | 9 | 4 | 7 | 5 | 1 | 7 | 4 | 3 |
| 55-64 | 29 | 19 | 10 | 20 | 13 | 7 | 9 | 6 | 3 |
| 65-74 | 25 | 19 | 6 | 13 | 12 | 1 | 12 | 7 | 5 |
| 75-84 | 27 | 23 | 4 | 14 | 11 | 3 | 13 | 12 | 1 |
| 85 and over | 25 | 21 | 4 | 6 | 5 | 1 | 19 | 16 | 3 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 12,502 |
| White | 8,377 |
| Black | 3,930 |
| Other | 195 |

Selected Death Statistics, Residents of Leake County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 259 | 168 | 80 | 1,116.7 | 1,447.7 | 822.3 |
| 1 | Heart Diseases | 70 | 44 | 25 | 301.8 | 379.1 | 257.0 |
| 2 | Malignant Neoplasms | 64 | 45 | 16 | 275.9 | 387.8 | 164.5 |
| 3 | Accidents | 18 | 13 | 3 | 77.6 | 112.0 | 30.8 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 18 | 17 | 1 | 77.6 | 146.5 | 10.3 |
| 5 | Cerebrovascular Disease | 11 | 2 | 8 | 47.4 | 17.2 | 82.2 |
| 6 | Alzheimer's Disease | 9 | 7 | 2 | 38.8 | 60.3 | 20.6 |
| 7 | Diabetes Mellitus | 6 | 2 | 3 | 25.9 | 17.2 | 30.8 |
| 8 | Homicide | 5 | 1 | 3 | 21.6 | 8.6 | 30.8 |
| 8 | Influenza and Pneumonia | 5 | 5 | 0 | 21.6 | 43.1 | 0.0 |
| 10 | Nephritis, Nephrotic Syndrome And Nephrosis | 4 | 2 | 2 | 17.2 | 17.2 | 20.6 |
| 10 | Hypertension/Hypertensive Renal Disease | 4 | 1 | 3 | 17.2 | 8.6 | 30.8 |
| 10 | Septicemia | 4 | 3 | 1 | 17.2 | 25.9 | 10.3 |
| 13 | Suicide | 3 | 3 | 0 | 12.9 | 25.9 | 0.0 |
| 13 | Chronic Liver Disease And Cirrhosis | 3 | 3 | 0 | 12.9 | 25.9 | 0.0 |
| 13 | Parkinson's Disease | 3 | 3 | 0 | 12.9 | 25.9 | 0.0 |
| 16 | Aortic Aneurysm and Dissection | 2 | 1 | 1 | 8.6 | 8.6 | 10.3 |
| 17 | Complications Medical/Surgical Care | 1 | 1 | 0 | 4.3 | 8.6 | 0.0 |
| 17 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 3.3 | 0.0 | 7.7 |
| 17 | In Situ, Benign and Unspecified Neoplasms | 1 | 0 | 1 | 4.3 | 0.0 | 10.3 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 1 | 0 | 1 | 4.3 | 0.0 | 10.3 |
| | All Other Causes | 26 | 15 | 9 | 112.1 | 129.3 | 92.5 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 259 | 168 | 80 | 139 | 88 | 46 | 120 | 80 | 34 |
| Under 1 | 2 | 0 | 2 | 1 | 0 | 1 | 1 | 0 | 1 |
| (Neonatal) | 2 | 0 | 2 | 1 | 0 | 1 | 1 | 0 | 1 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 2 | 0 | 2 | 2 | 0 | 2 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 2 | 2 | 0 | 2 | 2 | 0 | 0 | 0 | 0 |
| 15-24 | 4 | 2 | 0 | 4 | 2 | 0 | 0 | 0 | 0 |
| 25-34 | 6 | 4 | 2 | 3 | 2 | 1 | 3 | 2 | 1 |
| 35-44 | 18 | 9 | 7 | 10 | 4 | 6 | 8 | 5 | 1 |
| 45-54 | 18 | 7 | 10 | 14 | 6 | 7 | 4 | 1 | 3 |
| 55-64 | 48 | 31 | 16 | 30 | 17 | 12 | 18 | 14 | 4 |
| 65-74 | 54 | 36 | 15 | 29 | 19 | 9 | 25 | 17 | 6 |
| 75-84 | 53 | 37 | 14 | 25 | 20 | 5 | 28 | 17 | 9 |
| 85 and over | 52 | 40 | 12 | 19 | 16 | 3 | 33 | 24 | 9 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 23,193 |
| White | 11,605 |
| Black | 9,729 |
| Other | 1,859 |

Selected Death Statistics, Residents of Lee County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 926 | 752 | 168 | 1,086.3 | 1,278.7 | 686.9 |
| 1 | Malignant Neoplasms | 233 | 192 | 40 | 273.3 | 326.5 | 163.6 |
| 2 | Heart Diseases | 194 | 158 | 35 | 227.6 | 268.7 | 143.1 |
| 3 | Cerebrovascular Disease | 63 | 52 | 10 | 73.9 | 88.4 | 40.9 |
| 4 | Emphysema and Other Chronic Lower Resp. Diseases | 49 | 43 | 6 | 57.5 | 73.1 | 24.5 |
| 5 | Accidents | 46 | 35 | 10 | 54.0 | 59.5 | 40.9 |
| 6 | Alzheimer's Disease | 34 | 28 | 6 | 39.9 | 47.6 | 24.5 |
| 7 | Diabetes Mellitus | 21 | 14 | 7 | 24.6 | 23.8 | 28.6 |
| 8 | Hypertension/Hypertensive Renal Disease | 20 | 16 | 4 | 23.5 | 27.2 | 16.4 |
| 8 | Septicemia | 20 | 17 | 3 | 23.5 | 28.9 | 12.3 |
| 10 | Influenza and Pneumonia | 16 | 13 | 3 | 18.8 | 22.1 | 12.3 |
| 11 | Chronic Liver Disease And Cirrhosis | 14 | 13 | 1 | 16.4 | 22.1 | 4.1 |
| 12 | Nephritis, Nephrotic Syndrome And Nephrosis | 11 | 7 | 4 | 12.9 | 11.9 | 16.4 |
| 13 | Suicide | 7 | 6 | 1 | 8.2 | 10.2 | 4.1 |
| 13 | Pneumonitis Due to Solids and Liquids | 7 | 6 | 1 | 8.2 | 10.2 | 4.1 |
| 15 | Parkinson's Disease | 6 | 6 | 0 | 7.0 | 10.2 | 0.0 |
| 16 | Homicide | 5 | 2 | 2 | 5.9 | 3.4 | 8.2 |
| 16 | Conditions Originating in the Perinatal Period** | 5 | 0 | 5 | 4.3 | 0.0 | 11.6 |
| 18 | In Situ, Benign and Unspecified Neoplasms | 4 | 4 | 0 | 4.7 | 6.8 | 0.0 |
| 19 | Congenital Malformations and Chrom. Abnormalities | 3 | 3 | 0 | 3.5 | 5.1 | 0.0 |
| 19 | Malnutrition/Nutritional Deficiency | 3 | 1 | 2 | 3.5 | 1.7 | 8.2 |
| 21 | Complications Medical/Surgical Care | 1 | 1 | 0 | 1.2 | 1.7 | 0.0 |
| 21 | Legal Intervention | 1 | 1 | 0 | 1.2 | 1.7 | 0.0 |
| 21 | Infections of Kidney | 1 | 1 | 0 | 1.2 | 1.7 | 0.0 |
| 21 | Aortic Aneurysm and Dissection | 1 | 1 | 0 | 1.2 | 1.7 | 0.0 |
| 21 | Anemias | 1 | 1 | 0 | 1.2 | 1.7 | 0.0 |
| 21 | HIV Disease | 1 | 0 | 1 | 1.2 | 0.0 | 4.1 |
| 21 | Viral Hepatitis | 1 | 1 | 0 | 1.2 | 1.7 | 0.0 |
| 21 | Tuberculosis | 1 | 0 | 1 | 1.2 | 0.0 | 4.1 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 9 | 6 | 3 | 10.6 | 10.2 | 12.3 |
| | All Other Causes | 148 | 124 | 23 | 173.6 | 210.9 | 94.0 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 926 | 752 | 168 | 460 | 364 | 91 | 466 | 388 | 77 |
| Under 1 | 10 | 3 | 7 | 5 | 1 | 4 | 5 | 2 | 3 |
| (Neonatal) | 8 | 2 | 6 | 4 | 0 | 4 | 4 | 2 | 2 |
| (Postneonatal) | 2 | 1 | 1 | 1 | 1 | 0 | 1 | 0 | 1 |
| 1-4 | 2 | 2 | 0 | 1 | 1 | 0 | 1 | 1 | 0 |
| 5-9 | 2 | 1 | 1 | 1 | 1 | 0 | 1 | 0 | 1 |
| 10-14 | 2 | 1 | 1 | 1 | 0 | 1 | 1 | 1 | 0 |
| 15-24 | 11 | 8 | 3 | 9 | 6 | 3 | 2 | 2 | 0 |
| 25-34 | 14 | 10 | 3 | 8 | 5 | 3 | 6 | 5 | 0 |
| 35-44 | 24 | 13 | 10 | 13 | 6 | 6 | 11 | 7 | 4 |
| 45-54 | 75 | 55 | 19 | 48 | 33 | 14 | 27 | 22 | 5 |
| 55-64 | 134 | 101 | 32 | 83 | 60 | 22 | 51 | 41 | 10 |
| 65-74 | 186 | 152 | 33 | 98 | 80 | 17 | 88 | 72 | 16 |
| 75-84 | 243 | 209 | 33 | 126 | 110 | 15 | 117 | 99 | 18 |
| 85 and over | 223 | 197 | 26 | 67 | 61 | 6 | 156 | 136 | 20 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 85,246 |
| White | 58,809 |
| Black | 24,457 |
| Other | 1,980 |

Selected Death Statistics, Residents of Leflore County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 354 | 126 | 227 | 1,126.6 | 1,574.4 | 993.3 |
| 1 | Heart Diseases | 103 | 40 | 63 | 327.8 | 499.8 | 275.7 |
| 2 | Malignant Neoplasms | 86 | 27 | 58 | 273.7 | 337.4 | 253.8 |
| 3 | Cerebrovascular Disease | 22 | 5 | 17 | 70.0 | 62.5 | 74.4 |
| 4 | Diabetes Mellitus | 21 | 7 | 14 | 66.8 | 87.5 | 61.3 |
| 5 | Accidents | 15 | 2 | 13 | 47.7 | 25.0 | 56.9 |
| 6 | Emphysema and Other Chronic Lower Resp. Diseases | 11 | 7 | 4 | 35.0 | 87.5 | 17.5 |
| 6 | Hypertension/Hypertensive Renal Disease | 11 | 2 | 9 | 35.0 | 25.0 | 39.4 |
| 8 | Alzheimer's Disease | 9 | 8 | 1 | 28.6 | 100.0 | 4.4 |
| 9 | Nephritis, Nephrotic Syndrome And Nephrosis | 7 | 1 | 6 | 22.3 | 12.5 | 26.3 |
| 10 | Septicemia | 5 | 2 | 3 | 15.9 | 25.0 | 13.1 |
| 11 | Influenza and Pneumonia | 4 | 2 | 2 | 12.7 | 25.0 | 8.8 |
| 12 | Homicide | 3 | 1 | 2 | 9.5 | 12.5 | 8.8 |
| 12 | Congenital Malformations and Chrom. Abnormalities | 3 | 0 | 3 | 9.5 | 0.0 | 13.1 |
| 12 | Chronic Liver Disease And Cirrhosis | 3 | 1 | 2 | 9.5 | 12.5 | 8.8 |
| 12 | Anemias | 3 | 2 | 1 | 9.5 | 25.0 | 4.4 |
| 16 | HIV Disease | 2 | 0 | 2 | 6.4 | 0.0 | 8.8 |
| 17 | Complications Medical/Surgical Care | 1 | 0 | 1 | 3.2 | 0.0 | 4.4 |
| 17 | Conditions Originating in the Perinatal Period** | 1 | 1 | 0 | 2.0 | 8.9 | 0.0 |
| 17 | Parkinson's Disease | 1 | 1 | 0 | 3.2 | 12.5 | 0.0 |
| 17 | Malnutrition/Nutritional Deficiency | 1 | 1 | 0 | 3.2 | 12.5 | 0.0 |
| 17 | Viral Hepatitis | 1 | 1 | 0 | 3.2 | 12.5 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 3 | 0 | 3 | 9.5 | 0.0 | 13.1 |
| | All Other Causes | 38 | 15 | 23 | 120.9 | 187.4 | 100.6 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total* ----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|--------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 354 | 126 | 227 | 163 | 46 | 117 | 191 | 80 | 110 |
| Under 1 | 5 | 1 | 4 | 3 | 0 | 3 | 2 | 1 | 1 |
| (Neonatal) | 3 | 1 | 2 | 2 | 0 | 2 | 1 | 1 | 0 |
| (Postneonatal) | 2 | 0 | 2 | 1 | 0 | 1 | 1 | 0 | 1 |
| 1-4 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| 5-9 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 10-14 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 15-24 | 5 | 1 | 4 | 2 | 0 | 2 | 3 | 1 | 2 |
| 25-34 | 11 | 0 | 11 | 5 | 0 | 5 | 6 | 0 | 6 |
| 35-44 | 16 | 4 | 12 | 8 | 2 | 6 | 8 | 2 | 6 |
| 45-54 | 28 | 7 | 21 | 17 | 3 | 14 | 11 | 4 | 7 |
| 55-64 | 63 | 17 | 46 | 43 | 12 | 31 | 20 | 5 | 15 |
| 65-74 | 59 | 17 | 42 | 33 | 11 | 22 | 26 | 6 | 20 |
| 75-84 | 63 | 28 | 35 | 22 | 7 | 15 | 41 | 21 | 20 |
| 85 and over | 101 | 50 | 50 | 28 | 10 | 18 | 73 | 40 | 32 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 31,422 |
| White | 8,003 |
| Black | 22,853 |
| Other | 566 |

Selected Death Statistics, Residents of Lincoln County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 403 | 297 | 105 | 1,158.9 | 1,261.4 | 975.3 |
| 1 | Heart Diseases | 97 | 71 | 26 | 278.9 | 301.5 | 241.5 |
| 2 | Malignant Neoplasms | 84 | 58 | 26 | 241.6 | 246.3 | 241.5 |
| 3 | Accidents | 34 | 28 | 6 | 97.8 | 118.9 | 55.7 |
| 4 | Emphysema and Other Chronic Lower Resp. Diseases | 28 | 22 | 6 | 80.5 | 93.4 | 55.7 |
| 5 | Nephritis, Nephrotic Syndrome And Nephrosis | 16 | 12 | 4 | 46.0 | 51.0 | 37.2 |
| 6 | Influenza and Pneumonia | 14 | 12 | 2 | 40.3 | 51.0 | 18.6 |
| 7 | Cerebrovascular Disease | 13 | 11 | 2 | 37.4 | 46.7 | 18.6 |
| 8 | Hypertension/Hypertensive Renal Disease | 11 | 6 | 5 | 31.6 | 25.5 | 46.4 |
| 9 | Alzheimer's Disease | 10 | 10 | 0 | 28.8 | 42.5 | 0.0 |
| 10 | Diabetes Mellitus | 9 | 6 | 3 | 25.9 | 25.5 | 27.9 |
| 11 | Homicide | 4 | 0 | 4 | 11.5 | 0.0 | 37.2 |
| 12 | Parkinson's Disease | 3 | 2 | 1 | 8.6 | 8.5 | 9.3 |
| 13 | Chronic Liver Disease And Cirrhosis | 2 | 2 | 0 | 5.8 | 8.5 | 0.0 |
| 13 | Aortic Aneurysm and Dissection | 2 | 1 | 1 | 5.8 | 4.2 | 9.3 |
| 13 | In Situ, Benign and Unspecified Neoplasms | 2 | 1 | 1 | 5.8 | 4.2 | 9.3 |
| 13 | HIV Disease | 2 | 0 | 2 | 5.8 | 0.0 | 18.6 |
| 13 | Septicemia | 2 | 2 | 0 | 5.8 | 8.5 | 0.0 |
| 18 | Suicide | 1 | 1 | 0 | 2.9 | 4.2 | 0.0 |
| 18 | Congenital Malformations and Chrom. Abnormalities | 1 | 1 | 0 | 2.9 | 4.2 | 0.0 |
| 18 | Conditions Originating in the Perinatal Period** | 1 | 0 | 0 | 2.1 | 0.0 | 0.0 |
| 18 | Atherosclerosis | 1 | 1 | 0 | 2.9 | 4.2 | 0.0 |
| 18 | Malnutrition/Nutritional Deficiency | 1 | 0 | 1 | 2.9 | 0.0 | 9.3 |
| 18 | Anemias | 1 | 1 | 0 | 2.9 | 4.2 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 5 | 4 | 1 | 14.4 | 17.0 | 9.3 |
| | All Other Causes | 59 | 45 | 14 | 169.7 | 191.1 | 130.0 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 403 | 297 | 105 | 205 | 144 | 61 | 198 | 153 | 44 |
| Under 1 | 2 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 |
| (Neonatal) | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 |
| 1-4 | 2 | 2 | 0 | 1 | 1 | 0 | 1 | 1 | 0 |
| 5-9 | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 8 | 6 | 2 | 6 | 4 | 2 | 2 | 2 | 0 |
| 25-34 | 9 | 7 | 2 | 3 | 1 | 2 | 6 | 6 | 0 |
| 35-44 | 12 | 6 | 6 | 7 | 3 | 4 | 5 | 3 | 2 |
| 45-54 | 36 | 22 | 14 | 26 | 17 | 9 | 10 | 5 | 5 |
| 55-64 | 75 | 46 | 29 | 44 | 27 | 17 | 31 | 19 | 12 |
| 65-74 | 76 | 56 | 20 | 46 | 32 | 14 | 30 | 24 | 6 |
| 75-84 | 81 | 63 | 18 | 36 | 28 | 8 | 45 | 35 | 10 |
| 85 and over | 101 | 87 | 14 | 35 | 30 | 5 | 66 | 57 | 9 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 34,775 |
| White | 23,546 |
| Black | 10,766 |
| Other | 463 |

Selected Death Statistics, Residents of Lowndes County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 596 | 383 | 209 | 997.8 | 1,190.4 | 794.8 |
| 1 | Malignant Neoplasms | 130 | 75 | 55 | 217.6 | 233.1 | 209.2 |
| 2 | Heart Diseases | 116 | 79 | 37 | 194.2 | 245.5 | 140.7 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 52 | 38 | 14 | 87.1 | 118.1 | 53.2 |
| 4 | Cerebrovascular Disease | 33 | 21 | 12 | 55.2 | 65.3 | 45.6 |
| 5 | Alzheimer's Disease | 31 | 26 | 4 | 51.9 | 80.8 | 15.2 |
| 6 | Accidents | 26 | 19 | 7 | 43.5 | 59.1 | 26.6 |
| 7 | Nephritis, Nephrotic Syndrome And Nephrosis | 15 | 9 | 6 | 25.1 | 28.0 | 22.8 |
| 8 | Influenza and Pneumonia | 13 | 9 | 4 | 21.8 | 28.0 | 15.2 |
| 8 | Septicemia | 13 | 7 | 5 | 21.8 | 21.8 | 19.0 |
| 10 | Diabetes Mellitus | 11 | 3 | 8 | 18.4 | 9.3 | 30.4 |
| 11 | Chronic Liver Disease And Cirrhosis | 10 | 6 | 3 | 16.7 | 18.6 | 11.4 |
| 12 | Suicide | 6 | 6 | 0 | 10.0 | 18.6 | 0.0 |
| 13 | Homicide | 5 | 0 | 5 | 8.4 | 0.0 | 19.0 |
| 13 | Parkinson's Disease | 5 | 4 | 1 | 8.4 | 12.4 | 3.8 |
| 15 | Conditions Originating in the Perinatal Period** | 4 | 0 | 3 | 5.3 | 0.0 | 8.6 |
| 15 | HIV Disease | 4 | 0 | 4 | 6.7 | 0.0 | 15.2 |
| 15 | Viral Hepatitis | 4 | 2 | 2 | 6.7 | 6.2 | 7.6 |
| 18 | Hernia | 3 | 3 | 0 | 5.0 | 9.3 | 0.0 |
| 18 | Hypertension/Hypertensive Renal Disease | 3 | 2 | 1 | 5.0 | 6.2 | 3.8 |
| 20 | Congenital Malformations and Chrom. Abnormalities | 2 | 0 | 2 | 3.3 | 0.0 | 7.6 |
| 20 | Aortic Aneurysm and Dissection | 2 | 1 | 1 | 3.3 | 3.1 | 3.8 |
| 20 | Anemias | 2 | 0 | 2 | 3.3 | 0.0 | 7.6 |
| 23 | Maternal Causes** | 1 | 1 | 0 | 132.5 | 252.5 | 0.0 |
| 23 | Cholelithiasis/Gallbladder Disorder | 1 | 1 | 0 | 1.7 | 3.1 | 0.0 |
| 23 | Peptic Ulcer | 1 | 1 | 0 | 1.7 | 3.1 | 0.0 |
| 23 | Pneumonitis Due to Solids and Liquids | 1 | 0 | 1 | 1.7 | 0.0 | 3.8 |
| 23 | Malnutrition/Nutritional Deficiency | 1 | 0 | 1 | 1.7 | 0.0 | 3.8 |
| 23 | In Situ, Benign and Unspecified Neoplasms | 1 | 1 | 0 | 1.7 | 3.1 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 5 | 2 | 3 | 8.4 | 6.2 | 11.4 |
| | All Other Causes | 95 | 67 | 28 | 159.0 | 208.2 | 106.5 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total* ----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|--------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 596 | 383 | 209 | 281 | 171 | 108 | 315 | 212 | 101 |
| Under 1 | 7 | 1 | 5 | 5 | 1 | 3 | 2 | 0 | 2 |
| (Neonatal) | 4 | 0 | 3 | 3 | 0 | 2 | 1 | 0 | 1 |
| (Postneonatal) | 3 | 1 | 2 | 2 | 1 | 1 | 1 | 0 | 1 |
| 1-4 | 2 | 0 | 2 | 1 | 0 | 1 | 1 | 0 | 1 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 3 | 2 | 1 | 1 | 1 | 0 | 2 | 1 | 1 |
| 15-24 | 5 | 2 | 3 | 3 | 1 | 2 | 2 | 1 | 1 |
| 25-34 | 14 | 7 | 7 | 12 | 6 | 6 | 2 | 1 | 1 |
| 35-44 | 23 | 11 | 11 | 12 | 5 | 6 | 11 | 6 | 5 |
| 45-54 | 50 | 21 | 29 | 32 | 15 | 17 | 18 | 6 | 12 |
| 55-64 | 69 | 34 | 35 | 32 | 15 | 17 | 37 | 19 | 18 |
| 65-74 | 113 | 76 | 36 | 55 | 38 | 17 | 58 | 38 | 19 |
| 75-84 | 148 | 103 | 44 | 83 | 56 | 27 | 65 | 47 | 17 |
| 85 and over | 162 | 126 | 36 | 45 | 33 | 12 | 117 | 93 | 24 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 59,730 |
| White | 32,175 |
| Black | 26,295 |
| Other | 1,260 |

Selected Death Statistics, Residents of Madison County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 964 | 675 | 286 | 948.0 | 1,142.9 | 734.6 |
| 1 | Malignant Neoplasms | 262 | 173 | 87 | 257.7 | 292.9 | 223.5 |
| 2 | Heart Diseases | 194 | 139 | 54 | 190.8 | 235.4 | 138.7 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 58 | 51 | 7 | 57.0 | 86.4 | 18.0 |
| 4 | Accidents | 42 | 26 | 16 | 41.3 | 44.0 | 41.1 |
| 4 | Cerebrovascular Disease | 42 | 26 | 16 | 41.3 | 44.0 | 41.1 |
| 6 | Alzheimer's Disease | 39 | 32 | 7 | 38.4 | 54.2 | 18.0 |
| 7 | Diabetes Mellitus | 34 | 14 | 20 | 33.4 | 23.7 | 51.4 |
| 8 | Influenza and Pneumonia | 24 | 15 | 9 | 23.6 | 25.4 | 23.1 |
| 9 | Nephritis, Nephrotic Syndrome And Nephrosis | 14 | 7 | 7 | 13.8 | 11.9 | 18.0 |
| 9 | Septicemia | 14 | 8 | 6 | 13.8 | 13.5 | 15.4 |
| 11 | Parkinson's Disease | 12 | 12 | 0 | 11.8 | 20.3 | 0.0 |
| 12 | Homicide | 10 | 3 | 7 | 9.8 | 5.1 | 18.0 |
| 12 | Suicide | 10 | 10 | 0 | 9.8 | 16.9 | 0.0 |
| 12 | Chronic Liver Disease And Cirrhosis | 10 | 8 | 2 | 9.8 | 13.5 | 5.1 |
| 15 | Viral Hepatitis | 7 | 5 | 2 | 6.9 | 8.5 | 5.1 |
| 16 | Hypertension/Hypertensive Renal Disease | 6 | 3 | 3 | 5.9 | 5.1 | 7.7 |
| 17 | Pneumonitis Due to Solids and Liquids | 5 | 4 | 1 | 4.9 | 6.8 | 2.6 |
| 17 | HIV Disease | 5 | 1 | 4 | 4.9 | 1.7 | 10.3 |
| 19 | Conditions Originating in the Perinatal Period** | 3 | 1 | 2 | 2.2 | 1.3 | 3.7 |
| 19 | Anemias | 3 | 2 | 1 | 3.0 | 3.4 | 2.6 |
| 19 | In Situ, Benign and Unspecified Neoplasms | 3 | 3 | 0 | 3.0 | 5.1 | 0.0 |
| 22 | Congenital Malformations and Chrom. Abnormalities | 2 | 1 | 1 | 2.0 | 1.7 | 2.6 |
| 23 | Acute Bronchitis and Bronchiolitis | 1 | 1 | 0 | 1.0 | 1.7 | 0.0 |
| 23 | Atherosclerosis | 1 | 1 | 0 | 1.0 | 1.7 | 0.0 |
| 23 | Malnutrition/Nutritional Deficiency | 1 | 1 | 0 | 1.0 | 1.7 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 6 | 5 | 1 | 5.9 | 8.5 | 2.6 |
| | All Other Causes | 156 | 123 | 33 | 153.4 | 208.3 | 84.8 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 964 | 675 | 286 | 452 | 307 | 144 | 512 | 368 | 142 |
| Under 1 | 9 | 3 | 6 | 6 | 2 | 4 | 3 | 1 | 2 |
| (Neonatal) | 3 | 1 | 2 | 2 | 1 | 1 | 1 | 0 | 1 |
| (Postneonatal) | 6 | 2 | 4 | 4 | 1 | 3 | 2 | 1 | 1 |
| 1-4 | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 15-24 | 13 | 5 | 8 | 12 | 4 | 8 | 1 | 1 | 0 |
| 25-34 | 12 | 3 | 9 | 6 | 3 | 3 | 6 | 0 | 6 |
| 35-44 | 27 | 12 | 14 | 10 | 5 | 5 | 17 | 7 | 9 |
| 45-54 | 57 | 30 | 27 | 36 | 22 | 14 | 21 | 8 | 13 |
| 55-64 | 134 | 77 | 57 | 85 | 48 | 37 | 49 | 29 | 20 |
| 65-74 | 156 | 94 | 61 | 88 | 57 | 30 | 68 | 37 | 31 |
| 75-84 | 237 | 179 | 58 | 104 | 80 | 24 | 133 | 99 | 34 |
| 85 and over | 316 | 270 | 45 | 103 | 85 | 18 | 213 | 185 | 27 |
| Unknown | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|---------|
| Total | 101,688 |
| White | 59,059 |
| Black | 38,934 |
| Other | 3,695 |

Selected Death Statistics, Residents of Marion County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 306 | 212 | 93 | 1,182.9 | 1,241.9 | 1,107.9 |
| 1 | Heart Diseases | 64 | 45 | 19 | 247.4 | 263.6 | 226.4 |
| 2 | Malignant Neoplasms | 57 | 37 | 19 | 220.3 | 216.7 | 226.4 |
| 3 | Accidents | 28 | 21 | 7 | 108.2 | 123.0 | 83.4 |
| 4 | Emphysema and Other Chronic Lower Resp. Diseases | 22 | 19 | 3 | 85.0 | 111.3 | 35.7 |
| 5 | Cerebrovascular Disease | 19 | 15 | 4 | 73.4 | 87.9 | 47.7 |
| 6 | Influenza and Pneumonia | 16 | 15 | 1 | 61.9 | 87.9 | 11.9 |
| 7 | Alzheimer's Disease | 11 | 8 | 3 | 42.5 | 46.9 | 35.7 |
| 8 | Nephritis, Nephrotic Syndrome And Nephrosis | 10 | 4 | 6 | 38.7 | 23.4 | 71.5 |
| 9 | Septicemia | 8 | 7 | 1 | 30.9 | 41.0 | 11.9 |
| 10 | Homicide | 5 | 3 | 2 | 19.3 | 17.6 | 23.8 |
| 11 | Chronic Liver Disease And Cirrhosis | 4 | 3 | 1 | 15.5 | 17.6 | 11.9 |
| 12 | Suicide | 3 | 2 | 1 | 11.6 | 11.7 | 11.9 |
| 12 | Diabetes Mellitus | 3 | 1 | 2 | 11.6 | 5.9 | 23.8 |
| 14 | Conditions Originating in the Perinatal Period** | 2 | 0 | 2 | 6.5 | 0.0 | 18.2 |
| 14 | Parkinson's Disease | 2 | 2 | 0 | 7.7 | 11.7 | 0.0 |
| 14 | In Situ, Benign and Unspecified Neoplasms | 2 | 2 | 0 | 7.7 | 11.7 | 0.0 |
| 14 | HIV Disease | 2 | 0 | 2 | 7.7 | 0.0 | 23.8 |
| 18 | Complications Medical/Surgical Care | 1 | 0 | 1 | 3.9 | 0.0 | 11.9 |
| 18 | Congenital Malformations and Chrom. Abnormalities | 1 | 1 | 0 | 3.9 | 5.9 | 0.0 |
| 18 | Hernia | 1 | 0 | 1 | 3.9 | 0.0 | 11.9 |
| 18 | Pneumonitis Due to Solids and Liquids | 1 | 0 | 1 | 3.9 | 0.0 | 11.9 |
| 18 | Aortic Aneurysm and Dissection | 1 | 0 | 1 | 3.9 | 0.0 | 11.9 |
| 18 | Hypertension/Hypertensive Renal Disease | 1 | 1 | 0 | 3.9 | 5.9 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 4 | 2 | 2 | 15.5 | 11.7 | 23.8 |
| | All Other Causes | 38 | 24 | 14 | 146.9 | 140.6 | 166.8 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 306 | 212 | 93 | 168 | 111 | 57 | 138 | 101 | 36 |
| Under 1 | 4 | 2 | 2 | 3 | 1 | 2 | 1 | 1 | 0 |
| (Neonatal) | 3 | 1 | 2 | 2 | 0 | 2 | 1 | 1 | 0 |
| (Postneonatal) | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 1-4 | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 |
| 5-9 | 2 | 1 | 1 | 1 | 1 | 0 | 1 | 0 | 1 |
| 10-14 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 15-24 | 8 | 5 | 3 | 7 | 5 | 2 | 1 | 0 | 1 |
| 25-34 | 9 | 5 | 4 | 5 | 4 | 1 | 4 | 1 | 3 |
| 35-44 | 11 | 7 | 4 | 7 | 5 | 2 | 4 | 2 | 2 |
| 45-54 | 22 | 12 | 10 | 15 | 9 | 6 | 7 | 3 | 4 |
| 55-64 | 60 | 41 | 19 | 37 | 22 | 15 | 23 | 19 | 4 |
| 65-74 | 67 | 44 | 22 | 45 | 29 | 16 | 22 | 15 | 6 |
| 75-84 | 59 | 46 | 13 | 28 | 20 | 8 | 31 | 26 | 5 |
| 85 and over | 62 | 48 | 14 | 19 | 15 | 4 | 43 | 33 | 10 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 25,868 |
| White | 17,071 |
| Black | 8,394 |
| Other | 403 |

Selected Death Statistics, Residents of Marshall County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 410 | 255 | 153 | 1,131.5 | 1,383.9 | 887.2 |
| 1 | Malignant Neoplasms | 95 | 58 | 37 | 262.2 | 314.8 | 214.5 |
| 2 | Heart Diseases | 69 | 46 | 23 | 190.4 | 249.6 | 133.4 |
| 3 | Diabetes Mellitus | 38 | 20 | 18 | 104.9 | 108.5 | 104.4 |
| 4 | Emphysema and Other Chronic Lower Resp. Diseases | 31 | 29 | 2 | 85.6 | 157.4 | 11.6 |
| 5 | Accidents | 26 | 12 | 14 | 71.8 | 65.1 | 81.2 |
| 5 | Hypertension/Hypertensive Renal Disease | 26 | 15 | 11 | 71.8 | 81.4 | 63.8 |
| 7 | Cerebrovascular Disease | 17 | 13 | 4 | 46.9 | 70.6 | 23.2 |
| 8 | Alzheimer's Disease | 13 | 7 | 5 | 35.9 | 38.0 | 29.0 |
| 9 | Nephritis, Nephrotic Syndrome And Nephrosis | 8 | 3 | 5 | 22.1 | 16.3 | 29.0 |
| 10 | Influenza and Pneumonia | 7 | 4 | 3 | 19.3 | 21.7 | 17.4 |
| 10 | Septicemia | 7 | 4 | 3 | 19.3 | 21.7 | 17.4 |
| 12 | Homicide | 5 | 1 | 4 | 13.8 | 5.4 | 23.2 |
| 12 | HIV Disease | 5 | 2 | 3 | 13.8 | 10.9 | 17.4 |
| 14 | Suicide | 4 | 3 | 1 | 11.0 | 16.3 | 5.8 |
| 14 | Congenital Malformations and Chrom. Abnormalities | 4 | 2 | 2 | 11.0 | 10.9 | 11.6 |
| 14 | Chronic Liver Disease And Cirrhosis | 4 | 2 | 2 | 11.0 | 10.9 | 11.6 |
| 17 | Parkinson's Disease | 3 | 3 | 0 | 8.3 | 16.3 | 0.0 |
| 18 | Pneumonitis Due to Solids and Liquids | 2 | 2 | 0 | 5.5 | 10.9 | 0.0 |
| 19 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 2.2 | 0.0 | 4.6 |
| 19 | Hernia | 1 | 0 | 1 | 2.8 | 0.0 | 5.8 |
| 19 | Peptic Ulcer | 1 | 1 | 0 | 2.8 | 5.4 | 0.0 |
| 19 | Malnutrition/Nutritional Deficiency | 1 | 1 | 0 | 2.8 | 5.4 | 0.0 |
| 19 | Anemias | 1 | 1 | 0 | 2.8 | 5.4 | 0.0 |
| 19 | In Situ, Benign and Unspecified Neoplasms | 1 | 1 | 0 | 2.8 | 5.4 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 4 | 3 | 1 | 11.0 | 16.3 | 5.8 |
| | All Other Causes | 36 | 22 | 13 | 99.4 | 119.4 | 75.4 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 410 | 255 | 153 | 251 | 154 | 97 | 159 | 101 | 56 |
| Under 1 | 4 | 0 | 4 | 1 | 0 | 1 | 3 | 0 | 3 |
| (Neonatal) | 3 | 0 | 3 | 1 | 0 | 1 | 2 | 0 | 2 |
| (Postneonatal) | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| 1-4 | 3 | 0 | 3 | 3 | 0 | 3 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 6 | 1 | 5 | 3 | 1 | 2 | 3 | 0 | 3 |
| 25-34 | 15 | 6 | 9 | 12 | 5 | 7 | 3 | 1 | 2 |
| 35-44 | 16 | 7 | 9 | 13 | 5 | 8 | 3 | 2 | 1 |
| 45-54 | 37 | 23 | 13 | 22 | 16 | 6 | 15 | 7 | 7 |
| 55-64 | 78 | 41 | 37 | 56 | 28 | 28 | 22 | 13 | 9 |
| 65-74 | 97 | 68 | 28 | 60 | 43 | 17 | 37 | 25 | 11 |
| 75-84 | 85 | 59 | 26 | 51 | 34 | 17 | 34 | 25 | 9 |
| 85 and over | 69 | 50 | 19 | 30 | 22 | 8 | 39 | 28 | 11 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 36,234 |
| White | 18,426 |
| Black | 17,246 |
| Other | 562 |

Selected Death Statistics, Residents of Monroe County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 427 | 326 | 100 | 1,186.0 | 1,335.0 | 902.0 |
| 1 | Heart Diseases | 103 | 74 | 29 | 286.1 | 303.0 | 261.6 |
| 2 | Malignant Neoplasms | 98 | 74 | 24 | 272.2 | 303.0 | 216.5 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 26 | 24 | 2 | 72.2 | 98.3 | 18.0 |
| 3 | Cerebrovascular Disease | 26 | 21 | 5 | 72.2 | 86.0 | 45.1 |
| 5 | Accidents | 21 | 16 | 5 | 58.3 | 65.5 | 45.1 |
| 6 | Alzheimer's Disease | 19 | 16 | 3 | 52.8 | 65.5 | 27.1 |
| 7 | Nephritis, Nephrotic Syndrome And Nephrosis | 12 | 6 | 6 | 33.3 | 24.6 | 54.1 |
| 8 | Diabetes Mellitus | 11 | 9 | 1 | 30.6 | 36.9 | 9.0 |
| 9 | Hypertension/Hypertensive Renal Disease | 10 | 5 | 5 | 27.8 | 20.5 | 45.1 |
| 10 | Influenza and Pneumonia | 9 | 9 | 0 | 25.0 | 36.9 | 0.0 |
| 11 | Homicide | 7 | 4 | 3 | 19.4 | 16.4 | 27.1 |
| 11 | Septicemia | 7 | 4 | 3 | 19.4 | 16.4 | 27.1 |
| 13 | Pneumonitis Due to Solids and Liquids | 6 | 6 | 0 | 16.7 | 24.6 | 0.0 |
| 14 | Chronic Liver Disease And Cirrhosis | 5 | 5 | 0 | 13.9 | 20.5 | 0.0 |
| 15 | Suicide | 4 | 4 | 0 | 11.1 | 16.4 | 0.0 |
| 16 | Congenital Malformations and Chrom. Abnormalities | 3 | 2 | 1 | 8.3 | 8.2 | 9.0 |
| 17 | In Situ, Benign and Unspecified Neoplasms | 2 | 1 | 1 | 5.6 | 4.1 | 9.0 |
| 18 | Complications Medical/Surgical Care | 1 | 0 | 1 | 2.8 | 0.0 | 9.0 |
| 18 | Infections of Kidney | 1 | 1 | 0 | 2.8 | 4.1 | 0.0 |
| 18 | Aortic Aneurysm and Dissection | 1 | 1 | 0 | 2.8 | 4.1 | 0.0 |
| 18 | Malnutrition/Nutritional Deficiency | 1 | 1 | 0 | 2.8 | 4.1 | 0.0 |
| 18 | Anemias | 1 | 1 | 0 | 2.8 | 4.1 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 4 | 3 | 1 | 11.1 | 12.3 | 9.0 |
| | All Other Causes | 49 | 39 | 10 | 136.1 | 159.7 | 90.2 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 427 | 326 | 100 | 212 | 156 | 56 | 215 | 170 | 44 |
| Under 1 | 4 | 3 | 1 | 3 | 2 | 1 | 1 | 1 | 0 |
| (Neonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 4 | 3 | 1 | 3 | 2 | 1 | 1 | 1 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 7 | 4 | 3 | 5 | 3 | 2 | 2 | 1 | 1 |
| 25-34 | 5 | 3 | 2 | 4 | 3 | 1 | 1 | 0 | 1 |
| 35-44 | 6 | 5 | 1 | 5 | 4 | 1 | 1 | 1 | 0 |
| 45-54 | 34 | 22 | 12 | 17 | 10 | 7 | 17 | 12 | 5 |
| 55-64 | 61 | 36 | 25 | 42 | 23 | 19 | 19 | 13 | 6 |
| 65-74 | 91 | 67 | 24 | 56 | 39 | 17 | 35 | 28 | 7 |
| 75-84 | 113 | 99 | 13 | 50 | 45 | 5 | 63 | 54 | 8 |
| 85 and over | 105 | 86 | 19 | 30 | 27 | 3 | 75 | 59 | 16 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 36,003 |
| White | 24,419 |
| Black | 11,087 |
| Other | 497 |

Selected Death Statistics, Residents of Montgomery County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 135 | 80 | 55 | 1,298.1 | 1,444.8 | 1,162.8 |
| 1 | Heart Diseases | 32 | 16 | 16 | 307.7 | 289.0 | 338.3 |
| 2 | Malignant Neoplasms | 21 | 15 | 6 | 201.9 | 270.9 | 126.8 |
| 3 | Accidents | 11 | 7 | 4 | 105.8 | 126.4 | 84.6 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 11 | 9 | 2 | 105.8 | 162.5 | 42.3 |
| 5 | Cerebrovascular Disease | 8 | 5 | 3 | 76.9 | 90.3 | 63.4 |
| 6 | Homicide | 5 | 2 | 3 | 48.1 | 36.1 | 63.4 |
| 6 | Nephritis, Nephrotic Syndrome And Nephrosis | 5 | 3 | 2 | 48.1 | 54.2 | 42.3 |
| 8 | Diabetes Mellitus | 4 | 2 | 2 | 38.5 | 36.1 | 42.3 |
| 9 | Suicide | 3 | 3 | 0 | 28.8 | 54.2 | 0.0 |
| 9 | Influenza and Pneumonia | 3 | 2 | 1 | 28.8 | 36.1 | 21.1 |
| 9 | Hypertension/Hypertensive Renal Disease | 3 | 1 | 2 | 28.8 | 18.1 | 42.3 |
| 9 | Alzheimer's Disease | 3 | 3 | 0 | 28.8 | 54.2 | 0.0 |
| 13 | Septicemia | 2 | 1 | 1 | 19.2 | 18.1 | 21.1 |
| 14 | Chronic Liver Disease And Cirrhosis | 1 | 0 | 1 | 9.6 | 0.0 | 21.1 |
| 14 | Pneumonitis Due to Solids and Liquids | 1 | 1 | 0 | 9.6 | 18.1 | 0.0 |
| 14 | Parkinson's Disease | 1 | 1 | 0 | 9.6 | 18.1 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 6 | 2 | 4 | 57.7 | 36.1 | 84.6 |
| | All Other Causes | 15 | 7 | 8 | 144.2 | 126.4 | 169.1 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 135 | 80 | 55 | 69 | 40 | 29 | 66 | 40 | 26 |
| Under 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Neonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 4 | 1 | 3 | 2 | 0 | 2 | 2 | 1 | 1 |
| 25-34 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 35-44 | 6 | 2 | 4 | 3 | 1 | 2 | 3 | 1 | 2 |
| 45-54 | 12 | 7 | 5 | 5 | 3 | 2 | 7 | 4 | 3 |
| 55-64 | 28 | 15 | 13 | 18 | 10 | 8 | 10 | 5 | 5 |
| 65-74 | 21 | 12 | 9 | 12 | 5 | 7 | 9 | 7 | 2 |
| 75-84 | 28 | 15 | 13 | 14 | 9 | 5 | 14 | 6 | 8 |
| 85 and over | 35 | 27 | 8 | 14 | 11 | 3 | 21 | 16 | 5 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 10,400 |
| White | 5,537 |
| Black | 4,730 |
| Other | 133 |

Selected Death Statistics, Residents of Neshoba County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 370 | 254 | 60 | 1,255.7 | 1,428.0 | 961.2 |
| 1 | Heart Diseases | 118 | 88 | 14 | 400.5 | 494.7 | 224.3 |
| 2 | Malignant Neoplasms | 66 | 48 | 12 | 224.0 | 269.9 | 192.2 |
| 3 | Cerebrovascular Disease | 21 | 12 | 6 | 71.3 | 67.5 | 96.1 |
| 4 | Accidents | 18 | 12 | 4 | 61.1 | 67.5 | 64.1 |
| 5 | Emphysema and Other Chronic Lower Resp. Diseases | 14 | 13 | 0 | 47.5 | 73.1 | 0.0 |
| 5 | Alzheimer's Disease | 14 | 11 | 2 | 47.5 | 61.8 | 32.0 |
| 7 | Influenza and Pneumonia | 10 | 9 | 0 | 33.9 | 50.6 | 0.0 |
| 8 | Septicemia | 9 | 2 | 1 | 30.5 | 11.2 | 16.0 |
| 9 | Nephritis, Nephrotic Syndrome And Nephrosis | 8 | 2 | 3 | 27.2 | 11.2 | 48.1 |
| 10 | Chronic Liver Disease And Cirrhosis | 6 | 1 | 2 | 20.4 | 5.6 | 32.0 |
| 11 | Diabetes Mellitus | 5 | 1 | 0 | 17.0 | 5.6 | 0.0 |
| 11 | Homicide | 5 | 0 | 4 | 17.0 | 0.0 | 64.1 |
| 11 | Suicide | 5 | 3 | 1 | 17.0 | 16.9 | 16.0 |
| 14 | In Situ, Benign and Unspecified Neoplasms | 3 | 3 | 0 | 10.2 | 16.9 | 0.0 |
| 15 | Parkinson's Disease | 2 | 1 | 1 | 6.8 | 5.6 | 16.0 |
| 15 | Malnutrition/Nutritional Deficiency | 2 | 2 | 0 | 6.8 | 11.2 | 0.0 |
| 15 | Congenital Malformations and Chrom. Abnormalities | 2 | 2 | 0 | 6.8 | 11.2 | 0.0 |
| 18 | HIV Disease | 1 | 0 | 0 | 3.4 | 0.0 | 0.0 |
| 18 | Legal Intervention | 1 | 0 | 0 | 3.4 | 0.0 | 0.0 |
| 18 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 2.4 | 0.0 | 10.1 |
| 18 | Female Pelvic Inflammatory Disease | 1 | 0 | 0 | 6.5 | 0.0 | 0.0 |
| 18 | Cholelithiasis/Gallbladder Disorder | 1 | 1 | 0 | 3.4 | 5.6 | 0.0 |
| 18 | Pneumonitis Due to Solids and Liquids | 1 | 1 | 0 | 3.4 | 5.6 | 0.0 |
| 18 | Pneumoconioses and Chemical Effects | 1 | 1 | 0 | 3.4 | 5.6 | 0.0 |
| 18 | Aortic Aneurysm and Dissection | 1 | 1 | 0 | 3.4 | 5.6 | 0.0 |
| 18 | Hypertension/Hypertensive Renal Disease | 1 | 0 | 1 | 3.4 | 0.0 | 16.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 17 | 13 | 4 | 57.7 | 73.1 | 64.1 |
| | All Other Causes | 36 | 27 | 4 | 122.2 | 151.8 | 64.1 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 370 | 254 | 60 | 191 | 122 | 35 | 179 | 132 | 25 |
| Under 1 | 4 | 3 | 1 | 4 | 3 | 1 | 0 | 0 | 0 |
| (Neonatal) | 3 | 2 | 1 | 3 | 2 | 1 | 0 | 0 | 0 |
| (Postneonatal) | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 1-4 | 2 | 2 | 0 | 2 | 2 | 0 | 0 | 0 | 0 |
| 5-9 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 2 | 1 | 1 | 2 | 1 | 1 | 0 | 0 | 0 |
| 25-34 | 12 | 4 | 5 | 7 | 2 | 4 | 5 | 2 | 1 |
| 35-44 | 12 | 4 | 0 | 10 | 2 | 0 | 2 | 2 | 0 |
| 45-54 | 40 | 19 | 11 | 20 | 8 | 6 | 20 | 11 | 5 |
| 55-64 | 69 | 39 | 16 | 45 | 24 | 10 | 24 | 15 | 6 |
| 65-74 | 61 | 43 | 8 | 34 | 22 | 7 | 27 | 21 | 1 |
| 75-84 | 84 | 70 | 8 | 34 | 29 | 3 | 50 | 41 | 5 |
| 85 and over | 83 | 68 | 10 | 32 | 28 | 3 | 51 | 40 | 7 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 29,465 |
| White | 17,787 |
| Black | 6,242 |
| Other | 5,436 |

Selected Death Statistics, Residents of Newton County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 285 | 208 | 68 | 1,305.4 | 1,501.9 | 1,027.0 |
| 1 | Heart Diseases | 58 | 43 | 13 | 265.7 | 310.5 | 196.3 |
| 2 | Malignant Neoplasms | 47 | 35 | 10 | 215.3 | 252.7 | 151.0 |
| 3 | Accidents | 20 | 13 | 4 | 91.6 | 93.9 | 60.4 |
| 4 | Diabetes Mellitus | 16 | 10 | 6 | 73.3 | 72.2 | 90.6 |
| 5 | Emphysema and Other Chronic Lower Resp. Diseases | 15 | 13 | 2 | 68.7 | 93.9 | 30.2 |
| 5 | Influenza and Pneumonia | 15 | 11 | 4 | 68.7 | 79.4 | 60.4 |
| 5 | Cerebrovascular Disease | 15 | 10 | 4 | 68.7 | 72.2 | 60.4 |
| 8 | Alzheimer's Disease | 13 | 11 | 2 | 59.5 | 79.4 | 30.2 |
| 9 | Hypertension/Hypertensive Renal Disease | 12 | 10 | 2 | 55.0 | 72.2 | 30.2 |
| 10 | Nephritis, Nephrotic Syndrome And Nephrosis | 9 | 4 | 5 | 41.2 | 28.9 | 75.5 |
| 11 | Suicide | 5 | 3 | 2 | 22.9 | 21.7 | 30.2 |
| 11 | Chronic Liver Disease And Cirrhosis | 5 | 4 | 1 | 22.9 | 28.9 | 15.1 |
| 13 | Pneumonitis Due to Solids and Liquids | 4 | 2 | 2 | 18.3 | 14.4 | 30.2 |
| 13 | Septicemia | 4 | 2 | 1 | 18.3 | 14.4 | 15.1 |
| 15 | Homicide | 3 | 1 | 2 | 13.7 | 7.2 | 30.2 |
| 15 | Conditions Originating in the Perinatal Period** | 3 | 2 | 1 | 10.5 | 13.1 | 9.3 |
| 15 | In Situ, Benign and Unspecified Neoplasms | 3 | 2 | 1 | 13.7 | 14.4 | 15.1 |
| 18 | Anemias | 2 | 2 | 0 | 9.2 | 14.4 | 0.0 |
| 19 | Congenital Malformations and Chrom. Abnormalities | 1 | 0 | 1 | 4.6 | 0.0 | 15.1 |
| 19 | Aortic Aneurysm and Dissection | 1 | 1 | 0 | 4.6 | 7.2 | 0.0 |
| 19 | Parkinson's Disease | 1 | 1 | 0 | 4.6 | 7.2 | 0.0 |
| 19 | HIV Disease | 1 | 1 | 0 | 4.6 | 7.2 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 4 | 3 | 1 | 18.3 | 21.7 | 15.1 |
| | All Other Causes | 28 | 24 | 4 | 128.3 | 173.3 | 60.4 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 285 | 208 | 68 | 141 | 102 | 35 | 144 | 106 | 33 |
| Under 1 | 6 | 3 | 2 | 4 | 3 | 1 | 2 | 0 | 1 |
| (Neonatal) | 4 | 2 | 2 | 3 | 2 | 1 | 1 | 0 | 1 |
| (Postneonatal) | 2 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 |
| 1-4 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 2 | 2 | 0 | 1 | 1 | 0 | 1 | 1 | 0 |
| 25-34 | 10 | 2 | 7 | 8 | 2 | 6 | 2 | 0 | 1 |
| 35-44 | 13 | 8 | 3 | 9 | 4 | 3 | 4 | 4 | 0 |
| 45-54 | 18 | 11 | 5 | 9 | 6 | 1 | 9 | 5 | 4 |
| 55-64 | 45 | 29 | 15 | 27 | 18 | 9 | 18 | 11 | 6 |
| 65-74 | 57 | 44 | 12 | 36 | 28 | 8 | 21 | 16 | 4 |
| 75-84 | 61 | 53 | 8 | 25 | 20 | 5 | 36 | 33 | 3 |
| 85 and over | 72 | 56 | 16 | 22 | 20 | 2 | 50 | 36 | 14 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 21,832 |
| White | 13,849 |
| Black | 6,621 |
| Other | 1,362 |

Selected Death Statistics, Residents of Noxubee County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 110 | 31 | 79 | 989.7 | 1,035.4 | 991.3 |
| 1 | Heart Diseases | 33 | 8 | 25 | 296.9 | 267.2 | 313.7 |
| 2 | Malignant Neoplasms | 27 | 8 | 19 | 242.9 | 267.2 | 238.4 |
| 3 | Cerebrovascular Disease | 7 | 2 | 5 | 63.0 | 66.8 | 62.7 |
| 4 | Accidents | 5 | 1 | 4 | 45.0 | 33.4 | 50.2 |
| 4 | Alzheimer's Disease | 5 | 3 | 2 | 45.0 | 100.2 | 25.1 |
| 6 | Nephritis, Nephrotic Syndrome And Nephrosis | 4 | 0 | 4 | 36.0 | 0.0 | 50.2 |
| 6 | Emphysema and Other Chronic Lower Resp. Diseases | 4 | 2 | 2 | 36.0 | 66.8 | 25.1 |
| 8 | Conditions Originating in the Perinatal Period** | 2 | 1 | 1 | 11.5 | 20.0 | 8.2 |
| 8 | Influenza and Pneumonia | 2 | 0 | 2 | 18.0 | 0.0 | 25.1 |
| 8 | Hypertension/Hypertensive Renal Disease | 2 | 0 | 2 | 18.0 | 0.0 | 25.1 |
| 11 | Chronic Liver Disease And Cirrhosis | 1 | 0 | 1 | 9.0 | 0.0 | 12.5 |
| 11 | Aortic Aneurysm and Dissection | 1 | 1 | 0 | 9.0 | 33.4 | 0.0 |
| 11 | Parkinson's Disease | 1 | 1 | 0 | 9.0 | 33.4 | 0.0 |
| 11 | Diabetes Mellitus | 1 | 0 | 1 | 9.0 | 0.0 | 12.5 |
| 11 | HIV Disease | 1 | 1 | 0 | 9.0 | 33.4 | 0.0 |
| 11 | Septicemia | 1 | 1 | 0 | 9.0 | 33.4 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 3 | 1 | 2 | 27.0 | 33.4 | 25.1 |
| | All Other Causes | 10 | 1 | 9 | 90.0 | 33.4 | 112.9 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 110 | 31 | 79 | 63 | 19 | 44 | 47 | 12 | 35 |
| Under 1 | 2 | 1 | 1 | 1 | 1 | 0 | 1 | 0 | 1 |
| (Neonatal) | 2 | 1 | 1 | 1 | 1 | 0 | 1 | 0 | 1 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| 25-34 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 35-44 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| 45-54 | 6 | 1 | 5 | 5 | 0 | 5 | 1 | 1 | 0 |
| 55-64 | 15 | 2 | 13 | 11 | 2 | 9 | 4 | 0 | 4 |
| 65-74 | 20 | 4 | 16 | 15 | 3 | 12 | 5 | 1 | 4 |
| 75-84 | 32 | 8 | 24 | 16 | 5 | 11 | 16 | 3 | 13 |
| 85 and over | 32 | 15 | 17 | 14 | 8 | 6 | 18 | 7 | 11 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 11,115 |
| White | 2,994 |
| Black | 7,969 |
| Other | 152 |

Selected Death Statistics, Residents of Oktibbeha County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|-------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 303 | 165 | 138 | 613.2 | 573.9 | 751.6 |
| 1 | Malignant Neoplasms | 72 | 45 | 27 | 145.7 | 156.5 | 147.1 |
| 2 | Heart Diseases | 67 | 35 | 32 | 135.6 | 121.7 | 174.3 |
| 3 | Diabetes Mellitus | 18 | 8 | 10 | 36.4 | 27.8 | 54.5 |
| 4 | Cerebrovascular Disease | 14 | 8 | 6 | 28.3 | 27.8 | 32.7 |
| 5 | Emphysema and Other Chronic Lower Resp. Diseases | 12 | 9 | 3 | 24.3 | 31.3 | 16.3 |
| 6 | Accidents | 11 | 3 | 8 | 22.3 | 10.4 | 43.6 |
| 6 | Nephritis, Nephrotic Syndrome And Nephrosis | 11 | 4 | 7 | 22.3 | 13.9 | 38.1 |
| 8 | Chronic Liver Disease And Cirrhosis | 7 | 6 | 1 | 14.2 | 20.9 | 5.4 |
| 8 | Alzheimer's Disease | 7 | 3 | 4 | 14.2 | 10.4 | 21.8 |
| 8 | Septicemia | 7 | 2 | 5 | 14.2 | 7.0 | 27.2 |
| 11 | Influenza and Pneumonia | 5 | 3 | 2 | 10.1 | 10.4 | 10.9 |
| 12 | Homicide | 4 | 1 | 3 | 8.1 | 3.5 | 16.3 |
| 12 | Hypertension/Hypertensive Renal Disease | 4 | 1 | 3 | 8.1 | 3.5 | 16.3 |
| 14 | Parkinson's Disease | 3 | 3 | 0 | 6.1 | 10.4 | 0.0 |
| 15 | Suicide | 2 | 2 | 0 | 4.0 | 7.0 | 0.0 |
| 15 | Conditions Originating in the Perinatal Period** | 2 | 0 | 2 | 3.7 | 0.0 | 8.2 |
| 15 | Cholelithiasis/Gallbladder Disorder | 2 | 2 | 0 | 4.0 | 7.0 | 0.0 |
| 15 | Pneumonitis Due to Solids and Liquids | 2 | 1 | 1 | 4.0 | 3.5 | 5.4 |
| 15 | Aortic Aneurysm and Dissection | 2 | 1 | 1 | 4.0 | 3.5 | 5.4 |
| 15 | In Situ, Benign and Unspecified Neoplasms | 2 | 1 | 1 | 4.0 | 3.5 | 5.4 |
| 21 | Congenital Malformations and Chrom. Abnormalities | 1 | 1 | 0 | 2.0 | 3.5 | 0.0 |
| 21 | Hernia | 1 | 1 | 0 | 2.0 | 3.5 | 0.0 |
| 21 | Anemias | 1 | 0 | 1 | 2.0 | 0.0 | 5.4 |
| 21 | HIV Disease | 1 | 0 | 1 | 2.0 | 0.0 | 5.4 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 0 | 0 | 0 | 0.0 | 0.0 | 0.0 |
| | All Other Causes | 45 | 25 | 20 | 91.1 | 87.0 | 108.9 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total* ----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|--------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 303 | 165 | 138 | 155 | 82 | 73 | 148 | 83 | 65 |
| Under 1 | 4 | 1 | 3 | 3 | 1 | 2 | 1 | 0 | 1 |
| (Neonatal) | 3 | 0 | 3 | 2 | 0 | 2 | 1 | 0 | 1 |
| (Postneonatal) | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 2 | 1 | 1 | 1 | 1 | 0 | 1 | 0 | 1 |
| 25-34 | 5 | 2 | 3 | 2 | 1 | 1 | 3 | 1 | 2 |
| 35-44 | 7 | 2 | 5 | 6 | 1 | 5 | 1 | 1 | 0 |
| 45-54 | 31 | 12 | 19 | 21 | 10 | 11 | 10 | 2 | 8 |
| 55-64 | 55 | 24 | 31 | 30 | 11 | 19 | 25 | 13 | 12 |
| 65-74 | 63 | 40 | 23 | 34 | 21 | 13 | 29 | 19 | 10 |
| 75-84 | 64 | 32 | 32 | 31 | 13 | 18 | 33 | 19 | 14 |
| 85 and over | 72 | 51 | 21 | 27 | 23 | 4 | 45 | 28 | 17 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 49,414 |
| White | 28,751 |
| Black | 18,361 |
| Other | 2,302 |

Selected Death Statistics, Residents of Panola County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 411 | 257 | 154 | 1,193.2 | 1,531.9 | 899.3 |
| 1 | Heart Diseases | 136 | 89 | 47 | 394.8 | 530.5 | 274.5 |
| 2 | Malignant Neoplasms | 90 | 53 | 37 | 261.3 | 315.9 | 216.1 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 29 | 21 | 8 | 84.2 | 125.2 | 46.7 |
| 4 | Accidents | 22 | 16 | 6 | 63.9 | 95.4 | 35.0 |
| 5 | Cerebrovascular Disease | 18 | 12 | 6 | 52.3 | 71.5 | 35.0 |
| 6 | Alzheimer's Disease | 13 | 9 | 4 | 37.7 | 53.6 | 23.4 |
| 7 | Influenza and Pneumonia | 11 | 8 | 3 | 31.9 | 47.7 | 17.5 |
| 8 | Nephritis, Nephrotic Syndrome And Nephrosis | 9 | 2 | 7 | 26.1 | 11.9 | 40.9 |
| 9 | Diabetes Mellitus | 8 | 3 | 5 | 23.2 | 17.9 | 29.2 |
| 10 | Homicide | 6 | 2 | 4 | 17.4 | 11.9 | 23.4 |
| 10 | Hypertension/Hypertensive Renal Disease | 6 | 2 | 4 | 17.4 | 11.9 | 23.4 |
| 12 | Suicide | 5 | 4 | 1 | 14.5 | 23.8 | 5.8 |
| 13 | Pneumonitis Due to Solids and Liquids | 4 | 4 | 0 | 11.6 | 23.8 | 0.0 |
| 13 | Septicemia | 4 | 1 | 3 | 11.6 | 6.0 | 17.5 |
| 15 | Chronic Liver Disease And Cirrhosis | 2 | 2 | 0 | 5.8 | 11.9 | 0.0 |
| 15 | Viral Hepatitis | 2 | 1 | 1 | 5.8 | 6.0 | 5.8 |
| 17 | Conditions Originating in the Perinatal Period** | 1 | 1 | 0 | 1.9 | 4.6 | 0.0 |
| 17 | Aortic Aneurysm and Dissection | 1 | 1 | 0 | 2.9 | 6.0 | 0.0 |
| 17 | Parkinson's Disease | 1 | 1 | 0 | 2.9 | 6.0 | 0.0 |
| 17 | In Situ, Benign and Unspecified Neoplasms | 1 | 1 | 0 | 2.9 | 6.0 | 0.0 |
| 17 | HIV Disease | 1 | 0 | 1 | 2.9 | 0.0 | 5.8 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 1 | 0 | 1 | 2.9 | 0.0 | 5.8 |
| | All Other Causes | 40 | 24 | 16 | 116.1 | 143.1 | 93.4 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total* ----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|--------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 411 | 257 | 154 | 223 | 143 | 80 | 188 | 114 | 74 |
| Under 1 | 2 | 1 | 1 | 1 | 0 | 1 | 1 | 1 | 0 |
| (Neonatal) | 2 | 1 | 1 | 1 | 0 | 1 | 1 | 1 | 0 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 4 | 2 | 2 | 3 | 1 | 2 | 1 | 1 | 0 |
| 25-34 | 10 | 2 | 8 | 7 | 2 | 5 | 3 | 0 | 3 |
| 35-44 | 14 | 6 | 8 | 6 | 3 | 3 | 8 | 3 | 5 |
| 45-54 | 33 | 23 | 10 | 17 | 12 | 5 | 16 | 11 | 5 |
| 55-64 | 85 | 48 | 37 | 52 | 32 | 20 | 33 | 16 | 17 |
| 65-74 | 73 | 51 | 22 | 42 | 31 | 11 | 31 | 20 | 11 |
| 75-84 | 109 | 68 | 41 | 65 | 39 | 26 | 44 | 29 | 15 |
| 85 and over | 81 | 56 | 25 | 30 | 23 | 7 | 51 | 33 | 18 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 34,444 |
| White | 16,777 |
| Black | 17,124 |
| Other | 543 |

Selected Death Statistics, Residents of Pearl River County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 604 | 535 | 69 | 1,093.7 | 1,148.4 | 977.5 |
| 1 | Malignant Neoplasms | 146 | 134 | 12 | 264.4 | 287.6 | 170.0 |
| 2 | Heart Diseases | 134 | 116 | 18 | 242.6 | 249.0 | 255.0 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 55 | 49 | 6 | 99.6 | 105.2 | 85.0 |
| 4 | Accidents | 40 | 30 | 10 | 72.4 | 64.4 | 141.7 |
| 5 | Diabetes Mellitus | 26 | 20 | 6 | 47.1 | 42.9 | 85.0 |
| 6 | Cerebrovascular Disease | 21 | 18 | 3 | 38.0 | 38.6 | 42.5 |
| 7 | Nephritis, Nephrotic Syndrome And Nephrosis | 19 | 19 | 0 | 34.4 | 40.8 | 0.0 |
| 7 | Septicemia | 19 | 16 | 3 | 34.4 | 34.3 | 42.5 |
| 9 | Alzheimer's Disease | 16 | 13 | 3 | 29.0 | 27.9 | 42.5 |
| 10 | Suicide | 11 | 11 | 0 | 19.9 | 23.6 | 0.0 |
| 10 | Influenza and Pneumonia | 11 | 11 | 0 | 19.9 | 23.6 | 0.0 |
| 12 | Chronic Liver Disease And Cirrhosis | 8 | 6 | 2 | 14.5 | 12.9 | 28.3 |
| 13 | Viral Hepatitis | 5 | 5 | 0 | 9.1 | 10.7 | 0.0 |
| 14 | Parkinson's Disease | 4 | 4 | 0 | 7.2 | 8.6 | 0.0 |
| 15 | Conditions Originating in the Perinatal Period** | 3 | 3 | 0 | 4.8 | 5.6 | 0.0 |
| 15 | Pneumonitis Due to Solids and Liquids | 3 | 2 | 1 | 5.4 | 4.3 | 14.2 |
| 15 | Aortic Aneurysm and Dissection | 3 | 3 | 0 | 5.4 | 6.4 | 0.0 |
| 15 | Hypertension/Hypertensive Renal Disease | 3 | 3 | 0 | 5.4 | 6.4 | 0.0 |
| 19 | In Situ, Benign and Unspecified Neoplasms | 2 | 2 | 0 | 3.6 | 4.3 | 0.0 |
| 20 | Complications Medical/Surgical Care | 1 | 1 | 0 | 1.8 | 2.1 | 0.0 |
| 20 | Homicide | 1 | 1 | 0 | 1.8 | 2.1 | 0.0 |
| 20 | Congenital Malformations and Chrom. Abnormalities | 1 | 1 | 0 | 1.8 | 2.1 | 0.0 |
| 20 | Maternal Causes** | 1 | 1 | 0 | 158.7 | 185.5 | 0.0 |
| 20 | Cholelithiasis/Gallbladder Disorder | 1 | 1 | 0 | 1.8 | 2.1 | 0.0 |
| 20 | Hernia | 1 | 0 | 1 | 1.8 | 0.0 | 14.2 |
| 20 | Pneumoconioses and Chemical Effects | 1 | 1 | 0 | 1.8 | 2.1 | 0.0 |
| 20 | Anemias | 1 | 1 | 0 | 1.8 | 2.1 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 8 | 7 | 1 | 14.5 | 15.0 | 14.2 |
| | All Other Causes | 59 | 56 | 3 | 106.8 | 120.2 | 42.5 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 604 | 535 | 69 | 313 | 280 | 33 | 291 | 255 | 36 |
| Under 1 | 5 | 3 | 2 | 2 | 1 | 1 | 3 | 2 | 1 |
| (Neonatal) | 3 | 3 | 0 | 1 | 1 | 0 | 2 | 2 | 0 |
| (Postneonatal) | 2 | 0 | 2 | 1 | 0 | 1 | 1 | 0 | 1 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 10-14 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 15-24 | 13 | 11 | 2 | 9 | 7 | 2 | 4 | 4 | 0 |
| 25-34 | 13 | 10 | 3 | 5 | 4 | 1 | 8 | 6 | 2 |
| 35-44 | 8 | 5 | 3 | 5 | 4 | 1 | 3 | 1 | 2 |
| 45-54 | 57 | 47 | 10 | 31 | 26 | 5 | 26 | 21 | 5 |
| 55-64 | 92 | 77 | 15 | 49 | 43 | 6 | 43 | 34 | 9 |
| 65-74 | 139 | 129 | 10 | 85 | 80 | 5 | 54 | 49 | 5 |
| 75-84 | 154 | 138 | 16 | 77 | 68 | 9 | 77 | 70 | 7 |
| 85 and over | 121 | 114 | 7 | 48 | 46 | 2 | 73 | 68 | 5 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 55,224 |
| White | 46,587 |
| Black | 7,059 |
| Other | 1,578 |

Selected Death Statistics, Residents of Perry County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 136 | 116 | 20 | 1,112.3 | 1,217.7 | 808.4 |
| 1 | Heart Diseases | 30 | 24 | 6 | 245.4 | 251.9 | 242.5 |
| 2 | Malignant Neoplasms | 23 | 21 | 2 | 188.1 | 220.4 | 80.8 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 20 | 20 | 0 | 163.6 | 210.0 | 0.0 |
| 4 | Accidents | 10 | 9 | 1 | 81.8 | 94.5 | 40.4 |
| 5 | Cerebrovascular Disease | 6 | 4 | 2 | 49.1 | 42.0 | 80.8 |
| 6 | Septicemia | 5 | 4 | 1 | 40.9 | 42.0 | 40.4 |
| 7 | Alzheimer's Disease | 4 | 3 | 1 | 32.7 | 31.5 | 40.4 |
| 7 | Diabetes Mellitus | 4 | 3 | 1 | 32.7 | 31.5 | 40.4 |
| 9 | Suicide | 3 | 3 | 0 | 24.5 | 31.5 | 0.0 |
| 9 | Nephritis, Nephrotic Syndrome And Nephrosis | 3 | 3 | 0 | 24.5 | 31.5 | 0.0 |
| 9 | Influenza and Pneumonia | 3 | 2 | 1 | 24.5 | 21.0 | 40.4 |
| 12 | Chronic Liver Disease And Cirrhosis | 2 | 2 | 0 | 16.4 | 21.0 | 0.0 |
| 12 | Pneumonitis Due to Solids and Liquids | 2 | 2 | 0 | 16.4 | 21.0 | 0.0 |
| 12 | Viral Hepatitis | 2 | 2 | 0 | 16.4 | 21.0 | 0.0 |
| 15 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 7.4 | 0.0 | 43.5 |
| 15 | Cholelithiasis/Gallbladder Disorder | 1 | 1 | 0 | 8.2 | 10.5 | 0.0 |
| 15 | Hypertension/Hypertensive Renal Disease | 1 | 1 | 0 | 8.2 | 10.5 | 0.0 |
| 15 | Parkinson's Disease | 1 | 1 | 0 | 8.2 | 10.5 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 1 | 1 | 0 | 8.2 | 10.5 | 0.0 |
| | All Other Causes | 14 | 10 | 4 | 114.5 | 105.0 | 161.7 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 136 | 116 | 20 | 69 | 59 | 10 | 67 | 57 | 10 |
| Under 1 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| (Neonatal) | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 15-24 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 25-34 | 2 | 1 | 1 | 2 | 1 | 1 | 0 | 0 | 0 |
| 35-44 | 4 | 3 | 1 | 4 | 3 | 1 | 0 | 0 | 0 |
| 45-54 | 15 | 12 | 3 | 10 | 7 | 3 | 5 | 5 | 0 |
| 55-64 | 26 | 21 | 5 | 16 | 14 | 2 | 10 | 7 | 3 |
| 65-74 | 25 | 25 | 0 | 14 | 14 | 0 | 11 | 11 | 0 |
| 75-84 | 34 | 30 | 4 | 12 | 12 | 0 | 22 | 18 | 4 |
| 85 and over | 27 | 22 | 5 | 9 | 6 | 3 | 18 | 16 | 2 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 12,227 |
| White | 9,526 |
| Black | 2,474 |
| Other | 227 |

Selected Death Statistics, Residents of Pike County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 525 | 296 | 229 | 1,310.6 | 1,615.5 | 1,089.3 |
| 1 | Malignant Neoplasms | 121 | 57 | 64 | 302.1 | 311.1 | 304.4 |
| 2 | Heart Diseases | 78 | 51 | 27 | 194.7 | 278.3 | 128.4 |
| 3 | Diabetes Mellitus | 45 | 19 | 26 | 112.3 | 103.7 | 123.7 |
| 4 | Hypertension/Hypertensive Renal Disease | 33 | 21 | 12 | 82.4 | 114.6 | 57.1 |
| 5 | Cerebrovascular Disease | 28 | 14 | 14 | 69.9 | 76.4 | 66.6 |
| 6 | Emphysema and Other Chronic Lower Resp. Diseases | 24 | 19 | 5 | 59.9 | 103.7 | 23.8 |
| 7 | Accidents | 22 | 11 | 11 | 54.9 | 60.0 | 52.3 |
| 8 | Influenza and Pneumonia | 20 | 13 | 7 | 49.9 | 70.9 | 33.3 |
| 9 | Alzheimer's Disease | 15 | 14 | 1 | 37.4 | 76.4 | 4.8 |
| 10 | Suicide | 12 | 11 | 1 | 30.0 | 60.0 | 4.8 |
| 11 | Nephritis, Nephrotic Syndrome And Nephrosis | 9 | 1 | 8 | 22.5 | 5.5 | 38.1 |
| 12 | In Situ, Benign and Unspecified Neoplasms | 8 | 4 | 4 | 20.0 | 21.8 | 19.0 |
| 13 | Septicemia | 7 | 4 | 3 | 17.5 | 21.8 | 14.3 |
| 14 | Congenital Malformations and Chrom. Abnormalities | 4 | 3 | 1 | 10.0 | 16.4 | 4.8 |
| 15 | Conditions Originating in the Perinatal Period** | 3 | 1 | 2 | 5.6 | 4.9 | 6.0 |
| 15 | Chronic Liver Disease And Cirrhosis | 3 | 2 | 1 | 7.5 | 10.9 | 4.8 |
| 17 | Homicide | 2 | 1 | 1 | 5.0 | 5.5 | 4.8 |
| 17 | Pneumonitis Due to Solids and Liquids | 2 | 0 | 2 | 5.0 | 0.0 | 9.5 |
| 17 | Parkinson's Disease | 2 | 2 | 0 | 5.0 | 10.9 | 0.0 |
| 17 | Viral Hepatitis | 2 | 1 | 1 | 5.0 | 5.5 | 4.8 |
| 21 | Complications Medical/Surgical Care | 1 | 1 | 0 | 2.5 | 5.5 | 0.0 |
| 21 | Cholelithiasis/Gallbladder Disorder | 1 | 1 | 0 | 2.5 | 5.5 | 0.0 |
| 21 | Malnutrition/Nutritional Deficiency | 1 | 0 | 1 | 2.5 | 0.0 | 4.8 |
| 21 | Anemias | 1 | 1 | 0 | 2.5 | 5.5 | 0.0 |
| 21 | HIV Disease | 1 | 0 | 1 | 2.5 | 0.0 | 4.8 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 4 | 3 | 1 | 10.0 | 16.4 | 4.8 |
| | All Other Causes | 76 | 41 | 35 | 189.7 | 223.8 | 166.5 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 525 | 296 | 229 | 254 | 138 | 116 | 271 | 158 | 113 |
| Under 1 | 7 | 4 | 3 | 2 | 1 | 1 | 5 | 3 | 2 |
| (Neonatal) | 5 | 3 | 2 | 2 | 1 | 1 | 3 | 2 | 1 |
| (Postneonatal) | 2 | 1 | 1 | 0 | 0 | 0 | 2 | 1 | 1 |
| 1-4 | 2 | 1 | 1 | 2 | 1 | 1 | 0 | 0 | 0 |
| 5-9 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 5 | 3 | 2 | 4 | 2 | 2 | 1 | 1 | 0 |
| 25-34 | 12 | 7 | 5 | 5 | 3 | 2 | 7 | 4 | 3 |
| 35-44 | 22 | 11 | 11 | 16 | 9 | 7 | 6 | 2 | 4 |
| 45-54 | 42 | 14 | 28 | 22 | 6 | 16 | 20 | 8 | 12 |
| 55-64 | 101 | 46 | 55 | 57 | 25 | 32 | 44 | 21 | 23 |
| 65-74 | 103 | 60 | 43 | 59 | 37 | 22 | 44 | 23 | 21 |
| 75-84 | 91 | 52 | 39 | 46 | 27 | 19 | 45 | 25 | 20 |
| 85 and over | 139 | 98 | 41 | 41 | 27 | 14 | 98 | 71 | 27 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 40,058 |
| White | 18,323 |
| Black | 21,023 |
| Other | 712 |

Selected Death Statistics, Residents of Pontotoc County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 295 | 278 | 16 | 953.2 | 1,092.6 | 332.0 |
| 1 | Heart Diseases | 69 | 68 | 1 | 222.9 | 267.3 | 20.7 |
| 2 | Malignant Neoplasms | 51 | 45 | 6 | 164.8 | 176.9 | 124.5 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 20 | 20 | 0 | 64.6 | 78.6 | 0.0 |
| 3 | Cerebrovascular Disease | 20 | 20 | 0 | 64.6 | 78.6 | 0.0 |
| 5 | Accidents | 18 | 16 | 2 | 58.2 | 62.9 | 41.5 |
| 6 | Diabetes Mellitus | 9 | 7 | 2 | 29.1 | 27.5 | 41.5 |
| 7 | Nephritis, Nephrotic Syndrome And Nephrosis | 7 | 5 | 1 | 22.6 | 19.7 | 20.7 |
| 8 | Alzheimer's Disease | 6 | 6 | 0 | 19.4 | 23.6 | 0.0 |
| 8 | Septicemia | 6 | 6 | 0 | 19.4 | 23.6 | 0.0 |
| 10 | Chronic Liver Disease And Cirrhosis | 5 | 5 | 0 | 16.2 | 19.7 | 0.0 |
| 10 | Influenza and Pneumonia | 5 | 5 | 0 | 16.2 | 19.7 | 0.0 |
| 10 | Parkinson's Disease | 5 | 5 | 0 | 16.2 | 19.7 | 0.0 |
| 13 | Conditions Originating in the Perinatal Period** | 3 | 2 | 1 | 6.5 | 5.0 | 14.7 |
| 13 | Hypertension/Hypertensive Renal Disease | 3 | 3 | 0 | 9.7 | 11.8 | 0.0 |
| 15 | Suicide | 2 | 2 | 0 | 6.5 | 7.9 | 0.0 |
| 16 | Congenital Malformations and Chrom. Abnormalities | 1 | 1 | 0 | 3.2 | 3.9 | 0.0 |
| 16 | Pneumonitis Due to Solids and Liquids | 1 | 1 | 0 | 3.2 | 3.9 | 0.0 |
| 16 | Aortic Aneurysm and Dissection | 1 | 1 | 0 | 3.2 | 3.9 | 0.0 |
| 16 | In Situ, Benign and Unspecified Neoplasms | 1 | 1 | 0 | 3.2 | 3.9 | 0.0 |
| 16 | HIV Disease | 1 | 1 | 0 | 3.2 | 3.9 | 0.0 |
| 16 | Viral Hepatitis | 1 | 1 | 0 | 3.2 | 3.9 | 0.0 |
| 16 | Tuberculosis | 1 | 1 | 0 | 3.2 | 3.9 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 2 | 2 | 0 | 6.5 | 7.9 | 0.0 |
| | All Other Causes | 57 | 54 | 3 | 184.2 | 212.2 | 62.2 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 295 | 278 | 16 | 136 | 127 | 9 | 159 | 151 | 7 |
| Under 1 | 4 | 3 | 1 | 2 | 1 | 1 | 2 | 2 | 0 |
| (Neonatal) | 4 | 3 | 1 | 2 | 1 | 1 | 2 | 2 | 0 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 3 | 3 | 0 | 3 | 3 | 0 | 0 | 0 | 0 |
| 25-34 | 3 | 3 | 0 | 2 | 2 | 0 | 1 | 1 | 0 |
| 35-44 | 11 | 10 | 1 | 4 | 3 | 1 | 7 | 7 | 0 |
| 45-54 | 13 | 12 | 1 | 6 | 5 | 1 | 7 | 7 | 0 |
| 55-64 | 41 | 39 | 2 | 25 | 24 | 1 | 16 | 15 | 1 |
| 65-74 | 63 | 58 | 5 | 36 | 33 | 3 | 27 | 25 | 2 |
| 75-84 | 58 | 55 | 3 | 24 | 23 | 1 | 34 | 32 | 2 |
| 85 and over | 99 | 95 | 3 | 34 | 33 | 1 | 65 | 62 | 2 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 30,950 |
| White | 25,444 |
| Black | 4,820 |
| Other | 686 |

Selected Death Statistics, Residents of Prentiss County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 278 | 248 | 30 | 1,093.3 | 1,158.3 | 833.6 |
| 1 | Heart Diseases | 55 | 51 | 4 | 216.3 | 238.2 | 111.1 |
| 1 | Malignant Neoplasms | 55 | 50 | 5 | 216.3 | 233.5 | 138.9 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 30 | 27 | 3 | 118.0 | 126.1 | 83.4 |
| 4 | Accidents | 22 | 19 | 3 | 86.5 | 88.7 | 83.4 |
| 5 | Diabetes Mellitus | 14 | 11 | 3 | 55.1 | 51.4 | 83.4 |
| 6 | Septicemia | 11 | 11 | 0 | 43.3 | 51.4 | 0.0 |
| 7 | Nephritis, Nephrotic Syndrome And Nephrosis | 10 | 8 | 2 | 39.3 | 37.4 | 55.6 |
| 8 | Influenza and Pneumonia | 9 | 9 | 0 | 35.4 | 42.0 | 0.0 |
| 8 | Cerebrovascular Disease | 9 | 7 | 2 | 35.4 | 32.7 | 55.6 |
| 8 | Alzheimer's Disease | 9 | 9 | 0 | 35.4 | 42.0 | 0.0 |
| 11 | Chronic Liver Disease And Cirrhosis | 4 | 3 | 1 | 15.7 | 14.0 | 27.8 |
| 12 | Pneumonitis Due to Solids and Liquids | 3 | 3 | 0 | 11.8 | 14.0 | 0.0 |
| 13 | Suicide | 2 | 2 | 0 | 7.9 | 9.3 | 0.0 |
| 13 | Parkinson's Disease | 2 | 2 | 0 | 7.9 | 9.3 | 0.0 |
| 15 | Complications Medical/Surgical Care | 1 | 1 | 0 | 3.9 | 4.7 | 0.0 |
| 15 | Homicide | 1 | 0 | 1 | 3.9 | 0.0 | 27.8 |
| 15 | Congenital Malformations and Chrom. Abnormalities | 1 | 1 | 0 | 3.9 | 4.7 | 0.0 |
| 15 | Conditions Originating in the Perinatal Period** | 1 | 1 | 0 | 3.7 | 4.4 | 0.0 |
| 15 | Aortic Aneurysm and Dissection | 1 | 1 | 0 | 3.9 | 4.7 | 0.0 |
| 15 | Hypertension/Hypertensive Renal Disease | 1 | 1 | 0 | 3.9 | 4.7 | 0.0 |
| 15 | Anemias | 1 | 1 | 0 | 3.9 | 4.7 | 0.0 |
| 15 | HIV Disease | 1 | 1 | 0 | 3.9 | 4.7 | 0.0 |
| 15 | Viral Hepatitis | 1 | 1 | 0 | 3.9 | 4.7 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 3 | 2 | 1 | 11.8 | 9.3 | 27.8 |
| | All Other Causes | 31 | 26 | 5 | 121.9 | 121.4 | 138.9 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 278 | 248 | 30 | 135 | 123 | 12 | 143 | 125 | 18 |
| Under 1 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| (Neonatal) | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 25-34 | 7 | 6 | 1 | 3 | 2 | 1 | 4 | 4 | 0 |
| 35-44 | 14 | 11 | 3 | 6 | 6 | 0 | 8 | 5 | 3 |
| 45-54 | 25 | 18 | 7 | 11 | 9 | 2 | 14 | 9 | 5 |
| 55-64 | 38 | 36 | 2 | 27 | 26 | 1 | 11 | 10 | 1 |
| 65-74 | 45 | 42 | 3 | 27 | 25 | 2 | 18 | 17 | 1 |
| 75-84 | 67 | 61 | 6 | 34 | 33 | 1 | 33 | 28 | 5 |
| 85 and over | 79 | 73 | 6 | 24 | 21 | 3 | 55 | 52 | 3 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 25,428 |
| White | 21,411 |
| Black | 3,599 |
| Other | 418 |

Selected Death Statistics, Residents of Quitman County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 84 | 35 | 49 | 1,094.0 | 1,572.3 | 923.3 |
| 1 | Heart Diseases | 28 | 12 | 16 | 364.7 | 539.1 | 301.5 |
| 2 | Malignant Neoplasms | 21 | 6 | 15 | 273.5 | 269.5 | 282.6 |
| 3 | Accidents | 7 | 4 | 3 | 91.2 | 179.7 | 56.5 |
| 4 | Diabetes Mellitus | 4 | 1 | 3 | 52.1 | 44.9 | 56.5 |
| 5 | Influenza and Pneumonia | 3 | 2 | 1 | 39.1 | 89.8 | 18.8 |
| 6 | Alzheimer's Disease | 2 | 0 | 2 | 26.0 | 0.0 | 37.7 |
| 7 | Suicide | 1 | 1 | 0 | 13.0 | 44.9 | 0.0 |
| 7 | Congenital Malformations and Chrom. Abnormalities | 1 | 0 | 1 | 13.0 | 0.0 | 18.8 |
| 7 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 10.0 | 0.0 | 11.8 |
| 7 | Nephritis, Nephrotic Syndrome And Nephrosis | 1 | 0 | 1 | 13.0 | 0.0 | 18.8 |
| 7 | Chronic Liver Disease And Cirrhosis | 1 | 1 | 0 | 13.0 | 44.9 | 0.0 |
| 7 | Aortic Aneurysm and Dissection | 1 | 0 | 1 | 13.0 | 0.0 | 18.8 |
| 7 | Cerebrovascular Disease | 1 | 1 | 0 | 13.0 | 44.9 | 0.0 |
| 7 | Hypertension/Hypertensive Renal Disease | 1 | 0 | 1 | 13.0 | 0.0 | 18.8 |
| 7 | Anemias | 1 | 0 | 1 | 13.0 | 0.0 | 18.8 |
| 7 | Septicemia | 1 | 1 | 0 | 13.0 | 44.9 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 0 | 0 | 0 | 0.0 | 0.0 | 0.0 |
| | All Other Causes | 9 | 6 | 3 | 117.2 | 269.5 | 56.5 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 84 | 35 | 49 | 47 | 21 | 26 | 37 | 14 | 23 |
| Under 1 | 2 | 0 | 2 | 1 | 0 | 1 | 1 | 0 | 1 |
| (Neonatal) | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| (Postneonatal) | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 25-34 | 2 | 1 | 1 | 1 | 1 | 0 | 1 | 0 | 1 |
| 35-44 | 6 | 3 | 3 | 2 | 1 | 1 | 4 | 2 | 2 |
| 45-54 | 11 | 5 | 6 | 6 | 3 | 3 | 5 | 2 | 3 |
| 55-64 | 19 | 5 | 14 | 11 | 5 | 6 | 8 | 0 | 8 |
| 65-74 | 19 | 8 | 11 | 15 | 6 | 9 | 4 | 2 | 2 |
| 75-84 | 19 | 10 | 9 | 8 | 4 | 4 | 11 | 6 | 5 |
| 85 and over | 5 | 3 | 2 | 2 | 1 | 1 | 3 | 2 | 1 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|-------|
| Total | 7,678 |
| White | 2,226 |
| Black | 5,307 |
| Other | 145 |

Selected Death Statistics, Residents of Rankin County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|-------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 1,064 | 886 | 172 | 718.6 | 775.0 | 573.7 |
| 1 | Heart Diseases | 275 | 231 | 44 | 185.7 | 202.1 | 146.8 |
| 2 | Malignant Neoplasms | 223 | 196 | 24 | 150.6 | 171.5 | 80.1 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 69 | 63 | 6 | 46.6 | 55.1 | 20.0 |
| 4 | Cerebrovascular Disease | 57 | 43 | 13 | 38.5 | 37.6 | 43.4 |
| 5 | Accidents | 51 | 46 | 5 | 34.4 | 40.2 | 16.7 |
| 6 | Diabetes Mellitus | 39 | 31 | 8 | 26.3 | 27.1 | 26.7 |
| 7 | Influenza and Pneumonia | 28 | 20 | 8 | 18.9 | 17.5 | 26.7 |
| 8 | Septicemia | 22 | 17 | 4 | 14.9 | 14.9 | 13.3 |
| 9 | Alzheimer's Disease | 18 | 13 | 5 | 12.2 | 11.4 | 16.7 |
| 10 | Nephritis, Nephrotic Syndrome And Nephrosis | 16 | 11 | 5 | 10.8 | 9.6 | 16.7 |
| 11 | Hypertension/Hypertensive Renal Disease | 14 | 14 | 0 | 9.5 | 12.2 | 0.0 |
| 12 | Chronic Liver Disease And Cirrhosis | 13 | 11 | 2 | 8.8 | 9.6 | 6.7 |
| 13 | Suicide | 12 | 11 | 1 | 8.1 | 9.6 | 3.3 |
| 14 | Viral Hepatitis | 8 | 7 | 1 | 5.4 | 6.1 | 3.3 |
| 15 | Congenital Malformations and Chrom. Abnormalities | 6 | 4 | 2 | 4.1 | 3.5 | 6.7 |
| 15 | Anemias | 6 | 3 | 3 | 4.1 | 2.6 | 10.0 |
| 17 | Homicide | 5 | 3 | 2 | 3.4 | 2.6 | 6.7 |
| 17 | Pneumonitis Due to Solids and Liquids | 5 | 4 | 1 | 3.4 | 3.5 | 3.3 |
| 19 | Conditions Originating in the Perinatal Period** | 4 | 2 | 2 | 2.2 | 1.4 | 5.6 |
| 19 | Parkinson's Disease | 4 | 3 | 1 | 2.7 | 2.6 | 3.3 |
| 19 | In Situ, Benign and Unspecified Neoplasms | 4 | 3 | 1 | 2.7 | 2.6 | 3.3 |
| 22 | Aortic Aneurysm and Dissection | 3 | 1 | 2 | 2.0 | 0.9 | 6.7 |
| 23 | Malnutrition/Nutritional Deficiency | 2 | 1 | 1 | 1.4 | 0.9 | 3.3 |
| 23 | HIV Disease | 2 | 1 | 1 | 1.4 | 0.9 | 3.3 |
| 25 | Complications Medical/Surgical Care | 1 | 1 | 0 | 0.7 | 0.9 | 0.0 |
| 25 | Legal Intervention | 1 | 1 | 0 | 0.7 | 0.9 | 0.0 |
| 25 | Peptic Ulcer | 1 | 1 | 0 | 0.7 | 0.9 | 0.0 |
| 25 | Atherosclerosis | 1 | 1 | 0 | 0.7 | 0.9 | 0.0 |
| 25 | Tuberculosis | 1 | 0 | 1 | 0.7 | 0.0 | 3.3 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 10 | 7 | 3 | 6.8 | 6.1 | 10.0 |
| | All Other Causes | 163 | 136 | 26 | 110.1 | 119.0 | 86.7 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 1,064 | 886 | 172 | 550 | 455 | 91 | 514 | 431 | 81 |
| Under 1 | 9 | 5 | 3 | 5 | 2 | 2 | 4 | 3 | 1 |
| (Neonatal) | 6 | 3 | 3 | 4 | 2 | 2 | 2 | 1 | 1 |
| (Postneonatal) | 3 | 2 | 0 | 1 | 0 | 0 | 2 | 2 | 0 |
| 1-4 | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 |
| 5-9 | 2 | 2 | 0 | 1 | 1 | 0 | 1 | 1 | 0 |
| 10-14 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| 15-24 | 11 | 10 | 1 | 7 | 6 | 1 | 4 | 4 | 0 |
| 25-34 | 24 | 21 | 3 | 17 | 15 | 2 | 7 | 6 | 1 |
| 35-44 | 36 | 26 | 10 | 21 | 17 | 4 | 15 | 9 | 6 |
| 45-54 | 109 | 84 | 25 | 69 | 57 | 12 | 40 | 27 | 13 |
| 55-64 | 178 | 143 | 34 | 101 | 82 | 19 | 77 | 61 | 15 |
| 65-74 | 241 | 208 | 31 | 137 | 115 | 20 | 104 | 93 | 11 |
| 75-84 | 217 | 177 | 38 | 99 | 77 | 21 | 118 | 100 | 17 |
| 85 and over | 235 | 209 | 26 | 93 | 83 | 10 | 142 | 126 | 16 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|---------|
| Total | 148,070 |
| White | 114,317 |
| Black | 29,979 |
| Other | 3,774 |

Selected Death Statistics, Residents of Scott County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 282 | 180 | 101 | 990.8 | 1,071.9 | 931.0 |
| 1 | Heart Diseases | 77 | 55 | 22 | 270.5 | 327.5 | 202.8 |
| 2 | Malignant Neoplasms | 59 | 37 | 22 | 207.3 | 220.3 | 202.8 |
| 3 | Accidents | 27 | 17 | 10 | 94.9 | 101.2 | 92.2 |
| 4 | Influenza and Pneumonia | 11 | 7 | 4 | 38.6 | 41.7 | 36.9 |
| 5 | Cerebrovascular Disease | 10 | 7 | 3 | 35.1 | 41.7 | 27.7 |
| 5 | Diabetes Mellitus | 10 | 3 | 7 | 35.1 | 17.9 | 64.5 |
| 7 | Hypertension/Hypertensive Renal Disease | 8 | 3 | 5 | 28.1 | 17.9 | 46.1 |
| 8 | Emphysema and Other Chronic Lower Resp. Diseases | 7 | 6 | 1 | 24.6 | 35.7 | 9.2 |
| 8 | Alzheimer's Disease | 7 | 6 | 1 | 24.6 | 35.7 | 9.2 |
| 10 | Nephritis, Nephrotic Syndrome And Nephrosis | 5 | 4 | 1 | 17.6 | 23.8 | 9.2 |
| 11 | Septicemia | 4 | 3 | 1 | 14.1 | 17.9 | 9.2 |
| 12 | Homicide | 3 | 1 | 2 | 10.5 | 6.0 | 18.4 |
| 12 | Suicide | 3 | 2 | 1 | 10.5 | 11.9 | 9.2 |
| 14 | Conditions Originating in the Perinatal Period** | 2 | 1 | 1 | 4.1 | 3.5 | 5.2 |
| 14 | Pneumonitis Due to Solids and Liquids | 2 | 0 | 2 | 7.0 | 0.0 | 18.4 |
| 14 | Parkinson's Disease | 2 | 1 | 1 | 7.0 | 6.0 | 9.2 |
| 14 | In Situ, Benign and Unspecified Neoplasms | 2 | 2 | 0 | 7.0 | 11.9 | 0.0 |
| 14 | HIV Disease | 2 | 1 | 1 | 7.0 | 6.0 | 9.2 |
| 19 | Complications Medical/Surgical Care | 1 | 0 | 1 | 3.5 | 0.0 | 9.2 |
| 19 | Congenital Malformations and Chrom. Abnormalities | 1 | 1 | 0 | 3.5 | 6.0 | 0.0 |
| 19 | Cholelithiasis/Gallbladder Disorder | 1 | 0 | 1 | 3.5 | 0.0 | 9.2 |
| 19 | Chronic Liver Disease And Cirrhosis | 1 | 1 | 0 | 3.5 | 6.0 | 0.0 |
| 19 | Peptic Ulcer | 1 | 1 | 0 | 3.5 | 6.0 | 0.0 |
| 19 | Anemias | 1 | 0 | 1 | 3.5 | 0.0 | 9.2 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 3 | 2 | 1 | 10.5 | 11.9 | 9.2 |
| | All Other Causes | 32 | 19 | 12 | 112.4 | 113.1 | 110.6 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births.
 Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 282 | 180 | 101 | 164 | 98 | 65 | 118 | 82 | 36 |
| Under 1 | 6 | 4 | 1 | 4 | 3 | 0 | 2 | 1 | 1 |
| (Neonatal) | 3 | 2 | 1 | 2 | 2 | 0 | 1 | 0 | 1 |
| (Postneonatal) | 3 | 2 | 0 | 2 | 1 | 0 | 1 | 1 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 2 | 1 | 1 | 2 | 1 | 1 | 0 | 0 | 0 |
| 10-14 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 15-24 | 8 | 3 | 5 | 7 | 2 | 5 | 1 | 1 | 0 |
| 25-34 | 9 | 4 | 5 | 7 | 3 | 4 | 2 | 1 | 1 |
| 35-44 | 7 | 6 | 1 | 5 | 4 | 1 | 2 | 2 | 0 |
| 45-54 | 30 | 14 | 16 | 19 | 9 | 10 | 11 | 5 | 6 |
| 55-64 | 47 | 25 | 22 | 32 | 17 | 15 | 15 | 8 | 7 |
| 65-74 | 50 | 32 | 18 | 31 | 20 | 11 | 19 | 12 | 7 |
| 75-84 | 61 | 43 | 18 | 34 | 24 | 10 | 27 | 19 | 8 |
| 85 and over | 61 | 47 | 14 | 22 | 14 | 8 | 39 | 33 | 6 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 28,461 |
| White | 16,793 |
| Black | 10,848 |
| Other | 820 |

Selected Death Statistics, Residents of Sharkey County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 70 | 25 | 45 | 1,506.3 | 1,843.7 | 1,391.9 |
| 1 | Heart Diseases | 21 | 6 | 15 | 451.9 | 442.5 | 464.0 |
| 2 | Malignant Neoplasms | 8 | 1 | 7 | 172.2 | 73.7 | 216.5 |
| 3 | Alzheimer's Disease | 5 | 3 | 2 | 107.6 | 221.2 | 61.9 |
| 4 | Accidents | 4 | 2 | 2 | 86.1 | 147.5 | 61.9 |
| 4 | Influenza and Pneumonia | 4 | 0 | 4 | 86.1 | 0.0 | 123.7 |
| 6 | Emphysema and Other Chronic Lower Resp. Diseases | 3 | 2 | 1 | 64.6 | 147.5 | 30.9 |
| 7 | Nephritis, Nephrotic Syndrome And Nephrosis | 2 | 1 | 1 | 43.0 | 73.7 | 30.9 |
| 8 | Homicide | 1 | 0 | 1 | 21.5 | 0.0 | 30.9 |
| 8 | Chronic Liver Disease And Cirrhosis | 1 | 1 | 0 | 21.5 | 73.7 | 0.0 |
| 8 | Pneumonitis Due to Solids and Liquids | 1 | 1 | 0 | 21.5 | 73.7 | 0.0 |
| 8 | Atherosclerosis | 1 | 0 | 1 | 21.5 | 0.0 | 30.9 |
| 8 | Cerebrovascular Disease | 1 | 0 | 1 | 21.5 | 0.0 | 30.9 |
| 8 | Parkinson's Disease | 1 | 1 | 0 | 21.5 | 73.7 | 0.0 |
| 8 | Diabetes Mellitus | 1 | 1 | 0 | 21.5 | 73.7 | 0.0 |
| 8 | Anemias | 1 | 0 | 1 | 21.5 | 0.0 | 30.9 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 2 | 1 | 1 | 43.0 | 73.7 | 30.9 |
| | All Other Causes | 13 | 5 | 8 | 279.8 | 368.7 | 247.4 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 70 | 25 | 45 | 31 | 11 | 20 | 39 | 14 | 25 |
| Under 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Neonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 2 | 2 | 0 | 0 | 0 | 0 | 2 | 2 | 0 |
| 25-34 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 35-44 | 3 | 1 | 2 | 3 | 1 | 2 | 0 | 0 | 0 |
| 45-54 | 5 | 1 | 4 | 3 | 0 | 3 | 2 | 1 | 1 |
| 55-64 | 17 | 4 | 13 | 10 | 2 | 8 | 7 | 2 | 5 |
| 65-74 | 10 | 4 | 6 | 5 | 4 | 1 | 5 | 0 | 5 |
| 75-84 | 17 | 5 | 12 | 6 | 2 | 4 | 11 | 3 | 8 |
| 85 and over | 15 | 8 | 7 | 3 | 2 | 1 | 12 | 6 | 6 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|-------|
| Total | 4,647 |
| White | 1,356 |
| Black | 3,233 |
| Other | 58 |

Selected Death Statistics, Residents of Simpson County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 299 | 232 | 67 | 1,088.7 | 1,350.3 | 683.0 |
| 1 | Malignant Neoplasms | 73 | 57 | 16 | 265.8 | 331.7 | 163.1 |
| 2 | Heart Diseases | 72 | 59 | 13 | 262.2 | 343.4 | 132.5 |
| 3 | Accidents | 17 | 13 | 4 | 61.9 | 75.7 | 40.8 |
| 4 | Cerebrovascular Disease | 15 | 11 | 4 | 54.6 | 64.0 | 40.8 |
| 5 | Emphysema and Other Chronic Lower Resp. Diseases | 13 | 12 | 1 | 47.3 | 69.8 | 10.2 |
| 6 | Influenza and Pneumonia | 11 | 9 | 2 | 40.1 | 52.4 | 20.4 |
| 7 | Suicide | 5 | 5 | 0 | 18.2 | 29.1 | 0.0 |
| 7 | Nephritis, Nephrotic Syndrome And Nephrosis | 5 | 3 | 2 | 18.2 | 17.5 | 20.4 |
| 7 | Alzheimer's Disease | 5 | 4 | 1 | 18.2 | 23.3 | 10.2 |
| 7 | Septicemia | 5 | 4 | 1 | 18.2 | 23.3 | 10.2 |
| 11 | Chronic Liver Disease And Cirrhosis | 4 | 4 | 0 | 14.6 | 23.3 | 0.0 |
| 11 | Diabetes Mellitus | 4 | 4 | 0 | 14.6 | 23.3 | 0.0 |
| 13 | Hypertension/Hypertensive Renal Disease | 3 | 1 | 2 | 10.9 | 5.8 | 20.4 |
| 14 | Homicide | 2 | 1 | 1 | 7.3 | 5.8 | 10.2 |
| 14 | In Situ, Benign and Unspecified Neoplasms | 2 | 1 | 1 | 7.3 | 5.8 | 10.2 |
| 16 | Congenital Malformations and Chrom. Abnormalities | 1 | 1 | 0 | 3.6 | 5.8 | 0.0 |
| 16 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 3.3 | 0.0 | 8.8 |
| 16 | Cholelithiasis/Gallbladder Disorder | 1 | 0 | 1 | 3.6 | 0.0 | 10.2 |
| 16 | Peptic Ulcer | 1 | 1 | 0 | 3.6 | 5.8 | 0.0 |
| 16 | Aortic Aneurysm and Dissection | 1 | 1 | 0 | 3.6 | 5.8 | 0.0 |
| 16 | Parkinson's Disease | 1 | 1 | 0 | 3.6 | 5.8 | 0.0 |
| 16 | HIV Disease | 1 | 0 | 1 | 3.6 | 0.0 | 10.2 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 13 | 8 | 5 | 47.3 | 46.6 | 51.0 |
| | All Other Causes | 43 | 32 | 11 | 156.6 | 186.2 | 112.1 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 299 | 232 | 67 | 146 | 111 | 35 | 153 | 121 | 32 |
| Under 1 | 3 | 1 | 2 | 0 | 0 | 0 | 3 | 1 | 2 |
| (Neonatal) | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| (Postneonatal) | 2 | 1 | 1 | 0 | 0 | 0 | 2 | 1 | 1 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 5 | 4 | 1 | 1 | 1 | 0 | 4 | 3 | 1 |
| 25-34 | 3 | 2 | 1 | 2 | 1 | 1 | 1 | 1 | 0 |
| 35-44 | 8 | 5 | 3 | 5 | 3 | 2 | 3 | 2 | 1 |
| 45-54 | 26 | 20 | 6 | 9 | 8 | 1 | 17 | 12 | 5 |
| 55-64 | 56 | 37 | 19 | 36 | 23 | 13 | 20 | 14 | 6 |
| 65-74 | 66 | 49 | 17 | 41 | 30 | 11 | 25 | 19 | 6 |
| 75-84 | 75 | 63 | 12 | 36 | 33 | 3 | 39 | 30 | 9 |
| 85 and over | 56 | 51 | 5 | 15 | 12 | 3 | 41 | 39 | 2 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 27,463 |
| White | 17,182 |
| Black | 9,810 |
| Other | 471 |

Selected Death Statistics, Residents of Smith County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 177 | 143 | 33 | 1,093.4 | 1,170.4 | 863.4 |
| 1 | Heart Diseases | 68 | 57 | 11 | 420.1 | 466.5 | 287.8 |
| 2 | Malignant Neoplasms | 35 | 25 | 10 | 216.2 | 204.6 | 261.6 |
| 3 | Accidents | 12 | 9 | 2 | 74.1 | 73.7 | 52.3 |
| 4 | Cerebrovascular Disease | 10 | 10 | 0 | 61.8 | 81.8 | 0.0 |
| 5 | Alzheimer's Disease | 9 | 6 | 3 | 55.6 | 49.1 | 78.5 |
| 6 | Emphysema and Other Chronic Lower Resp. Diseases | 4 | 4 | 0 | 24.7 | 32.7 | 0.0 |
| 7 | Suicide | 3 | 2 | 1 | 18.5 | 16.4 | 26.2 |
| 7 | Influenza and Pneumonia | 3 | 2 | 1 | 18.5 | 16.4 | 26.2 |
| 7 | Hypertension/Hypertensive Renal Disease | 3 | 2 | 1 | 18.5 | 16.4 | 26.2 |
| 7 | Septicemia | 3 | 2 | 1 | 18.5 | 16.4 | 26.2 |
| 11 | Nephritis, Nephrotic Syndrome And Nephrosis | 2 | 2 | 0 | 12.4 | 16.4 | 0.0 |
| 11 | Diabetes Mellitus | 2 | 2 | 0 | 12.4 | 16.4 | 0.0 |
| 13 | Homicide | 1 | 1 | 0 | 6.2 | 8.2 | 0.0 |
| 13 | Pneumonitis Due to Solids and Liquids | 1 | 1 | 0 | 6.2 | 8.2 | 0.0 |
| 13 | Anemias | 1 | 1 | 0 | 6.2 | 8.2 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 2 | 1 | 1 | 12.4 | 8.2 | 26.2 |
| | All Other Causes | 18 | 16 | 2 | 111.2 | 131.0 | 52.3 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 177 | 143 | 33 | 93 | 71 | 21 | 84 | 72 | 12 |
| Under 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Neonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 |
| 15-24 | 2 | 1 | 1 | 1 | 0 | 1 | 1 | 1 | 0 |
| 25-34 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 35-44 | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 |
| 45-54 | 12 | 8 | 4 | 10 | 6 | 4 | 2 | 2 | 0 |
| 55-64 | 35 | 29 | 5 | 22 | 17 | 4 | 13 | 12 | 1 |
| 65-74 | 36 | 30 | 6 | 21 | 15 | 6 | 15 | 15 | 0 |
| 75-84 | 52 | 48 | 4 | 27 | 25 | 2 | 25 | 23 | 2 |
| 85 and over | 37 | 24 | 13 | 11 | 7 | 4 | 26 | 17 | 9 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 16,188 |
| White | 12,218 |
| Black | 3,822 |
| Other | 148 |

Selected Death Statistics, Residents of Stone County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 160 | 141 | 19 | 895.1 | 1,009.0 | 537.3 |
| 1 | Malignant Neoplasms | 36 | 30 | 6 | 201.4 | 214.7 | 169.7 |
| 2 | Heart Diseases | 32 | 29 | 3 | 179.0 | 207.5 | 84.8 |
| 3 | Accidents | 12 | 12 | 0 | 67.1 | 85.9 | 0.0 |
| 4 | Emphysema and Other Chronic Lower Resp. Diseases | 10 | 9 | 1 | 55.9 | 64.4 | 28.3 |
| 5 | Influenza and Pneumonia | 9 | 7 | 2 | 50.3 | 50.1 | 56.6 |
| 6 | Cerebrovascular Disease | 6 | 5 | 1 | 33.6 | 35.8 | 28.3 |
| 7 | Alzheimer's Disease | 5 | 4 | 1 | 28.0 | 28.6 | 28.3 |
| 8 | Suicide | 3 | 3 | 0 | 16.8 | 21.5 | 0.0 |
| 8 | Diabetes Mellitus | 3 | 3 | 0 | 16.8 | 21.5 | 0.0 |
| 10 | Nephritis, Nephrotic Syndrome And Nephrosis | 2 | 2 | 0 | 11.2 | 14.3 | 0.0 |
| 10 | In Situ, Benign and Unspecified Neoplasms | 2 | 2 | 0 | 11.2 | 14.3 | 0.0 |
| 10 | HIV Disease | 2 | 1 | 1 | 11.2 | 7.2 | 28.3 |
| 10 | Septicemia | 2 | 2 | 0 | 11.2 | 14.3 | 0.0 |
| 14 | Homicide | 1 | 1 | 0 | 5.6 | 7.2 | 0.0 |
| 14 | Congenital Malformations and Chrom. Abnormalities | 1 | 1 | 0 | 5.6 | 7.2 | 0.0 |
| 14 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 4.5 | 0.0 | 23.8 |
| 14 | Chronic Liver Disease And Cirrhosis | 1 | 1 | 0 | 5.6 | 7.2 | 0.0 |
| 14 | Pneumonitis Due to Solids and Liquids | 1 | 1 | 0 | 5.6 | 7.2 | 0.0 |
| 14 | Hypertension/Hypertensive Renal Disease | 1 | 1 | 0 | 5.6 | 7.2 | 0.0 |
| 14 | Parkinson's Disease | 1 | 1 | 0 | 5.6 | 7.2 | 0.0 |
| 14 | Viral Hepatitis | 1 | 1 | 0 | 5.6 | 7.2 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 0 | 0 | 0 | 0.0 | 0.0 | 0.0 |
| | All Other Causes | 28 | 25 | 3 | 156.6 | 178.9 | 84.8 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 160 | 141 | 19 | 86 | 76 | 10 | 74 | 65 | 9 |
| Under 1 | 2 | 1 | 1 | 2 | 1 | 1 | 0 | 0 | 0 |
| (Neonatal) | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| (Postneonatal) | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 3 | 3 | 0 | 3 | 3 | 0 | 0 | 0 | 0 |
| 25-34 | 3 | 3 | 0 | 2 | 2 | 0 | 1 | 1 | 0 |
| 35-44 | 5 | 5 | 0 | 4 | 4 | 0 | 1 | 1 | 0 |
| 45-54 | 15 | 11 | 4 | 8 | 6 | 2 | 7 | 5 | 2 |
| 55-64 | 28 | 23 | 5 | 17 | 13 | 4 | 11 | 10 | 1 |
| 65-74 | 38 | 34 | 4 | 25 | 23 | 2 | 13 | 11 | 2 |
| 75-84 | 40 | 38 | 2 | 19 | 18 | 1 | 21 | 20 | 1 |
| 85 and over | 26 | 23 | 3 | 6 | 6 | 0 | 20 | 17 | 3 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 17,875 |
| White | 13,974 |
| Black | 3,536 |
| Other | 365 |

Selected Death Statistics, Residents of Sunflower County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 284 | 108 | 175 | 1,032.9 | 1,523.5 | 872.0 |
| 1 | Heart Diseases | 68 | 28 | 40 | 247.3 | 395.0 | 199.3 |
| 2 | Malignant Neoplasms | 62 | 26 | 36 | 225.5 | 366.8 | 179.4 |
| 3 | Diabetes Mellitus | 24 | 6 | 18 | 87.3 | 84.6 | 89.7 |
| 4 | Cerebrovascular Disease | 15 | 8 | 7 | 54.6 | 112.9 | 34.9 |
| 5 | Nephritis, Nephrotic Syndrome And Nephrosis | 10 | 2 | 8 | 36.4 | 28.2 | 39.9 |
| 5 | Emphysema and Other Chronic Lower Resp. Diseases | 10 | 6 | 4 | 36.4 | 84.6 | 19.9 |
| 5 | Alzheimer's Disease | 10 | 2 | 8 | 36.4 | 28.2 | 39.9 |
| 8 | Accidents | 9 | 5 | 4 | 32.7 | 70.5 | 19.9 |
| 9 | Influenza and Pneumonia | 8 | 4 | 3 | 29.1 | 56.4 | 14.9 |
| 9 | HIV Disease | 8 | 2 | 6 | 29.1 | 28.2 | 29.9 |
| 11 | Hypertension/Hypertensive Renal Disease | 7 | 3 | 4 | 25.5 | 42.3 | 19.9 |
| 12 | Suicide | 3 | 1 | 2 | 10.9 | 14.1 | 10.0 |
| 12 | Congenital Malformations and Chrom. Abnormalities | 3 | 0 | 3 | 10.9 | 0.0 | 14.9 |
| 12 | Homicide | 3 | 0 | 3 | 10.9 | 0.0 | 14.9 |
| 15 | Chronic Liver Disease And Cirrhosis | 2 | 1 | 1 | 7.3 | 14.1 | 5.0 |
| 15 | Pneumonitis Due to Solids and Liquids | 2 | 1 | 1 | 7.3 | 14.1 | 5.0 |
| 17 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 3.1 | 0.0 | 3.8 |
| 17 | Hernia | 1 | 1 | 0 | 3.6 | 14.1 | 0.0 |
| 17 | Peptic Ulcer | 1 | 0 | 1 | 3.6 | 0.0 | 5.0 |
| 17 | Parkinson's Disease | 1 | 1 | 0 | 3.6 | 14.1 | 0.0 |
| 17 | Meningitis | 1 | 0 | 1 | 3.6 | 0.0 | 5.0 |
| 17 | Malnutrition/Nutritional Deficiency | 1 | 0 | 1 | 3.6 | 0.0 | 5.0 |
| 17 | Anemias | 1 | 0 | 1 | 3.6 | 0.0 | 5.0 |
| 17 | In Situ, Benign and Unspecified Neoplasms | 1 | 1 | 0 | 3.6 | 14.1 | 0.0 |
| 17 | Septicemia | 1 | 1 | 0 | 3.6 | 14.1 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 3 | 1 | 2 | 10.9 | 14.1 | 10.0 |
| | All Other Causes | 28 | 8 | 20 | 101.8 | 112.9 | 99.7 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 284 | 108 | 175 | 142 | 53 | 89 | 142 | 55 | 86 |
| Under 1 | 5 | 0 | 4 | 2 | 0 | 2 | 3 | 0 | 2 |
| (Neonatal) | 2 | 0 | 2 | 1 | 0 | 1 | 1 | 0 | 1 |
| (Postneonatal) | 3 | 0 | 2 | 1 | 0 | 1 | 2 | 0 | 1 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 2 | 0 | 2 | 2 | 0 | 2 | 0 | 0 | 0 |
| 15-24 | 5 | 2 | 3 | 3 | 0 | 3 | 2 | 2 | 0 |
| 25-34 | 4 | 0 | 4 | 2 | 0 | 2 | 2 | 0 | 2 |
| 35-44 | 15 | 3 | 12 | 12 | 3 | 9 | 3 | 0 | 3 |
| 45-54 | 32 | 6 | 26 | 17 | 4 | 13 | 15 | 2 | 13 |
| 55-64 | 58 | 15 | 43 | 39 | 13 | 26 | 19 | 2 | 17 |
| 65-74 | 60 | 29 | 31 | 34 | 15 | 19 | 26 | 14 | 12 |
| 75-84 | 53 | 28 | 25 | 18 | 13 | 5 | 35 | 15 | 20 |
| 85 and over | 50 | 25 | 25 | 13 | 5 | 8 | 37 | 20 | 17 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 27,496 |
| White | 7,089 |
| Black | 20,068 |
| Other | 339 |

Selected Death Statistics, Residents of Tallahatchie County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 146 | 69 | 76 | 989.1 | 1,130.6 | 913.4 |
| 1 | Heart Diseases | 33 | 16 | 16 | 223.6 | 262.2 | 192.3 |
| 1 | Malignant Neoplasms | 33 | 13 | 20 | 223.6 | 213.0 | 240.4 |
| 3 | Hypertension/Hypertensive Renal Disease | 10 | 4 | 6 | 67.7 | 65.5 | 72.1 |
| 3 | Diabetes Mellitus | 10 | 5 | 5 | 67.7 | 81.9 | 60.1 |
| 5 | Influenza and Pneumonia | 9 | 4 | 5 | 61.0 | 65.5 | 60.1 |
| 6 | Emphysema and Other Chronic Lower Resp. Diseases | 7 | 5 | 2 | 47.4 | 81.9 | 24.0 |
| 6 | Cerebrovascular Disease | 7 | 1 | 6 | 47.4 | 16.4 | 72.1 |
| 8 | Accidents | 6 | 4 | 2 | 40.6 | 65.5 | 24.0 |
| 8 | Alzheimer's Disease | 6 | 6 | 0 | 40.6 | 98.3 | 0.0 |
| 10 | Nephritis, Nephrotic Syndrome And Nephrosis | 3 | 1 | 2 | 20.3 | 16.4 | 24.0 |
| 11 | Septicemia | 2 | 1 | 1 | 13.5 | 16.4 | 12.0 |
| 12 | Complications Medical/Surgical Care | 1 | 1 | 0 | 6.8 | 16.4 | 0.0 |
| 12 | Congenital Malformations and Chrom. Abnormalities | 1 | 0 | 1 | 6.8 | 0.0 | 12.0 |
| 12 | Chronic Liver Disease And Cirrhosis | 1 | 1 | 0 | 6.8 | 16.4 | 0.0 |
| 12 | Pneumonitis Due to Solids and Liquids | 1 | 1 | 0 | 6.8 | 16.4 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 0 | 0 | 0 | 0.0 | 0.0 | 0.0 |
| | All Other Causes | 16 | 6 | 10 | 108.4 | 98.3 | 120.2 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births.

Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 146 | 69 | 76 | 77 | 38 | 38 | 69 | 31 | 38 |
| Under 1 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| (Neonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 15-24 | 2 | 1 | 1 | 1 | 1 | 0 | 1 | 0 | 1 |
| 25-34 | 4 | 1 | 3 | 1 | 1 | 0 | 3 | 0 | 3 |
| 35-44 | 3 | 0 | 3 | 1 | 0 | 1 | 2 | 0 | 2 |
| 45-54 | 17 | 4 | 13 | 10 | 3 | 7 | 7 | 1 | 6 |
| 55-64 | 22 | 6 | 16 | 16 | 5 | 11 | 6 | 1 | 5 |
| 65-74 | 28 | 13 | 15 | 13 | 6 | 7 | 15 | 7 | 8 |
| 75-84 | 34 | 21 | 13 | 20 | 13 | 7 | 14 | 8 | 6 |
| 85 and over | 34 | 23 | 10 | 13 | 9 | 3 | 21 | 14 | 7 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 14,761 |
| White | 6,103 |
| Black | 8,321 |
| Other | 337 |

Selected Death Statistics, Residents of Tate County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 271 | 198 | 70 | 960.9 | 1,051.8 | 793.0 |
| 1 | Heart Diseases | 96 | 61 | 34 | 340.4 | 324.1 | 385.2 |
| 2 | Malignant Neoplasms | 57 | 42 | 14 | 202.1 | 223.1 | 158.6 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 20 | 17 | 3 | 70.9 | 90.3 | 34.0 |
| 4 | Accidents | 14 | 10 | 4 | 49.6 | 53.1 | 45.3 |
| 5 | Alzheimer's Disease | 12 | 10 | 2 | 42.5 | 53.1 | 22.7 |
| 6 | Cerebrovascular Disease | 11 | 9 | 1 | 39.0 | 47.8 | 11.3 |
| 6 | Diabetes Mellitus | 11 | 9 | 2 | 39.0 | 47.8 | 22.7 |
| 8 | Suicide | 5 | 5 | 0 | 17.7 | 26.6 | 0.0 |
| 8 | Nephritis, Nephrotic Syndrome And Nephrosis | 5 | 2 | 3 | 17.7 | 10.6 | 34.0 |
| 8 | Influenza and Pneumonia | 5 | 4 | 1 | 17.7 | 21.2 | 11.3 |
| 11 | Chronic Liver Disease And Cirrhosis | 2 | 2 | 0 | 7.1 | 10.6 | 0.0 |
| 11 | Hypertension/Hypertensive Renal Disease | 2 | 2 | 0 | 7.1 | 10.6 | 0.0 |
| 11 | Septicemia | 2 | 1 | 1 | 7.1 | 5.3 | 11.3 |
| 14 | Homicide | 1 | 0 | 1 | 3.5 | 0.0 | 11.3 |
| 14 | Congenital Malformations and Chrom. Abnormalities | 1 | 1 | 0 | 3.5 | 5.3 | 0.0 |
| 14 | Cholelithiasis/Gallbladder Disorder | 1 | 1 | 0 | 3.5 | 5.3 | 0.0 |
| 14 | Anemias | 1 | 1 | 0 | 3.5 | 5.3 | 0.0 |
| 14 | Viral Hepatitis | 1 | 1 | 0 | 3.5 | 5.3 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 0 | 0 | 0 | 0.0 | 0.0 | 0.0 |
| | All Other Causes | 24 | 20 | 4 | 85.1 | 106.2 | 45.3 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births.
 Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 271 | 198 | 70 | 137 | 103 | 34 | 134 | 95 | 36 |
| Under 1 | 2 | 1 | 1 | 1 | 0 | 1 | 1 | 1 | 0 |
| (Neonatal) | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 |
| (Postneonatal) | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 15-24 | 4 | 3 | 1 | 4 | 3 | 1 | 0 | 0 | 0 |
| 25-34 | 5 | 3 | 2 | 3 | 2 | 1 | 2 | 1 | 1 |
| 35-44 | 6 | 4 | 2 | 2 | 2 | 0 | 4 | 2 | 2 |
| 45-54 | 15 | 6 | 9 | 9 | 4 | 5 | 6 | 2 | 4 |
| 55-64 | 43 | 28 | 13 | 24 | 16 | 8 | 19 | 12 | 5 |
| 65-74 | 64 | 53 | 10 | 35 | 31 | 4 | 29 | 22 | 6 |
| 75-84 | 64 | 49 | 15 | 37 | 28 | 9 | 27 | 21 | 6 |
| 85 and over | 67 | 50 | 17 | 21 | 16 | 5 | 46 | 34 | 12 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 28,204 |
| White | 18,824 |
| Black | 8,827 |
| Other | 553 |

Selected Death Statistics, Residents of Tippah County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 265 | 237 | 26 | 1,202.4 | 1,322.5 | 707.9 |
| 1 | Heart Diseases | 78 | 70 | 7 | 353.9 | 390.6 | 190.6 |
| 2 | Malignant Neoplasms | 43 | 40 | 3 | 195.1 | 223.2 | 81.7 |
| 3 | Accidents | 17 | 15 | 2 | 77.1 | 83.7 | 54.5 |
| 4 | Emphysema and Other Chronic Lower Resp. Diseases | 11 | 11 | 0 | 49.9 | 61.4 | 0.0 |
| 4 | Alzheimer's Disease | 11 | 10 | 1 | 49.9 | 55.8 | 27.2 |
| 6 | Influenza and Pneumonia | 9 | 8 | 0 | 40.8 | 44.6 | 0.0 |
| 6 | Cerebrovascular Disease | 9 | 6 | 3 | 40.8 | 33.5 | 81.7 |
| 8 | Nephritis, Nephrotic Syndrome And Nephrosis | 4 | 4 | 0 | 18.1 | 22.3 | 0.0 |
| 9 | Homicide | 3 | 3 | 0 | 13.6 | 16.7 | 0.0 |
| 9 | Conditions Originating in the Perinatal Period** | 3 | 1 | 2 | 10.5 | 4.0 | 52.6 |
| 9 | Chronic Liver Disease And Cirrhosis | 3 | 3 | 0 | 13.6 | 16.7 | 0.0 |
| 12 | Suicide | 2 | 2 | 0 | 9.1 | 11.2 | 0.0 |
| 12 | Parkinson's Disease | 2 | 2 | 0 | 9.1 | 11.2 | 0.0 |
| 12 | Septicemia | 2 | 2 | 0 | 9.1 | 11.2 | 0.0 |
| 15 | Congenital Malformations and Chrom. Abnormalities | 1 | 0 | 1 | 4.5 | 0.0 | 27.2 |
| 15 | Infections of Kidney | 1 | 1 | 0 | 4.5 | 5.6 | 0.0 |
| 15 | Hypertension/Hypertensive Renal Disease | 1 | 1 | 0 | 4.5 | 5.6 | 0.0 |
| 15 | Diabetes Mellitus | 1 | 1 | 0 | 4.5 | 5.6 | 0.0 |
| 15 | In Situ, Benign and Unspecified Neoplasms | 1 | 1 | 0 | 4.5 | 5.6 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 4 | 4 | 0 | 18.1 | 22.3 | 0.0 |
| | All Other Causes | 59 | 52 | 7 | 267.7 | 290.2 | 190.6 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 265 | 237 | 26 | 134 | 123 | 10 | 131 | 114 | 16 |
| Under 1 | 4 | 1 | 3 | 1 | 0 | 1 | 3 | 1 | 2 |
| (Neonatal) | 4 | 1 | 3 | 1 | 0 | 1 | 3 | 1 | 2 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 5 | 3 | 2 | 3 | 2 | 1 | 2 | 1 | 1 |
| 25-34 | 3 | 2 | 0 | 2 | 2 | 0 | 1 | 0 | 0 |
| 35-44 | 7 | 6 | 1 | 4 | 3 | 1 | 3 | 3 | 0 |
| 45-54 | 25 | 22 | 2 | 17 | 15 | 1 | 8 | 7 | 1 |
| 55-64 | 34 | 31 | 3 | 23 | 21 | 2 | 11 | 10 | 1 |
| 65-74 | 56 | 52 | 4 | 28 | 26 | 2 | 28 | 26 | 2 |
| 75-84 | 63 | 58 | 5 | 35 | 34 | 1 | 28 | 24 | 4 |
| 85 and over | 68 | 62 | 6 | 21 | 20 | 1 | 47 | 42 | 5 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 22,039 |
| White | 17,920 |
| Black | 3,673 |
| Other | 446 |

Selected Death Statistics, Residents of Tishomingo County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 297 | 290 | 6 | 1,529.4 | 1,559.9 | 1,096.9 |
| 1 | Heart Diseases | 88 | 86 | 2 | 453.1 | 462.6 | 365.6 |
| 2 | Malignant Neoplasms | 60 | 58 | 2 | 309.0 | 312.0 | 365.6 |
| 3 | Cerebrovascular Disease | 22 | 22 | 0 | 113.3 | 118.3 | 0.0 |
| 4 | Accidents | 20 | 20 | 0 | 103.0 | 107.6 | 0.0 |
| 5 | Alzheimer's Disease | 11 | 11 | 0 | 56.6 | 59.2 | 0.0 |
| 6 | Nephritis, Nephrotic Syndrome And Nephrosis | 10 | 10 | 0 | 51.5 | 53.8 | 0.0 |
| 6 | Septicemia | 10 | 10 | 0 | 51.5 | 53.8 | 0.0 |
| 8 | Emphysema and Other Chronic Lower Resp. Diseases | 9 | 9 | 0 | 46.3 | 48.4 | 0.0 |
| 9 | Influenza and Pneumonia | 8 | 7 | 1 | 41.2 | 37.7 | 182.8 |
| 10 | Diabetes Mellitus | 5 | 5 | 0 | 25.7 | 26.9 | 0.0 |
| 11 | Suicide | 4 | 4 | 0 | 20.6 | 21.5 | 0.0 |
| 11 | Parkinson's Disease | 4 | 3 | 1 | 20.6 | 16.1 | 182.8 |
| 13 | Pneumonitis Due to Solids and Liquids | 3 | 3 | 0 | 15.4 | 16.1 | 0.0 |
| 14 | Conditions Originating in the Perinatal Period** | 2 | 2 | 0 | 10.7 | 10.9 | 0.0 |
| 14 | Chronic Liver Disease And Cirrhosis | 2 | 2 | 0 | 10.3 | 10.8 | 0.0 |
| 16 | Peptic Ulcer | 1 | 1 | 0 | 5.1 | 5.4 | 0.0 |
| 16 | Aortic Aneurysm and Dissection | 1 | 1 | 0 | 5.1 | 5.4 | 0.0 |
| 16 | Atherosclerosis | 1 | 1 | 0 | 5.1 | 5.4 | 0.0 |
| 16 | Hypertension/Hypertensive Renal Disease | 1 | 1 | 0 | 5.1 | 5.4 | 0.0 |
| 16 | HIV Disease | 1 | 0 | 0 | 5.1 | 0.0 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 4 | 4 | 0 | 20.6 | 21.5 | 0.0 |
| | All Other Causes | 30 | 30 | 0 | 154.5 | 161.4 | 0.0 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 297 | 290 | 6 | 157 | 152 | 4 | 140 | 138 | 2 |
| Under 1 | 2 | 2 | 0 | 2 | 2 | 0 | 0 | 0 | 0 |
| (Neonatal) | 2 | 2 | 0 | 2 | 2 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 25-34 | 4 | 4 | 0 | 4 | 4 | 0 | 0 | 0 | 0 |
| 35-44 | 12 | 10 | 1 | 11 | 9 | 1 | 1 | 1 | 0 |
| 45-54 | 20 | 19 | 1 | 12 | 11 | 1 | 8 | 8 | 0 |
| 55-64 | 45 | 45 | 0 | 28 | 28 | 0 | 17 | 17 | 0 |
| 65-74 | 58 | 58 | 0 | 38 | 38 | 0 | 20 | 20 | 0 |
| 75-84 | 78 | 77 | 1 | 40 | 40 | 0 | 38 | 37 | 1 |
| 85 and over | 77 | 74 | 3 | 21 | 19 | 2 | 56 | 55 | 1 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 19,420 |
| White | 18,591 |
| Black | 547 |
| Other | 282 |

Selected Death Statistics, Residents of Tunica County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 97 | 37 | 60 | 915.3 | 1,573.1 | 749.2 |
| 1 | Heart Diseases | 34 | 17 | 17 | 320.8 | 722.8 | 212.3 |
| 2 | Malignant Neoplasms | 25 | 6 | 19 | 235.9 | 255.1 | 237.2 |
| 3 | Influenza and Pneumonia | 4 | 1 | 3 | 37.7 | 42.5 | 37.5 |
| 3 | Hypertension/Hypertensive Renal Disease | 4 | 1 | 3 | 37.7 | 42.5 | 37.5 |
| 3 | Diabetes Mellitus | 4 | 0 | 4 | 37.7 | 0.0 | 49.9 |
| 6 | Accidents | 3 | 1 | 2 | 28.3 | 42.5 | 25.0 |
| 6 | Emphysema and Other Chronic Lower Resp. Diseases | 3 | 2 | 1 | 28.3 | 85.0 | 12.5 |
| 6 | HIV Disease | 3 | 1 | 2 | 28.3 | 42.5 | 25.0 |
| 9 | Homicide | 2 | 0 | 2 | 18.9 | 0.0 | 25.0 |
| 9 | Nephritis, Nephrotic Syndrome And Nephrosis | 2 | 2 | 0 | 18.9 | 85.0 | 0.0 |
| 9 | Cerebrovascular Disease | 2 | 1 | 1 | 18.9 | 42.5 | 12.5 |
| 12 | Suicide | 1 | 1 | 0 | 9.4 | 42.5 | 0.0 |
| 12 | Chronic Liver Disease And Cirrhosis | 1 | 1 | 0 | 9.4 | 42.5 | 0.0 |
| 12 | Alzheimer's Disease | 1 | 0 | 1 | 9.4 | 0.0 | 12.5 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 2 | 0 | 2 | 18.9 | 0.0 | 25.0 |
| | All Other Causes | 6 | 3 | 3 | 56.6 | 127.6 | 37.5 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 97 | 37 | 60 | 45 | 18 | 27 | 52 | 19 | 33 |
| Under 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Neonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 3 | 0 | 3 | 2 | 0 | 2 | 1 | 0 | 1 |
| 25-34 | 2 | 0 | 2 | 0 | 0 | 0 | 2 | 0 | 2 |
| 35-44 | 3 | 0 | 3 | 0 | 0 | 0 | 3 | 0 | 3 |
| 45-54 | 11 | 4 | 7 | 3 | 1 | 2 | 8 | 3 | 5 |
| 55-64 | 18 | 8 | 10 | 12 | 6 | 6 | 6 | 2 | 4 |
| 65-74 | 21 | 10 | 11 | 13 | 6 | 7 | 8 | 4 | 4 |
| 75-84 | 21 | 6 | 15 | 11 | 4 | 7 | 10 | 2 | 8 |
| 85 and over | 18 | 9 | 9 | 4 | 1 | 3 | 14 | 8 | 6 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 10,598 |
| White | 2,352 |
| Black | 8,009 |
| Other | 237 |

Selected Death Statistics, Residents of Union County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 301 | 260 | 41 | 1,071.3 | 1,117.8 | 964.0 |
| 1 | Heart Diseases | 83 | 72 | 11 | 295.4 | 309.5 | 258.6 |
| 2 | Malignant Neoplasms | 63 | 51 | 12 | 224.2 | 219.3 | 282.2 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 25 | 22 | 3 | 89.0 | 94.6 | 70.5 |
| 4 | Cerebrovascular Disease | 22 | 17 | 5 | 78.3 | 73.1 | 117.6 |
| 5 | Accidents | 19 | 17 | 2 | 67.6 | 73.1 | 47.0 |
| 6 | Alzheimer's Disease | 16 | 15 | 1 | 56.9 | 64.5 | 23.5 |
| 7 | Influenza and Pneumonia | 9 | 9 | 0 | 32.0 | 38.7 | 0.0 |
| 8 | Nephritis, Nephrotic Syndrome And Nephrosis | 5 | 5 | 0 | 17.8 | 21.5 | 0.0 |
| 8 | Septicemia | 5 | 5 | 0 | 17.8 | 21.5 | 0.0 |
| 10 | Diabetes Mellitus | 4 | 3 | 1 | 14.2 | 12.9 | 23.5 |
| 11 | Suicide | 3 | 3 | 0 | 10.7 | 12.9 | 0.0 |
| 11 | Chronic Liver Disease And Cirrhosis | 3 | 3 | 0 | 10.7 | 12.9 | 0.0 |
| 13 | Congenital Malformations and Chrom. Abnormalities | 2 | 2 | 0 | 7.1 | 8.6 | 0.0 |
| 13 | Aortic Aneurysm and Dissection | 2 | 2 | 0 | 7.1 | 8.6 | 0.0 |
| 13 | Anemias | 2 | 2 | 0 | 7.1 | 8.6 | 0.0 |
| 13 | In Situ, Benign and Unspecified Neoplasms | 2 | 2 | 0 | 7.1 | 8.6 | 0.0 |
| 17 | Complications Medical/Surgical Care | 1 | 1 | 0 | 3.6 | 4.3 | 0.0 |
| 17 | Homicide | 1 | 0 | 1 | 3.6 | 0.0 | 23.5 |
| 17 | Pneumonitis Due to Solids and Liquids | 1 | 1 | 0 | 3.6 | 4.3 | 0.0 |
| 17 | Hypertension/Hypertensive Renal Disease | 1 | 1 | 0 | 3.6 | 4.3 | 0.0 |
| 17 | Parkinson's Disease | 1 | 1 | 0 | 3.6 | 4.3 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 5 | 4 | 1 | 17.8 | 17.2 | 23.5 |
| | All Other Causes | 26 | 22 | 4 | 92.5 | 94.6 | 94.1 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total* ----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|--------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 301 | 260 | 41 | 152 | 129 | 23 | 149 | 131 | 18 |
| Under 1 | 3 | 3 | 0 | 3 | 3 | 0 | 0 | 0 | 0 |
| (Neonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 3 | 3 | 0 | 3 | 3 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 15-24 | 2 | 0 | 2 | 1 | 0 | 1 | 1 | 0 | 1 |
| 25-34 | 9 | 9 | 0 | 8 | 8 | 0 | 1 | 1 | 0 |
| 35-44 | 8 | 7 | 1 | 4 | 3 | 1 | 4 | 4 | 0 |
| 45-54 | 21 | 17 | 4 | 10 | 8 | 2 | 11 | 9 | 2 |
| 55-64 | 30 | 26 | 4 | 21 | 18 | 3 | 9 | 8 | 1 |
| 65-74 | 66 | 58 | 8 | 35 | 31 | 4 | 31 | 27 | 4 |
| 75-84 | 88 | 72 | 16 | 51 | 41 | 10 | 37 | 31 | 6 |
| 85 and over | 73 | 68 | 5 | 18 | 17 | 1 | 55 | 51 | 4 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 28,097 |
| White | 23,260 |
| Black | 4,253 |
| Other | 584 |

Selected Death Statistics, Residents of Walthall County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 200 | 115 | 84 | 1,346.0 | 1,428.6 | 1,289.3 |
| 1 | Heart Diseases | 45 | 27 | 18 | 302.8 | 335.4 | 276.3 |
| 2 | Malignant Neoplasms | 37 | 26 | 11 | 249.0 | 323.0 | 168.8 |
| 3 | Accidents | 19 | 8 | 11 | 127.9 | 99.4 | 168.8 |
| 4 | Cerebrovascular Disease | 14 | 7 | 7 | 94.2 | 87.0 | 107.4 |
| 5 | Diabetes Mellitus | 10 | 3 | 7 | 67.3 | 37.3 | 107.4 |
| 6 | Emphysema and Other Chronic Lower Resp. Diseases | 9 | 6 | 3 | 60.6 | 74.5 | 46.0 |
| 7 | Influenza and Pneumonia | 8 | 7 | 0 | 53.8 | 87.0 | 0.0 |
| 8 | Alzheimer's Disease | 6 | 5 | 1 | 40.4 | 62.1 | 15.3 |
| 9 | Septicemia | 5 | 4 | 1 | 33.6 | 49.7 | 15.3 |
| 10 | Homicide | 4 | 1 | 3 | 26.9 | 12.4 | 46.0 |
| 10 | Nephritis, Nephrotic Syndrome And Nephrosis | 4 | 1 | 3 | 26.9 | 12.4 | 46.0 |
| 12 | Suicide | 3 | 2 | 1 | 20.2 | 24.8 | 15.3 |
| 13 | Pneumonitis Due to Solids and Liquids | 2 | 1 | 1 | 13.5 | 12.4 | 15.3 |
| 13 | Hypertension/Hypertensive Renal Disease | 2 | 2 | 0 | 13.5 | 24.8 | 0.0 |
| 15 | Complications Medical/Surgical Care | 1 | 0 | 1 | 6.7 | 0.0 | 15.3 |
| 15 | Congenital Malformations and Chrom. Abnormalities | 1 | 0 | 1 | 6.7 | 0.0 | 15.3 |
| 15 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 5.3 | 0.0 | 11.1 |
| 15 | Pneumoconioses and Chemical Effects | 1 | 1 | 0 | 6.7 | 12.4 | 0.0 |
| 15 | Aortic Aneurysm and Dissection | 1 | 1 | 0 | 6.7 | 12.4 | 0.0 |
| 15 | Anemias | 1 | 0 | 1 | 6.7 | 0.0 | 15.3 |
| 15 | In Situ, Benign and Unspecified Neoplasms | 1 | 0 | 1 | 6.7 | 0.0 | 15.3 |
| 15 | HIV Disease | 1 | 0 | 1 | 6.7 | 0.0 | 15.3 |
| 15 | Viral Hepatitis | 1 | 1 | 0 | 6.7 | 12.4 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 2 | 2 | 0 | 13.5 | 24.8 | 0.0 |
| | All Other Causes | 21 | 10 | 11 | 141.3 | 124.2 | 168.8 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 200 | 115 | 84 | 105 | 58 | 47 | 95 | 57 | 37 |
| Under 1 | 2 | 0 | 2 | 2 | 0 | 2 | 0 | 0 | 0 |
| (Neonatal) | 2 | 0 | 2 | 2 | 0 | 2 | 0 | 0 | 0 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| 10-14 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 15-24 | 4 | 1 | 3 | 4 | 1 | 3 | 0 | 0 | 0 |
| 25-34 | 6 | 3 | 3 | 5 | 2 | 3 | 1 | 1 | 0 |
| 35-44 | 3 | 1 | 2 | 1 | 1 | 0 | 2 | 0 | 2 |
| 45-54 | 18 | 8 | 10 | 10 | 3 | 7 | 8 | 5 | 3 |
| 55-64 | 30 | 15 | 15 | 20 | 9 | 11 | 10 | 6 | 4 |
| 65-74 | 47 | 27 | 20 | 29 | 18 | 11 | 18 | 9 | 9 |
| 75-84 | 44 | 32 | 11 | 21 | 15 | 6 | 23 | 17 | 5 |
| 85 and over | 44 | 27 | 17 | 12 | 8 | 4 | 32 | 19 | 13 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 14,859 |
| White | 8,050 |
| Black | 6,515 |
| Other | 294 |

Selected Death Statistics, Residents of Warren County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 509 | 299 | 204 | 1,060.8 | 1,255.9 | 881.8 |
| 1 | Heart Diseases | 144 | 78 | 63 | 300.1 | 327.6 | 272.3 |
| 2 | Malignant Neoplasms | 112 | 65 | 46 | 233.4 | 273.0 | 198.8 |
| 3 | Emphysema and Other Chronic Lower Resp. Diseases | 39 | 32 | 6 | 81.3 | 134.4 | 25.9 |
| 4 | Cerebrovascular Disease | 37 | 18 | 19 | 77.1 | 75.6 | 82.1 |
| 5 | Accidents | 20 | 11 | 9 | 41.7 | 46.2 | 38.9 |
| 6 | Nephritis, Nephrotic Syndrome And Nephrosis | 16 | 9 | 6 | 33.3 | 37.8 | 25.9 |
| 7 | Alzheimer's Disease | 14 | 13 | 1 | 29.2 | 54.6 | 4.3 |
| 8 | Diabetes Mellitus | 10 | 5 | 5 | 20.8 | 21.0 | 21.6 |
| 9 | Suicide | 9 | 8 | 1 | 18.8 | 33.6 | 4.3 |
| 9 | Influenza and Pneumonia | 9 | 5 | 4 | 18.8 | 21.0 | 17.3 |
| 11 | Hypertension/Hypertensive Renal Disease | 8 | 3 | 5 | 16.7 | 12.6 | 21.6 |
| 11 | Septicemia | 8 | 3 | 5 | 16.7 | 12.6 | 21.6 |
| 13 | Chronic Liver Disease And Cirrhosis | 6 | 6 | 0 | 12.5 | 25.2 | 0.0 |
| 14 | Parkinson's Disease | 5 | 2 | 3 | 10.4 | 8.4 | 13.0 |
| 15 | Homicide | 3 | 0 | 3 | 6.3 | 0.0 | 13.0 |
| 15 | Congenital Malformations and Chrom. Abnormalities | 3 | 1 | 2 | 6.3 | 4.2 | 8.6 |
| 15 | Pneumonitis Due to Solids and Liquids | 3 | 1 | 2 | 6.3 | 4.2 | 8.6 |
| 18 | Aortic Aneurysm and Dissection | 2 | 1 | 1 | 4.2 | 4.2 | 4.3 |
| 18 | In Situ, Benign and Unspecified Neoplasms | 2 | 2 | 0 | 4.2 | 8.4 | 0.0 |
| 20 | Complications Medical/Surgical Care | 1 | 1 | 0 | 2.1 | 4.2 | 0.0 |
| 20 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 1.6 | 0.0 | 2.8 |
| 20 | Peptic Ulcer | 1 | 0 | 1 | 2.1 | 0.0 | 4.3 |
| 20 | HIV Disease | 1 | 0 | 1 | 2.1 | 0.0 | 4.3 |
| 20 | Viral Hepatitis | 1 | 1 | 0 | 2.1 | 4.2 | 0.0 |
| 20 | Tuberculosis | 1 | 1 | 0 | 2.1 | 4.2 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 2 | 1 | 1 | 4.2 | 4.2 | 4.3 |
| | All Other Causes | 51 | 32 | 19 | 106.3 | 134.4 | 82.1 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 509 | 299 | 204 | 265 | 162 | 101 | 244 | 137 | 103 |
| Under 1 | 5 | 2 | 3 | 3 | 1 | 2 | 2 | 1 | 1 |
| (Neonatal) | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| (Postneonatal) | 4 | 2 | 2 | 2 | 1 | 1 | 2 | 1 | 1 |
| 1-4 | 2 | 1 | 1 | 1 | 1 | 0 | 1 | 0 | 1 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 15-24 | 3 | 1 | 2 | 1 | 0 | 1 | 2 | 1 | 1 |
| 25-34 | 5 | 2 | 3 | 2 | 1 | 1 | 3 | 1 | 2 |
| 35-44 | 15 | 10 | 5 | 9 | 7 | 2 | 6 | 3 | 3 |
| 45-54 | 41 | 23 | 18 | 20 | 14 | 6 | 21 | 9 | 12 |
| 55-64 | 98 | 41 | 54 | 53 | 24 | 29 | 45 | 17 | 25 |
| 65-74 | 109 | 61 | 48 | 71 | 39 | 32 | 38 | 22 | 16 |
| 75-84 | 111 | 78 | 31 | 57 | 39 | 16 | 54 | 39 | 15 |
| 85 and over | 119 | 80 | 38 | 47 | 36 | 11 | 72 | 44 | 27 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 47,983 |
| White | 23,808 |
| Black | 23,135 |
| Other | 1,040 |

Selected Death Statistics, Residents of Washington County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|--|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 589 | 226 | 359 | 1,203.1 | 1,711.6 | 1,026.5 |
| 1 | Malignant Neoplasms | 131 | 51 | 80 | 267.6 | 386.2 | 228.7 |
| 2 | Heart Diseases | 90 | 41 | 48 | 183.8 | 310.5 | 137.2 |
| 3 | Diabetes Mellitus | 53 | 18 | 35 | 108.3 | 136.3 | 100.1 |
| 4 | Cerebrovascular Disease | 37 | 12 | 25 | 75.6 | 90.9 | 71.5 |
| 5 | Emphysema and Other Chronic Lower Resp. Diseases | 30 | 20 | 10 | 61.3 | 151.5 | 28.6 |
| 6 | Accidents | 23 | 11 | 12 | 47.0 | 83.3 | 34.3 |
| 6 | Nephritis, Nephrotic Syndrome And Nephrosis | 23 | 6 | 17 | 47.0 | 45.4 | 48.6 |
| 8 | Hypertension/Hypertensive Renal Disease | 20 | 6 | 14 | 40.9 | 45.4 | 40.0 |
| 9 | Homicide | 10 | 0 | 10 | 20.4 | 0.0 | 28.6 |
| 10 | Alzheimer's Disease | 9 | 3 | 6 | 18.4 | 22.7 | 17.2 |
| 11 | Septicemia | 8 | 1 | 6 | 16.3 | 7.6 | 17.2 |
| 12 | Influenza and Pneumonia | 5 | 0 | 5 | 10.2 | 0.0 | 14.3 |
| 12 | HIV Disease | 5 | 0 | 5 | 10.2 | 0.0 | 14.3 |
| 14 | Chronic Liver Disease And Cirrhosis | 4 | 0 | 4 | 8.2 | 0.0 | 11.4 |
| 15 | Aortic Aneurysm and Dissection | 3 | 1 | 2 | 6.1 | 7.6 | 5.7 |
| 15 | Parkinson's Disease | 3 | 3 | 0 | 6.1 | 22.7 | 0.0 |
| 17 | Suicide | 2 | 1 | 1 | 4.1 | 7.6 | 2.9 |
| 17 | Hyperplasia of Prostate | 2 | 1 | 1 | 8.7 | 15.3 | 6.3 |
| 19 | Congenital Malformations and Chrom. Abnormalities | 1 | 1 | 0 | 2.0 | 7.6 | 0.0 |
| 19 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 1.3 | 0.0 | 1.6 |
| 19 | Cholelithiasis/Gallbladder Disorder | 1 | 0 | 1 | 2.0 | 0.0 | 2.9 |
| 19 | Anemias | 1 | 0 | 1 | 2.0 | 0.0 | 2.9 |
| 19 | In Situ, Benign and Unspecified Neoplasms | 1 | 0 | 1 | 2.0 | 0.0 | 2.9 |
| 19 | Viral Hepatitis | 1 | 0 | 1 | 2.0 | 0.0 | 2.9 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 12 | 7 | 5 | 24.5 | 53.0 | 14.3 |
| | All Other Causes | 113 | 43 | 68 | 230.8 | 325.7 | 194.4 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births.
 Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 589 | 226 | 359 | 306 | 118 | 185 | 283 | 108 | 174 |
| Under 1 | 3 | 1 | 2 | 1 | 1 | 0 | 2 | 0 | 2 |
| (Neonatal) | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 2 | 0 | 2 | 0 | 0 | 0 | 2 | 0 | 2 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 10 | 3 | 7 | 7 | 1 | 6 | 3 | 2 | 1 |
| 25-34 | 14 | 2 | 12 | 5 | 0 | 5 | 9 | 2 | 7 |
| 35-44 | 19 | 6 | 13 | 10 | 3 | 7 | 9 | 3 | 6 |
| 45-54 | 38 | 11 | 27 | 18 | 7 | 11 | 20 | 4 | 16 |
| 55-64 | 118 | 31 | 86 | 82 | 21 | 60 | 36 | 10 | 26 |
| 65-74 | 131 | 55 | 76 | 77 | 35 | 42 | 54 | 20 | 34 |
| 75-84 | 129 | 60 | 68 | 63 | 29 | 34 | 66 | 31 | 34 |
| 85 and over | 126 | 57 | 67 | 43 | 21 | 20 | 83 | 36 | 47 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 48,958 |
| White | 13,204 |
| Black | 34,974 |
| Other | 780 |

Selected Death Statistics, Residents of Wayne County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 218 | 151 | 67 | 1,063.9 | 1,245.5 | 829.3 |
| 1 | Heart Diseases | 70 | 51 | 19 | 341.6 | 420.7 | 235.2 |
| 2 | Malignant Neoplasms | 47 | 32 | 15 | 229.4 | 263.9 | 185.7 |
| 3 | Accidents | 14 | 10 | 4 | 68.3 | 82.5 | 49.5 |
| 4 | Emphysema and Other Chronic Lower Resp. Diseases | 13 | 11 | 2 | 63.4 | 90.7 | 24.8 |
| 5 | Cerebrovascular Disease | 9 | 6 | 3 | 43.9 | 49.5 | 37.1 |
| 6 | Homicide | 4 | 2 | 2 | 19.5 | 16.5 | 24.8 |
| 6 | Influenza and Pneumonia | 4 | 4 | 0 | 19.5 | 33.0 | 0.0 |
| 8 | Alzheimer's Disease | 3 | 2 | 1 | 14.6 | 16.5 | 12.4 |
| 8 | Diabetes Mellitus | 3 | 3 | 0 | 14.6 | 24.7 | 0.0 |
| 10 | Suicide | 2 | 2 | 0 | 9.8 | 16.5 | 0.0 |
| 10 | Nephritis, Nephrotic Syndrome And Nephrosis | 2 | 2 | 0 | 9.8 | 16.5 | 0.0 |
| 10 | Chronic Liver Disease And Cirrhosis | 2 | 1 | 1 | 9.8 | 8.2 | 12.4 |
| 10 | Hypertension/Hypertensive Renal Disease | 2 | 2 | 0 | 9.8 | 16.5 | 0.0 |
| 10 | Septicemia | 2 | 1 | 1 | 9.8 | 8.2 | 12.4 |
| 15 | Complications Medical/Surgical Care | 1 | 0 | 1 | 4.9 | 0.0 | 12.4 |
| 15 | Congenital Malformations and Chrom. Abnormalities | 1 | 0 | 1 | 4.9 | 0.0 | 12.4 |
| 15 | Peptic Ulcer | 1 | 0 | 1 | 4.9 | 0.0 | 12.4 |
| 15 | Pneumonitis Due to Solids and Liquids | 1 | 1 | 0 | 4.9 | 8.2 | 0.0 |
| 15 | Aortic Aneurysm and Dissection | 1 | 1 | 0 | 4.9 | 8.2 | 0.0 |
| 15 | Atherosclerosis | 1 | 0 | 1 | 4.9 | 0.0 | 12.4 |
| 15 | Parkinson's Disease | 1 | 1 | 0 | 4.9 | 8.2 | 0.0 |
| 15 | Meningitis | 1 | 0 | 1 | 4.9 | 0.0 | 12.4 |
| 15 | HIV Disease | 1 | 0 | 1 | 4.9 | 0.0 | 12.4 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 1 | 1 | 0 | 4.9 | 8.2 | 0.0 |
| | All Other Causes | 31 | 18 | 13 | 151.3 | 148.5 | 160.9 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 218 | 151 | 67 | 114 | 79 | 35 | 104 | 72 | 32 |
| Under 1 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| (Neonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 6 | 1 | 5 | 5 | 1 | 4 | 1 | 0 | 1 |
| 25-34 | 4 | 2 | 2 | 4 | 2 | 2 | 0 | 0 | 0 |
| 35-44 | 13 | 7 | 6 | 4 | 2 | 2 | 9 | 5 | 4 |
| 45-54 | 18 | 7 | 11 | 13 | 6 | 7 | 5 | 1 | 4 |
| 55-64 | 38 | 21 | 17 | 19 | 13 | 6 | 19 | 8 | 11 |
| 65-74 | 44 | 36 | 8 | 24 | 19 | 5 | 20 | 17 | 3 |
| 75-84 | 51 | 40 | 11 | 30 | 23 | 7 | 21 | 17 | 4 |
| 85 and over | 43 | 37 | 6 | 15 | 13 | 2 | 28 | 24 | 4 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 20,490 |
| White | 12,124 |
| Black | 8,079 |
| Other | 287 |

Selected Death Statistics, Residents of Webster County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 139 | 113 | 25 | 1,393.9 | 1,440.0 | 1,271.0 |
| 1 | Heart Diseases | 39 | 32 | 6 | 391.1 | 407.8 | 305.0 |
| 2 | Malignant Neoplasms | 35 | 27 | 8 | 351.0 | 344.1 | 406.7 |
| 3 | Alzheimer's Disease | 10 | 6 | 4 | 100.3 | 76.5 | 203.4 |
| 4 | Accidents | 7 | 7 | 0 | 70.2 | 89.2 | 0.0 |
| 5 | Septicemia | 6 | 5 | 1 | 60.2 | 63.7 | 50.8 |
| 6 | Nephritis, Nephrotic Syndrome And Nephrosis | 5 | 5 | 0 | 50.1 | 63.7 | 0.0 |
| 7 | Emphysema and Other Chronic Lower Resp. Diseases | 4 | 4 | 0 | 40.1 | 51.0 | 0.0 |
| 7 | Influenza and Pneumonia | 4 | 4 | 0 | 40.1 | 51.0 | 0.0 |
| 9 | Cerebrovascular Disease | 3 | 2 | 1 | 30.1 | 25.5 | 50.8 |
| 10 | Hypertension/Hypertensive Renal Disease | 2 | 1 | 1 | 20.1 | 12.7 | 50.8 |
| 10 | Diabetes Mellitus | 2 | 1 | 1 | 20.1 | 12.7 | 50.8 |
| 12 | Complications Medical/Surgical Care | 1 | 1 | 0 | 10.0 | 12.7 | 0.0 |
| 12 | Congenital Malformations and Chrom. Abnormalities | 1 | 1 | 0 | 10.0 | 12.7 | 0.0 |
| 12 | Pneumonitis Due to Solids and Liquids | 1 | 1 | 0 | 10.0 | 12.7 | 0.0 |
| 12 | Meningitis | 1 | 1 | 0 | 10.0 | 12.7 | 0.0 |
| 12 | Malnutrition/Nutritional Deficiency | 1 | 1 | 0 | 10.0 | 12.7 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 1 | 1 | 0 | 10.0 | 12.7 | 0.0 |
| | All Other Causes | 16 | 13 | 3 | 160.4 | 165.7 | 152.5 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 139 | 113 | 25 | 73 | 60 | 12 | 66 | 53 | 13 |
| Under 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Neonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 25-34 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 35-44 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 45-54 | 12 | 11 | 1 | 5 | 5 | 0 | 7 | 6 | 1 |
| 55-64 | 19 | 13 | 6 | 10 | 6 | 4 | 9 | 7 | 2 |
| 65-74 | 27 | 24 | 3 | 20 | 17 | 3 | 7 | 7 | 0 |
| 75-84 | 44 | 35 | 9 | 24 | 20 | 4 | 20 | 15 | 5 |
| 85 and over | 35 | 28 | 6 | 12 | 10 | 1 | 23 | 18 | 5 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|-------|
| Total | 9,972 |
| White | 7,847 |
| Black | 1,967 |
| Other | 158 |

Selected Death Statistics, Residents of Wilkinson County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 127 | 33 | 94 | 1,381.8 | 1,241.1 | 1,457.8 |
| 1 | Heart Diseases | 33 | 7 | 26 | 359.0 | 263.3 | 403.2 |
| 2 | Malignant Neoplasms | 25 | 7 | 18 | 272.0 | 263.3 | 279.2 |
| 3 | Accidents | 10 | 4 | 6 | 108.8 | 150.4 | 93.1 |
| 4 | Alzheimer's Disease | 7 | 3 | 4 | 76.2 | 112.8 | 62.0 |
| 5 | Emphysema and Other Chronic Lower Resp. Diseases | 6 | 1 | 5 | 65.3 | 37.6 | 77.5 |
| 5 | Diabetes Mellitus | 6 | 1 | 5 | 65.3 | 37.6 | 77.5 |
| 7 | Influenza and Pneumonia | 5 | 2 | 3 | 54.4 | 75.2 | 46.5 |
| 8 | Nephritis, Nephrotic Syndrome And Nephrosis | 4 | 1 | 3 | 43.5 | 37.6 | 46.5 |
| 8 | Cerebrovascular Disease | 4 | 3 | 1 | 43.5 | 112.8 | 15.5 |
| 10 | Homicide | 3 | 0 | 3 | 32.6 | 0.0 | 46.5 |
| 11 | Septicemia | 2 | 0 | 2 | 21.8 | 0.0 | 31.0 |
| 12 | Complications Medical/Surgical Care | 1 | 0 | 1 | 10.9 | 0.0 | 15.5 |
| 12 | Conditions Originating in the Perinatal Period** | 1 | 0 | 1 | 9.4 | 0.0 | 13.2 |
| 12 | Pneumonitis Due to Solids and Liquids | 1 | 0 | 1 | 10.9 | 0.0 | 15.5 |
| 12 | Parkinson's Disease | 1 | 1 | 0 | 10.9 | 37.6 | 0.0 |
| 12 | HIV Disease | 1 | 0 | 1 | 10.9 | 0.0 | 15.5 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 0 | 0 | 0 | 0.0 | 0.0 | 0.0 |
| | All Other Causes | 17 | 3 | 14 | 185.0 | 112.8 | 217.1 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 127 | 33 | 94 | 73 | 14 | 59 | 54 | 19 | 35 |
| Under 1 | 2 | 0 | 2 | 1 | 0 | 1 | 1 | 0 | 1 |
| (Neonatal) | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| (Postneonatal) | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 4 | 1 | 3 | 4 | 1 | 3 | 0 | 0 | 0 |
| 25-34 | 4 | 0 | 4 | 4 | 0 | 4 | 0 | 0 | 0 |
| 35-44 | 2 | 0 | 2 | 2 | 0 | 2 | 0 | 0 | 0 |
| 45-54 | 11 | 2 | 9 | 9 | 1 | 8 | 2 | 1 | 1 |
| 55-64 | 19 | 3 | 16 | 11 | 2 | 9 | 8 | 1 | 7 |
| 65-74 | 25 | 7 | 18 | 14 | 2 | 12 | 11 | 5 | 6 |
| 75-84 | 31 | 10 | 21 | 21 | 5 | 16 | 10 | 5 | 5 |
| 85 and over | 29 | 10 | 19 | 7 | 3 | 4 | 22 | 7 | 15 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|-------|
| Total | 9,191 |
| White | 2,659 |
| Black | 6,448 |
| Other | 84 |

Selected Death Statistics, Residents of Winston County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 242 | 151 | 90 | 1,309.7 | 1,588.3 | 1,051.9 |
| 1 | Heart Diseases | 64 | 46 | 18 | 346.4 | 483.9 | 210.4 |
| 2 | Malignant Neoplasms | 36 | 19 | 17 | 194.8 | 199.9 | 198.7 |
| 3 | Accidents | 26 | 16 | 9 | 140.7 | 168.3 | 105.2 |
| 4 | Cerebrovascular Disease | 18 | 14 | 4 | 97.4 | 147.3 | 46.8 |
| 5 | Emphysema and Other Chronic Lower Resp. Diseases | 12 | 10 | 2 | 64.9 | 105.2 | 23.4 |
| 6 | Alzheimer's Disease | 10 | 4 | 6 | 54.1 | 42.1 | 70.1 |
| 7 | Suicide | 8 | 7 | 1 | 43.3 | 73.6 | 11.7 |
| 7 | Influenza and Pneumonia | 8 | 4 | 4 | 43.3 | 42.1 | 46.8 |
| 7 | Diabetes Mellitus | 8 | 5 | 3 | 43.3 | 52.6 | 35.1 |
| 10 | Nephritis, Nephrotic Syndrome And Nephrosis | 5 | 2 | 3 | 27.1 | 21.0 | 35.1 |
| 10 | Chronic Liver Disease And Cirrhosis | 5 | 2 | 3 | 27.1 | 21.0 | 35.1 |
| 12 | Conditions Originating in the Perinatal Period** | 4 | 0 | 4 | 19.8 | 0.0 | 37.0 |
| 12 | Septicemia | 4 | 2 | 2 | 21.6 | 21.0 | 23.4 |
| 14 | Pneumonitis Due to Solids and Liquids | 2 | 1 | 1 | 10.8 | 10.5 | 11.7 |
| 15 | Homicide | 1 | 0 | 1 | 5.4 | 0.0 | 11.7 |
| 15 | Peptic Ulcer | 1 | 1 | 0 | 5.4 | 10.5 | 0.0 |
| 15 | Aortic Aneurysm and Dissection | 1 | 1 | 0 | 5.4 | 10.5 | 0.0 |
| 15 | Hypertension/Hypertensive Renal Disease | 1 | 0 | 1 | 5.4 | 0.0 | 11.7 |
| 15 | Meningitis | 1 | 1 | 0 | 5.4 | 10.5 | 0.0 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 5 | 3 | 2 | 27.1 | 31.6 | 23.4 |
| | All Other Causes | 22 | 13 | 9 | 119.1 | 136.7 | 105.2 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 242 | 151 | 90 | 113 | 74 | 39 | 129 | 77 | 51 |
| Under 1 | 5 | 1 | 4 | 0 | 0 | 0 | 5 | 1 | 4 |
| (Neonatal) | 4 | 0 | 4 | 0 | 0 | 0 | 4 | 0 | 4 |
| (Postneonatal) | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 |
| 25-34 | 7 | 3 | 4 | 3 | 1 | 2 | 4 | 2 | 2 |
| 35-44 | 10 | 6 | 4 | 6 | 4 | 2 | 4 | 2 | 2 |
| 45-54 | 22 | 12 | 10 | 12 | 7 | 5 | 10 | 5 | 5 |
| 55-64 | 39 | 21 | 18 | 20 | 11 | 9 | 19 | 10 | 9 |
| 65-74 | 39 | 23 | 15 | 28 | 17 | 11 | 11 | 6 | 4 |
| 75-84 | 59 | 40 | 19 | 30 | 22 | 8 | 29 | 18 | 11 |
| 85 and over | 59 | 44 | 15 | 12 | 11 | 1 | 47 | 33 | 14 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 18,478 |
| White | 9,507 |
| Black | 8,556 |
| Other | 415 |

Selected Death Statistics, Residents of Yalobusha County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|-------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 163 | 117 | 46 | 1,327.8 | 1,593.6 | 973.5 |
| 1 | Heart Diseases | 47 | 34 | 13 | 382.9 | 463.1 | 275.1 |
| 2 | Malignant Neoplasms | 37 | 26 | 11 | 301.4 | 354.1 | 232.8 |
| 3 | Nephritis, Nephrotic Syndrome And Nephrosis | 10 | 7 | 3 | 81.5 | 95.3 | 63.5 |
| 4 | Accidents | 8 | 4 | 4 | 65.2 | 54.5 | 84.7 |
| 5 | Emphysema and Other Chronic Lower Resp. Diseases | 6 | 5 | 1 | 48.9 | 68.1 | 21.2 |
| 5 | Influenza and Pneumonia | 6 | 5 | 1 | 48.9 | 68.1 | 21.2 |
| 5 | Alzheimer's Disease | 6 | 6 | 0 | 48.9 | 81.7 | 0.0 |
| 8 | Cerebrovascular Disease | 5 | 2 | 3 | 40.7 | 27.2 | 63.5 |
| 9 | Homicide | 2 | 0 | 2 | 16.3 | 0.0 | 42.3 |
| 9 | Chronic Liver Disease And Cirrhosis | 2 | 2 | 0 | 16.3 | 27.2 | 0.0 |
| 9 | Hypertension/Hypertensive Renal Disease | 2 | 1 | 1 | 16.3 | 13.6 | 21.2 |
| 9 | Septicemia | 2 | 1 | 1 | 16.3 | 13.6 | 21.2 |
| 13 | Suicide | 1 | 1 | 0 | 8.1 | 13.6 | 0.0 |
| 13 | Pneumonitis Due to Solids and Liquids | 1 | 1 | 0 | 8.1 | 13.6 | 0.0 |
| 13 | Diabetes Mellitus | 1 | 1 | 0 | 8.1 | 13.6 | 0.0 |
| 13 | Anemias | 1 | 1 | 0 | 8.1 | 13.6 | 0.0 |
| 13 | In Situ, Benign and Unspecified Neoplasms | 1 | 1 | 0 | 8.1 | 13.6 | 0.0 |
| 13 | HIV Disease | 1 | 0 | 1 | 8.1 | 0.0 | 21.2 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 0 | 0 | 0 | 0.0 | 0.0 | 0.0 |
| | All Other Causes | 24 | 19 | 5 | 195.5 | 258.8 | 105.8 |

* Total includes races other than white and black.

** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.

Less than 0.05.

| Age | ----- Total* ----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|--------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 163 | 117 | 46 | 81 | 56 | 25 | 82 | 61 | 21 |
| Under 1 | 2 | 0 | 2 | 1 | 0 | 1 | 1 | 0 | 1 |
| (Neonatal) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (Postneonatal) | 2 | 0 | 2 | 1 | 0 | 1 | 1 | 0 | 1 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 25-34 | 4 | 1 | 3 | 2 | 0 | 2 | 2 | 1 | 1 |
| 35-44 | 11 | 5 | 6 | 5 | 2 | 3 | 6 | 3 | 3 |
| 45-54 | 7 | 3 | 4 | 5 | 1 | 4 | 2 | 2 | 0 |
| 55-64 | 21 | 17 | 4 | 14 | 10 | 4 | 7 | 7 | 0 |
| 65-74 | 32 | 21 | 11 | 17 | 11 | 6 | 15 | 10 | 5 |
| 75-84 | 45 | 38 | 7 | 23 | 21 | 2 | 22 | 17 | 5 |
| 85 and over | 39 | 30 | 9 | 12 | 9 | 3 | 27 | 21 | 6 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| 2014 Population | |
|-----------------|--------|
| Total | 12,276 |
| White | 7,342 |
| Black | 4,725 |
| Other | 209 |

Selected Death Statistics, Residents of Yazoo County, 2014

| Rank | Leading Causes of Death Listed in Descending Order | -----Number----- | | | ----Mortality Rate**---- | | |
|------|---|------------------|-------|-------|--------------------------|---------|---------|
| | | Total* | White | Black | Total* | White | Black |
| | Total Of All Causes | 304 | 138 | 165 | 1,092.9 | 1,216.9 | 1,034.2 |
| 1 | Heart Diseases | 70 | 42 | 28 | 251.6 | 370.4 | 175.5 |
| 2 | Malignant Neoplasms | 58 | 23 | 35 | 208.5 | 202.8 | 219.4 |
| 3 | Diabetes Mellitus | 37 | 12 | 25 | 133.0 | 105.8 | 156.7 |
| 4 | Accidents | 19 | 7 | 12 | 68.3 | 61.7 | 75.2 |
| 5 | Cerebrovascular Disease | 18 | 7 | 10 | 64.7 | 61.7 | 62.7 |
| 6 | Emphysema and Other Chronic Lower Resp. Diseases | 15 | 9 | 6 | 53.9 | 79.4 | 37.6 |
| 7 | Septicemia | 12 | 8 | 4 | 43.1 | 70.5 | 25.1 |
| 8 | Homicide | 6 | 1 | 5 | 21.6 | 8.8 | 31.3 |
| 8 | Nephritis, Nephrotic Syndrome And Nephrosis | 6 | 2 | 4 | 21.6 | 17.6 | 25.1 |
| 8 | Alzheimer's Disease | 6 | 2 | 4 | 21.6 | 17.6 | 25.1 |
| 11 | Hypertension/Hypertensive Renal Disease | 3 | 0 | 3 | 10.8 | 0.0 | 18.8 |
| 11 | HIV Disease | 3 | 1 | 2 | 10.8 | 8.8 | 12.5 |
| 13 | Suicide | 2 | 1 | 1 | 7.2 | 8.8 | 6.3 |
| 13 | Congenital Malformations and Chrom. Abnormalities | 2 | 1 | 1 | 7.2 | 8.8 | 6.3 |
| 13 | Conditions Originating in the Perinatal Period** | 2 | 1 | 1 | 5.4 | 10.1 | 3.7 |
| 13 | Peptic Ulcer | 2 | 1 | 1 | 7.2 | 8.8 | 6.3 |
| 13 | Influenza and Pneumonia | 2 | 2 | 0 | 7.2 | 17.6 | 0.0 |
| 13 | Anemias | 2 | 0 | 2 | 7.2 | 0.0 | 12.5 |
| 19 | Pneumonitis Due to Solids and Liquids | 1 | 1 | 0 | 3.6 | 8.8 | 0.0 |
| 19 | Aortic Aneurysm and Dissection | 1 | 1 | 0 | 3.6 | 8.8 | 0.0 |
| 19 | Parkinson's Disease | 1 | 1 | 0 | 3.6 | 8.8 | 0.0 |
| 19 | In Situ, Benign and Unspecified Neoplasms | 1 | 1 | 0 | 3.6 | 8.8 | 0.0 |
| 19 | Viral Hepatitis | 1 | 0 | 1 | 3.6 | 0.0 | 6.3 |
| | --- Not Included in Ranking --- | | | | | | |
| | Symptoms, Signs, Ill-Defined Conditions | 2 | 0 | 2 | 7.2 | 0.0 | 12.5 |
| | All Other Causes | 32 | 14 | 18 | 115.0 | 123.5 | 112.8 |

* Total includes races other than white and black.
 ** Deaths per 100,000 population, except deaths from maternal causes are per 100,000 live births and deaths from conditions originating in the perinatal period are per 1,000 live births. Population figures are 2014 estimates.
 # Less than 0.05.

| Age | ----- Total*----- | | | ----- Male ----- | | | ----- Female ----- | | |
|----------------|-------------------|-------|-------|------------------|-------|-------|--------------------|-------|-------|
| | Total* | White | Black | Total* | White | Black | Total* | White | Black |
| Total | 304 | 138 | 165 | 148 | 65 | 82 | 156 | 73 | 83 |
| Under 1 | 3 | 1 | 1 | 1 | 0 | 0 | 2 | 1 | 1 |
| (Neonatal) | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| (Postneonatal) | 2 | 1 | 0 | 1 | 0 | 0 | 1 | 1 | 0 |
| 1-4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5-9 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-24 | 8 | 1 | 7 | 4 | 1 | 3 | 4 | 0 | 4 |
| 25-34 | 11 | 2 | 9 | 9 | 1 | 8 | 2 | 1 | 1 |
| 35-44 | 11 | 4 | 7 | 6 | 1 | 5 | 5 | 3 | 2 |
| 45-54 | 28 | 9 | 19 | 14 | 7 | 7 | 14 | 2 | 12 |
| 55-64 | 61 | 22 | 39 | 34 | 12 | 22 | 27 | 10 | 17 |
| 65-74 | 65 | 34 | 31 | 37 | 21 | 16 | 28 | 13 | 15 |
| 75-84 | 57 | 30 | 27 | 26 | 11 | 15 | 31 | 19 | 12 |
| 85 and over | 59 | 35 | 24 | 17 | 11 | 6 | 42 | 24 | 18 |
| Unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2014 Population

| | |
|-------|--------|
| Total | 27,817 |
| White | 11,340 |
| Black | 15,954 |
| Other | 523 |