

ANNUAL BULLETIN
OF
VITAL STATISTICS

1956

VOLUME I
STATE SUMMARY

MISSISSIPPI STATE BOARD OF HEALTH

ANNUAL BULLETIN

of

VITAL STATISTICS

1956

VOLUME 1

STATE SUMMARY

MISSISSIPPI STATE BOARD OF HEALTH

VOLUME 1
TABLE OF CONTENTS

	Page
General Summary	1
Table	
1. Estimated Population as of July 1, 1956, by County, by Race	21
2. Live Births, by County and City of Occurrence, Reallocated to Mother's Usual Residence, by Race, with Rates	23
3. Live Births, by County and City of Usual Residence of Mother, by Race and Attendant, and by Sex	26
4. Fetal Deaths, by County and City of Occurrence, Reallocated to Mother's Usual Residence, by Race and Attendant, and by Sex	30
5. Deaths, by Cause, by Race	34
6. Deaths, by County and City of Occurrence, Reallocated to Place of Residence, by Race, with Rates	45
7. Deaths from Selected Causes, by Race, Sex, and Age, with Rates	48
8. Deaths from Selected Causes, by Month of Occurrence	57
9. Deaths Under 1 Year, shown for Selected Causes, by Race and Age, with Rates	58
10. Deaths Due to Injuries which Occurred in Mississippi, by County of Injury, by Type of Injury	60
11. Marriages, by County of Occurrence and of Residence of Bride, and Divorces, by County of Occurrence, by Race.....	63
12. Divorces, by Cause, by Number of Minor Children, by Number of Years Married, by Race	65
13. Births, Deaths, and Marriages, by Month of Occurrence	66
14. Live Births, Deaths, Maternal Deaths, Deaths Under 1 Year of Age, and Deaths Under 1 Month of Age, 1947-1956, by Year, by Race, with Rates ...	67
15. Deaths from Heart Diseases, Nephritis (all forms), Vascular Lesions, Cancer, and Tuberculosis (all forms), 1947-1956, by Year, by Race, with Rates	68
16. Births, Fetal Deaths, Deaths Under 1 Year, Deaths Under 1 Month, and Deaths from Certain Selected Causes, on Basis of Place of Occurrence and Place of Residence, by Race, with Rates	69

SUMMARY OF 1956 VITAL STATISTICS

INTRODUCTION

The 1956 Vital Statistics Bulletin is the fourteenth consecutive issue prepared by the Public Health Statistical Service for the Division of Vital Statistics from data obtained from records of live births, fetal deaths, deaths, marriages, and divorces collected through the Division of Vital Statistics.

Volume 1 contains an analysis of the data and tables for the State, the 82 counties and in some instances the 15 cities with population of 10,000 or more. Summary tables for each county are included in Volume 2.

Population and Rates

Population figures used in this bulletin are estimates as of July 1, 1956, prepared by Dr. Harald A. Pedersen, formerly of the Sociology Department, Mississippi State College, according to Method #II described by the United States Bureau of the Census. Some changes were made by the Public Health Statistical Service to include military population and also to adjust the estimates by race to the total population of each county. Later adjustments by Dr. Pedersen equating the county totals to a recent state estimate by the Census Bureau were not incorporated into the population figures used in this volume.

No age or sex-specific rates (except for maternal and infant deaths) and no city rates were calculated for 1956 because current estimates by age, sex, and for cities are not available. Since rates for 1956 are based on the estimated population as of July 1, and those for 1955 were based on the now out-of-date 1950 census population, comparison of the rates for these two years should be made with caution. Some of the differences may be accounted for entirely by the change in population base.

Occurrence and Residence Allocation

Births, fetal deaths, deaths, and marriages were classified by occurrence and residence. Occurrence data were obtained from certificates filed in Mississippi at the place of birth, fetal death, death, or issuance of marriage license. Residence data include transcripts of these events (except marriages) occurring to Mississippi residents outside the State, and exclude records filed in Mississippi for nonresidents. There were 964 births to residents of Mississippi which occurred outside the State in 1956 and were reported to the Division of Vital Statistics; births occurring inside Mississippi to residents of other states numbered 1,592. Fetal death transcripts totaled only 19 and the nonresident certificates 62. During the year, 761 death transcripts were received and 631 nonresident deaths recorded. The total number of states sending transcripts was 36 including the District of Columbia. Birth records came from 29 states, fetal death records from 5 states, and death records from 34 states. Seventeen reports on Mississippi residents came from Alaska and 4 from foreign countries. The few transcripts received after completion of the tabulations were not included in these figures. In Table A the states furnishing the largest numbers of records are listed.

Some of the tables contain both residence and occurrence data; but, following recommended procedures, most of the tables and all of the rates are based on residence data only. Residence assignment for births and fetal deaths was determined on the basis of usual residence of mother, and for deaths, on the basis of usual residence of decedent. Deaths of inmates or employees of institutions were reallocated to previous residence regardless of length of stay in the institution.

Table A. Transcripts received, by place of occurrence

Place of Occurrence	Live Births	Deaths	Fetal Deaths
<u>Total</u>	<u>964</u>	<u>761</u>	<u>19</u>
Alabama	181	74	2
Arkansas	17	17	0
Florida	23	13	0
Illinois	20	30	0
Kentucky	30	7	0
Louisiana	134	168	5
Tennessee	362	347	10
Texas	21	26	0
Other places	176	79	2

Urban-Rural Classification

Towns with populations of 2,500 or more according to the 1950 census were considered as urban areas. All other areas were classified as rural.

Classification of Causes of Death

Causes of death were assigned according to the Sixth Revision (1948) of the International Statistical Classification of Causes of Death. Only underlying causes were coded. Changes in coding procedures since 1949 have been minor, but comparisons with earlier years should be made with caution, since revision of the code was responsible for several shifts in the distribution of deaths by cause.

LIVE BIRTHS

Resident Births

Live births to residents of Mississippi numbered 62,178 in 1956. This total was 1,066 less than that for 1955 and was the lowest since 1946. Declines of 268 for the whites and of 798 for the nonwhites were noted. The crude birth rate of 29.5 per 1,000 population was slightly higher than that for 1955. However, as mentioned in the section on population, rates for 1955 and 1956 are not entirely comparable. The state rate for the whites in 1956 was 23.8 and county rates varied from 12.1 in Issaquena to 32.0 in Harrison. For nonwhites the state rate was 36.2 and the range for the counties was from 25.7 in Tishomingo to 53.8 in Smith.

Characteristics

A breakdown of the 34,900 nonwhite births by specific race shows 34,731 were Negro, 126 were Indian, 33 were Chinese, 9 were Japanese, and one was another nonwhite race. Births of males numbered 31,685 and there were 30,493 females. These totals yielded a sex ratio of 1,039 males to 1,000 females. August was the peak month of occurrence for live births. There were 5,701 in this month in comparison with only 4,468 in April when the fewest occurred.

During 1956, 19,753 children were born to urban residents and 42,425 to rural residents. Over half (54 per cent) of the urban resident births were white, but 61 per cent of the rural were nonwhite. Comparison of the occurrence and residence

figures indicates that well over one-half of the white mothers but only a small percentage of the nonwhite mothers in rural areas went to cities for delivery.

The number of illegitimate births during 1956 was 7,791 or 12.5 per cent of the total. This percentage is the same as that reported in 1955, but in the period 1947-1955, for which comparable figures are available, there had been a steady increase from 9.1 to 12.5 per cent. White births classed as illegitimate totaled 294 or 1.1 per cent, and 7,497 or 21.5 per cent of the nonwhite births were illegitimate.

Liveborn infants classified as immature (birth weight $5\frac{1}{2}$ pounds or less) numbered 5,377 in 1956, or 8.6 per cent of the total. The incidence of immaturity was greater among nonwhites than among whites, 10.1 per cent in comparison with 6.6 per cent.

Classification by Attendant

In Table B which gives the distribution of live births by attendant and race, it may be seen that, in 1956, nearly 66 per cent of all births occurred in hospitals or clinics (including physicians' offices or clinics which provide in-patient care). A small proportion, 6.5 per cent, were delivered by physicians in places other than hospitals or clinics, and 27.8 per cent were delivered at home by nonmedical attendants. Almost half of the nonwhite mothers, but less than one per cent of the white mothers had no medical attendant.

Table B. Live births by attendant and race

Attendant	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
<u>Total</u>	<u>62,178</u>	<u>100.0</u>	<u>27,278</u>	<u>100.0</u>	<u>34,900</u>	<u>100.0</u>
Physician in hospital	40,832	65.7	26,510	97.2	14,322	41.0
Physician in home	4,066	6.5	574	2.1	3,492	10.0
Midwife	17,173	27.6	165	0.6	17,008	48.7
Other	107	0.2	29	0.1	78	0.2

There has been a steady increase in the proportion of live births occurring in hospitals in recent years (see Table C). The year 1951 was the first in which at least half of the deliveries took place in hospitals, and there has been much improvement since that time. The progress made by the nonwhites has been more rapid, but there is still a large difference between the percentages for the two racial groups.

Table C. Percentage of live births in hospitals or clinics, 1952-1956, by race

Year	Total	White	Nonwhite
1956	65.7	97.2	41.0
1955	62.4	96.0	36.4
1954	60.2	94.4	33.0
1953	58.2	93.3	30.5
1952	54.8	91.5	26.5

The proportions of live births attended by physicians during 1952-1956 are shown in Table D. The improvement shown in this table is less spectacular than that

in the preceding table, but is also steady. An important milestone for the nonwhites was reached in 1956 when, for the first time, more than half of the nonwhite mothers were delivered by physicians. Figures on deliveries in hospitals may soon be almost the same as those on deliveries by physicians, because in the 1950's there has been an uninterrupted decline in the percentage of births attended by physicians in homes.

Table D. Percentage of live births attended by physicians, 1952-1956, by race

Year	Total	White	Nonwhite
1956	72.2	99.3	51.0
1955	70.4	99.1	48.3
1954	70.4	98.9	47.7
1953	69.6	98.8	46.5
1952	67.7	98.7	43.9

Age of Mother

Table E gives the age distribution of mothers by race. Little variation in this distribution has occurred in recent years. Approximately half were 15-24 years of age and well over one-third were 25-34 years old. The period 25-34 years was the only one in which the percentage of white mothers was greater than the percentage of nonwhite.

Table E. Live births by age of mother and race, with per cent of total births

Age of Mother	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
10-14 years	338	0.5	31	0.1	307	0.9
15-19 years	61,550	99.0	27,186	99.7	34,364	98.5
15-24 years	30,976	49.8	13,545	49.6	17,431	49.9
25-34 years	23,503	37.8	10,925	40.0	12,578	36.0
35-44 years	7,071	11.4	2,716	10.0	4,355	12.5
45-54 years	151	0.2	37	0.1	114	0.3
Unknown	139	0.2	24	0.1	115	0.3

FETAL DEATHS

In 1952 the registration of all fetal deaths, regardless of length of gestation, was initiated in Mississippi. Prior to that time, only stillbirths of 20 or more weeks of gestation were reported. Registration of early fetal deaths is still far from complete, but there is no estimate of the degree of underregistration. There were 2,693 fetal death certificates filed during 1956 and 2,650 certificates for residents. The resident total was 50 more than the comparable figure for 1955, and the increase was observed in both races. For the State the fetal death ratio (fetal deaths per 1,000 live births) was 42.6. The ratio for whites was 37.0 with county ratios varying from zero in Franklin and Issaquena to 104.4 in Washington. The ratio for nonwhites was 47.0 with variations from zero in George County to 85.9 in Lawrence.

On 904 of the records the gestation period was stated to be less than 20 weeks, on 1,588 it was 20 weeks or more, and on 158 the length was not stated. In the period 1954-1956 there have been more fetal deaths after less than 20 weeks of gestation than

after 20 weeks or more reported among the whites, and the difference has grown larger each year. The reporting of late (20 or more weeks of gestation) fetal deaths for both racial groups has declined each year in this period.

Using the certificates with sex stated, a sex ratio of 1,419 males to 1,000 females was determined. This is the highest ratio calculated in the five years of all-inclusive fetal death registration and was a sharp jump from the 1,257 in 1955. As usual, the sex ratio for fetal deaths was much higher than that for live births, but was not so reliable a figure because the sex was not stated on 686 or 26 per cent of the fetal death certificates.

The distribution by race and attendant is given in Table F. As in 1954 and 1955, approximately 62 per cent of the total were nonwhite. Slightly over two-thirds of the reported fetal deaths were attended by physicians in hospitals and only one-fifth were attended by midwives. Although the proportion of white live births in hospitals was larger than the percentage of white fetal deaths in hospitals, the reverse was true for the nonwhites. This unusual situation in the nonwhite group may be accounted for by the underreporting of fetal deaths that occur at home and the tendency to transfer difficult and premature deliveries to a hospital.

Table F. Fetal deaths by attendant and race

<u>Attendant</u>	<u>Total</u>		<u>White</u>		<u>Nonwhite</u>	
	<u>No.</u>	<u>%</u>	<u>No.</u>	<u>%</u>	<u>No.</u>	<u>%</u>
<u>Total</u>	<u>2,650</u>	<u>100.0</u>	<u>1,008</u>	<u>100.0</u>	<u>1,642</u>	<u>100.0</u>
Physician in hospital	1,782	67.2	917	91.0	865	52.7
Physician in home	306	11.5	83	8.2	223	13.6
Midwife	535	20.2	3	0.3	532	32.4
Other	27	1.0	5	0.5	22	1.3

Table G shows the percentage of fetal deaths delivered in hospitals in the past five years. There has been very little change in the figures for the whites during this time, but the nonwhites have shown a steady improvement from 40.3 per cent in 1952 to 52.7 per cent in 1956, the first year in which more than half have occurred in hospitals. The gap between the two race groups is a large one, but not so large as that for live births.

Table G. Percentage of fetal deaths delivered in hospitals, 1952-1956, by race

<u>Year</u>	<u>Total</u>	<u>White</u>	<u>Nonwhite</u>
1956	67.2	91.0	52.7
1955	63.5	88.1	48.5
1954	61.6	90.2	44.6
1953	56.4	89.6	42.6
1952	54.9	90.0	40.3

DEATHS

Resident Deaths

In 1956 the resident death total was 19,780, and, although this figure represented an increase of about 380 over the number for 1955, it was still one of the lowest ever recorded in the State. The crude death rate, based on the new population estimate, was 9.4 per 1,000 population in comparison with a rate of 8.9 in 1955 based on the 1950 census population.

A major change took place in 1956 in the racial distribution of deaths. For the first time, more white than nonwhite deaths were registered. It was to be expected that this would eventually happen since the nonwhite death rate was approaching that for the whites and there continued to be a small majority of whites in the population, but this change was not expected as early as 1956. The total of 9,969 deaths of white persons was the largest recorded since 1937 and the sixth largest reported since 1913, the first year of death registration in Mississippi. According to preliminary figures from other states and the National Office of Vital Statistics, there was a general rise in the number of deaths in the South in 1956, but no information is available at present to indicate whether the increase occurred among whites only.

County rates for the white population ranged from 4.2 in Issaquena to 15.1 in Kemper; the state rate for whites was 8.7. The range for the nonwhites was from 2.2 in Itawamba County to 16.5 in Warren County. Although the number of deaths of nonwhites in 1956 was the lowest on record in Mississippi, the rate showed an increase from 10.0 in 1955 to 10.2 in 1956 because of the change in population base. Use of new population estimates appears to have affected the county rates for whites more than those for nonwhites. In 1956 there were 26 counties with a rate of 10 or more for the whites in comparison with only 7 such counties in 1955. The number of nonwhite rates in this category was nearly the same in both years. About one-third or 6,704 of the persons who died in 1956 were urban residents, but 47 per cent of the deaths occurred in urban areas.

The resident birth rate exceeded the death rate by 20.1 per 1,000 population. For whites the excess was 15.1 and for nonwhites it was 26.0. These figures were almost exactly the same as in 1955.

Age at Death

Table H shows the distribution of deaths by age and race. Despite the large increment over 1955 in the number of deaths of white persons, declines occurred in the 1-4, 5-9, 10-14, 35-44, and 45-54 year periods of this group. There were decreases in nearly all of the nonwhite age groups, but an important increase took place in the number of deaths among persons 65 years of age and over. Almost 12 per cent of all deaths were among persons under one year of age, but only a little over 13 per cent occurred in the groups 1-44 years old, that is, the next 44 years of life.

Comparison of the age distribution of the two races reveals significant differences. The proportion of deaths of nonwhites which occurred among children under 10 years of age was more than double that for whites, and in the groups from 10 to 55 years relatively more deaths of nonwhites than of whites were registered. "Retirement" age or 65 had been reached by 60 per cent of the whites who died in 1956 and only 44 per cent of the nonwhites. A new tabulation was prepared in 1956 and Table H has been expanded to show the number of deaths at 65-74, 75-84, and 85 years of age and over. This more detailed age breakdown made possible the calculation of the median age at death. For whites this figure was 69.1 years and for nonwhites, 60.8 years.

Table H. Age-race distribution of deaths

Age Group	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
All Ages	19,780	100.0	9,969	100.0	9,811	100.0
Under 1 year	2,353	11.9	734	7.4	1,619	16.5
1-4 years	353	1.8	95	1.0	258	2.6
5-9 years	151	0.8	51	0.5	100	1.0
10-14 years	136	0.7	55	0.6	81	0.8
15-24 years	466	2.4	212	2.1	254	2.6
25-34 years	543	2.7	226	2.3	317	3.2
35-44 years	958	4.8	424	4.2	534	5.4
45-54 years	1,763	8.9	820	8.2	943	9.6
55-64 years	2,776	14.0	1,406	14.1	1,370	14.0
65-74 years	4,358	22.0	2,353	23.6	2,005	20.4
75-84 years	4,106	20.8	2,595	26.0	1,511	15.4
85 years and over	1,784	9.0	993	10.0	791	8.1
Unknown	33	0.2	5	0.0	28	0.3

Leading Causes of Death

In Table I are listed the ten leading causes of death in the State in 1956. The first seven of these causes are the same as in 1955. Congenital malformations and general arteriosclerosis which ranked eighth and ninth respectively in 1955 tied for eighth place in 1956, and diabetes mellitus replaced homicide in tenth position. Provisional figures for the United States show nine of the causes listed for Mississippi, but cirrhosis of liver is included instead of chronic and unspecified nephritis and other renal sclerosis.

The leading cause, heart diseases, was responsible for nearly 31 per cent of the total number. Vascular lesions affecting the central nervous system, in second place, and malignant neoplasms which ranked third accounted for 12.9 and 12.3 per cent. Altogether the ten leading causes were responsible for 77 per cent of the deaths in 1956. For the whites the same ten causes were listed as for the total population. On the list for nonwhites, however, general arteriosclerosis, congenital malformations, and diabetes mellitus were replaced by homicide, gastritis including associated conditions, and tuberculosis, respectively.

Table I. Ten leading causes of death by race, with per cent of total deaths

Cause with International List Number	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
1. Heart diseases (410-443)	6,104	30.8	3,703	37.1	2,401	24.5
2. Vascular lesions affecting central nervous system (330-334)	2,550	12.9	1,229	12.3	1,321	13.5
3. Malignant neoplasms (140-205)	2,432	12.3	1,439	14.4	993	10.1
4. Accidents (800-962)	1,302	6.6	700	7.0	602	6.1
5. Certain diseases of early infancy (760-776)	1,191	6.0	472	4.7	719	7.3
6. Pneumonia and influenza (480-493)	678	3.4	250	2.5	428	4.4
7. Chronic & unspec. nephritis & other renal sclerosis (592-594)	329	1.7	130	1.3	199	2.0
8. General arteriosclerosis (450)	247	1.2	161	1.6	86	0.9
8. Congenital malformations (750-759)	247	1.2	142	1.4	105	1.1
10. Diabetes mellitus (260)	225	1.1	116	1.2	109	1.1

Table J. Five leading causes of death by age group

<u>Age and Cause of Death</u>	<u>No. of Deaths</u>	<u>% of Total</u>
1-4 years		
All Causes	<u>353</u>	<u>100.0</u>
1. Accidents	106	30.0
2. Pneumonia and influenza	66	18.7
3. Congenital malformations	30	8.5
4. Malignant neoplasms, including leukemia	19	5.4
5. Gastritis, duodenitis, enteritis, and colitis	15	4.2
5-14 years		
All Causes	<u>287</u>	<u>100.0</u>
1. Accidents	137	47.7
2. Malignant neoplasms, including leukemia	28	9.8
3. Pneumonia and influenza	11	3.8
4. Rheumatic fever	6	2.1
5. Tuberculosis	5	1.7
5. Nonmeningococcal meningitis	5	1.7
5. Heart diseases	5	1.7
5. Gastritis, duodenitis, enteritis, and colitis	5	1.7
5. Congenital malformations	5	1.7
15-24 years		
All Causes	<u>466</u>	<u>100.0</u>
1. Accidents	229	49.1
2. Homicide	44	9.4
3. Malignant neoplasms, including leukemia	26	5.6
4. Heart diseases	24	5.2
5. Maternal deaths	18	3.9
25-44 years		
All Causes	<u>1,501</u>	<u>100.0</u>
1. Accidents	309	20.6
2. Heart diseases	244	16.2
3. Malignant neoplasms, including leukemia	209	13.9
4. Vascular lesions affecting central nervous system	111	7.4
5. Homicide	110	7.3
45-64 years		
All Causes	<u>4,539</u>	<u>100.0</u>
1. Heart diseases	1,660	36.6
2. Malignant neoplasms, including leukemia	869	19.1
3. Vascular lesions affecting central nervous system	585	12.9
4. Accidents	205	4.5
5. Chronic & unspecified nephritis & other renal sclerosis	71	1.6
65 years and over		
All Causes	<u>10,248</u>	<u>100.0</u>
1. Heart diseases	4,154	40.5
2. Vascular lesions affecting central nervous system	1,828	17.8
3. Malignant neoplasms, including leukemia	1,272	12.4
4. Pneumonia and influenza	295	2.9
5. Accidents	244	2.4

Leading causes of death for each age group except infants are given in Table J. None of these lists was exactly like any other, but there were some causes which appeared repeatedly. Accidents not only was listed in every group but also was the chief cause in all groups under 45 years of age. Malignant neoplasms also appeared on every list and heart diseases on all except 1-4 years. Vascular lesions affecting the central nervous system was listed in the three groups 25 and over. Also appearing in three groups (1-4, 5-14, and 65 and over) was pneumonia and influenza. In both lists for children under 15, congenital malformations and gastritis including allied conditions were included, and homicide was one of the leading causes in the 15-24 and 25-44 year groups. Those causes which were on one list only were rheumatic fever (5-14), tuberculosis (5-14), nonmeningococcal meningitis (5-14), maternal deaths (15-24), and chronic and unspecified nephritis and other renal sclerosis (45-64). Actually nine causes were shown for the age group 5-14 because five tied for fifth place. Special tabulations on the age group 65 years and over show that for the period 65-80 years the five leading causes were the same as those for the group as a whole. However, in the periods 80-84 and 85 years and over, general arteriosclerosis replaced accidents.

Ill-Defined Causes

Certificates with no satisfactory statement of cause numbered 2,300 or 11.6 per cent of the total in 1956. Senility was the only cause given on 91, ill-defined diseases of early infancy were on 595, and other ill-defined and unknown causes on 1,614. For white persons the percentage of certificates in this category was 5.6 and for nonwhites, 17.8. Over 20 per cent of the certificates filed in Wilkinson, Hancock, and Issaquena Counties had no definite cause, and in Humphreys, Lawrence, Amite, DeSoto, Yazoo, Noxubee, Benton, Kemper, and Carroll Counties 15 to 20 per cent had ill-defined and unknown causes.

Deaths thus classified were so numerous that if placed on the list of ten leading causes they would rank fourth and would appear as one of the five chief causes in all age groups except 15-24 and 25-44 years. If the true causes of death for these persons could be determined and properly assigned, the order of the principal causes might not be the same and even the causes might be different.

Deaths by Cause

For the second consecutive year tuberculosis was not one of the ten leading causes of death in Mississippi. The number of deaths from this cause continued to decline and totaled only 199 in 1956. Both this total and the rate of 9.4 per 100,000 population were less than half those reported four years ago in 1952.

In Table K is given the age, race, sex distribution of tuberculosis deaths. Although there was continued improvement in tuberculosis mortality, an increase from 43 in 1955 to 62 in 1956 occurred in the number of deaths of white males, and the number doubled in the age group 65 years and over. Especially large decreases were noted among nonwhite females 35-64 years of age. At all ages under 55 years there were more deaths of nonwhites than of whites. Deaths among whites were concentrated in the older groups with 72 per cent occurring at 55 or later. Among nonwhites 70 per cent occurred before the age of 55. Males outnumbered females in each period except 15-24 and 25-34 years of age.

Table K. Tuberculosis deaths by age, race, and sex

Age Group	Total	White		Nonwhite	
		Male	Female	Male	Female
All Ages	199	62	18	76	43
Under 1 year	3	0	0	2	1
1-14 years	12	2	0	5	5
15-24 years	12	0	0	5	7
25-34 years	26	2	3	10	11
35-44 years	29	6	1	12	10
45-54 years	23	8	0	14	1
55-64 years	34	16	2	14	2
65 years and over	60	28	12	14	6

The number of syphilis deaths increased from 38 in 1955 to 44 in 1956 and both races were affected. As in 1955, no deaths from congenital syphilis were reported. Other types in 1956 were cardiovascular with 35 deaths, central nervous system with 8, and one unqualified.

No deaths from typhoid fever have been registered since 1953, and in 1956 there were none from paratyphoid fever.

Dysentery caused 54 deaths in 1956 as compared with 48 in 1955. Ten of these were classified as bacillary, 28 as amebic, and 16 were unspecified.

Only one death resulted from streptococcal sore throat in 1956, and 1952 was the last year in which there was a death from scarlet fever.

There were 5 deaths from diphtheria in 1956, as many as were reported in 1954 and 1955 combined. All 5 of these fatalities occurred among children 1-14 years of age.

Each year since 1953 the number of deaths due to whooping cough has risen. There were 14 in 1953 and 41 in 1956. All but one of the 1956 victims were under 5 years of age.

In 1956 meningococcal meningitis caused 14 deaths. Six of these were infants, but the rest were scattered throughout the other age groups.

Comparative figures on deaths from acute poliomyelitis in 1954, 1955, and 1956 are 16, 6, and 12 respectively. An increase over 1955 was also shown in the number of cases reported. One of those who died in 1956 was a colored female 56 years old, one was an 11 year old child, and the other 10 were less than 10 years of age. In addition, a child 5 years of age died of late (one year or more after onset) effects of the disease.

Smallpox has not appeared as a cause of death in Mississippi since 1945.

Another disease which caused nearly as many deaths in 1956 as in the two previous years was measles. Ten deaths from this cause were reported, and 2 of them were adults 35-54 years of age.

A nonwhite female 82 years of age died of rabies in 1956. This was the first death from rabies since 1953.

In the period 1953-1956 and also in 1951 no deaths from malaria were registered.

Thirteen deaths resulted from infectious hepatitis in 1956, 11 from tetanus,

8 from acute infectious encephalitis, 2 from mumps, and one each from brucellosis (undulant fever), rubella (German measles), and typhus.

In 1956, the cancer death totals for both racial groups were the highest on record in this State, and for the first time the rate for nonwhites exceeded 100 per 100,000 population. There were 2,432 deaths from this cause or 115 more than in 1955. The distribution of these deaths by age, sex, and race is given in Table L. Small declines from the numbers reported in 1955 were observed for males of both races, and the figures for females showed increases. There was an especially large increment in the number of deaths of nonwhite females 55 years of age and over.

More deaths occurred among whites than among nonwhites in all age groups except those under 15 years, and the excess was largest in the group 75-84 years. In most age groups, there were approximately the same number of males and females, but there were more than twice as many females in the 35-44 year period and females were in the majority in the 45-54 year period. The only groups in which males outnumbered the opposite sex were those 65 years and over.

Table L. Cancer deaths by age, sex, and race

Age Group	Total	Male		Female	
		White	Nonwhite	White	Nonwhite
All Ages	2,432	755	434	684	559
Under 10 years	39	12	9	8	10
10-14 years	15	4	3	3	5
15-24 years	26	9	4	8	5
25-34 years	55	17	11	17	10
35-44 years	154	32	15	53	54
45-54 years	347	82	52	103	110
55-64 years	522	169	91	140	122
65-74 years	665	226	132	165	142
75-84 years	480	169	94	147	70
85 years and over	127	35	23	39	30
Unknown	2	0	0	1	1

Table M lists cancer deaths by primary site in descending order according to the number in the total population. Deaths assigned to the leading site, digestive organs and peritoneum, exceeded those of the breast and genito-urinary system by 14 in 1956. These two sites tied for first place in 1955 and both showed an increase in 1956, but the increase in deaths from cancer of the breast and genito-urinary system occurred only among nonwhite females. The two leading sites accounted for 81 per cent of the cancer deaths among nonwhite females (51 per cent were assigned to breast and genito-urinary system alone), 72 per cent among nonwhite males, 70 per cent among white females, and only 49 per cent among white males. Cancer of the lymphatic and hematopoietic system also showed a large increment and moved up to third rank, but there was a decline in the number of deaths from cancer of the respiratory system.

The number of death certificates which showed diabetes mellitus as the underlying cause in 1956 was 225, or almost the same as in 1955. There was a small decrease for the whites and a slight increase for the nonwhites.

Pellagra, once a great killer in Mississippi, now appears to have been brought under control. In 1956, for the first time, no death from this cause was registered.

Table M. Cancer deaths by primary site, sex, and race, with percentage distribution

Primary Site and International List Number	Male				Female			
	White		Nonwhite		White		Nonwhite	
	No.	%	No.	%	No.	%	No.	%
Total	755	100.0	434	100.0	684	100.0	559	100.0
1. Digestive organs & peritoneum (150-159)	244	32.3	188	43.3	216	31.6	166	29.7
2. Breast & genito-urinary system (170-181)	126	16.7	124	28.6	263	38.4	287	51.3
3. Lymphatic & hematopoietic tissues (200-205)	105	13.9	37	8.5	79	11.5	27	4.8
4. Respiratory system (160-165)	148	19.6	38	8.8	24	3.5	14	2.5
5. Brain & nervous system (192, 193)	40	5.3	6	1.4	17	2.5	8	1.4
6. Skin (190, 191)	26	3.4	3	0.7	24	3.5	3	0.5
7. Buccal cavity & pharynx (140-148)	20	2.6	8	1.8	6	0.9	6	1.1
8. Bone & connective tissue (196, 197)	6	0.8	7	1.6	15	2.2	8	1.4
9. Other & unspec. sites (194, 195, 198, 199)	40	5.3	23	5.3	40	5.8	40	7.2

An increase of 103 was observed in deaths due to vascular lesions affecting the central nervous system, which brought the total for 1956 up to 2,550. Both races were affected by this change.

Only 13 certificates showed rheumatic fever as the cause of death in 1956. This was only about half as many as were reported in 1955.

The death toll from heart diseases reached a new high of 6,104 in 1956. However, there was no increase in the number of nonwhites who died from this cause. Arteriosclerotic and degenerative heart conditions were responsible for 4,097 deaths, hypertension with heart disease for 1,440, chronic rheumatic heart disease for 145, and other types for 422.

Hypertension without mention of heart involvement has caused approximately the same number of deaths in each of the past 3 years. The total for 1956 was 198.

In 1956 general arteriosclerosis caused 247 deaths and was again one of the ten leading causes. The total and the racial distribution for 1956 were almost identical with those for 1955.

Reported deaths from influenza also numbered about the same as in 1955. Of a total of 103, 78 were complicated by pneumonia. Approximately half of the deaths from this cause occurred during the first 3 months of the year.

The number of deaths with pneumonia as the primary cause rose from 530 in 1955 to 575 in 1956. However, small reductions were shown for the lobar and primary atypical types. Pneumonia of the newborn (infants under 28 days of age) caused 94 deaths in addition to the 575 already mentioned and this figure also was higher than in 1955.

There were 194 deaths from gastritis and associated conditions (duodenitis, enteritis, and colitis) in 1956. The majority of the deaths occurred among nonwhites, and this group of diseases ranked ninth among the leading causes of death for that racial group. For both whites and nonwhites, however, the totals were lower than in 1955.

Chronic nephritis was responsible for 329 deaths in 1956 or only 5 less than in 1955. Acute nephritis deaths among white persons, however, increased from 17 to 31 and brought the total number up to 81.

The number of deaths in 1956 which resulted from accidents was 1,302. Although there was a decline of 47 from 1955 in the total number of accidental deaths, those caused by motor vehicle accidents showed a small increase, from 541 to 558. Water transport, poisoning by solid and liquid substances, machinery, and obstruction or suffocation due to inhalation or ingestion of food and other objects were other types of accidents for which the number of deaths went up in 1956. There were fewer deaths from accidental fires among nonwhites and from accidental falls among whites. The largest number of accidental deaths occurred in December which was also the month when motor vehicle fatalities were highest. Table N shows the five leading causes of accidental death in 1956 by race. On the list for 1955 the same causes appeared in the same order.

Table N. Five leading causes of accidental deaths among residents, by race

Type of Accident	Total	White	Nonwhite
1. Motor vehicle	558	347	211
2. Fire	193	54	139
3. Fall	101	70	31
4. Drowning	87	42	45
5. Firearm	63	28	35

In 1956 there were 661 fatalities from transport accidents of all types - aircraft, watercraft, railway, motor and other road vehicle - which occurred in Mississippi. This is the largest total reported for this classification of accidents in the six years for which comparable figures are available. The site of 90 per cent of these accidents was a rural area. Counties in which as many as 15 persons were fatally injured in transport accidents were Hinds and Lauderdale with 32 each, Harrison with 29, Washington with 28, Jackson and Jones with 21 each, Grenada and Madison with 17 each, Coahoma with 16, and DeSoto with 15. Harrison, Hinds, and Lauderdale have been on this list in each of the six years, 1951-1956, this study has been carried out, and Coahoma, Jackson, Jones and Washington Counties have appeared on it several times.

Suicides numbered 113 and homicides 214 in 1956. There was a decline from 1955 of 12 (entirely among whites) in the number of suicides and a decrease of 13 (entirely among nonwhites) in the homicide total. The majority of suicides were committed by white persons and most of the homicide victims were nonwhites. The most common method of carrying out both of these types of violence was the use of firearms.

Infant Mortality

An infant mortality rate of 37.8 per 1,000 live births was recorded in 1956. This was slightly higher than the rate for 1955 and the rate for white infants was

also higher. The mortality rate for nonwhite infants remained the same although there was a small reduction in the actual number of deaths. Distribution of infant deaths by age and race is shown in Table 0. Usually the nonwhite rates for all age groups are larger than the corresponding rates for whites, but in 1956 there was an increase in the rate for white infants under 1 day old and also a decline for nonwhites of this age. The result was that in this one group the rate for whites exceeded that for nonwhites. The greatest difference between the two race groups occurred in the period 28 days to one year. Comparison of the figures in Table 0 with provisional data for the United States shows that Mississippi has not made as much progress in the field of infant mortality as the rest of the country. For the United States, the infant mortality rate in 1956 was 26.0 and the rate for infants 28 days to one year old was 7.2, only half that recorded in this State.

Table 0. Infant deaths by age and race, with rates

Age at Death	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
<u>Total under 1 year</u>	<u>2,353</u>	<u>37.8</u>	<u>734</u>	<u>26.9</u>	<u>1,619</u>	<u>46.4</u>
Under 1 day	667	10.7	301	11.0	366	10.5
One to 28 days	784	12.6	280	10.3	504	14.4
28 days to 1 year	902	14.5	153	5.6	749	21.5

* Deaths per 1,000 live births.

Table P gives the five leading causes of infant deaths. There was no change from 1955 in the causes listed or their rank, although there were fewer deaths from birth injuries and from gastritis and associated conditions. Immaturity, which is no longer considered a separate classification in ranking causes of infant deaths, was mentioned either in combination with another cause or as the only cause on 46 per cent of the certificates for white infants and 31 per cent of those for nonwhites.

Table P. Leading causes of infant deaths, by race

Cause of Death	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
<u>All Causes</u>	<u>2,353</u>	<u>100.0</u>	<u>734</u>	<u>100.0</u>	<u>1,619</u>	<u>100.0</u>
1. Postnatal asphyxia & atelectasis	246	10.4	145	19.8	101	6.2
2. Influenza & pneumonia, except pneumonia of newborn	206	8.8	29	4.0	177	10.9
3. Congenital malformations	192	8.2	115	15.7	77	4.8
4. Birth injuries	141	6.0	73	9.9	68	4.2
5. Gastritis, duodenitis, enteritis & colitis, except diarrhea of newborn	137	5.8	14	1.9	123	7.6

Infant death certificates which showed ill-defined causes or no cause at all numbered 920 or 39 per cent of the total in 1956. For 31 per cent of the white and 43 per cent of the nonwhite infants no definite cause of death was given, and the percentage for the whites was higher than in 1955. Determination of the true cause of death for this large group of infants might change the leading causes of death which were listed above.

Infant mortality rates by race for 1920-1956 are shown in Graph 1. In general the trend has been downward for both races, but in recent years there has been some upward swing in the curve for the nonwhites.

Graph 1. Infant Mortality
1920-1956



Perinatal Mortality

The recently coined term "perinatal mortality" includes fetal deaths of 20 or more weeks of gestation and infant deaths that occur during the first week of life. These two groups have been combined for study because many of the same factors are associated with the heavy loss of life during these periods. The formula for calculation of the perinatal mortality rate is given below:

$$\frac{\text{Fetal deaths of 20 or more weeks} + \text{postnatal deaths under 1 week}}{\text{Fetal deaths of 20 or more weeks} + \text{live births}} \times 1,000$$

Table Q shows the figures used for the calculations and the rates for 1956. The total rate of 43.8 was slightly lower than that for 1955. For nonwhites the rate also declined, but that for the whites increased. The nonwhite rate was much higher than the one for whites and the difference was probably due to the relatively large number of fetal deaths among nonwhites.

Table Q. Perinatal mortality rates, by race

	Total	White	Nonwhite
Fetal deaths, 20 weeks & over	1,588	419	1,169
Infant deaths, under 1 week	1,204	525	679
Live births	62,178	27,278	34,900
Rate*	43.8	34.1	51.2

* See formula above.

Maternal Mortality

Although the maternal mortality rates of 1.2 per 1,000 live births in the total population, 0.4 for white mothers, and 1.8 for nonwhites were the same in 1955 and 1956, there was a reduction of 2 in the number of deaths of nonwhite women. The total figure of 74 and that of 62 for nonwhites are the lowest on record; the lowest year for whites was 1954. There is still a large discrepancy between the rates by race, but the gap has narrowed considerably in recent years.

Toxemia was the leading cause of maternal deaths as shown in Table R, there was an increase over 1955 in the number caused by hemorrhage which ranked second, and these two conditions caused exactly the same number of deaths among white women. Numerical differences between the whites and nonwhites were greatest for toxemia and hemorrhage.

Table R. Maternal deaths by cause and race, with rates

Cause of Death	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
<u>All Causes</u>	<u>74</u>	<u>1.2</u>	<u>12</u>	<u>0.4</u>	<u>62</u>	<u>1.8</u>
Toxemia	25	0.4	3	0.1	22	0.6
Hemorrhage	19	0.3	3	0.1	16	0.4
Sepsis	8	0.1	2	0.1	6	0.2
Ectopic pregnancy	6	0.1	1	0.04	5	0.1
Abortion without sepsis or toxemia	1	0.02	0	-	1	0.03
Abortion with sepsis	2	0.03	0	-	2	0.06
<u>Other complications</u>	<u>13</u>	<u>0.2</u>	<u>3</u>	<u>0.1</u>	<u>10</u>	<u>0.3</u>

* Deaths per 1,000 live births.


Table S gives the age and race-specific maternal death rates. There was a large increment over 1955 in the rate for white women 35-44 years of age and small increases were noted for whites 25-34 and nonwhites 15-24 years old. Only one death occurred in the 10-14 year group and there were none in the 45-54 year group. At all ages the rates for nonwhite mothers were higher than those for white mothers.

Table S. Maternal deaths by age group and race, with rates

Age Group	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
All Ages	74	1.2	12	0.4	62	1.8
10-14 years	1	3.0	0	-	1	3.2
15-24 years	18	0.6	0	-	18	1.0
25-34 years	30	1.3	6	0.5	24	1.9
35-44 years	25	3.5	6	2.2	19	4.4
45-54 years	0	-	0	-	0	-

* Deaths per 1,000 live births to mothers in specific age-race group.

Maternal death rates by race for 1913-1956 are shown in Graph 2. A downward trend began in the 1930's and has been very rapid since 1940.



MARRIAGES

In 1956 the number of marriages occurring in Mississippi was 66,223, only 200 less than in the peak year of 1955, but over 2,000 more than the next largest number recorded in a single year. Among the whites there was an increase of over 500 which made 1956 the peak year for that race. The decrease among the nonwhites amounted to more than 700. The average daily number of marriages was 181, but in December, the month with the largest total, the average was 237.

Table T shows the age distribution of brides and grooms. The brides were relatively young; 30,461 or 46 per cent were teen-agers and 69 per cent were under 25 years of age. Comparison of the ages of resident and nonresident brides showed that the proportions of teen-agers were approximately equal for both groups and that there were only minor differences at other ages. Slightly more than 8 per cent of the grooms were teen-agers, but 46.4 per cent were 20-24 years of age.

Table T. Age distribution of brides and grooms

Age	Bride		Groom	
	No.	%	No.	%
<u>Total</u>	<u>66,223</u>	<u>100.0</u>	<u>66,223</u>	<u>100.0</u>
Under 15 years	352	0.5	1	0.0
15 years	1,065	1.6	11	0.0
16 years	2,270	3.4	121	0.2
17 years	2,634	4.0	694	1.0
18 years	16,979	25.6	2,151	3.2
19 years	7,161	10.8	2,564	3.9
20-24 years	15,070	22.8	30,699	46.4
25-34 years	10,234	15.4	15,655	23.6
35-44 years	5,783	8.7	6,754	10.2
45-54 years	2,976	4.5	3,995	6.0
55 years & over	1,640	2.5	3,525	5.3
Unknown	59	0.1	53	0.1

Although the total number of marriages in Mississippi in 1956 showed a decline, the number of out-of-state brides again increased and there were more than in any previous year. Since 1947 there has been a increment in the number of nonresident brides each year and in the eleven years, 1946-1956, over 345,000 have come to this State to be married. However, as the nonresident total has gone up, the resident total has declined. In 1956 there were 21,154 more out-of-state and 19,135 fewer Mississippi brides than in 1946.

In Table U are shown data on the residence status of both brides and grooms. Nonresident brides numbered 43,212 or 65.2 per cent of the total. By race, the proportions were 73.7 per cent for whites and 42.8 for nonwhites. In only 31 per cent of the marriages were both parties residents of this State. Large numbers of people come to Mississippi to be married because no waiting period or premarital examination is required, and, as a result, the State had the second highest marriage rate in the nation, according to the most recently published figures. Nevada was the only state with a higher rate in 1956 and the rate for Mississippi was more than triple that for the country as a whole.

Table U. Marriages in Mississippi by residence status of bride and groom

Groom	Total	Bride	
		Resident	Nonresident
<u>Total</u>	<u>66,223</u>	<u>23,011</u>	<u>43,212</u>
Resident	21,997	20,660	1,337
Nonresident	44,226	2,351	41,875

Brides came to Mississippi from every state in the union, the District of Columbia, and Alaska, and there were 43 from foreign countries. A little more than a third of all the women married in Mississippi in 1956 were from either Alabama (home of 12,005) or Tennessee (home of 10,847). Six other states furnished over 1,000 brides each: Arkansas - 4,590, Louisiana - 4,274, Illinois - 2,662, Missouri - 2,646, Florida - 2,034, and Kentucky - 1,337. There were also 474 from Texas, 386 from Indiana, and 378 from Michigan. The home states of the nonresident grooms were tabulated for the first time in 1956. Their distribution was similar to that for the nonresident brides, except that fewer came from states bordering on Mississippi and more came from the states farther away.

Each county in the State issued marriage licenses to at least a few out-of-state brides in 1956, but the counties adjacent to Alabama, Tennessee, and Arkansas issued the largest numbers. As a result, there were many differences, some of them quite large, between the occurrence and residence figures by county. Table V shows the counties to which the greatest numbers of out-of-state brides came. DeSoto and Alcorn were the two leading ones, and the brides who came from other states comprised 96 per cent of the total married there. In fact, over one-fourth of the brides married in Mississippi in 1956 were married in these two counties.

Table V. Number of out-of-state brides married in certain counties

DeSoto	8,831	Jackson	2,216
Alcorn	7,568	George	1,756
Tishomingo	3,684	Lauderdale	1,506
Lowndes	3,369	Itawamba	1,119
Washington	2,695	Harrison	1,075

On July 1, 1956, a new marriage record was put into use in Mississippi and more detailed information will be available for the year 1957. Preliminary analyses indicate that some remarriages appear to be taking place when participants are still legally married to other parties and also that Mississippi marriage licenses are being used in other states. Records filed during the last six months of 1956 showed that 81 Mississippi licenses were used for marriage ceremonies performed in other states. Most of these marriages were performed in Louisiana (31) and Tennessee (22) but they also occurred in nine other states, some as distant as Michigan and New Jersey. These 81 records have been omitted from the 1956 tabulations because of the questionable legality of the marriages.

DIVORCES

There were 5,027 divorces granted in Mississippi during 1956, 182 more than were registered in 1955 and 26 more than in 1954. However, during the nine-year period 1947-1955 there was a steady decline in the number of divorces in this State. Both racial groups shared in the current increase, the nonwhites to a greater degree. Cruel and inhuman treatment and desertion continued to rank as the two leading causes and together accounted for 85 per cent of the total granted. Of the divorces among whites, cruel and inhuman treatment was given as the main cause for 55 per cent; among nonwhites 57 per cent were granted on grounds of desertion. The wife was listed as the complainant on 62 per cent of the records. Only 221 suits or 4 per cent were contested.

No minor children were involved in 53 per cent of the divorces, and 21 per cent of the couples had only one minor child. Five of the families had 10 or more minor children. The total number of minor children affected was 4,646.

Forty-five per cent of the divorces in 1956 were granted to persons married less than 5 years, 8 per cent had been married less than one year, but 12 per cent had been married 20 years or longer. More divorces were granted to couples in their second year of marriage than in any other year. The distribution of divorces according to duration of marriage was very different for the two races. Of the white couples, 52 per cent were granted after less than 5 years of marriage and only 8 per cent after 20 years or longer; for the nonwhites, corresponding percentages were 29 and 21 respectively. The increment over 1955 in the number of divorces was proportionately greater among persons with three or more children and those married ten years or longer. There was also a large increase in the number who had been married only one year.

TABLE 1
ESTIMATED POPULATION*, MISSISSIPPI
AS OF JULY 1, 1956

County	Total	White	Nonwhite
<u>TOTAL</u>	<u>2,109,800</u>	<u>1,147,200</u>	<u>962,600</u>
Adams	36,300	19,300	17,000
Alcorn	25,100	21,500	3,600
Amite	16,900	7,200	9,700
Attala	23,200	12,300	10,900
Benton	8,500	4,300	4,200
Bolivar	59,100	17,600	41,500
Calhoun	15,100	10,500	4,600
Carroll	12,900	4,900	8,000
Chickasaw	17,200	9,800	7,400
Choctaw	9,400	6,200	3,200
Claiborne	11,200	3,100	8,100
Clarke	17,300	10,100	7,200
Clay	15,100	7,600	7,500
Coahoma	52,600	14,400	38,200
Copiah	27,400	12,400	15,000
Covington	15,300	9,500	5,800
DeSoto	24,100	8,400	15,700
Forrest	44,500	32,300	12,200
Franklin	8,800	5,300	3,500
George	10,500	9,200	1,300
Greene	8,000	6,400	1,600
Grenada	18,700	8,700	10,000
Hancock	13,100	11,800	1,300
Harrison	104,700	87,600	17,100
Hinds	170,700	98,400	72,300
Holmes	28,000	6,800	21,200
Humphreys	21,300	6,000	15,300
Issaquena	4,800	1,900	2,900
Itawamba	13,600	12,700	900
Jackson	42,500	34,000	8,500
Jasper	16,900	8,000	8,900
Jefferson	10,100	2,500	7,600
Jeff Davis	16,200	6,400	9,800
Jones	51,500	37,700	13,800
Kemper	11,900	3,700	8,200
Lafayette	16,200	8,100	8,100

Table 1 - Page 2
 Estimated Population*, Mississippi, as of July 1, 1956

County	Total	White	Nonwhite
Lamar	12,300	10,400	1,900
Lauderdale	62,100	39,400	22,700
Lawrence	11,600	6,900	4,700
Leake	15,500	8,200	7,300
Lee	37,400	26,000	11,400
Leflore	46,000	15,800	30,200
Lincoln	26,200	16,700	9,500
Lowndes	37,400	20,800	16,600
Madison	32,300	8,100	24,200
Marion	22,700	13,300	9,400
Marshall	22,900	6,500	16,400
Monroe	30,400	19,000	11,400
Montgomery	13,800	7,400	6,400
Neshoba	18,200	12,700	5,500
Newton	19,000	11,200	7,800
Noxubee	17,200	4,800	12,400
Oktibbeha	22,000	11,200	10,800
Panola	32,100	12,900	19,200
Pearl River	21,600	16,500	5,100
Perry	7,200	5,200	2,000
Pike	32,200	17,300	14,900
Pontotoc	18,200	14,700	3,500
Prentiss	18,100	15,900	2,200
Quitman	25,500	9,400	16,100
Rankin	27,400	15,300	12,100
Scott	19,900	10,900	9,000
Sharkey	12,500	3,400	9,100
Simpson	21,100	13,000	8,100
Smith	13,700	11,100	2,600
Stone	6,400	4,900	1,500
Sunflower	54,100	16,300	37,800
Tallahatchie	28,400	10,000	18,400
Tate	17,200	6,800	10,400
Tippah	16,600	12,900	3,700
Tishomingo	13,000	12,300	700
Tunica	22,700	4,400	18,300
Union	18,800	14,800	4,000
Walthall	14,300	7,900	6,400
Warren	39,500	22,800	16,700
Washington	76,500	31,100	45,400
Wayne	15,500	9,700	5,800
Webster	10,500	8,000	2,500
Wilkinson	13,200	3,400	9,800
Winston	19,500	11,100	8,400
Yalobusha	13,000	6,800	6,200
Yazoo	36,000	13,800	22,200

* Prepared by Dr. Harald A. Pedersen, Sociology Department,
 Mississippi State College, with adjustments by Statistical
 Service, Mississippi State Board of Health.

TABLE 2
LIVE BIRTHS
MISSISSIPPI
1956

By Place of Occurrence, Reallocated to Mother's Usual Residence,
by Race, with Rates

County	Place of Occurrence			Place of Residence			Rate*		
	Number		Non-white	Number		Non-white	Rate*		Non-white
	Total	White		Total	White		Total	White	
<u>STATE</u>	<u>62,806</u>	<u>27,785</u>	<u>35,021</u>	<u>62,178</u>	<u>27,278</u>	<u>34,900</u>	<u>29.5</u>	<u>23.8</u>	<u>36.2</u>
Adams	1,710	789	921	1,252	604	648	34.5	31.3	38.1
Alcorn	659	546	113	565	454	111	22.5	21.1	30.8
Amite	279	20	259	421	115	306	24.9	16.0	31.5
Attala	618	269	349	607	260	347	26.2	21.1	31.8
Benton	178	18	160	251	76	175	29.5	17.7	41.7
Bolivar	2,204	435	1,769	2,089	463	1,626	35.3	26.3	39.2
Calhoun	433	234	199	400	223	177	26.5	21.2	38.5
Carroll	279	36	243	347	84	263	26.9	17.1	32.9
Chickasaw	447	218	229	453	223	230	26.3	22.8	31.1
Choctaw	210	104	106	213	117	96	22.6	18.9	30.0
Claiborne	290	41	249	329	61	268	29.4	19.7	33.1
Clarke	290	142	148	442	198	244	25.5	19.6	33.9
Clay	548	180	368	561	197	364	37.2	25.9	48.5
Coahoma	1,630	377	1,253	1,664	342	1,322	31.6	23.8	34.6
Copiah	954	279	675	775	222	553	28.3	17.9	36.9
Covington	249	150	99	397	205	192	25.9	21.6	33.1
DeSoto	657	97	560	720	177	543	29.9	21.1	34.6
Forrest	1,577	1,149	428	1,329	860	469	29.9	26.6	38.4
Franklin	157	72	85	249	124	125	28.3	23.4	35.7
George	309	256	53	324	270	54	30.8	29.3	41.5
Greene	144	77	67	249	174	75	31.1	27.2	46.9
Grenada	583	244	339	532	194	338	28.4	22.3	33.8
Hancock	175	145	30	269	233	36	20.5	19.7	27.7
Harrison	3,645	2,988	657	3,436	2,803	633	32.8	32.0	37.0
Hinds	5,197	2,677	2,520	5,059	2,515	2,544	29.6	25.6	35.2
Holmes	958	158	800	957	153	804	34.2	22.5	37.9
Humphreys	726	114	612	749	141	608	35.2	23.5	39.7
Issaquena	45	1	44	105	23	82	21.9	12.1	28.3
Itawamba	107	89	18	282	257	25	20.7	20.2	27.8
Jackson	1,056	714	342	1,281	916	365	30.1	26.9	42.9
Jasper	193	67	126	440	140	300	26.0	17.5	33.7
Jefferson	152	0	152	344	44	300	34.0	17.6	39.5
Jeff Davis	412	156	256	382	115	267	23.6	18.0	27.2
Jones	2,133	1,123	1,010	1,452	885	567	28.2	23.5	41.1
Kemper	273	42	231	344	75	269	28.9	20.3	32.8
Lafayette	505	230	275	525	244	281	32.4	30.1	34.7

Table 2 - Page 2
Live Births - Mississippi - 1956

County	Place of Occurrence			Place of Residence					
	Number		Non-white	Number		Non-white	Rate*		Non-white
	Total	White		Total	White		Total	White	
Lamar	109	62	47	344	274	70	28.0	26.3	36.8
Lauderdale	2,065	1,086	979	1,613	858	755	26.0	21.8	33.2
Lawrence	188	30	158	279	116	163	24.0	16.8	34.7
Leake	441	170	271	467	170	297	30.1	20.7	40.7
Lee	1,274	889	385	1,059	682	377	28.3	26.2	33.1
Leflore	1,820	509	1,311	1,678	386	1,292	36.5	24.4	42.8
Lincoln	691	407	284	676	368	308	25.8	22.0	32.4
Lowndes	1,069	504	565	1,068	501	567	28.6	24.1	34.2
Madison	1,185	194	991	1,146	178	968	35.5	22.0	40.0
Marion	562	323	239	633	369	264	27.9	27.7	28.1
Marshall	770	90	680	817	144	673	35.7	22.2	41.0
Monroe	764	399	365	765	383	382	25.2	20.2	33.5
Montgomery	297	90	207	359	138	221	26.0	18.6	34.5
Neshoba	559	283	276	590	339	251	32.4	26.7	45.6
Newton	487	299	188	476	250	226	25.0	22.3	29.0
Noxubee	623	97	526	619	103	516	36.0	21.4	41.6
Oktibbeha	837	370	467	754	292	462	34.3	26.1	42.8
Panola	851	232	619	900	250	650	28.0	19.4	33.8
Pearl River	692	511	181	639	452	187	29.6	27.4	36.7
Perry	104	77	27	222	147	75	30.8	28.3	37.5
Pike	1,001	479	522	867	356	511	26.9	20.6	34.3
Pontotoc	319	217	102	375	266	109	20.6	18.1	31.1
Prentiss	487	411	76	410	343	67	22.6	21.6	30.4
Quitman	922	251	671	850	225	625	33.3	23.9	38.8
Rankin	251	24	227	638	295	343	23.3	19.3	28.3
Scott	596	299	297	585	257	328	29.4	23.6	36.4
Sharkey	274	53	221	422	91	331	33.8	26.8	36.4
Simpson	555	244	311	513	250	263	24.3	19.2	32.5
Smith	207	141	66	390	250	140	28.5	22.5	53.8
Stone	188	133	55	171	117	54	26.7	23.9	36.0
Sunflower	1,538	316	1,222	1,614	356	1,258	29.8	21.8	33.3
Tallahatchie	838	139	699	920	207	713	32.4	20.7	38.8
Tate	462	22	440	582	114	468	33.8	16.8	45.0
Tippah	336	218	118	331	215	116	19.9	16.7	31.4
Tishomingo	156	143	13	280	262	18	21.5	21.3	25.7
Tunica	654	82	572	674	89	585	29.7	20.2	32.0
Union	538	410	128	434	314	120	23.1	21.2	30.0
Walthall	644	355	289	383	153	230	26.8	19.4	35.9
Warren	1,476	685	791	1,172	538	634	29.7	23.6	38.0
Washington	3,032	1,229	1,803	2,586	881	1,705	33.8	28.3	37.6
Wayne	470	285	185	470	254	216	30.3	26.2	37.2
Webster	185	93	92	226	125	101	21.5	15.6	40.4

Table 2 - Page 3
Live Births - Mississippi - 1956

County	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Wilkinson	493	159	334	436	74	362	33.0	21.8	36.9
Winston	456	178	278	533	227	306	27.3	20.4	36.4
Yalobusha	309	95	214	327	119	208	25.2	17.5	33.5
Yazoo	1,049	265	784	1,091	313	778	30.3	22.7	35.0
Out-of-State	964	863	101	1,592	1,370	222	-	-	-

Cities over 10,000 Population**	Place of Occurrence			Place of Residence		
	Number			Number		
	Total	White	Non-white	Total	White	Non-white
Biloxi	2,287	2,053	234	1,631	1,433	198
Clarksdale	874	373	501	630	219	411
Columbus	794	473	321	532	259	273
Greenville	1,283	578	705	1,224	502	722
Greenwood	904	495	409	592	256	336
Gulfport	1,215	917	298	852	598	254
Hattiesburg	1,533	1,148	385	865	531	334
Jackson	4,509	2,665	1,844	3,211	1,545	1,666
Laurel	2,066	1,094	972	721	363	358
McComb	611	358	253	248	149	99
Meridian	1,952	1,081	871	1,110	641	469
Natchez	853	686	167	829	418	411
Pascagoula	102	5	97	437	310	127
Tupelo	894	705	189	338	220	118
Vicksburg	1,405	684	721	795	385	410
Total Urban#	32,914	20,237	12,677	19,753	10,735	9,018
Total Rural	29,892	7,548	22,344	42,425	16,543	25,882

* Rate per 1,000 population.

** Cities having over 10,000 population according to 1950 census.

Includes all towns of 2,500 population or more according to 1950 census.

TABLE 3
LIVE BIRTHS
MISSISSIPPI
1956

By County and Cities over 10,000, by Race, Attendant, and Sex
on Basis of Usual Residence of Mother

County	STATE	WHITE			NONWHITE			SEX		
		Total	Phys. in Hosp.	Other	Total	Phys. in Hosp.	Other	Male	Female	
TOTAL*	62,178	27,278	26,510	574	194	34,900	14,322	3,492	17,086	31,685
Adams	1,252	604	604	0	0	648	609	5	34	666
Alcorn	565	454	411	37	6	111	56	14	41	285
Amitte	421	115	108	7	0	306	70	46	190	212
Attala	607	260	245	13	2	347	128	43	176	207
Benton	251	76	69	6	1	175	19	22	134	129
Bolivar	2,089	463	432	18	13	1,626	645	122	859	1,085
Calhoun	400	223	212	9	2	177	115	11	51	204
Carroll	347	84	82	0	2	263	72	16	175	185
Chickasaw	453	223	205	18	0	230	83	43	104	162
Choctaw	213	117	107	8	2	96	33	0	63	237
Claiborne	329	61	61	0	0	268	40	5	223	118
Clarke	442	198	194	0	4	244	176	8	60	180
Clay	561	197	196	1	0	364	171	28	165	149
Coahoma	1,664	342	337	5	0	1,322	375	221	726	149
Copiah	775	222	219	2	1	553	352	24	177	357
Covington	397	205	202	2	1	192	141	9	42	357
DeSoto	720	177	171	3	3	543	199	31	313	357
Forrest	1,329	860	860	0	0	469	312	4	153	715
Franklin	249	124	123	1	0	125	69	0	56	121
George	324	270	260	9	1	54	16	6	32	163
Greene	249	174	156	14	4	75	22	9	44	128
Grenada	532	194	193	1	0	338	33	58	247	298
Hancock	269	233	231	2	0	36	6	5	25	135
Harrison	3,436	2,803	2,764	7	32	633	312	16	305	1,798
										1,638

Table 3 - Page 2
Live Births - Mississippi - 1956

County	TOTAL,*	WHITE			NONWHITE			SEX			
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female
Hinds	5,059	2,515	2,505	10	0	2,544	1,412	228	904	2,574	2,485
Holmes	957	153	144	8	1	804	103	90	611	478	479
Humphreys	749	141	136	3	2	608	97	15	496	387	362
Issaquena	105	23	22	0	1	82	40	2	40	57	48
Itawamba	282	257	228	28	1	25	17	2	6	159	123
Jackson	1,281	916	910	3	3	365	155	70	140	612	669
Jasper	440	140	139	1	0	300	200	16	84	219	221
Jefferson	344	44	44	0	0	300	148	4	148	188	156
Jeff Davis	382	115	105	10	0	267	102	72	93	188	194
Jones	1,452	885	881	1	3	567	504	3	60	726	726
Kosciusko	344	75	72	1	2	269	82	8	179	161	183
Lafayette	525	24	235	8	1	281	33	23	225	269	256
Lamar	344	274	263	11	0	70	43	20	7	185	159
Lauderdale	1,613	858	855	3	0	755	538	58	159	805	808
Lawrence	279	116	102	6	1	163	65	12	86	142	137
Leake	467	170	167	2	1	297	154	3	140	243	224
Lee	1,059	682	666	14	2	377	159	153	65	555	504
Leflore	1,678	386	375	9	2	1,292	161	122	1,009	830	848
Lincoln	676	368	365	0	3	308	130	83	95	347	329
Lowndes	1,068	501	463	37	1	567	174	135	258	538	530
Madison	1,146	178	173	4	1	968	165	68	735	599	547
Marion	633	369	355	8	6	264	107	3	154	318	315
Marshall	817	144	134	7	3	673	96	117	460	409	408
Monroe	765	383	372	8	3	382	142	48	192	405	360
Montgomery	359	138	136	1	1	221	48	5	168	184	175
Neshoba	590	239	233	3	3	251	132	10	109	304	286
Newton	476	250	246	3	1	226	137	30	59	245	231
Noxubee	619	103	100	2	1	516	90	39	387	317	302
Oktibbeha	754	292	290	1	1	462	278	10	174	389	365
Panola	900	250	241	6	3	650	276	105	269	484	416

Table 3 - Page 3
Live Births - Mississippi - 1956

County	TOTAL*	WHITE			NONWHITE			SEX		
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male
Pearl River	639	452	443	6	3	187	162	6	19	316
Perry	222	147	143	4	0	75	61	5	9	116
Pike	867	356	352	0	4	511	321	41	149	451
Pontotoc	375	266	254	9	3	109	36	14	59	191
Prentiss	410	343	338	4	1	67	35	9	23	201
Quitman	850	225	217	8	0	625	367	74	184	423
Rankin	638	295	291	2	2	343	171	24	148	322
Scott	585	257	253	3	1	328	164	7	157	297
Sharkey	422	91	88	1	2	331	168	4	159	205
Simpson	513	250	248	2	0	263	194	6	63	258
Smith	390	250	240	9	1	140	87	35	18	190
Stone	171	117	115	2	0	54	36	0	18	91
Sunflower	1,614	356	341	10	5	1,258	408	224	626	856
Tallahatchie	920	207	190	12	5	713	215	69	429	488
Tate	582	114	92	21	1	468	29	45	394	432
Tippah	331	215	190	23	2	116	38	42	36	290
Tishomingo	280	262	228	33	1	18	6	12	0	182
Tunica	674	89	76	11	2	585	162	37	386	149
Union	434	314	306	4	4	120	32	14	74	319
Walton	383	153	149	0	4	230	180	1	49	222
Warren	1,172	538	535	0	3	634	342	2	290	177
Washington	2,586	881	870	5	6	1,705	765	343	597	580
Wayne	470	254	249	0	5	216	146	4	66	592
Webster	226	125	121	4	0	101	35	15	51	1,264
Wilkinson	436	74	74	0	0	362	91	18	253	1,322
Winston	533	227	221	3	3	306	85	10	211	232
Yalobusha	327	119	110	8	1	208	38	112	58	238
Yazoo	1,091	313	277	7	7	778	106	19	653	111

Table 3 - Page 4
Live Births - Mississippi - 1956

Cities over 10,000 Population	TOTAL*	WHITE			NONWHITE			SEX			
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female
Biloxi	1,631	1,433	1,418	2	13	198	136	0	62	854	777
Clarksdale	630	219	218	1	0	411	218	92	101	326	304
Columbus	532	259	246	12	1	273	85	77	111	272	260
Greenville	1,224	502	499	1	2	722	250	196	276	610	614
Greenwood	592	256	255	0	1	336	74	3	259	286	306
Gulfport	852	598	591	3	4	254	121	9	124	427	425
Hattiesburg	865	531	531	0	0	334	225	3	106	465	400
Jackson	3,211	1,545	1,543	2	0	1,666	1,074	169	423	1,646	1,565
Laurel	721	363	363	0	0	358	328	2	28	351	370
McComb	248	149	149	0	0	99	77	10	12	132	116
Meridian	1,110	641	640	1	0	469	360	52	57	551	559
Natchez	829	418	418	0	0	411	389	2	20	462	367
Pascagoula	437	310	308	2	0	127	31	25	71	218	219
Tupelo	338	220	220	0	0	118	50	43	25	183	155
Vicksburg	795	385	383	0	2	410	216	2	192	396	399

* Includes transcripts furnished by the National Office of Vital Statistics of births to residents of Mississippi occurring out-of-state.

TABLE 4
FETAL DEATHS
MISSISSIPPI
1956

By Place of Occurrence, Reallocated to Mother's Usual Residence,
by Sex, Race and Attendant

County	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total	Phys. in Hosp.	Other	Total	Phys. in Hosp.	Other	Male	Female	Unde- termined
STATE	2,693	2,650	1,008	217	83	8	1,642	865	223	554	1,152
Adams	65	46	15	14	1	0	31	26	3	2	22
Alcorn	14	14	10	6	3	1	4	1	2	8	6
Amite	13	29	4	4	0	0	25	16	3	1	5
Attala	26	28	13	13	0	0	15	10	3	2	18
Benton	5	11	4	4	0	0	7	2	2	12	11
Bolivar	93	91	18	18	0	0	73	25	2	3	5
Calhoun	19	15	5	3	2	0	10	8	0	38	42
Carroll	12	16	3	0	0	0	13	5	2	42	34
Chickasaw	15	16	6	6	0	0	10	2	6	5	15
Choctaw	4	5	4	4	1	0	1	1	0	0	1
Claiborne	13	17	1	1	0	1	16	9	0	7	1
Clarke	10	17	3	2	1	0	14	10	3	1	1
Clay	18	20	5	5	0	0	11	11	0	18	21
Coahoma	83	93	23	22	6	0	70	31	6	3	42
Copiah	18	17	6	6	0	0	11	5	1	2	7
Covington	8	15	10	10	0	0	15	4	1	5	8
DeSoto	13	18	3	3	1	0	32	21	6	11	7
Forrest	72	68	36	35	0	1	0	0	1	10	21
Franklin	3	7	0	0	0	0	7	6	1	0	0
George	4	3	3	3	0	0	0	0	0	1	3
Greene	2	5	3	3	0	0	2	1	0	1	2
Grenada	35	34	10	10	9	1	0	0	1	10	6
Hancock	5	10	7	7	0	0	3	3	0	0	0
Harrison	87	77	56	53	3	0	21	13	0	8	40

Table 4 - Page 2
Fetal Deaths - Mississippi - 1956

County	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total	Phys. in Hosp.	Other	Total	Phys. in Hosp.	Other	Male	Female	Unde- termined
Hinds	344	304	160	116	43	1	144	84	21	39	100
Holmes	86	82	13	11	2	0	69	22	17	30	33
Humphreys	32	33	1	0	0	0	32	11	2	19	21
Issaquena	0	2	0	0	0	0	2	2	0	0	0
Itawamba	2	8	7	7	0	2	1	0	1	4	4
Jackson	47	56	43	39	0	2	13	6	1	31	9
Jasper	7	21	5	5	0	0	16	12	4	10	7
Jefferson	4	10	4	4	0	1	6	2	2	5	2
Jeff Davis	24	24	6	5	1	0	18	6	4	6	6
Jones	103	70	35	34	1	0	35	28	1	6	26
Kemper	5	11	1	1	6	0	10	8	1	5	4
Lafayette	15	14	6	6	0	0	8	1	5	7	2
Lamar	3	7	4	4	0	1	3	1	1	3	2
Lauderdale	102	81	30	29	5	0	51	37	1	44	20
Lawrence	13	20	6	5	5	0	14	5	1	12	6
Leake	12	19	5	5	1	0	14	10	1	9	10
Lee	27	23	14	14	1	0	9	5	3	14	6
Leflore	75	73	11	11	0	0	62	18	12	32	31
Lincoln	26	25	17	17	9	1	0	8	2	18	7
Lowndes	29	26	10	10	1	0	0	7	6	12	2
Madison	37	34	1	1	0	0	0	10	1	17	1
Marion	24	29	14	14	5	0	15	7	1	14	6
Marshall	50	39	18	18	9	0	10	6	0	13	3
Monroe	17	13	9	9	3	0	0	6	3	8	6
Montgomery	7	9	4	4	0	0	0	5	1	4	1
Neshoba	20	18	7	5	0	1	11	9	1	8	4
Newton	37	33	16	14	2	0	0	2	1	10	8
Noxubee	42	42	20	10	8	0	40	16	6	12	5
Oktibbeha	20	20	10	10	8	0	0	10	8	9	4
Panola	41	42	8	8	0	0	34	17	6	19	4

Table 4 - Page 3
Fetal Deaths - Mississippi - 1956

County	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total	Phys. in Hosp.	Phys. in Home	Total	Phys. in Hosp.	Phys. in Home	Male	Female	Undetermined
Pearl River	19	13	19	8	8	0	0	11	0	8	5
Perry	8	11	13	8	0	0	0	5	1	3	5
Pike	31	24	11	11	6	0	0	13	12	15	2
Pontotoc	19	13	11	6	8	0	0	7	4	1	7
Prentiss	12	11	8	0	0	0	0	3	2	1	0
Quitman	29	23	7	7	0	0	0	16	15	9	4
Rankin	12	30	17	14	3	0	0	13	8	5	16
Scott	25	30	12	12	1	0	0	18	7	13	9
Sharkey	10	16	8	7	1	0	0	15	8	6	9
Simpson	10	16	9	3	3	0	0	18	8	0	4
Smith	3	3	3	2	2	0	0	6	5	0	0
Stone	3	3	3	2	2	0	0	1	1	1	1
Sunflower	53	54	14	14	0	0	0	40	21	6	22
Tallahatchie	30	35	7	6	1	1	0	28	8	5	9
Tate	21	29	2	1	1	0	0	27	7	19	4
Tippah	30	23	15	15	2	0	0	8	5	0	0
Tishomingo	1	3	2	2	4	0	0	1	1	0	1
Tunica	27	30	4	4	4	0	0	26	14	10	14
Union	37	34	29	29	0	0	0	5	4	0	1
Walthall	25	15	5	5	0	0	0	10	9	1	7
Warren	79	44	13	13	1	0	0	31	19	0	3
Washington	203	193	92	88	4	1	0	101	70	12	11
Wayne	24	24	7	1	1	0	0	11	4	6	5
Webster	7	7	1	1	1	0	0	6	3	2	1
Wilkinson	27	19	4	3	2	0	0	15	7	0	8
Winston	14	17	1	1	1	0	0	15	6	2	1
Yalobusha	5	44	11	11	1	0	0	6	3	0	0
Yazoo	41	44	11	11	1	0	0	33	25	25	15

Table 4 - Page 4
Fetal Deaths - Mississippi - 1956

Cities over 10,000 Population	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male
Biloxi	58	46	35	32	3	0	11	10	0	1	23
Clarksdale	53	43	20	19	7	0	23	15	5	3	17
Columbus	23	16	8	7	1	0	8	2	4	2	8
Greenville	156	132	78	75	3	0	54	37	6	11	30
Greenwood	40	22	8	8	0	0	14	4	1	9	8
Gulfport	27	23	15	15	0	0	8	2	0	6	13
Hattiesburg	69	50	24	23	1	0	26	18	1	7	15
Jackson	302	210	115	87	27	1	95	56	19	20	66
Laurel	97	43	18	17	1	0	25	22	1	2	15
McComb	26	10	6	6	0	0	4	4	0	0	7
Meridian	96	56	23	22	1	0	33	25	4	4	30
Natchez	47	33	9	8	1	0	24	22	2	0	16
Pascagoula	11	20	16	15	1	0	4	3	1	0	11
Tupelo	22	7	6	6	0	0	1	0	1	0	4
Vicksburg	75	32	10	10	0	0	22	15	0	0	7

TABLE 5
DEATHS
MISSISSIPPI
1956

By Cause by Race on Basis of Place of Residence

International List Number	Cause of Death*	Total	White	Non-white
<u>001-999</u>	<u>TOTAL ALL CAUSES</u>	<u>19,780</u>	<u>9,969</u>	<u>9,811</u>
<u>001-019</u>	<u>Tuberculosis, all forms</u>	<u>199</u>	<u>80</u>	<u>119</u>
001	Respiratory, with mention of occupational disease of lung	1	0	1
002	Pulmonary	171	73	98
003	Pleural	2	1	1
004	Primary complex with symptoms	1	0	1
008	Unspecified site	1	1	0
010	Meninges and central nervous system	7	1	6
011	Intestines, peritoneum, and mesenteric glands	3	0	3
012	Bones and joints, active or unspecified	4	1	3
016	Genito-urinary system	1	1	0
017	Adrenal glands	1	1	0
018	Other organs	1	0	1
019	Disseminated	6	1	5
<u>020-029</u>	<u>Syphilis and its sequelae</u>	<u>44</u>	<u>13</u>	<u>31</u>
022	Aneurysm of aorta	18	10	8
023	Other cardiovascular	17	0	17
024	Tabes dorsalis	1	1	0
025	General paralysis of insane	3	2	1
026	Other central nervous system	4	0	4
029	Unqualified	1	0	1
042	Other Salmonella infections	3	2	1
044	Brucellosis (undulant fever)	1	1	0
<u>045-048</u>	<u>Dysentery, all forms</u>	<u>54</u>	<u>16</u>	<u>38</u>
045	Bacillary	10	4	6
046	Amebiasis	28	11	17
048	Unspecified forms	16	1	15
049	Food poisoning	6	0	6
<u>050,051</u>	<u>Scarlet fever and streptococcal sore throat</u>	<u>1</u>	<u>1</u>	<u>0</u>
051	Streptococcal sore throat	1	1	0
053	Septicemia and pyemia	20	6	14
055	Diphtheria	5	1	4
056	Whooping cough	41	2	39
057	Meningococcal infections	14	5	9
061	Tetanus	11	1	10
064	Other bacterial diseases	2	0	2
080	Acute poliomyelitis	12	4	8
081	Late effects of acute poliomyelitis	1	0	1
082	Acute infectious encephalitis	8	3	5
085	Measles	10	1	9
086	Rubella (German measles)	1	0	1
087	Chickenpox	3	2	1
089	Mumps	2	2	0
092	Infectious hepatitis	13	8	5
094	Rabies	1	0	1

Table 5 - Page 2
Deaths - Mississippi - 1956

35

International List Number	Cause of Death*	Total	White	Non-white
096	Other diseases attributable to viruses	5	3	2
<u>100-108</u>	<u>Typhus and other rickettsial diseases</u>	<u>1</u>	<u>1</u>	<u>0</u>
104	Tick-borne typhus	1	1	0
130	Infestation with worms of other, mixed, or unspecified type	3	0	3
134	Other fungus infections	9	6	3
138	Other infective and parasitic diseases	3	1	2
<u>140-205</u>	<u>Malignant neoplasm</u>	<u>2,432</u>	<u>1,439</u>	<u>993</u>
<u>140-148</u>	<u>Buccal cavity and pharynx</u>	<u>40</u>	<u>26</u>	<u>14</u>
140	Lip	4	4	0
141	Tongue	5	3	2
142	Salivary gland	3	2	1
143	Floor of mouth	4	2	2
144	Other parts of mouth and mouth unspecified	8	4	4
145	Oral mesopharynx	1	0	1
146	Nasopharynx	1	1	0
147	Hypopharynx	1	1	0
148	Pharynx, unspecified	13	9	4
<u>150-159</u>	<u>Digestive organs and peritoneum</u>	<u>814</u>	<u>460</u>	<u>354</u>
150	Esophagus	35	25	10
151	Stomach	265	105	160
152	Small intestine, including duodenum	10	7	3
153	Large intestine except rectum	207	136	71
154	Rectum	55	29	26
155	Biliary passages and liver (stated to be primary site)	29	24	5
156	Liver (secondary and unspecified)	72	42	30
157	Pancreas	126	83	43
158	Peritoneum	8	5	3
159	Unspecified digestive organs	7	4	3
<u>160-165</u>	<u>Respiratory system</u>	<u>224</u>	<u>172</u>	<u>52</u>
160	Nose, nasal cavities, middle ear, and accessory sinuses	8	5	3
161	Larynx	13	9	4
162	Trachea, and bronchus and lung specified as primary	85	67	18
163	Lung and bronchus, unspecified as to whether primary or secondary	115	89	26
164	Mediastinum	2	2	0
165	Thoracic organs (secondary)	1	0	1
<u>170-181</u>	<u>Breast and genito-urinary organs</u>	<u>800</u>	<u>389</u>	<u>411</u>
170	Breast	194	106	88
171	Cervix uteri	157	64	93
172	Corpus uteri	5	4	1
174	Uterus, unspecified	100	32	68
175	Ovary, Fallopian tube, and broad ligament	51	31	20
176	Other and unspecified female genital organs	6	4	2
177	Frostate	164	70	94
178	Testis	8	8	0
179	Other and unspecified male genital organs	5	2	3
180	Kidney	39	21	18
181	Bladder and other urinary organs	71	47	24

Table 5 - Page 3
Deaths - Mississippi - 1956

International List Number	Cause of Death*	Total	White	Non-white
<u>190-199</u>	<u>Other and unspecified sites</u>	<u>306</u>	<u>208</u>	<u>98</u>
190	Melanoma of skin	23	21	2
191	Skin, other	33	29	4
192	Eye	4	1	3
193	Brain and other parts of nervous system	67	56	11
194	Thyroid gland	13	9	4
195	Other endocrine glands	5	4	1
196	Bone including jaw bone	25	15	10
197	Connective tissue	11	6	5
198	Lymph nodes, secondary and unspecified	3	1	2
199	Other and unspecified sites	122	66	56
<u>200-205</u>	<u>Lymphatic and hematopoietic tissues</u>	<u>248</u>	<u>184</u>	<u>64</u>
200	Lymphosarcoma and reticulosarcoma	47	38	9
201	Hodgkin's disease	21	16	5
202	Other forms of lymphoma (reticulososis)	14	10	4
203	Multiple myeloma (plasmacytoma)	30	20	10
204	Leukemia and aleukemia	136	100	36
<u>210-229</u>	<u>Benign neoplasm</u>	<u>23</u>	<u>8</u>	<u>15</u>
210	Buccal cavity and pharynx	1	0	1
211	Other parts of digestive system	1	1	0
213	Breast	1	0	1
214	Uterine fibromyoma	8	1	7
216	Ovary	3	2	1
223	Brain and other parts of nervous system	8	4	4
228	Hemangioma and lymphangioma	1	0	1
<u>230-239</u>	<u>Neoplasm of unspecified nature</u>	<u>20</u>	<u>12</u>	<u>8</u>
230	Digestive organs	2	1	1
232	Breast	1	0	1
237	Brain and other parts of nervous system	12	9	3
238	Skin and musculoskeletal system	1	0	1
239	Other and unspecified organs	4	2	2
241	Asthma	49	30	19
243	Urticaria	1	0	1
245	Other allergic disorders	1	0	1
250	Simple goiter	3	1	2
252	Thyrotoxicosis with or without goiter	7	0	7
260	Diabetes mellitus	225	116	109
272	Diseases of pituitary gland	4	3	1
273	Diseases of thymus gland	1	1	0
274	Diseases of adrenal glands	6	5	1
277	Polyglandular dysfunction and other diseases of endocrine glands	2	1	1
280	Beriberi	1	0	1
285	Osteomalacia	1	1	0
286	Other avitaminoses and nutritional deficiency states	9	2	7
289	Other metabolic diseases	2	1	1
<u>290-293</u>	<u>Anemias</u>	<u>36</u>	<u>13</u>	<u>23</u>
290	Pernicious and other hyperchromic	5	4	1
291	Iron deficiency (hypochromic)	2	0	2
292	Other specified type	15	2	13
293	Unspecified type	14	7	7

Table 5 - Page 4
Deaths - Mississippi - 1956

37

International List Number	Cause of Death*	Total	White	Non-white
294	Polycythemia	6	6	0
296	Purpura and other hemorrhagic conditions	4	2	2
297	Agranulocytosis	1	1	0
298	Diseases of spleen	3	3	0
299	Other diseases of blood and blood-forming organs	2	1	1
300	Schizophrenic disorders (dementia praecox)	2	0	2
301	Manic-depressive reaction	1	1	0
304	Senile psychosis	5	2	3
307	Alcoholic psychosis	1	1	0
309	Other and unspecified psychoses	1	1	0
322	Alcoholism	15	8	7
323	Other drug addiction	1	1	0
<u>330-334</u>	<u>Vascular lesions affecting central nervous system</u>	<u>2,550</u>	<u>1,229</u>	<u>1,321</u>
330	Subarachnoid hemorrhage	56	29	27
331	Cerebral hemorrhage	1,639	721	918
332	Cerebral embolism and thrombosis	451	286	165
334	Other and ill-defined	404	193	211
340	Meningitis, except meningococcal and tuberculous	45	18	27
342	Intracranial and intraspinal abscess	7	3	4
343	Encephalitis, myelitis, and encephalomyelitis (except acute infectious)	8	2	6
344	Late effects of intracranial abscess or pyogenic infection	4	3	1
345	Multiple sclerosis	10	6	4
350	Paralysis agitans	18	14	4
351	Cerebral spastic infantile paralysis	6	2	4
352	Other cerebral paralysis	13	3	10
353	Epilepsy	32	12	20
355	Other diseases of brain	7	3	4
356	Motor neurone disease and muscular atrophy	7	6	1
357	Other diseases of spinal cord	4	4	0
364	Polyneuritis and polyradiculitis	3	2	1
366	Other and unspecified forms of neuralgia and neuritis	2	1	1
368	Other diseases of peripheral nerves except autonomic	1	1	0
391	Otitis media without mention of mastoiditis	2	1	1
<u>400-402</u>	<u>Rheumatic fever</u>	<u>13</u>	<u>6</u>	<u>7</u>
400	Without mention of heart involvement	1	0	1
401	With heart involvement	10	5	5
402	Chorea	2	1	1
<u>410-413</u>	<u>Diseases of heart</u>	<u>6,104</u>	<u>3,703</u>	<u>2,401</u>
410	Diseases of mitral valve	74	32	42
411	Diseases of aortic valve specified as rheumatic	10	8	2
414	Other endocarditis specified as rheumatic	4	3	1
415	Other myocarditis specified as rheumatic	5	0	5
416	Other heart disease specified as rheumatic	52	32	20
420	Arteriosclerotic heart disease including coronary disease	3,414	2,458	956
421	Chronic endocarditis not specified as rheumatic	119	47	72

Table 5 - Page 5
Deaths - Mississippi - 1956

38

International List Number	Cause of Death*	Total	White	Non-white
422	Other myocardial degeneration	564	281	283
430	Acute and subacute endocarditis	7	3	4
431	Acute myocarditis not specified as rheumatic	7	5	2
433	Functional disease of heart	45	29	16
434	Other and unspecified diseases of heart	363	188	175
441	Essential malignant hypertension with heart disease	14	4	10
442	Hypertensive heart disease with arteriolar nephrosclerosis	351	148	203
443	Other and unspecified hypertensive heart disease	1,075	465	610
<u>444-447</u>	<u>Hypertension without mention of heart</u>	<u>198</u>	<u>90</u>	<u>108</u>
444	Essential benign	37	14	23
445	Essential malignant	18	4	14
446	With arteriolar nephrosclerosis	119	58	61
447	Other hypertensive disease	24	14	10
450	General arteriosclerosis	247	161	86
<u>451-468</u>	<u>Other diseases of circulatory system</u>	<u>135</u>	<u>83</u>	<u>52</u>
451	Aortic aneurysm specified as nonsyphilitic, and dissecting aneurysm	49	38	11
452	Other aneurysm, except of heart and aorta	2	1	1
453	Peripheral vascular disease	2	1	1
454	Arterial embolism and thrombosis	5	4	1
455	Gangrene of unspecified cause	6	1	5
456	Other diseases of arteries	10	1	9
460	Varicose veins of lower extremities	2	0	2
462	Varicose veins of other specified sites	6	5	1
463	Phlebitis and thrombophlebitis of lower extremities	6	3	3
464	Phlebitis and thrombophlebitis of other sites	3	3	0
465	Pulmonary embolism and infarction	30	17	13
466	Other venous embolism and thrombosis	6	4	2
467	Other diseases of circulatory system	7	5	2
468	Certain diseases of lymph nodes and lymph channels	1	0	1
470	Acute nasopharyngitis (common cold)	3	0	3
471	Acute sinusitis	1	1	0
473	Acute tonsillitis	5	1	4
474	Acute laryngitis and tracheitis	2	0	2
475	Acute upper respiratory infection of multiple or unspecified sites	9	5	4
<u>480-483</u>	<u>Influenza</u>	<u>103</u>	<u>26</u>	<u>77</u>
480	With pneumonia	78	16	62
481	With other respiratory manifestations, and influenza unqualified	23	9	14
482	With digestive manifestations, but without respiratory symptoms	1	0	1
483	With nervous manifestations, but without digestive or respiratory symptoms	1	1	0
<u>490-493</u>	<u>Pneumonia</u>	<u>575</u>	<u>224</u>	<u>351</u>
490	Lobar	149	60	89

Table 5 - Page 6
Deaths - Mississippi - 1956

International List Number	Cause of Death*	Total	White	Non-white
491	Broncho	248	91	157
492	Primary atypical	41	17	24
493	Other and unspecified	137	56	81
<u>500-502</u>	<u>Bronchitis</u>	<u>29</u>	<u>15</u>	<u>14</u>
500	Acute	11	3	8
501	Unqualified	3	1	2
502	Chronic	15	11	4
510	Hypertrophy of tonsils and adenoids	5	1	4
513	Chronic sinusitis	2	2	0
517	Other diseases of upper respiratory tract	2	1	1
519	Pleurisy	3	0	3
520	Spontaneous pneumothorax	1	1	0
521	Abscess of lung	9	7	2
522	Pulmonary congestion and hypostasis	17	10	7
523	Pneumoconiosis due to silica and silicates (occupational)	1	0	1
524	Other specified pneumoconiosis and pulmonary fibrosis of occupational origin	1	1	0
525	Other chronic interstitial pneumonia	12	9	3
526	Bronchiectasis	25	22	3
527	Other diseases of lung and pleural cavity	56	44	12
532	Other inflammatory diseases of supporting structures of teeth	2	1	1
537	Diseases of salivary glands	1	1	0
539	Diseases of esophagus	2	1	1
<u>540, 541</u>	<u>Ulcer of stomach and duodenum</u>	<u>76</u>	<u>54</u>	<u>22</u>
540	Stomach	43	31	12
541	Duodenum	33	23	10
542	Gastrojejunal ulcer	1	1	0
543	Gastritis and duodenitis	4	3	1
545	Other diseases of stomach and duodenum	4	2	2
<u>550-553</u>	<u>Appendicitis</u>	<u>24</u>	<u>6</u>	<u>18</u>
550	Acute	21	5	16
551	Unqualified	3	1	2
<u>560, 561, 570</u>	<u>Hernia and intestinal obstruction</u>	<u>93</u>	<u>41</u>	<u>52</u>
560	Hernia of abdominal cavity without mention of obstruction	10	7	3
561	Hernia of abdominal cavity with obstruction	12	2	10
570	Intestinal obstruction, without mention of hernia	71	32	39
571	Gastro-enteritis and colitis, except ulcerative, age 4 weeks and over	178	30	148
572	Chronic enteritis and ulcerative colitis	12	7	5
573	Functional disorders of intestines	1	0	1
576	Peritonitis	7	3	4
577	Peritoneal adhesion	1	0	1
578	Other diseases of intestines and peritoneum	27	19	8
580	Acute and subacute yellow atrophy of liver	3	2	1
581	Cirrhosis of liver	95	68	27
583	Other diseases of liver	9	6	3
584	Cholelithiasis	31	25	6

Table 5 - Page 7
Deaths - Mississippi - 1956

40

International List Number	Cause of Death*	Total	White	Non-white
585	Cholecystitis without mention of calculi	18	14	4
586	Other diseases of gallbladder and biliary ducts	12	11	1
587	Diseases of pancreas	14	10	4
<u>590, 591</u>	<u>Acute nephritis and nephritis with edema</u>	<u>81</u>	<u>31</u>	<u>50</u>
590	Acute	53	23	30
591	With edema, including nephrosis	28	8	20
<u>529-594</u>	<u>Chronic and unspecified nephritis and other renal sclerosis</u>	<u>329</u>	<u>130</u>	<u>199</u>
592	Chronic nephritis	275	110	165
593	Nephritis not specified as acute or chronic	50	18	32
594	Other renal sclerosis	4	2	2
600	Infections of kidney	95	44	51
601	Hydronephrosis	2	1	1
602	Calculi of kidney and ureter	6	5	1
603	Other diseases of kidney and ureter	2	2	0
604	Calculi of other parts of urinary system	2	0	2
605	Cystitis	1	0	1
606	Other diseases of bladder	6	2	4
608	Stricture of urethra	6	0	6
609	Other diseases of urethra	5	0	5
610	Hyperplasia of prostate	77	31	46
611	Prostatitis	12	4	8
612	Other diseases of prostate	1	1	0
614	Orchitis and epididymitis	1	0	1
615	Redundant prepuce and phimosis	1	0	1
617	Other diseases of male genital organs	1	1	0
623	Chronic salpingitis and oophoritis	1	0	1
624	Salpingitis and oophoritis, unqualified	1	0	1
626	Diseases of parametrium and pelvic peritoneum (female)	4	0	4
630	Infective disease of uterus, vagina, and vulva	1	1	0
631	Uterovaginal prolapse	1	1	0
633	Other diseases of uterus	1	0	1
<u>640-689</u>	<u>Diseases of pregnancy, childbirth, and puerperium</u>	<u>74</u>	<u>12</u>	<u>62</u>
641	Other infections of genito-urinary tract during pregnancy	1	0	1
642	Toxemias of pregnancy	22	3	19
644	Other hemorrhage of pregnancy	1	0	1
645	Ectopic pregnancy	6	1	5
648	Other complications arising from pregnancy	3	0	3
650	Abortion without mention of sepsis or toxemia	1	0	1
651	Abortion with sepsis	2	0	2
660	Delivery without complication	4	0	4
670	Delivery complicated by placenta previa or antepartum hemorrhage	4	0	4
671	Delivery complicated by retained placenta	5	1	4
672	Delivery complicated by other postpartum hemorrhage	9	2	7
674	Delivery complicated by disproportion or mal-position of fetus	3	2	1
677	Delivery with other trauma	1	0	1

Table 5 - Page 8
Deaths - Mississippi - 1956

International List Number	Cause of Death*	Total	White	Non-white
678	Delivery with other complications of child-birth	1	1	0
681	Sepsis of childbirth and the puerperium	3	1	2
684	Puerperal pulmonary embolism	4	1	3
685	Puerperal eclampsia	3	0	3
688	Other and unspecified complications of the puerperium	1	0	1
692	Other cellulitis and abscess without mention of lymphangitis	3	3	0
698	Other local infections of skin and subcutaneous tissue	1	0	1
701	Eczema	1	0	1
702	Occupational dermatitis	1	0	1
704	Pemphigus	3	0	3
705	Erythematous conditions	7	5	2
710	Other hypertrophic and atrophic conditions of skin	5	4	1
715	Chronic ulcer of skin	3	1	2
720	Acute arthritis due to pyogenic organisms	2	0	2
722	Rheumatoid arthritis and allied conditions	11	9	2
723	Osteo-arthritis (arthrosis) and allied conditions	1	1	0
724	Other specified forms of arthritis	1	1	0
725	Arthritis, unspecified	10	7	3
733	Other diseases of bone	1	1	0
735	Displacement of intervertebral disc	1	0	1
744	Other diseases of muscle, tendon, and fascia	14	11	3
745	Curvature of spine	1	1	0
<u>750-759</u>	<u>Congenital malformations</u>	<u>247</u>	<u>142</u>	<u>105</u>
750	Monstrosity	12	8	4
751	Spina bifida and meningocele	21	13	8
752	Congenital hydrocephalus	21	8	13
753	Other nervous system and sense organs	6	1	5
754	Circulatory system	117	70	47
755	Cleft palate and harelip	2	1	1
756	Digestive system	36	22	14
757	Genito-urinary system	10	9	1
758	Bone and joint	4	1	3
759	Other and unspecified, not elsewhere classified	18	9	9
<u>760-762</u>	<u>Birth injuries, postnatal asphyxia, and atelectasis</u>	<u>387</u>	<u>218</u>	<u>169</u>
760	Intracranial and spinal injury at birth	73	36	37
761	Other birth injury	68	37	31
762	Postnatal asphyxia and atelectasis	246	145	101
<u>763-768</u>	<u>Infections of newborn</u>	<u>136</u>	<u>31</u>	<u>105</u>
763	Pneumonia	94	23	71
764	Diarrhea	27	3	24
767	Umbilical sepsis	2	0	2
768	Other sepsis	13	5	8
<u>769-776</u>	<u>Other diseases peculiar to early infancy, and immaturity unqualified</u>	<u>668</u>	<u>223</u>	<u>445</u>

Table 5 - Page 9
Deaths - Mississippi - 1956

42

International List Number	Cause of Death*	Total	White	Non-white
769	Neonatal disorders arising from maternal toxemia	15	5	10
770	Hemolytic disease of newborn (erythroblastosis)	23	9	14
771	Hemorrhagic disease of newborn	14	4	10
772	Nutritional maladjustment	21	4	17
773	Ill-defined diseases peculiar to early infancy	71	32	39
774	Immaturity with mention of any other subsidiary condition	30	10	20
776	Immaturity unqualified	494	159	335
780-795	<u>Symptoms, senility, and ill-defined conditions</u>	1,705	355	1,350
780	Certain symptoms referable to nervous system and special senses	7	2	5
782	Symptoms referable to cardiovascular and lymphatic system	46	22	24
783	Symptoms referable to respiratory system	3	1	2
784	Symptoms referable to upper gastro-intestinal tract	4	3	1
785	Symptoms referable to abdomen and lower gastro-intestinal system	5	1	4
786	Symptoms referable to genito-urinary system	2	1	1
788	Other general symptoms	5	1	4
792	Uremia unqualified	10	3	7
794	Senility without mention of psychosis	91	27	64
795	Ill-defined and unknown causes	1,532	294	1,238
800-962	<u>Accidents</u>	1,302	700	602
800-802	Railway	19	12	7
810-835	<u>Motor vehicle</u>	558	347	211
810	Collision with railway train, traffic	18	13	5
812	Traffic, to pedestrian	83	40	43
813	Traffic, to pedal cyclist	6	5	1
814	Traffic, to rider or passenger of motorcycle, in collision with nonmotor vehicle or object	1	0	1
815	Traffic, to rider or passenger of motorcycle in collision with other motor vehicle	2	2	0
816	Other traffic, two or more motor vehicles	189	135	54
818	Traffic, collision with animal or animal-drawn vehicle	3	2	1
819	Traffic, collision with fixed or unspecified object	23	12	11
821	Noncollision traffic, to rider of motorcycle	1	1	0
822	Traffic, overturning in roadway	36	21	15
823	Traffic, running off roadway	143	90	53
824	Other noncollision traffic	14	4	10
825	Traffic, unspecified nature	24	13	11
830	Nontraffic, to pedestrian	4	2	2
834	Nontraffic, while boarding and alighting	2	1	1
835	Nontraffic, other and unspecified nature	9	6	3
843	To rider of pedal cycle not involving collision with motor vehicle	1	1	0
845	Other nonmotor road vehicle	4	1	3
850	Submersion of occupant of small boat	26	16	10

Table 5 - Page 10
Deaths - Mississippi - 1956

43

International List Number	Cause of Death*	Total	White	Non-white
851	Other water transport injury by submersion	2	1	1
857	Other specified water transport	1	1	0
860	To personnel in military aircraft	8	8	0
866	Other and unspecified aircraft	3	3	0
871	Poisoning by barbituric acid and derivatives	3	3	0
872	Poisoning by aspirin and salicylates	1	1	0
874	Poisoning by other analgesic and soporific drugs	1	1	0
878	Poisoning by other and unspecified drugs	1	1	0
880	Poisoning by alcohol	1	0	1
881	Poisoning by petroleum products	6	1	5
882	Poisoning by industrial solvents	2	1	1
886	Poisoning by arsenic and antimony, and their compounds	2	1	1
888	Poisoning by other and unspecified solid and liquid substances	11	4	7
890	Poisoning by utility (illuminating) gas	3	3	0
891	Poisoning by motor vehicle exhaust gas	5	4	1
900	Fall on stairs	3	2	1
901	Fall from ladders	2	0	2
902	Other falls from one level to another	19	12	7
903	Fall on same level	12	10	2
904	Unspecified falls	65	46	19
910	Blow from falling object	29	16	13
912	Caused by machinery	28	16	12
913	Caused by cutting and piercing instruments	4	1	3
914	Caused by electric current	14	13	1
916	Caused by fire and explosion of combustible material	193	54	139
917	Caused by hot substance, corrosive liquid, and steam	13	3	10
919	Caused by firearm	63	28	35
921	Inhalation or ingestion of food causing obstruction or suffocation	23	8	15
922	Inhalation or ingestion of other object causing obstruction or suffocation	3	1	2
924	Mechanical suffocation in bed and cradle	22	1	21
925	Mechanical suffocation in other and unspecified circumstances	6	4	2
926	Lack of care of infants under 1 year of age	2	0	2
928	Other accidents caused by animals	5	1	4
929	Drowning and submersion	87	42	45
931	Excessive heat and insolation	2	1	1
932	Excessive cold	1	1	0
933	Hunger, thirst, and exposure	3	1	2
934	Cataclysm	3	1	2
935	Lightning	4	2	2
936	Other and unspecified accidents	15	9	6
950	Therapeutic misadventure in surgical treatment	10	8	2
953	Therapeutic misadventure in administration of drugs or biologicals	1	0	1
960	Late effect of motor vehicle accident	2	2	0

Table 5 - Page 11
 Deaths - Mississippi - 1956

International List Number	Cause of Death*	Total	White	Non-white
962	Late effect of other accidental injury	10	7	3
<u>963,970-979</u>	<u>Suicide</u>	<u>113</u>	<u>96</u>	<u>17</u>
970	By analgesic and soporific substances	2	2	0
971	By other solid and liquid substances	6	5	1
973	By other gases	1	1	0
974	By hanging and strangulation	11	7	4
975	By submersion (drowning)	2	1	1
976	By firearms and explosives	90	79	11
977	By cutting and piercing instruments	1	1	0
<u>964,980-985</u>	<u>Homicide</u>	<u>214</u>	<u>46</u>	<u>168</u>
981	Assault by firearm and explosive	139	35	104
982	Assault by cutting and piercing instruments	41	2	39
983	Assault by other means	18	9	9
984	Injury by intervention of police	9	0	9
985	Execution	7	0	7
<u>965,990-999</u>	<u>Injury resulting from operations of war</u>	<u>1</u>	<u>1</u>	<u>0</u>
965	Late effects of injuries due to war operations	1	1	0

* When cause of death and International List Numbers are omitted, no deaths were reported.

TABLE 6
DEATHS
MISSISSIPPI
1956

By Place of Occurrence, Reallocated to Place of Residence
By Race, with Rates

County	Place of Occurrence			Place of Residence					
	Number		Non-white	Number			Rate*		
	Total	White		Total	White	Non-white	Total	White	Non-white
STATE	19,650	9,912	9,738	19,780	9,969	9,811	9.4	8.7	10.2
Adams	454	173	281	382	137	245	10.5	7.1	14.4
Alcorn	242	200	42	237	196	41	9.4	9.1	11.4
Amite	98	45	53	147	75	72	8.7	10.4	7.4
Attala	198	115	83	206	122	84	8.9	9.9	7.7
Benton	38	16	22	56	30	26	6.6	7.0	6.2
Bolivar	612	102	510	634	127	507	10.7	7.2	12.2
Calhoun	138	99	39	168	130	38	11.1	12.4	8.3
Carroll	72	27	45	120	59	61	9.3	12.0	7.6
Chickasaw	157	99	58	172	108	64	10.0	11.0	8.6
Choctaw	73	58	15	83	61	22	8.8	9.8	6.9
Claiborne	115	19	96	147	38	109	13.1	12.2	13.4
Clarke	152	88	64	195	110	85	11.3	10.9	11.8
Clay	177	75	102	180	75	105	11.9	9.9	14.0
Coahoma	500	112	388	503	101	402	9.6	7.0	10.5
Copiah	263	115	148	302	133	169	11.0	10.7	11.3
Covington	93	67	26	127	86	41	8.3	9.0	7.1
DeSoto	173	52	121	212	64	148	8.8	7.6	9.4
Forrest	429	304	125	422	280	142	9.5	8.7	11.6
Franklin	72	47	25	99	65	34	11.2	12.3	9.7
George	79	69	10	91	78	13	8.7	8.5	10.0
Greene	59	41	18	72	51	21	9.0	8.0	13.1
Grenada	201	95	106	155	57	98	8.3	6.6	9.8
Hancock	74	61	13	99	86	13	7.6	7.3	10.0
Harrison	846	666	180	746	573	173	7.1	6.5	10.1
Hinds	1,860	973	887	1,345	674	671	7.4	6.8	9.3
Holmes	299	86	213	336	92	244	12.0	13.5	11.5
Humphreys	211	44	167	227	53	174	10.6	8.8	11.4
Issaquena	22	2	20	33	8	25	6.9	4.2	8.6
Itawamba	95	92	3	122	120	2	9.0	9.4	2.2
Jackson	285	212	73	331	250	81	7.8	7.4	9.5
Jasper	86	36	50	153	74	79	9.0	9.2	8.9
Jefferson	80	17	63	121	32	89	12.0	12.8	11.7
Jeff Davis	120	56	64	120	55	65	7.4	8.6	6.6
Jones	51.3	323	190	451	304	147	8.8	8.1	10.6
Kemper	107	45	62	125	56	69	10.5	15.1	8.4
Lafayette	161	88	73	163	94	69	10.1	11.6	8.5

Table 6 - Page 2
Deaths - Mississippi - 1956

County	Place of Occurrence			Place of Residence			Rate*		
	Number		Non-white	Number		Non-white	Rate*		Non-white
	Total	White		Total	White		Total	White	
Lamar	72	47	25	102	73	29	8.3	7.0	15.3
Lauderdale	882	552	330	674	381	293	10.8	9.7	12.9
Lawrence	57	34	23	96	65	31	8.3	9.4	6.6
Leake	115	70	45	148	92	56	9.5	11.2	7.7
Lee	402	301	101	375	272	103	10.0	10.5	9.0
Leflore	561	184	377	498	133	365	10.8	8.4	12.1
Lincoln	235	154	81	271	174	97	10.3	10.4	10.2
Lowndes	357	173	184	359	172	187	9.6	8.3	11.3
Madison	351	117	234	332	97	235	10.3	12.0	9.7
Marion	173	105	68	205	124	81	9.0	9.3	8.6
Marshall	197	64	133	207	69	138	9.0	10.6	8.4
Monroe	311	173	138	322	180	142	10.6	9.5	12.4
Montgomery	88	37	51	127	68	59	9.2	9.2	9.2
Neshoba	180	122	58	208	152	56	11.4	12.0	10.2
Newton	171	118	53	179	116	63	9.4	10.4	8.1
Noxubee	151	51	100	176	59	117	10.2	12.3	9.4
Oktibbeha	203	90	113	210	91	119	9.5	8.1	11.0
Panola	205	82	123	253	103	150	7.9	8.0	7.8
Pearl River	206	152	54	217	160	57	10.0	9.7	11.2
Perry	41	21	20	68	44	24	9.4	8.5	12.0
Pike	340	207	133	342	207	135	10.6	12.0	9.1
Pontotoc	142	117	25	165	133	32	9.1	9.0	9.1
Prentiss	151	125	26	157	129	28	8.7	8.1	12.7
Quitman	182	45	137	217	65	152	8.5	6.9	9.4
Rankin	417	204	213	233	141	92	8.5	9.2	7.6
Scott	144	88	56	160	90	70	8.0	8.2	7.8
Sharkey	98	24	74	112	25	87	9.0	7.4	9.6
Simpson	158	89	69	149	83	66	7.1	6.4	8.1
Smith	66	48	18	118	89	29	8.6	8.0	11.2
Stone	48	36	12	51	39	12	8.0	8.0	8.0
Sunflower	395	100	295	433	118	315	8.0	7.2	8.3
Tallahatchie	253	68	185	292	96	196	10.3	9.6	10.6
Tate	124	45	79	161	68	93	9.4	10.0	8.9
Tippah	114	87	27	124	96	28	7.5	7.4	7.6
Tishomingo	92	88	4	117	112	5	9.0	9.1	7.1
Tunica	221	33	188	231	34	197	10.2	7.7	10.8
Union	195	149	46	204	159	45	10.8	10.7	11.2
Walthall	118	67	51	114	65	49	8.0	8.2	7.6
Warren	635	253	382	477	201	276	12.1	8.8	16.5
Washington	742	225	517	781	215	566	10.2	6.9	12.5
Wayne	118	62	56	128	67	61	8.2	6.9	10.5
Webster	83	70	13	96	80	16	9.1	10.0	6.4

Table 6 - Page 3
Deaths - Mississippi - 1956

County	Place of Occurrence			Place of Residence					
	Number		Non-white	Number			Rate*		
	Total	White		Total	White	Non-white	Total	White	Non-white
Wilkinson	132	53	79	137	46	91	10.4	13.5	9.3
Winston	133	79	54	170	107	63	8.7	9.6	7.5
Yalobusha	127	76	51	158	97	61	12.2	14.3	9.8
Yazoo	310	108	202	344	128	216	9.6	9.3	9.7
Out-of-State	761	540	221	631	483	148	-	-	-

Cities over 10,000 Population**	Place of Occurrence			Place of Residence		
	Number		Non-white	Number		Non-white
	Total	White		Total	White	
Biloxi	257	201	56	242	186	56
Clarksdale	271	91	180	224	69	155
Columbus	240	128	112	199	92	107
Greenville	437	166	271	392	125	267
Greenwood	347	144	203	214	86	128
Gulfport	289	215	74	222	162	60
Hattiesburg	371	269	102	289	183	106
Jackson	1,606	888	718	966	536	430
Laurel	411	248	163	246	144	102
McComb	175	130	45	133	100	33
Meridian	752	487	265	467	271	196
Natchez	253	135	118	283	109	174
Pascagoula	36	24	12	99	64	35
Tupelo	239	176	63	139	92	47
Vicksburg	547	220	327	348	152	196
Total Urban#	9,312	5,243	4,069	6,704	3,564	3,140
Total Rural	10,338	4,669	5,669	13,076	6,405	6,671

* Rate per 1,000 population.

** Cities having over 10,000 population according to 1950 census.

Includes all towns of 2,500 population or more according to 1950 census.

TABLE 7
DEATHS FROM SELECTED CAUSES
MISSISSIPPI
1956

By Age, Race, and Sex, on Basis of Place of Residence, with Rates

Table 7 - Page 2
Deaths from Selected Causes - Mississippi - 1956

International List Number	Cause of Death	Total	Rate*	Total by Race	Total by Sex and Race	Total by Sex and Race	Total by Sex and Race	Age in Years										
								Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 and over	Unknown
050,051	Scarlet fever and streptococcal sore throat	1	0.05	W 0 C 0	M 1 F 0	W 1 M 0	M 1 F 0	0	0	0	0	0	0	0	0	0	0	0
055	Diphtheria	5	0.2	W 0 C 4	M 2 F 1	W 2 M 1	M 1 F 1	0	0	0	0	0	0	0	0	0	0	0
056	Whooping cough	41	1.9	W 39 C 0	M 20 F 19	W 20 M 19	M 13 F 12	6	7	0	0	0	0	0	0	0	0	0
057	Meningococcal infections	14	0.7	W 9 C 5	M 7 F 5	W 7 M 5	M 4 F 3	0	1	2	1	0	0	0	0	1	0	0
080	Acute poliomyelitis	12	0.6	W 8 C 4	M 5 F 3	W 8 M 5	M 3 F 2	0	1	1	1	0	0	0	0	1	0	0
085	Measles	10	0.5	W 9 C 0	M 4 F 4	W 9 M 4	M 4 F 3	0	1	1	1	0	0	0	0	0	0	0

Table 7 - Page 3
Deaths from Selected Causes - Mississippi - 1956

Inter-national List Number	Cause of Death	Total	Rate*	Total by Sex and Race		Age in Years		Unknown over 65 and under 15
				M	F	10-14	15-24	
100-108	Typhus and other rickettsial diseases	1	0.05	W 1	M 0	0 0	0 0	0 0
140-205	Malignant neoplasms, including neoplasms of lymphatic and hematopoietic tissues	2,432	115.3	W 1,429	M 559	11 9	7 7	1 1
210-239	Benign and unspecified neoplasms	43	2.0	W 20	M 23	16	8 1	0 0
240-289	Allergic, endocrine system, metabolic and nutritional diseases	312	14.8	W 161	M 151	51	65	13
260	Diabetes mellitus	225	10.7	W 116	M 109	74	12	12
290-299	Diseases of the blood and blood-forming organs	52	2.5	W 26	M 26	8	11	3

Table 7 - Page 4
Deaths from Selected Causes - Mississippi - 1956

International List Number	Cause of Death	Total	Rate*	Total by Race	Total by Sex and Race	Age in Years	Unknown	
							65 and over	65 and under
290-293	Anemias	36	1.7	W 13	M 7	0	6	0
	Mental, psycho-neurotic, and personality disorders	26	1.2	W 14	M 13	1	5	0
300-326	Vascular lesions affecting central nervous system	2,550	120.9	W 1,229	M 627	1	3	0
320-324	Other diseases of the nervous system and sense organs	169	8.0	W 81	M 53	1	1	0
340-398	Nonmeningococcal meningitis	45	2.1	W 27	M 13	1	0	0
	Diseases of the circulatory system	6,697	317.4	W 4,043	M 2,522	1	0	0
400-468				C 2,654	F 1,397	1	0	0

Table 7 - Page 5
Deaths from Selected Causes - Mississippi - 1956

International List Number	Cause of Death	Total	Rate*	Total by Sex and Race	Total by Sex and Race	Total by Sex and Race	Age in Years											
							5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 and over	Unknown			
400-402	Rheumatic fever	13	0.6	W 6 C 7	M 4 F 3	N 75 F 70	43	32	37	33	13	68	200	372	1,203	0		
	Chronic rheumatic heart disease	145	6.9	W 2,786 C 1,311	M 1,859 F 713	N 1,927 F 598	1,859	1,110	1,110	1,000	629	420	722	105	789	2		
410-416	Arteriosclerotic and degenerative heart disease	4,097	194.2	W 225 C 197	M 157 F 83	N 68 F 83	157	91	90	100	22	22	22	20	27	99	0	
420-422	Other diseases of the heart	422	20.0	W 617 C 823	M 294 F 323	N 407 F 416	294	200	200	200	5	5	5	1	44	233	0	
430-434	Hypertension with heart disease	1,440	68.2	W 47 F 43	M 41 F 55	N 53 F 53	47	0	0	0	0	0	0	0	365	265	0	
440-443	Hypertension without mention of heart	198	9.4	W 90 C 108	M 43 F 53	N 53 F 53	43	0	0	0	0	0	0	0	3	33	0	
444-447														4	6	7	34	0
														4	6	7	30	0

Table 7 - Page 6
Deaths from Selected Causes - Mississippi - 1956

Inter-national list number	Cause of Death	Total	Rate*	Total by Sex and Race	Total by Sex and Race	Age in Years	Unknown	
							5 over 50 and over 65	Under 5
70-527	Diseases of the respiratory system	861	40.8	W 370 M 491 C 0	M 217 F 269 C 222	23 153 20 113 89	2 1 5 1 1 1 1	2 1 5 1 1 1 1
80-483	Influenza	103	4.9	W 77 C 0	M 26 F 43 P 34	17 3 12 9 4	0 0 0 0 0	0 0 0 0 0
30-493	Pneumonia (excluding pneumonia of newborn)	575	27.2	W 351 C 0	M 120 F 104 M 184 F 167	17 8 20 26 20	0 1 3 2 1	0 1 3 2 1
10-502	Bronchitis	29	1.4	W 15 C 14	M 12 F 10 M 3	1 0 7 0 3	0 0 0 0 0	0 0 0 0 0
0-587	Diseases of the digestive system	615	22.1	W 305 M 310 C 0	M 172 F 133 M 169 F 141	10 8 20 54	1 1 2 6 3 2	1 1 2 6 3 2
0, 541	Ulcer of stomach and duodenum	76	3.6	W 54 C 22	M 43 F 11 M 15 F 7	0 0 0 0 0	0 0 0 0 0	0 0 0 0 0

Table 7 - Page 7
Deaths from Selected Causes - Mississippi - 1956

International List Number	Cause of Death	Total	Total by Sex and Race	Total by Sex and Race	Age in Years		Unknown 65 and over
					5-9	10-14	
550-553	Appendicitis	24	1.1	W 6 M 4	0 0	0 0	3 0
560, 561, 570	Intestinal obstruction and hernia	93	4.4	W 18 M 11	0 1	0 1	0 0
543, 71,572	Gastritis, duodenitis, enteritis & colitis, except diarrhea of the newborn	194	9.2	W 40 M 25	1 0	1 0	0 0
581	Cirrhosis of liver	95	4.5	W 68 M 48	0 0	0 0	0 0
20-637	Diseases of the genito-urinary system	637	20.2	W 254 M 162	5 1	1 1	1 1
10-594	Nephritis and nephrosis	470	19.4	W 161 M 76	0 0	0 1	0 0
				M 260 F 223	0 2	1 2	0 0
				M 101 F 148	1 5	1 1	1 1
				M 149 F 249	0 1	1 1	1 1

Table 7 - Page 8
Deaths from Selected Causes - Mississippi - 1956

International List Number	Cause of Death	Total	Rate*	Total by Sex and Race		Age in Years		Unknown
				Total by Race	Rate by Sex	Total	Rate*	
610	Hyperplasia of prostate	77	3.6	W 31 M 46	31 F 46 M 0	10-14	0.000	0
	Complications of pregnancy, childbirth, and the puerperium	74	1.2	W 12 C 62	12 F 62 M 0	15-24	0.000	0
640-689*	Diseases of the skin and cellular tissue	24	1.1	W 13 C 11	13 F 11 M 0	25-34	0.000	0
690-716	Diseases of the bones & organs of movement	42	2.0	W 31 C 11	31 F 11 M 0	35-44	0.000	0
720-749	Congenital malformations	247	11.7	W 142 C 105	142 F 105 M 0	45-54	0.000	0
"50-759	Certain diseases of early infancy	1,191	19.2	W 719 C 472	719 F 472 M 0	55-64	0.000	0
60-776*						65 and over	38 C 0	30 O 0

Table 7 - Page 9
Deaths from Selected Causes - Mississippi - 1956

International List Number	Cause of Death	Total by Sex and Race	Rate*	Total by Sex and Race	Age in Years									
					6	7-10	11-14	15-17	18-24	25-34	35-44	45-54	55-64	65 and over
780-795	Symptoms, senility, and ill-defined conditions	1,705	80.8	W 335 C 1,350	M 220 F 135	M 14 F 14	M 11 F 17	M 5 F 7	M 1 F 1	M 13 F 25	M 7 F 11	M 37 F 13	M 47 F 10	M 110 F 63
800-999	Accidents, poisonings and violence	1,630	77.2	W 843 C 787	M 643 F 569	M 7 F 2	M 14 F 22	M 15 F 26	M 15 F 22	M 105 F 20	M 115 F 20	M 54 F 15	M 103 F 22	M 367 F 312
810-835	Motor vehicle accidents	558	26.4	W 347 C 211	M 269 F 167	M 0 F 3	M 12 F 7	M 6 F 7	M 7 F 7	M 15 F 11	M 51 F 21	M 29 F 15	M 31 F 17	M 15 F 15
300-802, 340-962	All other accidents	744	35.3	W 353 C 391	M 252 F 101	M 7 F 11	M 10 F 10	M 8 F 8	M 10 F 8	M 12 F 11	M 44 F 39	M 33 F 30	M 19 F 16	M 49 F 40
963, 170-979	Suicide	113	5.4	W 96 C 17	M 79 F 17	M 0 F 0	M 0 F 0	M 0 F 0	M 0 F 0	M 0 F 0	M 0 F 4	M 9 F 4	M 5 F 5	M 33 F 33
964, 80-985	Homicide	214	10.1	C 168	M 42 F 4	M 0 F 0	M 2 F 0	M 0 F 0	M 0 F 0	M 0 F 1	M 6 F 1	M 15 F 1	M 10 F 1	M 3 F 3

Crude death rate per 1,000 population. Specific death rates per 100,000 population. Maternal death rate and rates for infant causes (760-776) per 1,000 live births.

TABLE 8
DEATHS FROM SELECTED CAUSES
MISSISSIPPI
1956

By Month of Occurrence on Basis of Place of Residence

Cause of Death and International List Number	Total	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
<u>TOTAL ALL CAUSES</u>	<u>19,780</u>	<u>1,888</u>	<u>1,609</u>	<u>1,704</u>	<u>1,588</u>	<u>1,596</u>	<u>1,578</u>	<u>1,525</u>	<u>1,558</u>	<u>1,601</u>	<u>1,731</u>	<u>1,831</u>	
Dysentery (045-048)	54	8	6	3	4	7	6	2	5	4	5	1	
Diphtheria (055)	5	1	0	0	0	0	1	0	1	0	0	0	
Whooping cough (C56)	41	0	2	5	3	6	5	10	2	4	1	1	
Acute poliomyelitis (080)	12	1	0	0	1	4	0	3	0	2	1	0	
Measles (085)	10	0	1	1	3	1	0	0	0	0	1	2	
Influenza (480-483)	103	17	22	12	10	9	7	2	4	2	7	9	
Lobar pneumonia (490)	149	28	20	14	10	8	13	10	6	7	14	13	
Bronchopneumonia (491)	248	30	27	27	19	18	13	13	16	17	14	26	
Other & unspec. pneumonia (492, 493)	178	24	18	27	12	11	12	4	5	5	12	16	
Diarrhea and enteritis (-2) (5710)	150	19	2	2	4	18	21	13	12	4	17	21	
Diarrhea and enteritis (-2) (5711)	28	2	4	1	3	1	3	1	4	3	1	1	
Accidents (800-962)	1,302	116	72	106	95	117	104	119	100	107	96	126	
Motor vehicle accidents (810-835)	558	43	29	40	30	52	37	47	42	58	54	52	

TABLE 9
DEATHS UNDER 1 YEAR
MISSISSIPPI
1956

For Selected Causes, by Race, by Age, on Basis of Place of Residence, with Rates

Cause of Death and International List Number	Total	Rate*	Total by cause	DAYS												MONTHS						
				Under 1 day	1- 6	7- 13	14- 20	21- 27	Under 28 days	1	2	3	4	5	6	7	8	9	10	11		
<u>ALL CAUSES</u>	<u>2,353</u>	<u>37.8</u>	<u>W C</u> <u>734 1,619</u>	<u>301</u>	<u>224</u>	<u>27</u>	<u>18</u>	<u>11</u>	<u>58</u>	<u>23</u>	<u>15</u>	<u>16</u>	<u>15</u>	<u>16</u>	<u>15</u>	<u>14</u>	<u>5</u>	<u>5</u>	<u>4</u>	<u>7</u>		
Syphilis (020-029)	0	-	W C 0 0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Diphtheria (055)	0	-	W C 2 0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Whooping cough (056)	27	0.4	W C 25 4	0	0	0	0	1	1	2	6	0	4	1	2	3	2	0	3	2		
Measles (085)	4	0.1	W C 4 0	0	0	0	0	1	1	0	0	0	0	1	0	0	0	0	0	0		
Influenza (480-483)	25	0.4	W C 4 21	0	1	0	0	0	0	1	2	2	3	2	1	3	1	0	0	0		
**Pneumonia (490-493)	181	2.9	W C 25 156	0	0	0	0	0	0	0	7	1	2	3	1	6	4	0	1	0		
Other respiratory diseases (470-475, 500-527)	4.9	0.8	W C 24 25	3	12	0	0	1	0	15	4	1	0	3	1	0	0	0	1	0		
**Diarrhea and enteritis (5710)	137	2.2	W C 14 123	0	0	0	0	0	0	0	3	1	0	2	3	2	0	1	1	0		

Table 9 - Page 2
Deaths Under 1 Year - Mississippi - 1956

Cause of Death and International List Number	Total	Rate*	Total by Race	Total by Race	DAYS										MONTHS						
					Under 1 day	1- 6	7- 13	14- 20	21- 27	Under 28 days	1	2	3	4	5	6	7	8	9	10	11
Congenital malfor- mations (750-759)	192	3.1	W C	115 77	22	33	11	6	5	77	9	7	6	5	2	4	1	2	2	0	0
Birth injuries (760, 761)	141	2.3	W C	73 68	42	30	1	0	0	73	0	0	0	0	0	0	0	0	1	1	2
Postnatal asphyxia & atelectasis (762)	246	4.0	W C	145 101	89	54	1	0	0	144	0	0	1	0	0	0	0	0	0	0	0
**Infections of the newborn (763-768)	136	2.2	W C	31 105	2	18	4	6	1	31	0	0	0	0	0	0	0	0	0	0	0
Hemolytic disease of the newborn (770)	23	0.4	W C	9 14	2	6	1	0	0	105	0	0	0	0	0	0	0	0	0	0	0
All other defined dis. of early infancy (769, 771, 772)	50	0.8	W C	13 37	6	2	2	0	2	10	2	0	1	2	0	0	0	0	0	0	0
Violent & accidental deaths (800-999)	74	1.2	W C	12 62	1	1	1	1	1	11	12	9	3	8	7	2	0	0	0	1	1
Other specified causes	148	2.4	W C	38 110	2	4	0	1	3	10	5	2	1	2	4	2	3	2	0	3	4
III-defined (773-776, 780-793, 795)	920	14.8	W C	229 691	132	64	6	4	2	208	6	6	3	3	1	0	0	0	1	0	1

* Rate per 1,000 live births.

** Pneumonia and diarrhea of the newborn (under 28 days) included in infections of the newborn (763-768).

TABLE 10
DEATHS DUE TO INJURIES WHICH OCCURRED IN MISSISSIPPI
1956

By Place of Injury by Type of Injury

County	Motor Vehicle Accidents (310-835)		Other Transport Accidents (800-8C2, 840-866)		Non-Transport Accidents (870-936)		Other and Violences (270-285)		Rural	
	Total	Urban	Total	Urban	Rural	Total	Urban	Rural	Total	Rural
STATE	588	56	532	73	9	64	14	8	167	489
Adams	8	4	4	1	0	1	7	3	4	0
Alcorn	4	1	3	1	0	0	3	5	2	2
Anite	1	0	1	0	0	0	0	0	1	0
Attala	7	0	7	0	0	0	1	3	1	0
Benton	0	0	0	0	0	0	0	0	0	0
Bolivar	8	0	5	0	5	25	0	2	23	0
Calhoun	3	0	3	0	0	0	3	3	2	1
Carroll	4	0	4	0	0	0	10	3	12	3
Chickasaw	5	0	5	1	0	0	4	4	8	0
Choctaw	1	0	1	0	0	0	0	0	0	0
Claiborne	5	0	5	0	0	0	6	0	6	0
Clarke	11	0	11	0	0	0	3	3	7	0
Clay	2	0	2	0	0	0	6	3	10	7
Coahoma	15	0	15	8	6	10	16	7	14	5
Copiah	10	2	10	0	0	0	13	4	14	0
Covington	6	0	6	1	0	0	6	0	6	0
DeSoto	9	0	9	0	0	0	14	0	14	0
Forrest	10	0	10	1	0	0	10	0	10	0
Franklin	4	0	4	0	0	0	5	0	5	0
George	6	0	6	4	2	2	0	2	4	0
Greene	4	0	4	2	0	0	3	0	3	0
Grenada	12	0	12	9	3	3	0	0	0	9
Hancock	9	0	9	0	0	0	0	0	0	0
Harrison	20	0	20	5	0	0	15	0	15	0

Table 10 - Page 2
Deaths Due to Injuries which Occurred in Mississippi - 1956

County	Motor Vehicle Accidents (810-835)		Other Transport Accidents (800-802, 840-866)		Non-Transport Accidents (870-936)		Other and Non-Transport Accidents (870-926)		Violences (970-985)	
	Total	Urban	Rural	Total	Urban	Rural	Total	Urban	Rural	Total
Hinds	30	13	17	2	0	2	20	15	5	8
Holmes	8	1	7	1	0	1	1	1	0	1
Humphreys	7	2	5	2	0	2	12	13	1	1
Issaquena	2	0	1	0	0	0	1	0	0	1
Itawamba	5	0	5	0	0	0	6	3	3	2
Jackson	13	12	1	13	10	3	21	19	2	2
Jasper	9	9	2	0	0	0	6	3	3	2
Jefferson	2	2	0	0	0	0	1	0	1	1
Jeff Davis	2	1	1	0	0	0	1	0	1	1
Jones	20	18	2	0	0	0	12	12	0	12
Kemper	5	5	4	1	0	1	6	2	4	6
Lafayette	4	4	0	0	0	0	23	22	5	22
Lamar	1	1	0	0	0	0	7	6	1	7
Lauderdale	29	23	6	1	1	0	13	12	1	13
Lawrence	3	3	0	0	0	0	1	1	0	1
Leake	6	6	0	0	0	0	1	1	0	1
Lee	6	6	0	0	0	0	1	1	0	1
Leflore	5	4	1	1	1	0	8	7	1	8
Lincoln	10	10	0	2	2	0	10	15	5	17
Townships	10	10	0	1	1	0	1	1	0	1
Madison	15	15	0	1	1	0	1	1	0	1
Marion	10	9	1	0	0	0	1	1	0	1
Marshall	9	9	0	0	0	0	1	1	0	1
Monroe	4	4	0	0	0	0	1	1	0	1
Montgomery	6	6	0	0	0	0	1	1	0	1
Neshoba	2	2	0	0	0	0	1	1	0	1
Newton	4	4	0	0	0	0	1	1	0	1
Noxubee	1	1	0	0	0	0	1	1	0	1
Oktibbeha	4	4	0	0	0	0	1	1	0	1
Panola	4	4	0	0	0	0	1	1	0	1

Table 10 - Page 3
Deaths Due to Injuries which Occurred in Mississippi - 1956

County	Motor Vehicle Accidents (810-835)			Other Transport Accidents (800-802, 840-866)			Non-Transport Accidents (870-936)			Other and Transport Accidents (870-936)			Violences (970-985)		
	Total	Urban	Rural	Total	Urban	Rural	Total	Urban	Rural	Total	Urban	Rural	Total	Urban	Rural
Pearl River	9	3	6	2	0	2	11	4	7	6	3	4	2	3	0
Perry	0	0	0	1	0	1	10	5	5	7	2	2	5	0	0
Pike	13	0	13	1	1	0	0	0	0	4	0	0	4	0	0
Pontotoc	6	0	6	0	0	0	6	0	6	4	3	3	0	0	0
Prentiss	7	0	7	0	0	0	3	0	3	0	3	0	0	0	0
Quitman	5	0	5	0	0	0	7	0	7	4	1	0	4	1	0
Rankin	11	0	11	0	0	0	18	0	18	8	0	0	8	3	3
Scott	4	4	0	0	0	0	10	3	7	6	3	3	3	0	0
Sharkey	5	5	0	0	0	0	10	3	7	4	0	0	4	0	0
Simpson	3	3	0	0	0	0	10	3	5	10	0	0	10	0	0
Smith	3	3	0	0	0	0	12	2	10	11	7	6	2	0	0
Stone	4	4	0	0	0	0	12	9	6	4	4	4	1	1	0
Sunflower	12	1	11	2	0	2	12	9	6	10	0	0	10	0	0
Tallahatchie	9	9	0	2	2	0	12	4	4	10	0	0	10	0	0
Tate	2	2	0	0	0	0	12	0	0	10	0	0	10	0	0
Tippah	3	3	0	0	0	0	12	0	0	10	0	0	10	0	0
Tishomingo	4	4	0	0	0	0	12	0	0	10	0	0	10	0	0
Tunica	12	0	12	0	0	0	12	0	0	10	0	0	10	0	0
Union	5	5	0	0	0	0	12	0	0	10	0	0	10	0	0
Walton	4	4	0	0	0	0	12	0	0	10	0	0	10	0	0
Warren	7	2	5	5	4	1	18	7	11	14	3	5	12	7	0
Washington	22	4	18	6	0	0	22	0	0	20	0	0	20	0	0
Wayne	7	0	7	0	0	0	7	0	0	6	3	3	5	2	3
Webster	7	7	0	0	0	0	7	0	0	6	1	0	6	1	0
Wilkinson	6	1	5	1	0	0	6	1	5	4	1	0	4	1	0
Winston	1	0	1	0	0	0	1	0	0	0	0	0	0	0	0
Yalobusha	5	0	5	0	0	0	5	0	0	4	1	0	4	1	0
Yazoo	7	0	7	0	0	0	7	0	0	6	2	1	6	1	0

TABLE 11
MARRIAGES AND DIVORCES
MISSISSIPPI
1956

By County, by Race

County	Marriages						Divorces		
	Occurrence			Residence of Bride			Total	White	Non-white
	Total	White	Non-white	Total	White	Non-white			
STATE	66,223	48,140	18,083	23,011	12,677	10,334	5,027	3,452	1,575
Adams	740	567	173	329	190	139	165	109	56
Alcorn	7,847	6,989	858	298	248	50	80	64	16
Amite	251	173	78	165	87	78	32	17	15
Attala	207	122	85	221	135	86	66	43	23
Benton	237	87	150	64	37	27	10	7	3
Bolivar	664	191	473	618	170	448	71	41	30
Calhoun	150	109	41	156	124	32	38	25	13
Carroll	103	60	43	112	61	51	15	9	6
Chickasaw	210	125	85	190	120	70	29	21	8
Choctaw	93	62	31	74	50	24	14	12	2
Claiborne	202	104	98	114	27	87	31	6	25
Clarke	407	314	93	201	118	83	31	20	11
Clay	315	189	126	167	73	94	35	17	18
Coahoma	924	388	536	571	145	426	64	40	24
Copiah	260	129	131	267	128	139	80	43	37
Covington	170	127	43	180	132	48	30	24	6
DeSoto	9,226	6,112	3,114	268	80	188	22	12	10
Forrest	595	439	156	618	448	170	237	167	70
Franklin	93	56	37	95	52	43	17	7	10
George	1,944	1,753	191	142	130	12	20	15	5
Greene	616	565	51	96	72	24	17	13	4
Grenada	212	113	99	210	109	101	56	35	21
Hancock	1,082	1,025	57	97	79	18	72	66	6
Harrison	2,129	1,843	286	1,165	927	238	552	451	101
Hinds	1,752	1,012	740	1,924	1,160	764	372	258	114
Holmes	336	80	256	357	63	294	26	8	18
Humphreys	191	61	130	187	64	123	29	22	7
Issaquena	40	26	14	28	11	17	4	3	1
Itawamba	1,286	1,259	27	89	83	6	28	24	4
Jackson	2,702	1,762	940	478	351	127	161	132	29
Jasper	151	66	85	173	78	95	26	16	10
Jefferson	306	253	53	88	27	61	14	4	10
Jeff Davis	181	102	79	122	67	55	21	9	12
Jones	701	510	191	634	460	174	165	121	44
Kemper	135	46	89	102	35	67	16	10	6
Lafayette	149	81	68	174	87	87	34	13	21

Table 11 - Page 2
 Marriages and Divorces - Mississippi - 1956

County	Marriages						Divorces		
	Occurrence			Residence of Bride			Total	White	Non-white
	Total	White	Non-white	Total	White	Non-white			
Lamar	274	229	45	179	137	42	28	23	5
Lauderdale	2,160	1,413	747	728	483	245	194	146	48
Lawrence	119	83	36	168	113	55	25	18	7
Leake	198	116	82	220	126	94	34	23	11
Lee	567	436	131	459	318	141	83	68	15
Leflore	624	191	433	612	196	416	65	36	29
Lincoln	341	219	122	245	155	90	65	34	31
Lowndes	3,777	3,042	735	445	233	212	83	54	29
Madison	357	110	247	284	70	214	66	16	50
Marion	335	191	144	301	185	116	84	61	23
Marshall	1,007	664	343	236	69	167	29	17	12
Monroe	542	409	133	359	225	134	85	68	17
Montgomery	170	95	75	148	68	80	27	17	10
Neshoba	261	188	73	225	162	63	45	35	10
Newton	204	140	64	185	124	61	28	15	13
Noxubee	318	115	203	178	59	119	10	4	6
Oktibbeha	416	285	131	213	82	131	27	14	13
Panola	439	242	197	366	148	218	46	22	24
Pearl River	836	664	172	284	212	72	120	93	27
Perry	129	109	20	82	61	21	24	16	8
Pike	709	437	272	440	237	203	100	66	34
Pontotoc	221	186	35	161	126	35	37	28	9
Prentiss	456	421	35	218	192	26	54	48	6
Quitman	417	168	249	317	90	227	44	34	10
Rankin	431	314	117	197	102	95	41	35	6
Scott	285	174	111	229	136	93	32	25	7
Sharkey	110	44	66	77	26	51	9	2	7
Simpson	212	143	69	175	118	57	57	42	15
Smith	109	97	12	148	107	41	34	31	3
Stone	158	140	18	73	61	12	11	9	2
Sunflower	608	308	300	497	146	351	37	28	9
Tallahatchie	408	108	300	311	76	235	40	30	10
Tate	700	548	152	180	67	113	19	7	12
Tippah	263	219	44	185	154	31	12	11	1
Tishomingo	3,833	3,554	279	187	177	10	34	33	1
Tunica	696	388	308	257	38	219	21	7	14
Union	254	212	42	209	170	39	34	29	5
Walthall	344	222	122	113	60	53	47	29	18
Warren	806	573	233	431	246	185	141	73	68
Washington	3,529	2,716	813	838	354	484	195	126	69
Wayne	421	277	144	176	109	67	36	23	13
Webster	106	85	21	103	77	26	11	6	5
Wilkinson	745	630	115	115	39	76	23	9	14
Winston	180	107	73	160	96	64	37	19	18
Yalobusha	165	94	71	158	86	72	20	15	5
Yazoo	276	134	242	365	133	232	53	23	30

TABLE 12
DIVORCES
MISSISSIPPI
1956

By Cause of Divorce

Cause	Total	White	Non-white
<u>TOTAL</u>	<u>5,027</u>	<u>3,452</u>	<u>1,575</u>
Cruel and inhuman treatment	2,370	1,909	461
Desertion	1,919	1,024	895
Adultery	277	145	132
Drunkenness	317	271	46
Penitentiary	37	32	5
Insanity	5	4	1
Annulment	29	22	7
Bigamy	22	12	10
Miscellaneous and other	20	13	7
Not stated	31	20	11

By Number of Minor Children

Minor Children	Total	White	Non-white
None	2,673	1,751	922
One	1,051	786	265
Two	601	442	159
Three or more	613	410	203
Not stated	89	63	26

By Number of Years Married

Number Years Married	Total	White	Non-white
<u>TOTAL</u>	<u>5,027</u>	<u>3,452</u>	<u>1,575</u>
Under one year	396	350	46
One year	658	563	95
Two years	522	395	127
Three years	380	278	102
Four years	307	227	80
5-9 years	1,096	729	367
10-19 years	978	588	390
Twenty years and over	612	282	330
Not stated	78	40	38

TABLE 13
 BIRTHS, DEATHS, MARRIAGES
 MISSISSIPPI
 1956

By Month of Occurrence

Month	Births	Deaths	Marriages
<u>TOTAL</u>	<u>62,178</u>	<u>19,780</u>	<u>66,223</u>
January	5,511	1,888	4,950
February	5,333	1,609	4,388
March	5,313	1,704	5,135
April	4,468	1,588	4,792
May	4,508	1,578	4,867
June	4,571	1,596	6,622
July	5,564	1,571	5,738
August	5,701	1,525	5,751
September	5,531	1,558	5,852
October	5,373	1,601	5,365
November	5,093	1,731	5,416
December	5,212	1,831	7,347

TABLE 14
LIVE BIRTHS, DEATHS, MATERNAL DEATHS, AND DEATHS UNDER 1 YEAR OF AGE
MISSISSIPPI
1947-1956

Shown by Year and Race, on Basis of Place of Residence, with Rates*

YEAR	LIVE BIRTHS			DEATHS			MATERNAL DEATHS			DEATHS UNDER 1 YEAR			DEATHS UNDER 1 MONTH		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
1947 No.	66291	32423	33868	20007	9238	10769	175	50	125	2428	1009	1419	1595	750	845
Rate	28.6	27.5	29.9	8.6	7.8	9.5	2.6	1.5	3.7	36.6	31.1	41.9	24.1	23.1	24.9
1948 No.	65291	29666	35625	20183	9216	10967	177	38	139	2470	988	1482	1674	773	901
Rate	28.0	24.9	31.2	8.6	7.7	9.6	2.7	1.3	3.9	37.8	33.3	41.6	25.6	26.0	25.3
1949 No.	66128	28785	37342	20489	9233	11256	147	32	115	2612	900	1712	1594	661	933
Rate	28.1	24.0	32.5	8.7	7.7	9.8	2.2	1.1	3.1	39.5	31.3	45.8	24.1	23.0	25.0
1950 No.	64427	27188	37239	20567	9466	11101	165	30	135	2348	763	1585	1446	560	886
Rate	29.6	22.9	37.6	9.4	8.0	11.2	2.6	1.1	3.6	36.4	28.1	42.6	22.4	20.6	23.8
1951 No.	65512	27798	37714	21007	9593	11414	143	28	115	2589	814	1775	1679	626	1053
Rate	30.1	23.4	38.1	9.6	8.1	11.5	2.2	1.0	3.0	39.5	29.3	47.1	25.6	22.5	27.9
1952 No.	64509	28073	36436	20657	9387	11270	119	15	104	2592	760	1832	1658	581	1077
Rate	29.6	23.6	36.8	9.5	7.9	11.4	1.8	0.5	2.8	40.2	27.1	50.3	25.7	20.7	29.6
1953 No.	64061	28312	35749	20492	9495	10997	119	24	95	2534	807	1727	1630	635	995
Rate	29.4	23.8	36.1	9.4	8.0	11.1	1.8	0.8	2.6	39.6	28.5	48.3	25.4	22.4	27.8
1954 No.	64748	28781	35967	19353	9453	9900	104	11	93	2360	744	1616	1521	601	920
Rate	29.7	24.2	36.3	8.9	8.0	10.0	1.6	0.4	2.6	36.4	25.8	44.9	23.5	20.9	25.6
1955 No.	63244	27546	35698	19403	9501	9902	76	12	64	2338	680	1658	1440	527	913
Rate	29.0	23.2	36.0	8.9	8.0	10.0	1.2	0.4	1.8	37.0	24.7	45.4	22.8	19.1	25.6
1956 No.	62178	27278	34900	19780	9969	9811	74	12	62	2353	734	1619	1451	581	870
Rate	29.5	23.8	36.2	9.4	8.7	10.2	1.2	0.4	1.8	37.8	26.9	46.4	23.3	21.3	24.9

* Population for 1947, 1948 and 1949, estimates as of July 1, based on 1930 and 1940 censuses; for 1950-1955, the April 1, 1950, United States Census figures; for 1956, estimate as of July 1, prepared by Dr. Harold A. Pedersen, Sociology Department, Mississippi State College, with adjustments by Statistical Service, Mississippi State Board of Health. Birth and death rates per 1,000 population. Maternal and infant death rates per 1,000 live births.

TABLE 15
DEATHS FROM HEART DISEASES, NEPHRITIS (ALL FORMS)
VASCULAR LESIONS, CANCER, AND TUBERCULOSIS (ALL FORMS)
MISSISSIPPI
1947-1956

Shown by Year and Race, on Basis of Place of Residence, with Rates*

YEAR	HEART DISEASES				NEPHRITIS (ALL FORMS)				VASCULAR LESIONS				CANCER				TUBERCULOSIS (ALL FORMS)			
	Non-white		White		Total		Non-white		White		Total		Non-white		White		Total		Non-white	
	Total	White	Total	White	Total	White	Total	White	Total	White	Total	Total	White	Total	White	Total	White	Total	White	Total
1947 No.	4,174	2,416	1,758	1,748	680	1,068	1,957	903	1,054	1,799	1,055	744	735	239	496	20.2	31.8	20.2	43.8	
Rate	180.4	204.7	155.1	175.6	57.6	94.2	84.6	76.5	93.0	77.8	89.4	65.6	668	206	211	462	17.3	28.6	17.3	40.5
1948 No.	4,339	2,460	1,879	1,646	666	959	1,956	878	1,078	1,906	1,148	758	668	206	211	40.5	44.0	15.8	34.0	
Rate	186.1	206.7	164.6	69.7	56.0	84.0	83.9	73.8	94.5	81.8	96.4	66.4	66.4	24.7	24.7	39.1	34.0	15.8	34.0	
1949 No.	5,496	3,060	2,436	729	236	493	2,093	976	1,117	2,039	1,205	834	581	190	190	40.5	40.5	15.8	34.0	
Rate	233.9	254.9	212.0	31.0	19.7	42.9	89.1	81.3	97.2	86.8	100.4	72.6	72.6	24.7	24.7	39.1	34.0	15.8	34.0	
1950 No.	5,546	3,158	2,388	733	246	487	2,200	971	1,229	2,143	1,333	810	558	179	179	38.3	38.3	15.1	38.3	
Rate	254.5	265.7	241.1	33.6	20.7	49.2	101.0	81.7	124.1	98.4	112.2	81.8	81.8	25.6	25.6	37.9	37.9	15.1	38.3	
1951 No.	5,626	3,134	2,492	681	210	471	2,344	1,089	1,255	2,186	1,326	860	497	158	158	34.2	34.2	13.3	34.2	
Rate	258.2	263.7	251.6	31.2	17.7	47.6	107.6	91.6	126.7	100.3	111.6	86.8	86.8	22.8	22.8	33.9	33.9	13.3	34.2	
1952 No.	5,876	3,256	2,620	658	214	444	2,386	1,098	1,283	2,189	1,285	904	418	133	133	28.8	28.8	11.2	28.8	
Rate	269.7	273.9	264.6	30.2	18.0	44.8	109.5	92.4	130.1	100.5	108.1	91.3	91.3	19.2	19.2	28.8	28.8	11.2	28.8	
1953 No.	5,758	3,183	2,575	474	164	310	2,404	1,125	1,279	2,230	1,348	882	308	85	85	22.5	22.5	7.2	22.5	
Rate	264.3	267.8	260.0	21.8	13.8	31.3	110.3	94.6	129.2	102.3	113.4	89.1	89.1	14.1	14.1	22.3	22.3	7.2	22.5	
1954 No.	5,708	3,325	2,383	448	169	279	2,422	1,176	1,246	2,332	1,420	912	226	80	80	14.7	14.7	6.7	14.7	
Rate	262.0	279.7	240.6	20.6	14.2	28.2	111.2	98.9	125.8	107.0	119.5	92.1	92.1	10.4	10.4	14.6	14.6	6.7	14.6	
1955 No.	5,969	3,517	2,452	401	144	257	2,447	1,169	1,278	2,317	1,402	915	211	66	66	14.6	14.6	5.6	14.6	
Rate	273.9	295.9	247.6	18.4	12.1	26.0	112.3	98.3	129.0	106.3	118.0	92.4	92.4	11.2	11.2	14.6	14.6	5.6	14.6	
1956 No.	6,104	3,703	2,401	410	161	249	2,550	1,229	1,321	2,432	1,439	993	199	80	80	11.9	11.9	7.0	12.4	
Rate	289.3	322.8	249.4	19.4	14.0	25.9	120.9	107.1	137.2	115.3	125.4	103.2	103.2	9.4	9.4	12.4	12.4	7.0	12.4	

* Rate per 100,000 population.

TABLE 16
BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
STATE OF MISSISSIPPI, 1956

69

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	62806	27785	35021	62178	27278	34900	29.5	23.8	36.2
Fetal deaths	2693	1038	1655	2650	1008	1642	42.6	37.0	47.0
Deaths from all causes	19650	9912	9738	19780	9969	9811	9.4	8.7	10.2
Deaths under 1 year	2338	717	1621	2353	734	1619	37.8	26.9	46.4
Deaths under 1 month	1439	569	870	1451	581	870	23.3	21.3	24.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	193	73	120	199	80	119	9.4	7.0	12.4
Tbc., respiratory (001-008)	172	69	103	176	75	101	8.3	6.5	10.5
Syphilis (020-029)	42	10	32	44	13	31	2.1	1.1	3.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	55	16	39	54	16	38	2.6	1.4	3.9
Scarlet fever & streptococcal sore throat (050, 051)	2	2	0	1	1	0	0.05	0.1	-
Diphtheria (055)	4	1	3	5	1	4	0.2	0.1	0.4
Whooping cough (056)	41	2	39	41	2	39	1.9	0.2	4.0
Meningococcal infections (057)	15	6	9	14	5	9	0.7	0.4	0.9
Acute poliomyelitis (080)	12	4	8	12	4	8	0.6	0.3	0.8
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	10	1	9	10	1	9	0.5	0.1	0.9
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	2338	1369	969	2432	1439	993	115.3	125.4	103.2
Diabetes (260)	224	119	105	225	116	109	10.7	10.1	11.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	2527	1217	1310	2550	1229	1321	120.9	107.1	137.2
Heart diseases (410-443)	6104	3731	2373	6104	3703	2401	289.3	322.8	249.4
Arteriosclerosis (450)	247	160	87	247	161	86	11.7	14.0	8.9
Influenza (480-483)	103	25	78	103	26	77	4.9	2.3	8.0
#Pneumonia (490-493)	581	227	354	575	224	351	27.2	19.5	36.5
#Diarrhea & enteritis (571)	178	30	148	178	30	148	8.4	2.6	15.4
Acute nephritis (590, 591)	81	30	51	81	31	50	3.8	2.7	5.2
Chronic nephritis, etc. (592-594)	320	124	196	329	130	199	15.6	11.3	20.7
Maternal deaths (640-689)	75	12	63	74	12	62	1.2	0.4	1.8
Cong. malformations (750-759)	231	132	99	247	142	105	11.7	12.4	10.9
Birth inj., atelect. (760-762)	392	220	172	387	218	169	6.2	8.0	4.8
#Other dis. early inf. (763-776)	798	248	550	804	254	550	12.9	9.3	15.8
Violent & acc. deaths (800-999)	1647	860	787	1630	843	787	77.2	73.5	81.8
Acc. deaths, total (800-962)	1326	719	607	1302	700	602	61.7	61.0	62.5
Motor vehicle acc. (810-835)	585	371	214	558	347	211	26.4	30.2	21.9
Ill-defined (780-795)	1729	367	1362	1705	355	1350	80.8	30.9	140.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

MISSISSIPPI STATE BOARD OF HEALTH
JACKSON, MISS.

MY 29 '58

ANNUAL BULLETIN
OF
VITAL STATISTICS
1956

VOLUME 2
COUNTY TABLES

MISSISSIPPI STATE BOARD OF HEALTH

ANNUAL BULLETIN

of

VITAL STATISTICS

1956

VOLUME 2

COUNTY TABLES

MISSISSIPPI STATE BOARD OF HEALTH

VOLUME 2

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 1 MONTH, AND DEATHS FROM CERTAIN SELECTED CAUSES, ON BASIS OF PLACE OF OCCURRENCE AND PLACE OF RESIDENCE, BY RACE, WITH RATES, FOR THE STATE OF MISSISSIPPI AND FOR EACH COUNTY.

COUNTY	PAGE
Adams	1
Alcorn	2
Amite	3
Attala	4
Benton	5
Bolivar	6
Calhoun	7
Carroll	8
Chickasaw	9
Choctaw	10
Claiborne	11
Clarke	12
Clay	13
Coahoma	14
Copiah	15
Covington	16
DeSoto	17
Forrest	18
Franklin	19
George	20
Greene	21
Grenada	22
Hancock	23
Harrison	24
Hinds	25
Holmes	26
Humphreys	27
Issaquena	28
Itawamba	29
Jackson	30
Jasper	31
Jefferson	32
Jeff Davis	33
Jones	34
Kemper	35
Lafayette	36
Lamar	37
Lauderdale	38
Lawrence	39
Leake	40
Lee	41
Leflore	42

COUNTY	PAGE
Lincoln	43
Lowndes	44
Madison	45
Marion	46
Marshall	47
Monroe	48
Montgomery	49
Neshoba	50
Newton	51
Noxubee	52
Oktibbeha	53
Panola	54
Pearl River	55
Perry	56
Pike	57
Pontotoc	58
Prentiss	59
Quitman	60
Rankin	61
Scott	62
Sharkey	63
Simpson	64
Smith	65
Stone	66
Sunflower	67
Tallahatchie	68
Tate	69
Tippah	70
Tishomingo	71
Tunica	72
Union	73
Walthall	74
Warren	75
Washington	76
Wayne	77
Webster	78
Wilkinson	79
Winston	80
Yalobusha	81
Yazoo	82
STATE	83

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ADAMS COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number		Non-white	Total	White	Non-white
				Total	White				
Live births	1,710	789	921	1,252	604	648	34.5	31.3	38.1
Fetal deaths	65	21	44	46	15	31	36.7	24.8	47.8
Deaths from all causes	454	173	281	382	137	245	10.5	7.1	14.4
Deaths under 1 year	75	26	49	58	25	33	46.3	41.4	50.9
Deaths under 1 month	54	21	33	39	20	19	31.2	33.1	29.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	2	1	4	3	1	11.0	15.5	5.9
Tbc., respiratory (001-008)	2	2	0	4	3	1	11.0	15.5	5.9
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	2.8	-	5.9
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	2.8	-	5.9
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (082)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	2.8	-	5.9
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	50	17	33	42	13	29	115.7	67.4	170.6
Diabetes (260)	4	2	2	2	0	2	5.5	-	11.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	57	17	40	51	11	40	140.5	57.0	235.3
Heart diseases (410-443)	156	66	90	130	52	78	358.1	269.4	458.8
Arteriosclerosis (450)	10	5	5	9	4	5	24.8	20.7	29.4
Influenza (480-483)	1	0	1	1	0	1	2.8	-	5.9
#Pneumonia (490-493)	7	3	4	5	3	2	13.8	15.5	11.8
#Diarrhea & enteritis (571)	5	0	5	5	0	5	13.8	-	29.4
Acute nephritis (590, 591)	0	0	0	1	1	0	2.8	5.2	-
Chronic nephritis, etc.(592-594)	3	1	2	1	0	1	2.8	-	5.9
Maternal deaths (640-689)	4	1	3	3	1	2	2.4	1.6	3.1
Cong. malformations (750-759)	12	8	4	11	8	3	30.3	41.4	17.6
Birth inj., atelect. (760-762)	19	8	11	9	6	3	7.2	9.9	4.6
#Other dis. early inf. (763-776)	25	8	17	22	9	13	17.6	14.9	20.1
Violent & acc. deaths (800-999)	32	17	15	27	9	18	74.4	46.6	105.9
Acc. deaths, total (800-962)	28	15	13	23	7	16	63.4	36.3	94.1
Motor vehicle acc. (810-835)	11	6	5	8	2	6	22.0	10.4	35.3
Ill-defined (780-795)	18	1	17	15	1	14	41.3	5.2	82.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ALCORN COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	659	546	113	565	454	111	22.5	21.1	30.8
Fetal deaths	14	10	4	14	10	4	24.8	22.0	36.0
Deaths from all causes	242	200	42	237	196	41	9.4	9.1	11.4
Deaths under 1 year	18	15	3	21	18	3	37.2	39.6	27.0
Deaths under 1 month	12	11	1	13	12	1	23.0	26.4	9.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	2	0	2	2	0	8.0	9.3	-
Tbc., respiratory (001-008)	2	2	0	2	2	0	8.0	9.3	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	33	28	5	28	23	5	111.6	107.0	138.9
Diabetes (260)	7	6	1	6	5	1	23.9	23.2	27.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	30	24	6	26	20	6	103.6	93.0	166.7
Heart diseases (410-443)	94	84	10	89	80	9	354.6	372.1	250.0
Arteriosclerosis (450)	6	5	1	7	6	1	27.9	27.9	27.8
Influenza (480-483)	2	0	2	2	0	2	8.0	-	55.6
#Pneumonia (490-493)	9	7	2	8	6	2	31.9	27.9	55.6
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	2	1	2	2	0	8.0	9.3	-
Maternal deaths (640-689)	1	0	1	1	0	1	1.8	-	9.0
Cong. malformations (750-759)	5	5	0	8	8	0	31.9	37.2	-
Birth inj., atelect. (760-762)	1	1	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	7	7	0	8	8	0	14.2	17.6	-
Violent & acc. deaths (800-999)	14	10	4	18	13	5	71.7	60.5	138.9
Acc. deaths, total (800-962)	13	9	4	15	11	4	59.8	51.2	111.1
Motor vehicle acc. (810-835)	4	3	1	5	5	0	19.9	23.2	-
Ill-defined (790-795)	8	4	4	7	4	3	27.9	18.6	83.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
 DEATHS FROM CERTAIN SELECTED CAUSES
 AMITE COUNTY, MISSISSIPPI, 1956

3

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	279	20	259	421	115	306	24.9	16.0	31.5
Fetal deaths	13	1	12	29	4	25	68.9	34.8	81.7
Deaths from all causes	98	45	53	147	75	72	8.7	10.4	7.4
Deaths under 1 year	11	1	10	16	2	14	38.0	17.4	45.8
Deaths under 1 month	5	0	5	9	1	8	21.4	8.7	26.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	2	0	2	11.8	-	20.6
Tbc., respiratory (001-008)	2	0	2	2	0	2	11.8	-	20.6
Syphilis (020-029)	0	0	0	1	1	0	5.9	13.9	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	5	5	18	11	7	106.5	152.8	72.2
Diabetes (260)	0	0	0	1	0	1	5.9	-	10.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	18	5	13	21	8	13	124.3	111.1	134.0
Heart diseases (410-443)	36	23	13	52	33	19	307.7	458.3	195.9
Arteriosclerosis (450)	4	4	0	4	4	0	23.7	55.6	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	0	2	4	1	3	23.7	13.9	30.9
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	0	1	5.9	-	10.3
Maternal deaths (640-689)	0	0	0	1	0	1	2.4	-	3.3
Cong. malformations (750-759)	1	0	1	1	0	1	5.9	-	10.3
Birth inj., atelect. (760-762)	1	0	1	1	0	1	2.4	-	3.3
#Other dis. early inf. (763-776)	3	1	2	7	2	5	16.6	17.4	16.3
Violent & acc. deaths (800-999)	5	2	3	11	8	3	65.1	111.1	30.9
Acc. deaths, total (800-962)	3	1	2	8	6	2	47.3	83.3	20.6
Motor vehicle acc. (810-835)	1	1	0	4	4	0	23.7	55.6	-
Un-defined (780-795)	16	5	11	16	5	11	94.7	69.4	113.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ATTALA COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	618	269	349	607	260	347	26.2	21.1	31.8
Fetal deaths	26	12	14	28	13	15	46.1	50.0	43.2
Deaths from all causes	198	115	83	206	122	84	8.9	9.9	7.7
Deaths under 1 year	27	10	17	26	10	16	42.8	38.5	46.1
Deaths under 1 month	16	7	9	15	7	8	24.7	26.9	23.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	2	1	1	8.6	8.1	9.2
Tbc., respiratory (001-008)	2	1	1	2	1	1	8.6	8.1	9.2
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	1	0	1	4.3	-	9.2
Whooping cough (056)	1	0	1	1	0	1	4.3	-	9.2
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia(140-205)	21	14	7	24	16	8	103.4	130.1	73.4
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	31	17	14	36	21	15	155.2	170.7	137.6
Heart diseases (410-443)	77	48	29	81	51	30	414.6	658.5	275.2
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	1	0	1	1	0	1	4.3	-	9.2
#Pneumonia (490-493)	7	2	5	7	2	5	30.2	16.3	45.9
#Diarrhea & enteritis (571)	2	1	1	2	1	1	8.6	8.1	9.2
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis,etc.(592-594)	2	2	0	3	3	0	12.9	24.4	-
Maternal deaths (640-689)	1	0	1	0	0	0	-	-	-
Cong. malformations (750-759)	6	4	2	5	4	1	21.6	32.5	9.2
Birth inj., atelect. (760-762)	2	2	0	1	1	0	1.6	3.8	-
#Other dis. early inf.(763-776)	12	3	9	12	4	8	19.8	15.4	23.0
Violent & acc. deaths (800-999)	11	7	4	7	4	3	30.2	32.5	27.5
Acc. deaths, total (800-962)	10	7	3	6	4	2	25.9	32.5	18.3
Motor vehicle acc. (810-835)	8	6	2	3	2	1	12.9	16.3	9.2
Un-defined (780-795)	7	3	4	7	3	4	30.2	24.4	36.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
BENTON COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	178	18	160	251	76	175	29.5	17.7	41.7
Fetal deaths	5	1	4	11	4	7	43.8	52.6	40.0
Deaths from all causes	38	16	22	56	30	26	6.6	7.0	6.2
Deaths under 1 year	3	0	3	8	3	5	31.9	39.5	28.6
Deaths under 1 month	2	0	2	5	2	3	19.9	26.3	17.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia(140-205)	8	3	5	11	6	5	129.4	139.5	119.0
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	4	1	3	7	2	5	82.4	46.5	119.0
Heart diseases (410-443)	10	6	4	12	8	4	141.2	186.0	95.2
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	0	1	2	1	1	23.5	23.2	23.8
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc.(592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	1	0	11.8	23.2	-
Birth inj., atelect. (760-762)	2	0	2	3	1	2	12.0	13.2	11.4
#Other dis. early inf.(763-776)	0	0	0	3	1	2	12.0	13.2	11.4
Violent & acc. deaths (800-999)	2	1	1	3	2	1	35.3	46.5	23.8
Acc. deaths, total (800-962)	2	1	1	3	2	1	35.3	46.5	23.8
Motor vehicle acc. (810-835)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	7	2	5	7	2	5	82.4	46.5	119.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
BOLIVAR COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	2,204	435	1,769	2,089	463	1,626	35.3	26.3	39.2
Fetal deaths	93	15	78	91	18	73	43.6	38.9	44.9
Deaths from all causes	612	102	510	634	127	507	10.7	7.2	12.2
Deaths under 1 year	74	8	66	78	10	68	37.3	21.6	41.8
Deaths under 1 month	45	8	37	47	8	39	22.5	17.3	24.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	1	2	7	3	4	11.8	17.0	9.6
Tbc., respiratory (001-008)	2	0	2	5	2	3	8.5	11.4	7.2
Syphilis (020-029)	4	1	3	5	1	4	8.5	5.7	9.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	0	2	4	0	4	6.8	-	9.6
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	4	0	4	3	0	3	5.1	-	7.2
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	76	19	57	90	30	60	152.3	170.4	144.6
Diabetes (260)	6	1	5	6	1	5	10.2	5.7	12.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	76	3	73	76	10	66	128.6	56.8	159.0
Heart diseases (410-443)	177	42	135	176	47	129	297.8	267.0	310.8
Arteriosclerosis (450)	8	0	8	7	0	7	11.8	-	16.9
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	9	1	8	12	3	9	20.3	17.0	21.7
#Diarrhea & enteritis (571)	10	0	10	9	0	9	15.2	-	21.7
Acute nephritis (590, 591)	5	1	4	5	1	4	8.5	5.7	9.6
Chronic nephritis, etc. (592-594)	9	2	7	9	3	6	15.2	17.0	14.4
Maternal deaths (640-689)	2	0	2	2	0	2	1.0	-	1.2
Cong. malformations (750-759)	2	1	1	2	1	1	3.4	5.7	2.4
Birth inj., atelect. (760-762)	16	3	13	17	3	14	8.1	6.5	8.6
#Other dis. early inf. (763-776)	22	4	18	23	4	19	11.0	8.6	11.7
Violent & acc. deaths (800-999)	50	17	33	45	12	33	76.1	68.2	79.5
Acc. deaths, total (800-962)	41	17	24	35	12	23	59.2	68.2	55.4
Motor vehicle acc. (810-835)	10	6	4	7	4	3	11.8	22.7	7.2
Undefined (780-795)	86	2	84	88	3	85	148.9	17.0	204.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CALHOUN COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	433	234	199	400	223	177	26.5	21.2	38.5
Fetal deaths	19	7	12	15	5	10	37.5	22.4	56.5
Deaths from all causes	138	99	39	168	130	38	11.1	12.4	8.3
Deaths under 1 year	14	8	6	17	11	6	42.5	49.3	33.9
Deaths under 1 month	12	8	4	13	9	4	32.5	40.4	22.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	1	0	6.6	9.5	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	24	20	4	25	21	4	165.6	200.0	87.0
Diabetes (260)	1	1	0	1	1	0	6.6	9.5	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	18	14	4	22	18	4	145.7	171.4	87.0
Heart diseases (410-443)	39	35	4	50	45	5	331.1	428.6	108.7
Arteriosclerosis (450)	2	0	2	2	0	2	13.2	-	43.5
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	13	10	3	14	11	3	92.7	104.8	65.2
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	2	0	3	2	1	19.9	19.0	21.7
Maternal deaths (640-689)	1	0	1	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	2	1	1	13.2	9.5	21.7
Birth inj., atelect. (760-762)	1	0	1	1	0	1	2.5	-	5.6
#Other dis. early inf. (763-776)	11	8	3	12	9	3	30.0	40.4	16.9
Violent & acc. deaths (800-999)	7	3	4	11	8	3	72.8	76.2	65.2
Acc. deaths, total (800-962)	5	3	2	8	7	1	53.0	66.7	21.7
Motor vehicle acc. (810-835)	2	2	0	4	4	0	26.5	38.1	-
Ill-defined (780-795)	6	0	6	7	1	6	46.4	9.5	130.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CARROLL COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	279	36	243	347	84	263	26.9	17.1	32.9
Fetal deaths	12	0	12	16	3	13	46.1	35.7	49.4
Deaths from all causes	72	27	45	120	59	61	9.3	12.0	7.6
Deaths under 1 year	3	0	3	11	4	7	31.7	47.6	26.6
Deaths under 1 month	1	0	1	7	3	4	20.2	35.7	15.2
DEATHS FROM SELECTED CAUSES AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	7.8	-	12.5
Tbc., respiratory (001-008)	1	0	1	1	0	1	7.8	-	12.5
Syphilis (020-029)	1	1	0	1	1	0	7.8	20.4	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	4	2	2	9	4	5	69.8	81.6	62.5
Diabetes (260)	1	0	1	1	0	1	7.8	-	12.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	9	2	7	13	5	8	100.8	102.0	100.0
Heart diseases (410-443)	20	9	11	36	22	14	279.1	449.0	175.0
Arteriosclerosis (450)	0	0	0	1	1	0	7.8	20.4	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	3	0	3	7	2	5	54.3	40.8	62.5
#Diarrhea & enteritis (571)	0	0	0	1	0	1	7.8	-	12.5
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	7.8	-	12.5
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	1	1	0	7.8	20.4	-
Birth inj., atelect. (760-762)	0	0	0	3	2	1	8.6	23.8	3.8
#Other dis. early inf. (763-776)	1	0	1	3	1	2	8.6	11.9	7.6
Violent & acc. deaths (800-999)	10	7	3	13	8	5	100.8	163.3	62.5
Acc. deaths, total (800-962)	6	3	3	10	5	5	77.5	102.0	62.5
Motor vehicle acc. (810-835)	4	3	1	7	5	2	54.3	102.0	25.0
Ill-defined (780-795)	14	3	11	13	3	10	100.8	61.2	125.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CHICKASAW COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	447	218	229	453	223	230	26.3	22.8	31.1
Fetal deaths	15	3	12	16	6	10	35.3	26.9	43.5
Deaths from all causes	157	99	58	172	108	64	10.0	11.0	8.6
Deaths under 1 year	19	6	13	17	6	11	37.5	26.9	47.8
Deaths under 1 month	9	6	3	9	6	3	19.9	26.9	13.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	4	1	3	23.2	10.2	40.5
Tbc., respiratory (001-008)	1	0	1	4	1	3	23.2	10.2	40.5
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	19	15	4	24	18	6	139.5	183.7	81.1
Diabetes (260)	3	2	1	2	2	0	11.6	20.4	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	7	8	16	8	8	93.0	81.6	108.1
Heart diseases (410-443)	58	44	14	66	49	17	383.7	500.0	229.7
Arteriosclerosis (450)	2	2	0	2	2	0	11.6	20.4	-
Influenza (480-483)	1	0	1	1	0	1	5.8	-	13.5
#Pneumonia (490-493)	2	1	1	3	2	1	17.4	20.4	13.5
#Diarrhea & enteritis (571)	6	0	6	4	0	4	23.2	-	54.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	11.6	10.2	13.5
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	4	4	0	3	3	0	6.6	13.4	-
#Other dis. early inf. (763-776)	1	0	1	2	1	1	4.4	4.5	4.3
Violent & acc. deaths (800-999)	16	12	4	14	11	3	81.4	112.2	40.5
Acc. deaths, total (800-962)	14	11	3	12	10	2	69.8	102.0	27.0
Motor vehicle acc. (810-835)	5	3	2	4	3	1	23.2	30.6	13.5
Ill-defined (780-795)	12	3	9	12	3	9	69.8	30.6	121.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CHOCTAW COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	210	104	106	213	117	96	22.6	18.9	30.0
Fetal deaths	4	4	0	5	4	1	23.5	34.2	10.4
Deaths from all causes	73	58	15	83	61	22	8.8	9.8	6.9
Deaths under 1 year	2	2	0	7	5	2	32.9	42.7	20.8
Deaths under 1 month	1	1	0	4	4	0	18.8	34.2	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	12	11	1	12	10	2	127.6	161.3	62.5
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	8	6	2	10	7	3	106.4	112.9	93.8
Heart diseases (410-443)	30	24	6	34	28	6	361.7	451.6	187.5
Arteriosclerosis (450)	4	3	1	3	2	1	31.9	32.2	31.2
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	0	0	0	0	0	0	-	-	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	2	0	2	21.3	-	62.5
#Other dis. early inf. (763-776)	1	1	0	2	2	0	-	-	-
Violent & acc. deaths (800-999)	5	4	1	7	4	3	74.5	64.5	93.8
Acc. deaths, total (800-962)	4	3	1	6	4	2	63.8	64.5	62.5
Motor vehicle acc. (810-835)	1	1	0	2	2	0	21.3	32.2	-
Ill-defined (780-795)	4	2	2	4	2	2	42.6	32.2	62.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CLAIBORNE COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	290	41	249	329	61	268	29.4	19.7	33.1
Fetal deaths	13	1	12	17	1	16	51.7	16.4	59.7
Deaths from all causes	115	19	96	147	38	109	13.1	12.2	13.4
Deaths under 1 year	12	0	12	15	1	14	45.6	16.4	52.2
Deaths under 1 month	7	0	7	8	0	8	24.3	-	29.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	8.9	-	12.3
Tbc., respiratory (001-008)	1	0	1	1	0	1	8.9	-	12.3
Syphilis (020-029)	1	0	1	1	0	1	8.9	-	12.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	2	8	15	4	11	133.9	129.0	135.8
Diabetes (260)	5	0	5	6	0	6	53.6	-	74.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	16	2	14	20	7	13	178.6	225.8	160.5
Heart diseases (410-443)	39	10	29	50	15	35	446.4	483.9	432.1
Arteriosclerosis (450)	2	1	1	1	0	1	8.9	-	12.3
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	5	0	5	5	0	5	44.6	-	61.7
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	8.9	32.2	-
Maternal deaths (640-689)	0	0	0	1	0	1	3.0	-	3.7
Cong. malformations (750-759)	0	0	0	1	0	1	17.8	32.2	12.3
Birth inj., atelect. (760-762)	1	0	1	2	1	1	9.1	-	11.2
#Other dis. early inf. (763-776)	4	0	4	4	0	4	12.2	-	14.9
Violent & acc. deaths (800-999)	9	1	8	13	4	9	116.1	129.0	111.1
Acc. deaths, total (800-962)	8	1	7	11	4	7	98.2	129.0	86.4
Motor vehicle acc. (810-835)	4	0	4	5	1	4	44.6	32.2	49.4
Ill-defined (780-795)	16	1	15	14	1	13	125.0	32.2	160.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CLARKE COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	290	142	148	442	198	244	25.5	19.6	33.9
Fetal deaths	10	2	8	17	3	14	38.5	15.2	57.4
Deaths from all causes	152	88	64	195	110	85	11.3	10.9	11.8
Deaths under 1 year	11	5	6	14	5	9	31.7	25.2	36.9
Deaths under 1 month	5	4	1	6	4	2	13.6	20.2	8.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	2	1	1	11.6	9.9	13.9
Tbc., respiratory (001-008)	1	0	1	2	1	1	11.6	9.9	13.9
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	21	10	11	21	10	11	121.4	99.0	152.8
Diabetes (260)	3	2	1	3	2	1	17.3	19.8	13.9
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	23	12	11	36	20	16	208.1	198.0	222.2
Heart diseases (410-443)	61	43	18	72	49	23	416.2	485.1	319.4
Arteriosclerosis (450)	1	1	0	1	1	0	5.8	9.9	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	5	1	4	7	1	6	40.5	9.9	83.3
#Diarrhea & enteritis (571)	1	0	1	1	0	1	5.8	-	13.9
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	6	2	4	6	2	4	34.7	19.8	55.6
Maternal deaths (640-689)	1	0	1	2	0	2	4.5	-	8.2
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	3	3	0	3	3	0	6.8	15.2	-
#Other dis. early inf. (763-776)	4	2	2	4	2	2	9.0	10.1	8.2
Violent & acc. deaths (800-999)	12	6	6	16	8	8	92.5	79.2	111.1
Acc. deaths, total (800-962)	9	5	4	13	7	6	75.1	69.3	83.3
Motor vehicle acc. (810-835)	6	4	2	10	6	4	57.8	59.4	55.6
Ill-defined (780-795)	7	4	3	8	4	4	46.2	39.6	55.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CLAY COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	548	180	368	561	197	364	37.2	25.9	48.5
Fetal deaths	18	4	14	20	5	15	35.6	25.4	41.2
Deaths from all causes	177	75	102	180	75	105	11.9	9.9	14.0
Deaths under 1 year	19	4	15	22	5	17	39.2	25.4	46.7
Deaths under 1 month	9	2	7	10	3	7	17.8	15.2	19.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	6.6	-	13.3
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	1	0	1	1	0	1	6.6	-	13.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	0	1	1	0	1	6.6	-	13.3
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	25	16	9	25	16	9	165.6	210.5	120.0
Diabetes (260)	3	2	1	3	2	1	19.9	26.3	13.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	17	10	7	18	10	8	119.2	131.6	106.7
Heart diseases (410-443)	65	28	37	64	27	37	423.8	355.3	493.3
Arteriosclerosis (450)	5	3	2	4	2	2	26.5	26.3	26.7
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	4	2	2	4	2	2	26.5	26.3	26.7
#Diarrhea & enteritis (571)	2	0	2	4	0	4	26.5	-	53.3
Acute nephritis (590, 591)	1	0	1	1	0	1	6.6	-	13.3
Chronic nephritis, etc. (592-594)	3	0	3	3	0	3	19.9	-	40.0
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	4	3	1	4	3	1	26.5	39.5	13.3
Birth inj., atelect. (760-762)	1	0	1	1	0	1	1.8	-	2.7
#Other dis. early inf. (763-776)	10	2	8	11	3	8	19.6	15.2	22.0
Violent & acc. deaths (800-999)	8	4	4	7	5	2	46.4	65.8	26.7
Acc. deaths, total (800-962)	8	4	4	7	5	2	46.4	65.8	26.7
Motor vehicle acc. (810-835)	2	1	1	3	2	1	19.9	26.3	13.3
Ill-defined (780-795)	15	0	15	16	0	16	106.0	-	213.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
COAHOMA COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,630	377	1,253	1,664	342	1,322	31.6	23.8	34.6
Fetal deaths	83	23	60	93	23	70	55.9	67.2	53.0
Deaths from all causes	500	112	388	503	101	402	9.6	7.0	10.5
Deaths under 1 year	85	8	77	81	5	76	48.7	14.6	57.5
Deaths under 1 month	47	5	42	44	3	41	26.4	8.8	31.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	0	3	5	2	3	9.5	13.9	7.8
Tbc., respiratory (001-008)	2	0	2	4	2	2	7.6	13.9	5.2
Syphilis (020-029)	1	0	1	1	0	1	1.9	-	2.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	3	0	3	3	0	3	5.7	-	7.8
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	1	0	1	1	0	1	1.9	-	2.6
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	50	19	31	50	17	33	95.0	118.0	86.4
Diabetes (260)	5	1	4	5	1	4	9.5	6.9	10.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	60	13	47	64	13	51	121.7	90.3	133.5
Heart diseases (410-443)	131	36	95	134	35	99	254.8	243.0	259.2
Arteriosclerosis (450)	6	5	1	6	5	1	11.4	34.7	2.6
Influenza (480-483)	8	0	8	8	0	8	15.2	-	20.9
#Pneumonia (490-493)	22	2	20	20	1	19	38.0	6.9	49.7
#Diarrhea & enteritis (571)	10	0	10	10	0	10	19.0	-	26.2
Acute nephritis (590, 591)	3	1	2	2	0	2	3.8	-	5.2
Chronic nephritis, etc. (592-594)	12	1	11	13	1	12	24.7	6.9	31.4
Maternal deaths (640-689)	2	0	2	4	0	4	2.4	-	3.0
Cong. malformations (750-759)	4	1	3	5	1	4	9.5	6.9	10.5
Birth inj., atelect. (760-762)	7	0	7	7	0	7	4.2	-	5.3
#Other dis. early inf. (763-776)	33	5	28	30	3	27	18.0	8.8	20.4
Violent & acc. deaths (800-999)	40	19	21	36	14	22	68.4	97.2	57.6
Acc. deaths, total (800-962)	32	17	15	27	12	15	51.3	83.3	39.3
Motor vehicle acc. (810-835)	14	10	4	8	4	4	15.2	27.8	10.5
Ill-defined (780-795)	63	0	63	65	0	65	123.6	-	170.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
COPIAH COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	954	279	675	775	222	553	28.3	17.9	36.9
Fetal deaths	18	4	14	17	6	11	21.9	27.0	19.9
Deaths from all causes	263	115	148	302	133	169	11.0	10.7	11.3
Deaths under 1 year	27	4	23	24	3	21	31.0	13.5	38.0
Deaths under 1 month	11	1	10	10	2	8	12.9	9.0	14.5
DEATHS FROM SELECTED CAUSES AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	3.6	-	6.7
Tbc., respiratory (001-008)	1	0	1	1	0	1	3.6	-	6.7
Syphilis (020-029)	1	0	1	1	0	1	3.6	-	6.7
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	3.6	-	6.7
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	0	0	0	-	-	-
Meningococcal infections (057)	1	1	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	27	18	9	34	23	11	124.1	185.5	73.3
Diabetes (260)	2	1	1	1	0	1	3.6	-	6.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	27	10	17	36	12	24	131.4	96.8	160.0
Heart diseases (410-443)	109	56	53	121	66	55	441.6	532.2	366.7
Arteriosclerosis (450)	3	2	1	3	2	1	10.9	16.1	6.7
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	7	1	6	9	2	7	32.8	16.1	46.7
#Diarrhea & enteritis (571)	1	1	0	2	0	2	7.3	-	13.3
Acute nephritis (590, 591)	1	0	1	1	0	1	3.6	-	6.7
Chronic nephritis, etc. (592-594)	5	1	4	6	2	4	21.9	16.1	26.7
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	2	0	2	7.3	-	13.3
Birth inj., atelect. (760-762)	1	0	1	1	0	1	1.3	-	1.8
#Other dis. early inf. (763-776)	12	1	11	10	2	8	12.9	9.0	14.5
Violent & acc. deaths (800-999)	24	9	15	28	8	20	102.2	64.5	133.3
Acc. deaths, total (800-962)	19	9	10	22	8	14	80.3	64.5	93.3
Motor vehicle acc. (810-835)	8	4	4	6	0	6	21.9	-	40.0
Ill-defined (780-795)	21	5	16	21	4	17	76.6	32.2	113.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
COVINGTON COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	249	150	99	397	205	192	25.9	21.6	33.1
Fetal deaths	8	4	4	15	10	5	37.8	48.8	26.0
Deaths from all causes	93	67	26	127	86	41	8.3	9.0	7.1
Deaths under 1 year	7	5	2	10	7	3	25.2	34.1	15.6
Deaths under 1 month	4	3	1	5	3	2	12.6	14.6	10.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	2	0	5	4	1	32.7	42.1	17.2
Tbc., respiratory (001-008)	2	2	0	4	3	1	26.1	31.6	17.2
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	7	3	14	9	5	91.5	94.7	86.2
Diabetes (260)	2	1	1	3	1	2	19.6	10.5	34.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	19	14	5	22	17	5	143.8	178.9	86.2
Heart diseases (410-443)	30	23	7	40	30	10	261.4	315.8	172.4
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	1	1	3	2	1	19.6	21.0	17.2
#Diarrhea & enteritis (571)	1	1	0	1	1	0	6.5	10.5	-
Acute nephritis (590, 591)	0	0	0	1	0	1	6.5	-	17.2
Chronic nephritis, etc. (592-594)	2	0	2	4	2	2	26.1	21.0	34.5
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	2	1	1	13.1	10.5	17.2
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	4	3	1	5	4	1	12.6	19.5	5.2
Violent & acc. deaths (800-999)	8	7	1	8	5	3	52.3	52.6	51.7
Acc. deaths, total (800-962)	7	6	1	7	4	3	45.8	42.1	51.7
Motor vehicle acc. (810-835)	4	4	0	2	1	1	13.1	10.5	17.2
Ill-defined (780-795)	7	3	4	8	3	5	52.3	31.6	86.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
DESOTO COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	657	97	560	720	177	543	29.9	21.1	34.6
Fetal deaths	13	1	12	18	3	15	25.0	16.9	27.6
Deaths from all causes	173	52	121	212	64	148	8.8	7.6	9.4
Deaths under 1 year	16	1	15	27	5	22	37.5	28.2	40.5
Deaths under 1 month	6	0	6	15	4	11	20.8	22.6	20.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	11	5	6	23	14	9	95.4	166.7	57.3
Diabetes (260)	1	0	1	1	0	1	4.1	-	6.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	19	2	17	26	4	22	107.9	47.6	140.1
Heart diseases (410-443)	45	19	26	53	20	33	219.9	238.1	210.2
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	13	4	9	13	4	9	53.9	47.6	57.3
#Diarrhea & enteritis (571)	1	0	1	1	0	1	4.1	-	6.4
Acute nephritis (590, 591)	2	0	2	2	0	2	8.3	-	12.7
Chronic nephritis, etc. (592-594)	1	0	1	4	1	3	16.6	11.9	19.1
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	5	1	4	20.7	11.9	25.5
Birth inj., atelect. (760-762)	1	0	1	2	1	1	2.8	5.6	1.8
#Other dis. early inf. (763-776)	4	0	4	8	2	6	11.1	11.3	11.0
Violent & acc. deaths (800-999)	32	15	17	25	7	18	103.7	83.3	114.6
Acc. deaths, total (800-962)	24	11	13	16	3	13	66.4	35.7	82.8
Motor vehicle acc. (810-835)	7	1	6	5	2	3	20.7	23.8	19.1
Ill-defined (780-795)	29	3	26	31	4	27	128.6	47.6	172.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
FORREST COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,577	1,149	428	1,329	860	469	29.9	26.6	38.4
Fetal deaths	72	46	26	68	36	32	51.2	41.9	68.2
Deaths from all causes	429	304	125	422	280	142	9.5	8.7	11.6
Deaths under 1 year	40	23	17	34	15	19	25.6	17.4	40.5
Deaths under 1 month	28	18	10	25	14	11	18.8	16.3	23.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	6	5	1	12	9	3	27.0	27.9	24.6
Tbc., respiratory (001-008)	5	4	1	12	9	3	27.0	27.9	24.6
Syphilis (020-029)	2	1	1	2	1	1	4.5	3.1	8.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	2	1	1	4.5	3.1	8.2
Scarlet fever & streptococcal sore throat (050, 051)	1	1	0	1	1	0	2.2	3.1	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	1	1	0	2	1	1	4.5	3.1	8.2
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	49	37	12	46	33	13	103.4	102.2	106.6
Diabetes (260)	4	4	0	6	4	2	13.5	12.4	16.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	55	34	21	59	35	24	132.6	108.4	196.7
Heart diseases (410-443)	139	113	26	127	100	27	285.4	309.6	221.3
Arteriosclerosis (450)	9	8	1	7	6	1	15.7	18.6	8.2
Influenza (480-483)	6	1	5	6	1	5	13.5	3.1	41.0
#Pneumonia (490-493)	17	13	4	15	11	4	33.7	34.0	32.8
#Diarrhea and enteritis (571)	1	0	1	0	0	0	-	-	-
Acute nephritis (590, 591)	1	1	0	2	1	1	4.5	3.1	8.2
Chronic nephritis, etc. (592-594)	14	5	9	14	5	9	31.5	15.5	73.8
Maternal deaths (640-689)	2	0	2	2	0	2	1.5	-	4.3
Cong. malformations (750-759)	7	5	2	7	4	3	15.7	12.4	24.6
Birth inj., atelect. (760-762)	6	5	1	5	3	2	3.8	3.5	4.3
#Other dis. early inf. (763-776)	17	10	7	16	9	7	12.0	10.5	14.9
Violent & acc. deaths (800-999)	31	19	12	39	23	16	87.6	71.2	131.1
Acc. deaths, total (800-962)	27	17	10	34	21	13	76.4	65.0	106.6
Motor vehicle acc. (810-835)	12	8	4	18	12	6	40.4	37.2	49.2
Ill-defined (780-795)	18	6	12	16	4	12	36.0	12.4	98.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
FRANKLIN COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	157	72	85	249	124	125	28.3	23.4	35.7
Fetal deaths	3	0	3	7	0	7	28.1	-	56.0
Deaths from all causes	72	47	25	99	65	34	11.2	12.3	9.7
Deaths under 1 year	4	3	1	6	5	1	24.1	40.3	8.0
Deaths under 1 month	3	3	0	4	4	0	16.1	32.2	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	11.4	18.9	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	11.4	18.9	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	7	3	15	9	6	170.4	169.8	171.4
Diabetes (260)	0	0	0	1	1	0	11.4	18.9	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	13	4	9	18	8	10	204.5	150.9	285.7
Heart diseases (410-443)	18	15	3	26	20	6	295.4	377.4	171.4
Arteriosclerosis (450)	1	1	0	1	1	0	11.4	18.9	-
Influenza (480-483)	2	0	2	2	0	2	22.7	-	57.1
#Pneumonia (490-493)	2	1	1	2	0	2	22.7	-	57.1
#Diarrhea & enteritis (571)	2	1	1	2	1	1	22.7	18.9	28.6
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	3	0	3	3	0	34.1	56.6	-
Maternal deaths (640-689)	1	0	1	1	0	1	4.0	-	8.0
Cong. malformations (750-759)	2	1	1	2	1	1	22.7	18.9	28.6
Birth inj., atelect. (760-762)	3	3	0	4	4	0	16.1	32.2	-
#Other dis. early inf. (763-776)	0	0	0	0	0	0	-	-	-
Violent & acc. deaths (800-999)	9	7	2	9	7	2	102.3	132.1	57.1
Acc. deaths, total (800-962)	7	6	1	7	5	2	79.5	94.3	57.1
Motor vehicle acc. (810-835)	3	2	1	3	2	1	34.1	37.7	28.6
Ill-defined (780-795)	2	1	1	3	1	2	34.1	18.9	57.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
GEORGE COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	309	256	53	324	270	54	30.8	29.3	41.5
Fetal deaths	4	4	0	3	3	0	9.2	11.1	-
Deaths from all causes	79	69	10	91	78	13	8.7	8.5	10.0
Deaths under 1 year	7	4	3	9	6	3	27.8	22.2	55.6
Deaths under 1 month	6	3	3	7	4	3	21.6	14.8	55.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	9.5	-	76.9
Tbc., respiratory (001-008)	0	0	0	1	0	1	9.5	-	76.9
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	1	0	9.5	10.9	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	1	1	0	9.5	10.9	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	15	13	2	15	13	2	142.8	141.3	153.8
Diabetes (260)	4	4	0	4	4	0	38.1	43.5	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	6	6	0	8	8	0	76.2	87.0	-
Heart diseases (410-443)	27	25	2	31	28	3	295.2	304.3	230.8
Arteriosclerosis (450)	4	3	1	4	3	1	38.1	32.6	76.9
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	1	0	1	1	0	9.5	10.9	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	2	0	2	2	0	19.0	21.7	-
Birth inj., atelect. (760-762)	3	2	1	3	2	1	9.2	7.4	18.5
#Other dis. early inf. (763-776)	3	1	2	3	1	2	9.2	3.7	37.0
Violent & acc. deaths (800-999)	6	5	1	6	4	2	57.1	43.5	153.8
Acc. deaths, total (800-962)	5	5	0	5	4	1	47.6	43.5	76.9
Motor vehicle acc. (810-835)	5	5	0	5	4	1	47.6	43.5	76.9
Ill-defined (780-795)	2	2	0	2	2	0	19.0	21.7	-

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
GREENE COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	144	77	67	249	174	75	31.1	27.2	46.9
Fetal deaths	2	1	1	5	3	2	20.1	17.2	26.7
Deaths from all causes	59	41	18	72	51	21	9.0	8.0	13.1
Deaths under 1 year	3	1	2	5	2	3	20.1	11.5	40.0
Deaths under 1 month	3	1	2	4	2	2	16.1	11.5	26.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	12.5	15.6	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	12.5	15.6	-
Syphilis (020-029)	1	1	0	1	1	0	12.5	15.6	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	7	5	2	11	9	2	137.5	140.6	125.0
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	8	3	5	9	4	5	112.5	62.5	312.5
Heart diseases (410-443)	21	18	3	24	20	4	300.0	312.5	250.0
Arteriosclerosis (450)	0	0	0	1	1	0	12.5	15.6	-
Influenza (480-483)	2	2	0	2	2	0	25.0	31.2	-
#Pneumonia (490-493)	2	1	1	4	3	1	50.0	46.9	62.5
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	1	1	3	1	2	37.5	15.6	125.0
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	2	0	2	2	0	2	8.0	-	26.7
Violent & acc. deaths (800-999)	6	5	1	6	5	1	75.0	78.1	62.5
Acc. deaths, total (800-962)	6	5	1	6	5	1	75.0	78.1	62.5
Motor vehicle acc. (810-835)	4	3	1	4	3	1	50.0	46.9	62.5
Ill-defined (780-795)	6	3	3	6	3	3	75.0	46.9	187.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
GRENADA COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	583	244	339	532	194	338	28.4	22.3	33.8
Fetal deaths	35	12	23	34	10	24	63.9	51.5	71.0
Deaths from all causes	201	95	106	155	57	98	8.3	6.6	9.8
Deaths under 1 year	26	9	17	21	6	15	39.5	30.9	44.4
Deaths under 1 month	17	5	12	13	3	10	24.4	15.5	29.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	5.3	-	10.0
Tbc., respiratory (001-008)	1	0	1	1	0	1	5.3	-	10.0
Syphilis (020-029)	0	0	0	1	1	0	5.3	11.5	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	16	7	9	17	7	10	90.9	80.4	100.0
Diabetes (260)	2	1	1	2	1	1	10.7	11.5	10.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	40	11	29	32	7	25	171.1	80.4	250.0
Heart diseases (410-443)	42	27	15	28	14	14	149.7	160.9	140.0
Arteriosclerosis (450)	11	7	4	11	7	4	58.8	80.4	40.0
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	6	1	5	5	0	5	26.7	-	50.0
#Diarrhea & enteritis (571)	2	1	1	2	1	1	10.7	11.5	10.0
Acute nephritis (590, 591)	2	2	0	1	1	0	5.3	11.5	-
Chronic nephritis, etc. (592-594)	2	1	1	1	0	1	5.3	-	10.0
Maternal deaths (640-689)	1	0	1	2	0	2	3.8	-	5.9
Cong. malformations (750-759)	2	2	0	1	1	0	5.3	11.5	-
Birth inj., atelect. (760-762)	1	1	0	1	1	0	1.9	5.2	-
#Other dis. early inf. (763-776)	15	4	11	12	3	9	22.6	15.5	26.6
Violent & acc. deaths (800-999)	25	16	9	10	3	7	53.5	34.5	70.0
Acc. deaths, total (800-962)	25	16	9	10	3	7	53.5	34.5	70.0
Motor vehicle acc. (810-835)	13	7	6	5	1	4	26.7	11.5	40.0
<u>Ill-defined (780-795)</u>	14	4	10	15	5	10	80.2	57.5	100.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HANCOCK COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	175	145	30	269	233	36	20.5	19.7	27.7
Fetal deaths	5	5	0	10	7	3	37.2	30.0	83.3
Deaths from all causes	74	61	13	99	86	13	7.6	7.3	10.0
Deaths under 1 year	6	5	1	10	9	1	37.2	38.6	27.8
Deaths under 1 month	4	4	0	8	8	0	29.7	34.3	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	2	1	1	15.3	8.5	76.9
Tbc., respiratory (001-008)	0	0	0	1	1	0	7.6	8.5	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	6	4	2	13	11	2	99.2	93.2	153.8
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	11	9	2	16	14	2	122.1	118.6	153.8
Heart diseases (410-443)	21	20	1	25	24	1	190.8	203.4	76.9
Arteriosclerosis (450)	1	1	0	2	2	0	15.3	16.9	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	2	0	4	3	1	30.5	25.4	76.9
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	7.6	8.5	-
Maternal deaths (640-689)	1	1	0	1	1	0	3.7	4.3	-
Cong. malformations (750-759)	2	2	0	3	3	0	22.9	25.4	-
Birth inj., atelect. (760-762)	1	1	0	3	3	0	11.2	12.9	-
#Other dis. early inf. (763-776)	2	2	0	3	3	0	11.2	12.9	-
Violent & acc. deaths (800-999)	8	5	3	3	2	1	22.9	16.9	76.9
Acc. deaths, total (800-962)	8	5	3	3	2	1	22.9	16.9	76.9
Motor vehicle acc. (810-835)	8	5	3	2	2	0	15.3	16.9	-
Ill-defined (780-795)	17	13	4	14	10	4	106.9	84.7	307.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HARRISON COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	3,645	2,988	657	3,436	2,803	633	32.8	32.0	37.0
Fetal deaths	87	61	26	77	56	21	22.4	20.0	33.2
Deaths from all causes	846	666	180	746	573	173	7.1	6.5	10.1
Deaths under 1 year	109	81	28	102	74	28	29.7	26.4	44.2
Deaths under 1 month	82	60	22	76	54	22	22.1	19.3	34.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	7	1	6	4	0	4	3.8	-	23.4
Tbc., respiratory (001-008)	6	1	5	3	0	3	2.9	-	17.5
Syphilis (020-029)	4	1	3	3	1	2	2.9	1.1	11.7
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	1	1	2	1	1	1.9	1.1	5.8
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	97	77	20	82	59	23	78.3	67.4	134.5
Diabetes (260)	7	5	2	8	6	2	7.6	6.8	11.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	57	47	10	52	42	10	49.7	47.9	58.5
Heart diseases (410-443)	324	274	50	286	239	47	273.2	272.8	274.8
Arteriosclerosis (450)	6	6	0	6	5	1	5.7	5.7	5.8
Influenza (480-483)	1	1	0	1	1	0	1.0	1.1	-
#Pneumonia (490-493)	23	23	0	16	15	1	15.3	17.1	5.8
#Diarrhea & enteritis (571)	4	3	1	4	3	1	3.8	3.4	5.8
Acute nephritis (590, 591)	2	1	1	2	1	1	1.9	1.1	5.8
Chronic nephritis, etc. (592-594)	2	2	0	2	1	1	1.9	1.1	5.8
Maternal deaths (640-689)	2	2	0	2	2	0	0.6	0.7	-
Cong. malformations (750-759)	13	13	0	12	12	0	11.5	13.7	-
Birth inj., atelect. (760-762)	28	21	7	23	16	7	6.7	5.7	11.0
#Other dis. early inf. (763-776)	35	24	11	34	23	11	9.9	8.2	17.4
Violent & acc. deaths (800-999)	70	53	17	67	52	15	64.0	59.4	87.7
Acc. deaths, total (800-962)	57	43	14	53	41	12	50.6	46.8	70.2
Motor vehicle acc. (810-835)	23	19	4	22	18	4	21.0	20.5	23.4
Ill-defined (780-795)	74	43	31	74	44	30	70.7	50.2	175.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HINDS COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	5,197	2,677	2,520	5,059	2,515	2,544	29.6	25.6	35.2
Fetal deaths	344	195	149	304	160	144	60.1	63.6	56.6
Deaths from all causes	1,860	973	887	1,345	674	671	7.9	6.8	9.3
Deaths under 1 year	227	78	149	168	48	120	33.2	19.1	47.2
Deaths under 1 month	164	65	99	127	41	86	25.1	16.3	33.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	24	5	19	10	3	7	5.8	3.0	9.7
Tbc., respiratory (001-008)	20	4	16	9	2	7	5.3	2.0	9.7
Syphilis (020-029)	7	2	5	5	2	3	2.9	2.0	4.1
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	1	1	3	1	2	1.8	1.0	2.8
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	3	1	2	0	0	0	-	-	-
Whooping cough (056)	2	0	2	3	0	3	1.8	-	4.1
Meningococcal infections (057)	3	2	1	0	0	0	-	-	-
Acute poliomyelitis (080)	2	2	0	3	2	1	1.8	2.0	1.4
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	2	0	2	1	0	1	0.6	-	1.4
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	307	191	116	201	125	76	117.8	127.0	105.1
Diabetes (260)	20	10	10	17	8	9	10.0	8.1	12.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	196	101	95	141	73	68	82.6	74.2	94.0
Heart diseases (410-443)	521	320	201	388	237	151	227.3	240.8	208.8
Arteriosclerosis (450)	23	12	11	22	11	11	12.9	11.2	15.2
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	33	8	25	31	9	22	18.2	9.1	30.4
#Diarrhea & enteritis (571)	10	2	8	5	2	3	2.9	2.0	4.1
Acute nephritis (590, 591)	16	4	12	10	1	9	5.8	1.0	12.4
Chronic nephritis, etc. (592-594)	33	12	21	21	5	16	12.3	5.1	22.1
Maternal deaths (640-689)	4	0	4	2	0	2	0.4	-	0.8
Cong. malformations (750-759)	40	16	24	19	10	9	11.1	10.2	12.4
Birth inj., atelect. (760-762)	35	18	17	31	14	17	6.1	5.6	6.7
#Other dis. early inf. (763-776)	91	32	59	71	16	55	14.0	6.4	21.6
Violent & acc. deaths (800-999)	131	65	66	114	55	59	66.8	55.9	81.6
Acc. deaths, total (800-962)	97	51	46	83	43	40	48.6	43.7	55.3
Motor vehicle acc. (810-835)	42	25	17	33	23	10	19.3	23.4	13.8
Ill-defined (780-795)	125	30	95	121	30	91	70.9	30.5	125.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HOLMES COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	958	158	800	957	153	804	34.2	22.5	37.9
Fetal deaths	86	17	69	82	13	69	85.7	85.0	85.8
Deaths from all causes	299	86	213	336	92	244	12.0	13.5	11.5
Deaths under 1 year	43	4	39	40	4	36	41.8	26.1	44.8
Deaths under 1 month	26	3	23	25	3	22	26.1	19.6	27.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	2	0	2	7.1	-	9.4
Tbc., respiratory (001-008)	1	0	1	1	0	1	3.6	-	4.7
Syphilis (020-029)	1	0	1	1	0	1	3.6	-	4.7
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	1	0	3.6	14.7	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	1	0	1	3.6	-	4.7
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	35	11	24	41	14	27	146.4	205.9	127.4
Diabetes (260)	5	1	4	6	1	5	21.4	14.7	23.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	47	16	31	51	14	37	182.1	205.9	174.5
Heart diseases (410-443)	71	34	37	83	36	47	296.4	529.4	221.7
Arteriosclerosis (450)	11	3	8	11	3	8	39.3	44.1	37.7
Influenza (480-483)	8	0	8	8	0	8	28.6	-	37.7
#Pneumonia (490-493)	13	3	10	13	3	10	46.4	44.1	47.2
#Diarrhea & enteritis (571)	3	0	3	1	0	1	3.6	-	4.7
Acute nephritis (590, 591)	1	0	1	1	0	1	3.6	-	4.7
Chronic nephritis, etc. (592-594)	3	0	3	6	0	6	21.4	-	28.3
Maternal deaths (640-689)	4	0	4	5	0	5	5.2	-	6.2
Cong. malformations (750-759)	3	3	0	3	3	0	10.7	44.1	-
Birth inj., atelect. (760-762)	8	0	8	8	0	8	8.4	-	10.0
#Other dis. early inf. (763-776)	14	2	12	13	2	11	13.6	13.1	13.7
Violent & acc. deaths (800-999)	26	4	22	30	5	25	107.1	73.5	117.9
Acc. deaths, total (800-962)	20	3	17	23	4	19	82.1	58.8	89.6
Motor vehicle acc. (810-835)	6	0	6	6	0	6	21.4	-	28.3
Ill-defined (780-795)	27	2	25	26	2	24	92.8	29.4	113.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HUMPHREYS COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	726	114	612	749	141	608	35.2	23.5	39.7
Fetal deaths	32	1	31	33	1	32	44.0	7.1	52.6
Deaths from all causes	211	44	167	227	53	174	10.6	8.8	11.4
Deaths under 1 year	36	4	32	35	5	30	46.7	35.5	49.3
Deaths under 1 month	16	2	14	16	2	14	21.4	14.2	23.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	4.7	-	6.5
Tbc., respiratory (001-008)	1	0	1	1	0	1	4.7	-	6.5
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	1	0	4.7	16.7	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	2	0	2	2	0	2	9.4	-	13.1
Meningococcal infections (057)	1	0	1	1	0	1	4.7	-	6.5
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	4	13	24	6	18	112.7	100.0	117.6
Diabetes (260)	1	0	1	2	1	1	9.4	16.7	6.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	26	6	20	31	7	24	145.5	116.7	156.9
Heart diseases (410-443)	49	17	32	51	19	32	239.4	316.7	209.2
Arteriosclerosis (450)	2	0	2	2	0	2	9.4	-	13.1
Influenza (480-483)	4	0	4	4	0	4	18.8	-	26.1
#Pneumonia (490-493)	6	2	4	7	2	5	32.9	33.3	32.7
#Diarrhea & enteritis (571)	8	1	7	7	1	6	32.9	16.7	39.2
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	7	0	7	5	0	5	23.5	-	32.7
Maternal deaths (640-689)	2	0	2	2	0	2	2.7	-	3.3
Cong. malformations (750-759)	1	0	1	1	0	1	4.7	-	6.5
Birth inj., atelect. (760-762)	2	1	1	2	1	1	2.7	7.1	1.6
#Other dis. early inf. (763-776)	10	0	10	8	0	8	10.7	-	13.2
Violent & acc. deaths (800-999)	23	5	18	20	3	17	93.9	50.0	111.1
Acc. deaths, total (800-962)	22	5	17	18	2	16	84.5	33.3	104.6
Motor vehicle acc. (810-835)	7	2	5	6	1	5	28.2	16.7	32.7
Illi-defined (780-795)	32	0	32	31	1	30	145.5	16.7	196.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ISSAQENA COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		Non-white
				Total	White	Non-white	Total	White	
Live births	45	1	44	105	23	82	21.9	12.1	28.3
Fetal deaths	0	0	0	2	0	2	19.0	-	24.4
Deaths from all causes	22	2	20	33	8	25	6.9	4.2	8.6
Deaths under 1 year	2	0	2	2	0	2	19.0	-	24.4
Deaths under 1 month	1	0	1	1	0	1	9.5	-	12.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	20.8	-	34.5
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	2	0	2	3	2	1	62.5	105.3	34.5
Diabetes (260)	0	0	0	1	0	1	20.8	-	34.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	2	0	2	3	0	3	62.5	-	103.4
Heart diseases (410-443)	6	0	6	12	4	8	250.0	210.5	275.9
Arteriosclerosis (450)	1	1	0	1	1	0	20.8	52.6	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	0	0	0	0	0	0	-	-	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	1	0	1	1	0	1	9.5	-	12.2
Violent & acc. deaths (800-999)	1	0	1	2	1	1	41.7	52.6	34.5
Acc. deaths, total (800-962)	1	0	1	1	0	1	20.8	-	34.5
Motor vehicle acc. (810-835)	1	0	1	0	0	0	-	-	-
Ill-defined (780-795)	7	1	6	6	0	6	125.0	-	206.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ITAWAMBA COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	107	89	18	282	257	25	20.7	20.2	27.8
Fetal deaths	2	1	1	8	7	1	28.4	27.2	40.0
Deaths from all causes	95	92	3	122	120	2	9.0	9.4	2.2
Deaths under 1 year	1	1	0	11	11	0	39.0	42.8	-
Deaths under 1 month	0	0	0	9	9	0	31.9	35.0	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	2	0	2	2	0	14.7	15.7	-
Tbc., respiratory (001-008)	2	2	0	2	2	0	14.7	15.7	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	1	0	7.4	7.9	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	18	18	0	20	20	0	147.0	157.5	-
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	13	13	0	15	15	0	110.3	118.1	-
Heart diseases (410-443)	37	36	1	43	42	1	316.2	330.7	111.1
Arteriosclerosis (450)	1	1	0	1	1	0	7.4	7.9	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	2	0	2	2	0	14.7	15.7	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	2	0	2	2	0	14.7	15.7	-
Maternal deaths (640-689)	2	2	0	2	2	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	1	1	0	3	3	0	22.0	23.6	-
#Other dis. early inf. (763-776)	0	0	0	4	4	0	14.2	15.6	-
Violent & acc. deaths (800-999)	6	4	2	9	8	1	66.2	63.0	111.1
Acc. deaths, total (800-962)	4	2	2	6	5	1	44.1	39.4	111.1
Motor vehicle acc. (810-835)	3	2	1	4	4	0	29.4	31.5	-
Ill-defined (780-795)	4	4	0	4	4	0	29.4	31.5	-

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JACKSON COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,056	714	342	1,281	916	365	30.1	26.9	42.9
Fetal deaths	47	36	11	56	43	13	43.7	46.9	35.6
Deaths from all causes	285	212	73	331	250	81	7.8	7.4	9.5
Deaths under 1 year	26	15	11	33	22	11	25.8	24.0	30.1
Deaths under 1 month	18	12	6	22	16	6	17.2	17.5	16.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	2	0	4	2	2	9.4	5.9	23.5
Tbc., respiratory (001-008)	2	2	0	4	2	2	9.4	5.9	23.5
Syphilis (020-029)	0	0	0	1	0	1	2.4	-	11.8
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	1	0	2	1	1	4.7	2.9	11.8
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	31	26	5	35	30	5	82.4	88.2	58.8
Diabetes (260)	7	6	1	8	7	1	18.8	20.6	11.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	40	28	12	41	29	12	96.5	85.3	141.2
Heart diseases (410-443)	87	68	19	109	89	20	256.5	261.8	235.3
Arteriosclerosis (450)	7	6	1	8	7	1	18.8	20.6	11.8
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	8	6	2	7	5	2	16.5	14.7	23.5
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	0	1	1	0	1	2.4	-	11.8
Chronic nephritis, etc.(592-594)	6	2	4	6	3	3	14.1	8.8	35.3
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	4	4	0	9.4	11.8	-
Birth inj., atelect. (760-762)	9	8	1	12	11	1	9.4	12.0	2.7
#Other dis. early inf. (763-776)	7	3	4	7	3	4	5.5	3.3	11.0
Violent & acc. deaths (800-999)	35	24	11	33	25	8	77.6	73.5	94.1
Acc. deaths, total (800-962)	31	22	9	30	24	6	70.6	70.6	70.6
Motor vehicle acc. (810-835)	11	9	2	16	14	2	37.6	41.2	23.5
Ill-defined (780-795)	24	17	7	22	15	7	51.8	44.1	82.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JASPER COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	193	67	126	440	140	300	26.0	17.5	33.7
Fetal deaths	7	1	6	21	5	16	47.7	35.7	53.3
Deaths from all causes	86	36	50	153	74	79	9.0	9.2	8.9
Deaths under 1 year	8	1	7	11	2	9	25.0	14.3	30.0
Deaths under 1 month	5	1	4	7	2	5	15.9	14.3	16.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	5.9	-	11.2
Tbc., respiratory (001-008)	0	0	0	1	0	1	5.9	-	11.2
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	0	1	5.9	-	11.2
Acute poliomyelitis (080)	0	0	0	1	0	1	5.9	-	11.2
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	11	6	5	19	11	8	112.4	137.5	89.9
Diabetes (260)	3	3	0	4	4	0	23.7	50.0	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	17	7	10	24	14	10	142.0	175.0	112.4
Heart diseases (410-443)	20	6	14	42	19	23	248.5	237.5	258.4
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	1	1	5	3	2	29.6	37.5	22.5
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	2	2	9	4	5	53.2	50.0	56.2
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	0	1	5.9	-	11.2
Birth inj., atelect. (760-762)	2	1	1	3	2	1	6.8	14.3	3.3
#Other dis. early inf. (763-776)	2	0	2	3	0	3	6.8	-	10.0
Violent & acc. deaths (800-999)	8	5	3	16	7	9	94.7	87.5	101.1
Acc. deaths, total (800-962)	8	5	3	13	6	7	76.9	75.0	78.6
Motor vehicle acc. (810-835)	4	3	1	6	4	2	35.5	50.0	22.5
Ill-defined (780-795)	11	2	9	11	2	9	65.1	25.0	101.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JEFFERSON COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	152	0	152	344	44	300	34.0	17.6	39.5
Fetal deaths	4	0	4	10	4	6	29.1	90.9	20.0
Deaths from all causes	80	17	63	121	32	89	12.0	12.8	11.7
Deaths under 1 year	7	0	7	17	0	17	49.4	-	56.7
Deaths under 1 month	4	0	4	13	0	13	37.8	-	43.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	2	0	2	2	0	2	19.8	-	26.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	9.9	-	13.2
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	3	1	2	10	3	7	99.0	120.0	92.1
Diabetes (260)	2	0	2	2	0	2	19.8	-	26.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	19	6	13	21	8	13	207.9	320.0	171.0
Heart diseases (410-443)	30	6	24	39	11	28	386.1	440.0	368.4
Arteriosclerosis (450)	0	0	0	2	1	1	19.8	40.0	13.2
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	4	0	4	4	0	4	39.6	-	52.6
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	1	0	1	1	0	9.9	40.0	-
Chronic nephritis, etc. (592-594)	1	0	1	2	1	1	19.8	40.0	13.2
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	0	1	9.9	-	13.2
Birth inj., atelect. (760-762)	0	0	0	4	0	4	11.6	-	13.3
#Other dis. early inf. (763-776)	3	0	3	5	0	5	14.5	-	16.7
Violent & acc. deaths (800-999)	4	2	2	10	4	6	99.0	160.0	78.9
Acc. deaths, total (800-962)	3	2	1	9	4	5	89.1	160.0	65.8
Motor vehicle acc. (810-835)	1	1	0	3	2	1	29.7	80.0	13.2
Ill-defined (780-795)	8	1	7	10	2	8	99.0	80.0	105.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JEFF DAVIS COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	412	156	256	382	115	267	23.6	18.0	27.2
Fetal deaths	24	7	17	24	6	18	62.8	52.2	67.4
Deaths from all causes	120	56	64	120	55	65	7.4	8.6	6.6
Deaths under 1 year	16	2	14	15	3	12	39.3	26.1	44.9
Deaths under 1 month	4	1	3	4	1	3	10.5	8.7	11.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	6.2	15.6	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	6.2	15.6	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	1	0	1	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	7	10	19	9	10	117.3	140.6	102.0
Diabetes (260)	2	2	0	2	1	1	12.3	15.6	10.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	18	9	9	13	5	8	80.2	78.1	81.6
Heart diseases (410-443)	35	20	15	37	22	15	228.4	343.8	153.1
Arteriosclerosis (450)	1	1	0	2	2	0	12.3	31.2	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	0	2	1	0	1	6.2	-	10.2
#Diarrhea & enteritis (571)	2	0	2	2	0	2	12.3	-	20.4
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	6	5	1	5	3	2	30.9	46.9	20.4
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	3	1	2	3	1	2	18.5	15.6	20.4
Birth inj., atelect. (760-762)	1	1	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	5	0	5	6	1	5	15.7	8.7	18.7
Violent & acc. deaths (800-999)	8	4	4	11	5	6	67.9	78.1	61.2
Acc. deaths, total (800-962)	8	4	4	11	5	6	67.9	78.1	61.2
Motor vehicle acc. (810-835)	1	0	1	3	1	2	18.5	15.6	20.4
Ill-defined (780-795)	7	0	7	7	0	7	43.2	-	71.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JONES COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	2,133	1,123	1,010	1,452	885	567	28.2	23.5	41.1
Fetal deaths	103	44	59	70	35	35	48.2	39.5	61.7
Deaths from all causes	513	323	190	451	304	147	8.8	8.1	10.6
Deaths under 1 year	53	20	33	35	13	22	24.1	14.7	38.8
Deaths under 1 month	32	14	18	21	9	12	14.5	10.2	21.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	7	2	5	11	5	6	21.4	13.3	43.5
Tbc., respiratory (001-008)	7	2	5	10	5	5	19.4	13.3	36.2
Syphilis (020-029)	2	2	0	2	2	0	3.9	5.3	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	1	0	3	2	1	5.8	5.3	7.2
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	1	0	1	1	0	1.9	2.6	-
Acute poliomyelitis (080)	3	0	3	1	0	1	1.9	-	7.2
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	1.9	-	7.2
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	72	56	16	65	51	14	126.2	135.3	101.4
Diabetes (260)	6	3	3	7	4	3	13.6	10.6	21.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	67	44	23	63	40	23	122.3	106.1	166.7
Heart diseases (410-443)	147	103	44	138	106	32	268.0	281.2	231.9
Arteriosclerosis (450)	5	4	1	5	4	1	9.7	10.6	7.2
Influenza (480-483)	1	0	1	1	0	1	1.9	-	7.2
#Pneumonia (490-493)	20	8	12	19	10	9	36.9	26.5	65.2
#Diarrhea & enteritis (571)	4	3	1	1	1	0	1.9	2.6	-
Acute nephritis (590, 591)	9	2	7	8	1	7	15.5	2.6	50.7
Chronic nephritis, etc. (592-594)	9	6	3	5	4	1	9.7	10.6	7.2
Maternal deaths (640-689)	2	0	2	1	0	1	0.7	-	1.8
Cong. malformations (750-759)	7	3	4	5	3	2	9.7	8.0	14.5
Birth inj., atelect. (760-762)	15	8	7	9	6	3	6.2	6.8	5.3
#Other dis. early inf. (763-776)	13	6	7	10	3	7	6.9	3.4	12.3
Violent & acc. deaths (800-999)	46	27	19	34	23	11	66.0	61.0	79.7
Acc. deaths, total (800-962)	38	25	13	28	20	8	54.4	53.0	58.0
Motor vehicle acc. (810-835)	24	18	6	16	11	5	31.1	29.2	36.2
Ill-defined (780-795)	23	9	14	16	8	8	31.1	21.2	58.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
KEMPER COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	273	42	231	344	75	269	28.9	20.3	32.8
Fetal deaths	5	0	5	11	1	10	32.0	13.3	37.2
Deaths from all causes	107	45	62	125	56	69	10.5	15.1	8.4
Deaths under 1 year	6	0	6	7	0	7	20.3	-	26.0
Deaths under 1 month	1	0	1	2	0	2	5.8	-	7.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	2	2	0	16.8	54.0	-
Tbc., respiratory (001-008)	1	1	0	2	2	0	16.8	54.0	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	8.4	-	12.2
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	4	6	18	10	8	151.3	270.3	97.6
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	18	8	10	20	10	10	168.1	270.3	122.0
Heart diseases (410-443)	24	13	11	29	16	13	243.7	432.4	158.5
Arteriosclerosis (450)	4	3	1	4	3	1	33.6	81.1	12.2
Influenza (480-483)	2	1	1	1	0	1	8.4	-	12.2
#Pneumonia (490-493)	2	1	1	5	3	2	42.0	81.1	24.4
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	0	3	3	0	3	25.2	-	36.6
Maternal deaths (640-689)	0	0	0	1	1	0	2.9	13.3	-
Cong. malformations (750-759)	0	0	0	1	0	1	8.4	-	12.2
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	1	0	1	1	0	1	2.9	-	3.7
Violent & acc. deaths (800-999)	10	6	4	6	3	3	50.4	81.1	36.6
Acc. deaths, total (800-962)	8	5	3	6	3	3	50.4	81.1	36.6
Motor vehicle acc. (810-835)	4	4	0	3	2	1	25.2	54.0	12.2
Ill-defined (780-795)	20	4	16	21	4	17	176.5	108.1	207.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LAFAYETTE COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	505	230	275	525	244	281	32.4	30.1	34.7
Fetal deaths	15	6	9	14	6	8	26.7	24.6	28.5
Deaths from all causes	161	88	73	163	94	69	10.1	11.6	8.5
Deaths under 1 year	19	6	13	18	8	10	34.3	32.8	35.6
Deaths under 1 month	15	6	9	15	8	7	28.6	32.8	24.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	6.2	-	12.3
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	11	9	2	11	9	2	67.9	111.1	24.7
Diabetes (260)	3	3	0	3	3	0	18.5	37.0	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	27	15	12	26	14	12	160.5	172.8	148.1
Heart diseases (410-443)	59	40	19	62	41	21	382.7	506.2	259.2
Arteriosclerosis (450)	1	0	1	1	0	1	6.2	-	12.3
Influenza (480-483)	2	0	2	2	0	2	12.3	-	24.7
#Pneumonia (490-493)	4	0	4	4	1	3	24.7	12.3	37.0
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	0	1	1	0	1	6.2	-	12.3
Chronic nephritis, etc. (592-594)	4	0	4	3	1	2	18.5	12.3	24.7
Maternal deaths (640-689)	2	1	1	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	2	1	1	12.3	12.3	12.3
Birth inj., atelect. (760-762)	7	4	3	6	5	1	11.4	20.5	3.6
#Other dis. early inf. (763-776)	6	1	5	7	2	5	13.3	8.2	17.8
Violent & acc. deaths (800-999)	7	4	3	9	6	3	55.6	74.1	37.0
Acc. deaths, total (800-962)	6	4	2	8	6	2	49.4	74.1	24.7
Motor vehicle acc. (810-835)	4	3	1	7	5	2	43.2	61.7	24.7
Ill-defined (780-795)	11	2	9	11	2	9	67.9	24.7	111.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LAMAR COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	109	62	47	344	274	70	28.0	26.3	36.8
Fetal deaths	3	0	3	7	4	3	20.3	14.6	42.8
Deaths from all causes	72	47	25	102	73	29	8.3	7.0	15.3
Deaths under 1 year	3	1	2	8	6	2	23.2	21.9	28.6
Deaths under 1 month	2	1	1	5	4	1	14.5	14.6	14.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	1	1	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	8	7	1	15	14	1	122.0	134.6	52.6
Diabetes (260)	1	0	1	1	0	1	8.1	-	52.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	14	11	3	14	10	4	113.8	96.2	210.5
Heart diseases (410-443)	24	18	6	29	23	6	235.8	221.2	315.8
Arteriosclerosis (450)	0	0	0	2	2	0	16.3	19.2	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	1	1	4	2	2	32.5	19.2	105.3
#Diarrhea & enteritis (571)	0	0	0	1	0	1	8.1	-	52.6
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	0	1	8.1	-	-
Maternal deaths (640-689)	0	0	0	1	0	1	8.1	-	52.6
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	1	1	0	8.1	9.6	-
#Other dis. early inf. (763-776)	1	0	1	1	0	1	2.9	-	14.3
Violent & acc. deaths (800-999)	9	5	4	15	9	6	11.6	10.9	14.3
Acc. deaths, total (800-962)	6	3	3	12	7	5	122.0	86.5	315.8
Motor vehicle acc. (810-835)	1	1	0	4	3	1	97.6	67.3	263.2
Ill-defined (780-795)	8	2	6	8	3	5	65.0	28.8	52.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LAUDERDALE COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	2,065	1,086	979	1,613	858	755	26.0	21.8	33.2
Fetal deaths	102	32	70	81	30	51	50.2	35.0	67.5
Deaths from all causes	882	552	330	674	381	293	10.8	9.7	12.9
Deaths under 1 year	61	25	36	56	22	34	34.7	25.6	45.0
Deaths under 1 month	38	17	21	34	15	19	21.1	17.5	25.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	5	3	2	4	1	3	6.4	2.5	13.2
Tbc., respiratory (001-008)	5	3	2	4	1	3	6.4	2.5	13.2
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	1.6	-	4.4
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	2	0	2	2	0	2	3.2	-	8.8
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	96	64	32	78	51	27	125.6	129.4	118.9
Diabetes (260)	10	6	4	6	3	3	9.7	7.6	13.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	141	105	36	97	65	32	156.2	165.0	141.0
Heart diseases (410-443)	308	197	111	237	136	101	381.6	345.2	444.9
Arteriosclerosis (450)	15	13	2	8	6	2	12.9	15.2	8.8
Influenza (480-483)	3	3	0	4	4	0	6.4	10.2	-
#Pneumonia (490-493)	20	11	9	13	5	8	20.9	12.7	35.2
#Diarrhea & enteritis (571)	6	4	2	5	4	1	8.0	10.2	4.4
Acute nephritis (590, 591)	4	2	2	2	2	0	3.2	5.1	-
Chronic nephritis, etc. (592-594)	11	5	6	9	4	5	14.5	10.2	22.0
Maternal deaths (640-689)	4	1	3	1	0	1	0.6	-	1.3
Cong. malformations (750-759)	6	5	1	6	4	2	9.7	10.2	8.8
Birth inj., atelect. (760-762)	12	8	4	10	6	4	6.2	7.0	5.3
#Other dis. early inf. (763-776)	21	6	15	19	6	13	11.8	7.0	17.2
Violent & acc. deaths (800-999)	75	38	37	53	23	30	85.3	58.4	132.2
Acc. deaths, total (800-962)	65	35	30	43	20	23	69.2	50.8	101.3
Motor vehicle acc. (810-835)	34	24	10	18	10	8	29.0	25.4	35.2
Ill-defined (780-795)	61	20	41	59	18	41	95.0	45.7	180.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LAWRENCE COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	188	30	158	279	116	163	24.0	16.8	34.7
Fetal deaths	13	0	13	20	6	14	71.7	51.7	85.9
Deaths from all causes	57	34	23	96	65	31	8.3	9.4	6.6
Deaths under 1 year	4	0	4	7	1	6	25.1	8.6	36.8
Deaths under 1 month	2	0	2	3	1	2	10.8	8.6	12.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	8.6	14.5	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	8.6	14.5	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	8.6	-	21.3
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	6	5	1	10	9	1	86.2	130.4	21.3
Diabetes (260)	2	1	1	3	3	0	25.9	43.5	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	8	5	3	12	8	4	103.4	115.9	85.1
Heart diseases (410-443)	23	15	8	33	23	10	284.5	333.3	212.8
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	1	1	0	1	1	0	8.6	14.5	-
#Pneumonia (490-493)	1	0	1	2	0	2	17.2	-	42.6
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	0	1	1	0	1	8.6	-	21.3
Chronic nephritis, etc. (592-594)	0	0	0	2	2	0	17.2	29.0	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	1	1	0	8.6	14.5	-
Birth inj., atelect. (760-762)	1	0	1	1	0	1	3.6	-	6.1
#Other dis. early inf. (763-776)	0	0	0	1	1	0	3.6	8.6	-
Violent & acc. deaths (800-999)	3	0	3	10	6	4	86.2	87.0	85.1
Acc. deaths, total (800-962)	3	0	3	10	6	4	86.2	87.0	85.1
Motor vehicle acc. (810-835)	1	0	1	8	5	3	69.0	72.5	63.8
Ill-defined (780-795)	9	5	4	10	5	5	86.2	72.5	106.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LEAKE COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	441	170	271	467	170	297	30.1	20.7	40.7
Fetal deaths	12	2	10	19	5	14	40.7	29.4	47.1
Deaths from all causes	115	70	45	148	92	56	9.5	11.2	7.7
Deaths under 1 year	15	6	9	20	5	15	42.8	29.4	50.5
Deaths under 1 month	7	4	3	11	4	7	23.6	23.5	23.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	6.4	12.2	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	6.4	12.2	-
Syphilis (020-029)	1	0	1	1	0	1	6.4	-	13.7
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	2	0	2	2	0	2	12.9	-	27.4
Meningococcal infections (057)	0	0	0	1	1	0	6.4	12.2	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	18	12	6	21	14	7	135.5	170.7	95.9
Diabetes (260)	3	1	2	4	2	2	25.8	24.4	27.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	17	11	6	20	13	7	129.0	158.5	95.9
Heart diseases (410-443)	44	31	13	50	37	13	322.6	451.2	178.1
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	1	0	1	0	0	0	-	-	-
#Pneumonia (490-493)	1	0	1	2	1	1	12.9	12.2	13.7
#Diarrhea & enteritis (571)	0	0	0	1	0	1	6.4	-	13.7
Acute nephritis (590, 591)	1	1	0	1	1	0	6.4	12.2	-
Chronic nephritis, etc. (592-594)	2	1	1	3	2	1	19.4	24.4	13.7
Maternal deaths (640-689)	0	0	0	2	0	2	4.3	-	6.7
Cong. malformations (750-759)	2	0	2	3	1	2	19.4	12.2	27.4
Birth inj., atelect (760-762)	3	2	1	3	2	1	6.4	11.8	3.4
#Other dis. early inf. (763-776)	2	1	1	6	1	5	12.8	5.9	16.8
Violent & acc. deaths (800-999)	9	6	3	16	11	5	103.2	134.1	68.5
Acc. deaths, total (800-962)	8	5	3	13	8	5	83.9	97.6	68.5
Motor vehicle acc. (810-835)	3	3	0	5	4	1	32.2	48.8	13.7
Ill-defined (780-795)	3	1	2	2	0	2	12.9	-	27.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LEE COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,274	889	385	1,059	682	377	28.3	26.2	33.1
Fetal deaths	27	18	9	23	14	9	21.7	20.5	23.9
Deaths from all causes	402	301	101	375	272	103	10.0	10.5	9.0
Deaths under 1 year	51	41	10	31	23	8	29.3	33.7	21.2
Deaths under 1 month	39	33	6	24	18	6	22.7	26.4	15.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	5	5	0	5	4	1	13.4	15.4	8.8
Tbc., respiratory (001-008)	5	5	0	5	4	1	13.4	15.4	8.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	1	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	2.7	-	8.8
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	57	43	14	58	45	13	155.1	173.1	114.0
Diabetes (260)	5	4	1	6	4	2	16.0	15.4	17.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	59	41	18	63	45	18	168.4	173.1	157.9
Heart diseases (410-443)	147	110	37	137	99	38	366.3	380.8	333.3
Arteriosclerosis (450)	3	3	0	3	3	0	8.0	11.5	-
Influenza (480-483)	1	1	0	1	1	0	2.7	3.8	-
#Pneumonia (490-493)	14	10	4	15	11	4	40.1	42.3	35.1
#Diarrhea & enteritis (571)	3	2	1	1	1	0	2.7	3.8	-
Acute nephritis (590, 591)	2	1	1	2	1	1	5.3	3.8	8.8
Chronic nephritis, etc. (592-594)	4	3	1	4	3	1	10.7	11.5	8.8
Maternal deaths (640-689)	1	1	0	1	1	0	0.9	1.5	-
Cong. malformations (750-759)	9	7	2	4	2	2	10.7	7.7	17.5
Birth inj., atelect. (760-762)	14	12	2	10	8	2	9.4	11.7	5.3
#Other dis. early inf. (763-776)	19	16	3	11	8	3	10.4	11.7	8.0
Violent & acc. deaths (800-999)	23	18	5	20	13	7	53.5	50.0	61.4
Acc. deaths, total (800-962)	18	15	3	16	11	5	42.8	42.3	43.8
Motor vehicle acc. (810-835)	11	9	2	7	4	3	18.7	15.4	26.3
Ill-defined (780-795)	13	9	4	13	9	4	34.8	34.6	35.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LEFLOR COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number		Non-white	Total	White	Non-white
				Total	White				
Live births	1,820	509	1,311	1,678	386	1,292	36.5	24.4	42.8
Fetal deaths	75	13	62	73	11	62	43.5	28.5	48.0
Deaths from all causes	561	184	377	498	133	365	10.8	8.4	12.1
Deaths under 1 year	101	13	88	87	6	81	51.8	15.5	62.7
Deaths under 1 month	53	11	42	43	5	38	25.6	13.0	29.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	0	4	5	0	5	10.9	-	16.6
Tbc., respiratory (001-008)	1	0	1	2	0	2	4.3	-	6.6
Syphilis (020-029)	1	1	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	2.2	-	3.3
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	2.2	-	3.3
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	40	18	22	35	15	20	76.1	94.9	66.2
Diabetes (260)	8	2	6	6	1	5	13.0	6.3	16.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	80	30	50	69	18	51	150.0	113.9	168.9
Heart diseases (410-443)	154	71	83	134	53	81	291.3	335.4	268.2
Arteriosclerosis (450)	6	5	1	5	4	1	10.9	25.3	3.3
Influenza (480-483)	4	1	3	4	1	3	8.7	6.3	9.9
#Pneumonia (490-493)	22	4	18	17	2	15	37.0	12.6	49.7
#Diarrhea & enteritis (571)	15	0	15	15	0	15	32.6	-	49.7
Acute nephritis (590, 591)	2	0	2	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	10	1	9	10	1	9	21.7	6.3	29.8
Maternal deaths (640-689)	5	0	5	5	0	5	3.0	-	3.9
Cong. malformations (750-759)	2	1	1	1	0	1	2.2	-	3.3
Birth inj., atelect. (760-762)	10	3	7	6	1	5	3.6	2.6	3.9
#Other dis. early inf. (763-776)	35	7	28	31	4	27	18.5	10.4	20.9
Violent & acc. deaths (800-999)	37	14	23	36	11	25	78.3	69.6	82.8
Acc. deaths, total (800-962)	26	11	15	26	8	18	56.5	50.6	59.6
Motor vehicle acc. (810-835)	7	4	3	7	2	5	15.2	12.6	16.6
Ill-defined (780-795)	81	7	74	76	5	71	165.2	31.6	235.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LINCOLN COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	691	407	284	676	368	308	25.8	22.0	32.4
Fetal deaths	26	19	7	25	17	8	37.0	46.2	26.0
Deaths from all causes	235	154	81	271	174	97	10.3	10.4	10.2
Deaths under 1 year	36	16	20	38	16	22	56.2	43.5	71.4
Deaths under 1 month	24	14	10	26	14	12	38.5	38.0	39.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	3.8	6.0	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	3.8	6.0	-
Syphilis (020-029)	0	0	0	1	0	1	3.8	-	10.5
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	3	0	3	3	0	3	11.4	-	31.6
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	26	18	8	31	19	12	118.3	113.8	126.3
Diabetes (260)	2	2	0	3	3	0	11.4	18.0	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	18	7	11	18	6	12	68.7	35.9	126.3
Heart diseases (410-443)	95	69	26	99	72	27	377.9	431.1	284.2
Arteriosclerosis (450)	1	1	0	2	2	0	7.6	12.0	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	6	1	5	6	2	4	22.9	12.0	42.1
#Diarrhea & enteritis (571)	1	1	0	2	1	1	7.6	6.0	10.5
Acute nephritis (590, 591)	1	1	0	2	1	1	7.6	6.0	10.5
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	3.8	6.0	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	3.8	6.0	-
Birth inj., atelect. (760-762)	13	7	6	12	6	6	17.8	16.3	19.5
#Other dis. early inf. (763-776)	12	7	5	14	7	7	20.7	19.0	22.7
Violent & acc. deaths (800-999)	20	16	4	37	29	8	141.2	173.6	84.2
Acc. deaths, total (800-962)	18	14	4	32	26	6	122.1	155.7	63.2
Motor vehicle acc. (810-835)	13	12	1	24	20	4	91.6	119.8	42.1
Ill-defined (780-795)	15	6	9	14	6	8	53.4	35.9	84.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LOWNDES COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,069	504	565	1,068	501	567	28.6	24.1	34.2
Fetal deaths	29	11	18	26	10	16	24.3	20.0	28.2
Deaths from all causes	357	173	184	359	172	187	9.6	8.3	11.3
Deaths under 1 year	40	14	26	38	10	28	35.6	20.0	49.4
Deaths under 1 month	19	10	9	17	7	10	15.9	14.0	17.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	3	1	7	3	4	18.7	14.4	24.1
Tbc., respiratory (001-008)	3	2	1	5	2	3	13.4	9.6	18.1
Syphilis (020-029)	0	0	0	1	0	1	2.7	-	6.0
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	38	18	20	39	20	19	104.3	96.2	114.4
Diabetes (260)	2	1	1	2	1	1	5.3	4.8	6.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	41	17	24	48	20	28	128.3	96.2	168.7
Heart diseases (410-443)	111	72	39	114	72	42	304.8	346.2	253.0
Arteriosclerosis (450)	9	7	2	10	8	2	26.7	38.5	12.0
Influenza (480-483)	2	0	2	2	0	2	5.3	-	12.0
#Pneumonia (490-493)	9	4	5	11	6	5	29.4	28.8	30.1
#Diarrhea & enteritis (571)	5	0	5	5	0	5	13.4	-	30.1
Acute nephritis (590, 591)	1	1	0	1	1	0	2.7	4.8	-
Chronic nephritis, etc. (592-594)	4	0	4	4	0	4	10.7	-	24.1
Maternal deaths (640-689)	2	0	2	2	0	2	1.9	-	3.5
Cong. malformations (750-759)	3	3	0	2	2	0	5.3	9.6	-
Birth inj., atelect. (760-762)	7	5	2	6	4	2	5.6	8.0	3.5
#Other dis. early inf. (763-776)	9	3	6	8	2	6	7.5	4.0	10.6
Violent & acc. deaths (800-999)	29	15	14	24	11	13	64.2	52.9	78.3
Acc. deaths, total (800-962)	22	11	11	18	7	11	48.1	33.6	66.3
Motor vehicle acc. (810-835)	9	7	2	5	3	2	13.4	14.4	12.0
Ill-defined (780-795)	45	5	40	41	4	37	109.6	19.2	222.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MADISON COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,185	194	991	1,146	178	968	35.5	22.0	40.0
Fetal deaths	37	2	35	34	1	33	29.7	5.6	34.1
Deaths from all causes	351	117	234	332	97	235	10.3	12.0	9.7
Deaths under 1 year	41	4	37	46	6	40	40.1	33.7	41.3
Deaths under 1 month	30	4	26	32	6	26	27.9	33.7	26.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	0	3	4	1	3	12.4	12.3	12.4
Tbc., respiratory (001-008)	3	0	3	4	1	3	12.4	12.3	12.4
Syphilis (020-029)	1	0	1	1	0	1	3.1	-	4.1
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	39	18	21	37	15	22	114.6	185.2	90.9
Diabetes (260)	5	3	2	4	2	2	12.4	24.7	8.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	56	15	41	52	13	39	161.0	160.5	161.2
Heart diseases (410-443)	86	38	48	81	34	47	250.8	419.8	194.2
Arteriosclerosis (450)	3	3	0	5	5	0	15.5	61.7	-
Influenza (480-483)	1	0	1	1	0	1	3.1	-	4.1
#Pneumonia (490-493)	8	1	7	9	1	8	27.9	12.3	33.0
#Diarrhea & enteritis (571)	4	1	3	4	1	3	12.4	12.3	12.4
Acute nephritis (590, 591)	2	1	1	2	1	1	6.2	12.3	4.1
Chronic nephritis, etc. (592-594)	8	4	4	8	4	4	24.8	49.4	16.5
Maternal deaths (640-689)	1	0	1	2	0	2	1.7	-	2.1
Cong. malformations (750-759)	1	0	1	5	1	4	15.5	12.3	16.5
Birth inj., atelect. (760-762)	7	0	7	8	1	7	7.0	5.6	7.2
#Other dis. early inf. (763-776)	19	3	16	19	3	16	16.6	16.8	16.5
Violent & acc. deaths (800-999)	41	15	26	24	3	21	74.3	37.0	86.8
Acc. deaths, total (800-962)	35	15	20	18	3	15	55.7	37.0	62.0
Motor vehicle acc. (810-835)	16	10	6	5	2	3	15.5	24.7	12.4
Ill-defined (780-795)	40	6	34	38	4	34	117.6	49.4	140.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MARION COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	562	323	239	633	369	264	27.9	27.7	28.1
Fetal deaths	24	11	13	29	14	15	45.8	37.9	56.8
Deaths from all causes	173	105	68	205	124	81	9.0	9.3	8.6
Deaths under 1 year	17	8	9	20	10	10	31.6	27.1	37.9
Deaths under 1 month	9	5	4	11	6	5	17.4	16.3	18.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	2	0	2	8.8	-	21.3
Tbc., respiratory (001-008)	0	0	0	2	0	2	8.8	-	21.3
Syphilis (020-029)	0	0	0	1	1	0	4.4	7.5	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	1	0	1	4.4	-	10.6
Whooping cough (056)	1	1	0	1	1	0	4.4	7.5	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	12	5	20	15	5	88.1	112.8	53.2
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	27	10	17	32	13	19	141.0	97.7	202.1
Heart diseases (410-443)	56	39	17	62	42	20	273.1	315.8	212.8
Arteriosclerosis (450)	4	4	0	3	3	0	13.2	22.6	-
Influenza (480-483)	1	0	1	1	0	1	4.4	-	10.6
#Pneumonia (490-493)	6	2	4	8	3	5	35.2	22.6	53.2
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	1	0	2	2	0	8.8	15.0	-
Chronic nephritis, etc. (592-594)	4	0	4	4	0	4	17.6	-	42.6
Maternal deaths (640-689)	1	1	0	1	1	0	1.6	2.7	-
Cong. malformations (750-759)	3	3	0	5	3	2	22.0	22.6	21.3
Birth inj., atelect. (760-762)	2	1	1	3	2	1	4.7	5.4	3.8
#Other dis. early inf. (763-776)	5	3	2	5	3	2	7.9	8.1	7.6
Violent & acc. deaths (800-999)	21	17	4	23	21	2	101.3	157.9	21.3
Acc. deaths, total (800-962)	19	15	4	21	20	1	92.5	150.4	10.6
Motor vehicle acc. (810-835)	10	7	3	13	13	0	57.3	97.7	-
Ill-defined (780-795)	8	4	4	11	4	7	48.4	30.1	74.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MARSHALL COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	770	90	680	817	144	673	35.7	22.2	41.0
Fetal deaths	50	5	45	39	5	34	47.7	34.7	50.5
Deaths from all causes	197	64	133	207	69	138	9.0	10.6	8.4
Deaths under 1 year	21	3	18	18	5	13	22.0	34.7	19.3
Deaths under 1 month	13	3	10	14	5	9	17.1	34.7	13.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	0	4	5	0	5	21.8	-	30.5
Tbc., respiratory (001-008)	4	0	4	5	0	5	21.8	-	30.5
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	1	0	4.4	15.4	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	1	0	1	4.4	-	6.1
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	19	8	11	24	10	14	104.8	153.8	85.4
Diabetes (260)	1	0	1	1	0	1	4.4	-	6.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	29	6	23	30	8	22	131.0	123.1	134.1
Heart diseases (410-443)	69	27	42	68	24	44	296.9	369.2	268.3
Arteriosclerosis (450)	1	1	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	6	3	3	7	5	2	30.6	76.9	12.2
#Diarrhea & enteritis (571)	1	0	1	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	3	1	2	2	0	8.7	30.8	-
Maternal deaths (640-689)	3	0	3	3	0	3	3.7	-	4.4
Cong. malformations (750-759)	3	0	3	2	0	2	8.7	-	12.2
Birth inj., atelect. (760-762)	2	1	1	2	1	1	2.4	6.9	1.5
#Other dis. early inf. (763-776)	9	2	7	7	3	4	8.6	20.8	5.9
Violent & acc. deaths (800-999)	20	5	15	18	6	12	78.6	92.3	73.2
Acc. deaths, total (800-962)	19	4	15	16	5	11	69.9	76.9	67.1
Motor vehicle acc. (810-835)	9	2	7	7	1	6	30.6	15.4	36.6
Ill-defined (780-795)	17	3	14	19	3	16	83.0	46.2	97.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MONROE COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	764	399	365	765	383	382	25.2	20.2	33.5
Fetal deaths	17	9	8	18	9	9	23.5	23.5	23.6
Deaths from all causes	311	173	138	322	180	142	10.6	9.5	12.4
Deaths under 1 year	34	9	25	33	7	26	43.1	18.3	68.1
Deaths under 1 month	18	6	12	17	5	12	22.2	13.0	31.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	3.3	5.3	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	3.3	5.3	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	0	2	3	1	2	9.9	5.3	17.5
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	0	1	1	0	1	3.3	-	8.8
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	38	23	15	40	25	15	131.6	131.6	131.6
Diabetes (260)	5	3	2	5	3	2	16.4	15.8	17.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	44	24	20	46	25	21	151.3	131.6	184.2
Heart diseases (410-443)	84	57	27	86	57	29	282.9	300.0	254.4
Arteriosclerosis (450)	8	6	2	10	8	2	32.9	42.1	17.5
Influenza (480-483)	2	1	1	2	1	1	6.6	5.3	8.8
#Pneumonia (490-493)	7	3	4	8	4	4	26.3	21.0	35.1
#Diarrhea & enteritis (571)	4	0	4	6	1	5	19.7	5.3	43.8
Acute nephritis (590, 591)	2	2	0	3	2	1	9.9	10.5	8.8
Chronic nephritis, etc. (592-594)	11	3	8	10	3	7	32.9	15.8	61.4
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	3	1	2	3	1	2	9.9	5.3	17.5
Birth inj., atelect. (760-762)	4	3	1	2	1	1	2.6	2.6	2.6
#Other dis. early inf. (763-776)	10	2	8	11	3	8	14.4	7.8	20.9
Violent & acc. deaths (800-999)	28	18	10	27	17	10	88.8	89.5	87.7
Acc. deaths, total (800-962)	20	13	7	17	11	6	55.9	57.9	52.6
Motor vehicle acc. (810-835)	9	6	3	7	5	2	23.0	26.3	17.5
Ill-defined (780-795)	28	8	20	26	7	19	85.5	36.8	166.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MONTGOMERY COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	297	90	207	359	138	221	26.0	18.6	34.5
Fetal deaths	7	2	5	9	4	5	25.1	29.0	22.6
Deaths from all causes	88	37	51	127	68	59	9.2	9.2	9.2
Deaths under 1 year	6	0	6	12	4	8	33.4	29.0	36.2
Deaths under 1 month	2	0	2	8	4	4	22.3	29.0	18.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	7.2	-	15.6
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	7	10	23	13	10	166.7	175.7	156.2
Diabetes (260)	0	0	0	1	1	0	7.2	13.5	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	5	2	3	9	5	4	65.2	67.6	62.5
Heart diseases (410-443)	36	20	16	51	33	18	369.6	445.9	281.2
Arteriosclerosis (450)	1	0	1	1	0	1	7.2	-	15.6
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	7	1	6	6	1	5	43.5	13.5	78.1
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	7.2	13.5	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	2	2	0	14.5	27.0	-
Birth inj., atelect. (760-762)	0	0	0	3	2	1	8.4	14.5	4.5
#Other dis. early inf. (763-776)	2	0	2	4	1	3	11.1	7.2	13.6
Violent & acc. deaths (800-999)	6	4	2	8	5	3	58.0	67.6	46.9
Acc. deaths, total (800-962)	6	4	2	6	3	3	43.5	40.5	46.9
Motor vehicle acc. (810-835)	2	2	0	2	0	2	14.5	-	31.2
Ill-defined (780-795)	7	0	7	7	0	7	50.7	-	109.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
NESHOBIA COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	559	283	276	590	339	251	32.4	26.7	45.6
Fetal deaths	20	9	11	18	7	11	30.5	20.6	43.8
Deaths from all causes	180	122	58	208	152	56	11.4	12.0	10.2
Deaths under 1 year	28	6	22	22	6	16	37.3	17.7	63.7
Deaths under 1 month	15	4	11	12	4	8	20.3	11.8	31.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	2	1	1	11.0	7.9	18.2
Tbc., respiratory (001-008)	2	1	1	2	1	1	11.0	7.9	18.2
Syphilis (020-029)	1	0	1	1	0	1	5.5	-	18.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	0	1	1	0	1	5.5	-	18.2
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	1	0	1	1	0	5.5	7.9	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	26	22	4	33	26	7	181.3	204.7	127.3
Diabetes (260)	1	1	0	2	2	0	11.0	15.7	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	27	20	7	35	27	8	192.3	212.6	145.4
Heart diseases (410-443)	57	46	11	63	49	14	346.2	385.8	254.5
Arteriosclerosis (450)	0	0	0	1	1	0	5.5	7.9	-
Influenza (480-483)	1	0	1	2	0	2	11.0	-	36.4
#Pneumonia (490-493)	5	2	3	7	4	3	38.5	31.5	54.5
#Diarrhea & enteritis (571)	1	0	1	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	1	1	0	5.5	7.9	-
Chronic nephritis, etc. (592-594)	2	1	1	1	1	0	5.5	7.9	-
Maternal deaths (640-689)	0	0	0	1	1	0	1.7	2.9	-
Cong. malformations (750-759)	4	0	4	4	2	2	22.0	15.7	36.4
Birth inj., atelect. (760-762)	4	4	0	3	3	0	5.1	8.8	-
#Other dis. early inf. (763-776)	5	0	5	3	0	3	5.1	-	12.0
Violent & acc. deaths (800-999)	15	7	8	16	10	6	87.9	78.7	109.1
Acc. deaths, total (800-962)	13	6	7	14	9	5	76.9	70.9	90.9
Motor vehicle acc. (810-835)	6	5	1	6	5	1	33.0	39.4	18.2
Ill-defined (780-795)	10	5	5	10	5	5	54.9	39.4	90.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
NEWTON COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	487	299	188	476	250	226	25.0	22.3	29.0
Fetal deaths	37	20	17	33	16	17	69.3	64.0	75.2
Deaths from all causes	171	118	53	179	116	63	9.4	10.4	8.1
Deaths under 1 year	19	8	11	19	7	12	39.9	28.0	53.1
Deaths under 1 month	16	7	9	17	7	10	35.7	28.0	44.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	1	0	1	1	0	5.3	8.9	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	5.3	-	12.8
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	26	16	10	25	14	11	131.6	125.0	141.0
Diabetes (260)	5	5	0	4	4	0	21.0	35.7	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	23	19	4	25	21	4	131.6	187.5	51.3
Heart diseases (410-443)	51	41	10	48	39	9	252.6	348.2	115.4
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	1	1	0	1	1	0	5.3	8.9	-
#Pneumonia (490-493)	8	3	5	7	2	5	36.8	17.8	64.1
#Diarrhea & enteritis (571)	1	1	0	1	0	1	5.3	-	12.8
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	3	2	1	15.8	17.8	12.8
Maternal deaths (640-689)	1	1	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	2	0	2	1	1	10.5	8.9	12.8
Birth inj., atelect. (760-762)	4	2	2	5	3	2	10.5	12.0	8.8
#Other dis. early inf. (763-776)	6	3	3	6	3	3	12.6	12.0	13.3
Violent & acc. deaths (800-999)	11	8	3	16	10	6	84.2	89.3	76.9
Acc. deaths, total (800-962)	6	5	1	11	7	4	57.9	62.5	51.3
Motor vehicle acc. (810-835)	2	2	0	6	5	1	31.6	44.6	12.8
Ill-defined (780-795)	13	1	12	16	1	15	84.2	8.9	192.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
NOXUBEE COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	623	97	526	619	103	516	36.0	21.4	41.6
Fetal deaths	42	3	39	42	2	40	67.8	19.4	77.5
Deaths from all causes	151	51	100	176	59	117	10.2	12.3	9.4
Deaths under 1 year	19	0	19	20	0	20	32.3	-	38.8
Deaths under 1 month	12	0	12	12	0	12	19.4	-	23.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	2	0	2	11.6	-	16.1
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	1	0	1	1	0	1	5.8	-	8.1
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	20	8	12	21	8	13	122.1	166.7	104.8
Diabetes (260)	1	1	0	2	1	1	11.6	20.8	8.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	23	14	9	29	17	12	168.6	354.2	96.8
Heart diseases (410-443)	38	22	16	42	23	19	244.2	479.2	153.2
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	9	3	6	8	2	6	46.5	41.7	48.4
#Diarrhea & enteritis (571)	2	0	2	2	0	2	11.6	-	16.1
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	2	2	0	11.6	41.7	-
Maternal deaths (640-689)	1	0	1	1	0	1	1.6	-	1.9
Cong. malformations (750-759)	0	0	0	1	0	1	5.8	-	8.1
Birth inj., atelect. (760-762)	1	0	1	1	0	1	1.6	-	1.9
#Other dis. early inf. (763-776)	9	0	9	9	0	9	14.5	-	17.4
Violent & acc. deaths (800-999)	12	1	11	15	2	13	87.2	41.7	104.8
Acc. deaths, total (800-962)	10	1	9	11	2	9	64.0	41.7	72.6
Motor vehicle acc. (810-835)	4	1	3	4	1	3	23.2	20.8	24.2
Ill-defined (780-795)	26	1	25	30	2	28	174.4	41.7	225.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
OKTIBBEHA COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	837	370	467	754	292	462	34.3	26.1	42.8
Fetal deaths	20	9	11	20	10	10	26.5	34.2	21.6
Deaths from all causes	203	90	113	210	91	119	9.5	8.1	11.0
Deaths under 1 year	33	7	26	32	6	26	42.4	20.5	56.3
Deaths under 1 month	22	6	16	21	5	16	27.8	17.1	34.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	2	1	1	9.1	8.9	9.2
Tbc., respiratory (001-008)	1	1	0	2	1	1	9.1	8.9	9.2
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	4	0	4	4	0	4	18.2	-	37.0
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	18	7	11	20	8	12	90.9	71.4	111.1
Diabetes (260)	2	0	2	2	0	2	9.1	-	18.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	31	14	17	32	14	18	145.4	125.0	166.7
Heart diseases (410-443)	80	44	36	75	40	35	340.9	357.1	324.1
Arteriosclerosis (450)	2	2	0	3	3	0	13.6	26.8	-
Influenza (480-483)	1	1	0	1	1	0	4.5	8.9	-
#Pneumonia (490-493)	6	1	5	6	1	5	27.3	8.9	46.3
#Diarrhea & enteritis (571)	2	0	2	2	0	2	9.1	-	18.5
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	3	1	4	2	2	18.2	17.8	18.5
Maternal deaths (640-689)	1	0	1	1	0	1	1.3	-	2.2
Cong. malformations (750-759)	3	0	3	3	0	3	13.6	-	27.8
Birth inj., atelect. (760-762)	7	5	2	7	5	2	9.3	17.1	4.3
#Other dis. early inf. (763-776)	8	1	7	7	0	7	9.3	-	15.2
Violent & acc. deaths (800-999)	15	6	9	18	7	11	81.8	62.5	101.8
Acc. deaths, total (800-962)	9	4	5	12	5	7	54.5	44.6	64.8
Motor vehicle acc. (810-835)	2	1	1	3	1	2	13.6	8.9	18.5
Ill-defined (780-795)	6	0	6	7	0	7	31.8	-	64.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PANOLA COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		Non-white
				Total	White	Non-white	Total	White	
Live births	851	232	619	900	250	650	28.0	19.4	33.8
Fetal deaths	41	8	33	42	8	34	46.7	32.0	52.3
Deaths from all causes	205	82	123	253	103	150	7.9	8.0	7.8
Deaths under 1 year	40	8	32	45	11	34	50.0	44.0	52.3
Deaths under 1 month	24	8	16	28	11	17	31.1	44.0	26.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	3.1	7.8	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	3.1	7.8	-
Syphilis (020-029)	1	0	1	1	0	1	3.1	-	5.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	1	1	1	0	1	3.1	-	5.2
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	3	0	3	3	0	3	9.3	-	15.6
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	19	9	10	22	9	13	68.5	69.8	67.7
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	32	13	19	39	18	21	121.5	139.5	109.4
Heart diseases (410-443)	72	34	38	85	42	43	264.8	325.6	224.0
Arteriosclerosis (450)	1	0	1	2	1	1	6.2	7.8	5.2
Influenza (480-483)	2	0	2	2	0	2	6.2	-	10.4
#Pneumonia (490-493)	7	5	2	6	4	2	18.7	31.0	10.4
#Diarrhea & enteritis (571)	6	1	5	6	1	5	18.7	7.8	26.0
Acute nephritis (590, 591)	0	0	0	2	2	0	6.2	15.5	-
Chronic nephritis, etc. (592-594)	4	1	3	5	1	4	15.6	7.8	20.8
Maternal deaths (640-689)	1	0	1	3	1	2	3.3	4.0	3.1
Cong. malformations (750-759)	4	3	1	5	4	1	15.6	31.0	5.2
Birth inj., atelect. (760-762)	9	4	5	9	4	5	10.0	16.0	7.7
#Other dis. early inf. (763-776)	6	1	5	8	2	6	8.9	8.0	9.2
Violent & acc. deaths (800-999)	11	7	4	16	5	11	49.8	38.8	57.3
Acc. deaths, total (800-962)	6	3	3	11	2	9	34.3	15.5	46.9
Motor vehicle acc. (810-835)	3	1	2	4	1	3	12.5	7.8	15.6
Ill-defined (780-795)	17	1	16	21	2	19	65.4	15.5	99.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PEARL RIVER COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	692	511	181	639	452	187	29.6	27.4	36.7
Fetal deaths	19	10	9	19	8	11	29.7	17.7	58.8
Deaths from all causes	206	152	54	217	160	57	10.0	9.7	11.2
Deaths under 1 year	26	20	6	28	22	6	43.8	48.7	32.1
Deaths under 1 month	24	20	4	26	22	4	40.7	48.7	21.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	2	2	0	9.2	12.1	-
Tbc., respiratory (001-008)	1	1	0	2	2	0	9.2	12.1	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	4.6	-	19.6
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	0	1	1	0	1	4.6	-	19.6
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	26	21	5	31	25	6	143.5	151.5	117.6
Diabetes (260)	4	3	1	3	2	1	13.9	12.1	19.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	20	13	7	19	12	7	88.0	72.7	137.2
Heart diseases (410-443)	78	63	15	79	63	16	365.7	381.8	313.7
Arteriosclerosis (450)	5	4	1	6	5	1	27.8	30.3	19.6
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	3	1	2	4	2	2	18.5	12.1	39.2
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	5	5	0	7	7	0	32.4	42.4	-
Birth inj., atelect. (760-762)	6	6	0	6	6	0	9.4	13.3	-
#Other dis. early inf. (763-776)	13	9	4	13	9	4	20.3	19.9	21.4
Violent & acc. deaths (800-999)	22	15	7	26	18	8	120.4	109.1	156.9
Acc. deaths, total (800-962)	17	13	4	21	16	5	97.2	97.0	98.0
Motor vehicle acc. (810-835)	7	5	2	9	6	3	41.7	36.4	58.8
Ill-defined (780-795)	8	2	6	6	1	5	27.8	6.1	98.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PERRY COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	104	77	27	222	147	75	30.8	28.3	37.5
Fetal deaths	8	3	5	13	8	5	58.6	54.4	66.7
Deaths from all causes	41	21	20	68	44	24	9.4	8.5	12.0
Deaths under 1 year	5	3	2	11	7	4	49.5	47.6	53.3
Deaths under 1 month	4	2	2	9	5	4	40.5	34.0	53.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	13.9	19.2	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	13.9	19.2	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	0	0	0	2	1	1	27.8	19.2	50.0
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	4	0	4	10	6	4	138.9	115.4	200.0
Heart diseases (410-443)	20	11	9	26	19	7	361.1	365.4	350.0
Arteriosclerosis (450)	0	0	0	1	1	0	13.9	19.2	-
Influenza (480-483)	1	1	0	1	1	0	13.9	19.2	-
#Pneumonia (490-493)	1	1	0	1	1	0	13.9	19.2	-
#Diarrhea & enteritis (571)	0	0	0	1	1	0	13.9	19.2	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	2	0	2	27.8	-	100.0
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	13.9	19.2	-
Birth inj., atelect. (760-762)	0	0	0	3	1	2	13.5	6.8	26.7
#Other dis. early inf. (763-776)	3	1	2	4	2	2	18.0	13.6	26.7
Violent & acc. deaths (800-999)	7	5	2	9	6	3	125.0	115.4	150.0
Acc. deaths, total (800-962)	4	3	1	6	4	2	83.3	76.9	100.0
Motor vehicle acc. (810-835)	0	0	0	2	2	0	27.8	38.5	-
Ill-defined (780-795)	2	1	1	2	1	1	27.8	19.2	50.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PIKE COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,001	479	522	867	356	511	26.9	20.6	34.3
Fetal deaths	31	13	18	24	11	13	27.7	30.9	25.4
Deaths from all causes	340	207	133	342	207	135	10.6	12.0	9.1
Deaths under 1 year	40	15	25	40	17	23	46.1	47.8	45.0
Deaths under 1 month	27	15	12	25	13	12	28.8	36.5	23.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	6	1	5	18.6	5.8	33.6
Tbc., respiratory (001-008)	2	0	2	5	1	4	15.5	5.8	26.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	1	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	1	0	3.1	5.8	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	39	25	14	46	29	17	142.8	167.6	114.1
Diabetes (260)	5	3	2	5	3	2	15.5	17.3	13.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	33	17	16	38	21	17	118.0	121.4	114.1
Heart diseases (410-443)	120	88	32	118	83	35	366.4	479.8	234.9
Arteriosclerosis (450)	5	2	3	4	1	3	12.4	5.8	20.1
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	8	3	5	8	4	4	24.8	23.1	26.8
#Diarrhea & enteritis (571)	4	2	2	5	3	2	15.5	17.3	13.4
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	6.2	5.8	6.7
Maternal deaths (640-689)	3	1	2	3	1	2	3.5	2.8	3.9
Cong. malformations (750-759)	4	2	2	8	4	4	24.8	23.1	26.8
Birth inj., atelect. (760-762)	7	6	1	6	5	1	6.9	14.0	2.0
#Other dis. early inf. (763-776)	13	6	7	11	5	6	12.7	14.0	11.7
Violent & acc. deaths (800-999)	33	25	8	22	17	5	68.3	98.3	33.6
Acc. deaths, total (800-962)	26	20	6	18	13	5	55.9	75.1	33.6
Motor vehicle acc. (810-835)	16	12	4	9	6	3	28.0	34.7	20.1
Ill-defined (780-795)	26	7	19	23	7	16	71.4	40.5	107.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PONTOTOC COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	319	217	102	375	266	109	20.6	18.1	31.1
Fetal deaths	19	10	9	13	6	7	34.7	22.6	64.2
Deaths from all causes	142	117	25	165	133	32	9.1	9.0	9.1
Deaths under 1 year	10	4	6	13	6	7	34.7	22.6	64.2
Deaths under 1 month	7	3	4	9	5	4	24.0	18.8	36.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	5.5	6.8	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	5.5	6.8	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	20	18	2	25	22	3	137.4	149.6	85.7
Diabetes (260)	1	1	0	2	2	0	11.0	13.6	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	13	11	2	14	12	2	76.9	81.6	57.1
Heart diseases (410-443)	68	59	9	70	60	10	384.6	408.2	285.7
Arteriosclerosis (450)	2	2	0	2	2	0	11.0	13.6	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	3	2	1	3	2	1	16.5	13.6	28.6
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	11.0	6.8	28.6
Maternal deaths (640-689)	1	0	1	1	0	1	2.7	-	9.2
Cong. malformations (750-759)	1	0	1	1	0	1	5.5	-	28.6
Birth inj., atelect. (760-762)	2	2	0	2	2	0	5.3	7.5	-
#Other dis. early inf. (763-776)	4	1	3	6	3	3	16.0	11.3	27.5
Violent & acc. deaths (800-999)	12	9	3	16	12	4	87.9	81.6	114.3
Acc. deaths, total (800-962)	8	5	3	13	9	4	71.4	61.2	114.3
Motor vehicle acc. (810-835)	3	3	0	6	5	1	33.0	34.0	28.6
Ill-defined (780-795)	6	4	2	6	4	2	33.0	27.2	57.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PRENTISS COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	487	411	76	410	343	67	22.6	21.6	30.4
Fetal deaths	12	8	4	11	8	3	26.8	23.3	44.8
Deaths from all causes	151	125	26	157	129	28	8.7	8.1	12.7
Deaths under 1 year	12	8	4	11	7	4	26.8	20.4	59.7
Deaths under 1 month	6	4	2	5	3	2	12.2	8.7	29.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	5.5	6.3	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	5.5	6.3	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	15	14	1	17	16	1	93.9	100.6	45.4
Diabetes (260)	3	2	1	4	3	1	22.1	18.9	45.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	12	9	3	15	10	5	82.9	62.9	227.3
Heart diseases (410-443)	53	48	5	53	47	6	292.8	295.6	272.7
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	5	3	2	5	3	2	27.6	18.9	90.9
#Pneumonia (490-493)	6	5	1	5	4	1	27.6	25.2	45.4
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	5.5	6.3	-
Maternal deaths (640-689)	1	0	1	1	0	1	2.4	-	14.9
Cong. malformations (750-759)	2	1	1	3	2	1	16.6	12.6	45.4
Birth inj., atelect. (760-762)	1	0	1	1	0	1	2.4	-	14.9
#Other dis. early inf. (763-776)	3	3	0	2	2	0	4.9	5.8	-
Violent & acc. deaths (800-999)	13	10	3	12	10	2	66.3	62.9	90.9
Acc. deaths, total (800-962)	9	7	2	9	8	1	49.7	50.3	45.4
Motor vehicle acc. (810-835)	6	6	0	7	7	0	38.7	44.0	-
Ill-defined (780-795)	16	12	4	16	12	4	88.4	75.5	181.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
QUITMAN COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	922	251	671	850	225	625	33.3	23.9	38.8
Fetal deaths	29	10	19	23	7	16	27.0	31.1	25.6
Deaths from all causes	182	45	137	217	65	152	8.5	6.9	9.4
Deaths under 1 year	46	11	35	48	12	36	56.5	53.3	57.6
Deaths under 1 month	24	10	14	25	11	14	29.4	48.9	22.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	0	4	6	1	5	23.5	10.6	31.0
Tbc., respiratory (001-008)	4	0	4	5	1	4	19.6	10.6	24.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	2	0	2	2	0	2	7.8	-	12.4
Meningococcal infections (057)	0	0	0	2	1	1	7.8	10.6	6.2
Acute poliomyelitis (080)	0	0	0	1	0	1	3.9	-	6.2
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	5	12	23	8	15	90.2	85.1	93.2
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	2	13	20	4	16	78.4	42.6	99.4
Heart diseases (410-443)	46	16	30	54	21	33	211.8	223.4	205.0
Arteriosclerosis (450)	1	1	0	1	1	0	3.9	10.6	-
Influenza (480-483)	1	0	1	1	0	1	3.9	-	6.2
#Pneumonia (490-493)	9	2	7	9	2	7	35.3	21.3	43.5
#Diarrhea & enteritis (571)	3	0	3	4	0	4	15.7	-	24.8
Acute nephritis (590, 591)	1	1	0	2	2	0	7.8	21.3	-
Chronic nephritis, etc. (592-594)	5	0	5	5	0	5	19.6	-	31.0
Maternal deaths (640-689)	1	0	1	2	0	2	2.4	-	3.2
Cong. malformations (750-759)	5	4	1	4	4	0	15.7	42.6	-
Birth inj., atelect. (760-762)	8	4	4	7	3	4	8.2	13.3	6.4
#Other dis. early inf. (763-776)	7	1	6	8	2	6	9.4	8.9	9.6
Violent & acc. deaths (800-999)	16	4	12	21	9	12	82.4	95.7	74.5
Acc. deaths, total (800-962)	12	4	8	17	9	8	66.7	95.7	49.7
Motor vehicle acc. (810-835)	5	2	3	8	4	4	31.4	42.6	24.8
Ill-defined (780-795)	25	1	24	22	1	21	86.3	10.6	130.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
RANKIN COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	251	24	227	638	295	343	23.3	19.3	28.3
Fetal deaths	12	3	9	30	17	13	47.0	57.6	37.9
Deaths from all causes	417	204	213	233	141	92	8.5	9.2	7.6
Deaths under 1 year	10	1	9	28	14	14	43.9	47.4	40.8
Deaths under 1 month	6	0	6	20	12	8	31.3	40.7	23.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	13	3	10	1	1	0	3.6	6.5	-
Tbc., respiratory (001-008)	12	3	9	1	1	0	3.6	6.5	-
Syphilis (020-029)	4	0	4	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	23	6	17	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	47	25	22	48	26	22	175.2	169.9	181.8
Diabetes (260)	2	1	1	1	1	0	3.6	6.5	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	70	26	44	26	13	13	94.9	85.0	107.4
Heart diseases (410-443)	100	65	35	60	41	19	219.0	268.0	157.0
Arteriosclerosis (450)	3	1	2	0	0	0	-	-	-
Influenza (480-483)	1	0	1	1	0	1	3.6	-	8.3
#Pneumonia (490-493)	23	17	6	7	2	5	25.5	13.1	41.3
#Diarrhea & enteritis (571)	0	0	0	2	1	1	7.3	6.5	8.3
Acute nephritis (590, 591)	1	1	0	2	2	0	7.3	13.1	-
Chronic nephritis, etc. (592-594)	1	0	1	3	3	0	10.9	19.6	-
Maternal deaths (640-689)	1	0	1	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	4	3	1	14.6	19.6	8.3
Birth inj., atelect. (760-762)	0	0	0	4	3	1	6.3	10.2	2.9
#Other dis. early inf. (763-776)	6	0	6	12	6	6	18.8	20.3	17.5
Violent & acc. deaths (800-999)	37	19	18	27	17	10	98.5	111.1	82.6
Acc. deaths, total (800-962)	29	14	15	21	14	7	76.6	91.5	57.8
Motor vehicle acc. (810-835)	12	7	5	12	10	2	43.8	65.4	16.5
Ill-defined (780-795)	24	10	14	14	6	8	51.1	39.2	66.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SCOTT COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	596	299	297	585	257	328	29.4	23.6	36.4
Fetal deaths	25	8	17	30	12	18	51.3	46.7	54.9
Deaths from all causes	144	88	56	160	90	70	8.0	8.2	7.8
Deaths under 1 year	10	5	5	16	8	8	27.4	31.1	24.4
Deaths under 1 month	7	4	3	12	7	5	20.5	27.2	15.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	5.0	-	11.1
Tbc., respiratory (001-008)	1	0	1	1	0	1	5.0	-	11.1
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	5.0	-	11.1
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	1	0	5.0	9.2	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	5.0	-	11.1
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	18	11	7	20	13	7	100.5	119.3	77.8
Diabetes (260)	0	0	0	1	1	0	5.0	9.2	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	20	9	11	21	9	12	105.5	82.6	133.3
Heart diseases (410-443)	44	33	11	49	30	19	246.2	275.2	211.1
Arteriosclerosis (450)	3	1	2	3	1	2	15.1	9.2	22.2
Influenza (480-483)	1	0	1	1	0	1	5.0	-	11.1
#Pneumonia (490-493)	1	1	0	0	0	0	-	-	-
#Diarrhea & enteritis (571)	3	1	2	3	1	2	15.1	9.2	22.2
Acute nephritis (590, 591)	1	1	0	1	1	0	5.0	9.2	-
Chronic nephritis, etc. (592-594)	8	7	1	6	5	1	30.2	45.9	11.1
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	2	0	2	10.0	-	22.2
Birth inj., atelect. (760-762)	3	3	0	5	5	0	8.5	19.4	-
#Other dis. early inf. (763-776)	3	1	2	6	2	4	10.2	7.8	12.2
Violent & acc. deaths (800-999)	11	6	5	9	5	4	45.2	45.9	44.4
Acc. deaths, total (800-962)	6	5	1	5	4	1	25.1	36.7	11.1
Motor vehicle acc. (810-835)	4	3	1	4	3	1	20.1	27.5	11.1
Ill-defined (780-795)	8	1	7	8	1	7	40.2	9.2	77.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SHARKEY COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	274	53	221	422	91	331	33.8	26.8	36.4
Fetal deaths	10	0	10	16	1	15	37.9	11.0	45.3
Deaths from all causes	98	24	74	112	25	87	9.0	7.4	9.6
Deaths under 1 year	17	1	16	17	1	16	40.3	11.0	48.3
Deaths under 1 month	9	1	8	11	1	10	26.1	11.0	30.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	2	2	0	16.0	58.8	-
Tbc., respiratory (001-008)	1	1	0	2	2	0	16.0	58.8	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	8.0	-	11.0
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	3	7	10	2	8	80.0	58.8	87.9
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	13	3	10	14	5	9	112.0	147.0	98.9
Heart diseases (410-443)	31	9	22	34	10	24	272.0	294.1	263.7
Arteriosclerosis (450)	1	0	1	1	0	1	8.0	-	11.0
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	3	0	3	3	0	3	24.0	-	33.0
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	1	0	1	8.0	-	11.0
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	16.0	29.4	11.0
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	1	0	1	8.0	-	11.0
Birth inj., atelect. (760-762)	3	1	2	3	1	2	7.1	11.0	6.0
#Other dis. early inf. (763-776)	7	0	7	9	0	9	21.3	-	27.2
Violent & acc. deaths (800-999)	14	6	8	15	4	11	120.0	117.6	120.9
Acc. deaths, total (800-962)	12	4	8	12	3	9	96.0	88.2	98.9
Motor vehicle acc. (810-835)	6	3	3	8	3	5	64.0	88.2	54.9
Ill-defined (780-795)	8	0	8	8	0	8	64.0	-	87.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SIMPSON COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	555	244	311	513	250	263	24.3	19.2	32.5
Fetal deaths	10	2	8	16	8	8	31.2	32.0	30.4
Deaths from all causes	158	89	69	149	83	66	7.1	6.4	8.1
Deaths under 1 year	17	4	13	23	6	17	44.8	24.0	64.6
Deaths under 1 month	8	2	6	11	4	7	21.4	16.0	26.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	32	15	17	1	0	1	4.7	-	12.3
Tbc., respiratory (001-008)	31	15	16	1	0	1	4.7	-	12.3
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	1	1	0	4.7	7.7	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	1	0	1	4.7	-	12.3
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	14	11	3	17	12	5	80.6	92.3	61.7
Diabetes (260)	0	0	0	1	1	0	4.7	7.7	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	18	11	7	19	11	8	90.0	84.6	98.8
Heart diseases (410-443)	46	31	15	47	29	18	222.7	223.1	222.2
Arteriosclerosis (450)	1	0	1	2	1	1	9.5	7.7	12.3
Influenza (480-483)	2	0	2	1	0	1	4.7	-	12.3
#Pneumonia (490-493)	5	2	3	4	2	2	19.0	15.4	24.7
#Diarrhea & enteritis (571)	1	0	1	1	0	1	4.7	-	12.3
Acute nephritis (590, 591)	1	1	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	4	1	3	19.0	7.7	37.0
Birth inj., atelect. (760-762)	2	1	1	3	2	1	5.8	8.0	3.8
#Other dis. early inf. (763-776)	6	1	5	7	2	5	13.6	8.0	19.0
Violent & acc. deaths (800-999)	14	8	6	22	14	8	104.3	107.7	98.8
Acc. deaths, total (800-962)	11	6	5	18	11	7	85.3	84.6	86.4
Motor vehicle acc. (810-835)	1	0	1	4	2	2	19.0	15.4	24.7
Ill-defined (780-795)	4	0	4	4	0	4	19.0	-	49.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SMITH COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	207	141	66	390	250	140	28.5	22.5	53.8
Fetal deaths	3	2	1	9	3	6	23.1	12.0	42.8
Deaths from all causes	66	48	18	118	89	29	8.6	8.0	11.2
Deaths under 1 year	4	2	2	10	5	5	25.6	20.0	35.7
Deaths under 1 month	1	1	0	3	3	0	7.7	12.0	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	7.3	9.0	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	7.3	9.0	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	6	4	16	11	5	116.8	99.1	192.3
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	11	10	1	19	16	3	138.7	144.1	115.4
Heart diseases (410-443)	26	22	4	43	38	5	313.9	342.3	192.3
Arteriosclerosis (450)	1	1	0	1	1	0	7.3	9.0	-
Influenza (480-483)	0	0	0	1	0	1	7.3	-	38.5
#Pneumonia (490-493)	3	2	1	4	2	2	29.2	18.0	76.9
#Diarrhea & enteritis (571)	0	0	0	2	1	1	14.6	9.0	38.5
Acute nephritis (590, 591)	1	0	1	2	1	1	14.6	9.0	38.5
Chronic nephritis, etc. (592-594)	3	3	0	5	5	0	36.5	45.0	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	1	1	2	1	1	14.6	9.0	38.5
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	1	1	0	3	3	0	7.7	12.0	-
Violent & acc. deaths (800-999)	4	2	2	10	5	5	73.0	45.0	192.3
Acc. deaths, total (800-962)	3	2	1	9	5	4	65.7	45.0	153.8
Motor vehicle acc. (810-835)	1	1	0	5	2	3	36.5	18.0	115.4
Ill-defined (780-795)	3	0	3	4	0	4	29.2	-	153.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
STONE COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	188	133	55	171	117	54	26.7	23.9	36.0
Fetal deaths	3	2	1	3	2	1	17.5	17.1	18.5
Deaths from all causes	48	36	12	51	39	12	8.0	8.0	8.0
Deaths under 1 year	2	1	1	2	1	1	11.7	8.5	18.5
Deaths under 1 month	2	1	1	2	1	1	11.7	8.5	18.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	15.6	20.4	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	15.6	20.4	-
Syphilis (020-029)	0	0	0	1	1	0	15.6	20.4	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	4	3	1	7	5	2	109.4	102.0	133.3
Diabetes (260)	1	1	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	3	3	0	3	3	0	46.9	61.2	-
Heart diseases (410-443)	24	19	5	25	19	6	390.6	387.8	400.0
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	0	0	0	0	0	0	-	-	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	2	1	1	2	1	1	11.7	8.5	18.5
#Other dis. early inf. (763-776)	0	0	0	0	0	0	-	-	-
Violent & acc. deaths (800-999)	10	6	4	5	3	2	78.1	61.2	133.3
Acc. deaths, total (800-962)	10	6	4	5	3	2	78.1	61.2	133.3
Motor vehicle acc. (810-835)	4	4	0	2	2	0	31.2	40.8	-
Ill-defined (780-795)	1	0	1	1	0	1	15.6	-	66.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SUNFLOWER COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,538	316	1,222	1,614	356	1,258	29.8	21.8	33.3
Fetal deaths	53	13	40	54	14	40	33.4	39.3	31.8
Deaths from all causes	395	100	295	433	118	315	8.0	7.2	8.3
Deaths under 1 year	67	8	59	74	10	64	45.8	28.1	50.9
Deaths under 1 month	39	5	34	42	7	35	26.0	19.7	27.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	0	3	5	0	5	9.2	-	13.2
Tbc., respiratory (001-008)	3	0	3	4	0	4	7.4	-	10.6
Syphilis (020-029)	1	0	1	1	0	1	1.8	-	2.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	1.8	-	2.6
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	2	0	2	3.7	-	5.3
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	41	18	23	47	19	28	86.9	116.6	74.1
Diabetes (260)	7	2	5	10	2	8	18.5	12.3	21.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	58	8	50	61	13	48	112.8	79.8	127.0
Heart diseases (410-443)	79	34	45	93	40	53	171.9	245.4	140.2
Arteriosclerosis (450)	3	1	2	3	0	3	5.5	-	7.9
Influenza (480-483)	3	1	2	2	1	1	3.7	6.1	2.6
#Pneumonia (490-493)	14	4	10	13	3	10	24.0	18.4	26.4
#Diarrhea & enteritis (571)	5	0	5	6	0	6	11.1	-	15.9
Acute nephritis (590, 591)	2	0	2	1	0	1	1.8	-	2.6
Chronic nephritis, etc. (592-594)	8	2	6	9	1	8	16.6	6.1	21.2
Maternal deaths (640-689)	1	0	1	1	0	1	0.6	-	0.8
Cong. malformations (750-759)	1	0	1	4	3	1	7.4	18.4	2.6
Birth inj., atelect. (760-762)	6	2	4	5	2	3	3.1	5.6	2.4
#Other dis. early inf. (763-776)	29	2	27	33	2	31	20.4	5.6	24.6
Violent & acc. deaths (800-999)	38	11	27	32	13	19	59.1	79.8	50.3
Acc. deaths, total (800-962)	22	8	14	25	11	14	46.2	67.5	37.0
Motor vehicle acc. (810-835)	9	2	7	12	5	7	22.2	30.7	18.5
Ill-defined (780-795)	58	6	52	58	6	52	107.2	36.8	137.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TALLAHATCHIE COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	838	139	699	920	207	713	32.4	20.7	38.8
Fetal deaths	30	4	26	35	7	28	38.0	33.8	39.3
Deaths from all causes	253	68	185	292	96	196	10.3	9.6	10.6
Deaths under 1 year	48	8	40	50	10	40	54.3	48.3	56.1
Deaths under 1 month	28	7	21	32	9	23	34.8	43.5	32.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	0	4	8	1	7	28.2	10.0	38.0
Tbc., respiratory (001-008)	3	0	3	7	1	6	24.6	10.0	32.6
Syphilis (020-029)	1	0	1	1	0	1	3.5	-	5.4
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	1	1	1	1	0	3.5	10.0	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	3.5	-	5.4
Meningococcal infections (057)	1	1	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	3.5	-	5.4
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	21	9	12	32	18	14	112.7	180.0	76.1
Diabetes (260)	2	1	1	2	1	1	7.0	10.0	5.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	26	8	18	32	13	19	112.7	130.0	103.3
Heart diseases (410-443)	67	29	38	76	35	41	267.6	350.0	222.8
Arteriosclerosis (450)	4	1	3	5	2	3	17.6	20.0	16.3
Influenza (480-483)	2	0	2	2	0	2	7.0	-	10.9
#Pneumonia (490-493)	7	0	7	8	1	7	28.2	10.0	38.0
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	0	1	1	0	1	3.5	-	5.4
Chronic nephritis, etc. (592-594)	4	0	4	4	0	4	14.1	-	21.7
Maternal deaths (640-689)	6	1	5	5	1	4	5.4	4.8	5.6
Cong. malformations (750-759)	2	1	1	2	1	1	7.0	10.0	5.4
Birth inj., atelect. (760-762)	11	5	6	14	7	7	15.2	33.8	9.8
#Other dis. early inf. (763-776)	13	0	13	14	1	13	15.2	4.8	18.2
Violent & acc. deaths (800-999)	24	7	17	23	7	16	81.0	70.0	87.0
Acc. deaths, total (800-962)	20	6	14	19	6	13	66.9	60.0	70.6
Motor vehicle acc. (810-835)	9	3	6	9	4	5	31.7	40.0	27.2
Ill-defined (780-795)	33	1	32	35	2	33	123.2	20.0	179.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TATE COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	462	22	440	582	114	468	33.8	16.8	45.0
Fetal deaths	21	1	20	29	2	27	49.8	17.5	57.7
Deaths from all causes	124	45	79	161	68	93	9.4	10.0	8.9
Deaths under 1 year	15	0	15	23	2	21	39.5	17.5	44.9
Deaths under 1 month	10	0	10	14	2	12	24.0	17.5	25.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	1	0	5.8	14.7	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	14	3	11	17	7	10	98.8	102.9	96.2
Diabetes (260)	1	1	0	1	1	0	5.8	14.7	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	14	4	10	18	7	11	104.6	102.9	105.8
Heart diseases (410-443)	48	25	23	56	33	23	325.6	485.3	221.2
Arteriosclerosis (450)	1	1	0	1	1	0	5.8	14.7	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	3	0	3	4	0	4	23.2	-	38.5
#Diarrhea & enteritis (571)	3	0	3	4	0	4	23.2	-	38.5
Acute nephritis (590, 591)	1	0	1	1	0	1	5.8	-	9.6
Chronic nephritis, etc. (592-594)	2	0	2	4	1	3	23.2	14.7	28.8
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	0	1	5.8	-	9.6
Birth inj., atelect. (760-762)	1	0	1	3	2	1	5.2	17.5	2.1
#Other dis. early inf. (763-776)	8	0	8	11	0	11	18.9	-	23.5
Violent & acc. deaths (800-999)	7	2	5	12	3	9	69.8	44.1	86.5
Acc. deaths, total (800-962)	5	1	4	9	2	7	52.3	29.4	67.3
Motor vehicle acc. (810-835)	1	0	1	3	1	2	17.4	14.7	19.2
Ill-defined (780-795)	14	2	12	15	2	13	87.2	29.4	125.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TIPPAH COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	336	218	118	331	215	116	19.9	16.7	31.4
Fetal deaths	30	22	8	23	15	8	69.5	69.8	69.0
Deaths from all causes	114	87	27	124	96	28	7.5	7.4	7.6
Deaths under 1 year	12	6	6	8	3	5	24.2	14.0	43.1
Deaths under 1 month	8	6	2	5	3	2	15.1	14.0	17.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	6.0	-	27.0
Tbc., respiratory (001-008)	1	0	1	1	0	1	6.0	-	27.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	19	16	3	21	18	3	126.5	139.5	81.1
Diabetes (260)	1	1	0	1	1	0	6.0	7.8	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	9	7	2	11	9	2	66.3	69.8	54.0
Heart diseases (410-443)	55	44	11	59	49	10	355.4	379.8	270.3
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	2	0	2	2	0	12.0	15.5	-
#Diarrhea & enteritis (571)	1	0	1	1	0	1	6.0	-	27.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	2	1	4	2	2	24.1	15.5	54.0
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	1	1	0	6.0	7.8	-
Birth inj., atelect. (760-762)	6	4	2	3	1	2	9.1	4.6	17.2
#Other dis. early inf. (763-776)	2	1	1	1	1	0	3.0	4.6	-
Violent & acc. deaths (800-999)	6	3	3	9	4	5	54.2	31.0	135.1
Acc. deaths, total (800-962)	6	3	3	9	4	5	54.2	31.0	135.1
Motor vehicle acc. (810-835)	3	0	3	4	1	3	24.1	7.8	81.1
Ill-defined (780-795)	3	2	1	3	2	1	18.1	15.5	27.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TISHOMINGO COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	156	143	13	280	262	18	21.5	21.3	25.7
Fetal deaths	1	1	0	3	2	1	10.7	7.6	55.6
Deaths from all causes	92	88	4	117	112	5	9.0	9.1	7.1
Deaths under 1 year	3	2	1	6	5	1	21.4	19.1	55.6
Deaths under 1 month	1	1	0	2	2	0	7.1	7.6	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	7.7	-	142.8
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	6	6	0	11	11	0	84.6	89.4	-
Diabetes (260)	1	1	0	1	1	0	7.7	8.1	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	24	21	3	25	22	3	192.3	178.9	428.6
Heart diseases (410-443)	27	27	0	34	34	0	261.5	276.4	-
Arteriosclerosis (450)	1	1	0	1	1	0	7.7	8.1	-
Influenza (480-483)	1	1	0	1	1	0	7.7	8.1	-
#Pneumonia (490-493)	4	4	0	6	6	0	46.2	48.8	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	2	2	0	2	2	0	15.4	16.3	-
Chronic nephritis, etc. (592-594)	3	3	0	4	3	1	30.8	24.4	142.8
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	1	1	0	2	2	0	7.1	7.6	-
#Other dis. early inf. (763-776)	1	1	0	1	1	0	3.6	3.8	-
Violent & acc. deaths (800-999)	7	7	0	11	11	0	84.6	89.4	-
Acc. deaths, total (800-962)	6	6	0	10	10	0	76.9	81.3	-
Motor vehicle acc. (810-835)	3	3	0	3	3	0	23.1	24.4	-
Ill-defined (780-795)	3	3	0	3	3	0	23.1	24.4	-

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TUNICA COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	654	82	572	674	89	585	29.7	20.2	32.0
Fetal deaths	27	4	23	30	4	26	44.5	44.9	44.4
Deaths from all causes	221	33	188	231	34	197	10.2	7.7	10.8
Deaths under 1 year	44	2	42	44	2	42	65.3	22.5	71.8
Deaths under 1 month	21	2	19	20	2	18	29.7	22.5	30.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	2	0	2	8.8	-	10.9
Tbc., respiratory (001-008)	1	0	1	1	0	1	4.4	-	5.5
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	3	0	3	3	0	3	13.2	-	16.4
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	2	0	2	2	0	2	8.8	-	10.9
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	4.4	-	5.5
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	21	3	18	28	5	23	123.3	113.6	125.7
Diabetes (260)	1	0	1	1	0	1	4.4	-	5.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	23	3	20	25	3	22	110.1	68.2	120.2
Heart diseases (410-443)	53	15	38	55	15	40	242.3	340.9	218.6
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	1	0	1	1	0	1	4.4	-	5.5
#Pneumonia (490-493)	7	0	7	9	1	8	39.6	22.7	43.7
#Diarrhea & enteritis (571)	6	0	6	6	0	6	26.4	-	32.8
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc.(592-594)	10	2	8	9	1	8	39.6	22.7	43.7
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	3	0	3	2	0	2	8.8	-	10.9
Birth inj., atelect. (760-762)	4	2	2	4	2	2	5.9	22.5	3.4
#Other dis. early inf. (763-776)	11	0	11	11	0	11	16.3	-	18.8
Violent & acc. deaths (800-999)	34	5	29	27	3	24	118.9	68.2	131.1
Acc. deaths, total (800-962)	26	4	22	18	2	16	79.3	45.4	87.4
Motor vehicle acc. (810-835)	15	3	12	7	0	7	30.8	-	38.2
Ill-defined (780-795)	27	1	26	26	0	26	114.5	-	142.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
UNION COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	538	410	128	434	314	120	23.1	21.2	30.0
Fetal deaths	37	31	6	34	29	5	78.3	92.4	41.7
Deaths from all causes	195	149	46	204	159	45	10.8	10.7	11.2
Deaths under 1 year	21	10	11	21	11	10	48.4	35.0	83.3
Deaths under 1 month	15	9	6	15	10	5	34.6	31.8	41.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	2	2	0	10.6	13.5	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	5.3	6.8	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	0	1	1	0	1	5.3	-	25.0
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	15	2	21	18	3	111.7	121.6	75.0
Diabetes (260)	1	1	0	1	1	0	5.3	6.8	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	44	36	8	41	34	7	218.1	229.7	17.5
Heart diseases (410-443)	61	51	10	65	54	11	345.7	364.9	275.0
Arteriosclerosis (450)	0	0	0	1	1	0	5.3	6.8	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	0	0	0	0	0	0	-	-	-
#Diarrhea & enteritis (571)	3	2	1	3	2	1	16.0	13.5	25.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	6	3	3	6	3	3	31.9	20.3	75.0
Maternal deaths (640-689)	1	1	0	1	1	0	2.3	3.2	-
Cong. malformations (750-759)	1	0	1	1	0	1	5.3	-	25.0
Birth inj., atelect. (760-762)	4	4	0	4	4	0	9.2	12.7	-
#Other dis. early inf. (763-776)	11	5	6	11	6	5	25.3	19.1	41.7
Violent & acc. deaths (800-999)	6	6	0	11	8	3	58.5	54.0	75.0
Acc. deaths, total (800-962)	5	5	0	10	7	3	53.2	47.3	75.0
Motor vehicle acc. (810-835)	4	4	0	8	5	3	42.6	33.8	75.0
Ill-defined (780-795)	11	5	6	11	5	6	58.5	33.8	150.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WALTHALL COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	644	355	289	383	153	230	26.8	19.4	35.9
Fetal deaths	25	11	14	15	5	10	39.2	32.7	43.5
Deaths from all causes	118	67	51	114	65	49	8.0	8.2	7.6
Deaths under 1 year	17	6	11	13	2	11	33.9	13.1	47.8
Deaths under 1 month	10	4	6	7	1	6	18.3	6.5	26.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	13	9	4	15	11	4	104.9	139.2	62.5
Diabetes (260)	6	4	2	5	2	3	35.0	25.3	46.9
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	13	8	5	16	10	6	111.9	126.6	93.8
Heart diseases (410-443)	34	25	9	34	24	10	237.8	303.8	156.2
Arteriosclerosis (450)	1	0	1	1	0	1	7.0	-	15.6
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	5	0	5	6	1	5	42.0	12.6	78.1
#Diarrhea & enteritis (571)	1	0	1	1	0	1	7.0	-	15.6
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	4	3	1	2	1	1	14.0	12.6	15.6
Birth inj., atelect. (760-762)	3	2	1	2	1	1	5.2	6.5	4.3
#Other dis. early inf. (763-776)	5	1	4	4	0	4	10.4	-	17.4
Violent & acc. deaths (800-999)	14	6	8	13	7	6	90.9	88.6	93.8
Acc. deaths, total (800-962)	13	5	8	11	5	6	76.9	63.3	93.8
Motor vehicle acc. (810-835)	6	3	3	5	4	1	35.0	50.6	15.6
Ill-defined (780-795)	8	3	5	7	3	4	49.0	38.0	62.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WARREN COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,476	685	791	1,172	538	634	29.7	23.6	38.0
Fetal deaths	79	25	54	44	13	31	37.5	24.2	48.9
Deaths from all causes	635	253	382	477	201	276	12.1	8.8	16.5
Deaths under 1 year	71	17	54	49	13	36	41.8	24.2	56.8
Deaths under 1 month	44	14	30	32	12	20	27.3	22.3	31.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	1	3	4	1	3	10.1	4.4	18.0
Tbc., respiratory (001-008)	2	1	1	4	1	3	10.1	4.4	18.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	1	1	1	0	1	2.5	-	6.0
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	2.5	-	6.0
Meningococcal infections (057)	3	1	2	1	0	1	2.5	-	6.0
Acute poliomyelitis (080)	4	1	3	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	97	51	46	66	41	25	167.1	179.8	149.7
Diabetes (260)	10	1	9	2	1	1	5.1	4.4	6.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	77	31	46	68	26	42	172.2	114.0	251.5
Heart diseases (410-443)	171	87	84	145	76	69	367.1	333.3	413.2
Arteriosclerosis (450)	7	4	3	4	3	1	10.1	13.2	6.0
Influenza (480-483)	6	0	6	6	0	6	15.2	-	35.9
#Pneumonia (490-493)	14	2	12	10	2	8	25.3	8.8	47.9
#Diarrhea & enteritis (571)	4	0	4	2	0	2	5.1	-	12.0
Acute nephritis (590, 591)	4	1	3	1	0	1	2.5	-	6.0
Chronic nephritis, etc. (592-594)	10	3	7	9	3	6	22.8	13.2	35.9
Maternal deaths (640-689)	1	0	1	0	0	0	-	-	-
Cong. malformations (750-759)	9	4	5	1	1	0	2.5	4.4	-
Birth inj., atelect. (760-762)	12	5	7	10	5	5	8.5	9.3	7.9
#Other dis. early inf. (763-776)	28	7	21	19	5	14	16.2	9.3	22.1
Violent & acc. deaths (800-999)	41	19	22	24	14	10	60.8	61.4	59.9
Acc. deaths, total (800-962)	36	16	20	21	12	9	53.2	52.6	53.9
Motor vehicle acc. (810-835)	11	7	4	7	5	2	17.7	21.9	12.0
Ill-defined (780-795)	62	8	54	61	8	53	154.4	35.1	317.4

* Crude birth and death rate per 1,000 population. Rate for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WASHINGTON COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	3,032	1,229	1,803	2,586	881	1,705	33.8	28.3	37.6
Fetal deaths	203	100	103	193	92	101	74.6	104.4	59.2
Deaths from all causes	742	225	517	781	215	566	10.2	6.9	12.5
Deaths under 1 year	130	32	98	136	31	105	52.6	35.2	61.6
Deaths under 1 month	70	30	40	70	28	42	27.1	31.8	24.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	8	1	7	10	0	10	13.1	-	22.0
Tbc., respiratory (001-008)	5	1	4	8	0	8	10.4	-	17.6
Syphilis (020-029)	2	0	2	2	0	2	2.6	-	4.4
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	5	0	5	6	0	6	7.8	-	13.2
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	2	1	1	2	1	1	2.6	3.2	2.2
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	88	34	54	88	35	53	115.0	112.5	116.7
Diabetes (260)	5	1	4	8	0	8	10.4	-	17.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	80	22	58	92	22	70	120.3	70.7	154.2
Heart diseases (410-443)	214	72	142	220	67	153	287.6	215.4	337.0
Arteriosclerosis (450)	1	1	0	4	3	1	5.2	9.6	2.2
Influenza (480-483)	7	0	7	7	0	7	9.2	-	15.4
#Pneumonia (490-493)	23	6	17	24	5	19	31.4	16.1	41.8
#Diarrhea & enteritis (571)	12	1	11	14	1	13	18.3	3.2	28.6
Acute nephritis (590, 591)	0	0	0	3	0	3	3.9	-	6.6
Chronic nephritis, etc. (592-594)	6	3	3	9	4	5	11.8	12.9	11.0
Maternal deaths (640-689)	2	0	2	1	0	1	0.4	-	0.6
Cong. malformations (750-759)	6	3	3	7	3	4	9.2	9.6	8.8
Birth inj., atelect. (760-762)	17	12	5	17	11	6	6.6	12.5	3.5
#Other dis. early inf. (763-776)	42	12	30	43	12	31	16.6	13.6	18.2
Violent & acc. deaths (800-999)	66	22	44	69	22	47	90.2	70.7	103.5
Acc. deaths, total (800-962)	54	18	36	57	19	38	74.5	61.1	83.7
Motor vehicle acc. (810-835)	22	11	11	19	7	12	24.8	22.5	26.4
Ill-defined (780-795)	93	9	84	92	8	84	120.3	25.7	185.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WAYNE COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	470	285	185	470	254	216	30.3	26.2	37.2
Fetal deaths	24	12	12	24	13	11	51.1	51.2	50.9
Deaths from all causes	118	62	56	128	67	61	8.2	6.9	10.5
Deaths under 1 year	12	4	8	15	6	9	31.9	23.6	41.7
Deaths under 1 month	6	4	2	9	6	3	19.1	23.6	13.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	1	0	1	6.4	-	17.2
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	11	6	5	14	8	6	90.3	82.5	103.4
Diabetes (260)	1	0	1	1	0	1	6.4	-	17.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	14	5	9	15	6	9	96.8	61.8	155.2
Heart diseases (410-443)	37	24	13	38	25	13	245.2	257.7	224.1
Arteriosclerosis (450)	3	1	2	3	1	2	19.4	10.3	34.5
Influenza (480-483)	1	0	1	1	0	1	6.4	-	17.2
#Pneumonia (490-493)	7	3	4	6	3	3	38.7	30.9	51.7
#Diarrhea & enteritis (571)	1	0	1	1	0	1	6.4	-	17.2
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	1	3	4	1	3	25.8	10.3	51.7
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	2	0	2	12.9	-	34.5
Birth inj., atelect. (760-762)	4	3	1	6	4	2	12.8	15.7	9.2
#Other dis. early inf. (763-776)	2	1	1	3	2	1	6.4	7.9	4.6
Violent & acc. deaths (800-999)	14	12	2	12	9	3	77.4	92.8	51.7
Acc. deaths, total (800-962)	11	9	2	8	6	2	51.6	61.8	34.5
Motor vehicle acc. (810-835)	8	7	1	4	3	1	25.8	30.9	17.2
Ill-defined (780-795)	6	1	5	7	1	6	45.2	10.3	103.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WEBSTER COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	185	93	92	226	125	101	21.5	15.6	40.4
Fetal deaths	7	2	5	7	1	6	31.0	8.0	59.4
Deaths from all causes	83	70	13	96	80	16	9.1	10.0	6.4
Deaths under 1 year	5	4	1	9	6	3	39.8	48.0	29.7
Deaths under 1 month	5	4	1	7	6	1	31.0	48.0	9.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	2	1	1	19.0	12.5	40.0
Tbc., respiratory (001-008)	2	1	1	2	1	1	19.0	12.5	40.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	8	2	16	14	2	152.4	175.0	80.0
Diabetes (260)	1	1	0	1	1	0	9.5	12.5	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	22	20	2	18	16	2	171.4	200.0	80.0
Heart diseases (410-443)	16	15	1	20	18	2	190.5	225.0	80.0
Arteriosclerosis (450)	3	3	0	1	1	0	9.5	12.5	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	1	0	4	2	2	38.1	25.0	80.0
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	2	1	1	19.0	12.5	40.0
Maternal deaths (640-689)	0	0	0	1	0	1	4.4	-	9.9
Cong. malformations (750-759)	3	3	0	3	3	0	28.6	37.5	-
Birth inj., atelect. (760-762)	1	1	0	3	3	0	13.3	24.0	-
#Other dis. early inf. (763-776)	1	1	0	1	1	0	4.4	8.0	-
Violent & acc. deaths (800-999)	11	9	2	10	9	1	95.2	112.5	40.0
Acc. deaths, total (800-962)	9	8	1	9	8	1	85.7	100.0	40.0
Motor vehicle acc. (810-835)	5	5	0	3	3	0	28.6	37.5	-
Ill-defined (780-795)	6	3	3	6	3	3	57.1	37.5	120.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WILKINSON COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	493	159	334	436	74	362	33.0	21.8	36.9
Fetal deaths	27	10	17	19	4	15	43.6	54.0	41.4
Deaths from all causes	132	53	79	137	46	91	10.4	13.5	9.3
Deaths under 1 year	15	4	11	16	2	14	36.7	27.0	38.7
Deaths under 1 month	10	3	7	11	1	10	25.2	13.5	27.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	1	0	1	7.6	-	10.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	11	5	6	14	6	8	106.1	176.5	81.6
Diabetes (260)	3	2	1	2	1	1	15.2	29.4	10.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	6	9	12	2	10	90.9	58.8	102.0
Heart diseases (410-443)	47	26	21	46	23	23	348.5	676.5	234.7
Arteriosclerosis (450)	1	0	1	1	0	1	7.6	-	10.2
Influenza (480-483)	1	0	1	1	0	1	7.6	-	10.2
#Pneumonia (490-493)	5	2	3	5	2	3	37.9	58.8	30.6
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	1	0	1	2	0	2	4.6	-	5.5
#Other dis. early inf. (763-776)	4	3	1	4	1	3	9.2	13.5	8.3
Violent & acc. deaths (800-999)	9	5	4	11	5	6	83.3	147.0	61.2
Acc. deaths, total (800-962)	6	4	2	8	4	4	60.6	117.6	40.8
Motor vehicle acc. (810-835)	3	2	1	3	2	1	22.7	58.8	10.2
Ill-defined (780-795)	28	2	26	27	2	25	204.5	58.8	255.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WINSTON COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	456	178	278	533	227	306	27.3	20.4	36.4
Fetal deaths	14	2	12	17	2	15	31.9	8.8	49.0
Deaths from all causes	133	79	54	170	107	63	8.7	9.6	7.5
Deaths under 1 year	14	4	10	14	5	9	26.3	22.0	29.4
Deaths under 1 month	7	3	4	8	4	4	15.0	17.6	13.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	1	2	4	2	2	20.5	18.0	23.8
Tbc., respiratory (001-008)	3	1	2	4	2	2	20.5	18.0	23.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	16	11	5	25	18	7	128.2	162.2	83.3
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	17	10	7	21	12	9	107.7	108.1	107.1
Heart diseases (410-443)	35	28	7	41	33	8	210.2	297.3	95.2
Arteriosclerosis (450)	1	0	1	1	0	1	5.1	-	11.9
Influenza (480-483)	4	4	0	4	4	0	20.5	36.0	-
#Pneumonia (490-493)	8	0	8	8	0	8	41.0	-	95.2
#Diarrhea & enteritis (571)	2	0	2	2	0	2	10.2	-	23.8
Acute nephritis (590, 591)	1	0	1	1	0	1	5.1	-	11.9
Chronic nephritis, etc. (592-594)	2	1	1	3	2	1	15.4	18.0	11.9
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	2	0	2	10.2	-	23.8
Birth inj., atelect. (760-762)	2	2	0	2	2	0	3.8	8.8	-
#Other dis. early inf. (763-776)	5	1	4	6	2	4	11.2	8.8	13.1
Violent & acc. deaths (800-999)	11	9	2	19	15	4	97.4	135.1	47.6
Acc. deaths, total (800-962)	8	7	1	14	12	2	71.8	108.1	23.8
Motor vehicle acc. (810-835)	1	1	0	6	6	0	30.8	54.0	-
Ill-defined (780-795)	12	3	9	12	4	8	61.5	36.0	95.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
YALOBUSHA COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	309	95	214	327	119	208	25.2	17.5	33.5
Fetal deaths	5	0	5	7	1	6	21.4	8.4	28.8
Deaths from all causes	127	76	51	158	97	61	12.2	14.3	9.8
Deaths under 1 year	3	0	3	7	3	4	21.4	25.2	19.2
Deaths under 1 month	0	0	0	4	3	1	12.2	25.2	4.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	7.7	14.7	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	7.7	14.7	-
Syphilis (020-029)	1	0	1	1	0	1	7.7	-	16.1
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	1	1	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	15	9	6	18	11	7	138.5	161.8	112.9
Diabetes (260)	1	0	1	1	0	1	7.7	-	16.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	17	10	7	22	12	10	169.2	176.5	161.3
Heart diseases (410-443)	41	27	14	53	36	17	407.7	529.4	274.2
Arteriosclerosis (450)	1	0	1	1	0	1	7.7	-	16.1
Influenza (480-483)	1	1	0	1	1	0	7.7	14.7	-
#Pneumonia (490-493)	4	2	2	5	3	2	38.5	44.1	32.2
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	9	6	3	10	7	3	76.9	102.9	48.4
Maternal deaths (640-689)	1	0	1	1	0	1	3.0	-	4.8
Cong. malformations (750-759)	1	0	1	1	0	1	7.7	-	16.1
Birth inj., atelect. (760-762)	0	0	0	1	1	0	3.0	8.4	-
#Other dis. early inf. (763-776)	0	0	0	3	2	1	9.2	16.8	4.8
Violent & acc. deaths (800-999)	13	9	4	15	11	4	115.4	161.8	64.5
Acc. deaths, total (800-962)	12	8	4	14	10	4	107.7	147.0	64.5
Motor vehicle acc. (810-835)	5	3	2	5	4	1	38.5	58.8	16.1
Ill-defined (780-795)	6	1	5	6	1	5	46.2	14.7	80.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
YAZOO COUNTY, MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,049	265	784	1,091	313	778	30.3	22.7	35.0
Fetal deaths	41	6	35	44	11	33	40.3	35.1	42.4
Deaths from all causes	310	108	202	344	128	216	9.6	9.3	9.7
Deaths under 1 year	31	6	25	38	8	30	34.8	25.6	38.6
Deaths under 1 month	20	6	14	22	6	16	20.2	19.2	20.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	2	0	2	5.6	-	9.0
Tbc., respiratory (001-008)	1	0	1	2	0	2	5.6	-	9.0
Syphilis (020-029)	0	0	0	1	0	1	2.8	-	4.5
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	45	15	30	55	23	32	152.8	166.7	144.1
Diabetes (260)	5	1	4	5	1	4	13.9	7.2	18.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	40	14	26	42	14	28	116.7	101.4	126.1
Heart diseases (410-443)	73	36	37	77	39	38	213.9	282.6	171.2
Arteriosclerosis (450)	3	2	1	3	2	1	8.3	14.5	4.5
Influenza (480-483)	3	0	3	4	1	3	11.1	7.2	13.5
#Pneumonia (490-493)	8	2	6	7	2	5	19.4	14.5	22.5
#Diarrhea & enteritis (571)	3	0	3	5	0	5	13.9	-	22.5
Acute nephritis (590, 591)	2	0	2	4	1	3	11.1	7.2	13.5
Chronic nephritis, etc. (592-594)	3	0	3	3	0	3	8.3	-	13.5
Maternal deaths (640-689)	2	0	2	1	0	1	0.9	-	1.3
Cong. malformations (750-759)	3	1	2	4	0	4	11.1	-	18.0
Birth inj., atelect. (760-762)	3	1	2	2	1	1	1.8	3.2	1.3
#Other dis. early inf. (763-776)	11	4	7	13	5	8	11.9	16.0	10.3
Violent & acc. deaths (800-999)	23	12	11	23	12	11	63.9	87.0	49.5
Acc. deaths, total (800-962)	17	10	7	16	9	7	44.4	65.2	31.5
Motor vehicle acc. (810-835)	6	3	3	4	1	3	11.1	7.2	13.5
Ill-defined (780-795)	53	10	43	59	12	47	163.9	87.0	211.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
STATE OF MISSISSIPPI, 1956

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	62806	27785	35021	62178	27278	34900	29.5	23.8	36.2
Fetal deaths	2693	1038	1655	2650	1008	1642	42.6	37.0	47.0
Deaths from all causes	19650	9912	9738	19780	9969	9811	9.4	8.7	10.2
Deaths under 1 year	2338	717	1621	2353	734	1619	37.8	26.9	46.4
Deaths under 1 month	1439	569	870	1451	581	870	23.3	21.3	24.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	193	73	120	199	80	119	9.4	7.0	12.4
Tbc., respiratory (001-008)	172	69	103	176	75	101	8.3	6.5	10.5
Syphilis (020-029)	42	10	32	44	13	31	2.1	1.1	3.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	55	16	39	54	16	38	2.6	1.4	3.9
Scarlet fever & streptococcal sore throat (050, 051)	2	2	0	1	1	0	0.05	0.1	-
Diphtheria (055)	4	1	3	5	1	4	0.2	0.1	0.4
Whooping cough (056)	41	2	39	41	2	39	1.9	0.2	4.0
Meningococcal infections (057)	15	6	9	14	5	9	0.7	0.4	0.9
Acute poliomyelitis (080)	12	4	8	12	4	8	0.6	0.3	0.8
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	10	1	9	10	1	9	0.5	0.1	0.9
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	2338	1369	969	2432	1439	993	115.3	125.4	103.2
Diabetes (260)	224	119	105	225	116	109	10.7	10.1	11.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	2527	1217	1310	2550	1229	1321	120.9	107.1	137.2
Heart diseases (410-443)	6104	3731	2373	6104	3703	2401	289.3	322.8	249.4
Arteriosclerosis (450)	247	160	87	247	161	86	11.7	14.0	8.9
Influenza (480-483)	103	25	78	103	26	77	4.9	2.3	8.0
#Pneumonia (490-493)	581	227	354	575	224	351	27.2	19.5	36.5
#Diarrhea & enteritis (571)	178	30	148	178	30	148	8.4	2.6	15.4
Acute nephritis (590, 591)	81	30	51	81	31	50	3.8	2.7	5.2
Chronic nephritis, etc. (592-594)	320	124	196	329	130	199	15.6	11.3	20.7
Maternal deaths (640-689)	75	12	63	74	12	62	1.2	0.4	1.8
Cong. malformations (750-759)	231	132	99	247	142	105	11.7	12.4	10.9
Birth inj., atelect. (760-762)	392	220	172	387	218	169	6.2	8.0	4.8
#Other dis. early inf. (763-776)	798	248	550	804	254	550	12.9	9.3	15.8
Violent & acc. deaths (800-999)	1647	860	787	1630	843	787	77.2	73.5	81.8
Acc. deaths, total (800-962)	1326	719	607	1302	700	602	61.7	61.0	62.5
Motor vehicle acc. (810-835)	585	371	214	558	347	211	26.4	30.2	21.9
Ill-defined (780-795)	1729	367	1362	1705	355	1350	30.8	30.9	140.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.