


PUBLIC HEALTH STATISTICS

STATE OF
MISSISSIPPI

1957



Part I VITAL STATISTICS STATE SUMMARY

Mississippi State Board of Health
Jackson, Mississippi

A. L. Gray, M. D., Executive Officer

ANNUAL BULLETIN

of

VITAL STATISTICS

1957

VOLUME 1

STATE SUMMARY

MISSISSIPPI STATE BOARD OF HEALTH

VOLUME 1
TABLE OF CONTENTS

	Page
General Summary	1
Table	
1. Estimated Population as of July 1, 1956, by County, by Race	19
2. Live Births, by County and City of Occurrence, Reallocated to Mother's Usual Residence, by Race, with Rates	21
3. Live Births, by County and City of Usual Residence of Mother, by Race and Attendant, and by Sex	24
4. Fetal Deaths, by County and City of Occurrence, Reallocated to Mother's Usual Residence, by Race and Attendant, and by Sex	28
5. Deaths, by Cause, by Race	32
6. Deaths, by County and City of Occurrence, Reallocated to Place of Residence, by Race, with Rates	43
7. Deaths from Selected Causes, by Race, Sex, and Age, with Rates	46
8. Deaths from Selected Causes, by Month of Occurrence	55
9. Deaths Under 1 Year, shown for Selected Causes, by Race and Age, with Rates	56
10. Deaths Due to Injuries which Occurred in Mississippi, by County of Injury, by Type of Injury	58
11. Marriages, by County of Occurrence and of Residence of Bride, and Divorces, by County of Occurrence, by Race	61
12. Divorces, by Cause, by number of Minor Children, by Number of Years Married, by Race	63
13. Births, Deaths, and Marriages, by Month of Occurrence	64
14. Live Births, Deaths, Maternal Deaths, Deaths Under 1 Year of Age, and Deaths Under 1 Month of Age, 1948-1957, by Year, by Race, with Rates ..	65
15. Deaths from Heart Diseases, Nephritis (all forms), Vascular Lesions, Cancer, and Tuberculosis (all forms), 1948-1957, by Year, by Race, with Rates	66
16. Births, Fetal Deaths, Deaths Under 1 Year, Deaths Under 1 Month, and Deaths from Certain Selected Causes, on Basis of Place of Occurrence and Place of Residence, by Race, with Rates	67

SUMMARY OF 1957 VITAL STATISTICS

INTRODUCTION

The 1957 Vital Statistics Bulletin is the fifteenth consecutive issue prepared by the Public Health Statistical Service for the Division of Vital Statistics* from data obtained from records of live births, fetal deaths, deaths, marriages and divorces collected through the Division of Vital Statistics.

Volume 1 contains an analysis of the data and tables for the State, the 82 counties and in some instances the 15 cities with population of 10,000 or more. Summary tables for each county are included in Volume 2.

Population and Rates

Population figures used in this bulletin are estimates as of July 1, 1956, prepared by Dr. Harald A. Pedersen, formerly of the Sociology Department, Mississippi State College**, according to Method #II described by the United States Bureau of the Census. Some changes were made by the Public Health Statistical Service to include military population and also to adjust the estimates by race to the total population of each county. Later adjustments by Dr. Pedersen equaling the county totals to a recent state estimate by the Census Bureau were not incorporated into the population figures used in this volume.

No age or sex-specific rates (except for maternal and infant deaths) and no city rates were calculated for 1957 because current estimates by age, sex, and for cities are not available.

Occurrence and Residence Allocation

Births, fetal deaths, deaths, and marriages were classified by occurrence and residence. Occurrence data were obtained from certificates filed in Mississippi at the place of birth, fetal death, death, or issuance of marriage license. Residence data include transcripts of these events (except marriages) occurring to Mississippi residents outside the State, and exclude records filed in Mississippi for nonresidents. There were 1,065 births to residents of Mississippi which occurred outside the State in 1957 and were reported to the Division of Vital Statistics; births occurring inside Mississippi to residents of other states numbered 1,644. Fetal death transcripts totaled only 20 and the nonresident certificates 52. During the year, 805 death transcripts were received and 562 nonresident deaths recorded. The total number of states sending transcripts was 38 including the District of Columbia. Birth records came from 32 states, fetal death records from 7 states, and death records from 31 states. Births to 17 Mississippi residents were reported from Alaska. In Table A the states furnishing the largest numbers of records are listed.

Some of the tables contain both residence and occurrence data; but, following recommended procedures, most of the tables and all of the rates are based on residence data only. Residence assignment for births and fetal deaths was determined on the basis of usual residence of mother; and for deaths, on the basis of usual residence of decedent. Deaths of inmates or employees of institutions were reallocated to previous residence regardless of length of stay in the institution.

* Changed July 1, 1958 to Division of Public Health Statistics.

** Changed July 1, 1957 to Mississippi State University.

Table A. Transcripts received, by place of occurrence

<u>Place of Occurrence</u>	<u>Live Births</u>	<u>Deaths</u>	<u>Fetal Deaths</u>
<u>Total</u>	<u>1,065</u>	<u>805</u>	<u>20</u>
Alabama	169	89	3
Arkansas	17	12	0
Colorado	26	3	0
Florida	16	12	0
Illinois	79	26	2
Louisiana	159	183	1
Missouri	19	12	0
Tennessee	433	382	11
Texas	21	27	0
<u>Other places</u>	<u>126</u>	<u>59</u>	<u>3</u>

Urban-Rural Classification

Towns with populations of 2,500 or more according to the 1950 census were considered as urban areas. All other areas were classified as rural.

Classification of Causes of Death

Causes of death were assigned according to the Sixth Revision (1948) of the International Statistical Classification of Causes of Death. Only underlying causes were coded. Changes in coding procedures since 1949 have been minor, but comparisons with earlier years should be made with caution, since revision of the code was responsible for several shifts in the distribution of deaths by cause.

LIVE BIRTHS

Resident Births

Live births to residents of Mississippi numbered 60,712 in 1957. This total was 1,466 less than that for 1956 and was the lowest since 1945. Declines of 474 for the whites and of 992 for the nonwhites were noted. The crude birth rate of 28.8 was the lowest since 1949. The state rate for the whites in 1957 was 23.4 and the county rates varied from 9.8 in Carroll County to 33.6 in Washington County. For nonwhites, the state rate was 35.2 and the range for the counties was from 24.0 in Union County to 60.0 in Smith County.

Characteristics

A breakdown of the 33,908 nonwhite births by specific race shows 33,741 were Negro, 132 were Indian, 22 were Chinese, 11 were Japanese, and 2 were other nonwhites. Births of males numbered 30,893 and there were 29,819 females. These totals yielded a sex ratio of 1,036 males to 1,000 females. August was the peak month of occurrence for live births. There were 5,689 in this month in comparison with only 4,388 in April when the fewest occurred.

During 1957, 20,056 children were born to urban residents and 40,656 to rural residents. Over half (54 per cent) of the urban resident births were white,

but 61 per cent of the rural were nonwhite. This percentage distribution was the same as for 1956. Of the births to rural residents, 25,299 or 62 per cent were born to farm residents; 12,566 were born to nonfarm residents; and on 2,791 certificates information pertaining to farm residency was not stated. Comparison of the occurrence and residence figures indicates that well over one-half of the white mothers living in rural areas went to cities for delivery as compared with only a small percentage of the nonwhite mothers.

The number of illegitimate births during 1957 was 7,815 or 12.9 per cent of the total. This was a slight increase over 1956 following the trend which has been upward since 1947 when the percentage was 9.1. White births classified as illegitimate totaled 272 or 1.0 per cent, and 7,543 or 22.2 per cent of the nonwhite births were illegitimate.

Liveborn infants classified as immature (birth weight $5\frac{1}{2}$ pounds or less) numbered 5,462 in 1957, or 8.9 per cent of the total. The incidence of immaturity was greater among the nonwhites than among the whites, 10.6 per cent in comparison with 6.7 per cent.

Classification by Attendant

In Table B which gives the distribution of live births by attendant and race, it may be seen that, in 1957, over 68 per cent of all births occurred in hospitals or clinics (including physicians' offices or clinics which provide in-patient care). A small proportion, 5.3 per cent, was delivered by physicians in places other than hospitals or clinics; and 26.3 per cent were delivered at home by nonmedical attendants. Almost half of the nonwhite mothers, but less than one per cent of the white mothers, had no medical attendant.

Table B. Live births by attendant and race

Attendant	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
<u>Total</u>	<u>60,712</u>	<u>100.0</u>	<u>26,804</u>	<u>100.0</u>	<u>33,908</u>	<u>100.0</u>
Physician in hospital	41,458	68.3	26,259	98.0	15,199	44.8
Physician in home	3,239	5.3	393	1.5	2,846	8.4
Midwife	15,936	26.2	138	0.5	15,798	46.6
Other	79	0.1	14	0.05	65	0.2

There has been a steady increase in the proportion of live births occurring in hospitals in recent years (see Table C). The year 1951 was the first in which at least half of the deliveries took place in hospitals, and the proportion of hospital deliveries has increased each year since. The progress made by the nonwhites has been more rapid, but there is still a large difference between the percentages for the two racial groups.

Table C. Percentage of live births in hospitals or clinics, 1953-1957, by race

<u>Year</u>	<u>Total</u>	<u>White</u>	<u>Nonwhite</u>
1957	68.3	98.0	44.8
1956	65.7	97.2	41.0
1955	62.4	96.0	36.4
1954	60.2	94.4	33.0
<u>1953</u>	<u>58.2</u>	<u>93.3</u>	<u>30.5</u>

The proportions of live births attended by physicians during 1953-1957 are shown in Table D. The improvement shown in this table is less spectacular than that in the preceding table, but is also steady. This is the second year in which more than half the nonwhite mothers were attended by physicians.

Table D. Percentage of live births attended by physicians, 1953-1957, by race

<u>Year</u>	<u>Total</u>	<u>White</u>	<u>Nonwhite</u>
1957	73.6	99.5	53.2
1956	72.2	99.3	51.0
1955	70.4	99.1	48.3
1954	70.4	98.9	47.7
<u>1953</u>	<u>69.6</u>	<u>98.8</u>	<u>46.5</u>

Age of Mother

Table E gives the age distribution of mothers by race. Little variation in this distribution has occurred in recent years. Approximately half of the mothers were 15-24 years of age and well over one-third were 25-34 years old. In the periods 15-24 and 25-34 years the percentages of white mothers were slightly larger than the nonwhites.

Table E. Live births by age of mother and race, with per cent of total births

<u>Age of Mother</u>	<u>Total</u>		<u>White</u>		<u>Nonwhite</u>	
	<u>No.</u>	<u>%</u>	<u>No.</u>	<u>%</u>	<u>No.</u>	<u>%</u>
10-14 years	333	0.5	30	0.1	303	0.9
<u>15-44 years</u>	<u>60,137</u>	<u>99.0</u>	<u>26,724</u>	<u>99.7</u>	<u>33,413</u>	<u>98.5</u>
15-24 years	30,205	49.8	13,414	50.0	16,791	49.5
25-34 years	22,955	37.8	10,629	39.6	12,326	36.4
35-44 years	6,977	11.5	2,681	10.0	4,296	12.7
45-54 years	138	0.2	32	0.1	106	0.3
<u>Unknown</u>	<u>104</u>	<u>0.2</u>	<u>18</u>	<u>0.1</u>	<u>86</u>	<u>0.2</u>

FETAL DEATHS

In 1952 the registration of all fetal deaths, regardless of length of gestation, was initiated in Mississippi. Prior to that time, only stillbirths of 20 or more weeks of gestation were reported. Registration of early fetal deaths is still far from complete, but there is no estimate of the degree of underregistration. There were 2,486 fetal death certificates filed during 1957 and 2,454 certificates

for residents. The residents total was 196 less than the comparable figure for 1956, and a decrease occurred in both races. For the State the fetal death ratio (fetal deaths per 1,000 live births) was 40.4. The ratio for whites was 36.0 with county ratios varying from zero in Amite, Franklin, Issaquena, and Tunica Counties to 118.5 in Washington County. The ratio for nonwhites was 43.9 with variations from zero in Tishomingo to 87.0 in Itawamba.

On 830 of the records the gestation period was stated to be less than 20 weeks; on 1,485 it was 20 weeks or more; and on 139 the length was not stated. In the period 1954-1957 there have been reported among the whites more fetal deaths after less than 20 weeks of gestation than after 20 weeks or more and the difference has grown larger each year. The reporting of late (20 or more weeks of gestation) fetal deaths for both racial groups has declined each year in this period.

Using the certificates with sex stated, a sex ratio of 1,361 males to 1,000 females was determined. This ratio is not so high as the 1956 figure, but is in line with the previous years. As usual, the sex ratio for fetal deaths was much higher than that for live births, but was not so reliable a figure because the sex was not stated on 667 or 27 per cent of the fetal death certificates.

The distribution by race and attendant is given in Table F. Approximately 61 per cent of the total were nonwhite. More than 69 per cent of the reported fetal deaths were attended by physicians in hospitals and about 19 per cent were attended by midwives. Although the proportion of white live births in hospitals was larger than the percentage of white fetal deaths in hospitals, the reverse was true for the nonwhites. This situation in the nonwhite group may be accounted for by the under-reporting of fetal deaths that occur at home and the tendency to transfer difficult and premature deliveries to a hospital.

Table F. Fetal deaths by attendant and race

<u>Attendant</u>	<u>Total</u>		<u>White</u>		<u>Nonwhite</u>	
	<u>No.</u>	<u>%</u>	<u>No.</u>	<u>%</u>	<u>No.</u>	<u>%</u>
<u>Total</u>	<u>2,454</u>	<u>100.0</u>	<u>965</u>	<u>100.0</u>	<u>1,489</u>	<u>100.0</u>
Physicians in hospital	1,706	69.5	890	92.2	816	54.8
Physicians in home	252	10.3	67	6.9	185	12.4
Midwife	477	19.4	6	0.6	471	31.6
Other	19	0.8	2	0.2	17	1.1

Table G shows the percentage of fetal deaths delivered in hospitals in the past five years. There has been very little change in the figures for the whites during this time, but the nonwhites have shown a steady improvement from 42.6 per cent in 1953 to 54.8 per cent in 1957. The gap between the two race groups is a large one, but not so large as that for live births.

Table G. Percentage of fetal deaths delivered in hospitals, 1953-1957, by race

<u>Year</u>	<u>Total</u>	<u>White</u>	<u>Nonwhite</u>
1957	69.5	92.2	54.8
1956	67.2	91.0	52.7
1955	63.5	88.1	48.5
1954	61.6	90.2	44.6
1953	56.4	89.6	42.6

DEATHS

Resident Deaths

In 1957 the resident death total was 20,476 which is an increase of 696 over the number for 1956. The 1957 figure, however, is only 408 above the mean for the past 14 years, the years for which resident figures are available. The crude death rate, based on the 1956 population estimate, was 9.7 per 1,000 population in comparison with a rate of 9.4 in 1956 based on the same estimate. Slightly over one-third or 7,045 of the persons who died in 1957 were urban residents, but 48 per cent of the deaths occurred in urban areas.

Although the number of deaths of nonwhites was 202 less than that of the whites, the death rate of the nonwhites was 1.5 per 1,000 population more than that of the whites. The total of 10,339 deaths of white persons was the largest recorded since 1936 and the second largest reported since 1913, the first year of death registration in Mississippi. The number of deaths of nonwhites in 1957 (10,137) was higher than in 1956 (9,811), 1955 (9,902), or 1954 (9,900). Rates for both races increased 0.3 per 1,000 over the 1956 figure as did the rate for the state as a whole.

County rates for the white population ranged from 3.7 in Issaquena to 16.6 in Leake with the State rate of 8.7, while county rates for the nonwhites ranged from 6.4 in Jefferson Davis to 18.4 in Warren with the State rate of 10.5.

The resident birth rate exceeded the death rate by 19.1 per 1,000 population. For whites the excess was 14.4 and for the nonwhites it was 24.7.

Age at Death

Table H shows the distribution of deaths by age and race. There were increases in all the white groups except groups under one and from 5 to 9 and the 15-24 year group which showed a decrease of approximately 9 per cent under 1956 figures. Despite the large increment over 1956 in the number of deaths of nonwhite persons, declines occurred in the groups under 15 years and in the 35-44 year period. For deaths of all races there was a decrease in 1957 for all age groups under 25 years of age and for the group 35-44 years. Increases were seen in age groups 25-34 and in all age groups 45 years and over, with the major increase in the age group over 65 years of age.

Comparison of the age distribution of the two races reveals significant differences. The percentage of deaths of nonwhites which occurred among children under 10 years of age was more than double that for the whites, and in the groups from 10 through 64 years relatively more deaths of nonwhites than of whites were registered. "Retirement" age or 65 had been reached by nearly 60 per cent of the whites who died in 1957 and only 45 per cent of the nonwhites. The median age at death for the State was 66.1; for the whites, 69.1; and for the nonwhites, 61.8 years.

Table H. Age-race distribution of deaths

Age Group	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
All Ages	20,476	100.0	10,339	100.0	10,137	100.0
Under 1 year	2,257	11.0	694	6.7	1,563	15.4
1-4 years	342	1.7	107	1.0	235	2.3
5-9 years	140	0.7	51	0.5	89	0.9
10-14 years	132	0.6	64	0.6	68	0.7
15-24 years	456	2.2	193	1.9	263	2.6
25-34 years	581	2.8	246	2.4	335	3.3
35-44 years	953	4.6	435	4.2	518	5.1
45-54 years	1,913	9.3	914	8.8	999	9.8
55-64 years	2,944	14.4	1,483	14.3	1,461	14.4
65-74 years	4,528	22.1	2,402	23.2	2,126	21.0
75-84 years	4,232	20.7	2,649	25.6	1,583	15.6
85 years and over	1,970	9.6	1,098	10.6	872	8.6
Unknown	28	0.1	3	0.03	25	0.2

Leading Causes of Death

In Table I are listed the ten leading causes of death in the State in 1957. The first seven of these causes are the same as in 1955 and in 1956. Congenital malformations and general arteriosclerosis which tied for eighth place in 1956, rank in ninth and tenth places respectively in 1957. Diabetes mellitus, which was in tenth place in 1956, moved up to eighth place in 1957.

Altogether the ten leading causes were responsible for 78 per cent of the deaths. For the whites the same ten causes were listed as for the total population with the exception that suicide ranked in eighth place and chronic and unspecified nephritis and other renal sclerosis was not on the list. For nonwhites, however, homicide and gastritis including associated conditions replaced congenital malformations and general arteriosclerosis respectively. On further breakdown, it was noted that tuberculosis dropped from twelfth place to sixteenth.

Table I. Ten leading causes of death by race, with per cent of total deaths

Cause with International List Number	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
1. Heart diseases (410-443)	6,367	31.1	3,706	35.8	2,661	26.2
2. Vascular lesions affecting central nervous system (330-334)	2,800	13.7	1,420	13.7	1,380	13.6
3. Malignant neoplasms (140-205)	2,496	12.2	1,482	14.3	1,014	10.0
4. Accidents (800-962)	1,302	6.4	707	6.8	595	5.9
5. Certain diseases of early infancy (760-776)	1,142	5.6	414	4.0	728	7.2
6. Pneumonia and influenza (480-493)	794	3.9	301	2.9	493	4.9
7. Chronic & unspec. nephritis & other renal sclerosis (592-594)	304	1.5	119	1.2	185	1.8
8. Diabetes mellitus (260)	260	1.3	140	1.4	120	1.2
9. Congenital malformations (750-759)	259	1.3	168	1.6	91	0.9
10. General arteriosclerosis (450)	234	1.1	139	1.3	95	0.9

Heart disease, the leading cause of death, was responsible for nearly one-third of all deaths in 1957. During the year 6,367 deaths were attributed to some form of heart disease. On Table J, it may be seen that a large number of these (4,282 deaths) were specified as arteriosclerotic and degenerative heart diseases and that the next largest group was hypertensive heart disease with 1,458 deaths. The total death rate was 301.8 per 100,000 population, an increase from 289.3 for 1956. The increase was almost entirely among the nonwhites.

Table J. Deaths from heart disease, by race by type of disease

Cause of Death and International List Number	Race		
	Total	White	Nonwhite
Diseases of the heart	<u>6,367</u>	<u>3,706</u>	<u>2,661</u>
Chronic rheumatic heart disease (410-416)	146	78	68
Arteriosclerotic and degenerative heart disease (420-422)	4,282	2,878	1,404
Other diseases of heart (430-434)	481	210	271
Hypertension with heart disease (440-443)	1,458	540	918

For the second leading cause of death, vascular lesions affecting the central nervous system, there were 2,800 deaths representing 13.7 per cent of the total. The components of this category are listed on Table K showing that a majority of the deaths were due to cerebral hemorrhage (1,757), and 570 were due to cerebral embolism or thrombosis. The rate continued to increase; and, following the pattern of past years, the rate for nonwhites exceeded that for whites although in 1957, for the first time, the number of deaths of white persons was greater than the number of nonwhites who died from these causes.

Table K. Deaths from vascular lesions affecting the central nervous system by race by type of lesion

Cause of Death and International List Number	Race		
	Total	White	Nonwhite
Vascular lesions affecting central nervous system (330-334)	<u>2,800</u>	<u>1,420</u>	<u>1,380</u>
Subarachnoid hemorrhage (330)	63	36	27
Cerebral hemorrhage (331)	1,757	843	914
Cerebral embolism and thrombosis (332)	570	354	216
Other and ill-defined (334)	410	187	223

In 1957, the cancer death totals for both racial groups were the highest on record in this State. There were 2,496 deaths from this cause or 64 more than in 1956. The distribution of these deaths by age, sex, and race is given in Table L. A small decline from the number reported in 1956 was observed for nonwhite females, and the figures for white females and for males of both races showed increases. White males showed decreases in the age groups under 15 and over 65; nonwhite males showed decreases in groups under 15, 25-34, 35-44, and 75 years and over; white females showed decreases in ages under 45; nonwhite females showed decreases in age groups under 15, 35-44, 45-54, and 65-74 with the major decreases in the older groups. All four of the race-sex groups showed decreases in the age groups under 15 and all four showed increases in the age group 55-64.

More deaths occurred among the whites than among nonwhites in all age groups except 15-24 in which group the same number of deaths occurred for both races, and the excess was largest in the group 75-84 years. There were more males than females in the 55-64 and 65-74 year age groups and more females in the age groups 25-34, 35-44, 45-54, and 85 years and over.

Table L. Cancer deaths by age, sex, and race

Age Group	Total	Male		Female	
		White	Nonwhite	White	Norwhite
All Ages	2,496	762	498	720	516
Under 10 years	32	12	7	8	5
10-14 years	5	2	1	2	0
15-24 years	28	9	7	5	7
25-34 years	58	13	6	16	18
35-44 years	139	40	13	47	39
45-54 years	379	93	83	109	94
55-64 years	570	176	129	142	123
65-74 years	656	214	146	182	114
75-84 years	496	158	90	165	83
85 years & over	132	40	15	44	33
Unknown	1	0	1	0	0

Table M lists cancer deaths by primary site in descending order according to the number in the total population. The number of deaths assigned to the leading site, breast and genito-urinary system is one less than in 1956, but this site moved up to first position due to a decrease of 21 in deaths from cancer of the digestive organs and peritoneum. These two leading sites accounted for 79 per cent of the cancer deaths among nonwhite females, 70 per cent among white females and among non-white males, and only 44 per cent among white males. Cancer of the respiratory system showed an increment of 58 and moved to third rank, but there was a decrease of 57 in the number of deaths from cancer of the lymphatic and hematopoietic tissues.

Table M. Cancer deaths by primary site, sex, and race, with percentage distribution

Primary site and International List Number	Male				Female			
	White		Nonwhite		White		Nonwhite	
	No.	%	No.	%	No.	%	No.	%
Total	762	100.0	498	100.0	720	100.0	516	100.0
1. Breast & genito-urinary system (170-181)	134	17.6	139	27.9	277	38.5	249	48.2
2. Digestive organs & peritoneum (150-159)	199	26.1	207	41.6	230	31.9	157	30.4
3. Respiratory system (160-165)	178	23.4	51	10.2	32	4.4	21	4.1
4. Lymphatic & hematopoietic tissues (200-205)	73	9.6	32	6.4	71	9.9	15	2.9
5. Brain & nervous system (192,193)	38	5.0	11	2.2	20	2.8	5	1.0
6. Buccal cavity & pharynx (140-148)	26	3.4	10	2.0	15	2.1	6	1.2
7. Skin (190,191)	32	4.2	3	0.6	18	2.5	3	0.6
8. Bone & connective tissue (196,197)	13	1.7	5	1.0	9	1.2	6	1.2
9. Other & unspecified sites (194,195,198,199)	69	9.0	40	8.0	48	6.7	54	10.5

The number of deaths in 1957 which resulted from accidents was 1,302. Although this was the same number of accidental deaths as in 1956, those caused by motor vehicle accidents showed a small decrease, from 558 to 551. Aircraft, falls, electric current, firearm, obstruction or suffocation due to inhalation or ingestion of food and other objects, and drowning were types of accidents for which the number of deaths went up in 1957. The type of accident resulting in the greatest increase over 1956 in the number of deaths among the whites was falls and among the nonwhites, drowning. Showing the greatest decrease for both races was fire and explosion. The largest number of accidental deaths occurred in December which was also the month when motor vehicle fatalities were highest. July moved from third place in the number of deaths from all accidents and sixth place in the number of deaths from motor vehicle accidents to the month with the second largest number of deaths from each of these causes. Table N shows the six leading causes of accidental death in 1957 by race. The first five of these causes appeared on the 1955 and 1956 lists in the same order.

Table N. Six leading causes of accidental deaths among residents, by race

Type of Accident	Total	White	Nonwhite
1. Motor vehicle	551	346	205
2. Fire	134	34	100
3. Fall	121	85	36
4. Drowning except water transport	96	41	55
5. Firearm	71	38	33
6. Suffocation or obstruction	37	11	26

Leading causes of death for each age group except infants are given in Table O. None of these lists was exactly like any other, but there were some causes which appeared repeatedly. Accidents not only was listed in every group, but also was the chief cause in all groups under 45 years of age. Malignant neoplasms also appeared on every list and heart diseases on all except 1-4 years. Vascular lesions affecting the central nervous system was listed in the three groups 25 and over. Also appearing in four groups (1-4, 5-14, 45-64, and 65 years and over) was pneumonia and influenza. In both lists for children under 15, congenital malformations was included, and for 15-24 and 25-44 year groups homicide and maternal deaths were listed. The gastritis group was listed only for the 1-4 year group. Special tabulations of the age group 65 years and over show that the lists for age groups 70-74 and 80-84 are the same as the list for the age group 65 years and over as a whole. In age group 65-69 diabetes ranks in fourth place moving pneumonia and influenza to fifth place and dropping accidents from the list of five causes. In age groups 75-79 and 85 years and over arteriosclerosis replaced accidents.

Ill-defined Causes

Certificates with no satisfactory statement of cause numbered 2,357 or 11.5 per cent of the total in 1957. Senility was the only cause given on 74; ill-defined diseases of early infancy on 594; and other ill-defined and unknown causes on 1,689. For white persons the percentage of certificates in this category was 5.4 and for non-whites, 17.8. Over 20 per cent of the certificates filed in DeSoto and Carroll Counties had no definite cause, and in Washington, Marshall, Issaquena, Hancock, Tunica, Claiborne, Humphreys, Leflore, and Yazoo Counties 15 to 20 per cent had ill-defined and unknown causes. Of the counties listed in 1956, six remain on the list in 1957.

Table 0. Five leading causes of death by age group

<u>Age and Cause of Death</u>	<u>No. of Deaths</u>	<u>% of Total</u>
1-4 years		
All Causes	<u>342</u>	<u>100.0</u>
1. Accidents	101	29.5
2. Pneumonia and influenza	48	14.0
3. Congenital malformations	33	9.6
4. Gastritis, duodenitis, enteritis, and colitis	16	4.7
5. Malignant neoplasms, including leukemia	13	3.8
5-14 years		
All Causes	<u>272</u>	<u>100.0</u>
1. Accidents	127	46.7
2. Malignant neoplasms, including leukemia	19	7.0
3. Pneumonia and influenza	18	6.6
4. Congenital malformations	12	4.4
5. Heart diseases	8	2.9
15-24 years		
All Causes	<u>456</u>	<u>100.0</u>
1. Accidents	211	46.3
2. Homicide	39	8.6
3. Heart diseases	30	6.6
4. Malignant neoplasms, including leukemia	28	6.1
5. Maternal deaths	18	3.9
25-44 years		
All Causes	<u>1,534</u>	<u>100.0</u>
1. Accidents	295	19.2
2. Heart diseases	280	18.2
3. Malignant neoplasms, including leukemia	197	12.8
4. Vascular lesions affecting central nervous system	98	6.4
Homicide	98	6.4
5. Maternal deaths	55	3.6
45-64 years		
All Causes	<u>4,857</u>	<u>100.0</u>
1. Heart diseases	1,711	35.2
2. Malignant neoplasms, including leukemia	949	19.5
3. Vascular lesions affecting central nervous system	621	12.8
4. Accidents	238	4.9
5. Pneumonia	92	1.9
65 years and over		
All Causes	<u>10,730</u>	<u>100.0</u>
1. Heart diseases	4,323	40.3
2. Vascular lesions affecting central nervous system	2,063	19.2
3. Malignant neoplasms, including leukemia	1,284	12.0
4. Pneumonia and influenza	385	3.6
5. Accidents	241	2.2

Deaths thus classified were so numerous that if placed on the list of ten leading causes they would rank fourth and would appear as one of the five chief causes in all age groups. If the true causes of death for these persons could be determined and properly assigned, the order of the principal causes might not be the same and even the causes might be different.

Infant Mortality

An infant mortality rate of 37.2 per 1,000 live births was recorded in 1957, slightly lower than the rate for 1956. The rate for both races was lower than that of 1956, but the white rate showed the greater decrease. Distribution of infant deaths by age and race is shown in Table P. In each age group, the rate for nonwhites exceeded that for whites with the greatest difference between the two race groups occurring in the period 28 days to one year.

Table P. Infant deaths by age and race, with rates

Age at Death	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
Total under 1 year	2,257	37.2	694	25.9	1,563	46.1
Under 1 day	665	11.0	266	9.9	399	11.8
One to 28 days	760	12.5	274	10.2	486	14.3
28 days to 1 year	832	13.7	154	5.7	678	20.0

* Deaths per 1,000 live births

Table Q gives the five leading causes of infant deaths. There was no change from 1956 in the causes listed or their rank. Immaturity, which is no longer considered a separate classification in ranking causes of infant deaths, was mentioned either in combination with another cause or as the only cause on 42 per cent of the certificates for white infants and on 33 per cent of those for nonwhites.


Table Q. Leading causes of infant deaths, by race

Cause of Death	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
All Causes	2,257	100.0	694	100.0	1,563	100.0
1. Postnatal asphyxia and atelectasis	224	9.9	111	16.0	113	7.3
2. Influenza & pneumonia, except pneumonia of newborn	207	9.2	30	4.3	177	11.3
3. Congenital malformations	191	8.5	126	18.2	65	4.2
4. Birth injuries	141	6.2	62	8.9	79	5.0
5. Gastritis, duodenitis, enteritis & colitis except diarrhea of newborn	124	5.5	10	1.4	114	7.3

Infant death certificates which showed ill-defined causes or no cause at all numbered 935 or 41 per cent of the total in 1957. For 32 per cent of the white and 46 per cent of the nonwhite infants no definite cause of death was given. Determination of the true cause of death for this large group of infants might change the leading causes of death which are listed above.

Infant mortality rates by race for 1920-1957 are shown in Graph 1. In general, the trend has been downward for both races, but in recent years there has been some upward swing in the curve for the nonwhites.

Graph 1. Infant Mortality
1920-1957



Maternal Mortality

The maternal mortality rate for 1957 was 1.2 per 1,000 live births in the total population, 0.5 for white mothers and 1.8 for nonwhites. The total of 75 maternal deaths was an increase of one over 1956. The white increase was 2 deaths and the nonwhite decrease was one. There is still a large discrepancy between the rates by race, but the gap has narrowed considerably in recent years.

In 1957, as in each year from 1947 through 1956, the two leading causes were hemorrhage and toxemia. In 1957, hemorrhage ranks as leading cause, while in each of the 10 preceding years toxemia was the leading cause.

Table R. Maternal deaths by cause and race, with rates

<u>Cause of Death</u>	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
<u>All Causes</u>	<u>75</u>	<u>1.2</u>	<u>14</u>	<u>0.5</u>	<u>61</u>	<u>1.8</u>
Toxemia	21	0.3	5	0.2	16	0.5
Hemorrhage	31	0.5	8	0.3	23	0.7
Sepsis	4	0.06	0	-	4	0.1
Ectopic pregnancy	2	0.03	0	-	2	0.06
Abortion without sepsis or toxemia	1	0.02	0	-	1	0.03
Abortion with sepsis	4	0.06	0	-	4	0.1
Abortion with toxemia, without sepsis	1	0.02	0	-	1	0.03
<u>Other complications</u>	<u>11</u>	<u>0.2</u>	<u>1</u>	<u>0.04</u>	<u>10</u>	<u>0.3</u>

* Deaths per 1,000 live births.

Table S gives the age and race-specific maternal death rates. There were no deaths in the 10-14 year group and only 1 death in the 45-54 year group and 1 death in the groups above 54 years. At all ages the rates for nonwhite mothers were higher than those for white mothers.


Table S. Maternal deaths by age group and race, with rates

<u>Age Group</u>	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
<u>All Ages</u>	<u>75</u>	<u>1.2</u>	<u>14</u>	<u>0.5</u>	<u>61</u>	<u>1.8</u>
10-14 years	0	-	0	-	0	-
15-24 years	18	0.6	2	0.1	16	1.0
25-34 years	27	1.2	2	0.2	25	2.0
35-44 years	28	4.0	9	3.4	19	4.4
45-54 years	1	7.2	0	-	1	9.4
55 years & over	1	#	1	#	0	-

* Deaths per 1,000 live births to mothers in specific age-race group.

No live births in this age group. Death was late effects of maternal causes.

Maternal death rates by race for 1913-1957 are shown in Graph 2. A downward trend began in the 1930's and has been rapid since 1940.



In 1957 the number of marriages occurring in Mississippi was 62,495. This was a decrease of 3,728 from the 1956 figure. There were 1,272 fewer marriages among the whites and 2,456 among the nonwhites. The average daily number of marriages was 171, ten less than the average for 1956; but in June, the month with the largest total, the daily average was 213. A new marriage law passed in a special session of the legislature in December 1957 apparently was responsible for the few marriages recorded in December. In recent years, the average daily number of marriages in December has far exceeded the number in any other month. Although the law (with a three day waiting period, blood tests, and proof of age) did not become effective until July 1, 1958, there was widespread publicity when it was signed and many out-of-state couples evidently did not attempt to secure licenses in Mississippi.

This was the first marriage for 42,192 or 67.5 per cent of the brides and 42,815 or 68.5 per cent of the grooms; and for about a fourth of each group there had been one previous marriage. Approximately 500 brides and the same number of grooms did not state their previous marital status.

Table T shows the age distribution of brides and grooms. The brides were relatively young; 29,060 or 46.5 per cent were teen-agers and nearly 69 per cent were under 25 years of age. There was a much smaller percentage of grooms in the teen-age group -- less than 10 per cent. Of all marriages, 45.9 per cent or 28,711

of the brides were in the 15-19 year age group and 45.4 per cent or 28,366 of the grooms were in the 20-24 year age group. A cross tabulation of the ages of brides and grooms shows marriages of one couple with both parties under 15 years of age and 19 couples with both 75 years of age and over. While the youngest groom of a bride 75 years or more was 21 years of age and the youngest bride of a groom of 75 was 30 years of age, most marriages were between persons of approximately the same age.

Table T. Age distribution of brides and grooms

<u>Age</u>	<u>Bride</u>		<u>Groom</u>	
	<u>No.</u>	<u>%</u>	<u>No.</u>	<u>%</u>
<u>Total</u>	<u>62,495</u>	<u>100.0</u>	<u>62,495</u>	<u>100.0</u>
Under 15 years	349	0.5	2	0.003
15 years	1,151	1.8	9	0.01
16 years	2,427	3.9	137	0.2
17 years	2,679	4.3	808	1.3
18 years	15,689	25.1	2,347	3.7
19 years	6,765	10.8	2,867	4.6
20-24 years	13,910	22.2	28,366	45.4
25-34 years	9,357	15.0	14,351	23.0
35-44 years	5,328	8.5	6,142	9.8
45-54 years	3,088	4.9	3,790	6.1
55 years & over	1,735	2.8	3,662	5.8
Unknown	17	0.02	14	0.02

Among the marriages between nonresident brides and grooms there were 171 brides and 2 grooms under 15 years of age; and 18,484 or 64.4 per cent of the brides 15-19 years of age were nonresidents.

In Table U are shown data on the residence status of both brides and grooms. In only 30 per cent of the marriages were both parties residents of this State. In the period 1946-1957, 386,624 nonresident brides were married in Mississippi. For the first time since 1947 a decrease was seen in the number of out-of-state brides with the 1957 figure being 1,702 lower than that of 1956. Nonresident brides in 1957 numbered 41,510 or 66.4 per cent of the total. This figure is expected to drop much lower in 1958 and the following years due to the new marriage laws which are effective beginning July 1, 1958.

Table U. Marriages in Mississippi by residence status of bride and groom

<u>Groom</u>	<u>Total</u>	<u>Bride</u>		
		<u>Resident</u>	<u>Nonresident</u>	<u>Not Stated</u>
<u>Total</u>	<u>62,495</u>	<u>20,864</u>	<u>41,510</u>	<u>121</u>
Resident	19,911	18,572	1,315	24
Nonresident	42,462	2,265	40,176	21
Not stated	122	27	19	76

Brides came to Mississippi from every state in the union, the District of Columbia, Hawaii, and Puerto Rico, and there were 38 from other foreign countries.

Bordering states furnished 48.6 per cent of the brides married in Mississippi and 47 per cent of the grooms with totals as follows: Alabama, 11,218 brides and 10,905 grooms; Arkansas, 4,412 brides and 4,060 grooms; Louisiana, 4,825 brides and 4,877 grooms; and Tennessee, 9,894 brides and 9,511 grooms. Four other states, three of which were only one state removed from Mississippi, furnished over 1,000 brides each and over 1,000 grooms each: Florida, 1,798 brides and 1,771 grooms; Illinois, 2,870 brides and 3,079 grooms; Kentucky, 1,234 brides and 1,245 grooms; and Missouri, 2,577 brides and 2,580 grooms. These 8 states furnished 62.1 per cent of the brides married in Mississippi and 60.8 per cent of the grooms.

Each county in the State issued marriage licenses to at least a few out-of-state brides in 1957, but the counties adjacent to Alabama, Tennessee, and Arkansas issued the largest numbers. Table V shows the counties to which the greatest numbers of out-of-state brides came. In Alcorn, DeSoto and Tishomingo Counties over 95 per cent of the licenses were for nonresident brides, and in George this percentage was over 90. Nearly 30 per cent of all the brides married in Mississippi were nonresidents who came to DeSoto, Alcorn, and Tishomingo Counties -- counties bordering Tennessee.

Table V. Number of out-of-state brides obtaining marriage license in certain counties

DeSoto	8,430	George	1,872
Alcorn	7,058	Lauderdale	1,368
Tishomingo	3,177	Itawamba	1,210
Lowndes	3,020	Harrison	1,051
Washington	2,396	Hancock	1,033
Jackson	2,006	Marshall	818

DIVORCES

Divorces granted in Mississippi during 1957 numbered 4,974 - 53 less than in 1956. There was a decrease of 102 among the whites and an increase of 49 among the nonwhites. Cruel and inhuman treatment and desertion continued to rank as the two leading causes and together accounted for nearly 87 per cent of the total granted. Of the divorces among the whites, cruel and inhuman treatment was given as the main cause for 58 per cent; among nonwhites 60 per cent were granted on grounds of desertion. The wife was listed as the complainant on 63 per cent of the records. Only 249 suits or 5 per cent were contested.

No minor children were involved in 54 per cent of the divorces, and 21 per cent of the couples had only one minor child. Seven of the families had 10 or more minor children. The total number of minor children affected was 4,541.

Forty-three per cent of the divorces in 1957 were granted to persons married less than 5 years; 8 per cent had been married less than one year; but 14 per cent

had been married 20 years or longer. More divorces were granted to couples in their second year of marriage than in any other year. The distribution of divorces according to duration of marriage was different for the two races. Of the white couples, 51 per cent were granted after less than 5 years of marriage and only 8 per cent after 20 years or longer; for the nonwhites, corresponding percentages were 25 and 27 respectively.

TABLE 1
ESTIMATED POPULATION*, MISSISSIPPI
AS OF JULY 1, 1956

County	Total	White	Nonwhite
<u>TOTAL</u>	<u>2,109,800</u>	<u>1,147,200</u>	<u>962,600</u>
Adams	36,300	19,300	17,000
Alcorn	25,100	21,500	3,600
Amite	16,900	7,200	9,700
Attala	23,200	12,300	10,900
Benton	8,500	4,300	4,200
Bolivar	59,100	17,600	41,500
Calhoun	15,100	10,500	4,600
Carroll	12,900	4,900	8,000
Chickasaw	17,200	9,800	7,400
Choctaw	9,400	6,200	3,200
Claiborne	11,200	3,100	8,100
Clarke	17,300	10,100	7,200
Clay	15,100	7,600	7,500
Coahoma	52,600	14,400	38,200
Copiah	27,400	12,400	15,000
Covington	15,300	9,500	5,800
DeSoto	24,100	8,400	15,700
Forrest	44,500	32,300	12,200
Franklin	8,800	5,300	3,500
George	10,500	9,200	1,300
Greene	8,000	6,400	1,600
Grenada	18,700	8,700	10,000
Hancock	13,100	11,800	1,300
Harrison	104,700	87,600	17,100
Hinds	170,700	98,400	72,300
Holmes	28,000	6,800	21,200
Humphreys	21,300	6,000	15,300
Issaquena	4,800	1,900	2,900
Itawamba	13,600	12,700	900
Jackson	42,500	34,000	8,500
Jasper	16,900	8,000	8,900
Jefferson	10,100	2,500	7,600
Jeff Davis	16,200	6,400	9,800
Jones	51,500	37,700	13,800
Kemper	11,900	3,700	8,200
Lafayette	16,200	8,100	8,100

Table 1 - Page 2
Estimated Population*, Mississippi, as of July 1, 1956

20

<u>County</u>	<u>Total</u>	<u>White</u>	<u>Nonwhite</u>
Lamar	12,300	10,400	1,900
Lauderdale	62,100	39,400	22,700
Lawrence	11,600	6,900	4,700
Leake	15,500	8,200	7,300
Lee	37,400	26,000	11,400
Leflore	46,000	15,800	30,200
Lincoln	26,200	16,700	9,500
Lowndes	37,400	20,800	16,600
Madison	32,300	8,100	24,200
Marion	22,700	13,300	9,400
Marshall	22,900	6,500	16,400
Monroe	30,400	19,000	11,400
Montgomery	13,800	7,400	6,400
Neshoba	18,200	12,700	5,500
Newton	19,000	11,200	7,800
Noxubee	17,200	4,800	12,400
Oktibbeha	22,000	11,200	10,800
Panola	32,100	12,900	19,200
Pearl River	21,600	16,500	5,100
Perry	7,200	5,200	2,000
Pike	32,200	17,300	14,900
Pontotoc	18,200	14,700	3,500
Prentiss	18,100	15,900	2,200
Quitman	25,500	9,400	16,100
Rankin	27,400	15,300	12,100
Scott	19,900	10,900	9,000
Sharkey	12,500	3,400	9,100
Simpson	21,100	13,000	8,100
Smith	13,700	11,100	2,600
Stone	6,400	4,900	1,500
Sunflower	54,100	16,300	37,800
Tallahatchie	28,400	10,000	18,400
Tate	17,200	6,800	10,400
Tippah	16,600	12,900	3,700
Tishomingo	13,000	12,300	700
Tunica	22,700	4,400	18,300
Union	18,800	14,800	4,000
Walthall	14,300	7,900	6,400
Warren	39,500	22,800	16,700
Washington	76,500	31,100	45,400
Wayne	15,500	9,700	5,800
Webster	10,500	8,000	2,500
Wilkinson	13,200	3,400	9,800
Winston	19,500	11,100	8,400
Yalobusha	13,000	6,800	6,200
Yazoo	36,000	13,800	22,200

* Prepared by Dr. Harald A. Pedersen, Sociology Department,
Mississippi State College, with adjustments by Public
Health Statistics, Mississippi State Board of Health.

TABLE 2
LIVE BIRTHS
MISSISSIPPI
1957

By Place of Occurrence, Reallocated to Mother's Usual Residence,
by Race, with Rates

County	Place of Occurrence			Place of Residence			Rate*		
	Number		Non-white	Number		Non-white	Rate*		Non-white
	Total	White		Total	White		Total	White	
STATE	61,291	27,320	33,971	60,712	26,804	33,908	28.8	23.4	35.2
Adams	1,766	727	1,039	1,246	544	702	34.3	28.2	41.3
Alcorn	607	485	122	545	434	111	21.7	20.2	30.8
Amite	274	30	244	418	125	293	24.7	17.4	30.2
Attala	570	228	342	531	200	331	22.9	16.3	30.4
Benton	119	9	110	216	78	138	25.4	18.1	32.8
Bolivar	1,953	410	1,543	1,811	414	1,397	30.6	23.5	33.7
Calhoun	436	182	254	405	186	219	26.8	17.7	47.6
Carroll	231	23	208	268	48	220	20.8	9.8	27.5
Chickasaw	439	207	232	458	207	251	26.6	21.1	33.9
Choctaw	207	103	104	214	112	102	22.8	18.1	31.9
Claiborne	256	26	230	305	42	263	27.2	13.5	32.5
Clarke	264	137	127	437	201	236	25.3	19.9	32.8
Clay	555	209	346	579	223	356	38.3	29.3	47.5
Coahoma	1,597	390	1,207	1,616	353	1,263	30.7	24.5	33.1
Copiah	910	275	635	756	238	518	27.6	19.2	34.5
Covington	193	130	63	347	189	158	22.7	19.9	27.2
DeSoto	692	98	594	750	178	572	31.1	21.2	36.4
Forrest	1,538	1,141	397	1,292	839	453	29.0	26.0	37.1
Franklin	122	54	68	246	103	143	28.0	19.4	40.8
George	305	257	48	311	262	49	29.6	28.5	37.7
Greene	126	81	45	209	137	72	26.1	21.4	45.0
Grenada	578	243	335	545	212	333	29.1	24.4	33.3
Hancock	101	66	35	262	213	49	20.0	18.0	37.7
Harrison	3,529	2,820	709	3,225	2,549	676	30.8	29.1	39.5
Hinds	5,596	2,977	2,619	5,336	2,743	2,593	31.2	27.9	35.9
Holmes	810	153	657	825	145	680	29.5	21.3	32.1
Humphreys	700	105	595	716	119	597	33.6	19.8	39.0
Issaquena	49	2	47	110	25	85	22.9	13.2	29.3
Itawamba	98	78	20	284	261	23	20.9	20.6	25.6
Jackson	1,100	750	350	1,303	934	369	30.6	27.5	43.4
Jasper	148	45	103	434	145	289	25.7	18.1	32.5
Jefferson	119	4	115	308	35	273	30.5	14.0	35.9
Jeff Davis	384	172	212	383	136	247	23.6	21.2	25.2
Jones	2,211	1,117	1,094	1,423	870	553	27.6	23.1	40.1
Kemper	236	48	188	330	74	256	27.7	20.0	31.2
Lafayette	537	278	259	529	251	278	32.6	31.0	34.3

Table 2 - Page 2
Live Births - Mississippi - 1957

22

County	Place of Occurrence			Place of Residence			Rate*		
	Number		Non-white	Number		Non-white	Rate*		Non-white
	Total	White		Total	White		Total	White	
Lamar	130	78	52	345	263	82	28.0	25.3	43.2
Lauderdale	2,123	1,088	1,035	1,654	868	786	26.6	22.0	34.6
Lawrence	173	21	152	292	142	150	25.2	20.6	31.9
Leake	429	191	238	466	205	261	30.1	25.0	35.8
Lee	1,256	878	378	1,013	648	365	27.1	24.9	32.0
Leflore	1,701	439	1,262	1,647	355	1,292	35.8	22.5	42.8
Lincoln	693	396	297	669	360	309	25.5	21.6	32.5
Lowndes	1,175	548	627	1,154	523	631	30.8	25.1	38.0
Madison	1,099	188	911	1,088	199	889	33.7	24.6	36.7
Marion	540	298	242	611	333	278	26.9	25.0	29.6
Marshall	769	101	668	814	152	662	35.5	23.4	40.4
Monroe	807	415	392	778	378	400	25.6	19.9	35.1
Montgomery	319	108	211	347	124	223	25.1	16.8	34.8
Neshoba	567	311	256	593	349	244	32.6	27.5	44.4
Newton	460	275	185	459	236	223	24.2	21.1	28.6
Noxubee	557	87	470	544	84	460	31.6	17.5	37.1
Oktibbeha	894	383	511	801	314	487	36.4	28.0	45.1
Panola	887	257	630	909	256	653	28.3	19.8	34.0
Pearl River	674	494	180	582	398	184	26.9	24.1	36.1
Perry	97	69	28	223	146	77	31.0	28.1	38.5
Pike	1,009	528	481	851	373	478	26.4	21.6	32.1
Pontotoc	267	183	84	317	224	93	17.4	15.2	26.6
Prentiss	509	428	81	444	372	72	24.5	23.4	32.7
Quitman	823	226	597	787	215	572	30.9	22.9	35.5
Rankin	230	31	199	610	273	337	22.3	17.8	27.8
Scott	528	264	264	556	251	305	27.9	23.0	33.9
Sharkey	284	57	227	403	71	332	32.2	20.9	36.5
Simpson	531	236	295	500	248	252	23.7	19.1	31.1
Smith	186	128	58	364	208	156	26.6	18.7	60.0
Stone	145	99	46	138	86	52	21.6	17.6	34.7
Sunflower	1,417	266	1,151	1,523	319	1,204	28.2	19.6	31.8
Tallahatchie	812	140	672	849	175	674	29.9	17.5	36.6
Tate	393	8	385	552	130	422	32.1	19.1	40.6
Tippah	333	233	100	301	202	99	18.1	15.6	26.8
Tishomingo	138	131	7	264	243	21	20.3	19.8	30.0
Tunica	698	84	614	714	84	630	31.4	19.1	34.4
Union	473	369	104	398	302	96	21.2	20.4	24.0
Walthall	628	366	262	345	151	194	24.1	19.1	30.3
Warren	1,488	664	824	1,192	575	617	30.2	25.2	36.9
Washington	2,862	1,133	1,729	2,709	1,046	1,663	35.4	33.6	36.6
Wayne	457	279	178	453	244	209	29.2	25.2	36.0
Webster	153	62	91	252	146	106	24.0	18.2	42.4

Table 2 - Page 3
Live Births - Mississippi - 1957

23

County	Place of Occurrence			Place of Residence			Rate*		
	Number		Non-white	Number		Non-white	Rate*		Non-white
	Total	White		Total	White		Total	White	
Wilkinson	497	190	307	412	89	323	31.2	26.2	33.0
Winston	474	196	278	515	214	301	26.4	19.3	35.8
Yalobusha	291	78	213	326	124	202	25.1	18.2	32.6
Yazoo	959	256	703	975	281	694	27.1	20.4	31.3
Out-of-State	1,065	874	191	1,644	1,390	254	-	-	-

Cities over 10,000 Population**	Place of Occurrence			Place of Residence		
	Number		Non-white	Number		Non-white
	Total	White		Total	White	
Biloxi	1,997	1,755	242	1,521	1,324	197
Clarksdale	837	375	462	615	241	374
Columbus	900	519	381	586	267	319
Greenville	1,283	600	683	1,265	561	704
Greenwood	859	433	426	576	222	354
Gulfport	1,390	1,049	341	798	525	273
Hattiesburg	1,509	1,141	368	844	506	338
Jackson	5,014	2,971	2,043	3,350	1,640	1,710
Laurel	2,148	1,087	1,061	722	366	356
McComb	648	403	245	271	155	116
Meridian	2,016	1,080	936	1,130	626	504
Natchez	766	636	130	842	380	462
Pascagoula	88	1	87	492	343	149
Tupelo	924	714	210	395	261	134
Vicksburg	1,443	664	779	844	419	425
Total Urban#	33,412	20,267	13,145	20,056	10,751	9,305
Total Rural	27,879	7,053	20,826	40,656	16,053	24,603

* Rate per 1,000 population.

** Cities having over 10,000 population according to 1950 census.

Includes all towns of 2,500 population or more according to 1950 census.

TABLE 3
LIVE BIRTHS
MISSISSIPPI
1957

By County and Cities over 10,000, by Race, Attendant, and Sex
on Basis of Usual Residence of Mother

County	TOTAL*	WHITE			NONWHITE			SEX		
		Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female
STATE	60,712	26,804	26,259	393	152	33,908	15,199	2,846	15,863	30,893
Adams	1,246	544	543	1	0	702	666	6	30	627
Alcorn	545	424	406	21	7	111	53	23	35	276
Amité	418	125	121	4	0	293	82	48	163	209
Attala	531	200	191	8	1	331	87	52	192	272
Benton	216	78	70	6	2	138	28	11	99	94
Bolivar	1,811	414	390	15	9	1,397	575	86	736	892
Calhoun	405	186	180	6	0	219	149	5	65	186
Carroll	268	48	47	0	1	220	60	12	148	130
Chickasaw	458	207	193	14	0	251	108	38	105	241
Choctaw	214	112	105	6	1	102	38	3	61	114
Claiborne	305	42	41	0	1	263	52	2	209	159
Clarke	437	201	199	1	1	236	178	10	48	195
Clay	579	223	221	1	1	356	178	16	162	291
Coahoma	1,616	353	338	13	2	1,263	325	242	696	841
Copiah	756	238	238	0	0	518	339	12	167	390
Covington	347	189	185	3	1	158	118	6	34	179
DeSoto	750	178	172	3	3	572	238	23	311	366
Forrest	1,292	839	838	0	1	453	293	2	158	661
Franklin	246	103	102	1	0	143	102	1	40	130
George	311	262	257	4	1	49	23	1	25	159
Greene	209	137	122	12	3	72	29	9	34	110
Grenada	545	212	210	0	2	333	46	46	241	280
Hancock	262	213	211	1	1	49	20	1	28	143
Harrison	3,225	2,549	2,525	4	4	676	348	29	299	1,630
										1,595

Table 3 - Page 2
Live Births - Mississippi - 1957

County	TOTAL*	WHITE			NONWHITE			SEX		
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male
Hinds	5,336	2,743	2,736	7	0	2,593	1,652	185	756	2,711
Holmes	825	145	139	5	1	680	131	68	481	405
Humphreys	716	119	113	0	6	597	107	3	487	361
Issaquena	110	25	23	0	2	85	44	2	39	57
Itawamba	284	261	242	18	1	23	13	5	5	142
Jackson	1,303	934	927	3	4	369	172	66	131	727
Jasper	434	145	142	3	0	289	215	4	70	208
Jefferson	308	35	34	1	0	273	157	0	116	151
Jeff Davis	383	136	131	5	0	247	135	43	69	186
Jones	1,423	870	868	1	1	553	515	1	37	750
Kemper	330	74	73	0	1	256	83	0	173	163
Lafayette	529	251	242	3	6	278	57	9	212	255
Lamar	345	263	247	15	1	82	45	14	23	188
Lauderdale	1,654	868	866	2	0	786	641	33	112	876
Lawrence	292	142	138	2	2	150	59	11	80	161
Leake	466	205	199	1	5	261	120	5	136	235
Lee	1,013	648	641	7	0	365	186	122	57	520
Leflore	1,647	355	349	4	2	1,292	200	111	981	812
Lincoln	669	360	358	0	2	309	133	120	56	351
Lowndes	1,154	523	490	29	4	631	208	123	300	585
Madison	1,088	199	196	2	1	889	190	50	649	571
Marion	611	333	326	6	1	278	134	4	140	319
Marshall	814	152	149	2	1	662	119	130	413	402
Monroe	778	378	374	2	2	400	157	52	191	391
Montgomery	347	124	122	2	0	223	49	3	171	163
Neshoba	593	349	340	4	5	244	139	1	104	315
Newton	459	236	229	6	1	223	131	33	59	219
Noxubee	544	84	81	1	2	460	89	23	348	275
Oktibbeha	801	314	314	0	0	487	291	0	440	40
Panola	909	256	247	7	2	653	295	95	263	470

Table 3 - Page 3
Live Births - Mississippi - 1957

County	TOTAL*	WHITE			NONWHITE			Male	Female	SEX
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home		
Pearl River	582	398	395	2	1	184	170	7	7	282
Perry	223	146	143	2	1	77	63	5	9	114
Pike	851	373	372	1	0	478	301	15	162	439
Pontotoc	317	224	218	4	2	93	37	6	41	163
Prentiss	444	372	367	5	0	72	43	6	23	237
Quitman	787	215	205	6	4	572	334	61	177	390
Rankin	610	273	269	3	1	337	211	29	97	301
Scott	556	251	243	6	2	305	143	0	162	286
Sharkey	403	71	69	1	1	332	172	2	158	208
Simpson	500	248	248	0	0	252	169	4	79	255
Smith	364	208	200	7	1	156	104	24	28	200
Stone	138	86	86	0	0	86	32	0	20	76
Sunflower	1,523	319	314	3	2	1,204	408	150	646	769
Tallahatchie	849	175	167	6	2	674	165	74	435	444
Tate	552	130	121	9	0	422	37	43	342	272
Tippah	301	202	188	11	3	99	44	33	22	160
Tishomingo	264	243	215	26	6	21	14	7	0	139
Tunica	714	84	75	3	3	630	173	47	410	362
Union	398	302	298	1	3	96	21	13	62	201
Walthall	345	151	150	0	1	194	144	0	50	161
Warren	1,192	575	575	0	0	617	400	5	212	597
Washington	2,709	1,046	1,041	4	1	1,663	900	191	572	1,395
Wayne	453	244	239	3	2	209	135	0	74	218
Webster	252	146	143	3	0	106	26	7	73	129
Wilkinson	412	89	88	0	1	323	92	12	219	202
Winston	515	214	213	0	1	301	75	3	223	246
Yalobusha	326	124	115	9	0	202	65	74	63	173
Yazoo	975	281	261	13	7	694	119	19	556	490

Table 3 - Page 4
Live Births - Mississippi - 1957

Cities over 10,000 Population	TOTAL*	WHITE			NONWHITE			SEX			
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female
Biloxi	1,521	1,324	1,318	2	4	197	141	1	55	766	755
Clarksdale	615	241	241	0	0	374	185	96	93	315	300
Columbus	586	267	260	7	0	319	100	76	143	308	278
Greenville	1,265	561	561	0	0	704	366	73	265	640	625
Greenwood	576	222	220	2	0	354	80	1	273	296	280
Gulfport	798	525	522	1	2	273	133	21	119	411	387
Hattiesburg	844	506	505	0	1	338	212	2	124	425	419
Jackson	3,350	1,640	1,635	5	0	1,710	1,253	132	325	1,718	1,632
Laurel	722	366	365	1	0	356	340	1	15	383	339
McComb	271	155	155	0	0	116	85	4	27	138	133
Meridian	1,130	626	625	1	0	504	450	28	26	612	518
Matchez	842	380	379	1	0	462	445	5	12	421	421
Pascagoula	492	343	342	1	0	149	61	23	65	264	228
Tupelo	395	261	261	0	0	134	68	31	35	212	183
Vicksburg	844	419	419	0	0	425	264	4	157	425	419

* Includes transcripts furnished by the National Office of Vital Statistics of births to residents of Mississippi occurring out-of-state.

TABLE 4
FETAL DEATHS
MISSISSIPPI
1957
By Place of Occurrence, Reallocated to Mother's Usual Residence
by Sex, Race and Attendant

County	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male
STATE	<u>2,486</u>	<u>2,454</u>	<u>890</u>	<u>67</u>	<u>8</u>	<u>1,489</u>	<u>816</u>	<u>185</u>	<u>488</u>	<u>1,030</u>	<u>757</u>
Adams	51	33	5	0	0	28	27	0	1	16	12
Alcorn	12	12	9	0	0	3	1	2	0	5	7
Amitte	5	7	0	0	0	7	3	1	0	0	0
Attala	14	13	2	2	0	11	6	0	5	6	1
Benton	3	15	7	7	0	8	5	1	2	2	0
Bolivar	68	68	14	14	0	54	25	8	21	32	5
Calhoun	21	16	5	0	0	11	10	0	1	9	4
Carroll	11	21	5	0	0	16	7	3	1	10	3
Chickasaw	15	18	4	1	0	14	11	2	1	6	4
Choctaw	2	3	1	0	0	2	1	0	0	0	0
Claiborne	9	12	3	3	0	9	1	0	1	6	3
Clarke	8	6	2	2	0	4	2	1	1	5	0
Clay	12	12	2	2	0	10	8	0	2	9	3
Coahoma	69	67	20	20	3	47	16	11	3	32	23
Copiah	23	22	8	4	0	19	11	3	5	13	8
Covington	3	8	4	4	0	4	3	0	1	3	1
DeSoto	21	24	1	0	1	23	13	3	7	13	4
Forrest	54	49	23	23	0	26	19	0	7	26	11
Franklin	1	6	0	0	0	6	0	0	0	0	0
George	6	8	7	0	0	1	0	0	1	2	4
Greene	3	5	2	2	0	3	0	0	0	3	2
Grenada	29	25	3	3	0	22	6	3	0	13	8
Hancock	4	8	7	6	0	1	1	0	0	4	2
Harrison	80	76	46	45	0	30	19	3	1	32	5

Table 4 - Page 2
Fetal Deaths - Mississippi - 1957

County	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male
Hinds	360	329	193	153	40	0	136	81	22	33	87
Holmes	41	44	5	5	0	0	39	13	8	18	23
Humphreys	24	29	3	2	0	1	26	7	1	18	10
Issaquena	0	1	0	0	0	1	1	0	0	1	1
Itawamba	3	12	10	9	1	0	2	0	2	0	15
Jackson	62	63	37	35	1	1	26	12	8	6	32
Jasper	8	22	2	2	1	0	20	13	0	7	11
Jefferson	5	14	1	1	2	0	13	8	5	5	8
Jeff Davis	14	14	2	2	0	0	12	3	1	4	6
Jones	92	44	24	24	0	1	20	18	1	23	11
Kemper	13	19	5	3	0	0	14	7	0	4	10
Lafayette	19	16	3	3	0	0	13	1	3	3	6
Lamar	0	4	2	2	0	0	2	2	0	2	2
Lauderdale	77	58	21	19	2	0	37	33	0	4	25
Lawrence	4	9	2	2	1	0	7	3	2	1	19
Leake	4	8	1	1	0	0	7	3	0	3	7
Lee	55	43	24	23	0	0	19	16	3	0	20
Leflore	86	81	23	23	0	0	0	7	3	0	25
Lincoln	18	21	10	9	0	0	11	8	2	1	14
Lowndes	41	39	17	17	7	0	0	6	7	9	15
Madison	52	51	11	11	4	4	44	12	11	4	30
Marion	20	26	4	4	0	1	15	11	0	8	11
Marshall	55	44	13	13	4	1	40	12	8	0	20
Monroe	18	18	7	6	4	0	11	8	5	0	13
Montgomery	12	14	4	4	1	0	0	0	0	0	6
Neshoba	12	10	1	1	0	0	0	0	0	0	4
Newton	36	39	20	18	1	0	0	0	0	0	10
Noxubee	30	28	9	3	0	0	0	0	0	0	9
Oktibbeha	14	13	3	3	0	0	0	0	0	0	6
Panola	24	29	13	12	1	0	0	0	0	0	10

Table 4 - Page 3
Fetal Deaths - Mississippi - 1957

County	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total	Phys. in Hosp.	Other	Total	Phys. in Hosp.	Other	Male	Female	Unde- termined
Pearl River	31	25	14	14	0	11	10	1	0	7	10
Perry	0	9	5	5	0	4	4	0	0	3	8
Pike	31	27	10	10	0	17	16	0	1	13	2
Pontotoc	10	16	12	12	0	4	3	1	0	8	6
Prentiss	11	11	7	7	0	4	4	0	0	6	4
Quitman	17	18	3	3	0	9	15	3	1	10	1
Rankin	10	26	13	8	4	13	8	2	3	9	8
Scott	10	14	4	0	0	10	6	0	4	6	4
Sharkey	15	19	5	5	0	14	10	1	3	5	2
Simpson	12	14	4	2	0	10	7	1	2	8	2
Smith	3	7	5	5	0	2	2	0	0	1	1
Stone	2	2	1	1	0	0	0	0	0	0	1
Sunflower	56	61	16	16	0	45	23	3	19	28	11
Tallahatchie	32	32	7	7	0	25	7	8	10	14	7
Tate	21	36	1	1	0	35	14	4	17	21	0
Tippah	20	14	2	2	0	2	2	0	0	4	9
Tishomingo	2	2	0	0	0	0	0	0	0	0	0
Tunica	20	17	0	0	0	0	0	0	0	0	0
Union	20	22	20	20	0	0	2	1	1	5	15
Walthall	28	17	8	8	0	9	9	0	0	3	6
Warren	79	51	21	19	2	30	19	2	0	19	18
Washington	234	230	124	120	4	106	67	11	1	67	132
Wayne	13	23	5	5	0	0	18	1	0	3	3
Webster	4	9	7	7	1	0	2	1	1	5	0
Wilkinson	5	5	1	1	0	0	4	2	0	2	0
Winston	16	19	7	7	0	0	12	3	0	6	6
Yalobusha	7	8	3	2	1	0	5	2	0	5	0
Yazoo	44	44	9	6	1	2	35	11	4	20	4

Table 4 - Page 4
Fetal Deaths - Mississippi - 1957

Cities over 10,000 Population	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total	Phys. in Hosp.	Phys. in Home	Total	Phys. in Hosp.	Phys. in Home	Male	Female	Unde- termined
Biloxi	39	31	22	22	0	9	8	0	1	16	13
Clarksdale	44	30	17	17	0	13	7	4	2	15	9
Columbus	32	30	17	17	0	13	4	3	6	12	13
Greenville	175	160	103	100	3	57	41	3	13	37	17
Greenwood	64	41	15	15	0	26	18	0	8	18	7
Gulfport	39	27	9	9	0	18	9	3	6	9	16
Hattiesburg	54	40	17	17	0	23	16	0	7	20	8
Jackson	320	218	127	100	27	0	91	59	19	55	39
Laurel	90	26	11	11	0	0	15	14	1	0	13
McComb	19	9	7	7	0	2	2	0	0	0	1
Meridian	75	39	16	15	1	0	23	20	0	3	15
Natchez	19	25	4	4	0	0	21	21	0	0	13
Pascagoula	3	20	15	14	1	0	5	3	1	1	12
Tupelo	51	22	10	10	0	0	12	9	3	0	12
Vicksburg	73	35	19	17	2	0	16	13	1	2	10

TABLE 5
DEATHS
MISSISSIPPI
1957

By Cause by Race on Basis of Place of Residence

International List Number	Cause of Death*	Total	White	Non-white
001-999	<u>TOTAL ALL CAUSES</u>	<u>20,476</u>	<u>10,339</u>	<u>10,137</u>
<u>001-019</u>	<u>Tuberculosis, all forms</u>	<u>161</u>	<u>61</u>	<u>100</u>
002	Pulmonary	144	56	88
003	Pleural	1	0	1
008	Unspecified site	5	2	3
010	Meninges and central nervous system	7	3	4
011	Intestines, peritoneum, and mesenteric glands	1	0	1
012	Bones and joints, active or unspecified	1	0	1
016	Genito-urinary system	1	0	1
019	Disseminated	1	0	1
<u>020-029</u>	<u>Syphilis and its sequelae</u>	<u>50</u>	<u>15</u>	<u>35</u>
020	Congenital	1	0	1
022	Aneurysm of aorta	18	10	8
023	Other cardiovascular	27	4	23
026	Other central nervous system	3	1	2
029	Unqualified	1	0	1
042	Other salmonella infections	2	2	0
<u>045-048</u>	<u>Dysentery, all forms</u>	<u>39</u>	<u>5</u>	<u>34</u>
045	Bacillary	10	0	10
046	Amebiasis	19	5	14
048	Unspecified forms	10	0	10
049	Food poisoning	7	1	6
<u>050,051</u>	<u>Scarlet fever and streptococcal sore throat</u>	<u>3</u>	<u>2</u>	<u>1</u>
051	Streptococcal sore throat	3	2	1
053	Septicemia and pyemia	21	13	8
056	Whooping cough	14	0	14
057	Meningococcal infections	10	6	4
061	Tetanus	13	3	10
064	Other bacterial diseases	1	0	1
072	Leptospirosis icterohaemorrhagica (Weil's disease)	2	0	2
080	Acute poliomyelitis	3	1	2
082	Acute infectious encephalitis	8	7	1
085	Measles	5	4	1
087	Chickenpox	1	0	1
092	Infectious hepatitis	19	12	7
096	Other diseases attributable to viruses	3	1	2
<u>100-108</u>	<u>Typhus and other rickettsial diseases</u>	<u>1</u>	<u>1</u>	<u>0</u>
107	Typhus, unspecified	1	1	0
130	Infestation with worms of other, mixed, or unspecified type	4	0	4
132	Actinomycosis	1	1	0
134	Other fungus infections	8	4	4

Table 5 - Page 2
 Deaths - Mississippi - 1957

International List Number	Cause of Death*	Total	White	Non-white
138	Other infective and parasitic diseases	4	2	2
<u>140-205</u>	<u>Malignant neoplasm</u>	<u>2,496</u>	<u>1,482</u>	<u>1,014</u>
<u>140-148</u>	<u>Buccal cavity and pharynx</u>	<u>57</u>	<u>41</u>	<u>16</u>
140	Lip	3	3	0
141	Tongue	10	5	5
142	Salivary gland	6	6	0
143	Floor of mouth	2	2	0
144	Other parts of mouth and mouth, unspecified	18	14	4
145	Oral mesopharynx	1	1	0
146	Nasopharynx	5	3	2
148	Pharynx, unspecified	12	7	5
<u>150-159</u>	<u>Digestive organs and peritoneum</u>	<u>793</u>	<u>429</u>	<u>364</u>
150	Esophagus	35	21	14
151	Stomach	230	95	135
152	Small intestine, including duodenum	12	4	8
153	Large intestine except rectum	180	110	70
154	Rectum	64	41	23
155	Biliary passages and liver (stated to be primary site)	37	20	17
156	Liver (secondary and unspecified)	85	51	34
157	Pancreas	125	76	49
158	Peritoneum	7	3	4
159	Unspecified digestive organs	18	8	10
<u>160-165</u>	<u>Respiratory system</u>	<u>282</u>	<u>210</u>	<u>72</u>
160	Nose, nasal cavities, middle ear, and accessory sinuses	6	5	1
161	Larynx	18	13	5
162	Trachea, and bronchus and lung specified as primary	101	81	20
163	Lung and bronchus, unspecified as to whether primary or secondary	146	104	42
164	Mediastinum	2	0	2
165	Thoracic organs (secondary)	9	7	2
<u>170-181</u>	<u>Breast and genito-urinary organs</u>	<u>799</u>	<u>411</u>	<u>388</u>
170	Breast	183	111	72
171	Cervix uteri	130	56	74
172	Corpus uteri	16	9	7
174	Uterus, unspecified	86	29	57
175	Ovary, Fallopian tube, and broad ligament	63	45	18
176	Other and unspecified female genital organs	9	5	4
177	Prostate	187	89	98
178	Testis	6	6	0
179	Other and unspecified male genital organs	3	1	2
180	Kidney	39	27	12
181	Bladder and other urinary organs	77	33	44
<u>190-199</u>	<u>Other and unspecified sites</u>	<u>374</u>	<u>247</u>	<u>127</u>
190	Melanoma of skin	20	18	2
191	Skin, other	36	32	4
192	Eye	3	2	1
193	Brain and other parts of nervous system	71	56	15
194	Thyroid gland	7	4	3

Table 5 - Page 3
Deaths - Mississippi - 1957

34

International List Number	Cause of Death*	Total	White	Non-white
195	Other endocrine glands	7	6	1
196	Bone including jaw bone	22	18	4
197	Connective tissue	11	4	7
198	Lymph nodes, secondary and unspecified	3	1	2
199	Other and unspecified sites	194	106	88
<u>200-205</u>	<u>Lymphatic and hematopoietic tissues</u>	<u>191</u>	<u>144</u>	<u>47</u>
200	Lymphosarcoma and reticulosarcoma	38	32	6
201	Hodgkin's disease	18	16	2
202	Other forms of lymphoma (reticulososis)	9	7	2
203	Multiple myeloma (plasmacytoma)	25	16	9
204	Leukemia and aleukemia	99	72	27
205	Mycosis fungoides	2	1	1
<u>210-229</u>	<u>Benign neoplasm</u>	<u>24</u>	<u>6</u>	<u>18</u>
211	Other parts of digestive system	2	1	1
214	Uterine fibromyoma	7	0	7
215	Uterus, other	1	0	1
216	Ovary	2	1	1
223	Brain and other parts of nervous system	9	3	6
224	Endocrine glands	2	0	2
228	Hemangioma and lymphangioma	1	1	0
<u>230-239</u>	<u>Neoplasm of unspecified nature</u>	<u>26</u>	<u>21</u>	<u>5</u>
230	Digestive organs	4	4	0
231	Respiratory organs	3	3	0
233	Uterus	1	1	0
234	Ovary	1	1	0
237	Brain and other parts of nervous system	15	10	5
239	Other and unspecified organs	2	2	0
241	Asthma	59	30	29
242	Angioneurotic edema	1	0	1
250	Simple goiter	1	1	0
252	Thyrotoxicosis with or without goiter	8	2	6
253	Myxedema and cretinism	4	2	2
254	Other diseases of thyroid gland	1	0	1
260	Diabetes mellitus	260	140	120
271	Diseases of parathyroid gland	1	1	0
272	Diseases of pituitary gland	5	3	2
273	Diseases of thymus gland	1	1	0
281	Pellagra	3	1	2
286	Other avitaminoses and nutritional deficiency states	20	7	13
289	Other metabolic diseases	3	1	2
<u>290-293</u>	<u>Anemias</u>	<u>37</u>	<u>12</u>	<u>25</u>
290	Pernicious and other hyperchromic	3	1	2
291	Iron deficiency (hyperchromic)	1	0	1
292	Other specified type	22	7	15
293	Unspecified type	11	4	7
294	Polycythemia	5	5	0
295	Hemophilia	2	1	1
296	Purpura and other hemorrhagic conditions	7	7	0
297	Agranulocytosis	1	1	0

Table 5 - Page 4
Deaths - Mississippi - 1957

International List Number	Cause of Death*	Total	White	Non-white
298	Diseases of spleen	4	4	0
300	Schizophrenic disorders (dementia praecox)	3	2	1
301	Manic-depressive reaction	1	0	1
304	Senile psychosis	5	2	3
322	Alcoholism	19	11	8
325	Mental deficiency	2	2	0
<u>330-334</u>	<u>Vascular lesions affecting central nervous system</u>	<u>2,800</u>	<u>1,420</u>	<u>1,380</u>
330	Subarachnoid hemorrhage	63	36	27
331	Cerebral hemorrhage	1,757	843	914
332	Cerebral embolism and thrombosis	570	354	216
334	Other and ill-defined	410	187	223
340	Meningitis, except meningococcal and tuberculous	31	11	20
342	Intracranial and intraspinal abscess	8	5	3
343	Encephalitis, myelitis, and encephalomyelitis (except acute infectious)	6	2	4
344	Late effects of intracranial abscess or pyogenic infection	8	2	6
345	Multiple sclerosis	9	6	3
350	Paralysis agitans	15	13	2
351	Cerebral spastic infantile paralysis	7	6	1
352	Other cerebral paralysis	7	2	5
353	Epilepsy	36	15	21
355	Other diseases of brain	8	7	1
356	Motor neurone disease and muscular atrophy	8	5	3
357	Other diseases of spinal cord	5	2	3
364	Polyneuritis and polyradiculitis	3	1	2
366	Other and unspecified forms of neuralgia and neuritis	1	1	0
385	Cataract	1	1	0
388	Other diseases of eye	1	1	0
391	Otitis media without mention of mastoiditis	2	2	0
<u>400-402</u>	<u>Rheumatic fever</u>	<u>11</u>	<u>5</u>	<u>6</u>
400	Without mention of heart involvement	5	3	2
401	With heart involvement	6	2	4
<u>410-443</u>	<u>Diseases of heart</u>	<u>6,367</u>	<u>3,706</u>	<u>2,661</u>
410	Diseases of mitral valve	70	38	32
411	Diseases of aortic valve specified as rheumatic	7	6	1
414	Other endocarditis specified as rheumatic	12	4	8
415	Other myocarditis specified as rheumatic	5	2	3
416	Other heart diseases specified as rheumatic	52	28	24
420	Arteriosclerotic heart disease including coronary disease	3,575	2,511	1,064
421	Chronic endocarditis not specified as rheumatic	104	56	48
422	Other myocardial degeneration	603	311	292
430	Acute and subacute endocarditis	10	3	7
431	Acute myocarditis not specified as rheumatic	16	7	9

Table 5 - Page 5
Deaths - Mississippi - 1957

International List Number	Cause of Death*	Total	White	Non-white
432	Acute pericarditis specified as nonrheumatic	2	1	1
433	Functional disease of heart	34	20	14
434	Other and unspecified diseases of heart	419	179	240
441	Essential malignant hypertension with heart disease	5	1	4
442	Hypertensive heart disease with arteriolar nephrosclerosis	326	129	197
443	Other and unspecified hypertensive heart disease	1,127	410	717
<u>444-447</u>	<u>Hypertension without mention of heart</u>	<u>208</u>	<u>89</u>	<u>119</u>
444	Essential benign	37	11	26
445	Essential malignant	20	6	14
446	With arteriolar nephrosclerosis	124	61	63
447	Other hypertensive disease	27	11	16
450	General arteriosclerosis	234	139	95
<u>451-468</u>	<u>Other diseases of circulatory system</u>	<u>173</u>	<u>110</u>	<u>63</u>
451	Aortic aneurysm specified as nonsyphilitic, and dissecting aneurysm	68	47	21
452	Other aneurysm, except of heart and aorta	2	1	1
453	Peripheral vascular disease	4	2	2
454	Arterial embolism and thrombosis	3	1	2
455	Gangrene of unspecified cause	6	3	3
456	Other diseases of arteries	8	7	1
460	Varicose veins of lower extremities	1	1	0
461	Hemorrhoids	1	1	0
462	Varicose veins of other specified sites	4	4	0
463	Phlebitis and thrombophlebitis of lower extremities	8	7	1
464	Phlebitis and thrombophlebitis of other sites	4	2	2
465	Pulmonary embolism and infarction	33	18	15
466	Other venous embolism and thrombosis	17	10	7
467	Other diseases of circulatory system	13	6	7
468	Certain diseases of lymph nodes and lymph channels	1	0	1
470	Acute nasopharyngitis (common cold)	4	0	4
473	Acute sinusitis	4	1	3
474	Acute laryngitis and tracheitis	2	0	2
475	Acute upper respiratory infection of multiple or unspecified site	8	2	6
<u>480-483</u>	<u>Influenza</u>	<u>173</u>	<u>53</u>	<u>120</u>
480	With pneumonia	112	29	83
481	With other respiratory manifestations, and influenza unqualified	52	19	33
482	With digestive manifestations, but without respiratory symptoms	4	3	1
483	With nervous manifestations, but without digestive or respiratory symptoms	5	2	3
<u>490-493</u>	<u>Pneumonia</u>	<u>621</u>	<u>248</u>	<u>373</u>
490	Lobar	155	64	91

Table 5 - Page 6
Deaths - Mississippi - 1957

International List Number	Cause of Death*	Total	White	Non-white
491	Broncho	258	114	144
492	Primary atypical	41	22	19
493	Other and unspecified	167	48	119
<u>500-502</u>	<u>Bronchitis</u>	<u>32</u>	<u>17</u>	<u>15</u>
500	Acute	7	5	2
501	Unqualified	12	3	9
502	Chronic	13	9	4
510	Hypertrophy of tonsils and adenoids	4	2	2
513	Chronic sinusitis	2	0	2
517	Other diseases of upper respiratory tract	3	2	1
518	Empyema	3	0	3
519	Pleurisy	6	5	1
520	Spontaneous pneumothorax	2	0	2
521	Abscess of lung	8	3	5
522	Pulmonary congestion and hypostasis	15	10	5
525	Other chronic interstitial pneumonia	21	16	5
526	Bronchiectasis	28	25	3
527	Other diseases of lung and pleural cavity	82	63	19
532	Other inflammatory diseases of supporting structures of teeth	1	0	1
539	Diseases of esophagus	4	3	1
<u>540, 541</u>	<u>Ulcer of stomach and duodenum</u>	<u>68</u>	<u>54</u>	<u>14</u>
540	Stomach	35	24	11
541	Duodenum	33	30	3
542	Gastrojejunal ulcer	1	1	0
543	Gastritis and duodenitis	4	2	2
544	Disorders of function of stomach	5	2	3
545	Other diseases of stomach and duodenum	6	5	1
<u>550-553</u>	<u>Appendicitis</u>	<u>17</u>	<u>3</u>	<u>14</u>
550	Acute	16	3	13
551	Unqualified	1	0	1
<u>560, 561, 570</u>	<u>Hernia and intestinal obstruction</u>	<u>102</u>	<u>47</u>	<u>55</u>
560	Hernia of abdominal cavity without mention of obstruction	12	7	5
561	Hernia of abdominal cavity with obstruction	30	14	16
570	Intestinal obstruction, without mention of hernia	60	26	34
571	Gastro-enteritis and colitis, except ulcerative, age 4 weeks and over	162	30	132
572	Chronic enteritis and ulcerative colitis	19	12	7
573	Functional disorders of intestines	1	0	1
574	Anal fissure and fistula	2	0	2
575	Abscess of anal and rectal regions	2	0	2
576	Peritonitis	4	0	4
577	Peritoneal adhesions	1	0	1
578	Other diseases of intestines and peritoneum	22	13	9
580	Acute and subacute yellow atrophy of liver	3	2	1
581	Cirrhosis of liver	70	51	19
583	Other diseases of liver	6	3	3
584	Cholelithiasis	27	22	5

Table 5 - Page 7
Deaths - Mississippi - 1957

International List Number	Cause of Death*	Total	White	Non-white
585	Cholecystitis without mention of calculi	15	10	5
586	Other diseases of gallbladder and biliary ducts	8	7	1
587	Diseases of pancreas	26	19	7
<u>590,591</u>	<u>Acute nephritis and nephritis with edema</u>	<u>58</u>	<u>24</u>	<u>34</u>
590	Acute	34	12	22
591	With edema, including nephrosis	24	12	12
<u>592-594</u>	<u>Chronic and unspecified nephritis and other renal sclerosis</u>	<u>304</u>	<u>119</u>	<u>185</u>
592	Chronic nephritis	243	97	146
593	Nephritis not specified as acute or chronic	56	21	35
594	Other renal sclerosis	5	1	4
600	Infections of kidney	83	52	31
601	Hydronephrosis	3	1	2
602	Calculi of kidney and ureter	14	11	3
603	Other diseases of kidney and ureter	10	6	4
605	Cystitis	3	2	1
606	Other diseases of bladder	6	3	3
608	Stricture of urethra	5	1	4
609	Other diseases of urethra	4	1	3
610	Hyperplasia of prostate	77	34	43
611	Prostatitis	10	2	8
612	Other diseases of prostate	2	2	0
624	Salpingitis and oophoritis, unqualified	1	0	1
626	Diseases of parametrium and pelvic peritoneum (female)	2	0	2
633	Other diseases of uterus	3	0	3
637	Other diseases of female genital organs	1	1	0
<u>640-689</u>	<u>Diseases of pregnancy, childbirth, and puerperium</u>	<u>75</u>	<u>14</u>	<u>61</u>
641	Other infections of genito-urinary tract during pregnancy	1	0	1
642	Toxemias of pregnancy	19	5	14
645	Ectopic pregnancy	2	0	2
648	Other complications arising from pregnancy	1	1	0
650	Abortion without mention of sepsis or toxemia	1	0	1
651	Abortion with sepsis	4	0	4
652	Abortion with toxemia, without mention of sepsis	1	0	1
660	Delivery without stated complication	2	0	2
670	Delivery complicated by placenta previa or antepartum hemorrhage	14	2	12
671	Delivery complicated by retained placenta	7	3	4
672	Delivery complicated by other postpartum hemorrhage	10	3	7
674	Delivery complicated by disproportion or malposition of fetus	1	0	1
675	Delivery complicated by prolonged labor of other origin	1	0	1
680	Puerperal urinary infection without other sepsis	1	0	1

Table 5 - Page 8
Deaths - Mississippi - 1957

39

International List Number	Cause of Death*	Total	White	Non-white
681	Sepsis of childbirth and puerperium	1	0	1
682	Puerperal phlebitis and thrombosis	1	0	1
684	Puerperal pulmonary embolism	1	0	1
685	Puerperal eclampsia	2	0	2
687	Other forms of puerperal toxemia	5	0	5
692	Other cellulitis and abscess without mention of lymphangitis	3	0	3
698	Other local infections of skin and subcutaneous tissue	2	1	1
704	Pemphigus	1	1	0
705	Erythematous conditions	7	3	4
710	Other hypotrophic and atrophic conditions of skin	4	3	1
715	Chronic ulcer of skin	4	2	2
716	Other diseases of skin	1	1	0
720	Acute arthritis due to pyogenic organisms	1	1	0
722	Rheumatoid arthritis and allied conditions	12	12	0
723	Osteo-arthritis (arthrosis) and allied conditions	6	6	0
725	Arthritis, unspecified	13	8	5
727	Rheumatism unspecified	1	1	0
730	Osteomyelitis and periostitis	1	1	0
743	Infective myositis and other inflammatory diseases of tendon and fascia	1	1	0
744	Other diseases of muscle, tendon, and fascia	12	9	3
<u>750-759</u>	<u>Congenital malformations</u>	<u>259</u>	<u>168</u>	<u>91</u>
750	Monstrosity	11	5	6
751	Spina bifida and meningocele	17	16	1
752	Congenital hydrocephalus	29	11	18
753	Other nervous system and sense organs	6	3	3
754	Circulatory system	114	78	36
755	Cleft palate and harelip	2	0	2
756	Digestive system	38	25	13
757	Genito-urinary system	10	7	3
758	Bone and joint	6	4	2
759	Other and unspecified, not elsewhere classified	26	19	7
<u>760-762</u>	<u>Birth injuries, postnatal asphyxia, and atelectasis</u>	<u>365</u>	<u>173</u>	<u>192</u>
760	Intracranial and spinal injury at birth	59	29	30
761	Other birth injury	82	33	49
762	Postnatal asphyxia and atelectasis	224	111	113
<u>763-768</u>	<u>Infections of newborn</u>	<u>97</u>	<u>23</u>	<u>74</u>
763	Pneumonia	75	17	58
764	Diarrhea	11	1	10
767	Umbilical sepsis	1	1	0
768	Other sepsis	10	4	6
<u>769-776</u>	<u>Other diseases peculiar to early infancy, and immaturity unqualified</u>	<u>680</u>	<u>218</u>	<u>462</u>
769	Neonatal disorders arising from maternal toxemia	20	8	12

Table 5 - Page 9
Deaths - Mississippi - 1957

International List Number	Cause of Death*	Total	White	Non-white
770	Hemolytic disease of newborn (erythroblastosis)	28	17	11
771	Hemorrhagic disease of newborn	11	4	7
772	Nutritional maladjustment	27	0	27
773	Ill-defined diseases peculiar to early infancy	74	31	43
774	Immaturity with mention of any other subsidiary condition	32	19	13
776	Immaturity unqualified	488	139	349
<u>780-795</u>	<u>Symptoms, senility, and ill-defined conditions</u>	<u>1,763</u>	<u>366</u>	<u>1,397</u>
780	Certain symptoms referable to nervous system and special senses	8	2	6
781	Other symptoms referable to nervous system and special senses	2	0	2
782	Symptoms referable to cardiovascular and lymphatic system	43	15	28
783	Symptoms referable to respiratory system	5	3	2
784	Symptoms referable to upper gastro-intestinal tract	2	1	1
785	Symptoms referable to abdomen and lower gastro-intestinal system	6	1	5
786	Symptoms referable to genito-urinary system	1	0	1
788	Other general symptoms	8	1	7
790	Nervousness and debility	1	0	1
792	Uremia unqualified	8	4	4
794	Senility without mention of psychosis	74	20	54
795	Ill-defined and unknown causes	1,605	319	1,286
<u>800-962</u>	<u>Accidents</u>	<u>1,302</u>	<u>707</u>	<u>595</u>
800-802	Railway	17	6	11
<u>810-835</u>	<u>Motor vehicle</u>	<u>551</u>	<u>346</u>	<u>205</u>
810	Collision with railway train, traffic	20	19	1
812	Traffic, to pedestrian	103	47	56
813	Traffic, to pedal cyclist	4	3	1
814	Traffic, to rider or passenger of motorcycle, in collision with nonmotor vehicle or object	1	1	0
815	Traffic, to rider or passenger of motorcycle in collision with other motor vehicle	7	6	1
816	Other traffic, two or more motor vehicles	180	126	54
818	Traffic, collision with animal or animal-drawn vehicle	4	3	1
819	Traffic, collision with fixed or unspecified object	27	19	8
821	Noncollision traffic, to rider of motorcycle	1	1	0
822	Traffic, overturning in roadway	23	12	11
823	Traffic, running off roadway	123	78	45
824	Other noncollision traffic	18	5	13
825	Traffic, unspecified nature	26	19	7
830	Nontraffic, to pedestrian	12	6	6
835	Nontraffic, other and unspecified nature	2	1	1
<u>842</u>	<u>To pedestrian caused by pedal cycle</u>	<u>1</u>	<u>1</u>	<u>0</u>

Table 5 - Page 10
Deaths - Mississippi - 1957

International List Number	Cause of Death*	Total	White	Non-white
843	To rider of pedal cycle not involving collision with motor vehicle	1	1	0
845	Other nonmotor road vehicle	5	2	3
850	Submersion of occupant of small boat	19	11	8
851	Other water transport injury by submersion	3	2	1
857	Other specified water transport	4	4	0
860	To personnel in military aircraft	11	11	0
866	Other and unspecified aircraft	9	9	0
870	Poisoning by morphine and other opium derivatives	1	0	1
871	Poisoning by barbituric acid and derivatives	1	1	0
872	Poisoning by aspirin and salicylates	2	1	1
881	Poisoning by petroleum products	2	0	2
882	Poisoning by industrial solvents	1	0	1
883	Poisoning by corrosive aromatics, acids, and caustic alkalis	1	0	1
888	Poisoning by other and unspecified solid and liquid substances	4	1	3
890	Poisoning by utility (illuminating) gas	2	2	0
891	Poisoning by motor vehicle exhaust gas	1	1	0
894	Poisoning by other specified gases and vapours	3	2	1
900	Fall on stairs	6	5	1
901	Fall from ladders	1	1	0
902	Other falls from one level to another	34	21	13
903	Fall on same level	18	10	8
904	Unspecified falls	62	48	14
910	Blow from falling object	28	15	13
912	Caused by machinery	22	12	10
913	Caused by cutting and piercing instruments	4	0	4
914	Caused by electric current	20	17	3
916	Caused by fire and explosion of combustible material	134	34	100
917	Caused by hot substance, corrosive liquid, and steam	11	5	6
919	Caused by firearm	71	38	33
921	Inhalation and ingestion of food causing obstruction or suffocation	32	9	23
922	Inhalation and ingestion of other object causing obstruction or suffocation	5	2	3
923	Foreign body entering other orifice	1	1	0
924	Mechanical suffocation in bed and cradle	18	4	14
925	Mechanical suffocation in other and unspecified circumstances	17	9	8
926	Lack of care of infants under 1 year of age	4	0	4
928	Other accidents caused by animals	3	0	3
929	Drowning and submersion	96	41	55
931	Excessive heat and insolation	1	1	0
932	Excessive cold	3	2	1
933	Hunger, thirst, and exposure	4	0	4
934	Cataclysm	7	5	2
935	Lightning	11	5	6

Table 5 - Page 11
Deaths - Mississippi - 1957

42

International List Number	Cause of Death*	Total	White	Non-white
936	Other and unspecified accidents	28	13	15
941	Postvaccinal encephalitis	1	1	0
950	Therapeutic misadventure in surgical treatment	9	3	6
953	Therapeutic misadventure in administration of drugs or biologicals	2	0	2
954	Therapeutic misadventure in anesthesia	1	0	1
960	Late effects of motor vehicle accident	1	0	1
962	Late effects of other accidental injury	8	4	4
<u>963, 970-979</u>	<u>Suicide</u>	<u>163</u>	<u>143</u>	<u>20</u>
970	By analgesic and soporific substances	2	2	0
971	By other solid and liquid substances	14	11	3
973	By other gases	1	1	0
974	By hanging and strangulation	13	11	2
975	By submersion (drowning)	3	2	1
976	By firearms and explosives	126	112	14
977	By cutting and piercing instruments	1	1	0
979	By other and unspecified means	3	3	0
<u>964, 980-985</u>	<u>Homicide</u>	<u>195</u>	<u>148</u>	<u>147</u>
964	Late effect of injury purposely inflicted by another person (not in war)	1	0	1
980	Nonaccidental poisoning by another person	1	0	1
981	Assault by firearm and explosive	119	38	81
982	Assault by cutting and piercing instruments	47	6	41
983	Assault by other means	21	3	18
984	Injury by intervention of police	4	1	3
985	Execution	2	0	2

* When cause of death and International List Numbers are omitted, no deaths were reported.

TABLE 6
DEATHS
MISSISSIPPI
1957

By Place of Occurrence, Reallocated to Place of Residence
by Race, with Rates

County	Place of Occurrence			Place of Residence			Rate*		
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
STATE	20,233	10,173	10,060	20,476	10,339	10,137	9.7	9.0	10.5
Adams	453	160	293	415	154	261	11.4	8.0	15.3
Alcorn	242	194	48	248	199	49	9.9	9.2	13.6
Amite	115	45	70	179	81	98	10.6	11.2	10.1
Attala	215	130	85	231	148	83	9.9	12.0	7.6
Benton	48	25	23	80	46	34	9.4	10.7	8.1
Bolivar	548	102	446	574	145	429	9.7	8.2	10.3
Calhoun	123	90	33	141	104	37	9.3	9.9	8.0
Carroll	69	29	40	113	56	57	8.7	11.4	7.1
Chickasaw	178	93	85	179	96	83	10.4	9.8	11.2
Choctaw	92	57	35	110	74	36	11.7	11.9	11.2
Claiborne	98	24	74	124	27	97	11.1	8.7	12.0
Clarke	155	83	72	195	113	82	11.3	11.2	11.4
Clay	179	84	95	199	87	112	13.2	11.4	14.9
Coahoma	469	130	339	488	113	375	9.3	7.8	9.8
Copiah	258	108	150	302	137	165	11.0	11.0	11.0
Covington	116	71	45	152	96	56	9.9	10.1	9.6
DeSoto	179	53	126	231	87	144	9.6	10.3	9.2
Forrest	457	321	136	448	287	161	10.1	8.9	13.2
Franklin	65	33	32	87	45	42	9.9	8.5	12.0
George	92	78	14	98	84	14	9.3	9.1	10.8
Greene	62	44	18	77	57	20	9.6	8.9	12.5
Grenada	202	90	112	184	70	114	9.8	8.0	11.4
Hancock	76	61	15	111	88	23	8.5	7.4	17.7
Harrison	949	724	225	839	613	226	8.0	7.0	13.2
Hinds	1,981	1,055	926	1,433	709	724	8.4	7.2	10.0
Holmes	303	79	224	348	99	249	12.4	14.5	11.7
Humphreys	156	35	121	195	44	151	9.1	7.3	9.9
Issaquena	26	3	23	34	7	27	7.1	3.7	9.3
Itawamba	96	91	5	121	113	8	8.9	8.9	8.9
Jackson	309	198	111	340	226	114	8.0	6.6	13.4
Jasper	94	41	53	162	79	83	9.6	9.9	9.3
Jefferson	88	27	61	108	33	75	10.7	13.2	9.9
Jeff Davis	128	66	62	128	65	63	7.9	10.1	6.4
Jones	626	389	237	500	340	160	9.7	9.0	11.6
Kemper	93	38	55	122	53	69	10.2	14.3	8.4
Lafayette	179	95	84	199	117	82	12.3	14.4	10.1

Table 6 - Page 2
Deaths - Mississippi - 1957

44

County	Place of Occurrence			Place of Residence					
	Number		Non-white	Number		Non-white	Rate*		Non-white
	Total	White		Total	White		Total	White	
Lamar	72	58	14	107	85	22	8.7	8.2	11.6
Lauderdale	840	536	304	671	409	262	10.8	10.4	11.5
Lawrence	58	34	24	109	68	41	9.4	9.8	8.7
Leake	177	110	67	213	136	77	13.7	16.6	10.5
Lee	381	266	115	363	247	116	9.7	9.5	10.2
Leflore	568	207	361	517	153	364	11.2	9.7	12.0
Lincoln	232	144	88	263	165	98	10.0	9.9	10.3
Lowndes	375	177	198	393	186	207	10.5	8.9	12.5
Madison	298	87	211	315	87	228	9.7	10.7	9.4
Marion	180	106	74	204	121	83	9.0	9.1	8.8
Marshall	215	60	155	216	60	156	9.4	9.2	9.5
Monroe	262	159	103	265	165	100	8.7	8.7	8.8
Montgomery	119	62	57	147	82	65	10.6	11.1	10.1
Neshoba	175	113	62	191	128	63	10.5	10.1	11.4
Newton	156	105	51	167	106	61	8.8	9.5	7.8
Noxubee	172	58	114	187	65	122	10.9	13.5	9.8
Oktibbeha	189	79	110	200	87	113	9.1	7.8	10.5
Panola	249	92	157	303	127	176	9.4	9.8	9.2
Pearl River	170	115	55	197	134	63	9.1	8.1	12.3
Perry	37	25	12	67	49	18	9.3	9.4	9.0
Pike	387	214	173	395	224	171	12.3	12.9	11.5
Pontotoc	154	123	31	170	137	33	9.3	9.3	9.4
Prentiss	150	131	19	154	136	18	8.5	8.5	8.2
Quitman	160	31	129	198	54	144	7.8	5.7	8.9
Rankin	402	185	217	217	105	112	7.9	6.9	9.2
Scott	187	119	68	202	125	77	10.1	11.5	8.5
Sharkey	101	23	78	109	21	88	8.7	6.2	9.7
Simpson	181	109	72	180	124	56	8.5	9.5	6.9
Smith	71	55	16	94	75	19	6.9	6.7	7.3
Stone	50	32	18	46	29	17	7.2	5.9	11.3
Sunflower	366	91	275	431	124	307	8.0	7.6	8.1
Tallahatchie	260	73	187	288	90	198	10.1	9.0	10.8
Tate	123	50	73	175	78	97	10.2	11.5	9.3
Tippah	120	97	23	128	104	24	7.7	8.1	6.5
Tishomingo	110	102	8	143	133	10	11.0	10.8	14.3
Tunica	174	25	149	193	30	163	8.5	6.8	8.9
Union	205	152	53	211	156	55	11.2	10.5	13.7
Walthall	135	76	59	137	72	65	9.6	9.1	10.1
Warren	680	250	430	520	213	307	13.2	9.3	18.4
Washington	789	239	550	824	222	602	10.8	7.1	13.2
Wayne	123	81	42	160	106	54	10.3	10.9	9.3
Webster	80	61	19	102	77	25	9.7	9.6	10.0

Table 6 - Page 3
Deaths - Mississippi - 1957

County	Place of Occurrence			Place of Residence			Rate*		
	Number		Non-white	Number		Non-white	Rate*		Non-white
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Wilkinson	143	49	94	125	36	89	9.5	10.6	9.1
Winston	158	85	73	182	104	78	9.3	9.4	9.3
Yalobusha	127	67	60	147	84	63	11.3	12.3	10.2
Yazoo	353	114	239	355	128	227	9.9	9.3	10.2
Out-of-State	805	602	203	562	436	126	-	-	-

Cities over 10,000 Population**	Place of Occurrence			Place of Residence		
	Number		Non-white	Number		Non-white
	Total	White	Non-white	Total	White	Non-white
Biloxi	284	214	70	288	215	73
Clarksdale	253	101	152	210	71	139
Columbus	254	137	117	211	105	106
Greenville	455	178	277	407	138	269
Greenwood	311	137	174	223	89	134
Gulfport	356	253	103	268	180	88
Hattiesburg	376	261	115	295	172	123
Jackson	1,722	968	754	1,017	546	471
Laurel	499	293	206	254	149	105
McComb	198	133	65	143	105	38
Meridian	720	456	264	461	277	184
Natchez	258	117	141	307	120	187
Pascagoula	45	26	19	102	60	42
Tupelo	209	146	63	124	78	46
Vicksburg	580	221	359	376	164	212
Total Urban#	9,669	5,405	4,264	7,045	3,744	3,301
Total Rural	10,564	4,768	5,796	13,431	6,595	6,836

* Rate per 1,000 population.

** Cities having over 10,000 population according to 1950 census.

Includes all towns of 2,500 population or more according to 1950 census.

TABLE 7
DEATHS FROM SELECTED CAUSES
MISSISSIPPI
1957

By Age, Race, and Sex, on Basis of Place of Residence, with Rates

Inter-national List Number	Cause of Death	Total	Rate*	Total by Sex and Race	Total by Sex and Race	Age in Years	Age in Years							
							0-5 5-10 10-14 15-20 21-25 26-30 31-35 36-40 41-45 46-50 51-55 56-60 61-65 66-70 71-75 76-80 81-85 86-90 91-95 96-100 Unknown							
001-999	<u>ALL CAUSES</u>	20,476	2.7	W 10,329 C 10,137	M 6,041 F 4,298 M 5,373 F 4,764	299 295 122 724	56 51 146 117	140 21 46 22	165 149 150 185	286 281 243 275	1,031 452 800 661	3,285 2,864 2,435 2,146	1/2 1/6 9	
001-128	<u>Infective and parasitic diseases</u>	380	18.0	C 239	W 61	49	0	0	1	3	6	15	18	0
001-019	Tuberculosis (all forms)	161	7.6	C 100	M 43	12	0	3	0	1	2	3	5	0
001-008	Tuberculosis (respiratory system)	150	7.1	C 92	M 58	46	0	0	0	5	6	15	18	0
020-029	Syphilis & sequelae	50	2.4	C 35	M 15	12	0	0	0	0	0	3	4	5
045-048	Dysentery (all forms)	39	1.8	C 34	M 5	27	1	0	0	0	0	0	2	0

Table 7 - Page 2
Deaths from Selected Causes - Mississippi - 1957

Inter-national List Number	Cause of Death	Total	Rate*	Race	Total by Sex and Race	Age in Years	Unknown	
							65 and over	65 and over
050,051	Scarlet fever and streptococcal sore throat	3	0.1	W C	2 M	65-64	00000	00000
056	Whooping cough	14	0.7	W C	2 M	65-64	00000	00000
057	Meningococcal infections	10	0.5	W C	2 M	65-64	00000	00000
080	Acute poliomyelitis	3	0.1	W C	2 M	65-64	00000	00000
085	Measles	5	0.2	W C	1 M	65-64	00000	00000
100-108	Typhus and other rickettsial diseases	1	0.05	C	0 F	65-64	00000	00000

Table 7 - Page 3
Deaths from Selected Causes - Mississippi - 1957

Inter-national List Number	Cause of Death	Total	Rate*	Race	Sex	Age in Years	Total by Sex and Race		65 and over		Unknown	
							Male	Female	Male	Female	Male	Female
140-205	Malignant neoplasms, including neoplasms of lymphatic and hematopoietic tissues	2,496	118.3	W 1,482 C 1,014	M 762 F 720	1-4	5 2 2 1 1	4 2 1 1 1	18 16 6 18	93 109 82 24	412 391 251 230	0 0 0 0
210-239	Benign and Unspecified neoplasms	50	2.4	C 23	M 27 F 21	5-9	6 4 2 1 1	4 2 1 1 1	14 13 9 10 10	3 6 5 9 18	30 20 16 11 20	0 0 0 0 0
240-289	Allergic, endocrine system, metabolic and nutritional diseases	367	17.4	C 189	M 178 F 110	10-14	1 1 1 1 1	1 1 1 1 1	85 104 68 103	20 16 11 20	64 28 49 0	0 0 0 0 0
260	Diabetes mellitus	260	12.3	C 120	M 140 F 86	15-19	0 0 0 0 0	0 0 0 0 0	54 40 20 17 12	12 13 7 18 18	30 27 41 3 1	0 0 0 0 0
290-299	Diseases of the blood and blood forming organs	56	2.6	C 26	M 20 F 12	20-24	0 0 0 0 0	0 0 0 0 0	4 8 11 14	3 1 2 2	9 10 2 0	1 1 0 0 0
290-293	Anemias	37	1.8	C 25	M 26 F 21	25-29	0 0 0 0 0	0 0 0 0 0	14 11 2 2	0 0 0 0 0	0 0 0 0 0	1 1 0 0 0

Table 7 - Page 4
Deaths from Selected Causes - Mississippi - 1957

Inter-national List Number	Cause of Death	Total	Rate*	Total by Race	Total by Sex and Race	Total by Age and Race	Age in Years						
							1-4	5-9	10-14	15-24	25-34	35-44	45-54
300-326	Mental, psycho-neurotic, and personality disorders	30	1.4	W 17 C 13	M F	116 94	116 94	116 94	116 94	116 94	116 94	116 94	116 94
330-334	Vascular lesions affecting central nervous system	2,800	132.7	W 1,420 C 1,380	M F	741 679	741 679	741 679	741 679	741 679	741 679	741 679	741 679
340-398	Other diseases of the nervous system and sense organs	156	7.4	W 82 C 74	M F	44 33	44 33	44 33	44 33	44 33	44 33	44 33	44 33
400-468	Nonmeningococcal meningitis	31	1.5	W 11 C 20	M F	6 5	6 5	7 7	7 7	7 7	7 7	7 7	7 7
400-468	Diseases of the circulatory system	6,923	331.4	W 4,042 C 2,944	M F	2,428 1,621	2,428 1,621	1,557 1,387	1,557 1,387	2 0	0 0	0 0	0 0
400-468	Rheumatic fever	11	0.5	C 6	M F	5 2	5 2	2 0	2 0	1 0	0 0	0 0	0 0

Table 7 - Page 5
Deaths from Selected Causes - Mississippi - 1957

Inter-national List Number	Cause of Death	Total	Rate*	Total by Sex and Race	Total by Sex and Race	Age in Years							
						1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64
410-416	Chronic rheumatic heart disease	146	6.9	W 78 M 40	M 38 F 40	0 0	0 0	0 0	1 1	8 7	4 3	8 7	11 0
				C 68 F 35	L 33 F 35	0 0	0 0	0 0	1 1	2 1	9 3	6 4	23 0
420-422	Arteriosclerotic and degenerative heart disease	4,282	203.0	W 2,878 C 1,404	M 1,805 F 1,073	0 0	0 0	0 0	0 0	2 2	3 2	33 28	125 111
				F 633	M 771	0 0	0 0	0 0	1 0	4 11	29 29	74 74	916 156
430-434	Other diseases of the heart	481	22.8	W 210 C 271	M 133 F 77	0 2	0 2	1 1	1 1	2 1	5 5	13 11	89 53
				C 271	M 139 F 132	0 1	0 1	0 0	2 2	2 2	6 6	22 12	26 15
440-443	Hypertension with heart disease	1,458	69.1	W 540 C 918	M 259 F 281	0 0	0 0	0 0	0 0	0 0	0 0	4 3	21 12
				C 918	M 477 F 441	0 0	0 0	0 0	0 0	0 0	0 0	22 20	43 41
444-447	Hypertension without mention of heart	208	9.8	W 89 C 119	M 50 F 39	0 0	0 0	0 0	0 0	0 0	0 0	1 1	4 2
				C 119	M 61 F 58	0 0	0 0	0 0	0 0	0 0	0 0	8 7	6 5
470-527	Diseases of the respiratory system	1,018	48.2	W 447 C 571	M 272 F 273	26 21	21 20	3 2	3 2	3 2	3 2	2 1	176 124
				C 571	F 273	104 102	102 22	17 13	17 13	17 13	17 13	83 92	83 92

Table 7 - Page 6
Deaths from Selected Causes - Mississippi - 1957

International List Number	Cause of Death	Total	Rate*	Race	Total by Sex and Race	Age in Years	Unknown							
							5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 and over
480-483	Influenza	173	8.2	C	53	25	3	0	0	0	0	0	17	0
						28	2	0	0	1	1	0	22	0
						58	18	4	0	3	4	7	22	0
						62	16	3	0	2	0	4	28	0
490-493	Pneumonia (excluding pneumonia of newborn)	621	29.4	C	248	141	15	0	1	1	1	5	102	0
						107	10	1	1	2	1	4	82	1
						66	20	3	1	1	1	13	54	0
						77	18	1	1	2	3	10	58	0
500-502	Bronchitis	32	1.5	C	173	12	0	1	0	0	0	1	5	0
						5	2	2	0	0	0	0	0	0
						9	6	1	0	0	0	1	0	0
						6	2	1	0	0	0	0	1	0
530-587	Diseases of the digestive system	276	27.3	C	286	172	4	2	1	3	1	41	20	0
						114	5	2	1	3	1	15	58	0
						156	67	5	1	2	1	22	37	0
						134	56	6	1	3	1	7	10	28
540,541	Ulcer of stomach and duodenum	68	3.2	C	290	43	0	0	0	0	0	9	12	0
						11	0	0	0	0	0	3	3	4
						3	0	0	0	0	0	0	1	2
550-553	Appendicitis	17	0.8	C	14	1	1	0	0	0	0	0	0	0
						1	2	0	0	0	0	0	0	0
						9	0	0	0	0	0	0	0	0
						5	0	0	0	0	0	0	0	0

Table 7 - Page 7
Deaths from Selected Causes - Mississippi - 1957

Table 7 - Page 8
Deaths from Selected Causes - Mississippi - 1957

International List Number	Cause of Death	Total	Rate*	Race	Total by Sex and Race		Total by Age in Years		Total by Sex and Race		Total by Age in Years	
					M	F	W	M	Yr	65 and over		
640-689*	Complications of pregnancy, childbirth, and the puerperium	75	1.2	W	14	M	14	1	23	10	134	67
690-716	Diseases of the skin and cellular tissue	22	1.0	C	6	F	6	2	23	10	2354	327
720-749	Diseases of the bones & organs of movement	47	2.2	W	11	M	11	2	21	10	110	57
750-759	Congenital malformations	259	12.3	C	29	F	29	1	21	10	42	21
760-776*	Certain diseases of early infancy	1,142	18.8	W	414	M	414	1	21	10	354	172
780-795	Symptoms, senility, and ill-defined conditions	1,763	83.6	C	728	F	728	1	21	10	134	67

Table 7 - Page 9
Deaths from Selected Causes - Mississippi - 1957

International List Number	Cause of Death	Total	Rate*	Total by Sex and Race	Total by Sex and Race	Age in Years						
						5-9	10-14	15-24	25-34	35-44	45-54	55-64
800-929	<u>Accidents, poisonings and violence</u>	1,660	78.7	W 898	M 662	17	20	16	34	102	91	78
				F 236	M 225	6	13	8	25	27	19	18
				F 237	M 237	32	20	22	24	86	67	26
				F 22	M 22	22	29	14	5	23	16	12
810-835	Motor vehicle accidents	551	26.1	W 346	M 257	2	12	5	12	59	53	28
				F 89	M 89	1	2	3	7	19	11	13
				F 149	M 149	0	5	8	4	31	18	11
				F 56	M 56	0	6	2	0	13	8	6
800-802, 840-962	All other accidents	751	35.6	W 361	M 259	15	8	11	20	42	31	39
				F 102	M 102	5	11	5	1	2	4	5
				F 239	M 239	32	24	13	20	34	21	21
				F 151	M 151	32	33	12	4	11	3	7
963, 970-979	Suicide	163	7.7	W 143	M 110	0	0	0	1	4	8	15
				F 33	M 33	0	0	0	0	3	3	7
				F 19	M 19	1	0	0	0	2	1	1
				F 1	M 1	0	0	0	0	0	0	0
964, 980-985	Homicide	195	9.2	W 48	M 36	0	0	0	1	5	10	9
				F 12	M 12	0	0	0	0	4	4	4
				F 118	M 118	0	1	1	0	22	32	27
				F 29	M 29	0	0	0	1	12	11	1

* Crude death rate per 1,000 population. Specific death rates per 100,000 population. Infant causes (760-776) per 1,000 live births.

TABLE 8
DEATHS FROM SELECTED CAUSES
MISSISSIPPI I
1957

By Month of Occurrence on Basis of Place of Residence

Cause of Death and International List Number	Total	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
<u>TOTAL ALL CAUSES</u>	<u>20,476</u>	<u>1,903</u>	<u>1,469</u>	<u>1,778</u>	<u>1,622</u>	<u>1,570</u>	<u>1,610</u>	<u>1,654</u>	<u>1,566</u>	<u>1,483</u>	<u>1,904</u>	<u>1,830</u>	<u>2,087</u>
Dysentery (045-048)	39	4	4	3	2	4	3	3	1	3	7	2	3
Whooping cough (056)	14	6	0	0	0	1	2	0	0	4	0	0	1
Acute poliomyelitis (080)	3	0	1	0	0	0	1	0	1	0	0	0	0
Measles (085)	5	0	0	1	1	0	1	0	0	0	0	1	1
Influenza (480-483)	173	16	10	12	5	4	3	5	8	8	10	17	18
Lobar pneumonia (490)	155	28	9	21	8	10	8	10	8	5	5	17	11
Bronchopneumonia (491)	258	40	23	26	21	16	21	12	18	10	10	24	21
Other & unspec. pneumonia (492,493)	208	32	22	16	18	5	13	11	9	11	11	17	24
Diarrhea & enteritis (-2) (5710)	132	18	11	8	6	10	19	11	8	10	5	10	16
Diarrhea & enteritis (f2) (5711)	30	5	3	1	1	4	3	6	2	2	1	0	2
Accidents (800-962)	1,302	115	73	98	106	97	117	132	100	96	99	122	147
Motor vehicle accidents (810-835)	551	42	38	36	41	36	48	60	28	55	46	50	71

TABLE 9
DEATHS UNDER 1 YEAR
MISSISSIPPI
1957

For Selected Causes, by Race, by Age, on Basis of Place of Residence, with Rates

Cause of Death and International List Number	Total	Rate*	Total by Race	DAYS							MONTHS										
				Under 1 day	1- 6	7- 13	14- 20	21- 27	28 days	Under 1 day	1	2	3	4	5	6	7	8			
<u>All causes</u>	<u>2,257</u>	<u>37.2</u>	<u>W</u> <u>C</u> <u>694</u> <u>1,563</u>	<u>266</u> <u>329</u>	<u>221</u> <u>203</u>	<u>27</u> <u>23</u>	<u>17</u> <u>51</u>	<u>9</u> <u>39</u>	<u>Under 28 days</u>	<u>540</u> <u>385</u>	<u>36</u> <u>150</u>	<u>23</u> <u>112</u>	<u>19</u> <u>21</u>	<u>18</u> <u>23</u>	<u>11</u> <u>52</u>	<u>19</u> <u>48</u>	<u>7</u> <u>34</u>	<u>3</u> <u>24</u>	<u>6</u> <u>28</u>	<u>5</u> <u>28</u>	<u>7</u> <u>28</u>
Syphilis (020-029)	1	0.02	W C 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	
Diphtheria (055)	0	-	W C 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	
Whooping cough (056)	9	0.1	W C 9	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 1 1	0 1 1	0 1 2	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	
Measles (085)	1	0.02	W C 1	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	
Influenza (480-483)	39	0.6	W C 5	34 1 0	1 5 0	1 1 0	0 1 0	1 1 0	1 1 0	9 5 0	2 2 0	0 1 0	1 1 0	0 1 0	1 3 0	3 2 0	2 3 0	2 1 0	1 1 0	1 1 0	
**Pneumonia (490-493)	168	2.8	W C 25	143 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	4 28 18	4 20 23	4 11 11	4 1 9	1 3 7	1 3 7	2 2 5	2 1 5	1 1 8	0 0 7	
Other respiratory diseases (470-475, 500-527)	46	0.8	W C 10	29 3 0	2 8 0	3 0 0	0 1 0	1 1 0	1 1 0	13 12 0	0 2 1	0 3 1	0 3 1	0 1 0	0 3 2	0 2 0	2 1 0	2 1 0	2 2 0		
**Diarrhea and enteritis (5710)	123	2.0	W C 113	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 1 0	1 0 0	1 1 0	1 0 0	1 1 0	1 0 0	1 1 0	1 0 0		

Table 9 - Page 2
Deaths Under 1 Year - Mississippi - 1957

Cause of Death and International List Number	Total	Rate*	Total by Race	Total by Race	DAYS							MONTHS								
					Under 1 day	1- 6	7- 13	14- 20	21- 27	28 days	Under	1	2	3	4	5	6	7	8	
Congenital malfor- mations (750-759)	191	3.1	W C	126 65	37	30	7	8	4	41	86	9	6	2	5	5	2	1	1	2
Birth injuries (760, 761)	141	2.3	W C	62 79	32	26	2	0	1	0	60	1	1	0	0	0	0	1	2	0
Postnatal asphyxia & atelectasis (762)	224	3.7	W C	111 113	64	39	4	0	2	1	109	2	0	0	0	0	0	0	0	0
** Infections of the newborn (763-768)	97	1.6	W C	23 74	2	13	5	3	0	10	23	0	0	0	0	0	0	0	0	0
Hemolytic disease of the newborn (770)	28	0.5	W C	17 11	8	5	0	0	0	2	16	1	0	0	0	0	0	0	0	0
All other defined dis. of early infancy (769, 771, 772)	58	1.0	W C	12 46	5	7	0	0	1	24	7	7	1	2	3	0	0	0	0	0
Violent & accidental deaths (800-999)	87	1.4	W C	23 64	1	4	1	1	0	2	14	5	2	2	2	0	1	1	0	1
Other specified causes	109	1.8	W C	42 67	3	7	3	2	2	17	4	2	5	2	3	3	1	2	1	1
III-defined (773-776, 780-793, 795)	935	15.4	W C	221 714	112	76	5	2	1	196	7	8	4	1	2	1	0	0	0	1
					258	169	43	15	14	499	53	47	31	15	14	22	7	11	4	6

* Rate per 1,000 live births.

** Pneumonia and diarrhea of the newborn (under 28 days) included in infections of the newborn (763-768).

TABLE 10
DEATHS DUE TO INJURIES WHICH OCCURRED IN MISSISSIPPI
1957

By Place of Injury by Type of Injury

County	Motor Vehicle Accidents (810-835)			Other Transport Accidents (800-802, 840-866)			Non-Transport Accidents (870-936)			Other and Violences (970-985)			Total	Urban	Rural
	Total	Urban	Rural	Total	Urban	Rural	Total	Urban	Rural	Total	Urban	Rural			
STATE	575	66	509	58	10	48	72	25	37	357	134	223			
Adams	10	7	3	8	0	0	0	0	0	0	1	14	12	2	6
Alcorn	7	2	5	6	0	0	0	0	0	0	0	7	1	0	4
Anite	5	0	1	5	0	0	0	0	0	0	0	4	0	2	2
Attala	3	1	0	2	0	0	0	0	0	0	1	6	0	0	4
Benton	5	8	5	8	0	1	0	0	1	0	1	5	1	0	4
Bolivar	8	0	0	0	0	0	0	0	0	0	0	1	0	0	1
Calhoun	3	5	0	3	0	0	0	0	0	0	1	0	0	0	0
Carroll	5	2	3	2	1	0	0	0	0	0	0	3	0	0	3
Chickasaw	2	1	1	1	0	1	0	0	0	0	0	0	0	0	0
Choctaw	1	1	0	1	0	1	0	0	0	0	0	0	0	0	0
Claiborne	3	3	0	3	0	1	0	0	1	0	1	0	0	0	0
Clarke	7	7	0	7	0	1	0	0	1	0	1	3	0	0	3
Clay	5	5	0	5	0	5	0	0	0	0	1	8	0	0	3
Coahoma	13	2	11	11	0	11	0	0	1	0	1	5	0	0	5
Copiah	7	0	7	2	0	0	0	0	0	0	0	0	0	0	0
Covington	2	0	2	2	0	0	0	0	0	0	0	0	0	0	0
DeSoto	9	7	9	6	0	1	0	0	1	0	0	3	0	0	2
Forrest	7	0	7	0	0	1	0	0	1	0	0	0	0	0	0
Franklin	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0
George	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Greene	3	0	3	0	0	0	0	0	0	0	0	0	0	0	0
Grenada	6	5	1	5	1	1	0	0	0	0	0	4	2	0	2
Hancock	2	1	1	1	1	0	0	0	0	0	0	1	0	0	1
Harrison	23	6	17	3	0	0	0	0	0	0	0	0	0	0	0
													3	0	3

Table 10 - Page 2
Deaths Due to Injuries which Occurred in Mississippi - 1957

County	Motor Vehicle Accidents (810-835)			Other Transport Accidents (800-802, 840-866)			Non-Transport Accidents (870-926)			Other and Vicencies (970-985)		
	Total	Urban	Rural	Total	Urban	Rural	Total	Urban	Rural	Total	Urban	Rural
Hinds	26	10	16	5	2	4	1	2	1	31	7	24
Holmes	10	1	9	1	0	1	0	0	0	0	3	2
Humphreys	5	1	4	1	0	1	0	0	0	0	4	2
Issaquena	1	0	1	0	0	0	0	0	0	0	0	0
Itawamba	1	0	1	0	0	0	0	0	0	0	0	0
Jackson	13	5	8	0	0	0	0	0	0	0	0	0
Jasper	2	0	2	0	0	0	0	0	0	0	0	0
Jefferson	1	0	1	0	0	0	0	0	0	0	0	0
Jeff Davis	4	0	4	0	0	0	0	0	0	0	0	0
Jones	24	5	19	0	0	0	0	0	0	0	0	0
Kemper	5	0	5	0	0	0	0	0	0	0	0	0
Lafayette	10	0	10	0	0	0	0	0	0	0	0	0
Lamar	4	0	4	0	0	0	0	0	0	0	0	0
Lauderdale	17	3	14	8	3	5	0	0	0	0	0	0
Lawrence	8	0	8	0	0	0	0	0	0	0	0	0
Leake	3	0	3	0	0	0	0	0	0	0	0	0
Lee	9	3	6	1	0	1	0	0	0	0	0	0
Leflore	11	0	10	9	0	9	0	0	0	0	0	0
Lincoln	9	0	9	0	0	0	0	0	0	0	0	0
Lowndes	14	6	13	5	2	3	0	0	0	0	0	0
Madison	21	6	20	5	1	4	0	0	0	0	0	0
Marion	9	0	9	0	0	0	0	0	0	0	0	0
Marshall	11	0	11	0	0	0	0	0	0	0	0	0
Monroe	0	0	0	0	0	0	0	0	0	0	0	0
Montgomery	5	2	5	2	1	2	0	0	0	0	0	0
Neshoba	7	2	5	2	1	2	0	0	0	0	0	0
Newton	9	2	8	2	1	2	0	0	0	0	0	0
Noxubee	2	1	1	0	0	0	0	0	0	0	0	0
Oktibbeha	5	2	4	2	1	2	0	0	0	0	0	0
Panola	6	2	4	2	1	2	0	0	0	0	0	0

Table 10 - Page 3
Deaths Due to Injuries which Occurred in Mississippi - 1957

County	Motor Vehicle Accidents (810-835)		Other Transport Accidents (800-802, 840-866)		Non-Transport Accidents (870-936)		Other and Violences (970-985)		Violences (970-985)	
	Total	Urban	Rural	Total	Urban	Rural	Total	Urban	Total	Rural
Pearl River	9	1	8	1	0	1	2	1	2	0
Perry	4	0	4	0	0	0	1	0	1	1
Pike	11	2	9	1	0	1	2	1	6	4
Pontotoc	7	0	7	0	0	0	4	2	4	1
Prentiss	7	1	6	0	0	0	2	0	2	0
Quitman	3	0	3	0	0	0	4	0	4	4
Rankin	11	6	11	0	0	0	5	0	5	0
Scott	6	5	6	0	0	0	0	0	0	0
Sharkey	5	5	6	0	0	0	0	0	0	0
Simpson	6	6	5	0	0	0	0	0	0	0
Smith	5	5	6	0	0	0	0	0	0	0
Stone	6	6	0	0	0	0	0	0	0	0
Sunflower	17	10	8	2	2	2	16	9	16	1
Tallahatchie	8	6	6	1	1	1	1	1	1	1
Tate	2	0	2	0	0	0	0	0	0	0
Tippah	2	0	2	0	0	0	0	0	0	0
Tishomingo	6	0	6	0	0	0	0	0	0	0
Tunica	9	9	0	0	0	0	0	0	0	0
Union	6	6	0	0	0	0	0	0	0	0
Walton	2	2	0	0	0	0	0	0	0	0
Warren	3	1	2	1	0	1	3	3	3	3
Washington	17	2	15	8	1	0	2	0	2	0
Wayne	8	1	0	0	0	0	0	0	0	0
Webster	1	0	1	0	0	0	2	0	2	0
Wilkinson	2	0	2	0	0	0	0	0	0	0
Winston	8	4	3	0	0	0	3	3	3	3
Yalobusha	3	0	3	0	0	0	0	0	0	0
Yazoo	6	3	3	0	0	0	0	0	0	0

TABLE 11
MARRIAGES AND DIVORCES
MISSISSIPPI
1957

By County, by Race of Bride

County	Marriages						Divorces		
	Place of Licensure			Residence of Bride			Total	White	Non-white
	Total	White	Non-white	Total	White	Non-white			
STATE	62,495	46,868	15,627	20,864	12,045	8,819	4,974	3,350	1,624
Adams	816	621	195	344	187	157	162	100	62
Alcorn	7,314	6,598	716	264	220	44	60	49	11
Amite	260	186	74	124	68	56	28	17	11
Attala	217	104	113	243	131	112	53	33	20
Benton	201	70	131	68	37	31	4	2	2
Bolivar	573	215	358	545	226	319	82	41	41
Calhoun	112	86	26	117	84	33	17	11	6
Carroll	90	48	42	90	47	43	17	7	10
Chickasaw	197	121	76	178	93	85	28	22	6
Choctaw	78	44	34	70	39	31	14	9	5
Claiborne	161	92	69	72	22	50	29	8	21
Clarke	406	333	73	144	96	48	28	18	10
Clay	286	175	111	156	67	89	33	15	18
Coahoma	786	340	446	510	145	365	71	45	26
Copiah	237	115	122	249	123	126	58	25	33
Covington	183	141	42	173	128	45	36	26	10
DeSoto	8,773	6,174	2,599	237	84	153	29	18	11
Forrest	526	399	127	513	378	135	189	136	53
Franklin	82	62	20	67	45	22	17	8	9
George	2,046	1,837	209	113	102	11	12	11	1
Greene	506	448	58	88	60	28	15	9	6
Grenada	222	118	104	182	89	93	29	18	11
Hancock	1,224	1,163	61	104	82	22	51	44	7
Harrison	1,993	1,721	272	1,053	828	225	558	484	74
Hinds	1,569	929	640	1,782	1,112	670	414	288	126
Holmes	286	73	213	330	83	247	24	10	14
Humphreys	183	71	112	195	71	124	25	14	11
Issaquena	25	11	14	29	4	25	5	3	2
Itawamba	1,394	1,373	21	102	96	6	15	15	0
Jackson	2,559	1,716	843	536	419	117	133	115	18
Jasper	135	70	65	186	92	94	12	5	7
Jefferson	336	296	40	62	20	42	25	6	19
Jeff Davis	202	121	81	105	55	50	20	14	6
Jones	666	510	156	586	460	126	177	125	52
Kemper	134	66	68	129	64	65	13	8	5
Lafayette	125	80	45	132	80	52	29	17	12

Table 11 - Page 2
Marriages and Divorces - Mississippi - 1957

62

County	Marriages						Divorces		
	Place of Licensure			Residence of Bride			Total	White	Non-white
	Total	White	Non-white	Total	White	Non-white			
Lamar	282	245	37	162	129	33	33	26	7
Lauderdale	1,976	1,292	684	687	446	241	185	136	49
Lawrence	111	72	39	170	98	72	24	17	7
Leake	183	118	65	191	113	78	36	18	18
Lee	552	442	110	444	331	113	95	80	15
Leflore	527	177	350	494	158	336	95	49	46
Lincoln	292	216	76	226	168	58	55	38	17
Lowndes	3,370	2,718	652	398	202	196	101	63	38
Madison	396	117	279	292	51	241	88	19	69
Marion	362	238	124	290	194	96	83	58	25
Marshall	1,023	746	277	181	61	120	34	9	25
Mohrue	477	354	123	299	181	118	86	60	26
Montgomery	126	87	39	112	69	43	23	16	7
Neshoba	258	192	66	221	172	49	55	39	16
Newton	197	128	69	198	111	87	32	16	16
Noxubee	285	97	188	179	57	122	26	7	19
Oktibbeha	469	330	139	211	81	130	25	13	12
Panola	374	230	144	255	118	137	54	32	22
Pearl River	968	790	178	284	218	66	121	100	21
Perry	131	106	25	85	76	9	18	13	5
Pike	600	400	200	374	201	173	106	67	39
Pontotoc	203	165	38	155	129	26	43	33	10
Prentiss	584	554	30	173	148	25	49	39	10
Quitman	300	148	152	246	97	149	40	23	17
Rankin	440	349	91	211	139	72	43	31	12
Scott	272	175	97	211	132	79	51	41	10
Sharkey	124	39	85	84	27	57	11	7	4
Simpson	226	163	53	190	138	52	47	37	10
Smith	133	113	20	139	112	27	31	27	4
Stone	122	108	14	59	47	12	14	9	5
Sunflower	640	366	274	442	144	298	51	34	17
Tallahatchie	344	127	217	244	108	136	38	29	9
Tate	835	709	126	172	71	101	21	9	12
Tippah	185	153	32	144	118	26	30	23	7
Tishomingo	3,284	3,008	286	142	137	5	41	39	2
Tunica	511	296	215	186	31	155	16	9	7
Union	277	238	39	157	125	32	34	28	6
Walthall	335	202	133	119	51	68	23	20	3
Warren	689	506	183	373	222	151	165	83	82
Washington	3,210	2,504	706	776	344	432	165	80	85
Wayne	354	238	116	157	113	44	33	22	11
Webster	111	81	30	117	89	28	13	10	3
Wilkinson	762	651	111	112	39	73	24	8	16
Winston	218	129	89	187	108	79	31	20	11
Yalobusha	133	80	53	124	64	60	12	8	4
Yazoo	341	144	197	313	140	173	61	29	32
Unstated				121	66	55			
Cut-of-state residents				41,510	34,757	6,753			

TABLE 12
DIVORCES
MISSISSIPPI
1957

By Cause of Divorce

Cause	Total	White	Non-white
<u>TOTAL</u>	<u>4,974</u>	<u>3,350</u>	<u>1,624</u>
Cruel and inhuman treatment	2,405	1,947	458
Desertion	1,911	934	977
Adultery	235	129	106
Drunkenness	293	246	47
Penitentiary	25	19	6
Insanity	2	2	0
Annulment	25	21	4
Bigamy	9	7	2
Miscellaneous and other	41	26	15
Not stated	28	19	9

By Number of Minor Children

Minor Children	Total	White	Non-white
None	2,683	1,685	998
One	1,044	794	250
Two	596	445	151
Three or more	587	384	203
Not stated	64	42	22

By Number of Years Married

Number Years Married	Total	White	Non-white
<u>TOTAL</u>	<u>4,974</u>	<u>3,350</u>	<u>1,624</u>
Under one year	398	367	31
One year	581	480	101
Two years	470	365	105
Three years	368	276	92
Four years	317	236	81
5-9 years	1,040	695	345
10-19 years	1,007	605	402
Twenty years and over	712	274	438
Not stated	81	52	29

TABLE 13
BIRTHS, DEATHS, MARRIAGES
MISSISSIPPI
1957

By Month of Occurrence

<u>Month</u>	<u>Births</u>	<u>Deaths</u>	<u>Marriages</u>
<u>TOTAL</u>	<u>60,712</u>	<u>20,476</u>	<u>62,495</u>
January	5,320	1,903	4,598
February	4,541	1,469	4,382
March	5,108	1,778	5,093
April	4,388	1,622	4,745
May	4,424	1,570	5,181
June	4,904	1,610	6,391
July	5,563	1,654	5,552
August	5,689	1,566	6,063
September	5,478	1,483	5,024
October	5,289	1,904	4,491
November	4,849	1,830	5,115
<u>December</u>	<u>5,159</u>	<u>2,087</u>	<u>5,860</u>

TABLE 14
LIVE BIRTHS, DEATHS, MATERNAL DEATHS, AND DEATHS UNDER 1 YEAR OF AGE
MISSISSIPPI
1948-1957
Shown by Year and Race, on Basis of Place of Residence, with Rates*

YEAR	LIVE BIRTHS				DEATHS				MATERNAL DEATHS				DEATHS UNDER 1 YEAR				DEATHS UNDER 1 MONTH			
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White
1948 No. Rate	65291 28.0	29666 24.9	35625 31.2	20183 8.6	9216 7.7	10967 9.6	177 2.7	38 1.3	139 3.9	2470 37.8	988 33.3	1482 41.6	1674 25.6	773 25.6	901 26.0	25.3	25.3	25.3	25.3	
1949 No. Rate	66128 28.1	28786 24.0	37342 32.5	20489 8.7	9233 7.7	11256 9.8	147 2.2	32 1.1	115 3.1	2612 39.5	900 31.3	1712 45.8	1594 24.1	661 23.0	933 25.0	933 25.0	933 25.0	933 25.0		
1950 No. Rate	64427 29.6	27188 22.9	37239 37.6	20567 9.4	9466 8.0	11101 11.2	165 2.6	30 1.1	135 3.6	2348 36.4	763 28.1	1585 42.6	1446 22.4	560 20.6	886 23.8	886 23.8	886 23.8	886 23.8		
1951 No. Rate	65512 30.1	27798 23.4	37714 38.1	21007 9.6	9593 8.1	11414 11.5	143 2.2	28 1.0	115 3.0	2589 39.5	814 29.3	1775 47.1	1679 25.6	626 22.5	1053 22.5	1053 22.5	1053 22.5	1053 22.5		
1952 No. Rate	64509 29.6	28073 23.6	36436 36.8	20657 9.5	9387 7.9	11270 11.4	119 1.8	15 0.5	104 2.8	2592 40.2	760 27.1	1832 50.3	1658 25.7	581 20.7	1077 20.7	1077 20.7	1077 20.7	1077 20.7		
1953 No. Rate	64061 29.4	28312 23.8	35749 36.1	20492 9.4	9495 8.0	10997 11.1	119 1.8	24 0.8	95 2.6	2534 39.6	807 28.5	1727 48.3	1630 25.4	635 22.4	995 22.4	995 22.4	995 22.4	995 22.4		
1954 No. Rate	64748 29.7	28781 24.2	35967 36.3	19353 8.9	9453 8.0	9900 10.0	104 1.6	11 0.4	93 2.6	2360 36.4	744 25.8	1616 44.9	1521 23.5	601 20.9	920 20.9	920 20.9	920 20.9	920 20.9		
1955 No. Rate	63244 29.0	27546 23.2	35698 36.0	19403 8.9	9501 8.0	9902 10.0	76 1.2	12 0.4	64 1.8	2338 37.0	680 24.7	1658 46.4	1440 22.8	527 19.1	913 19.1	913 19.1	913 19.1	913 19.1		
1956 No. Rate	62178 29.5	27278 23.8	34900 36.2	19780 9.4	9969 8.7	9811 10.2	74 1.2	12 0.4	62 1.8	2353 37.8	734 26.9	1619 46.4	1451 23.3	581 21.3	870 21.3	870 21.3	870 21.3	870 21.3		
1957 No. Rate	60712 28.8	26804 22.4	33908 35.2	20476 9.7	10339 9.0	10137 10.5	75 1.2	14 0.5	61 1.8	2257 27.2	694 25.9	1563 46.1	1425 23.5	540 20.1	885 20.1	885 20.1	885 20.1	885 20.1		

* Population for 1947, 1948 and 1949, estimates as of July 1, based on 1930 and 1940 censuses; for 1950-1955, the April 1, 1950, United States Census figures; 1956-57, estimate as of July 1, prepared by Dr. Harold Pedersen, Sociology Department, Mississippi State College, with adjustments by Statistical Service, Mississippi State Board of Health. Birth and death rates per 1,000 population. Maternal and infant death rates per 1,000 live births.

TABLE 15
DEATHS FROM HEART DISEASES, NEPHRITIS (ALL FORMS),
VASCULAR LESIONS, CANCER, AND TUBERCULOSIS (ALL FORMS)
MISSISSIPPI
1948-1957

Shown by Year and Race, on Basis of Place of Residence, with Rates*

YEAR	HEART DISEASES			NEPHRITIS (ALL FORMS)			VASCULAR LESIONS			CANCER			TUBERCULOSIS (ALL FORMS)		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
1948 No. Rate	4,339 186.1	2,460 206.7	1,879 164.6	1,625 69.7	666 56.0	959 83.9	1,956 73.8	878 94.5	1,078 81.8	1,906 1,205	1,148 86.8	758 72.6	668 24.7	206 15.8	462 17.3
1949 No. Rate	5,496 233.9	3,060 254.9	2,436 212.0	729 31.0	236 19.7	493 42.9	2,093 89.1	976 81.3	1,117 97.2	2,039 86.8	1,205 100.4	834 72.6	581 24.7	190 15.8	391 34.0
1950 No. Rate	5,546 254.5	3,158 265.7	2,388 241.1	733 33.6	246 20.7	487 49.2	2,200 101.0	971 81.7	1,229 124.1	2,143 98.4	1,333 112.2	810 81.8	558 25.6	179 15.1	379 38.3
1951 No. Rate	5,626 258.2	3,134 263.7	2,492 251.6	681 31.2	210 17.7	471 47.6	2,344 107.6	1,089 91.6	1,255 126.7	2,186 100.3	1,326 111.6	860 86.8	497 22.8	158 22.8	339 13.3
1952 No. Rate	5,876 269.7	3,256 273.9	2,620 264.6	658 30.2	214 18.0	444 44.8	2,386 109.5	1,098 92.4	1,288 130.1	2,189 100.5	1,285 108.1	904 91.3	418 19.2	133 11.2	285 28.8
1953 No. Rate	5,758 264.3	3,183 267.8	2,575 260.0	474 21.8	164 13.8	310 31.3	2,404 110.3	1,125 94.6	1,279 129.2	2,230 102.3	1,348 113.4	882 89.1	308 14.1	85 7.2	223 22.5
1954 No. Rate	5,708 262.0	3,325 279.7	2,383 240.6	448 20.6	169 14.2	279 28.2	2,422 111.2	1,176 98.9	1,246 125.8	2,332 107.0	1,420 119.5	912 92.1	226 10.4	80 6.7	146 14.7
1955 No. Rate	5,969 273.9	3,517 295.9	2,452 247.6	401 18.4	144 12.1	257 26.0	2,447 112.3	1,169 98.3	1,278 129.0	2,317 106.3	1,402 118.0	915 92.4	211 9.7	66 5.6	145 14.6
1956 No. Rate	6,104 289.3	3,703 322.8	2,401 249.4	410 19.4	161 14.0	249 25.9	2,550 120.9	1,229 107.1	1,321 137.2	2,432 115.3	1,439 125.4	993 103.2	199 9.4	80 7.0	119 12.4
1957 No. Rate	6,367 301.8	3,706 323.0	2,661 276.4	362 17.1	143 12.5	219 22.7	2,800 132.7	1,420 123.8	1,380 132.2	2,496 118.3	1,482 129.2	1,014 105.3	161 7.6	61 5.3	100 10.4

* Rate per 100,000 population.

1,434

TABLE 16
BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
STATE OF MISSISSIPPI, 1957

67


Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non- white	Number			Total	White	Non- white
				Total	White	Non- white			
Live births	61291	27320	33971	60712	26804	33908	28.8	23.4	35.2
Fetal deaths	2486	994	1492	2454	965	1489	40.4	36.0	43.9
Deaths from all causes	20233	10173	10060	20476	10339	10137	9.7	9.0	10.5
Deaths under 1 year	2236	677	1559	2257	694	1563	37.2	25.9	46.1
Deaths under 1 month	1408	528	880	1425	540	885	23.5	20.1	26.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	149	52	97	161	61	100	7.6	5.3	10.4
Tbc., respiratory (001-008)	138	49	89	150	58	92	7.1	5.0	9.5
Syphilis (020-029)	45	10	35	50	15	35	2.4	1.3	3.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	38	5	33	39	5	34	1.8	0.4	3.5
Scarlet fever & streptococcal sore throat (050, 051)	4	3	1	3	2	1	0.1	0.2	0.1
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	14	0	14	14	0	14	0.7	-	1.4
Meningococcal infections (057)	7	3	4	10	6	4	0.5	0.5	0.4
Acute poliomyelitis (080)	2	0	2	3	1	2	0.1	0.1	0.2
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	5	4	1	5	4	1	0.2	0.3	0.1
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	2408	1423	985	2496	1482	1014	118.3	129.2	105.3
Diabetes (260)	254	136	118	260	140	120	12.3	12.2	12.5
Pellagra (281)	3	1	2	3	1	2	0.1	0.1	0.2
Intracranial lesions (330-334)	2774	1404	1370	2800	1420	1380	132.7	123.8	143.4
Heart diseases (410-443)	6356	3716	2640	6367	3706	2661	301.8	323.0	276.4
Arteriosclerosis (450)	231	138	93	234	139	95	11.1	12.1	9.9
Influenza (480-483)	173	54	119	173	53	120	8.2	4.6	12.5
#Pneumonia (490-493)	611	241	370	621	248	373	29.4	21.6	38.7
#Diarrhea & enteritis (571)	170	34	136	162	30	132	7.7	2.6	13.7
Acute nephritis (590, 591)	56	21	35	58	24	34	2.7	2.1	3.5
Chronic nephritis, etc. (592-594)	303	116	187	304	119	185	14.4	10.4	19.2
Maternal deaths (640-689)	77	15	62	75	14	61	1.2	0.5	1.8
Cong. malformations (750-759)	232	143	89	259	168	91	12.3	14.6	9.4
Birth inj., atelect. (760-762)	368	176	192	365	173	192	6.0	6.4	5.7
#Other dis. early inf. (763-776)	772	237	535	777	241	536	12.8	9.0	15.8
Violent & acc. deaths (800-999)	1642	884	758	1660	898	762	78.7	78.3	79.2
Acc. deaths, total (800-962)	1291	694	597	1302	707	595	61.7	61.6	61.8
Motor vehicle acc. (810-835)	567	355	212	551	346	205	26.1	30.2	21.3
Ill-defined (780-795)	1772	373	1399	1763	366	1397	83.6	31.9	145.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

**PUBLIC HEALTH STATISTICS
STATE OF
MISSISSIPPI
1957**



**Part II
VITAL STATISTICS
COUNTY SUMMARIES**

Mississippi State Board of Health

Jackson, Mississippi

A. L. Gray, M. D., Executive Officer

ANNUAL BULLETIN

of

VITAL STATISTICS

1957

VOLUME 2

COUNTY TABLES

MISSISSIPPI STATE BOARD OF HEALTH

VOLUME 2

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 1 MONTH, AND DEATHS FROM CERTAIN SELECTED CAUSES, ON BASIS OF PLACE OF OCCURRENCE AND PLACE OF RESIDENCE, BY RACE, WITH RATES, FOR THE STATE OF MISSISSIPPI AND FOR EACH COUNTY.

COUNTY	PAGE
Adams	1
Alcorn	2
Amite	3
Attala	4
Benton	5
Bolivar	6
Calhoun	7
Carroll	8
Chickasaw	9
Choctaw	10
Claiborne	11
Clarke	12
Clay	13
Coahoma	14
Copiah	15
Covington	16
DeSoto	17
Forrest	18
Franklin	19
George	20
Greene	21
Grenada	22
Hancock	23
Harrison	24
Hinds	25
Holmes	26
Humphreys	27
Issaquena	28
Itawamba	29
Jackson	30
Jasper	31
Jefferson	32
Jeff Davis	33
Jones	34
Kemper	35
Lafayette	36
Lamar	37
Lauderdale	38
Lawrence	39
Leake	40
Lee	41
Leflore	42

COUNTY	PAGE
Lincoln	43
Lowndes	44
Madison	45
Marion	46
Marshall	47
Monroe	48
Montgomery	49
Neshoba	50
Newton	51
Noxubee	52
Oktibbeha	53
Panola	54
Pearl River	55
Perry	56
Pike	57
Pontotoc	58
Prentiss	59
Quitman	60
Rankin	61
Scott	62
Sharkey	63
Simpson	64
Smith	65
Stone	66
Sunflower	67
Tallahatchie	68
Tate	69
Tippah	70
Tishomingo	71
Tunica	72
Union	73
Walthall	74
Warren	75
Washington	76
Wayne	77
Webster	78
Wilkinson	79
Winston	80
Yalobusha	81
Yazoo	82
STATE	83

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ADAMS COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,766	727	1,039	1,246	544	702	34.3	28.2	41.3
Fetal deaths	51	8	43	33	5	28	26.5	9.2	39.9
Deaths from all causes	453	160	293	415	154	261	11.4	8.0	15.3
Deaths under 1 year	54	17	37	44	15	29	35.3	27.6	41.3
Deaths under 1 month	34	13	21	28	13	15	22.5	23.9	21.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	3	0	3	8.3	-	17.6
Tbc., respiratory (001-008)	0	0	0	2	0	2	5.5	-	11.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	2.8	-	5.9
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	46	13	33	45	15	30	124.0	77.7	176.5
Diabetes (260)	14	4	10	11	3	8	30.3	15.5	47.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	62	21	41	56	19	37	154.3	98.4	217.6
Heart diseases (410-443)	175	60	115	161	55	106	443.5	285.0	623.5
Arteriosclerosis (450)	4	1	3	1	1	0	2.8	5.2	-
Influenza (480-483)	1	0	1	1	0	1	2.8	-	5.9
#Pneumonia (490-493)	10	4	6	11	5	6	30.3	25.9	35.3
#Diarrhea & enteritis (571)	3	1	2	3	1	2	8.3	5.2	11.8
Acute nephritis (590, 591)	1	0	1	1	0	1	2.8	-	5.9
Chronic nephritis, etc. (592-594)	5	2	3	4	2	2	11.0	10.4	11.8
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	10	7	3	8	6	2	22.0	31.1	11.8
Birth inj., atelect. (760-762)	5	3	2	3	3	0	2.4	5.5	-
#Other dis. early inf. (763-776)	25	4	21	20	5	15	16.0	9.2	21.4
Violent & acc. deaths (800-999)	33	21	12	34	22	12	93.7	114.0	70.6
Acc. deaths, total (800-962)	21	12	9	23	14	9	63.4	72.5	52.9
Motor vehicle acc. (810-835)	9	6	3	14	10	4	38.6	51.8	23.5
Ill-defined (780-795)	16	0	16	15	1	14	41.3	5.2	82.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ALCORN COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	607	485	122	545	434	111	21.7	20.2	30.8
Fetal deaths	12	8	4	12	9	3	22.0	20.7	27.0
Deaths from all causes	242	194	48	248	199	49	9.9	9.2	13.6
Deaths under 1 year	18	14	4	18	13	5	33.0	30.0	45.0
Deaths under 1 month	14	11	3	13	10	3	23.8	23.0	27.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	1	1	0	1	1	0	4.0	4.6	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	1	0	1	1	0	4.0	4.6	-
Acute poliomyelitis (080)	1	0	1	1	0	1	4.0	-	27.8
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	28	23	5	26	21	5	103.6	97.7	138.9
Diabetes (260)	5	5	0	4	4	0	15.9	18.6	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	39	25	14	36	22	14	143.4	102.3	38.9
Heart diseases (410-443)	88	73	15	90	76	14	358.6	353.5	38.9
Arteriosclerosis (450)	9	8	1	10	9	1	39.8	41.9	27.8
Influenza (480-483)	1	1	0	1	1	0	4.0	4.6	-
#Pneumonia (490-493)	7	6	1	9	8	1	35.8	37.2	27.8
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	1	1	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	5	3	2	4	2	2	7.3	4.6	18.0
#Other dis. early inf. (763-776)	7	6	1	7	6	1	12.8	13.8	9.0
Violent & acc. deaths (800-999)	23	22	1	26	25	1	103.6	116.3	27.8
Acc. deaths, total (800-962)	16	15	1	19	18	1	75.7	83.7	27.8
Motor vehicle acc. (810-835)	7	7	0	8	8	0	31.9	37.2	-
Ill-defined (780-795)	3	1	2	4	2	2	15.9	9.3	55.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
AMITE COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non white	Number			Total	White	Non white
				Total	White	Non white			
Live births	274	30	244	418	125	293	24.7	17.4	30.2
Fetal deaths	5	0	5	7	0	7	16.7	-	23.9
Deaths from all causes	115	45	70	179	81	98	10.6	11.3	10.1
Deaths under 1 year	14	1	13	27	6	21	64.6	48.0	71.7
Deaths under 1 month	8	1	7	13	4	9	31.1	32.0	30.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	2	0	2	11.8	-	20.6
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	4	5	18	10	8	106.5	138.9	82.5
Diabetes (260)	3	2	1	3	2	1	17.8	27.8	10.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	22	10	12	32	18	14	189.3	250.0	144.3
Heart diseases (410-443)	30	15	15	44	23	21	260.4	319.4	216.5
Arteriosclerosis (450)	1	0	1	1	0	1	5.9	-	10.3
Influenza (480-483)	2	0	2	2	0	2	11.8	-	20.6
#Pneumonia (490-493)	5	1	4	5	1	4	29.6	13.9	41.2
#Diarrhea & enteritis (571)	1	0	1	4	0	4	23.7	-	41.2
Acute nephritis (590, 591)	0	0	0	1	0	1	5.9	-	10.3
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	1	0	1	1	0	1	2.4	-	3.4
Cong. malformations (750-759)	1	0	1	3	2	1	17.8	27.8	10.3
Birth inj., atelect. (760-762)	2	1	1	3	2	1	7.2	16.0	3.4
#Other dis. early inf. (763-776)	1	0	1	4	1	3	9.6	8.0	10.2
Violent & acc. deaths (800-999)	1	0	1	12	7	5	71.0	97.2	51.5
Acc. deaths, total (800-962)	9	7	2	8	4	4	47.3	55.6	41.2
Motor vehicle acc. (810-835)	6	4	2	3	1	1	23.7	41.7	10.3
Ill-defined (780-795)	3	3	0	4	3	10	94.7	83.3	103.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ATTALA COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	570	228	342	531	200	331	22.9	16.3	30.4
Fetal deaths	14	1	13	13	2	11	24.5	10.0	33.2
Deaths from all causes	215	130	85	231	148	83	10.0	12.0	7.6
Deaths under 1 year	17	6	11	19	8	11	35.8	40.0	33.2
Deaths under 1 month	12	5	7	13	6	7	24.5	30.0	21.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	1	0	1	4.3	-	9.2
Tbc., respiratory (001-008)	1	0	1	1	0	1	4.3	-	9.2
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	33	18	15	35	21	14	150.9	170.7	128.4
Diabetes (260)	3	3	0	3	3	0	12.9	24.4	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	35	25	10	35	25	10	150.9	203.2	91.7
Heart diseases (410-443)	76	48	28	82	52	30	353.4	422.8	275.2
Arteriosclerosis (450)	2	1	1	2	1	1	8.6	8.1	9.2
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	8	3	5	9	4	5	38.8	32.5	45.9
#Diarrhea & enteritis (571)	3	1	2	3	1	2	12.9	8.1	18.3
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	0	2	4	2	2	17.2	16.3	18.3
Birth inj., atelect. (760-762)	5	4	1	5	4	1	9.4	20.0	3.0
#Other dis. early inf. (763-776)	6	1	5	6	1	5	11.3	5.0	15.1
Violent & acc. deaths (800-999)	15	10	5	21	16	5	90.5	130.1	45.9
Acc. deaths, total (800-962)	9	7	2	15	13	2	64.6	105.7	18.3
Motor vehicle acc. (810-835)	3	3	0	5	4	1	21.6	32.5	9.2
Ill-defined (780-795)	14	6	8	12	6	6	51.7	48.8	55.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
BENTON COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	119	9	110	216	78	138	25.4	18.1	32.8
Fetal deaths	3	0	3	15	7	8	69.4	89.7	58.0
Deaths from all causes	48	25	23	80	46	34	9.4	10.7	8.1
Deaths under 1 year	4	0	4	11	3	8	50.9	38.5	58.0
Deaths under 1 month	1	0	1	8	3	5	37.0	38.5	36.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	11.8	-	23.8
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	4	2	2	9	7	2	105.9	162.8	47.6
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	8	5	3	10	7	3	117.6	162.8	71.4
Heart diseases (410-443)	20	10	10	28	17	11	329.4	395.3	261.9
Arteriosclerosis (450)	0	0	0	1	0	1	11.8	-	23.8
Influenza (480-483)	0	0	0	1	1	0	11.8	23.2	-
#Pneumonia (490-493)	0	0	0	0	0	0	-	-	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	2	1	1	23.5	23.2	23.8
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	2	2	0	23.5	46.5	-
Birth inj., atelect. (760-762)	0	0	0	3	3	0	13.9	38.5	-
#Other dis. early inf. (763-776)	1	0	1	5	0	5	23.1	-	36.2
Violent & acc. deaths (800-999)	5	3	2	4	2	2	47.0	46.5	47.6
Acc. deaths, total (800-962)	4	3	1	2	1	1	23.5	23.2	23.8
Motor vehicle acc. (810-835)	2	2	0	0	0	0	-	-	-
Ill-defined (780-795)	4	2	2	4	1	3	47.0	23.2	71.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
BOLIVAR COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,953	410	1,543	1,811	414	1,397	30.6	23.5	33.7
Fetal deaths	68	10	58	68	14	54	37.5	33.8	38.6
Deaths from all causes	548	102	446	574	145	429	9.7	8.2	10.3
Deaths under 1 year	68	8	60	68	10	58	37.5	24.2	41.5
Deaths under 1 month	35	7	28	36	9	27	19.9	21.7	19.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	0	3	2	0	2	3.4	-	4.8
Tbc., respiratory (001-008)	2	0	2	2	0	2	3.4	-	4.8
Syphilis (020-029)	3	0	3	3	0	3	5.1	-	7.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	1.7	-	2.4
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	1	0	1.7	5.7	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	70	17	53	77	23	54	130.3	130.7	130.1
Diabetes (260)	6	1	5	6	2	4	10.2	11.4	9.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	69	10	59	70	14	56	118.4	79.5	134.9
Heart diseases (410-443)	176	48	128	178	61	117	301.2	346.6	281.9
Arteriosclerosis (450)	4	2	2	6	4	2	10.2	22.7	4.8
Influenza (480-483)	5	0	5	4	0	4	6.8	-	9.6
#Pneumonia (490-493)	14	1	13	14	3	11	23.7	17.0	26.5
#Diarrhea & enteritis (571)	15	1	14	14	1	13	23.7	5.7	31.3
Acute nephritis (590, 591)	2	0	2	2	0	2	3.4	-	4.8
Chronic nephritis, etc. (592-594)	11	1	10	10	2	8	16.9	11.4	19.3
Maternal deaths (640-689)	2	0	2	4	0	4	2.2	-	2.9
Cong. malformations (750-759)	3	1	2	1	0	1	1.7	-	2.4
Birth inj., atelect. (760-762)	8	0	8	9	0	9	5.0	-	6.4
#Other dis. early inf. (763-776)	26	4	22	27	7	20	14.9	16.9	14.3
Violent & acc. deaths (800-999)	28	8	20	35	12	23	59.2	68.2	55.4
Acc. deaths, total (800-962)	22	6	16	29	10	19	49.1	56.8	45.8
Motor vehicle acc. (810-835)	5	2	3	6	4	2	10.2	22.7	4.8
Ill-defined (780-795)	70	1	69	71	2	69	120.1	11.4	166.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CALHOUN COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	436	182	254	405	186	219	26.8	17.7	47.6
Fetal deaths	21	7	14	16	5	11	39.5	26.9	50.2
Deaths from all causes	123	90	33	141	104	37	9.3	9.9	8.0
Deaths under 1 year	9	1	8	8	1	7	19.8	5.4	32.0
Deaths under 1 month	7	1	6	7	1	6	17.3	5.4	27.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	6.6	9.5	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	6.6	9.5	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	15	14	1	15	14	1	99.3	133.3	21.7
Diabetes (260)	3	1	2	4	2	2	26.5	19.0	43.5
Fellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	14	11	3	18	14	4	119.2	133.3	87.0
Heart diseases (410-443)	43	35	8	44	34	10	291.4	323.8	217.4
Arteriosclerosis (450)	2	2	0	2	2	0	13.2	19.0	-
Influenza (480-483)	1	1	0	1	1	0	6.6	9.5	-
#Pneumonia (490-493)	6	4	2	7	5	2	46.4	47.6	43.5
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	6.6	-	21.7
Maternal deaths (640-689)	1	0	1	1	0	1	2.5	-	4.6
Cong. malformations (750-759)	0	0	0	1	1	0	6.6	9.5	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	5	1	4	5	1	4	12.3	5.4	18.3
Violent & acc. deaths (800-999)	7	4	3	13	9	4	86.1	85.7	87.0
Acc. deaths, total (800-962)	6	3	3	12	8	4	79.5	76.2	87.0
Motor vehicle acc. (810-835)	3	2	1	6	4	2	39.7	38.1	43.5
Ill-defined (780-795)	15	10	5	16	11	5	106.0	104.8	108.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CARROLL COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	231	23	208	268	48	220	20.8	9.8	27.5
Fetal deaths	11	1	10	21	5	16	78.4	104.2	72.7
Deaths from all causes	69	29	40	113	56	57	8.8	11.4	7.1
Deaths under 1 year	7	0	7	9	0	9	33.6	-	40.9
Deaths under 1 month	4	0	4	5	0	5	18.6	-	22.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	6	5	1	12	9	3	93.0	183.7	37.5
Diabetes (260)	0	0	0	1	0	1	7.8	-	12.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	7	3	4	15	9	6	116.3	183.7	75.0
Heart diseases (410-443)	17	9	8	36	22	14	279.1	449.0	175.0
Arteriosclerosis (450)	0	0	0	1	1	0	7.8	20.4	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	0	0	0	2	0	2	15.5	-	25.0
#Diarrhea & enteritis (571)	1	0	1	1	0	1	7.8	-	12.5
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	2	1	1	15.5	20.4	12.5
Maternal deaths (640-689)	0	0	0	2	1	1	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	1	0	1	7.8	-	12.5
#Other dis. early inf. (763-776)	1	0	1	2	0	2	7.5	-	9.1
Violent & acc. deaths (800-999)	5	2	3	4	1	3	31.0	20.4	37.5
Acc. deaths, total (800-962)	5	2	3	4	1	3	31.0	20.4	37.5
Motor vehicle acc. (810-835)	4	2	2	3	1	2	23.2	20.4	25.0
Ill-defined (780-795)	24	4	20	25	5	20	193.8	102.0	250.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CHICKASAW COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	439	207	232	458	207	251	26.6	21.1	33.9
Fetal deaths	15	3	12	18	4	14	39.3	19.3	55.8
Deaths from all causes	178	93	85	179	96	83	10.4	9.8	11.2
Deaths under 1 year	22	9	13	24	9	15	52.4	43.5	59.8
Deaths under 1 month	16	5	11	17	5	12	37.1	24.2	47.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	1	0	1	1	0	1	5.8	-	13.5
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	14	11	3	16	12	4	93.0	122.4	54.0
Diabetes (260)	1	0	1	2	1	1	11.6	10.2	13.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	27	10	17	24	9	15	139.5	91.8	202.7
Heart diseases (410-443)	71	44	27	66	43	23	383.7	438.8	310.8
Arteriosclerosis (450)	1	1	0	3	1	2	17.4	10.2	27.0
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	6	3	3	7	4	3	40.7	40.8	40.5
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	5.8	-	13.5
Maternal deaths (640-689)	1	0	1	2	1	1	4.4	4.8	4.0
Cong. malformations (750-759)	2	2	0	1	1	0	5.8	10.2	-
Birth inj., atelect. (760-762)	3	2	1	2	1	1	4.4	4.8	4.0
#Other dis. early inf. (763-776)	11	2	9	13	3	10	28.4	14.5	39.8
Violent & acc. deaths (800-999)	16	5	11	17	6	11	98.8	61.2	148.6
Acc. deaths, total (800-962)	15	4	11	17	6	11	98.8	61.2	148.6
Motor vehicle acc. (810-835)	2	1	1	3	2	1	17.4	10.2	27.0
Ill-defined (780-795)	8	3	5	9	4	5	52.3	40.8	67.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CHOCTAW COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Number			Residence		
	Total	White	Non-white	Total	White	Non-white	Rate*		
							Total	White	
Live births	207	103	104	214	112	102	22.8	18.1	31.9
Fetal deaths	2	0	2	3	1	2	14.0	8.9	19.6
Deaths from all causes	92	57	35	110	74	36	11.7	11.9	11.2
Deaths under 1 year	7	3	4	6	2	4	28.0	17.8	39.2
Deaths under 1 month	6	3	3	5	2	3	23.4	17.8	29.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	1	0	1	1	0	10.6	16.1	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	14	7	7	16	8	8	170.2	129.0	250.0
Diabetes (260)	1	1	0	1	1	0	10.6	16.1	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	18	13	5	21	16	5	223.4	258.1	156.2
Heart diseases (410-443)	28	18	10	33	25	8	351.1	403.2	250.0
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	1	0	1	10.6	-	31.2
#Pneumonia (490-493)	2	0	2	1	0	1	10.6	-	31.2
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	10.6	16.1	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	1	0	1	10.6	-	31.2
Birth inj., atelect. (760-762)	2	1	1	2	1	1	9.3	8.9	9.8
#Other dis. early inf. (763-776)	0	0	0	0	0	0	-	-	-
Violent & acc. deaths (800-999)	9	8	1	13	11	2	138.3	177.4	62.5
Acc. deaths, total (800-962)	6	5	1	10	8	2	106.4	129.0	62.5
Motor vehicle acc. (810-835)	1	1	0	6	6	0	63.8	96.8	-
Ill-defined (780-795)	5	2	3	5	2	3	53.2	32.2	93.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CLAIBORNE COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	256	26	230	305	42	263	27.2	13.5	32.5
Fetal deaths	9	0	9	12	3	9	39.3	71.4	34.2
Deaths from all causes	98	24	74	124	27	97	11.1	8.7	12.0
Deaths under 1 year	9	1	8	15	1	14	49.2	23.8	53.2
Deaths under 1 month	5	1	4	10	1	9	32.8	23.8	34.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	2	0	2	17.8	-	24.7
Tbc., respiratory (001-008)	2	0	2	2	0	2	17.8	-	24.7
Syphilis (020-029)	0	0	0	1	0	1	8.9	-	12.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	2	0	2	17.8	-	24.7
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	8	0	8	13	0	13	116.1	-	160.5
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	16	2	14	17	2	15	151.8	64.5	185.2
Heart diseases (410-443)	33	14	19	40	16	24	357.1	516.1	296.3
Arteriosclerosis (450)	2	1	1	2	1	1	17.8	32.2	12.3
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	4	1	3	5	1	4	44.6	32.2	49.4
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	0	2	2	0	2	17.8	-	24.7
Maternal deaths (640-689)	1	1	0	1	1	0	3.3	23.8	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	1	0	1	3.3	-	3.8
#Other dis. early inf. (763-776)	6	1	5	10	1	9	32.8	23.8	34.2
Violent & acc. deaths (800-999)	5	1	4	6	1	5	53.6	32.2	61.7
Acc. deaths, total (800-962)	4	1	3	5	1	4	44.6	32.2	49.4
Motor vehicle acc. (810-835)	2	0	2	2	0	2	17.8	-	24.7
<u>Ill-defined (780-795)</u>	<u>17</u>	<u>3</u>	<u>14</u>	<u>17</u>	<u>3</u>	<u>14</u>	<u>151.8</u>	<u>96.8</u>	<u>172.8</u>

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CLARKE COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	264	137	127	437	201	236	25.3	19.9	32.8
Fetal deaths	8	3	5	6	2	4	13.7	10.0	16.9
Deaths from all causes	155	83	72	195	113	82	11.3	11.2	11.4
Deaths under 1 year	12	9	3	16	9	7	36.6	44.8	29.7
Deaths under 1 month	6	5	1	7	5	2	16.0	24.9	8.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	3	1	2	17.3	9.9	27.8
Tbc., respiratory (001-008)	2	0	2	3	1	2	17.3	9.9	27.8
Syphilis (020-029)	0	0	0	1	1	0	5.8	9.9	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	5.8	-	13.9
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	24	18	6	34	26	8	196.5	257.4	111.1
Diabetes (260)	3	3	0	3	3	0	17.3	29.7	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	16	3	13	25	11	14	144.5	108.9	194.4
Heart diseases (410-443)	61	35	26	64	39	25	369.9	386.1	347.2
Arteriosclerosis (450)	4	0	4	4	0	4	23.1	-	55.6
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	1	0	2	1	1	11.6	9.9	13.9
#Diarrhea & enteritis (571)	1	1	0	3	2	1	17.3	19.8	13.9
Acute nephritis (590, 591)	0	0	0	1	0	1	5.8	-	13.9
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	11.6	9.9	13.9
Maternal deaths (640-689)	1	1	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	2	1	1	11.6	9.9	13.9
Birth inj., atelect. (760-762)	3	3	0	4	3	1	9.2	14.9	4.2
#Other dis. early inf. (763-776)	2	1	1	2	1	1	4.6	5.0	4.2
Violent & acc. deaths (800-999)	16	9	7	22	13	9	127.2	128.7	125.0
Acc. deaths, total (800-962)	16	9	7	22	13	9	127.2	128.7	125.0
Motor vehicle acc. (810-835)	6	2	4	8	4	4	46.2	39.6	55.6
Ill-defined (780-795)	7	2	5	7	3	4	40.5	29.7	55.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CLAY COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	555	209	346	579	223	356	38.3	29.3	47.5
Fetal deaths	12	3	9	12	2	10	20.7	9.0	28.1
Deaths from all causes	179	84	95	199	87	112	13.2	11.4	14.9
Deaths under 1 year	24	8	16	28	8	20	48.4	35.9	56.2
Deaths under 1 month	13	5	8	14	5	9	24.2	22.4	25.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	1	1	0	6.6	13.2	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	10	7	20	13	7	132.4	171.0	93.3
Diabetes (260)	2	1	1	2	1	1	13.2	13.2	13.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	21	10	11	22	10	12	145.7	131.6	160.0
Heart diseases (410-443)	76	37	39	82	38	44	543.0	500.0	586.7
Arteriosclerosis (450)	1	1	0	1	1	0	6.6	13.2	-
Influenza (480-483)	2	1	1	2	1	1	13.2	13.2	13.3
#Pneumonia (490-493)	8	5	3	7	3	4	46.4	39.5	53.3
#Diarrhea & enteritis (571)	3	0	3	6	0	6	39.7	-	80.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	6.6	-	13.3
Maternal deaths (640-689)	3	1	2	2	1	1	3.4	4.5	2.8
Cong. malformations (750-759)	2	2	0	2	2	0	13.2	26.3	-
Birth inj., atelect. (760-762)	3	2	1	4	2	2	6.9	9.0	5.6
#Other dis. early inf. (763-776)	7	1	6	7	1	6	12.1	4.5	16.8
Violent & acc. deaths (800-999)	15	10	5	17	9	8	112.6	118.4	106.7
Acc. deaths, total (800-962)	13	9	4	14	8	6	92.7	105.3	80.0
Motor vehicle acc. (810-835)	3	3	0	3	1	2	19.9	13.2	26.7
Ill-defined (780-795)	13	2	11	14	1	13	92.7	13.2	173.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
COAHOMA COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,597	390	1,207	1,616	353	1,263	30.7	24.5	33.1
Fetal deaths	69	22	47	67	20	47	41.5	56.6	37.2
Deaths from all causes	469	130	339	488	113	375	9.3	7.8	9.8
Deaths under 1 year	67	5	62	67	4	63	41.5	11.3	49.9
Deaths under 1 month	41	4	37	42	4	38	26.0	11.3	30.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	6	0	6	8	0	8	15.2	-	20.9
Tbc., respiratory (001-008)	6	0	6	7	0	7	13.3	-	18.3
Syphilis (020-029)	1	1	0	1	1	0	1.9	6.9	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	61	21	40	67	20	47	127.4	138.9	123.0
Diabetes (260)	3	1	2	3	1	2	5.7	6.9	5.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	48	13	35	53	11	42	100.8	76.4	109.9
Heart diseases (410-443)	135	45	90	140	40	100	266.2	277.8	261.8
Arteriosclerosis (450)	5	2	3	5	2	3	9.5	13.9	7.8
Influenza (480-483)	4	0	4	5	0	5	9.5	-	13.1
#Pneumonia (490-493)	13	2	11	13	2	11	24.7	13.9	28.8
#Diarrhea & enteritis (571)	10	1	9	9	0	9	17.1	-	23.6
Acute nephritis (590, 591)	1	1	0	1	0	1	1.9	-	2.6
Chronic nephritis, etc. (592-594)	9	2	7	12	2	10	22.8	13.9	26.2
Maternal deaths (640-689)	3	0	3	3	0	3	1.8	-	2.4
Cong. malformations (750-759)	2	0	2	5	2	3	9.5	13.9	7.8
Birth inj., atelect. (760-762)	7	1	6	7	1	6	4.3	2.8	4.8
#Other dis. early inf. (763-776)	26	3	23	26	2	24	16.1	5.7	19.0
Violent & acc. deaths (800-999)	48	20	28	38	10	28	72.2	69.4	73.3
Acc. deaths, total (800-962)	41	19	22	31	9	22	58.9	62.5	57.6
Motor vehicle acc. (810-835)	14	11	3	8	5	3	15.2	34.7	7.8
Un-defined (780-795)	50	4	46	48	3	45	91.2	20.8	117.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
COPIAH COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	910	275	635	756	238	513	27.6	19.2	34.5
Fetal deaths	23	3	20	22	3	19	29.1	12.6	36.7
Deaths from all causes	258	108	150	302	137	165	11.0	11.0	11.0
Deaths under 1 year	32	4	28	31	4	27	41.0	16.8	52.1
Deaths under 1 month	18	4	14	16	3	13	21.2	12.6	25.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	2	1	1	7.3	8.1	6.7
Tbc., respiratory (001-008)	1	0	1	2	1	1	7.3	8.1	6.7
Syphilis (020-029)	3	1	2	4	2	2	14.6	16.1	13.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	25	17	8	31	19	12	113.1	153.2	80.0
Diabetes (260)	2	2	0	2	2	0	7.3	16.1	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	29	8	21	34	11	23	124.1	88.7	153.3
Heart diseases (410-443)	100	51	49	103	53	50	375.9	427.4	333.3
Arteriosclerosis (450)	3	2	1	3	2	1	10.9	16.1	6.7
Influenza (480-483)	3	0	3	3	0	3	10.9	-	20.0
#Pneumonia (490-493)	11	2	9	16	6	10	58.4	48.4	66.7
#Diarrhea & enteritis (571)	1	0	1	1	0	1	3.6	-	6.7
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	2	1	5	3	2	18.2	24.2	13.3
Maternal deaths (640-689)	2	0	2	1	0	1	1.3	-	1.9
Cong. malformations (750-759)	2	0	2	5	1	4	18.2	8.1	26.7
Birth inj., atelect. (760-762)	4	0	4	3	0	3	4.0	-	5.8
#Other dis. early inf. (763-776)	8	3	5	7	2	5	9.2	8.4	9.6
Violent & acc. deaths (800-999)	23	10	13	32	15	17	116.8	121.0	113.3
Acc. deaths, total (800-962)	18	7	11	26	12	14	94.9	96.8	93.3
Motor vehicle acc. (810-835)	6	5	1	10	8	2	36.5	64.5	13.3
Ill-defined (780-795)	15	1	14	15	1	14	54.7	8.1	93.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
COVINGTON COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	193	130	63	347	189	158	22.7	19.9	27.2
Fetal deaths	3	0	3	8	4	4	23.0	21.2	25.3
Deaths from all causes	116	71	45	152	96	56	9.9	10.1	9.6
Deaths under 1 year	7	3	4	14	8	6	40.3	42.3	38.0
Deaths under 1 month	5	3	2	12	7	5	34.6	37.0	31.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	1	0	6.5	10.5	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	0	1	2	1	1	13.1	10.5	17.2
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	15	10	5	16	11	5	104.6	115.8	86.2
Diabetes (260)	1	1	0	2	2	0	13.1	21.0	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	23	11	12	33	18	15	215.7	189.5	258.6
Heart diseases (410-443)	46	32	14	53	36	17	346.4	378.9	293.1
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	4	1	3	7	3	4	45.8	31.6	69.0
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	3	1	2	2	0	2	13.1	-	34.5
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	6.5	-	17.2
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	6.5	10.5	-
Birth inj., atelect. (760-762)	2	2	0	5	4	1	14.4	21.2	6.3
#Other dis. early inf. (763-776)	1	1	0	2	2	0	5.8	10.6	-
Violent & acc. deaths (800-999)	6	4	2	9	6	3	58.8	63.2	51.7
Acc. deaths, total (800-962)	4	3	1	7	5	2	45.8	52.6	34.5
Motor vehicle acc. (810-835)	3	2	1	5	4	1	32.7	42.1	17.2
<u>Ill-defined (780-795)</u>	8	4	4	10	4	6	65.4	42.1	103.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
DESOTO COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	692	98	594	750	178	572	31.1	21.2	36.4
Fetal deaths	21	1	20	24	1	23	32.0	5.6	40.2
Deaths from all causes	179	53	126	231	87	144	9.6	10.4	9.2
Deaths under 1 year	18	1	17	29	8	21	38.7	44.9	36.7
Deaths under 1 month	7	1	6	17	7	10	22.7	39.3	17.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	2	0	2	8.3	-	12.7
Tbc., respiratory (001-008)	2	0	2	2	0	2	8.3	-	12.7
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	16	4	12	24	9	15	99.6	107.1	95.5
Diabetes (260)	1	0	1	2	0	2	8.3	-	12.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	22	8	14	30	12	18	124.5	142.8	114.6
Heart diseases (410-443)	42	17	25	52	25	27	215.8	297.6	172.0
Arteriosclerosis (450)	4	2	2	4	2	2	16.6	23.8	12.7
Influenza (480-483)	2	1	1	2	1	1	8.3	11.9	6.4
#Pneumonia (490-493)	11	3	8	13	4	9	53.9	47.6	57.3
#Diarrhea & enteritis (571)	1	0	1	1	0	1	4.1	-	6.4
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	8.3	11.9	6.4
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	4	3	1	16.6	35.7	6.4
Birth inj., atelect. (760-762)	1	0	1	2	1	1	2.7	5.6	1.7
#Other dis. early inf. (763-776)	0	0	0	5	3	2	6.7	16.8	3.5
Violent & acc. deaths (800-999)	20	8	12	19	7	12	78.8	83.3	76.4
Acc. deaths, total (800-962)	18	7	11	17	6	11	70.5	71.4	70.1
Motor vehicle acc. (810-835)	8	5	3	6	4	2	24.9	47.6	12.7
Ill-defined (780-795)	41	4	37	44	5	39	182.6	59.5	248.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
FORREST COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,538	1,141	397	1,292	839	453	29.0	26.0	37.1
Fetal deaths	54	34	20	49	23	26	37.9	27.4	57.4
Deaths from all causes	457	321	136	448	287	161	10.1	8.9	13.2
Deaths under 1 year	33	17	16	37	16	21	28.6	19.1	46.4
Deaths under 1 month	24	14	10	27	13	14	20.9	15.5	30.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	7	4	3	15.7	12.4	24.6
Tbc., respiratory (001-008)	1	0	1	6	3	3	13.5	9.3	24.6
Syphilis (020-029)	0	0	0	1	1	0	2.2	3.1	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	53	33	20	55	32	23	123.6	99.1	188.5
Diabetes (260)	7	6	1	9	6	3	20.2	18.6	24.6
Pellagra (281)	1	0	1	1	0	1	2.2	-	8.2
Intracranial lesions (330-334)	63	44	19	63	40	23	141.6	123.8	188.5
Heart diseases (410-443)	172	135	37	156	113	43	350.6	349.8	352.5
Arteriosclerosis (450)	7	6	1	6	6	0	13.5	18.6	-
Influenza (480-483)	10	4	6	10	4	6	22.5	12.4	49.2
#Pneumonia (490-493)	10	9	1	8	6	2	18.0	18.6	16.4
#Diarrhea & enteritis (571)	1	1	0	1	1	0	2.2	3.1	-
Acute nephritis (590, 591)	2	2	0	2	2	0	4.5	6.2	-
Chronic nephritis, etc. (592-594)	6	4	2	6	3	3	13.5	9.3	24.6
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	3	3	0	2	2	0	4.5	6.2	-
Birth inj., atelect. (760-762)	8	6	2	11	7	4	8.5	8.3	8.8
#Other dis. early inf. (763-776)	14	6	8	15	5	10	11.6	6.0	22.1
Violent & acc. deaths (800-999)	36	20	16	33	19	14	74.2	58.8	114.8
Acc. deaths, total (800-962)	29	15	14	27	15	12	60.7	46.4	98.4
Motor vehicle acc. (810-835)	11	6	5	8	5	3	18.0	15.5	24.6
Ill-defined (780-795)	18	6	12	17	6	11	38.2	18.6	90.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
FRANKLIN COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	122	54	68	246	103	143	28.0	19.4	40.8
Fetal deaths	1	0	1	6	0	6	24.4	-	42.0
Deaths from all causes	65	33	32	87	45	42	9.9	8.5	12.0
Deaths under 1 year	6	1	5	10	2	8	40.6	19.4	55.9
Deaths under 1 month	3	1	2	6	2	4	24.4	19.4	28.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	2	2	0	22.7	37.7	-
Tbc., respiratory (001-008)	1	1	0	2	2	0	22.7	37.7	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	11.4	-	28.6
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	11	4	7	15	6	9	170.4	113.2	257.1
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	11	5	6	13	5	8	147.7	94.3	228.6
Heart diseases (410-443)	20	13	7	25	17	8	284.1	320.8	228.6
Arteriosclerosis (450)	2	2	0	6	3	3	68.2	56.6	85.7
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	0	1	1	0	1	11.4	-	28.6
#Diarrhea & enteritis (571)	1	0	1	1	0	1	11.4	-	28.6
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	11.4	18.9	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	1	0	1	11.4	-	28.6
Birth inj., atelect. (760-762)	2	0	2	3	0	3	12.2	-	21.0
#Other dis. early inf. (763-776)	1	1	0	2	1	1	8.1	9.7	7.0
Violent & acc. deaths (800-999)	4	3	1	5	4	1	56.8	75.5	28.6
Acc. deaths, total (800-962)	3	2	1	3	2	1	34.1	37.7	28.6
Motor vehicle acc. (810-835)	1	1	0	0	0	0	-	-	-
Ill-defined (780-795)	2	1	1	2	1	1	22.7	18.9	28.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
GEORGE COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	305	257	48	311	262	49	29.6	28.5	37.7
Fetal deaths	6	5	1	8	7	1	25.7	26.7	20.4
Deaths from all causes	92	78	14	98	84	14	9.3	9.1	10.8
Deaths under 1 year	9	6	3	11	8	3	35.4	30.5	61.2
Deaths under 1 month	4	4	0	5	5	0	16.1	19.1	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	9.5	10.9	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	9.5	10.9	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	1	1	0	9.5	10.9	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	9	0	13	13	0	123.8	141.3	-
Diabetes (260)	1	1	0	1	1	0	9.5	10.9	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	5	3	2	4	2	2	38.1	21.7	153.8
Heart diseases (410-443)	55	48	7	52	45	7	495.2	489.1	538.5
Arteriosclerosis (450)	1	1	0	1	1	0	9.5	10.9	-
Influenza (480-483)	2	1	1	2	1	1	19.0	10.9	76.9
#Pneumonia (490-493)	1	0	1	1	0	1	9.5	-	76.9
#Diarrhea & enteritis (571)	1	0	1	1	0	1	9.5	-	76.9
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	2	0	3	3	0	28.6	32.6	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	2	0	2	2	0	19.0	21.7	-
Birth inj., atelect. (760-762)	1	1	0	1	1	0	3.2	3.8	-
#Other dis. early inf. (763-776)	1	1	0	1	1	0	3.2	3.8	-
Violent & acc. deaths (800-999)	7	5	2	7	6	1	66.7	65.2	76.9
Acc. deaths, total (800-962)	4	2	2	4	3	1	38.1	32.6	76.9
Motor vehicle acc. (810-835)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	1	1	0	1	1	0	9.5	10.9	-

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
GREENE COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	126	81	45	209	137	72	26.1	21.4	45.0
Fetal deaths	3	0	3	5	2	3	23.9	14.6	41.7
Deaths from all causes	62	44	18	77	57	20	9.6	8.9	12.5
Deaths under 1 year	5	3	2	5	3	2	23.9	21.9	27.8
Deaths under 1 month	5	3	2	5	3	2	23.9	21.9	27.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	3	2	1	6	5	1	75.0	78.1	62.5
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	11	7	4	13	9	4	162.5	140.6	250.0
Heart diseases (410-443)	28	21	7	33	25	8	412.5	390.6	500.0
Arteriosclerosis (450)	1	0	1	1	0	1	12.5	-	62.5
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	3	3	0	3	3	0	37.5	46.9	-
#Diarrhea & enteritis (571)	0	0	0	1	1	0	12.5	15.6	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	1	1	0	1	1	0	4.8	7.3	-
#Other dis. early inf. (763-776)	2	1	1	2	1	1	9.6	7.3	13.9
Violent & acc. deaths (800-999)	7	6	1	8	6	2	100.0	93.8	125.0
Acc. deaths, total (800-962)	5	4	1	5	4	1	62.5	62.5	62.5
Motor vehicle acc. (810-835)	3	3	0	2	2	0	25.0	31.2	-
Ill-defined (780-795)	5	2	3	6	3	3	75.0	46.9	187.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
GRENADA COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	578	243	335	545	212	333	29.1	24.4	33.3
Fetal deaths	29	4	25	25	3	22	45.9	14.2	66.1
Deaths from all causes	202	90	112	184	70	114	9.8	8.0	11.4
Deaths under 1 year	28	4	24	27	4	23	49.5	18.9	69.1
Deaths under 1 month	19	4	15	18	3	15	33.0	14.2	45.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	5.3	-	10.0
Tbc., respiratory (001-008)	0	0	0	1	0	1	5.3	-	10.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	5.3	-	10.0
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	27	15	12	25	13	12	133.7	149.4	120.0
Diabetes (260)	1	0	1	1	0	1	5.3	-	10.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	50	22	28	45	16	29	240.6	183.9	290.0
Heart diseases (410-443)	44	27	17	37	22	15	197.9	252.9	150.0
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	4	0	4	5	1	4	26.7	11.5	40.0
#Pneumonia (490-493)	2	0	2	2	0	2	10.7	-	20.0
#Diarrhea & enteritis (571)	5	0	5	4	0	4	21.4	-	40.0
Acute nephritis (590, 591)	2	0	2	2	0	2	10.7	-	20.0
Chronic nephritis, etc. (592-594)	3	3	0	1	1	0	5.3	11.5	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	2	2	0	10.7	23.0	-
Birth inj., atelect. (760-762)	2	1	1	2	1	1	3.7	4.7	3.0
#Other dis. early inf. (763-776)	12	1	11	12	1	11	22.0	4.7	33.0
Violent & acc. deaths (800-999)	19	10	9	16	6	10	85.6	69.0	100.0
Acc. deaths, total (800-962)	14	7	7	12	4	8	64.2	46.0	80.0
Motor vehicle acc. (810-835)	6	3	3	5	2	3	26.7	23.0	30.0
Ill-defined (780-795)	19	4	15	18	3	15	96.2	34.5	150.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HANCOCK COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	101	66	35	262	213	49	20.0	18.0	37.7
Fetal deaths	4	3	1	8	7	1	30.5	32.9	20.4
Deaths from all causes	76	61	15	111	88	23	8.5	7.4	17.7
Deaths under 1 year	1	1	0	6	5	1	22.9	23.5	20.4
Deaths under 1 month	1	1	0	5	4	1	19.1	18.8	20.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	5	5	0	11	9	2	84.0	76.3	153.8
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	18	13	5	21	15	6	160.3	127.1	461.5
Heart diseases (410-443)	28	23	5	43	35	8	328.2	296.6	615.4
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	2	0	4	4	0	30.5	33.9	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	1	1	0	2	1	1	7.6	4.7	20.4
Violent & acc. deaths (800-999)	6	6	0	9	8	1	68.7	67.8	76.9
Acc. deaths, total (800-962)	5	5	0	7	6	1	53.4	50.8	76.9
Motor vehicle acc. (810-835)	2	2	0	0	0	0	-	-	-
Ill-defined (780-795)	13	10	3	12	9	3	91.6	76.3	230.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HARRISON COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	3,529	2,820	709	3,225	2,549	676	30.8	29.1	39.5
Fetal deaths	80	49	31	76	46	30	23.6	18.0	44.4
Deaths from all causes	949	724	225	839	613	226	8.0	7.0	13.2
Deaths under 1 year	99	66	33	95	65	30	29.4	25.5	44.4
Deaths under 1 month	68	51	17	64	49	15	19.8	19.2	22.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	6	4	2	9	5	4	8.6	5.7	23.4
Tbc., respiratory (001-008)	6	4	2	9	5	4	8.6	5.7	23.4
Syphilis (020-029)	3	1	2	1	0	1	1.0	-	5.8
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	1.0	-	5.8
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	1.0	-	5.8
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	115	100	15	102	84	18	97.4	95.9	105.3
Diabetes (260)	3	1	2	3	1	2	2.9	1.1	11.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	100	73	27	95	66	29	90.7	75.3	169.6
Heart diseases (410-443)	345	283	62	294	232	62	280.8	264.8	362.6
Arteriosclerosis (450)	5	4	1	5	5	0	4.8	5.7	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	24	17	7	25	19	6	23.9	21.7	35.1
#Diarrhea & enteritis (571)	4	2	2	4	2	2	3.8	2.3	11.7
Acute nephritis (590, 591)	3	0	3	3	0	3	2.9	-	17.5
Chronic nephritis, etc. (592-594)	8	3	5	9	2	7	8.6	2.3	40.9
Maternal deaths (640-689)	2	2	0	2	2	0	0.6	0.8	-
Cong. malformations (750-759)	9	4	5	14	9	5	13.4	10.3	29.2
Birth inj., atelect. (760-762)	27	20	7	24	19	5	7.4	7.4	7.4
#Other dis. early inf. (763-776)	30	24	6	30	24	6	9.3	9.4	8.9
Violent & acc. deaths (800-999)	85	59	26	69	46	23	65.9	52.5	134.5
Acc. deaths, total (800-962)	60	43	17	48	33	15	45.8	37.7	87.7
Motor vehicle acc. (810-835)	24	20	4	18	13	5	17.2	14.8	29.2
Ill-defined (780-795)	72	44	28	62	35	27	59.2	40.0	157.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HINDS COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	5,596	2,977	2,619	5,336	2,743	2,593	31.2	27.9	35.9
Fetal deaths	360	216	144	329	193	136	61.6	70.4	52.4
Deaths from all causes	1,981	1,055	926	1,433	709	724	8.4	7.2	10.0
Deaths under 1 year	238	112	126	169	68	101	31.7	24.8	39.0
Deaths under 1 month	168	94	74	121	61	60	22.7	22.2	23.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	18	4	14	20	8	12	11.7	8.1	16.6
Tbc., respiratory (001-008)	15	3	12	20	8	12	11.7	8.1	16.6
Syphilis (020-029)	7	2	5	4	0	4	2.3	-	5.5
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	3	0	3	2	0	2	1.2	-	2.8
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	1	0	1	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	321	181	140	205	113	92	120.1	114.8	127.2
Diabetes (260)	19	8	11	13	6	7	7.6	6.1	9.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	203	96	107	157	66	91	92.0	67.1	125.9
Heart diseases (410-443)	541	324	217	434	251	183	254.2	255.1	253.1
Arteriosclerosis (450)	20	11	9	19	10	9	11.1	10.2	12.4
Influenza (480-483)	8	3	5	5	0	5	2.9	-	6.9
#Pneumonia (490-493)	52	25	27	37	14	23	21.7	14.2	31.8
#Diarrhea & enteritis (571)	18	6	12	12	2	10	7.0	2.0	13.8
Acute nephritis (590, 591)	4	1	3	5	2	3	2.9	2.0	4.1
Chronic nephritis, etc. (592-594)	34	13	21	23	5	18	13.5	5.1	24.9
Maternal deaths (640-689)	3	0	3	3	0	3	0.6	-	1.2
Cong. malformations (750-759)	60	40	20	25	15	10	14.6	15.2	13.8
Birth inj., atelect. (760-762)	34	15	19	32	12	20	6.0	4.4	7.7
#Other dis. early inf. (763-776)	88	45	43	66	34	32	12.4	12.4	12.3
Violent & acc. deaths (800-999)	153	83	70	112	61	51	65.6	62.0	70.5
Acc. deaths, total (800-962)	115	65	50	82	46	36	48.0	46.7	49.8
Motor vehicle acc. (810-835)	50	33	17	39	29	10	22.8	29.5	13.8
Ill-defined (780-795)	137	41	96	124	35	89	72.6	35.6	123.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HOLMES COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	810	153	657	825	145	680	29.5	21.3	32.1
Fetal deaths	41	3	38	44	5	39	53.3	34.5	57.4
Deaths from all causes	303	79	224	348	99	249	12.4	14.6	11.7
Deaths under 1 year	49	6	43	49	5	44	59.4	34.5	64.7
Deaths under 1 month	30	5	25	29	4	25	35.2	27.6	36.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	0	3	5	1	4	17.8	14.7	18.9
Tbc., respiratory (001-008)	3	0	3	5	1	4	17.8	14.7	18.9
Syphilis (020-029)	2	0	2	2	0	2	7.1	-	9.4
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	0	2	3	0	3	10.7	-	14.2
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	0	1	1	0	1	3.6	-	4.7
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	34	11	23	39	14	25	139.3	205.9	117.9
Diabetes (260)	5	2	3	6	2	4	21.4	29.4	18.9
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	41	7	34	46	10	36	164.3	147.0	169.8
Heart diseases (410-443)	87	32	55	95	33	62	339.3	485.3	292.4
Arteriosclerosis (450)	6	1	5	7	2	5	25.0	29.4	23.6
Influenza (480-483)	4	2	2	5	2	3	17.8	29.4	14.2
#Pneumonia (490-493)	9	2	7	11	3	8	39.3	44.1	37.7
#Diarrhea & enteritis (571)	2	0	2	3	0	3	10.7	-	14.2
Acute nephritis (590, 591)	1	1	0	1	1	0	3.6	14.7	-
Chronic nephritis, etc. (592-594)	2	0	2	2	0	2	7.1	-	9.4
Maternal deaths (640-689)	0	0	0	1	0	1	1.2	-	1.5
Cong. malformations (750-759)	2	1	1	1	0	1	3.6	-	4.7
Birth inj., atelect. (760-762)	8	2	6	8	2	6	9.7	13.8	8.8
#Other dis. early inf. (763-776)	14	3	11	13	2	11	15.8	13.8	16.2
Violent & acc. deaths (800-999)	20	6	14	27	14	13	96.4	205.9	61.3
Acc. deaths, total (800-962)	19	5	14	24	12	12	85.7	176.5	56.6
Motor vehicle acc. (810-835)	6	0	6	9	6	3	32.1	88.2	14.2
Ill-defined (780-795)	40	1	39	40	1	39	142.8	14.7	184.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HUMPHREYS COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	700	105	595	716	119	597	33.6	19.8	39.0
Fetal deaths	24	2	22	29	3	26	40.5	25.2	43.6
Deaths from all causes	156	35	121	195	44	151	9.2	7.3	9.9
Deaths under 1 year	28	1	27	32	1	31	44.7	8.4	51.9
Deaths under 1 month	13	1	12	17	1	16	23.7	8.4	26.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	4.7	-	6.5
Tbc., respiratory (001-008)	0	0	0	1	0	1	4.7	-	6.5
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	4.7	-	6.5
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	13	5	8	19	6	13	89.2	100.0	85.0
Diabetes (260)	1	0	1	2	0	2	9.4	-	13.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	20	4	16	25	6	19	117.4	100.0	124.2
Heart diseases (410-443)	43	19	24	51	22	29	239.4	366.7	189.5
Arteriosclerosis (450)	1	0	1	1	0	1	4.7	-	6.5
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	3	0	3	5	0	5	23.5	-	32.7
#Diarrhea & enteritis (571)	6	0	6	6	0	6	28.2	-	39.2
Acute nephritis (590, 591)	1	0	1	1	0	1	4.7	-	6.5
Chronic nephritis, etc. (592-594)	7	0	7	8	0	8	37.6	-	52.3
Maternal deaths (640-689)	0	0	0	1	0	1	1.4	-	1.7
Cong. malformations (750-759)	1	0	1	1	0	1	4.7	-	6.5
Birth inj., atelect. (760-762)	1	0	1	1	0	1	1.4	-	1.7
#Other dis. early inf. (763-776)	10	1	9	14	1	13	19.6	8.4	21.8
Violent & acc. deaths (800-999)	12	2	10	18	5	13	84.5	83.3	85.0
Acc. deaths, total (800-962)	9	1	8	13	3	10	61.0	50.0	65.4
Motor vehicle acc. (810-835)	4	1	3	5	2	3	23.5	33.3	19.6
Ill-defined (780-795)	27	0	27	28	0	28	131.4	-	183.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ISSAQENA COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	49	2	47	110	25	85	22.9	13.2	29.3
Fetal deaths	0	0	0	1	0	1	9.1	-	11.8
Deaths from all causes	26	3	23	34	7	27	7.1	3.7	9.3
Deaths under 1 year	4	0	4	8	3	5	72.7	120.0	58.8
Deaths under 1 month	3	0	3	7	3	4	63.6	120.0	47.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	3	0	3	5	1	4	104.2	52.6	137.9
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	4	0	4	6	1	5	125.0	52.6	172.4
Heart diseases (410-443)	7	2	5	7	1	6	145.8	52.6	206.9
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	0	0	0	0	0	0	-	-	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	3	0	3	7	3	4	63.6	120.0	47.0
Violent & acc. deaths (800-999)	4	1	3	4	1	3	83.3	52.6	103.4
Acc. deaths, total (800-962)	4	1	3	4	1	3	83.3	52.6	103.4
Motor vehicle acc. (810-835)	0	0	0	1	1	0	20.8	52.6	-
<u>Ill-defined</u> (780-795)	4	0	4	4	0	4	83.3	-	137.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ITAWAMBA COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	98	78	20	284	261	23	20.9	20.6	25.6
Fetal deaths	3	2	1	12	10	2	42.2	38.3	87.0
Deaths from all causes	96	91	5	121	113	8	8.9	8.9	8.9
Deaths under 1 year	3	2	1	3	2	1	10.6	7.7	43.5
Deaths under 1 month	1	1	0	1	1	0	3.5	3.8	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	7.4	7.9	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	7.4	7.9	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	12	11	1	16	15	1	117.6	118.1	111.1
Diabetes (260)	2	2	0	3	3	0	22.0	23.6	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	17	16	1	20	19	1	147.0	149.6	111.1
Heart diseases (410-443)	32	32	0	40	39	1	294.1	307.1	111.1
Arteriosclerosis (450)	1	1	0	1	1	0	7.4	7.9	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	9	8	1	8	7	1	58.8	55.1	111.1
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	1	0	1	1	0	7.4	7.9	-
Chronic nephritis, etc. (592-594)	3	3	0	3	3	0	22.0	23.6	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	1	1	0	1	1	0	3.5	3.8	-
#Other dis. early inf. (763-776)	0	0	0	0	0	0	-	-	-
Violent & acc. deaths (800-999)	3	3	0	8	7	1	58.8	55.1	111.1
Acc. deaths, total (800-962)	1	1	0	5	4	1	36.8	31.5	111.1
Motor vehicle acc. (810-835)	1	1	0	4	3	1	29.4	23.6	111.1
Ill-defined (780-795)	8	7	1	9	7	2	66.2	55.1	222.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JACKSON COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,100	750	350	1,303	934	369	30.6	27.5	43.4
Fetal deaths	62	36	26	63	37	26	48.3	39.6	70.5
Deaths from all causes	309	198	111	340	226	114	8.0	6.6	13.4
Deaths under 1 year	37	13	24	43	17	26	33.0	18.2	70.5
Deaths under 1 month	27	9	18	31	12	19	23.8	12.8	51.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	2	1	1	4.7	2.9	11.8
Tbc., respiratory (001-008)	2	1	1	2	1	1	4.7	2.9	11.8
Syphilis (020-029)	0	0	0	1	0	1	2.4	-	11.8
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	40	28	12	44	33	11	103.5	97.0	129.4
Diabetes (260)	6	5	1	5	4	1	11.8	11.8	11.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	36	26	10	40	28	12	94.1	82.4	141.2
Heart diseases (410-443)	96	72	24	104	83	21	244.7	244.1	247.0
Arteriosclerosis (450)	6	5	1	7	5	2	16.5	14.7	23.5
Influenza (480-483)	7	6	1	7	6	1	16.5	17.6	11.8
#Pneumonia (490-493)	3	0	3	6	2	4	14.1	5.9	47.0
#Diarrhea & enteritis (571)	2	1	1	2	1	1	4.7	2.9	11.8
Acute nephritis (590, 591)	2	1	1	1	0	1	2.4	-	11.8
Chronic nephritis, etc. (592-594)	1	1	0	2	2	0	4.7	5.9	-
Maternal deaths (640-689)	1	1	0	2	1	1	1.5	1.1	2.7
Cong. malformations (750-759)	8	4	4	9	5	4	21.2	14.7	47.0
Birth inj., atelect. (760-762)	6	2	4	8	3	5	6.1	3.2	13.6
#Other dis. early inf. (763-776)	15	4	11	15	4	11	11.5	4.3	29.8
Violent & acc. deaths (800-999)	36	18	18	35	18	17	82.4	52.9	200.0
Acc. deaths, total (800-962)	29	16	13	28	15	13	65.9	44.1	152.9
Motor vehicle acc. (810-835)	11	6	5	11	5	6	25.9	14.7	70.6
Ill-defined (780-795)	24	14	10	23	13	10	54.1	38.2	117.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JASPER COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	148	45	103	434	145	289	25.7	18.1	32.5
Fetal deaths	8	0	8	22	2	20	50.7	13.8	69.2
Deaths from all causes	94	41	53	162	79	83	9.6	9.9	9.3
Deaths under 1 year	13	0	13	20	1	19	46.1	6.9	65.7
Deaths under 1 month	6	0	6	12	1	11	27.6	6.9	38.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	1	0	1	1	0	5.9	12.5	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	8	6	2	14	9	5	82.8	112.5	56.2
Diabetes (260)	2	0	2	4	1	3	23.7	12.5	33.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	12	6	6	20	13	7	118.3	162.5	78.6
Heart diseases (410-443)	36	20	16	59	34	25	349.1	425.0	280.9
Arteriosclerosis (450)	1	0	1	3	2	1	17.8	25.0	11.2
Influenza (480-483)	4	1	3	4	1	3	23.7	12.5	33.7
#Pneumonia (490-493)	1	0	1	1	0	1	5.9	-	11.2
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	1	0	2	2	0	11.8	25.0	-
Chronic nephritis, etc. (592-594)	1	0	1	2	0	2	11.8	-	22.5
Maternal deaths (640-689)	1	1	0	3	1	2	6.9	6.9	6.9
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	3	0	3	8	0	8	18.4	-	27.7
Violent & acc. deaths (800-999)	11	3	8	19	8	11	112.4	100.0	123.6
Acc. deaths, total (800-962)	8	2	6	16	7	9	94.7	87.5	101.1
Motor vehicle acc. (810-835)	2	1	1	5	3	2	29.6	37.5	22.5
Ill-defined (780-795)	8	1	7	9	2	7	53.2	25.0	78.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JEFFERSON COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence				Rate*		
	Total	White	Non-white	Number			Total	White	Non-white	Rate*
				Total	White	Non-white				
Live births	119	4	115	308	35	273	30.5	14.0	35.9	
Fetal deaths	5	0	5	14	1	13	45.4	28.6	47.6	
Deaths from all causes	88	27	61	108	33	75	10.7	13.2	9.9	
Deaths under 1 year	4	0	4	10	1	9	32.5	28.6	33.0	
Deaths under 1 month	2	0	2	6	1	5	19.5	28.6	18.3	
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.										
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-	-
Cancer, incl. leukemia (140-205)	18	6	12	19	4	15	188.1	160.0	197.4	
Diabetes (260)	1	0	1	1	0	1	9.9	-	13.2	
Pellagra (281)	0	0	0	0	0	0	-	-	-	
Intracranial lesions (330-334)	11	2	9	13	4	9	128.7	160.0	118.4	
Heart diseases (410-443)	34	11	23	38	14	24	376.2	560.0	315.8	
Arteriosclerosis (450)	4	2	2	4	2	2	39.6	80.0	26.3	
Influenza (480-483)	2	0	2	2	0	2	19.8	-	26.3	
#Pneumonia (490-493)	1	0	1	1	0	1	9.9	-	13.2	
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-	
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-	
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	9.9	40.0	-	
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-	
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-	
Birth inj., atelect. (760-762)	0	0	0	1	0	1	3.2	-	3.7	
#Other dis. early inf. (763-776)	3	0	3	8	1	7	26.0	28.6	25.6	
Violent & acc. deaths (800-999)	6	3	3	5	1	4	49.5	40.0	52.6	
Acc. deaths, total (800-962)	4	2	2	3	0	3	29.7	-	39.5	
Motor vehicle acc. (810-835)	1	1	0	0	0	0	-	-	-	
Ill-defined (780-795)	3	1	2	5	1	4	49.5	40.0	52.6	

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JEFF DAVIS COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	384	172	212	383	136	247	23.6	21.2	25.2
Fetal deaths	14	2	12	14	2	12	36.6	14.7	48.6
Deaths from all causes	128	66	62	128	65	63	7.9	10.2	6.4
Deaths under 1 year	11	2	9	9	3	6	23.5	22.0	24.3
Deaths under 1 month	6	1	5	4	2	2	10.4	14.7	8.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	26	13	13	25	13	12	154.3	203.1	122.4
Diabetes (260)	1	1	0	1	1	0	6.2	15.6	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	13	5	8	11	4	7	67.9	62.5	71.4
Heart diseases (410-443)	42	31	11	42	30	12	259.2	468.8	122.4
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	1	0	1	1	0	1	6.2	-	10.2
#Pneumonia (490-493)	2	0	2	3	0	3	18.5	-	30.6
#Diarrhea & enteritis (571)	2	1	1	2	1	1	12.3	15.6	10.2
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	6.2	15.6	-
Maternal deaths (640-689)	1	0	1	1	0	1	2.6	-	4.0
Cong. malformations (750-759)	1	1	0	2	2	0	12.3	31.2	-
Birth inj., atelect. (760-762)	1	0	1	0	0	0	-	-	-
#Other dis. early inf. (763-776)	5	0	5	4	0	4	10.4	-	16.2
Violent & acc. deaths (800-999)	9	7	2	8	5	3	49.4	78.1	30.6
Acc. deaths, total (800-962)	8	6	2	7	4	3	43.2	62.5	30.6
Motor vehicle acc. (810-835)	6	4	2	3	1	2	18.5	15.6	20.4
Ill-defined (780-795)	7	0	7	8	0	8	49.4	-	81.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JONES COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence				Rate*		
	Total	White	Non-white	Number		Non-white	Total	White	Non-white	
				Total	White					
Live births	2,211	1,117	1,094	1,423	870	553	27.6	23.1	40.1	
Fetal deaths	92	36	56	44	24	20	30.9	27.6	36.2	
Deaths from all causes	626	389	237	500	340	160	9.7	9.0	11.6	
Deaths under 1 year	96	38	58	54	17	37	37.9	19.5	66.9	
Deaths under 1 month	65	29	36	31	14	17	21.8	16.1	30.7	
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.										
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-	
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-	
Syphilis (020-029)	0	0	0	0	0	0	-	-	-	
Typhoid fever (040)	0	0	0	0	0	0	-	-	-	
Dysentery (045-048)	0	0	0	0	0	0	-	-	-	
Scarlet fever & streptococcal sore throat (050, 051)	2	1	1	2	1	1	3.9	2.6	7.2	
Diphtheria (055)	0	0	0	0	0	0	-	-	-	
Whooping cough (056)	0	0	0	0	0	0	-	-	-	
Meningococcal infections (057)	1	1	0	0	0	0	-	-	-	
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-	
Smallpox (084)	0	0	0	0	0	0	-	-	-	
Measles (085)	0	0	0	0	0	0	-	-	-	
Malaria (110-117)	0	0	0	0	0	0	-	-	-	
Cancer, incl. leukemia (140-205)	74	55	19	70	56	14	135.9	148.5	101.4	
Diabetes (260)	10	6	4	4	3	1	7.8	8.0	7.2	
Pellagra (281)	0	0	0	0	0	0	-	-	-	
Intracranial lesions (330-334)	83	56	27	64	43	21	124.3	114.0	152.2	
Heart diseases (410-443)	178	125	53	159	120	39	308.7	318.3	282.6	
Arteriosclerosis (450)	5	3	2	4	2	2	7.8	5.3	14.5	
Influenza (480-483)	7	3	4	6	2	4	11.6	5.3	29.0	
#Pneumonia (490-493)	21	2	19	20	5	15	38.8	13.3	108.7	
#Diarrhea & enteritis (571)	7	0	7	8	1	7	15.5	2.6	50.7	
Acute nephritis (590, 591)	5	3	2	3	2	1	5.8	5.3	7.2	
Chronic nephritis, etc. (592-594)	13	5	8	9	6	3	17.5	15.9	21.7	
Maternal deaths (640-689)	7	0	7	2	0	2	1.4	-	3.6	
Cong. malformations (750-759)	9	8	1	4	4	0	7.8	10.6	-	
Birth inj., atelect. (760-762)	24	12	12	13	6	7	9.1	6.9	12.6	
#Other dis. early inf. (763-776)	35	13	22	15	5	10	10.5	5.7	18.1	
Violent & acc. deaths (800-999)	65	50	15	62	47	15	120.4	124.7	108.7	
Acc. deaths, total (800-962)	52	43	9	48	39	9	93.2	103.4	65.2	
Motor vehicle acc. (810-835)	24	21	3	23	19	4	44.7	50.4	29.0	
Ill-defined (780-795)	20	11	9	16	10	6	31.1	26.5	43.5	

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
KEMPER COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	236	48	188	330	74	256	27.7	20.0	31.2
Fetal deaths	13	3	10	19	5	14	57.6	67.6	54.7
Deaths from all causes	93	38	55	122	53	69	10.2	14.3	8.4
Deaths under 1 year	4	0	4	6	0	6	18.2	-	23.4
Deaths under 1 month	1	0	1	3	0	3	9.1	-	11.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	3	1	2	25.2	27.0	24.4
Tbc., respiratory (001-008)	2	0	2	3	1	2	25.2	27.0	24.4
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	8.4	-	12.2
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	12	8	4	15	11	4	126.0	297.3	48.8
Diabetes (260)	5	2	3	5	2	3	42.0	54.0	36.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	11	6	5	20	11	9	168.1	297.3	109.8
Heart diseases (410-443)	22	8	14	31	13	18	260.5	351.4	219.5
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	1	1	0	1	1	0	8.4	27.0	-
#Pneumonia (490-493)	2	0	2	2	0	2	16.8	-	24.4
#Diarrhea & enteritis (571)	2	0	2	2	0	2	16.8	-	24.4
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	1	3	4	1	3	33.6	27.0	36.6
Maternal deaths (640-689)	1	1	0	1	1	0	3.0	13.5	-
Cong. malformations (750-759)	0	0	0	1	1	0	3.0	-	24.4
Birth inj., atelect. (760-762)	0	0	0	2	0	2	16.8	-	24.4
#Other dis. early inf. (763-776)	1	0	1	0	0	0	-	-	-
Violent & acc. deaths (800-999)	10	5	5	13	6	7	109.2	162.2	85.4
Acc. deaths, total (800-962)	8	3	5	10	4	6	84.0	108.1	73.2
Motor vehicle acc. (810-835)	4	3	1	4	2	2	33.6	54.0	24.4
Ill-defined (780-795)	13	2	11	14	2	12	117.6	54.0	146.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LAFAYETTE COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	537	278	259	529	251	278	32.6	31.0	34.3
Fetal deaths	19	4	15	16	3	13	30.2	12.0	46.8
Deaths from all causes	179	95	84	199	117	82	12.3	14.4	10.1
Deaths under 1 year	19	7	12	21	8	13	39.7	31.9	46.8
Deaths under 1 month	13	6	7	16	8	8	30.2	31.9	28.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	0	1	1	0	1	6.2	-	12.3
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	14	8	6	21	15	6	129.6	185.2	74.1
Diabetes (260)	5	2	3	3	2	1	18.5	24.7	12.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	32	21	11	38	26	12	234.6	321.0	148.1
Heart diseases (410-443)	52	40	12	55	42	13	339.5	518.5	160.5
Arteriosclerosis (450)	1	1	0	1	1	0	6.2	12.3	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	8	2	6	8	2	6	49.4	24.7	74.1
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	3	0	3	3	0	3	18.5	-	37.0
Chronic nephritis, etc. (592-594)	6	2	4	5	1	4	30.9	12.3	49.4
Maternal deaths (640-689)	2	0	2	1	0	1	1.9	-	3.6
Cong. malformations (750-759)	0	0	0	3	3	0	18.5	37.0	-
Birth inj., atelect. (760-762)	3	1	2	3	1	2	5.7	4.0	7.2
#Other dis. early inf. (763-776)	12	5	7	14	6	8	26.5	23.9	28.8
Violent & acc. deaths (800-999)	21	4	17	22	7	15	135.8	86.4	185.2
Acc. deaths, total (800-962)	16	1	15	16	3	13	98.8	37.0	160.5
Motor vehicle acc. (810-835)	10	0	10	12	3	9	74.1	37.0	111.1
Ill-defined (780-795)	6	1	5	8	2	6	49.4	24.7	74.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LAMAR COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	130	78	52	345	263	82	28.0	25.3	43.2
Fetal deaths	0	0	0	4	2	2	11.6	7.6	24.4
Deaths from all causes	72	58	14	107	85	22	8.7	8.2	11.6
Deaths under 1 year	2	2	0	7	6	1	20.3	22.8	12.2
Deaths under 1 month	1	1	0	5	4	1	14.5	15.2	12.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	1	1	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	11	11	0	21	19	2	170.7	182.7	105.3
Diabetes (260)	2	2	0	2	2	0	16.3	19.2	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	8	8	0	11	10	1	89.4	96.2	52.6
Heart diseases (410-443)	24	15	9	38	29	9	308.9	278.8	473.7
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	2	1	1	2	1	1	16.3	9.6	52.6
#Pneumonia (490-493)	3	2	1	4	3	1	32.5	28.8	52.6
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	2	1	1	16.3	9.6	52.6
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	1	1	0	1	1	0	2.9	3.8	-
#Other dis. early inf. (763-776)	0	0	0	4	3	1	11.6	11.4	12.2
Violent & acc. deaths (800-999)	8	7	1	6	4	2	48.8	38.5	105.3
Acc. deaths, total (800-962)	6	5	1	3	2	1	24.4	19.2	52.6
Motor vehicle acc. (810-835)	4	3	1	1	0	1	8.1	-	52.6
Ill-defined (780-795)	5	3	2	4	2	2	32.5	19.2	105.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LAUDERDALE COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	2,123	1,088	1,035	1,654	868	786	26.6	22.0	34.6
Fetal deaths	77	29	48	58	21	37	35.1	24.2	47.1
Deaths from all causes	840	536	304	671	409	262	10.8	10.4	11.5
Deaths under 1 year	88	38	50	73	35	38	44.1	40.3	48.3
Deaths under 1 month	67	33	34	54	29	25	32.6	33.4	31.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	3	1	3	1	2	4.8	2.5	8.8
Tbc., respiratory (001-008)	4	3	1	3	1	2	4.8	2.5	8.8
Syphilis (020-029)	2	0	2	2	0	2	3.2	-	8.8
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	1	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	1.6	-	4.4
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	88	66	22	80	61	19	128.8	154.8	83.7
Diabetes (260)	9	5	4	9	5	4	14.5	12.7	17.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	129	87	42	107	71	36	172.3	180.2	158.6
Heart diseases (410-443)	263	181	82	219	141	78	352.7	357.9	343.6
Arteriosclerosis (450)	23	22	1	14	13	1	22.5	33.0	4.4
Influenza (480-483)	3	2	1	2	1	1	3.2	2.5	4.4
#Pneumonia (490-493)	23	18	5	9	5	4	14.5	12.7	17.6
#Diarrhea & enteritis (571)	7	2	5	4	0	4	6.4	-	17.6
Acute nephritis (590, 591)	3	2	1	1	1	0	1.6	2.5	-
Chronic nephritis, etc. (592-594)	2	2	0	3	2	1	4.8	5.1	4.4
Maternal deaths (640-689)	2	0	2	1	0	1	0.6	-	1.3
Cong. malformations (750-759)	10	8	2	9	9	0	14.5	22.8	-
Birth inj., atelect. (760-762)	27	12	15	22	10	12	13.3	11.5	15.3
#Other dis. early inf. (763-776)	29	16	13	25	15	10	15.1	17.3	12.7
Violent & acc. deaths (800-999)	59	31	28	36	19	17	58.0	48.2	74.9
Acc. deaths, total (800-962)	49	29	20	27	17	10	43.5	43.1	44.0
Motor vehicle acc. (810-835)	22	10	12	13	7	6	20.9	17.8	26.4
Ill-defined (780-795)	60	17	43	54	12	42	87.0	30.4	185.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LAWRENCE COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	173	21	152	292	142	150	25.2	20.6	31.9
Fetal deaths	4	0	4	9	2	7	30.8	14.1	46.7
Deaths from all causes	58	34	24	109	68	41	9.4	9.8	8.7
Deaths under 1 year	4	0	4	9	2	7	30.8	14.1	46.7
Deaths under 1 month	2	0	2	6	2	4	20.5	14.1	26.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	2	0	2	17.2	-	42.6
Tbc., respiratory (001-008)	1	0	1	2	0	2	17.2	-	42.6
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	11	6	5	19	12	7	163.8	173.9	148.9
Diabetes (260)	0	0	0	1	1	0	8.6	14.5	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	5	2	3	9	3	6	77.6	43.5	127.6
Heart diseases (410-443)	20	16	4	33	26	7	284.5	376.8	148.9
Arteriosclerosis (450)	1	0	1	1	0	1	8.6	-	21.3
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	6	2	4	8	3	5	69.0	43.5	106.4
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	1	0	1	1	0	8.6	14.5	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	1	0	1	3.4	-	6.7
Cong. malformations (750-759)	1	1	0	3	2	1	25.9	29.0	21.3
Birth inj., atelect. (760-762)	0	0	0	1	0	1	3.4	-	6.7
#Other dis. early inf. (763-776)	1	0	1	4	2	2	13.7	14.1	13.3
Violent & acc. deaths (800-999)	5	4	1	12	10	2	103.4	144.9	42.6
Acc. deaths, total (800-962)	5	4	1	12	10	2	103.4	144.9	42.6
Motor vehicle acc. (810-835)	4	3	1	9	7	2	77.6	101.4	42.6
Ill-defined (780-795)	3	2	1	4	3	1	34.5	43.5	21.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LEAKE COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence				Rate*		
	Total	White	Non-white	Number			Non-white	Total	White	Non-white
				Total	White	Non-white				
Live births	429	191	238	466	205	261	30.1	25.0	35.8	
Fetal deaths	4	0	4	8	1	7	17.2	4.9	26.8	
Deaths from all causes	177	110	67	213	136	77	13.7	16.6	10.5	
Deaths under 1 year	23	8	15	27	9	18	57.9	43.9	69.0	
Deaths under 1 month	13	5	8	17	6	11	36.5	29.3	42.1	
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.										
Tbc., all forms (001-019)	1	1	0	2	1	1	12.9	12.2	13.7	
Tbc., respiratory (001-008)	1	1	0	1	1	0	6.4	12.2	-	
Syphilis (020-029)	0	0	0	0	0	0	-	-	-	
Typhoid fever (040)	0	0	0	0	0	0	-	-	-	
Dysentery (045-048)	0	0	0	0	0	0	-	-	-	
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-	
Diphtheria (055)	0	0	0	0	0	0	-	-	-	
Whooping cough (056)	0	0	0	0	0	0	-	-	-	
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-	
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-	
Smallpox (084)	0	0	0	0	0	0	-	-	-	
Measles (085)	0	0	0	0	0	0	-	-	-	
Malaria (110-117)	0	0	0	0	0	0	-	-	-	
Cancer, incl. leukemia (140-205)	17	12	5	23	17	6	148.4	207.3	82.2	
Diabetes (260)	3	2	1	4	2	2	25.8	24.4	27.4	
Pellagra (281)	0	0	0	0	0	0	-	-	-	
Intracranial lesions (330-334)	37	24	13	39	25	14	251.6	304.9	191.8	
Heart diseases (410-443)	56	39	17	61	44	17	393.5	536.6	232.9	
Arteriosclerosis (450)	3	1	2	4	2	2	25.8	24.4	27.4	
Influenza (480-483)	2	1	1	2	1	1	12.9	12.2	13.7	
#Pneumonia (490-493)	7	2	5	8	3	5	51.6	36.6	68.5	
#Diarrhea & enteritis (571)	2	1	1	2	1	1	12.9	12.2	13.7	
Acute nephritis (590, 591)	2	1	1	2	1	1	12.9	12.2	13.7	
Chronic nephritis, etc. (592-594)	3	2	1	3	2	1	19.4	24.4	13.7	
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-	
Cong. malformations (750-759)	4	4	0	5	5	0	32.2	61.0	-	
Birth inj., atelect. (760-762)	3	1	2	5	1	4	10.7	4.9	15.3	
#Other dis. early inf. (763-776)	8	3	5	9	3	6	19.3	14.6	23.0	
Violent & acc. deaths (800-999)	10	6	4	18	13	5	116.1	158.5	68.5	
Acc. deaths, total (800-962)	7	4	3	13	9	4	83.9	109.8	54.8	
Motor vehicle acc. (810-835)	2	1	1	5	3	2	32.2	36.6	27.4	
Ill-defined (780-795)	9	2	7	11	3	8	71.0	36.6	109.6	

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LEE COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,256	878	378	1,013	648	365	27.1	24.9	32.0
Fetal deaths	55	34	21	43	24	19	42.4	37.0	52.0
Deaths from all causes	381	266	115	363	247	116	9.7	9.5	10.2
Deaths under 1 year	38	25	13	29	18	11	28.6	27.8	30.1
Deaths under 1 month	26	20	6	19	14	5	18.8	21.6	13.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	1	2	5	1	4	13.4	3.8	35.1
Tbc., respiratory (001-008)	3	1	2	5	1	4	13.4	3.8	35.1
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	60	46	14	59	44	15	157.8	169.2	131.6
Diabetes (260)	4	2	2	4	2	2	10.7	7.7	17.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	46	33	13	45	31	14	120.3	119.2	122.8
Heart diseases (410-443)	121	89	32	114	80	34	304.8	307.7	298.2
Arteriosclerosis (450)	2	2	0	3	3	0	8.0	11.5	-
Influenza (480-483)	6	2	4	6	2	4	16.0	7.7	35.1
#Pneumonia (490-493)	24	11	13	23	11	12	61.5	42.3	105.3
#Diarrhea & enteritis (571)	2	1	1	1	0	1	2.7	-	8.8
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	7	6	1	5	4	1	13.4	15.4	8.8
Maternal deaths (640-689)	2	1	1	2	1	1	2.0	1.5	2.7
Cong. malformations (750-759)	3	3	0	4	4	0	10.7	15.4	-
Birth inj., atelect. (760-762)	10	8	2	7	5	2	6.9	7.7	5.5
#Other dis. early inf. (763-776)	13	11	2	10	8	2	9.9	12.3	5.5
Violent & acc. deaths (800-999)	34	24	10	34	25	9	90.9	96.2	78.9
Acc. deaths, total (800-962)	26	20	6	26	22	4	69.5	84.6	35.1
Motor vehicle acc. (810-835)	11	8	3	10	9	1	26.7	34.6	8.8
Ill-defined (780-795)	14	5	9	12	4	8	32.1	15.4	70.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LEFLORE COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
					White	Non-white			
Live births	1,701	439	1,262	1,647	355	1,292	35.8	22.5	42.8
Fetal deaths	86	25	61	81	23	58	49.2	64.8	44.9
Deaths from all causes	568	207	361	517	153	364	11.2	9.7	12.0
Deaths under 1 year	76	7	69	72	5	67	43.7	14.1	51.8
Deaths under 1 month	43	4	39	40	3	37	24.3	8.4	28.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	3	0	3	6.5	-	9.9
Tbc., respiratory (001-008)	0	0	0	2	0	2	4.3	-	6.6
Syphilis (020-029)	0	0	0	1	0	1	2.2	-	3.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	0	2	3	0	3	6.5	-	9.9
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	57	27	30	56	23	33	121.7	145.6	109.3
Diabetes (260)	6	1	5	4	1	3	8.7	6.3	9.9
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	96	47	49	78	29	49	169.6	183.5	162.2
Heart diseases (410-443)	135	63	72	119	45	74	258.7	284.8	245.0
Arteriosclerosis (450)	7	5	2	3	1	2	6.5	6.3	6.6
Influenza (480-483)	2	0	2	2	0	2	4.3	-	6.6
#Pneumonia (490-493)	24	7	17	22	5	17	47.8	31.6	56.3
#Diarrhea & enteritis (571)	1	0	1	1	0	1	2.2	-	3.3
Acute nephritis (590, 591)	3	1	2	3	1	2	6.5	6.3	6.6
Chronic nephritis, etc. (592-594)	7	3	4	6	2	4	13.0	12.7	13.2
Maternal deaths (640-689)	3	0	3	2	0	2	1.2	-	1.5
Cong. malformations (750-759)	4	2	2	5	4	1	10.9	25.3	3.3
Birth inj., atelect. (760-762)	4	1	3	4	1	3	2.4	2.8	2.3
#Other dis. early inf. (763-776)	29	2	27	27	1	26	16.4	2.8	20.1
Violent & acc. deaths (800-999)	37	14	23	36	12	24	78.3	75.9	79.5
Acc. deaths, total (800-962)	26	9	17	24	7	17	52.2	44.3	56.3
Motor vehicle acc. (810-835)	11	5	6	9	3	6	19.6	19.0	19.9
Ill-defined (780-795)	103	13	90	99	10	89	215.2	63.3	294.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LINCOLN COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	693	396	297	669	360	309	25.5	21.6	32.5
Fetal deaths	18	10	8	21	10	11	31.4	27.8	35.6
Deaths from all causes	232	144	88	263	165	98	10.0	9.9	10.3
Deaths under 1 year	25	10	15	27	10	17	40.4	27.8	55.0
Deaths under 1 month	15	8	7	18	9	9	26.9	25.0	29.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	3.8	-	10.5
Tbc., respiratory (001-008)	1	0	1	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	1	1	0	3.8	6.0	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	3.8	-	10.5
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	0	1	1	0	1	3.8	-	10.5
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	20	17	3	21	17	4	80.2	101.8	42.1
Diabetes (260)	8	6	2	8	5	3	30.5	29.9	31.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	29	13	16	35	16	19	133.6	95.8	200.0
Heart diseases (410-443)	101	69	32	101	70	31	385.5	419.2	326.3
Arteriosclerosis (450)	1	1	0	1	1	0	3.8	6.0	-
Influenza (480-483)	1	0	1	2	1	1	7.6	6.0	10.5
#Pneumonia (490-493)	12	4	8	12	5	7	45.8	29.9	73.7
#Diarrhea & enteritis (571)	2	1	1	1	0	1	3.8	-	10.5
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	2	1	4	3	1	15.3	18.0	10.5
Maternal deaths (640-689)	2	1	1	1	1	0	1.5	2.8	-
Cong. malformations (750-759)	5	4	1	4	3	1	15.3	18.0	10.5
Birth inj., atelect. (760-762)	5	2	3	6	3	3	9.0	8.3	9.7
#Other dis. early inf. (763-776)	5	3	2	6	3	3	9.0	8.3	9.7
Violent & acc. deaths (800-999)	19	12	7	28	19	9	106.9	113.8	94.7
Acc. deaths, total (800-962)	16	9	7	25	16	9	95.4	95.8	94.7
Motor vehicle acc. (810-835)	9	7	2	13	10	3	49.6	59.9	31.6
Ill-defined (780-795)	7	2	5	6	1	5	22.9	6.0	52.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LOWNDES COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,175	548	627	1,154	523	631	30.8	25.1	38.0
Fetal deaths	41	18	23	39	17	22	33.8	32.5	34.9
Deaths from all causes	375	177	198	393	186	207	10.5	8.9	12.5
Deaths under 1 year	46	18	28	40	16	24	34.7	30.6	38.0
Deaths under 1 month	26	14	12	23	12	11	19.9	22.9	17.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	2	1	1	5.3	4.8	6.0
Tbc., respiratory (001-008)	1	0	1	2	1	1	5.3	4.8	6.0
Syphilis (020-029)	2	1	1	1	1	0	2.7	4.8	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	1	1	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	35	24	11	43	31	12	115.0	149.0	72.3
Diabetes (260)	3	2	1	3	2	1	8.0	9.6	6.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	35	13	22	44	18	26	117.6	86.5	156.6
Heart diseases (410-443)	118	63	55	122	65	57	326.2	312.5	343.4
Arteriosclerosis (450)	6	5	1	5	4	1	13.4	19.2	6.0
Influenza (480-483)	7	1	6	8	1	7	21.4	4.8	42.2
#Pneumonia (490-493)	12	4	8	13	4	9	34.8	19.2	54.2
#Diarrhea & enteritis (571)	6	0	6	3	0	3	8.0	-	18.1
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	5	1	4	6	1	5	16.0	4.8	30.1
Maternal deaths (640-689)	3	0	3	3	0	3	2.6	-	4.8
Cong. malformations (750-759)	3	3	0	4	4	0	10.7	19.2	-
Birth inj., atelect. (760-762)	15	9	6	13	8	5	11.3	15.3	7.9
#Other dis. early inf. (763-776)	9	2	7	9	2	7	7.8	3.8	11.1
Violent & acc. deaths (800-999)	32	18	14	25	10	15	66.8	48.1	90.4
Acc. deaths, total (800-962)	23	12	11	16	5	11	42.8	24.0	66.3
Motor vehicle acc. (810-835)	15	9	6	9	3	6	24.1	14.4	36.1
Ill-defined (780-795)	42	7	35	44	8	36	117.6	38.5	216.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MADISON COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,099	188	911	1,088	199	889	33.7	24.6	36.7
Fetal deaths	52	5	47	51	7	44	46.9	35.2	49.5
Deaths from all causes	298	87	211	315	87	228	9.8	10.7	9.4
Deaths under 1 year	32	3	29	38	7	31	34.9	35.2	34.9
Deaths under 1 month	20	3	17	26	6	20	23.9	30.2	22.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	2	0	2	2	0	2	6.2	-	8.3
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	31	16	15	30	12	18	92.9	148.1	74.4
Diabetes (260)	5	0	5	6	0	6	18.6	-	24.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	51	12	39	55	12	43	170.3	148.1	177.7
Heart diseases (410-443)	84	32	52	88	34	54	272.4	419.8	223.1
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	3	1	2	3	1	2	9.3	12.3	8.3
#Pneumonia (490-493)	8	0	8	8	0	8	24.8	-	33.0
#Diarrhea & enteritis (571)	3	1	2	3	1	2	9.3	12.3	8.3
Acute nephritis (590, 591)	1	0	1	1	0	1	3.1	-	4.1
Chronic nephritis, etc. (592-594)	10	3	7	10	3	7	31.0	37.0	28.9
Maternal deaths (640-689)	2	0	2	2	0	2	1.8	-	2.2
Cong. malformations (750-759)	2	1	1	5	3	2	15.5	37.0	8.3
Birth inj., atelect. (760-762)	3	0	3	3	0	3	2.8	-	3.4
#Other dis. early inf. (763-776)	11	2	9	13	3	10	11.9	15.1	11.2
Violent & acc. deaths (800-999)	27	12	15	26	9	17	80.5	111.1	70.2
Acc. deaths, total (800-962)	20	9	11	18	6	12	55.7	74.1	49.6
Motor vehicle acc. (810-835)	18	9	9	12	4	8	37.2	49.4	33.0
Ill-defined (780-795)	39	4	35	39	4	35	120.7	49.4	144.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MARION COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	540	298	242	611	333	278	26.9	25.0	29.6
Fetal deaths	20	10	10	26	11	15	42.6	33.0	54.0
Deaths from all causes	180	106	74	204	121	83	9.0	9.1	8.8
Deaths under 1 year	15	4	11	16	5	11	26.2	15.0	39.6
Deaths under 1 month	10	3	7	11	4	7	18.0	12.0	25.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	4	2	2	17.6	15.0	21.3
Tbc., respiratory (001-008)	1	1	0	4	2	2	17.6	15.0	21.3
Syphilis (020-029)	3	1	2	3	1	2	13.2	7.5	21.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)-	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	1	0	1	1	0	4.4	7.5	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	38	24	14	35	22	13	154.2	165.4	138.3
Diabetes (260)	2	2	0	3	3	0	13.2	22.6	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	30	21	9	28	20	8	123.3	150.4	85.1
Heart diseases (410-443)	44	23	21	51	26	25	224.7	195.5	266.0
Arteriosclerosis (450)	1	1	0	1	1	0	4.4	7.5	-
Influenza (480-483)	1	0	1	1	0	1	4.4	-	10.6
#Pneumonia (490-493)	6	0	6	8	2	6	35.2	15.0	63.8
#Diarrhea & enteritis (571)	1	0	1	1	0	1	4.4	-	10.6
Acute nephritis (590, 591)	1	1	0	1	1	0	4.4	7.5	-
Chronic nephritis, etc. (592-594)	3	2	1	4	2	2	17.6	15.0	21.3
Maternal deaths (640-689)	0	0	0	1	0	1	1.6	-	3.6
Cong. malformations (750-759)	2	1	1	2	1	1	8.8	7.5	10.6
Birth inj., atelect. (760-762)	4	2	2	4	2	2	6.5	6.0	7.2
#Other dis. early inf. (763-776)	3	1	2	4	2	2	6.5	6.0	7.2
Violent & acc. deaths (800-999)	13	7	6	21	14	7	92.5	105.3	74.5
Acc. deaths, total (800-962)	11	7	4	18	13	5	79.3	97.7	53.2
Motor vehicle acc. (810-835)	6	4	2	9	6	3	39.6	45.1	31.9
Ill-defined (780-795)	13	7	6	12	6	6	52.9	45.1	63.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MARSHALL COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence				Rate*		
	Total	White	Non-white	Number			Total	White	Non-white	Rate*
Live births	769	101	668	814	152	662	35.5	23.4	40.4	
Fetal deaths	55	5	50	44	4	40	54.0	26.3	60.4	
Deaths from all causes	215	60	155	216	60	156	9.4	9.2	9.5	
Deaths under 1 year	30	1	29	29	3	26	35.6	19.7	39.3	
Deaths under 1 month	20	1	19	17	2	15	20.9	13.2	22.6	
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.										
Tbc., all forms (001-019)	0	0	0	1	1	0	4.4	15.4	-	
Tbc., respiratory (001-008)	0	0	0	1	1	0	4.4	15.4	-	
Syphilis (020-029)	2	0	2	1	0	1	4.4	-	6.1	
Typhoid fever (040)	0	0	0	0	0	0	-	-	-	
Dysentery (045-048)	0	0	0	0	0	0	-	-	-	
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-	
Diphtheria (055)	0	0	0	0	0	0	-	-	-	
Whooping cough (056)	0	0	0	0	0	0	-	-	-	
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-	
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-	
Smallpox (084)	0	0	0	0	0	0	-	-	-	
Measles (085)	0	0	0	0	0	0	-	-	-	
Malaria (110-117)	0	0	0	0	0	0	-	-	-	
Cancer, incl. leukemia (140-205)	19	9	10	23	10	13	100.4	153.8	79.3	
Diabetes (260)	4	0	4	3	0	3	13.1	-	18.3	
Pellagra (281)	0	0	0	0	0	0	-	-	-	
Intracranial lesions (330-334)	34	10	24	35	10	25	152.8	153.8	152.4	
Heart diseases (410-443)	56	21	35	61	22	39	266.4	338.5	237.8	
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-	
Influenza (480-483)	1	1	0	1	1	0	4.4	15.4	-	
#Pneumonia (490-493)	4	0	4	4	0	4	17.5	-	24.4	
#Diarrhea & enteritis (571)	3	0	3	3	0	3	13.1	-	18.3	
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-	
Chronic nephritis, etc. (592-594)	2	1	1	1	1	0	4.4	15.4	-	
Maternal deaths (640-689)	3	0	3	1	0	1	1.2	-	1.5	
Cong. malformations (750-759)	4	1	3	4	0	4	17.5	-	24.4	
Birth inj., atelect. (760-762)	4	1	3	4	1	3	4.9	6.6	4.5	
#Other dis. early inf. (763-776)	11	0	11	8	1	7	9.8	6.6	10.6	
Violent & acc. deaths (800-999)	20	8	12	15	4	11	65.5	61.5	67.1	
Acc. deaths, total (800-962)	15	5	10	11	1	10	48.0	15.4	61.0	
Motor vehicle acc. (810-835)	11	5	6	7	1	6	30.6	15.4	36.6	
Ill-defined (780-795)	33	2	31	34	3	31	148.5	46.2	189.0	

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MONROE COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence				Rate*		
	Total	White	Non-white	Number			Total	White	Non-white	
				Total	White	Non-white				
Live births	807	415	392	778	378	400	25.6	19.9	35.1	
Fetal deaths	18	7	11	18	7	11	23.1	18.5	27.5	
Deaths from all causes	262	159	103	265	165	100	8.7	8.7	8.8	
Deaths under 1 year	23	10	13	20	8	12	25.7	21.2	30.0	
Deaths under 1 month	10	5	5	9	4	5	11.6	10.6	12.5	
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.										
Tbc., all forms (001-019)	0	0	0	1	1	0	3.3	5.3	-	
Tbc., respiratory (001-008)	0	0	0	1	1	0	3.3	5.3	-	
Syphilis (020-029)	0	0	0	0	0	0	-	-	-	
Typhoid fever (040)	0	0	0	0	0	0	-	-	-	
Dysentery (045-048)	0	0	0	0	0	0	-	-	-	
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-	
Diphtheria (055)	0	0	0	0	0	0	-	-	-	
Whooping cough (056)	0	0	0	0	0	0	-	-	-	
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-	
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-	
Smallpox (084)	0	0	0	0	0	0	-	-	-	
Measles (085)	0	0	0	0	0	0	-	-	-	
Malaria (110-117)	0	0	0	0	0	0	-	-	-	
Cancer, incl. leukemia (140-205)	35	30	5	33	28	5	108.6	147.4	43.8	
Diabetes (260)	8	5	3	7	4	3	23.0	21.0	26.3	
Pellagra (281)	0	0	0	0	0	0	-	-	-	
Intracranial lesions (330-334)	32	18	14	37	22	15	121.7	115.8	131.6	
Heart diseases (410-443)	81	53	28	80	54	26	263.2	284.2	228.1	
Arteriosclerosis (450)	2	1	1	2	1	1	6.6	5.3	8.8	
Influenza (480-483)	1	0	1	1	0	1	3.3	-	8.8	
#Pneumonia (490-493)	9	5	4	10	7	3	32.9	36.8	26.3	
#Diarrhea & enteritis (571)	1	1	0	1	1	0	3.3	5.3	-	
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-	
Chronic nephritis, etc. (592-594)	13	4	9	12	4	8	39.5	21.0	70.2	
Maternal deaths (640-689)	0	0	0	1	0	1	1.3	-	2.5	
Cong. malformations (750-759)	2	1	1	1	0	1	3.3	-	8.8	
Birth inj., atelect. (760-762)	4	3	1	4	3	1	5.1	7.9	2.5	
#Other dis. early inf. (763-776)	8	2	6	5	1	4	6.4	2.6	10.0	
Violent & acc. deaths (800-999)	21	16	5	22	17	5	72.4	89.5	43.8	
Acc. deaths, total (800-962)	17	14	3	18	15	3	59.2	78.9	26.3	
Motor vehicle acc. (810-835)	9	8	1	10	9	1	32.9	47.4	8.8	
Ill-defined (780-795)	26	9	17	26	9	17	85.5	47.4	149.1	

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MONTGOMERY COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	319	108	211	347	124	223	25.1	16.8	34.8
Fetal deaths	12	4	8	14	4	10	40.3	32.2	44.8
Deaths from all causes	119	62	57	147	82	65	10.6	11.1	10.2
Deaths under 1 year	8	1	7	10	1	9	28.8	8.1	40.4
Deaths under 1 month	6	1	5	7	1	6	20.2	8.1	26.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	7.2	13.5	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	7.2	13.5	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	18	10	8	21	12	9	152.2	162.2	140.6
Diabetes (260)	2	2	0	4	2	2	29.0	27.0	31.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	19	10	9	25	17	8	181.2	229.7	125.0
Heart diseases (410-443)	41	26	15	46	29	17	333.3	391.9	265.6
Arteriosclerosis (450)	0	0	0	1	1	0	7.2	13.5	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	7	2	5	10	4	6	72.5	54.0	93.8
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	0	1	7.2	-	15.6
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	3	0	3	21.7	-	46.9
Birth inj., atelect. (760-762)	4	0	4	4	0	4	11.5	-	17.9
#Other dis. early inf. (763-776)	3	1	2	3	1	2	8.6	8.1	9.0
Violent & acc. deaths (800-999)	9	6	3	10	6	4	72.5	81.1	62.5
Acc. deaths, total (800-962)	7	4	3	8	4	4	58.0	54.0	62.5
Motor vehicle acc. (810-835)	4	2	2	4	2	2	29.0	27.0	31.2
Ill-defined (780-795)	6	0	6	6	1	5	43.5	13.5	78.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
NESHOBIA COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence				Rate*		
	Total	White	Non-white	Number			Non-white	Total	White	Non-white
Live births	567	311	256	593	349	244	32.6	27.5	44.4	
Fetal deaths	12	2	10	10	1	9	16.9	2.9	36.9	
Deaths from all causes	175	113	62	191	128	63	10.5	10.1	11.4	
Deaths under 1 year	20	5	15	23	8	15	38.8	22.9	61.5	
Deaths under 1 month	12	4	8	14	7	7	23.6	20.0	28.7	
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.										
Tbc., all forms (001-019)	1	1	0	2	2	0	11.0	15.7	-	
Tbc., respiratory (001-008)	0	0	0	1	1	0	5.5	7.9	-	
Syphilis (020-029)	0	0	0	0	0	0	-	-	-	
Typhoid fever (040)	0	0	0	0	0	0	-	-	-	
Dysentery (045-048)	0	0	0	1	0	1	5.5	-	18.2	
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-	
Diphtheria (055)	0	0	0	0	0	0	-	-	-	
Whooping cough (056)	0	0	0	0	0	0	-	-	-	
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-	
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-	
Smallpox (084)	0	0	0	0	0	0	-	-	-	
Measles (085)	0	0	0	0	0	0	-	-	-	
Malaria (110-117)	0	0	0	0	0	0	-	-	-	
Cancer, incl. leukemia (140-205)	19	14	5	20	16	4	109.9	126.0	72.7	
Diabetes (260)	4	2	2	2	2	0	11.0	15.7	-	
Pellagra (281)	0	0	0	0	0	0	-	-	-	
Intracranial lesions (330-334)	31	26	5	37	30	7	203.3	236.2	127.3	
Heart diseases (410-443)	56	40	16	55	41	14	302.2	322.8	254.5	
Arteriosclerosis (450)	1	1	0	1	1	0	5.5	7.9	-	
Influenza (480-483)	0	0	0	1	1	0	5.5	7.9	-	
#Pneumonia (490-493)	4	2	2	6	1	5	33.0	7.9	90.9	
#Diarrhea & enteritis (571)	2	0	2	2	0	2	11.0	-	36.4	
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-	
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-	
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-	
Cong. malformations (750-759)	1	1	0	4	2	2	22.0	15.7	36.4	
Birth inj., atelect. (760-762)	6	1	5	5	2	3	8.4	5.7	12.3	
#Other dis. early inf. (763-776)	5	2	3	5	3	2	8.4	8.6	8.2	
Violent & acc. deaths (800-999)	22	10	12	17	5	12	93.4	39.4	218.2	
Acc. deaths, total (800-962)	17	6	11	12	3	9	65.9	23.6	163.6	
Motor vehicle acc. (810-835)	9	3	6	6	0	6	33.0	-	109.1	
Ill-defined (780-795)	14	7	7	17	9	8	93.4	70.9	145.4	

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
NEWTON COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence				Rate*		
	Total	White	Non-white	Number			Total	White	Non-white	
				Total	White	Non-white				
Live births	460	275	185	459	236	223	24.2	21.1	28.6	
Fetal deaths	36	21	15	39	20	19	85.0	84.7	85.2	
Deaths from all causes	156	105	51	167	106	61	8.8	9.5	7.8	
Deaths under 1 year	10	4	6	14	3	11	30.5	12.7	49.3	
Deaths under 1 month	7	2	5	8	2	6	17.4	8.5	26.9	
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.										
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-	
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-	
Syphilis (020-029)	1	0	1	1	0	1	5.3	-	12.8	
Typhoid fever (040)	0	0	0	0	0	0	-	-	-	
Dysentery (045-048)	0	0	0	0	0	0	-	-	-	
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-	
Diphtheria (055)	0	0	0	0	0	0	-	-	-	
Whooping cough (056)	0	0	0	0	0	0	-	-	-	
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-	
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-	
Smallpox (084)	0	0	0	0	0	0	-	-	-	
Measles (085)	0	0	0	0	0	0	-	-	-	
Malaria (110-117)	0	0	0	0	0	0	-	-	-	
Cancer, incl. leukemia (140-205)	20	14	6	23	14	9	121.0	125.0	115.4	
Diabetes (260)	1	1	0	0	0	0	-	-	-	
Pellagra (281)	0	0	0	0	0	0	-	-	-	
Intracranial lesions (330-334)	26	18	8	26	19	7	136.8	169.6	89.7	
Heart diseases (410-443)	48	35	13	49	34	15	257.9	303.6	192.3	
Arteriosclerosis (450)	2	0	2	5	3	2	26.3	26.8	25.6	
Influenza (480-483)	0	0	0	0	0	0	-	-	-	
#Pneumonia (490-493)	2	1	1	5	4	1	26.3	35.7	12.8	
#Diarrhea & enteritis (571)	0	0	0	1	0	1	5.3	-	12.8	
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-	
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-	
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-	
Cong. malformations (750-759)	3	3	0	3	3	0	15.8	26.8	-	
Birth inj., atelect. (760-762)	2	0	2	4	0	4	8.7	-	17.9	
#Other dis. early inf. (763-776)	2	0	2	1	0	1	2.2	-	4.5	
Violent & acc. deaths (800-999)	19	13	6	19	11	8	100.0	98.2	102.6	
Acc. deaths, total (800-962)	17	11	6	15	9	6	78.9	80.4	76.9	
Motor vehicle acc. (810-835)	8	7	1	4	3	1	21.0	26.8	12.8	
Ill-defined (780-795)	10	4	6	13	6	7	68.4	53.6	89.7	

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
NOXUBEE COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	557	87	470	544	84	460	31.6	17.5	37.1
Fetal deaths	30	11	19	28	9	19	51.5	107.1	41.3
Deaths from all causes	172	58	114	187	65	122	10.9	13.5	9.8
Deaths under 1 year	17	3	14	19	5	14	34.9	59.5	30.4
Deaths under 1 month	14	3	11	16	5	11	29.4	59.5	23.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	2	1	1	11.6	20.8	8.1
Tbc., respiratory (001-008)	0	0	0	2	1	1	11.6	20.8	8.1
Syphilis (020-029)	0	0	0	1	0	1	5.8	-	8.1
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	1	0	1	1	0	5.8	20.8	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	22	8	14	24	8	16	139.5	166.7	129.0
Diabetes (260)	2	2	0	3	2	1	17.4	41.7	8.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	32	15	17	35	15	20	203.5	312.5	161.3
Heart diseases (410-443)	45	19	26	49	23	26	284.9	479.2	209.7
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	4	0	4	4	0	4	23.2	-	32.2
#Pneumonia (490-493)	4	0	4	4	0	4	23.2	-	32.2
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	1	1	0	5.8	20.8	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	5.8	-	8.1
Maternal deaths (640-689)	1	0	1	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	1	0	1	5.8	-	8.1
Birth inj., atelect. (760-762)	1	1	0	2	2	0	3.7	23.8	-
#Other dis. early inf. (763-776)	10	2	8	10	2	8	18.4	23.8	17.4
Violent & acc. deaths (800-999)	9	4	5	9	3	6	52.3	62.5	48.4
Acc. deaths, total (800-962)	8	4	4	8	3	5	46.5	62.5	40.3
Motor vehicle acc. (810-835)	2	1	1	2	0	2	11.6	-	16.1
Ill-defined (780-795)	25	2	23	23	1	22	133.7	20.8	177.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
OKTIBBEHA COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	894	383	511	801	314	487	36.4	28.0	45.1
Fetal deaths	14	5	9	13	3	10	16.2	9.6	20.5
Deaths from all causes	189	79	110	200	87	113	9.1	7.8	10.5
Deaths under 1 year	27	8	19	27	8	19	33.7	25.5	39.0
Deaths under 1 month	17	6	11	17	6	11	21.2	19.1	22.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	2	1	1	9.1	8.9	9.2
Tbc., respiratory (001-008)	2	1	1	2	1	1	9.1	8.9	9.2
Syphilis (020-029)	1	0	1	1	0	1	4.5	-	9.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	4.5	-	9.2
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	22	16	6	24	17	7	109.1	151.8	64.8
Diabetes (260)	3	1	2	3	1	2	13.6	8.9	18.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	33	13	20	32	15	17	145.4	133.9	157.4
Heart diseases (410-443)	51	23	28	53	24	29	240.9	214.3	268.5
Arteriosclerosis (450)	1	0	1	1	0	1	4.5	-	9.2
Influenza (480-483)	4	1	3	4	1	3	18.2	8.9	27.8
#Pneumonia (490-493)	1	0	1	2	0	2	9.1	-	18.5
#Diarrhea & enteritis (571)	1	0	1	1	0	1	4.5	-	9.2
Acute nephritis (590, 591)	1	1	0	1	1	0	4.5	8.9	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	1	1	0	1	1	0	1.2	3.2	-
Cong. malformations (750-759)	2	1	1	3	2	1	13.6	17.8	9.2
Birth inj., atelect. (760-762)	2	0	2	2	0	2	2.5	-	4.1
#Other dis. early inf. (763-776)	11	6	5	10	5	5	12.5	15.9	10.3
Violent & acc. deaths (800-999)	23	10	13	27	13	14	122.7	116.1	129.6
Acc. deaths, total (800-962)	22	10	12	26	13	13	118.2	116.1	120.4
Motor vehicle acc. (810-835)	4	4	0	9	8	1	40.9	71.4	9.2
Ill-defined (780-795)	19	0	19	18	0	18	81.8	-	166.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PANOLA COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	887	257	630	909	256	653	28.3	19.8	34.0
Fetal deaths	24	9	15	29	13	16	31.9	50.8	24.5
Deaths from all causes	249	92	157	303	127	176	9.4	9.8	9.2
Deaths under 1 year	29	2	27	33	3	30	36.3	11.7	45.9
Deaths under 1 month	15	1	14	17	1	16	18.7	3.9	24.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	3.1	-	5.2
Tbc., respiratory (001-008)	1	0	1	1	0	1	3.1	-	5.2
Syphilis (020-029)	0	0	0	2	1	1	6.2	7.8	5.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	0	2	1	0	1	3.1	-	5.2
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	2	2	0	2	2	0	6.2	15.5	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	32	12	20	38	16	22	118.4	124.0	114.6
Diabetes (260)	2	1	1	5	2	3	15.6	15.5	15.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	37	16	21	43	21	22	134.0	162.8	114.6
Heart diseases (410-443)	84	40	44	96	49	47	299.1	379.8	244.8
Arteriosclerosis (450)	4	1	3	4	1	3	12.5	7.8	15.6
Influenza (480-483)	1	0	1	1	0	1	3.1	-	5.2
#Pneumonia (490-493)	7	2	5	9	3	6	28.0	23.2	31.2
#Diarrhea & enteritis (571)	3	0	3	3	0	3	9.3	-	15.6
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	0	4	6	1	5	18.7	7.8	26.0
Maternal deaths (640-689)	1	0	1	2	1	1	2.2	3.9	1.5
Cong. malformations (750-759)	3	1	2	4	2	2	12.5	15.5	10.4
Birth inj., atelect. (760-762)	3	0	3	3	0	3	3.3	-	4.6
#Other dis. early inf. (763-776)	7	0	7	7	0	7	7.7	-	10.7
Violent & acc. deaths (800-999)	22	11	11	26	16	10	81.0	124.0	52.1
Acc. deaths, total (800-962)	16	8	8	19	12	7	59.2	93.0	36.4
Motor vehicle acc. (810-835)	4	4	0	6	6	0	18.7	46.5	-
Ill-defined (780-795)	21	2	19	26	4	22	81.0	31.0	114.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PEARL RIVER COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	674	494	180	582	398	184	26.9	24.1	36.1
Fetal deaths	31	20	11	25	14	11	43.0	35.2	59.8
Deaths from all causes	170	115	55	197	134	63	9.1	8.1	12.4
Deaths under 1 year	20	11	9	24	14	10	41.2	35.2	54.3
Deaths under 1 month	14	10	4	16	12	4	27.5	30.2	21.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	4.6	-	19.6
Tbc., respiratory (001-008)	0	0	0	1	0	1	4.6	-	19.6
Syphilis (020-029)	0	0	0	1	1	0	4.6	6.1	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	16	9	7	19	12	7	88.0	72.7	137.2
Diabetes (260)	1	1	0	1	1	0	4.6	6.1	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	11	4	17	13	4	78.7	78.8	78.4
Heart diseases (410-443)	63	50	13	69	55	14	319.4	333.3	274.5
Arteriosclerosis (450)	8	6	2	8	6	2	37.0	36.4	39.2
Influenza (480-483)	1	1	0	1	1	0	4.6	6.1	-
#Pneumonia (490-493)	8	2	6	5	0	5	23.1	-	98.0
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	2	0	2	2	0	2	9.2	-	39.2
Chronic nephritis, etc. (592-594)	6	5	1	7	6	1	32.4	36.4	19.6
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	2	0	3	2	1	13.9	12.1	19.6
Birth inj., atelect. (760-762)	10	7	3	9	6	3	15.5	15.1	16.3
#Other dis. early inf. (763-776)	1	0	1	2	1	1	3.4	2.5	5.4
Violent & acc. deaths (800-999)	18	11	7	20	12	8	92.6	72.7	156.9
Acc. deaths, total (800-962)	16	10	6	16	9	7	74.1	54.5	137.2
Motor vehicle acc. (810-835)	10	6	4	8	4	4	37.0	24.2	78.4
Ill-defined (780-795)	9	3	6	10	3	7	46.3	18.2	137.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PERRY COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	97	69	28	223	146	77	31.0	28.1	38.5
Fetal deaths	0	0	0	9	5	4	40.4	34.2	51.9
Deaths from all causes	37	25	12	67	49	18	9.3	9.4	9.0
Deaths under 1 year	2	1	1	9	5	4	40.4	34.2	51.9
Deaths under 1 month	0	0	0	5	2	3	22.4	13.7	39.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	13.9	19.2	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	13.9	19.2	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	4	3	1	6	5	1	83.3	96.2	50.0
Diabetes (260)	3	2	1	4	3	1	55.6	57.7	50.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	5	2	3	7	4	3	97.2	76.9	150.0
Heart diseases (410-443)	17	12	5	26	20	6	361.1	384.6	300.0
Arteriosclerosis (450)	0	0	0	1	0	1	13.9	-	50.0
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	1	0	1	1	0	13.9	19.2	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	13.9	19.2	-
Birth inj., atelect. (760-762)	0	0	0	3	2	1	13.4	13.7	13.0
#Other dis. early inf. (763-776)	0	0	0	3	1	2	13.4	6.8	26.0
Violent & acc. deaths (800-999)	2	1	1	6	4	2	83.3	76.9	100.0
Acc. deaths, total (800-962)	1	0	1	5	3	2	69.4	57.7	100.0
Motor vehicle acc. (810-835)	0	0	0	2	1	1	27.8	19.2	50.0
Ill-defined (780-795)	3	2	1	3	2	1	41.7	38.5	50.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PIKE COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,009	528	481	851	373	478	26.4	21.6	32.1
Fetal deaths	31	11	20	27	10	17	31.7	26.8	35.6
Deaths from all causes	387	214	173	395	224	171	12.3	12.9	11.5
Deaths under 1 year	44	10	34	43	12	31	50.5	32.2	64.8
Deaths under 1 month	26	4	22	25	2	23	29.4	5.4	48.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	0	3	6	3	3	18.6	17.3	20.1
Tbc., respiratory (001-008)	3	0	3	6	3	3	18.6	17.3	20.1
Syphilis (020-029)	0	0	0	2	2	0	6.2	11.6	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	2	0	2	1	0	1	3.1	-	6.7
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	45	35	10	49	36	13	152.2	208.1	87.2
Diabetes (260)	6	5	1	7	6	1	21.7	34.7	6.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	58	34	24	50	31	19	155.3	179.2	127.5
Heart diseases (410-443)	114	73	41	118	73	45	366.4	422.0	302.0
Arteriosclerosis (450)	6	5	1	5	4	1	15.5	23.1	6.7
Influenza (480-483)	2	0	2	2	0	2	6.2	-	13.4
#Pneumonia (490-493)	3	3	0	4	3	1	12.4	17.3	6.7
#Diarrhea & enteritis (571)	6	2	4	4	2	2	12.4	11.6	13.4
Acute nephritis (590, 591)	3	1	2	3	2	1	9.3	11.6	6.7
Chronic nephritis, etc. (592-594)	4	2	2	5	3	2	15.5	17.3	13.4
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	4	3	1	6	5	1	18.6	28.9	6.7
Birth inj., atelect. (760-762)	13	3	10	12	2	10	14.1	5.4	20.9
#Other dis. early inf. (763-776)	11	1	10	11	0	11	12.9	-	23.0
Violent & acc. deaths (800-999)	38	15	23	37	15	22	114.9	86.7	147.6
Acc. deaths, total (800-962)	28	12	16	29	13	16	90.1	75.1	107.4
Motor vehicle acc. (810-835)	11	5	6	9	7	2	28.0	40.5	13.4
Ill-defined (780-795)	30	7	23	29	5	24	90.1	28.9	161.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PONTOTOC COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence				Rate*	
	Total	White	Non-white	Number		Non-white	Total	White	Non-white
				Total	White				
Live births	267	183	84	317	224	93	17.4	15.2	26.6
Fetal deaths	10	8	2	16	12	4	50.5	53.6	43.0
Deaths from all causes	154	123	31	170	137	33	9.3	9.3	9.4
Deaths under 1 year	7	5	2	7	6	1	22.1	26.8	10.8
Deaths under 1 month	5	3	2	4	3	1	12.6	13.4	10.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	2	0	2	11.0	-	57.1
Tbc., respiratory (001-008)	2	0	2	2	0	2	11.0	-	57.1
Syphilis (020-029)	1	0	1	1	0	1	5.5	-	28.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	5.5	-	28.6
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	5.5	-	28.6
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	19	13	6	23	17	6	126.4	115.6	171.4
Diabetes (260)	5	5	0	5	5	0	27.5	34.0	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	13	2	16	14	2	87.9	95.2	57.1
Heart diseases (410-443)	63	52	11	66	55	11	362.6	374.1	314.3
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	8	7	1	10	8	2	54.9	54.4	57.1
#Diarrhea & enteritis (571)	2	2	0	2	2	0	11.0	13.6	-
Acute nephritis (590, 591)	1	1	0	1	1	0	5.5	6.8	-
Chronic nephritis, etc. (592-594)	1	0	1	2	1	1	11.0	6.8	28.6
Maternal deaths (640-689)	1	1	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	2	2	0	2	2	0	6.3	8.9	-
#Other dis. early inf. (763-776)	2	1	1	1	1	0	3.2	4.5	-
Violent & acc. deaths (800-999)	14	13	1	16	14	2	87.9	95.2	57.1
Acc. deaths, total (800-962)	10	10	0	12	11	1	65.9	74.8	28.6
Motor vehicle acc. (810-835)	4	4	0	7	6	1	38.5	40.8	28.6
Ill-defined (780-795)	3	1	2	3	1	2	16.5	6.8	57.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PRENTISS COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	509	428	81	444	372	72	24.5	23.4	32.7
Fetal deaths	11	8	3	11	7	4	24.8	18.8	55.6
Deaths from all causes	150	131	19	154	136	18	8.5	8.5	8.2
Deaths under 1 year	9	5	4	7	5	2	15.8	13.4	27.8
Deaths under 1 month	8	5	3	7	5	2	15.8	13.4	27.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	2	0	4	4	0	22.1	25.2	-
Tbc., respiratory (001-008)	2	2	0	4	4	0	22.1	25.2	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	16	1	20	19	1	110.5	119.5	45.4
Diabetes (260)	1	1	0	2	2	0	11.0	12.6	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	26	24	2	24	22	2	132.6	138.4	90.9
Heart diseases (410-443)	37	29	8	44	36	8	243.1	226.4	363.6
Arteriosclerosis (450)	1	1	0	1	1	0	5.5	6.3	-
Influenza (480-483)	7	7	0	7	7	0	38.7	44.0	-
#Pneumonia (490-493)	6	6	0	5	5	0	27.6	31.4	-
#Diarrhea & enteritis (571)	0	0	0	1	1	0	5.5	6.3	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	2	0	2	2	0	11.0	12.6	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	6	4	2	4	3	1	9.0	8.1	13.9
#Other dis. early inf. (763-776)	2	1	1	3	2	1	6.8	5.4	13.9
Violent & acc. deaths (800-999)	12	11	1	11	10	1	60.8	62.9	45.4
acc. deaths, total (800-962)	10	10	0	9	9	0	49.7	56.6	-
Motor vehicle acc. (810-835)	8	8	0	7	7	0	38.7	44.0	-
Ill-defined (780-795)	8	7	1	8	7	1	44.2	44.0	45.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
QUITMAN COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	823	226	597	787	215	572	30.9	22.9	35.5
Fetal deaths	17	5	12	18	3	15	22.9	14.0	26.2
Deaths from all causes	160	31	129	198	54	144	7.8	5.7	8.9
Deaths under 1 year	29	4	25	33	5	28	41.9	23.2	49.0
Deaths under 1 month	15	4	11	17	4	13	21.6	18.6	22.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	3	0	3	11.8	-	18.6
Tbc., respiratory (001-008)	2	0	2	3	0	3	11.8	-	18.6
Syphilis (020-029)	2	0	2	1	0	1	3.9	-	6.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	2	0	2	7.8	-	12.4
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	16	3	13	19	7	12	74.5	74.5	74.5
Diabetes (260)	1	0	1	3	1	2	11.8	10.6	12.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	29	4	25	34	6	28	133.3	63.8	173.9
Heart diseases (410-443)	32	9	23	41	14	27	160.8	148.9	167.7
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	1	0	1	1	0	1	3.9	-	6.2
#Pneumonia (490-493)	7	2	5	8	3	5	31.4	31.9	31.0
#Diarrhea & enteritis (571)	4	0	4	5	1	4	19.6	10.6	24.8
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	6	0	6	7	1	6	27.4	10.6	37.3
Maternal deaths (640-689)	1	1	0	2	1	1	2.5	4.6	1.7
Cong. malformations (750-759)	3	2	1	3	2	1	11.8	21.3	6.2
Birth inj., atelect. (760-762)	1	0	1	2	0	2	2.5	-	3.5
#Other dis. early inf. (763-776)	12	2	10	13	2	11	16.5	9.3	19.2
Violent & acc. deaths (800-999)	13	3	10	19	7	12	74.5	74.5	74.5
Acc. deaths, total (800-962)	9	2	7	14	6	8	54.9	63.8	49.7
Motor vehicle acc. (810-835)	3	1	2	6	4	2	23.5	42.6	12.4
Ill-defined (780-795)	22	1	21	22	1	21	86.3	10.6	130.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
RANKIN COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	230	31	199	610	273	337	22.3	17.8	27.8
Fetal deaths	10	5	5	26	13	13	42.6	47.6	38.6
Deaths from all causes	402	185	217	217	105	112	7.9	6.9	9.2
Deaths under 1 year	9	0	9	15	4	11	24.6	14.6	32.6
Deaths under 1 month	0	0	0	6	4	2	9.8	14.6	5.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	12	4	8	1	1	0	3.6	6.5	-
Tbc., respiratory (001-008)	9	3	6	1	1	0	3.6	6.5	-
Syphilis (020-029)	3	0	3	1	0	1	3.6	-	8.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	13	1	12	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	1	0	3.6	6.5	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	32	19	13	31	14	17	113.1	91.5	140.5
Diabetes (260)	3	3	0	1	1	0	3.6	6.5	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	77	39	38	29	13	16	105.8	85.0	132.2
Heart diseases (410-443)	130	68	62	69	44	25	251.8	287.6	206.6
Arteriosclerosis (450)	9	3	6	7	2	5	25.5	13.1	41.3
Influenza (480-483)	2	0	2	2	0	2	7.3	-	16.5
#Pneumonia (490-493)	18	7	11	9	3	6	32.8	19.6	49.6
#Diarrhea & enteritis (571)	1	1	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	1	1	0	3.6	6.5	-
Chronic nephritis, etc. (592-594)	3	0	3	5	1	4	18.2	6.5	33.0
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	3.6	6.5	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	2	0	2	7	3	4	11.5	11.0	11.9
Violent & acc. deaths (800-999)	20	7	13	17	6	11	62.0	39.2	90.9
Acc. deaths, total (800-962)	15	4	11	12	4	8	43.8	26.1	66.1
Motor vehicle acc. (810-835)	5	3	2	4	2	2	14.6	13.1	16.5
Ill-defined (780-795)	12	2	10	14	2	12	51.1	13.1	99.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SCOTT COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence				Rate*		
	Total	White	Non-white	Number			Total	White	Non-white	
				Total	White	Non-white				
Live births	528	264	264	556	251	305	27.9	23.0	33.9	
Fetal deaths	10	3	7	14	4	10	25.2	15.9	32.8	
Deaths from all causes	187	119	68	202	125	77	10.2	11.5	8.6	
Deaths under 1 year	16	8	8	20	10	10	36.0	39.8	32.8	
Deaths under 1 month	11	6	5	15	8	7	27.0	31.9	23.0	
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.										
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-	
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-	
Syphilis (020-029)	0	0	0	0	0	0	-	-	-	
Typhoid fever (040)	0	0	0	0	0	0	-	-	-	
Dysentery (045-048)	0	0	0	0	0	0	-	-	-	
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-	
Diphtheria (055)	0	0	0	0	0	0	-	-	-	
Whooping cough (056)	0	0	0	0	0	0	-	-	-	
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-	
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-	
Smallpox (084)	0	0	0	0	0	0	-	-	-	
Measles (085)	0	0	0	0	0	0	-	-	-	
Malaria (110-117)	0	0	0	0	0	0	-	-	-	
Cancer, incl. leukemia (140-205)	21	16	5	26	19	7	130.6	174.3	77.8	
Diabetes (260)	3	0	3	5	2	3	25.1	18.3	33.3	
Pellagra (281)	0	0	0	0	0	0	-	-	-	
Intracranial lesions (330-334)	34	19	15	33	18	15	165.8	165.1	166.7	
Heart diseases (410-443)	68	51	17	71	49	22	356.8	449.5	244.4	
Arteriosclerosis (450)	3	2	1	3	2	1	15.1	18.3	11.1	
Influenza (480-483)	2	1	1	2	1	1	10.0	9.2	11.1	
#Pneumonia (490-493)	3	2	1	2	1	1	10.0	9.2	11.1	
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-	
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-	
Chronic nephritis, etc. (592-594)	3	1	2	2	0	2	10.0	-	22.2	
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-	
Cong. malformations (750-759)	1	1	0	2	2	0	10.0	18.3	-	
Birth inj., atelect. (760-762)	1	1	0	1	1	0	1.8	4.0	-	
#Other dis. early inf. (763-776)	5	3	2	8	4	4	14.4	15.9	13.1	
Violent & acc. deaths (800-999)	14	6	8	17	9	8	85.4	82.6	88.9	
Acc. deaths, total (800-962)	14	6	8	17	9	8	85.4	82.6	88.9	
Motor vehicle acc. (810-835)	6	3	3	9	5	4	45.2	45.9	44.4	
Ill-defined (780-795)	20	7	13	19	6	13	95.5	55.0	144.4	

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SHARKEY COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	284	57	227	403	71	332	32.2	20.9	36.5
Fetal deaths	15	5	10	19	5	14	47.1	70.4	42.2
Deaths from all causes	101	23	78	109	21	88	8.7	6.2	9.7
Deaths under 1 year	15	4	11	10	1	9	24.8	14.1	27.1
Deaths under 1 month	8	3	5	3	0	3	7.4	-	9.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	8.0	-	11.0
Tbc., respiratory (001-008)	0	0	0	1	0	1	8.0	-	11.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	8.0	-	11.0
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	2	7	10	1	9	80.0	29.4	98.9
Diabetes (260)	1	0	1	1	0	1	8.0	-	11.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	13	1	12	16	1	15	128.0	29.4	164.8
Heart diseases (410-443)	35	10	25	41	12	29	328.0	352.9	318.7
Arteriosclerosis (450)	0	0	0	1	0	1	8.0	-	11.0
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	0	2	4	1	3	32.0	29.4	33.0
#Diarrhea & enteritis (571)	1	0	1	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	0	2	1	0	1	8.0	-	11.0
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	9	3	6	5	0	5	12.4	-	15.1
Violent & acc. deaths (800-999)	9	4	5	8	3	5	64.0	88.2	54.9
Acc. deaths, total (800-962)	9	4	5	7	2	5	56.0	58.8	54.9
Motor vehicle acc. (810-835)	4	2	2	2	0	2	16.0	-	22.0
Ill-defined (780-795)	12	1	11	10	1	9	80.0	29.4	98.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SIMPSON COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	531	236	295	500	248	252	23.7	19.1	31.1
Fetal deaths	12	2	10	14	4	10	28.0	16.1	39.7
Deaths from all causes	181	109	72	180	124	56	8.5	9.5	6.9
Deaths under 1 year	7	3	4	17	10	7	34.0	40.3	27.8
Deaths under 1 month	5	2	3	14	8	6	28.0	32.2	23.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	40	21	19	2	1	1	9.5	7.7	12.3
Tbc., respiratory (001-008)	40	21	19	2	1	1	9.5	7.7	12.3
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	19	16	3	20	16	4	94.8	123.1	49.4
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	1	1	0	1	1	0	4.7	7.7	-
Intracranial lesions (330-334)	20	10	10	25	16	9	118.5	123.1	111.1
Heart diseases (410-443)	51	40	11	64	52	12	303.3	400.0	148.1
Arteriosclerosis (450)	1	1	0	1	1	0	4.7	7.7	-
Influenza (480-483)	2	1	1	2	1	1	9.5	7.7	12.3
#Pneumonia (490-493)	12	6	6	16	9	7	75.8	69.2	86.4
#Diarrhea & enteritis (571)	1	0	1	2	1	1	9.5	7.7	12.3
Acute nephritis (590, 591)	0	0	0	2	2	0	9.5	15.4	-
Chronic nephritis, etc. (592-594)	3	0	3	2	0	2	9.5	-	24.7
Maternal deaths (640-689)	0	0	0	1	0	1	2.0	-	4.0
Cong. malformations (750-759)	2	1	1	6	5	1	28.4	38.5	12.3
Birth inj., atelect. (760-762)	2	1	1	3	3	0	6.0	12.1	-
#Other dis. early inf. (763-776)	3	1	2	8	3	5	16.0	12.1	19.8
Violent & acc. deaths (800-999)	8	6	2	8	7	1	37.9	53.8	12.3
Acc. deaths, total (800-962)	6	4	2	6	5	1	28.4	38.5	12.3
Motor vehicle acc. (810-835)	4	3	1	4	4	0	19.0	30.8	-
Ill-defined (780-795)	6	2	4	7	2	5	33.2	15.4	61.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SMITH COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	186	128	58	364	208	156	26.6	18.7	60.0
Fetal deaths	3	1	2	7	5	2	19.2	24.0	12.8
Deaths from all causes	71	55	16	94	75	19	6.9	6.8	7.3
Deaths under 1 year	1	1	0	9	8	1	24.7	38.5	6.4
Deaths under 1 month	1	1	0	7	7	0	19.2	33.6	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	8	1	10	8	2	73.0	72.1	76.9
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	12	9	3	15	12	3	109.5	108.1	115.4
Heart diseases (410-443)	29	25	4	35	30	5	255.5	270.3	192.3
Arteriosclerosis (450)	1	1	0	1	1	0	7.3	9.0	-
Influenza (480-483)	2	1	1	2	1	1	14.6	9.0	38.5
#Pneumonia (490-493)	0	0	0	1	0	1	7.3	-	38.5
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	3	3	0	21.9	27.0	-
Maternal deaths (640-689)	0	0	0	1	0	1	2.7	-	6.4
Cong. malformations (750-759)	1	1	0	6	6	0	43.8	54.0	-
Birth inj., atelect. (760-762)	1	1	0	2	2	0	5.5	9.6	-
#Other dis. early inf. (763-776)	0	0	0	1	1	0	2.7	4.8	-
Violent & acc. deaths (800-999)	8	3	5	7	5	2	51.1	45.0	76.9
Acc. deaths, total (800-962)	7	3	4	7	5	2	51.1	45.0	76.9
Motor vehicle acc. (810-835)	4	2	2	4	3	1	29.2	27.0	38.5
Ill-defined (780-795)	4	3	1	3	2	1	21.9	18.0	38.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
STONE COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	145	99	46	138	52	86	21.6	10.6	57.3
Fetal deaths	2	1	1	2	1	1	14.5	19.2	11.6
Deaths from all causes	50	32	18	46	29	17	7.2	5.9	11.3
Deaths under 1 year	2	1	1	3	1	2	21.7	19.2	23.2
Deaths under 1 month	1	1	0	2	1	1	14.5	19.2	11.6
SELECTED CAUSES OF DEATH									
AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	4	3	1	6	5	1	93.8	102.0	66.7
Diabetes (260)	1	0	1	1	0	1	15.6	-	66.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	5	2	3	4	2	2	62.5	40.8	133.3
Heart diseases (410-443)	23	16	7	21	13	8	328.1	265.3	533.3
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	1	1	1	0	1	15.6	-	66.7
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	1	1	0	15.6	20.4	-
Birth inj., atelect. (760-762)	0	0	0	1	0	1	7.2	-	11.6
#Other dis. early inf. (763-776)	0	0	0	0	0	0	-	-	-
Violent & acc. deaths (800-999)	7	5	2	3	2	1	46.9	40.8	66.7
Acc. deaths, total (800-962)	7	5	2	3	2	1	46.9	40.8	66.7
Motor vehicle acc. (810-835)	6	4	2	2	1	1	31.2	20.4	66.7
Ill-defined (780-795)	2	1	1	4	3	1	62.5	61.2	66.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SUNFLOWER COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,417	266	1,151	1,523	319	1,204	28.2	19.6	31.8
Fetal deaths	56	14	42	61	16	45	40.0	50.2	37.4
Deaths from all causes	366	91	275	431	124	307	8.0	7.6	8.1
Deaths under 1 year	57	5	52	62	8	54	40.7	25.1	44.8
Deaths under 1 month	29	4	25	32	6	26	21.0	18.8	21.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	4	1	3	7.4	6.1	7.9
Tbc., respiratory (001-008)	2	1	1	3	1	2	5.5	6.1	5.3
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	1	0	1.8	6.1	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	1.8	-	2.6
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	39	18	21	53	27	26	98.0	165.6	68.8
Diabetes (260)	8	2	6	8	1	7	14.8	6.1	18.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	50	11	39	60	15	45	110.9	92.0	119.0
Heart diseases (410-443)	94	31	63	114	40	74	210.7	245.4	195.8
Arteriosclerosis (450)	1	1	0	3	2	1	5.5	12.3	2.6
Influenza (480-483)	4	0	4	4	0	4	7.4	-	10.6
#Pneumonia (490-493)	10	2	8	12	4	8	22.2	24.5	21.2
#Diarrhea & enteritis (571)	4	0	4	3	0	3	5.5	-	7.9
Acute nephritis (590, 591)	1	0	1	2	1	1	3.7	6.1	2.6
Chronic nephritis, etc. (592-594)	7	5	2	8	5	3	14.8	30.7	7.9
Maternal deaths (640-689)	3	0	3	2	0	2	1.3	-	1.7
Cong. malformations (750-759)	5	0	5	6	0	6	11.1	-	15.9
Birth inj., atelect. (760-762)	6	3	3	8	4	4	5.2	12.5	3.3
#Other dis. early inf. (763-776)	19	1	18	19	2	17	12.5	6.3	14.1
Violent & acc. deaths (800-999)	30	9	21	31	9	22	57.3	55.2	58.2
Acc. deaths, total (800-962)	25	6	19	26	6	20	48.0	36.8	52.9
Motor vehicle acc. (810-835)	15	4	11	17	5	12	31.4	30.7	31.7
Ill-defined (780-795)	54	2	52	52	2	50	96.1	12.3	132.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TALLAHATCHIE COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	812	140	672	849	175	674	29.9	17.5	36.6
Fetal deaths	32	7	25	32	7	25	37.7	40.0	37.1
Deaths from all causes	260	73	187	288	90	198	10.1	9.0	10.8
Deaths under 1 year	46	6	40	47	7	40	55.4	40.0	59.3
Deaths under 1 month	28	3	25	28	3	25	33.0	17.1	37.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	1	0	1	3.5	-	5.4
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	3	0	3	2	0	2	7.0	-	10.9
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	27	12	15	34	15	19	119.7	150.0	103.3
Diabetes (260)	5	2	3	6	2	4	21.1	20.0	21.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	37	12	25	41	15	26	144.4	150.0	141.3
Heart diseases (410-443)	68	24	44	74	29	45	260.6	290.0	244.6
Arteriosclerosis (450)	3	2	1	3	2	1	10.6	20.0	5.4
Influenza (480-483)	5	1	4	5	1	4	17.6	10.0	21.7
#Pneumonia (490-493)	8	0	8	9	0	9	31.7	-	48.9
#Diarrhea & enteritis (571)	1	1	0	1	1	0	3.5	10.0	-
Acute nephritis (590, 591)	1	0	1	1	0	1	3.5	-	5.4
Chronic nephritis, etc. (592-594)	4	1	3	5	1	4	17.6	10.0	21.7
Maternal deaths (640-689)	3	1	2	3	1	2	3.5	5.7	3.0
Cong. malformations (750-759)	4	2	2	4	2	2	14.1	20.0	10.9
Birth inj., atelect. (760-762)	6	1	5	6	1	5	7.1	5.7	7.4
#Other dis. early inf. (763-776)	15	1	14	14	1	13	16.5	5.7	19.3
Violent & acc. deaths (800-999)	18	7	11	23	10	13	81.0	100.0	70.6
Acc. deaths, total (800-962)	15	6	9	20	9	11	70.4	90.0	59.8
Motor vehicle acc. (810-835)	6	4	2	8	5	3	28.2	50.0	16.3
Ill-defined (780-795)	35	1	34	35	1	34	123.2	10.0	184.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TATE COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	393	8	385	552	130	422	32.1	19.1	40.6
Fetal deaths	21	0	21	36	1	35	65.2	7.7	82.9
Deaths from all causes	123	50	73	175	78	97	10.2	11.5	9.3
Deaths under 1 year	9	0	9	15	4	11	27.2	30.8	26.1
Deaths under 1 month	7	0	7	13	4	9	23.6	30.8	21.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	1	0	1	2	0	2	11.6	-	19.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	16	9	7	22	13	9	127.9	191.2	86.5
Diabetes (260)	1	0	1	2	1	1	11.6	14.7	9.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	11	5	6	16	9	7	93.0	132.4	67.3
Heart diseases (410-443)	56	26	30	58	35	33	395.3	514.7	317.3
Arteriosclerosis (450)	2	1	1	2	1	1	11.6	14.7	9.6
Influenza (480-483)	1	0	1	1	0	1	5.8	-	9.6
#Pneumonia (490-493)	0	0	0	1	1	0	5.8	14.7	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	5	2	3	6	3	3	34.9	44.1	28.8
Maternal deaths (640-689)	0	0	0	3	0	3	5.4	-	7.1
Cong. malformations (750-759)	1	0	1	1	0	1	5.8	-	9.6
Birth inj., atelect. (760-762)	0	0	0	1	0	1	1.8	-	2.4
#Other dis. early inf. (763-776)	4	0	4	9	4	5	16.3	30.8	11.8
Violent & acc. deaths (800-999)	7	1	6	11	2	9	64.0	29.4	86.5
Acc. deaths, total (800-962)	7	1	6	10	2	8	58.1	29.4	76.9
Motor vehicle acc. (810-835)	2	1	1	2	1	1	11.6	14.7	9.6
Ill-defined (780-795)	12	3	9	13	3	10	75.6	44.1	96.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TIPPAH COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	333	233	100	301	202	99	18.1	15.6	26.8
Fetal deaths	20	16	4	14	12	2	46.5	59.4	20.2
Deaths from all causes	120	97	23	128	104	24	7.7	8.1	6.5
Deaths under 1 year	8	6	2	7	5	2	23.2	24.8	20.2
Deaths under 1 month	5	4	1	4	3	1	13.3	14.8	10.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	2	1	3	2	1	18.1	15.5	27.0
Tbc., respiratory (001-008)	3	2	1	3	2	1	18.1	15.5	27.0
Syphilis (020-029)	0	0	0	1	0	1	6.0	-	27.0
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	16	14	2	17	15	2	102.4	116.3	54.0
Diabetes (260)	1	1	0	1	1	0	6.0	7.8	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	11	10	1	14	13	1	84.3	100.8	27.0
Heart diseases (410-443)	58	46	12	56	44	12	337.3	341.1	324.3
Arteriosclerosis (450)	1	1	0	2	2	0	12.0	15.5	-
Influenza (480-483)	2	2	0	1	1	0	6.0	7.8	-
#Pneumonia (490-493)	1	1	0	2	2	0	12.0	15.5	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	0	3	3	0	3	18.1	-	81.1
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	1	0	1	6.0	-	27.0
Birth inj., atelect. (760-762)	5	4	1	3	2	1	10.0	9.9	10.1
#Other dis. early inf. (763-776)	1	1	0	1	1	0	3.3	5.0	-
Violent & acc. deaths (800-999)	7	6	1	7	6	1	42.2	46.5	27.0
Acc. deaths, total (800-962)	6	5	1	6	5	1	36.1	38.8	27.0
Motor vehicle acc. (810-835)	2	1	1	1	0	1	6.0	-	27.0
Ill-defined (780-795)	0	0	0	2	2	0	12.0	15.5	-

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TISHOMINGO COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	138	131	7	264	243	21	20.3	19.8	30.0
Fetal deaths	2	2	0	2	2	0	7.6	8.2	-
Deaths from all causes	110	102	8	143	133	10	11.0	10.8	14.3
Deaths under 1 year	9	5	4	16	11	5	60.6	45.3	238.1
Deaths under 1 month	6	4	2	13	10	3	49.2	41.2	142.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	7.7	8.1	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	7.7	8.1	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	9	1	15	14	1	115.4	113.8	142.8
Diabetes (260)	2	2	0	3	3	0	23.1	24.4	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	34	33	1	39	37	2	300.0	300.8	285.7
Heart diseases (410-443)	23	23	0	29	28	1	223.1	227.6	142.8
Arteriosclerosis (450)	4	3	1	5	4	1	38.5	32.5	142.8
Influenza (480-483)	1	1	0	1	1	0	7.7	8.1	-
#Pneumonia (490-493)	9	8	1	10	9	1	76.9	73.2	142.8
#Diarrhea & enteritis (571)	1	1	0	1	1	0	7.7	8.1	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	3	0	3	3	0	23.1	24.4	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	1	1	2	1	1	15.4	8.1	142.8
Birth inj., atelect. (760-762)	2	2	0	6	5	1	22.7	20.6	47.6
#Other dis. early inf. (763-776)	3	1	2	5	3	2	18.9	12.3	95.2
Violent & acc. deaths (800-999)	7	7	0	8	8	0	61.5	65.0	-
Acc. deaths, total (800-962)	7	7	0	8	8	0	61.5	65.0	-
Motor vehicle acc. (810-835)	5	5	0	6	6	0	46.2	48.8	-
Ill-defined (780-795)	2	1	1	2	2	0	15.4	16.3	-

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TUNICA COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	698	84	614	714	84	630	31.4	19.1	34.4
Fetal deaths	20	2	18	17	0	17	23.8	-	27.0
Deaths from all causes	174	25	149	193	30	163	8.5	6.8	8.9
Deaths under 1 year	28	1	27	33	2	31	46.2	23.8	49.2
Deaths under 1 month	14	0	14	17	1	16	23.8	11.9	25.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	4.4	-	5.5
Tbc., respiratory (001-008)	0	0	0	1	0	1	4.4	-	5.5
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	4.4	-	5.5
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	4.4	-	5.5
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	4	6	16	7	9	70.5	159.1	49.2
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	8	1	7	9	1	8	39.6	22.7	43.7
Heart diseases (410-443)	57	9	48	62	9	53	273.1	204.5	289.6
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	5	0	5	5	0	5	22.0	-	27.3
#Pneumonia (490-493)	11	1	10	11	1	10	48.4	22.7	54.6
#Diarrhea & enteritis (571)	1	0	1	1	0	1	4.4	-	5.5
Acute nephritis (590, 591)	1	0	1	1	0	1	4.4	-	5.5
Chronic nephritis, etc. (592-594)	9	0	9	8	0	8	35.2	-	43.7
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	1	1	3	2	1	13.2	45.4	5.5
Birth inj., atelect. (760-762)	2	0	2	2	0	2	2.8	-	3.2
#Other dis. early inf. (763-776)	7	0	7	10	1	9	14.0	11.9	14.3
Violent & acc. deaths (800-999)	17	3	14	16	3	13	70.5	68.2	71.0
Acc. deaths, total (800-962)	13	3	10	13	3	10	57.3	68.2	54.6
Motor vehicle acc. (810-835)	7	2	5	6	2	4	26.4	45.4	21.8
Ill-defined (780-795)	30	2	28	29	1	28	127.8	22.7	153.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
UNION COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	473	369	104	398	302	96	21.2	20.4	24.0
Fetal deaths	30	27	3	22	20	2	55.3	66.2	20.8
Deaths from all causes	205	152	53	211	156	55	11.2	10.5	13.8
Deaths under 1 year	10	7	3	11	7	4	27.6	23.2	41.7
Deaths under 1 month	6	4	2	5	3	2	12.6	9.9	20.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	1	0	1	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	1	1	0	1	1	0	5.3	6.8	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	22	15	7	25	17	8	133.0	114.9	200.0
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	43	31	12	46	33	13	244.7	223.0	325.0
Heart diseases (410-443)	77	60	17	80	61	19	425.5	412.2	475.0
Arteriosclerosis (450)	4	0	4	3	0	3	16.0	-	75.0
Influenza (480-483)	2	1	1	2	1	1	10.6	6.8	25.0
#Pneumonia (490-493)	1	0	1	3	2	1	16.0	13.5	25.0
#Diarrhea & enteritis (571)	2	2	0	2	2	0	10.6	13.5	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	3	2	1	16.0	13.5	25.0
Maternal deaths (640-689)	1	0	1	1	0	1	2.5	-	10.4
Cong. malformations (750-759)	2	2	0	2	2	0	10.6	13.5	-
Birth inj., atelect. (760-762)	4	3	1	3	2	1	7.5	6.6	10.4
#Other dis. early inf. (763-776)	2	1	1	2	1	1	5.0	3.3	10.4
Violent & acc. deaths (800-999)	13	11	2	12	10	2	63.8	67.6	50.0
Acc. deaths, total (800-962)	9	7	2	8	6	2	42.6	40.5	50.0
Motor vehicle acc. (810-835)	7	6	1	5	4	1	26.6	27.0	25.0
Ill-defined (780-795)	6	4	2	6	4	2	31.9	27.0	50.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WALTHALL COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence				Rate*		
	Total	White	Non-white	Number			Total	White	Non-white	
				Total	White	Non-white				
Live births	628	366	262	345	151	194	24.1	19.1	30.3	
Fetal deaths	28	14	14	17	8	9	49.3	53.0	46.4	
Deaths from all causes	135	76	59	137	72	65	9.6	9.1	10.2	
Deaths under 1 year	16	6	10	12	2	10	34.8	13.2	51.5	
Deaths under 1 month	6	3	3	5	2	3	14.5	13.2	15.5	
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.										
Tbc., all forms (001-019)	1	0	1	2	0	2	14.0	-	31.2	
Tbc., respiratory (001-008)	1	0	1	2	0	2	14.0	-	31.2	
Syphilis (020-029)	0	0	0	0	0	0	-	-	-	
Typhoid fever (040)	0	0	0	0	0	0	-	-	-	
Dysentery (045-048)	0	0	0	0	0	0	-	-	-	
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-	
Diphtheria (055)	0	0	0	0	0	0	-	-	-	
Whooping cough (056)	0	0	0	0	0	0	-	-	-	
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-	
Acute poliomyelitis (080)	0	0	0	1	0	1	7.0	-	15.6	
Smallpox (084)	0	0	0	0	0	0	-	-	-	
Measles (085)	0	0	0	0	0	0	-	-	-	
Malaria (110-117)	0	0	0	0	0	0	-	-	-	
Cancer, incl. leukemia (140-205)	18	11	7	18	11	7	125.9	139.2	109.4	
Diabetes (260)	4	2	2	4	2	2	28.0	25.3	31.2	
Pellagra (281)	0	0	0	0	0	0	-	-	-	
Intracranial lesions (330-334)	18	10	8	19	10	9	132.9	126.6	140.6	
Heart diseases (410-443)	48	29	19	46	27	19	321.7	341.8	296.9	
Arteriosclerosis (450)	2	0	2	2	0	2	14.0	-	31.2	
Influenza (480-483)	0	0	0	0	0	0	-	-	-	
#Pneumonia (490-493)	6	2	4	6	2	4	42.0	25.3	62.5	
#Diarrhea & enteritis (571)	1	1	0	0	0	0	-	-	-	
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-	
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-	
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-	
Cong. malformations (750-759)	1	1	0	1	0	1	7.0	-	15.6	
Birth inj., atelect. (760-762)	1	0	1	1	0	1	2.9	-	5.2	
#Other dis. early inf. (763-776)	3	2	1	2	1	1	5.8	6.6	5.2	
Violent & acc. deaths (800-999)	4	3	1	14	7	7	97.9	88.6	109.4	
Acc. deaths, total (800-962)	2	1	1	11	5	6	76.9	63.3	93.8	
Motor vehicle acc. (810-835)	1	0	1	6	2	4	42.0	25.3	62.5	
Ill-defined (780-795)	8	1	7	7	1	6	49.0	12.6	93.8	

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WARREN COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,488	664	824	1,192	575	617	30.2	25.2	36.9
Fetal deaths	79	24	55	51	21	30	42.8	36.5	48.6
Deaths from all causes	680	250	430	520	213	307	13.2	9.3	18.4
Deaths under 1 year	84	21	63	57	18	39	47.8	31.3	63.2
Deaths under 1 month	63	18	45	43	16	27	36.1	27.8	43.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	6	0	6	8	3	5	20.2	13.2	29.9
Tbc., respiratory (001-003)	6	0	6	7	2	5	17.7	8.8	29.9
Syphilis (020-029)	2	1	1	1	0	1	2.5	-	6.0
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	1	1	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	2.5	-	6.0
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	92	35	57	62	24	38	157.0	105.3	227.5
Diabetes (260)	4	2	2	3	2	1	7.6	8.8	6.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	85	30	55	78	31	47	197.5	136.0	281.4
Heart diseases (410-443)	186	89	97	154	76	78	389.9	333.3	467.1
Arteriosclerosis (450)	10	6	4	9	6	3	22.8	26.3	18.0
Influenza (480-483)	5	0	5	5	0	5	12.6	-	29.9
#Pneumonia (490-493)	14	4	10	7	2	5	17.7	8.8	29.9
#Diarrhea & enteritis (571)	5	0	5	4	0	4	10.1	-	24.0
Acute nephritis (590, 591)	1	0	1	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	17	4	13	9	4	5	22.8	17.5	29.9
Maternal deaths (640-689)	2	0	2	1	0	1	0.8	-	1.6
Cong. malformations (750-759)	12	4	8	11	4	7	27.8	17.5	41.9
Birth inj., atelect. (760-762)	10	3	7	6	3	3	5.0	5.2	4.9
#Other dis. early inf. (763-776)	41	13	28	28	11	17	23.5	19.1	27.6
Violent & acc. deaths (800-999)	44	16	28	24	11	13	60.8	48.2	77.8
Acc. deaths, total (800-962)	36	13	23	19	9	10	48.1	39.5	59.9
Motor vehicle acc. (810-835)	6	2	4	2	1	1	5.1	4.4	6.0
Ill-defined (780-795)	59	11	48	57	10	47	144.3	43.8	281.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WASHINGTON COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	2,862	1,133	1,729	2,709	1,046	1,663	35.4	33.6	36.6
Fetal deaths	234	133	101	230	124	106	84.9	118.5	63.7
Deaths from all causes	789	239	550	824	222	602	10.8	7.1	13.2
Deaths under 1 year	113	32	81	114	31	83	42.1	29.6	49.9
Deaths under 1 month	62	26	36	62	24	38	22.9	22.9	22.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	6	0	6	7.8	-	13.2
Tbc., respiratory (001-008)	0	0	0	4	0	4	5.2	-	8.8
Syphilis (020-029)	0	0	0	1	1	0	1.3	3.2	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	3	0	3	5	1	4	6.5	3.2	8.8
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	1.3	-	2.2
Meningococcal infections (057)	1	1	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	92	38	54	100	35	65	130.7	112.5	143.2
Diabetes (260)	4	1	3	5	1	4	6.5	3.2	8.8
Pellagra (281)	1	0	1	1	0	1	1.3	-	2.2
Intracranial lesions (330-334)	74	17	57	78	15	63	102.0	48.2	138.8
Heart diseases (410-443)	247	83	164	239	72	167	312.4	231.5	367.8
Arteriosclerosis (450)	12	3	9	11	2	9	14.4	6.4	19.8
Influenza (480-483)	6	0	6	6	0	6	7.8	-	13.2
#Pneumonia (490-493)	28	8	20	25	5	20	32.7	16.1	44.0
#Diarrhea & enteritis (571)	6	0	6	7	0	7	9.2	-	15.4
Acute nephritis (590, 591)	2	0	2	2	0	2	2.6	-	4.4
Chronic nephritis, etc. (592-594)	9	1	8	11	1	10	14.4	3.2	22.0
Maternal deaths (640-689)	5	0	5	5	0	5	1.8	-	3.0
Cong. malformations (750-759)	7	5	2	11	8	3	14.4	25.7	6.6
Birth inj., atelect. (760-762)	12	8	4	11	8	3	4.1	7.6	1.8
#Other dis. early inf. (763-776)	46	13	33	44	10	34	16.2	9.6	20.4
Violent & acc. deaths (800-999)	62	26	36	67	29	38	87.6	93.2	83.7
Acc. deaths, total (800-962)	49	25	24	54	28	26	70.6	90.0	57.3
Motor vehicle acc. (810-835)	19	11	8	20	10	10	26.1	32.2	22.0
Ill-defined (780-795)	118	9	109	120	9	111	156.9	28.9	244.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WAYNE COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	457	279	178	453	244	209	29.2	25.2	36.0
Fetal deaths	13	5	8	23	5	18	50.8	20.5	86.1
Deaths from all causes	123	81	42	160	106	54	10.3	10.9	9.3
Deaths under 1 year	7	2	5	14	6	8	30.9	24.6	38.3
Deaths under 1 month	3	2	1	9	5	4	19.9	20.5	19.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	6.4	10.3	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	6.4	10.3	-
Syphilis (020-029)	1	0	1	1	0	1	6.4	-	17.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	14	9	5	19	13	6	122.6	134.0	103.4
Diabetes (260)	4	3	1	5	4	1	32.2	41.2	17.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	19	14	5	23	17	6	148.4	175.2	103.4
Heart diseases (410-443)	37	29	8	40	31	9	258.1	319.6	155.2
Arteriosclerosis (450)	1	1	0	2	2	0	12.9	20.6	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	0	2	3	1	2	19.4	10.3	34.5
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	4	0	5	4	1	32.2	41.2	17.2
Maternal deaths (640-689)	0	0	0	1	0	1	2.2	-	4.8
Cong. malformations (750-759)	0	0	0	3	3	0	19.4	30.9	-
Birth inj., atelect. (760-762)	2	1	1	2	1	1	4.4	4.1	4.8
#Other dis. early inf. (763-776)	0	0	0	4	1	3	8.8	4.1	14.4
Violent & acc. deaths (800-999)	13	10	3	15	12	3	96.8	123.7	51.7
acc. deaths, total (800-962)	12	9	3	13	10	3	83.9	103.1	51.7
Motor vehicle acc. (810-835)	9	8	1	9	8	1	58.1	82.5	17.2
Ill-defined (780-795)	12	3	9	16	6	10	103.2	61.8	172.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WEBSTER COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	153	62	91	252	146	106	24.0	18.2	42.4
Fetal deaths	4	2	2	9	7	2	35.7	47.9	18.9
Deaths from all causes	80	61	19	102	77	25	9.7	9.6	10.0
Deaths under 1 year	5	1	4	9	5	4	35.7	34.2	37.7
Deaths under 1 month	2	1	1	4	3	1	15.9	20.5	9.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	9.5	12.5	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	9.5	12.5	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	16	15	1	18	17	1	171.4	212.5	40.0
Diabetes (260)	0	0	0	1	1	0	9.5	12.5	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	20	14	6	20	13	7	190.5	162.5	280.0
Heart diseases (410-443)	28	23	5	32	25	7	304.8	312.5	280.0
Arteriosclerosis (450)	2	0	2	2	0	2	19.0	-	80.0
Influenza (480-483)	1	1	0	2	2	0	19.0	25.0	-
#Pneumonia (490-493)	2	1	1	2	1	1	19.0	12.5	40.0
#Diarrhea & enteritis (571)	0	0	0	2	1	1	19.0	12.5	40.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	9.5	12.5	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	2	1	1	19.0	12.5	40.0
Birth inj., atelect. (760-762)	0	0	0	1	1	0	4.0	6.8	-
#Other dis. early inf. (763-776)	2	1	1	3	2	1	11.9	13.7	9.4
Violent & acc. deaths (800-999)	2	2	0	6	5	1	57.1	62.5	40.0
Acc. deaths, total (800-962)	2	2	0	5	4	1	47.6	50.0	40.0
Motor vehicle acc. (810-835)	1	1	0	3	2	1	28.6	25.0	40.0
Ill-defined (780-795)	4	2	2	4	2	2	38.1	25.0	80.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WILKINSON COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	497	190	307	412	89	323	31.2	26.2	33.0
Fetal deaths	5	2	3	5	1	4	12.1	11.2	12.4
Deaths from all causes	143	49	94	125	36	89	9.5	10.6	9.1
Deaths under 1 year	27	8	19	19	4	15	46.1	44.9	46.4
Deaths under 1 month	21	8	13	14	4	10	34.0	44.9	31.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	7.6	29.4	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	7.6	29.4	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	7.6	-	10.2
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	1	0	7.6	29.4	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	12	5	7	10	5	5	75.8	147.0	51.0
Diabetes (260)	0	0	0	2	0	2	15.2	-	20.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	22	11	11	18	7	11	136.4	205.9	112.2
Heart diseases (410-443)	39	14	25	37	11	26	280.3	323.5	265.3
Arteriosclerosis (450)	2	0	2	2	0	2	15.2	-	20.4
Influenza (480-483)	1	0	1	1	0	1	7.6	-	10.2
#Pneumonia (490-493)	2	0	2	2	0	2	15.2	-	20.4
#Diarrhea & enteritis (571)	3	0	3	2	0	2	15.2	-	20.4
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	15.2	29.4	10.2
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	4	3	1	4	3	1	30.3	88.2	10.2
Birth inj., atelect. (760-762)	6	2	4	5	1	4	12.1	11.2	12.4
#Other dis. early inf. (763-776)	10	3	7	5	0	5	12.1	-	15.5
Violent & acc. deaths (800-999)	9	5	4	8	2	6	60.6	58.8	61.2
Acc. deaths, total (800-962)	7	4	3	6	2	4	45.4	58.8	40.8
Motor vehicle acc. (810-835)	3	1	2	3	1	2	22.7	29.4	20.4
Ill-defined (780-795)	15	0	15	17	1	16	128.8	29.4	163.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WINSTON COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	474	196	278	515	214	301	26.4	19.3	35.8
Fetal deaths	16	4	12	19	7	12	36.9	32.7	39.9
Deaths from all causes	158	85	73	182	104	78	9.3	9.4	9.3
Deaths under 1 year	13	1	12	19	4	15	36.9	18.7	49.8
Deaths under 1 month	9	1	8	13	3	10	25.2	14.0	33.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	5.1	-	11.9
Tbc., respiratory (001-008)	1	0	1	1	0	1	5.1	-	11.9
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	5.1	-	11.9
Meningococcal infections (057)	0	0	0	1	1	0	5.1	9.0	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	6	3	13	10	3	66.7	90.1	35.7
Diabetes (260)	3	3	0	4	3	1	20.5	27.0	11.9
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	24	17	7	25	18	7	128.2	162.2	83.3
Heart diseases (410-443)	58	38	20	64	43	21	328.2	387.4	250.0
Arteriosclerosis (450)	0	0	0	1	1	0	5.1	9.0	-
Influenza (480-483)	4	1	3	3	1	2	15.4	9.0	23.8
#Pneumonia (490-493)	5	1	4	6	2	4	30.8	18.0	47.6
#Diarrhea & enteritis (571)	2	1	1	2	1	1	10.2	9.0	11.9
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	5.1	-	11.9
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	2	1	1	10.2	9.0	11.9
Birth inj., atelect. (760-762)	3	1	2	4	1	3	7.8	4.7	10.0
#Other dis. early inf. (763-776)	3	0	3	5	1	4	9.7	4.7	13.3
Violent & acc. deaths (800-999)	19	8	11	21	9	12	107.7	81.1	142.8
acc. deaths, total (800-962)	15	6	9	16	6	10	82.0	54.0	119.0
Motor vehicle acc. (810-835)	8	5	3	7	5	2	35.9	45.0	23.8
Ill-defined (780-795)	14	2	12	14	2	12	71.8	18.0	142.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
YALOBUSH_A COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	291	78	213	326	124	202	25.1	18.2	32.6
Fetal deaths	7	3	4	8	3	5	24.5	24.2	24.8
Deaths from all causes	127	67	60	147	84	63	11.3	12.4	10.2
Deaths under 1 year	8	0	8	9	1	8	27.6	8.1	39.6
Deaths under 1 month	5	0	5	5	1	4	15.3	8.1	19.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	7.7	14.7	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	7.7	14.7	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	15	10	5	17	11	6	130.8	161.8	96.8
Diabetes (260)	2	2	0	2	2	0	15.4	29.4	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	26	15	11	28	19	9	215.4	279.4	145.2
Heart diseases (410-443)	36	20	16	40	22	18	307.7	323.5	290.3
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	3	1	2	3	1	2	23.1	14.7	32.2
#Pneumonia (490-493)	3	2	1	4	2	2	30.8	29.4	32.2
#Diarrhea & enteritis (571)	2	0	2	2	0	2	15.4	-	32.2
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	8	5	3	10	7	3	76.9	102.9	48.4
Maternal deaths (640-689)	1	0	1	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	1	0	1	0	0	0	-	-	-
#Other dis. early inf. (763-776)	4	0	4	4	0	4	12.3	-	19.8
Violent & acc. deaths (800-999)	9	6	3	11	7	4	84.6	102.9	64.5
Acc. deaths, total (800-962)	7	5	2	8	5	3	61.5	73.5	48.4
Motor vehicle acc. (810-835)	3	2	1	1	1	0	7.7	14.7	-
Ill-defined (780-795)	6	2	4	7	3	4	53.8	44.1	64.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
YAZOO COUNTY, MISSISSIPPI, 1957

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	959	256	703	975	281	694	27.1	20.4	31.3
Fetal deaths	44	6	38	44	9	35	45.1	32.0	50.4
Deaths from all causes	353	114	239	355	128	227	9.9	9.3	10.2
Deaths under 1 year	46	5	41	54	8	46	55.4	28.5	66.3
Deaths under 1 month	33	4	29	42	7	35	43.1	24.9	50.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	2	0	2	5.6	-	9.0
Tbc., respiratory (001-008)	2	0	2	2	0	2	5.6	-	9.0
Syphilis (020-029)	1	0	1	1	0	1	2.8	-	4.5
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	2.8	-	4.5
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox. (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	31	14	17	32	20	12	88.9	144.9	54.1
Diabetes (260)	6	2	4	3	1	2	8.3	7.2	9.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	56	19	37	50	19	31	138.9	137.7	139.6
Heart diseases (410-443)	75	33	42	81	38	43	225.0	275.4	193.7
Arteriosclerosis (450)	1	1	0	1	1	0	2.8	7.2	-
Influenza (480-483)	5	0	5	4	0	4	11.1	-	18.0
#Pneumonia (490-493)	12	5	7	13	5	8	36.1	36.2	36.0
#Diarrhea & enteritis (571)	3	0	3	2	0	2	5.6	-	9.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	0	4	3	0	3	8.3	-	13.5
Maternal deaths (640-689)	6	0	6	5	0	5	5.1	-	7.2
Cong. malformations (750-759)	1	1	0	3	3	0	8.3	21.7	-
Birth inj., atelect. (760-762)	4	3	1	5	2	3	5.1	7.1	4.3
#Other dis. early inf. (763-776)	18	1	17	23	2	21	23.6	7.1	30.2
Violent & acc. deaths (800-999)	32	20	12	28	17	11	77.8	123.2	49.5
Acc. deaths, total (800-962)	22	13	9	20	12	8	55.6	87.0	36.0
Motor vehicle acc. (810-835)	8	5	3	7	4	3	19.4	29.0	13.5
Ill-defined (780-795)	66	5	61	64	5	59	177.8	36.2	265.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
STATE OF MISSISSIPPI, 1957

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	61291	27320	33971	60712	26804	33908	28.8	23.4	35.2
Fetal deaths	2486	994	1492	2454	965	1489	40.4	36.0	43.9
Deaths from all causes	20233	10173	10060	20476	10339	10137	9.7	9.0	10.5
Deaths under 1 year	2236	677	1559	2257	694	1563	37.2	25.9	46.1
Deaths under 1 month	1408	528	880	1425	540	885	23.5	20.1	26.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	149	52	97	161	61	100	7.6	5.3	10.4
Tbc., respiratory (001-008)	138	49	89	150	58	92	7.1	5.0	9.5
Syphilis (020-029)	45	10	35	50	15	35	2.4	1.3	3.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	38	5	33	39	5	34	1.8	0.4	3.5
Scarlet fever & streptococcal sore throat (050, 051)	4	3	1	3	2	1	0.1	0.2	0.1
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	14	0	14	14	0	14	0.7	-	1.4
Meningococcal infections (057)	7	3	4	10	6	4	0.5	0.5	0.4
Acute poliomyelitis (080)	2	0	2	3	1	2	0.1	0.1	0.2
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	5	4	1	5	4	1	0.2	0.3	0.1
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	2408	1423	985	2496	1482	1014	118.3	129.2	105.3
Diabetes (260)	254	136	118	260	140	120	12.3	12.2	12.5
Pellagra (281)	3	1	2	3	1	2	0.1	0.1	0.2
Intracranial lesions (330-334)	2774	1404	1370	2800	1420	1380	132.7	123.8	143.4
Heart diseases (410-443)	6356	3716	2640	6367	3706	2661	301.8	323.0	276.4
Arteriosclerosis (450)	231	138	93	234	139	95	11.1	12.1	9.9
Influenza (480-483)	173	54	119	173	53	120	8.2	4.6	12.5
#Pneumonia (490-493)	611	241	370	621	248	373	29.4	21.6	38.7
#Diarrhea & enteritis (571)	170	34	136	162	30	132	7.7	2.6	13.7
Acute nephritis (590, 591)	56	21	35	58	24	34	2.7	2.1	3.5
Chronic nephritis, etc. (592-594)	303	116	187	304	119	185	14.4	10.4	19.2
Maternal deaths (640-689)	77	15	62	75	14	61	1.2	0.5	1.8
Cong. malformations (750-759)	232	143	89	259	168	91	12.3	14.6	9.4
Birth inj., atelect. (760-762)	368	176	192	365	173	192	6.0	6.4	5.7
#Other dis. early inf. (763-776)	772	237	535	777	241	536	12.8	9.0	15.8
Violent & acc. deaths (800-999)	1642	884	758	1660	898	762	78.7	78.3	79.2
Acc. deaths, total (800-962)	1291	694	597	1302	707	595	61.7	61.6	61.8
Motor vehicle acc. (810-835)	567	355	212	551	346	205	26.1	30.2	21.3
All-defined (780-795)	1772	373	1399	1763	366	1397	83.6	31.9	145.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.