


PUBLIC HEALTH STATISTICS
STATE OF
MISSISSIPPI
1958



VITAL STATISTICS
SUMMARIES

Mississippi State Board of Health
Jackson, Mississippi

A. L. Gray, M. D., Executive Officer

ANNUAL BULLETIN

of

VITAL STATISTICS

1958

SUMMARIES

MISSISSIPPI STATE BOARD OF HEALTH

General

Table
1.

2.

3.

4.

5.

6.

7.

8.

9.

10.

11.

12.

13.

14.

15.


16.

17.

18.

TABLE OF CONTENTS

	Page
General Summary	1
Table	
1. Estimated Population as of January 1, 1958, by County, by Race	19
2. Live Births, by County and City of Occurrence, Reallocated to Mother's Usual Residence, by Race, with Rates	21
3. Live Births, by County and City of Usual Residence of Mother, by Race and Attendant, and by Sex	24
4. Fetal Deaths, by County and City of Occurrence, Reallocated to Mother's Usual Residence, by Race and Attendant, and by Sex	28
5. Deaths, by Cause, by Race	32
6. Deaths, by County and City of Occurrence, Reallocated to Place of Residence, by Race, with Rates	43
7. Deaths from Selected Causes, by Race, Sex, and Age, with Rates	46
8. Deaths Under 1 Year, Shown for Selected Causes, by Race and Age, with Rates	54
9. Deaths from Selected Causes, by Month of Occurrence	56
10. Marriages, by County of Occurrence and of Residence of Bride, and Divorces, by County of Occurrence, by Race	57
11. Divorces, by Cause, by Number of Minor Children, by Number of Years Married, by Race	59
12. Live Births, 1913-1958, by Year, by Race, with Rates	60
13. Deaths, 1913-1958, by Year, by Race, with Rates	62
14. Maternal Deaths, 1913-1958, by Year, by Race, with Rates	64
15. Deaths Under 1 Year, 1917-1958, by Year, by Race, with Rates	65
16. Deaths Under 1 Month, 1924-1958, by Year, by Race, with Rates	66
17. Deaths from Heart Diseases, Nephritis (all forms), Vascular Lesions, Cancer, and Tuberculosis (all forms), 1949-1958, by Year, by Race, with Rates	67
18. Births, Deaths, and Marriages, by Month of Occurrence	68



COUNTY TABLES AND STATE SUMMARY TABLE

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 1 MONTH, AND
DEATHS FROM CERTAIN SELECTED CAUSES, ON BASIS OF PLACE OF OCCURRENCE
AND PLACE OF RESIDENCE, BY RACE, WITH RATES:

	Page
For each County	71
For the State of Mississippi	153

prepa
marri

analy

Popul

1958,
Depar

city
citiel

Occur

and r
at th
dence
Missi
nonre
outsi
birth
Fetal
the y
The t
Colum
death
Alask

recom
dence
the b
dence
to pr

SUMMARY OF 1958 VITAL STATISTICS

INTRODUCTION

The 1958 Vital Statistics Bulletin is the sixteenth consecutive issue prepared from data obtained from records of live births, fetal deaths, deaths, marriages and divorces collected by the Division of Public Health Statistics.

In this issue all tables, both for the State and for each county, with an analysis of the data, have been included in one volume.

Population and Rates

Population figures used in this bulletin are estimates as of January 1, 1958, prepared by Dr. George L. Wilber and Mr. Carl T. Robbins, of the Sociology Department, Mississippi State University, according to the "vital rates" method.

No age or sex-specific rates (except for maternal and infant deaths) and no city rates were calculated for 1958 because current estimates by age, sex, and for cities are not available.

Occurrence and Residence Allocation

Births, fetal deaths, deaths, and marriages were classified by occurrence and residence. Occurrence data were obtained from certificates filed in Mississippi at the place of birth, fetal death, death, or issuance of marriage license. Residence data include transcripts of these events (except marriages) occurring to Mississippi residents outside the State, and exclude records filed in Mississippi for nonresidents. There were 1,071 births to residents of Mississippi which occurred outside the State in 1958 and were reported to the Division of Public Health Statistics; births occurring inside Mississippi to residents of other states numbered 1,555. Fetal death transcripts totaled only 12 and the nonresident certificates 64. During the year, 806 death transcripts were received and 637 nonresident deaths were recorded. The total number of states sending transcripts was 40, including the District of Columbia. Birth records came from 29 states, fetal death records from 6 states, and death records from 32 states. Births to 12 Mississippi residents were reported from Alaska. In Table A the states furnishing the largest numbers of records are listed.

Some of the tables contain both residence and occurrence data; but, following recommended procedures, most of the tables and all of the rates are based on residence data only. Residence assignment for births and fetal deaths was determined on the basis of usual residence of mother; and for deaths, on the basis of usual residence of decedent. Deaths of inmates or employees of institutions were reallocated to previous residence regardless of length of stay in the institution.

Table A. Transcripts received, by place of occurrence

<u>Place of Occurrence</u>	<u>Live Births</u>	<u>Deaths</u>	<u>Fetal Deaths</u>
<u>Total</u>	<u>1,071</u>	<u>806</u>	<u>12</u>
Alabama	215	89	2
Arkansas	11	15	0
Colorado	19	1	0
Florida	9	13	0
Illinois	71	14	1
Kentucky	19	4	0
Louisiana	182	184	1
Missouri	26	12	0
Tennessee	416	391	6
Texas	11	24	1
Other places	92	59	1

Urban-Rural Classification

Towns with populations of 2,500 or more according to the 1950 census were considered as urban areas. All other areas were classified as rural.

Classification of Causes of Death

For the first time, causes of death were assigned according to the Seventh Revision (1955) of the International Statistical Classification of Causes of Death. This revision differs only slightly from the Sixth Revision used from 1949 through 1957 and rates for the years prior to 1958 have not been adjusted for comparability with the Seventh Revision. Only underlying causes were coded.

LIVE BIRTHS

Resident Births

Live births to residents of Mississippi numbered 59,189 in 1958. This total was 1,523 less than that for 1957 and was the lowest since 1945. There was a slight increase in the number of births among whites, while there was a decrease of 1,531 for nonwhites. The crude birth rate of 27.1 per 1,000 population was the lowest since 1946. The State rate for whites in 1958 was 20.9 and county rates varied from 10.6 in Issaquena County to 40.6 in Washington County. For nonwhites, the State rate was 35.9 and the range for the counties was from 14.4 in Pontotoc County to 56.2 in Bolivar County.

Characteristics

A breakdown of the 32,377 nonwhite births by specific race shows 32,212 were Negro, 117 were Indian, 30 were Chinese, 13 were Japanese, and 5 were other nonwhites. Births of males numbered 30,043 and there were 29,146 females. These totals yielded a sex ratio of 1,031 males to 1,000 females. August was the peak month of occurrence for live births. There were 5,544 in this month in comparison with only 4,230 in April when the fewest occurred.

During 1958, 20,122 children were born to urban residents and 39,067 to rural residents. Over half (53 per cent) of the infants born to urban residents were white, but 59 per cent of the rural were nonwhite. Of the births to rural residents, 23,873 or 61 per cent were born to farm residents; 13,404 were born to nonfarm residents; and on 1,790 certificates information pertaining to farm residency was not stated. Comparison of the occurrence and residence figures indicates that a large majority of the white mothers living in rural areas went to cities for delivery as compared with a much smaller percentage of the nonwhite mothers.

The number of illegitimate births during 1958 was 7,581 or 12.8 per cent of the total. This was 234 less than in 1957, but the decrease in percentage was only 0.1. White births classified as illegitimate totaled 310 or 1.2 per cent, and 7,271 or 22.4 per cent of the nonwhite births were illegitimate.

Liveborn infants classified as immature (birth weight $5\frac{1}{2}$ pounds or less) numbered 5,499, or 9.2 per cent of the total number of births registered. The incidence of immaturity was greater among the nonwhites than among the whites, 11.4 per cent in comparison with 6.6 per cent.

Classification by Attendant

In Table B, which gives the distribution of live births by attendant and race, it may be seen that, in 1958, over 70 per cent of all births occurred in hospitals or clinics (including physicians' offices or clinics which provide inpatient care). A small proportion 3.9 per cent, was delivered by physicians in places other than hospitals or clinics; and 25.8 per cent were delivered at home by nonmedical attendants. Almost half of the nonwhite mothers, but less than one per cent of the white mothers, had no medical attendant.

Table B. Live births by attendant and race

<u>Attendant</u>	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
<u>Total</u>	<u>59,189</u>	<u>100.0</u>	<u>26,812</u>	<u>100.0</u>	<u>32,377</u>	<u>100.0</u>
Physician in hospital	41,644	70.4	26,439	98.6	15,205	47.0
Physician in home	2,288	3.9	246	0.9	2,042	6.3
Midwife	15,190	25.7	119	0.4	15,071	46.5
Other	67	0.1	8	0.03	59	0.2

There has been a steady increase in the proportion of live births occurring in hospitals in recent years (see Table C). The year 1951 was the first in which at least half of the deliveries took place in hospitals, and the proportion of hospital deliveries has increased each year since. The progress made by the nonwhites has been more rapid, but there is still a large difference between the percentages for two racial groups.

7
1ee
n
9S.
a

ril

Table C. Percentage of live births in hospitals or clinics, 1954-1958, by race

<u>Year</u>	<u>Total</u>	<u>White</u>	<u>Nonwhite</u>
1958	70.4	98.6	47.0
1957	68.3	98.0	44.8
1956	65.7	97.2	41.0
1955	62.4	96.0	36.4
<u>1954</u>	<u>60.2</u>	<u>94.4</u>	<u>33.0</u>

The proportions of live births attended by physicians during 1954-1958 are shown in Table D. The improvement shown in this table is less pronounced than that in the preceding table, but is also steady. This is the third year in which more than half the nonwhite mothers were attended by physicians.

Table D. Percentage of live births attended by physicians, 1954-1958, by race

<u>Year</u>	<u>Total</u>	<u>White</u>	<u>Nonwhite</u>
1958	74.2	99.5	53.3
1957	73.6	99.5	53.2
1956	72.2	99.3	51.0
1955	70.4	99.1	48.3
<u>1954</u>	<u>70.4</u>	<u>98.9</u>	<u>47.7</u>

Age of Mother

Table E gives the age distribution of mothers by race. Little variation in this distribution has occurred in recent years. Approximately half of the mothers were 15-24 years of age and more than one-third were 25-34 years old. In the periods 15-24 and 25-34 years the percentage of white mothers were slightly larger than those of nonwhites.

Table E. Live births by age of mother and race, with per cent of total births

<u>Age of Mother</u>	<u>Total</u>		<u>White</u>		<u>Nonwhite</u>	
	<u>No.</u>	<u>%</u>	<u>No.</u>	<u>%</u>	<u>No.</u>	<u>%</u>
10-14 years	306	0.5	36	0.1	270	0.8
<u>15-44 years</u>	<u>58,646</u>	<u>99.1</u>	<u>26,725</u>	<u>99.7</u>	<u>31,921</u>	<u>98.6</u>
15-24 years	30,099	50.8	13,761	51.3	16,338	50.5
25-34 years	22,036	37.2	10,534	39.3	11,502	35.5
35-44 years	6,511	11.0	2,430	9.1	4,081	12.6
45-54 years	135	0.2	32	0.1	103	0.3
55-64 years	1	0.002	0	-	1	0.003
Unknown	101	0.2	19	0.1	82	0.2

FETAL DEATHS

In 1952 the registration of all fetal deaths, regardless of length of gestation, was initiated in Mississippi. Prior to that time, only stillbirths of 20 or more weeks of gestation were reported. Registration of early fetal deaths is still far

from complete, but there is no estimate of the degree of nonregistration. There were 2,572 fetal death certificates filed during 1958 and 2,520 certificates for residents. The resident total was 66 more than the comparable figure for 1957, the change being entirely among nonwhites. For the State the fetal death ratio (fetal deaths per 1,000 live births) was 42.6. The ratio for whites was 36.0 with county ratios varying from zero in Issaquena, Tate, and Wilkinson Counties to 104.1 in Washington County. The ratio for nonwhites was 48.0 with variations from zero in Hancock, Prentiss, and Tishomingo Counties to 100.0 in Stone.

On 897 of the records the gestation period was stated to be less than 20 weeks; on 1,488 it was 20 weeks or more; and on 135 the length was not stated. In the period 1954-1958 there have been reported among the whites more fetal deaths after less than 20 weeks of gestation than after 20 weeks or more.

Using the certificates with sex stated, a sex ratio of 1,310 males to 1,000 females was determined. This ratio is lower than those for 1956 and 1957, but is comparable to sex ratios for some previous years. As usual, the sex ratio for fetal deaths was much higher than that for live births, but was not so reliable a figure because the sex was not stated on 732 or 29 per cent of the fetal death certificates.

The distribution by race and attendant is given in Table F. Approximately 62 per cent of the total were nonwhite. Sixty-nine per cent of the reported fetal deaths were attended by physicians in hospitals and about 19 per cent were attended by midwives. Although the proportion of white live births in hospitals was larger than the percentage of white fetal deaths in hospitals, the reverse was true for the nonwhites. This situation in the nonwhite group may be accounted for by the nonreporting of fetal deaths that occur at home and the tendency to transfer difficult and premature deliveries to a hospital.

Table F. Fetal deaths by attendant and race

Attendant	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
<u>Total</u>	<u>2,520</u>	<u>100.0</u>	<u>965</u>	<u>100.0</u>	<u>1,555</u>	<u>100.0</u>
Physician in hospital	1,740	69.0	857	88.8	883	56.8
Physician in home	279	11.1	105	10.9	174	11.2
Midwife	477	18.9	2	0.2	475	30.5
<u>Other</u>	<u>24</u>	<u>1.0</u>	<u>1</u>	<u>0.1</u>	<u>23</u>	<u>1.5</u>

Table G shows the percentage of fetal deaths delivered in hospitals in the last five years. There has been very little change in the figures for the whites during this time, but the nonwhites have shown a steady increase from 44.6 per cent in 1954 to 56.8 per cent in 1958. The gap between the two race groups is a large one, but not so large as that for live births.

Table G. Percentage of fetal deaths delivered in hospitals, 1954-1958, by race

<u>Year</u>	<u>Total</u>	<u>White</u>	<u>Nonwhite</u>
1958	69.0	88.8	56.8
1957	69.5	92.2	54.8
1956	67.2	91.0	52.7
1955	63.5	88.1	48.5
1954	61.6	90.2	44.6

DEATHS**Resident Deaths**

In 1958 the resident death total was 21,541, which is an increase of 1,065 over the number for 1957. The crude death rate, based on the 1958 population estimate, was 9.9 per 1,000 population in comparison with a rate of 9.7 in 1957 based on the 1956 population estimate. Slightly over one-third or 7,408 of the persons who died in 1958 were urban residents, but 48 per cent of the deaths occurred in urban areas.

Although the number of deaths of nonwhites was 81 less than that of the whites, the death rate of the nonwhites was 3.5 per 1,000 population greater than that of the whites. The total of 10,811 deaths of white persons was the largest reported since 1913, the first complete year of death registration in Mississippi. The number of deaths of nonwhites in 1958 (10,730) was higher than in 1957, 1956, 1955, 1954, 1946 or 1945. The death rate of whites was slightly less than that in 1957, while the rate for nonwhites increased over the 1957 figure.

County rates for the white population ranged from 4.5 in Harrison to 18.2 in Madison, with a rate for the State of 8.4, while county rates for the nonwhites ranged from 4.4 in Covington to 19.5 in Adams and Bolivar, with a State rate of 11.9.

The resident birth rate exceeded the death rate by 17.2 per 1,000 population. For whites the excess was 12.5 and for the nonwhites it was 24.0.

Age at Death

Table H shows the distribution of deaths by age and race. In comparison with the 1957 figures there were increases in all the white groups except the 25-34 year group which showed a decrease of approximately 4 per cent. Increases occurred in all nonwhite groups except those 15-44 years and the 55-64 year period. For deaths of all races combined there were decreases from 1957 in the groups 15-34 years. Increases were seen in age groups under 15 and 45 years and over, with the major increases in the age groups 65 years and over.

Comparison of the age distribution of the two races reveals significant differences. The percentage of deaths of nonwhites which occurred among children under 10 years of age was more than double that for the whites, and in the groups from 10 through 54 years relatively more deaths of nonwhites than of whites were registered. "Retirement" age or 65 years had been reached by 60 per cent of the whites who died in 1958 and only 46 per cent of the nonwhites. The median age at death for the State was 66.4; for the whites, 69.3; and for the nonwhites, 62.2 years.

Table H. Age-race distribution of deaths

Age Group	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
All Ages	21,541	100.0	10,811	100.0	10,730	100.0
Under 1 year	2,424	11.2	706	6.5	1,718	16.0
1-4 years	454	2.1	136	1.2	318	3.0
5-9 years	163	0.8	64	0.6	99	0.9
10-14 years	142	0.6	67	0.6	75	0.7
15-24 years	438	2.0	204	1.9	234	2.2
25-34 years	553	2.6	237	2.2	316	2.9
35-44 years	953	4.4	441	4.1	512	4.8
45-54 years	1,988	9.2	919	8.5	1,069	10.0
55-64 years	2,972	13.8	1,540	14.2	1,432	13.3
65-74 years	4,769	22.1	2,525	23.4	2,244	20.9
75-84 years	4,478	20.8	2,739	25.3	1,739	16.2
85 years and over	2,188	10.2	1,230	11.4	958	8.9
Unknown	19	0.09	3	0.03	16	0.1

Leading Causes of Death

In Table I are listed the ten leading causes of death in the State in 1958. The first six of these causes are the same as in the three preceding years. Chronic and unspecified nephritis and other renal sclerosis, which was in seventh place in 1956 and 1957, moved to ninth place in 1958. Diabetes mellitus, which was in tenth place in 1956, and in eighth place in 1957, moved to seventh place in 1958. Congenital malformations, which was in eighth place in 1956, and in ninth place in 1957, moved to tenth place in 1958. General arteriosclerosis, which was in eighth place in 1956, and in tenth place in 1957, was again in eighth place in 1958.

Altogether the ten leading causes were responsible for 77 per cent of the deaths. For the whites the same ten causes were listed as for the total population with the exception that suicide ranked in tenth place and chronic and unspecified nephritis and other renal sclerosis was not on the list. For nonwhites, gastritis ranked in eighth place and homicide in ninth place, while general arteriosclerosis and congenital malformations were not on the list.

Table I. Ten leading causes of death by race, with per cent of total deaths

Cause and International List Number	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
1. Diseases of heart (410-443)	6,515	30.2	3,924	36.3	2,591	24.1
2. Vascular lesions affecting central nervous system (330-334)	2,867	13.3	1,426	13.2	1,441	13.4
3. Malignant neoplasms (140-205)	2,525	11.7	1,524	14.1	1,001	9.3
4. Accidents (800-962)	1,379	6.4	729	6.7	650	6.0
5. Certain diseases of early infancy (760-776)	1,248	5.8	451	4.2	797	7.4
6. Pneumonia and influenza (480-493)	909	4.2	344	3.2	565	5.3
7. Diabetes mellitus (260)	317	1.5	163	1.5	154	1.4
8. General arteriosclerosis (450)	268	1.2	178	1.6	90	0.8
9. Chronic & unspec. nephritis & other renal sclerosis (592-594)	264	1.2	89	0.8	175	1.6
10. Congenital malformations (750-759)	254	1.2	141	1.3	113	1.0

Heart disease, the leading cause of death, was responsible for nearly one-third of all deaths in 1958. During the year 6,515 deaths were attributed to some form of heart disease. On Table J it may be seen that a large number of these (4,520 deaths) were specified as arteriosclerotic and degenerative heart disease, and that the next largest group was hypertensive heart disease with 1,381 deaths. The total death rate was 298.3 per 100,000 population, a decrease from 301.8 for 1957. The decrease was entirely among the whites, while an increase occurred among nonwhites.

Table J. Deaths from heart disease, by race by type of disease

Cause of Death and International List Number	Race		
	Total	White	Nonwhite
Diseases of the heart	<u>6,515</u>	<u>3,924</u>	<u>2,591</u>
Chronic rheumatic heart disease (410-416)	129	72	57
Arteriosclerotic and degenerative heart disease (420-422)	4,520	3,098	1,422
Other diseases of heart (430-434)	485	206	279
Hypertension with heart disease (440-443)	1,381	548	833

For the second leading cause of death, vascular lesions affecting the central nervous system, there were 2,867 deaths representing 13.3 per cent of the total. The components of this category are listed in Table K showing that a majority of the deaths were due to cerebral hemorrhage (1,783), and 601 were due to cerebral embolism or thrombosis. There was little change in rate from that of the previous year; and, following the pattern of the years, the rate for nonwhites exceeded that for whites.

Table K. Deaths from vascular lesions affecting the central nervous system, by race by type of lesion

Cause of Death and International List Number	Race		
	Total	White	Nonwhite
Vascular lesions affecting central nervous system (330-334)	<u>2,867</u>	<u>1,426</u>	<u>1,441</u>
Subarachnoid hemorrhage (330)	65	37	28
Cerebral hemorrhage (331)	1,783	821	962
Cerebral embolism and thrombosis (332)	601	374	227
Other and ill-defined (334)	418	194	224

In 1958, the cancer death total was the highest on record in this State, due to an increase in the number for whites. There were 2,525 deaths from this cause or 29 more than in 1957. The distribution of these deaths by age, sex, and race is given in Table L. In comparison with 1957, there were decreases in the number of deaths reported for white females and nonwhite males, while the figures for white males and nonwhite females showed increases. There were changes from 1957 in almost every age group with the largest decreases occurring in deaths of nonwhite males from 45 to 64 years and the largest increases in white males in the age groups from 55 through 84 years.

More deaths occurred among the whites than among nonwhites in all age groups, and the excess was largest in the age group 75-84. There were more males than females in the age groups 55-64, 65-74 and 75-84, and more females in the age groups 25-34, 35-44, 45-54, and 85 years and over.

Table L. Cancer deaths by age, sex, and race

<u>Age Group</u>	Total	Male		Female	
		White	Nonwhite	White	Nonwhite
<u>All Ages</u>	<u>2,525</u>	<u>835</u>	<u>457</u>	<u>689</u>	<u>544</u>
Under 10 years	37	10	7	15	5
10-14 years	18	4	3	7	4
15-24 years	31	14	6	7	4
25-34 years	52	13	5	16	18
35-44 years	156	38	23	48	47
45-54 years	370	89	47	117	117
55-64 years	560	202	106	131	121
65-74 years	677	233	149	164	131
75-84 years	490	191	90	143	66
85 years & over	134	41	21	41	31

Table M lists cancer deaths by primary site in descending order according to the number in the total population. The number of deaths assigned to the leading site, digestive organs and peritoneum, is two more than in 1957, and this site moved to first position due to a decrease of 15 in deaths from cancer of the breast and genito-urinary system. These two leading sites accounted for 77 per cent of the cancer deaths among nonwhite females, 68 per cent among white females and among nonwhite males, and only 46 per cent among white males. Cancer of the respiratory system showed an increase of 36 deaths, remaining in third rank. By far the largest increase was that of 63 in the number of deaths from cancer of the lymphatic and hematopoietic tissues.

Table M. Cancer deaths by primary site, sex, and race, with percentage distribution

Primary site and International List Number	Male				Female			
	White		Nonwhite		White		Nonwhite	
	No.	%	No.	%	No.	%	No.	%
<u>Total</u>	<u>835</u>	<u>100.0</u>	<u>457</u>	<u>100.0</u>	<u>689</u>	<u>100.0</u>	<u>544</u>	<u>100.0</u>
1. Digestive organs and peritoneum (150-159)	249	29.8	176	38.5	209	30.3	161	29.6
2. Breast & genito-urinary system (170-181)	134	16.0	134	29.3	257	37.3	259	47.6
3. Respiratory system (160-165)	194	23.2	67	14.7	35	5.1	22	4.0
4. Lymphatic & hematopoietic tissues (200-205)	101	12.1	37	8.1	85	12.3	31	5.7
5. Brain & nervous system (192, 193)	38	4.6	6	1.3	22	3.2	4	0.7
6. Skin (190, 191)	33	4.0	3	0.6	19	2.8	5	0.9
7. Buccal cavity & pharynx (140-148)	21	2.5	4	0.9	10	1.4	5	0.9
8. Bone & connective tissue (196, 197)	20	2.4	1	0.2	7	1.0	4	0.7
9. Other & unspecified sites (194, 195, 198, 199)	45	5.4	29	6.3	45	6.5	53	9.7

The number of deaths in 1958 which resulted from accidents was 1,379, an increase of 77 over the number in 1957. The greatest increases were in the number of deaths from fire and from drowning. Among whites the greatest increase in accidental deaths was in motor vehicle accidents, and among nonwhites, from fire and explosion. The greatest decrease for whites was in deaths from firearms, and for nonwhites in deaths from motor vehicle accidents. The largest number of deaths from all accidents occurred in December in both 1957 and 1958. October moved from eighth place to second place in the number of deaths from all accidents, and from sixth place to first place in the number of deaths from motor vehicle accidents. Table N shows the six leading causes of accidental death in 1958 by race. The causes and rank order are the same as on the 1957 list except that in 1957 falls caused a larger number of deaths than drownings.

Table N. Six leading causes of accidental deaths among residents, by race

Type of Accident	Total	White	Nonwhite
1. Motor vehicle	554	381	173
2. Fire and explosion	217	44	173
3. Drowning except water transport	125	49	76
4. Falls	123	89	34
5. Firearm	36	19	17
6. Suffocation or obstruction	31	8	23

Leading causes of death for each age group except infants are given in Table O. None of these lists was exactly like any other, but there were some causes which appeared repeatedly. Accidents was listed in every group, and appeared as the chief cause in all groups under 45 years of age. Malignant neoplasms also appeared on every list and heart diseases on all except 1-4 years. Vascular lesions affecting the central nervous system was listed in the three groups 25 and over. Also appearing in five groups (1-4, 5-14, 15-24, 45-64, and 65 years and over) were pneumonia and influenza. In both lists for children under 15, congenital malformations was included, and for the 15-24 and 25-44 year groups homicide was listed. The gastritis group was listed only for the 1-4 year group. Special tabulations of the age group 65 years and over show that the list for age group 75-79 is the same as the list for the age group 65 years and over as a whole. In the age groups 65-74 and 80 and over, accidents does not appear on the list of five leading causes. In the age group 65-69 both diabetes and pneumonia and influenza are in fourth place. In the age group 70-74 diabetes is in fifth place, and in the age group 80-84 arteriosclerosis is in fifth place. In the age group 85 and over pneumonia and influenza moved to third place, malignant neoplasms to fourth place, and arteriosclerosis to fifth place.

Ill-defined Causes

Of the 21,372 certificates filed for deaths that occurred in Mississippi in 1958, those with no satisfactory statement of cause numbered 2,605 or 12.2 per cent of the total. Senility was the only cause given on 67 of these certificates; ill-defined diseases of early infancy and those due to immaturity unqualified on 649; and other ill-defined and unknown causes on 1,889. For white persons the percentage of certificates in this category was 5.3 per cent and for nonwhites 19.1.

Table 0. Five leading causes of death by age group

<u>Age and Cause of Death</u>	<u>No. of Deaths</u>	<u>% of Total</u>
1-4 years		
All Causes	<u>454</u>	<u>100.0</u>
1. Accidents	130	28.6
2. Pneumonia and influenza	64	14.1
3. Gastritis, duodenitis, enteritis and colitis	27	5.9
4. Malignant neoplasms, including leukemia	24	5.3
5. Congenital malformations	23	5.1
5-14 years		
All Causes	<u>305</u>	<u>100.0</u>
1. Accidents	141	46.2
2. Malignant neoplasms, including leukemia	29	9.5
3. Pneumonia and influenza	16	5.2
4. Congenital malformations	12	3.9
5. Heart disease	9	3.0
15-24 years		
All Causes	<u>438</u>	<u>100.0</u>
1. Accidents	219	50.0
2. Homicide	48	11.0
3. Malignant neoplasms, including leukemia	31	7.1
4. Heart diseases	23	5.2
5. Pneumonia and influenza	15	3.4
25-44 years		
All Causes	<u>1,506</u>	<u>100.0</u>
1. Accidents	284	18.8
2. Heart diseases	270	17.9
3. Malignant neoplasms, including leukemia	208	13.8
4. Vascular lesions affecting central nervous system	103	6.8
5. Homicide	88	5.8
45-64 years		
All Causes	<u>4,960</u>	<u>100.0</u>
1. Heart diseases	1,710	34.5
2. Malignant neoplasms, including leukemia	930	18.8
3. Vascular lesions affecting central nervous system	574	11.6
4. Accidents	246	5.0
5. Pneumonia and influenza	122	2.4
65 years and over		
All Causes	<u>11,435</u>	<u>100.0</u>
1. Heart diseases	4,483	39.2
2. Vascular lesions affecting central nervous system	2,172	19.0
3. Malignant neoplasms, including leukemia	1,301	11.4
4. Pneumonia and influenza	437	3.8
5. Accidents	257	2.2

Deaths with unsatisfactory statement of cause were so numerous that if placed on the list of ten leading causes they would rank third and would appear as one of the five chief causes of death in every age group except 15-24. If the true causes of death for these persons could be determined and properly assigned, the order of principal causes might not be the same and even the causes might be different.

If deaths from ill-defined diseases peculiar to early infancy and deaths due to immaturity are excluded from the group of deaths without definite cause, this group is 9.2 per cent of all deaths recorded in Mississippi. In Claiborne and Yazoo Counties they were more than 20 per cent of the total deaths, and in Bolivar, Carroll, Hancock, Humphreys, Leflore, Noxubee, Quitman, Sharkey, and Sunflower Counties 15 to 20 per cent had no definite cause.

Infant Mortality

An infant mortality rate of 41.0 per 1,000 live births was recorded in 1958, higher than the 1957 rate by 3.8 per 1,000 births. The nonwhite rate showed a greater increase than the white rate. Distribution of infant deaths by age and race is shown in Table P. In each age group, the rate for nonwhites exceeded that for whites, with the greatest difference between the two racial groups occurring in the period 28 days to one year.

Table P. Infant deaths by age and race, with rates

Age at Death	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
Total under 1 year	2,424	41.0	706	26.3	1,718	53.1
Under 1 day	693	11.7	293	10.9	400	12.4
One to 28 days	782	13.2	247	9.2	535	16.5
28 days to 1 year	949	16.0	166	6.2	783	24.2

* Deaths per 1,000 live births.

Table Q gives the five leading causes of infant deaths. Gastritis, duodenitis, enteritis and colitis, which was in fifth place in 1957, moved to fourth place in 1958, and birth injuries moved from fourth to fifth place. Immaturity, which is no longer considered a separate classification in ranking causes of infant deaths, was mentioned either in combination with another cause or as the only cause on 45 per cent of the certificates for white infants and on 33 per cent of those for nonwhites.


Table Q. Leading causes of infant deaths, by race

Cause of Death	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
All Causes	2,424	100.0	706	100.0	1,718	100.0
1. Postnatal asphyxia & atelectasis	262	10.8	133	18.8	129	7.5
2. Influenza & pneumonia, except pneumonia of newborn	221	9.1	31	4.4	190	11.0
3. Congenital malformations	191	7.9	98	13.9	93	5.4
4. Gastritis, duodenitis, enteritis & colitis except diarrhea of newborn	133	5.5	11	15.6	122	7.1
5. Birth injuries	128	5.3	62	8.8	66	3.8

Infant death certificates which showed ill-defined causes or no cause at all numbered 965 or 40 per cent of the total in 1958. For 30 per cent of the white and 44 per cent of the nonwhite infants no definite cause of death was given. Determination of the true cause of death for this large group of infants might change the leading causes of death which are listed above.

Infant mortality rates by race for 1920-1958 are shown in Graph 1. In general, the trend has been downward for both races, but in recent years there has been some upward swing in the curve for the nonwhites.

Graph 1. Infant Mortality
1920-1958



Maternal Mortality

The maternal mortality rate for 1958 was 0.9 per 1,000 live births in the total population, 0.4 for white mothers and 1.3 for nonwhite. The total of 53 maternal deaths was 22 less than in 1957. The decrease among the white mothers was 4 deaths and among the nonwhite, 18. There is still a large discrepancy between the rates by race, but the gap has narrowed considerably in recent years.

In 1958, as in each year from 1947 through 1957, the two leading causes were toxemia and hemorrhage. Toxemia has been the leading cause in each year of this period except 1957.

Table R. Maternal deaths by cause and race, with rates

<u>Cause of Death</u>	<u>Total</u>		<u>White</u>		<u>Nonwhite</u>	
	No.	Rate*	No.	Rate*	No.	Rate*
<u>All Causes</u>	<u>53</u>	<u>0.9</u>	<u>10</u>	<u>0.4</u>	<u>43</u>	<u>1.3</u>
Toxemia	20	0.3	4	0.1	16	0.5
Hemorrhage	10	0.2	1	0.04	9	0.3
Sepsis	8	0.1	2	0.1	6	0.2
Ectopic pregnancy	3	0.05	0	-	3	0.1
Abortion without sepsis or toxemia	3	0.05	1	0.04	2	0.1
Abortion with sepsis	1	0.02	0	-	1	0.03
<u>Other complications</u>	<u>8</u>	<u>0.1</u>	<u>2</u>	<u>0.1</u>	<u>6</u>	<u>0.2</u>

* Deaths per 1,000 live births.


Table S gives the age and race-specific maternal death rates. There were no deaths in the 10-14 year group and only one death in the 45-54 year group. At all ages the rates for nonwhite mothers were higher than those for white mothers.

Table S. Maternal deaths by age group and race, with rates

<u>Age Group</u>	<u>Total</u>		<u>White</u>		<u>Nonwhite</u>	
	No.	Rate*	No.	Rate*	No.	Rate*
<u>All Ages</u>	<u>53</u>	<u>0.9</u>	<u>10</u>	<u>0.4</u>	<u>43</u>	<u>1.3</u>
15-24 years	12	0.4	2	0.1	10	0.6
25-34 years	24	1.1	4	0.4	20	1.7
35-44 years	16	2.4	4	1.6	12	2.9
45-54 years	1	7.4	0	-	1	9.7

* Deaths per 1,000 live births to mothers in specific
age-race group.

Maternal death rates by race for 1913-1958 are shown in Graph 2. A downward trend began in the 1930's and has been rapid since 1940.



MARRIAGES

A new marriage law was passed at a special session of the legislature in December 1957 and apparently this law was responsible for the small number of marriages in this state during 1958. Although the law (with a three day waiting period, blood tests and proof of age) did not become effective until July 1, 1958, there was widespread publicity when it was signed and many out-of-state couples evidently did not attempt to secure licenses in Mississippi even as early as December 1957. In 1958 the number of marriages occurring in Mississippi was 36,402. This was a decrease of 26,093 from the 1957 figure. There were 19,870 fewer marriages among the whites and 6,223 fewer among the nonwhites. A small part of the decrease (probably less than 100) is due to the fact that the circuit clerk in Tallahatchie County failed to file records for July through December in one district.

The average daily number of marriages was 100, 71 less than the average for 1957; but in June, the month with the largest total, the daily average was 205. In the first six months of 1958 the number of marriages was 26,642, a decrease of 3,748 from the number in the same period of 1957. In the second six months of 1958 the number of marriages was 9,760, a decrease of 22,345 from the number in the same period of 1957.

This was the first marriage for 25,388 or 69.7 per cent of the brides and 25,424 or 69.8 per cent of the grooms; and for less than one fourth of each group there had been one previous marriage. Two hundred sixty one brides and 218 grooms did not state the number of previous marriages.

Table T shows the age distribution of brides and grooms. Of the brides, 17,313 or 47.6 per cent were under 20 years of age, and 70.6 per cent were under 25 years of age. There was a much smaller percentage of grooms in the teen-age group -- 11.4 per cent -- while 45.8 per cent of the grooms were in the 20-24 year age group. A cross tabulation of the ages of brides and grooms shows marriages of one couple with both parties under 15 years of age and 16 couples with both 75 years of age and over. While the youngest bride of a groom of 75 years of age or more was 15 years of age, most marriages were between persons of approximately the same age.

Table T. Age distribution of brides and grooms

Age	Bride		Groom	
	No.	%	No.	%
<u>Total</u>	<u>36,402</u>	<u>100.0</u>	<u>36,402</u>	<u>100.0</u>
Under 15 years	167	0.4	2	0.005
15 years	873	2.4	6	0.02
16 years	1,621	4.4	76	0.2
17 years	2,077	5.7	550	1.5
18 years	8,661	23.8	1,546	4.2
19 years	3,914	10.8	1,975	5.4
20-24 years	8,372	23.0	16,657	45.8
25-34 years	4,896	13.4	7,884	21.6
35-44 years	2,954	8.1	3,332	9.2
45-54 years	1,744	4.8	2,122	5.8
55 years & over	1,109	3.0	2,243	6.2
Unknown	14	0.04	9	0.02

Among the marriages between nonresident brides and grooms there were 87 brides and no grooms under 15 years of age. Nonresident brides in the 15-19 year age group numbered 8,894 or 51.8 per cent of all brides in this age group.

In Table U are shown data on the residence status of both brides and grooms. In 1958 both parties were residents of Mississippi in 43 per cent of the marriages, with a percentage of 34 during the first six months and of almost 67 during the second six months. In 1957 both parties were residents in only 30 per cent of the marriages. There were 18,920 out-of-state brides or 52 per cent of the total in 1958 and in 1957 there were 41,510 nonresident brides or 66.4 per cent of the total. An analysis of the 1958 marriages by month shows that 16,486 or 87 per cent of the nonresident brides were married during the first six months, before the new law became effective.

Table U. Marriages in Mississippi by residence status of bride and groom

Groom	Total	Bride		
		Resident	Nonresident	Not Stated
<u>Total</u>	<u>36,402</u>	<u>17,370</u>	<u>18,920</u>	<u>112</u>
Resident	16,522	15,569	926	27
Nonresident	19,766	1,776	17,982	8
Not stated	114	25	12	77

A comparison during the last twelve years of the resident status of brides is shown in Table V. It may be seen that in each year 1947 through 1956 the number of out-of-state brides has increased and, with the exception of 1955, the number of resident brides has decreased. The declining trend among resident brides continued in the years 1957 and 1958 but the increase in nonresidents ended abruptly when the new marriage law was passed. Evidently the influx of out-of-state brides began to slow down in December 1957 since, for the first time, December was not the peak month.

Table V. Marriages in Mississippi by residence status of bride 1947-1958

Year	Total	Resident	Nonresident	Not Stated
1958	36,402	17,370	18,920	112
1957	62,495	20,864	41,510	121
1956	66,223	23,011	43,212	*
1955	66,423	23,683	42,740	*
1954	61,381	23,555	37,826	*
1953	61,751	24,497	37,254	*
1952	58,116	24,868	33,248	*
1951	57,321	25,713	31,608	*
1950	56,738	26,859	29,879	*
1949	52,765	27,717	25,048	*
1948	53,645	30,921	22,724	*
1947	53,584	34,067	19,517	*

* Not classified.

Brides came to Mississippi from every state in the union, the District of Columbia, Alaska and Hawaii, and there were 21 from foreign countries.

Bordering states furnished 38.6 per cent of the brides married in Mississippi and 38 per cent of the grooms, with totals as follows: Alabama, 4,971 brides and 4,917 grooms; Arkansas, 1,678 brides and 1,581 grooms; Louisiana, 2,393 brides and 2,461 grooms; and Tennessee, 5,019 brides and 4,886 grooms. Two other states furnished over 1,000 brides and grooms: Illinois, 1,212 brides and 1,376 grooms; and Missouri, 1,057 brides and 1,049 grooms. These 6 states furnished 44.8 per cent of the brides married in Mississippi and 44.6 per cent of the grooms.

Each county in the state issued marriage licenses to one or more out-of-state brides in 1958, but the counties adjacent to Alabama, Tennessee, and Arkansas issued the largest numbers. Table W shows the counties to which the greatest numbers of out-of-state brides came. In Alcorn, DeSoto and Tishomingo Counties over 90 per cent of the licenses were for nonresident brides, and in George, Lowndes and Itawamba Counties this percentage was over 80.. Twenty three per cent of all the brides married in Mississippi were nonresidents who came to DeSoto, Alcorn, and Tishomingo Counties -- counties bordering Tennessee.

Table W. Number of out-of-state brides obtaining marriage license
in certain counties

DeSoto	3,707	George	746
Alcorn	3,476	Harrison	609
Lowndes	1,269	Itawamba	582
Tishomingo	1,265	Hancock	543
Washington	966	Lauderdale	525
Jackson	933	Tate	408

DIVORCES

Divorces granted in Mississippi during 1958 numbered 5,016 - 42 more than in 1957. There was an increase of 135 among the whites and a decrease of 93 among the nonwhites. Cruel and inhuman treatment and desertion continued to rank as the two leading causes and together accounted for 87 per cent of the total granted. Of the divorces among the whites, cruel and inhuman treatment was given as the main cause for 58 per cent; among nonwhites 64 per cent were granted on grounds of desertion. The wife was listed as the complainant on 62.5 per cent of the records. Only 226 suits or 4.5 per cent were contested.

No minor children were involved in 52 per cent of the divorces, and 20 per cent of the couples had only one minor child. Six of the families had 9 or more minor children. The total number of minor children affected was 4,744.

Forty-two per cent of the divorces in 1958 were granted to persons married less than 5 years; 8 per cent had been married less than one year; but 15.5 per cent had been married 20 years or longer. More divorces were granted to couples in their second year of marriage than in any other year. The distribution of divorces according to duration of marriage was different for the two races. Of the white couples, 51.5 per cent were granted after less than 5 years of marriage and only 8 per cent after 20 years or longer; for the nonwhites, corresponding percentages were 21 and 31 respectively.

TABLE 1
ESTIMATED POPULATION*, MISSISSIPPI
AS OF JANUARY 1, 1958

County	Total	White	Nonwhite
<u>TOTAL</u>	<u>2,184,000</u>	<u>1,282,700</u>	<u>901,300</u>
Adams	33,000	20,000	13,000
Alcorn	27,000	21,000	6,000
Amite	20,000	8,100	11,900
Attala	20,000	11,500	8,500
Benton	11,000	5,400	5,600
Bolivar	39,000	15,000	24,000
Calhoun	17,000	11,000	6,000
Carroll	10,000	4,100	5,900
Chickasaw	17,000	8,100	8,900
Choctaw	11,000	7,500	3,500
Claiborne	10,000	3,100	6,900
Clarke	16,000	9,700	6,300
Clay	18,000	10,000	8,000
Coahoma	40,000	15,000	25,000
Copiah	28,000	14,000	14,000
Covington	18,000	10,500	7,500
DeSoto	23,000	9,500	13,500
Forrest	51,000	35,000	16,000
Franklin	8,600	4,800	3,800
George	16,000	14,000	2,000
Greene	7,800	6,100	1,700
Grenada	17,000	8,100	8,900
Hancock	12,000	9,200	2,800
Harrison	165,000	133,000	32,000
Hinds	176,000	107,000	69,000
Holmes	24,000	6,800	17,200
Humphreys	16,000	4,900	11,100
Issaquena	3,700	1,800	1,900
Itawamba	21,000	19,900	1,100
Jackson	59,000	40,500	18,500
Jasper	17,000	9,200	7,800
Jefferson	10,000	2,800	7,200
Jeff Davis	17,000	9,800	7,200
Jones	66,000	48,000	18,000
Kemper	12,000	5,300	6,700
Lafayette	28,000	19,000	9,000

<u>County</u>	<u>Total</u>	<u>White</u>	<u>Nonwhite</u>
Lamar	19,000	15,500	3,500
Lauderdale	59,000	37,000	22,000
Lawrence	12,000	8,400	3,600
Leake	22,000	13,000	9,000
Lee	44,000	32,000	12,000
Leflore	40,000	13,000	27,000
Lincoln	24,000	16,000	8,000
Lowndes	41,000	23,000	18,000
Madison	25,000	6,000	19,000
Marion	26,000	16,000	10,000
Marshall	23,000	6,400	16,600
Monroe	30,000	18,000	12,000
Montgomery	14,000	7,600	6,400
Neshoba	25,000	16,400	8,600
Newton	18,000	12,000	6,000
Noxubee	14,000	5,500	8,500
Oktibbeha	27,000	15,000	12,000
Panola	29,000	13,000	16,000
Pearl River	28,000	20,000	8,000
Perry	7,700	5,500	2,200
Pike	31,000	18,000	13,000
Pontotoc	21,000	16,000	5,000
Prentiss	19,000	16,000	3,000
Quitman	19,000	6,700	12,300
Rankin	36,000	22,000	14,000
Scott	20,000	12,000	8,000
Sharkey	9,200	3,000	6,200
Simpson	24,000	16,000	8,000
Smith	15,000	10,500	4,500
Stone	6,000	4,500	1,500
Sunflower	43,000	16,000	27,000
Tallahatchie	22,000	9,000	13,000
Tate	17,000	9,000	8,000
Tippah	17,000	12,500	4,500
Tishomingo	17,000	16,500	500
Tunica	16,000	2,000	14,000
Union	24,000	19,000	5,000
Walthall	15,000	9,000	6,000
Warren	37,000	19,000	18,000
Washington	63,000	26,000	37,000
Wayne	16,000	10,000	6,000
Webster	13,000	10,000	3,000
Wilkinson	10,000	3,000	7,000
Winston	20,000	11,000	9,000
Yalobusha	13,000	7,000	6,000
Yazoo	28,000	11,000	17,000

* Prepared by Dr. George L. Wilber and Mr. Carl T. Robbins, Sociology Department, Mississippi State University.

TABLE 2
LIVE BIRTHS
MISSISSIPPI
1958

By Place of Occurrence, Reallocated to Mother's Usual Residence,
by Race, with Rates

County	Place of Occurrence			Place of Residence					
	Number		Non-white	Number			Non-white	Rate*	
	Total	White		Total	White	Non-white		Total	White
STATE	59,673	27,231	32,442	59,189	26,812	32,377	27.1	20.9	35.9
Adams	1,703	704	999	1,223	549	674	37.1	27.4	51.8
Alcorn	599	481	118	513	403	110	19.0	19.2	18.3
Amite	257	41	216	384	127	257	19.2	15.7	21.6
Attala	527	228	299	490	201	289	24.5	17.5	34.0
Benton	116	5	111	232	86	146	21.1	15.9	26.1
Bolivar	1,895	404	1,491	1,775	425	1,350	45.5	28.3	56.2
Calhoun	382	203	179	337	190	147	19.8	17.3	24.5
Carroll	238	40	198	284	77	207	28.4	18.8	35.1
Chickasaw	421	199	222	467	219	248	27.5	27.0	27.9
Choctaw	189	90	99	190	91	99	17.3	12.1	28.3
Claiborne	236	31	205	288	39	249	28.8	12.6	36.1
Clarke	229	112	117	430	201	229	26.9	20.7	36.3
Clay	474	191	283	495	202	293	27.5	20.2	36.6
Coahoma	1,496	363	1,133	1,531	327	1,204	38.3	21.8	48.2
Copiah	877	248	629	716	221	495	25.6	15.8	35.4
Covington	221	158	63	382	216	166	21.2	20.6	22.1
DeSoto	643	60	583	703	170	533	30.6	17.9	39.5
Forrest	1,471	1,081	390	1,271	787	484	24.9	22.5	30.2
Franklin	133	60	73	232	96	136	27.0	20.0	35.8
George	337	282	55	345	288	57	21.6	20.6	28.5
Greene	112	62	50	189	120	69	24.2	19.7	40.6
Grenada	555	230	325	516	194	322	30.4	24.0	36.2
Hancock	135	103	32	266	217	49	22.2	23.6	17.5
Harrison	3,691	2,991	700	3,394	2,722	672	20.6	20.5	21.0
Hinds	5,546	2,926	2,620	5,146	2,568	2,578	29.2	24.0	37.4
Holmes	804	135	669	786	120	666	32.8	17.6	38.7
Humphreys	662	109	553	664	115	549	41.5	23.5	49.4
Issaquena	36	0	36	104	19	85	28.1	10.6	44.7
Itawamba	62	53	9	241	224	17	11.5	11.2	15.4
Jackson	1,277	889	388	1,532	1,120	412	26.0	27.6	22.3
Jasper	124	45	79	438	152	286	25.8	16.5	36.7
Jefferson	117	0	117	320	40	280	32.0	14.3	38.9
Jeff Davis	346	147	199	390	129	261	22.9	13.2	36.2
Jones	2,365	1,208	1,157	1,506	968	538	22.8	20.2	29.9
Kemper	206	37	169	294	60	234	24.5	11.3	34.9
Lafayette	505	259	246	524	261	263	18.7	13.7	29.2

Table 2 - Page 2
Live Births - Mississippi - 1958

County	Place of Occurrence			Place of Residence			Rate*		
	Number		Non-white	Number		Non-white	Rate*		Non-white
	Total	White		Total	White		Total	White	
Lamar	114	77	37	327	255	72	17.2	16.4	20.6
Lauderdale	2,088	1,061	1,027	1,591	851	740	27.0	23.0	33.6
Lawrence	180	23	157	256	124	132	21.3	14.8	36.7
Leake	345	147	198	411	168	243	18.7	12.9	27.0
Lee	1,135	830	305	976	658	318	22.2	20.6	26.5
Leflore	1,557	395	1,162	1,522	340	1,182	38.0	26.2	43.8
Lincoln	697	421	276	671	369	302	28.0	23.1	37.8
Lowndes	1,176	581	595	1,158	558	600	28.2	24.3	33.3
Madison	1,048	181	867	1,033	184	849	41.3	30.7	44.7
Marion	525	310	215	593	354	239	22.8	22.1	23.9
Marshall	719	111	608	742	137	605	32.3	21.4	36.4
Monroe	849	435	414	795	408	387	26.5	22.7	32.2
Montgomery	311	136	175	305	124	181	21.8	16.3	28.3
Neshoba	457	254	203	482	277	205	19.3	16.9	23.8
Newton	388	245	143	424	204	220	23.6	17.0	36.7
Noxubee	550	103	447	519	87	432	37.1	15.8	50.8
Oktibbeha	824	362	462	759	312	447	28.1	20.8	37.2
Panola	798	254	544	831	264	567	28.6	20.3	35.4
Pearl River	686	497	189	582	393	189	20.8	19.6	23.6
Perry	79	58	21	220	158	62	28.6	28.7	28.2
Pike	1,021	537	484	895	407	488	28.9	22.6	37.5
Pontotoc	292	216	76	301	229	72	14.3	14.3	14.4
Prentiss	457	396	61	395	338	57	20.8	21.1	19.0
Quitman	780	202	578	728	179	549	38.3	26.7	44.6
Rankin	248	44	204	699	385	314	19.4	17.5	22.4
Scott	555	319	236	558	288	270	27.9	24.0	33.8
Sharkey	233	39	194	355	65	290	38.6	21.7	46.8
Simpson	527	263	264	486	262	224	20.2	16.4	28.0
Smith	127	90	37	342	207	135	22.8	19.7	30.0
Stone	159	111	48	150	100	50	25.0	22.2	33.3
Sunflower	1,354	263	1,091	1,459	311	1,148	33.9	19.4	42.5
Tallahatchie	790	132	658	826	174	652	37.5	19.3	50.2
Tate	450	22	428	584	138	446	34.4	15.3	55.8
Tippah	348	265	83	313	234	79	18.4	18.7	17.6
Tishomingo	98	90	8	237	219	18	13.9	13.3	36.0
Tunica	655	63	592	661	71	590	41.3	35.5	42.1
Union	461	365	96	402	306	96	16.8	16.1	19.2
Walton	619	362	257	350	137	213	23.3	15.2	35.5
Warren	1,467	594	873	1,165	504	661	31.5	26.5	36.7
Washington	2,820	1,124	1,696	2,707	1,057	1,650	43.0	40.6	44.6
Wayne	452	314	138	430	256	174	26.9	25.6	29.0
Webster	125	67	58	203	130	73	15.6	13.0	24.3

Table 2 - Page 3
Live Births - Mississippi - 1958

County	Place of Occurrence			Place of Residence					
	Number		Non-white	Number			Rate*		
	Total	White		Total	White	Non-white	Total	White	Non-white
Wilkinson	441	188	253	387	73	314	38.7	24.3	44.8
Winston	442	182	260	470	188	282	23.5	17.1	31.3
Yalobusha	305	86	219	313	105	208	24.1	15.0	34.7
Yazoo	934	241	693	978	280	698	34.9	25.4	41.0
Out-of-State	1,071	886	186	1,555	1,304	251	-	-	-

Cities over 10,000 Population**	Place of Occurrence			Place of Residence		
	Number		Non-white	Number		Non-white
	Total	White		Total	White	
Biloxi	2,073	1,831	242	1,519	1,320	199
Clarksdale	840	357	483	599	228	371
Columbus	943	569	374	584	268	316
Greenville	1,427	749	678	1,305	582	723
Greenwood	796	386	410	574	212	362
Gulfport	1,475	1,139	336	868	608	260
Hattiesburg	1,451	1,081	370	851	480	371
Jackson	5,093	2,925	2,168	3,227	1,476	1,751
Laurel	2,302	1,173	1,129	802	437	365
McComb	670	402	268	309	190	119
Meridian	2,019	1,054	965	1,062	593	469
Natchez	778	647	131	811	347	464
Pascagoula	77	0	77	545	392	153
Tupelo	867	687	180	371	256	115
Vicksburg	1,435	594	841	820	357	463
Total Urban#	33,449	20,280	13,169	20,122	10,731	9,391
Total Rural	26,224	6,951	19,273	39,067	16,081	22,986

* Rate per 1,000 population.

** Cities having 10,000 population and over according to 1950 census.

Includes all towns of 2,500 population or more according to 1950 census.

TABLE 3
LIVE BIRTHS
MISSISSIPPI
1958

By County and Cities over 10,000, by Race, Attendant, and Sex
on Basis of Usual Residence of Mother

County	TOTAL*	WHITE			NONWHITE			SEX			
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female
STATE	59,189	26,812	26,439	246	127	32,377	15,205	2,042	15,130	30,043	29,146
Adams	1,243	549	549	0	0	674	648	1	25	612	611
Alcorn	513	403	386	16	1	110	50	34	26	250	263
Amitte	384	127	125	2	0	257	80	11	166	203	181
Attala	490	201	196	5	0	289	78	37	174	235	255
Benton	232	86	82	2	2	146	37	10	99	104	128
Bolivar	1,775	425	399	15	11	1,350	567	75	708	924	851
Calhoun	337	190	185	4	1	147	103	7	37	151	186
Carroll	284	77	74	1	2	207	39	6	162	145	139
Chickasaw	467	219	216	3	0	248	102	30	116	238	229
Choctaw	190	91	90	0	1	99	46	2	51	96	94
Claiborne	288	39	39	0	0	249	57	2	190	140	148
Clarke	430	201	200	0	1	229	163	7	59	209	221
Clay	495	202	201	1	0	293	169	23	101	243	252
Coahoma	1,531	327	320	5	2	1,204	343	192	669	763	768
Copiah	716	221	220	1	0	495	330	20	145	390	326
Covington	382	216	215	0	1	166	140	6	20	192	190
DeSoto	703	170	166	1	3	533	210	15	308	346	357
Forrest	1,271	787	786	1	0	484	339	4	141	639	632
Franklin	232	96	96	0	0	136	104	2	30	121	111
George	345	288	286	1	1	57	34	2	21	190	155
Greene	189	120	113	6	1	69	26	7	36	112	77
Grenada	516	194	194	0	0	322	36	59	227	252	264
Hancock	266	217	216	1	0	49	20	0	29	131	135
Harrison	3,394	2,722	2,697	5	20	672	355	23	294	1,763	1,631

Table 3 - Page 2
Live Births - Mississippi - 1958

County	TOTAL*	WHITE			NONWHITE			SEX			
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female
Hinds	5,146	2,568	2,567	1	0	2,578	1,790	133	655	2,613	2,533
Holmes	786	120	117	3	0	666	104	48	514	386	400
Humphreys	664	115	111	0	4	549	93	5	451	331	333
Issaquena	104	19	19	0	0	85	48	3	34	53	51
Itawamba	241	224	221	3	0	17	13	2	2	122	119
Jackson	1,532	1,120	1,110	4	6	412	236	48	128	783	749
Jasper	438	152	148	4	0	286	225	5	56	209	229
Jefferson	320	40	40	0	0	280	161	0	119	150	170
Jeff Davis	390	129	121	6	2	261	137	23	101	177	213
Jones	1,506	968	965	2	1	538	494	2	42	778	728
Kemper	294	60	60	0	0	234	88	1	145	139	155
Lafayette	524	261	257	2	2	263	40	8	215	287	237
Lamar	327	255	247	8	0	72	49	13	10	176	151
Lauderdale	1,591	851	850	1	0	740	643	18	79	836	755
Lawrence	256	124	117	3	4	132	57	4	71	126	130
Leake	411	168	165	3	0	243	113	4	126	240	171
Lee	976	658	653	5	0	318	187	68	63	483	493
Leflore	1,522	340	336	3	1	1,182	155	46	981	822	700
Lincoln	671	369	365	1	3	302	209	30	63	333	338
Lowndes	1,158	558	541	13	4	600	204	51	345	561	597
Madison	1,033	184	184	0	0	849	212	35	602	524	509
Marion	593	354	346	6	2	239	93	5	141	294	299
Marshall	742	137	133	0	4	605	105	92	408	380	362
Monroe	795	408	403	4	1	387	167	47	173	396	399
Montgomery	305	124	122	0	2	181	36	0	145	153	152
Neshoba	482	277	274	1	2	205	140	3	62	241	241
Newton	424	204	200	4	0	220	164	19	37	206	218
Noxubee	519	87	86	0	1	432	107	12	313	283	236
Oktibbeha	759	312	309	2	1	447	285	4	158	369	390
Panola	831	264	259	5	0	567	247	60	260	413	418

Table 3 - Page 3
Live Births - Mississippi - 1958

County	TOTAL*	WHITE			NONWHITE			SEX		
		Total	Phys. in Hosp.	Phys. in Home	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female
Pearl River	582	393	392	0	189	181	1	7	292	290
Perry	220	158	156	2	0	62	56	3	117	103
Pike	895	407	405	1	1	488	337	21	130	428
Pontotoc	301	229	224	5	0	72	24	8	40	133
Prentiss	395	338	337	1	0	57	35	5	17	213
Quitman	728	179	176	3	0	549	305	30	214	341
Rankin	699	385	384	1	0	314	254	7	53	328
Scott	558	288	282	5	1	270	132	2	136	291
Sharkey	355	65	65	0	0	290	161	7	122	190
Simpson	486	262	262	0	0	224	152	5	67	248
Smith	342	207	204	2	1	135	105	14	16	169
Stone	150	100	100	0	0	50	20	0	30	68
Sunflower	1,459	311	300	6	5	1,148	335	150	663	724
Tallahatchie	826	174	160	7	7	652	107	71	474	405
Tate	584	138	129	9	0	446	46	25	375	283
Tippah	313	234	214	16	4	79	28	33	18	152
Tishomingo	237	219	214	5	0	18	12	6	0	118
Tunica	661	71	65	3	3	590	177	28	385	351
Union	402	306	297	3	6	96	24	3	69	213
Walton	350	137	136	0	1	213	165	1	47	164
Warren	1,165	504	503	0	1	661	479	1	181	575
Washington	2,707	1,057	1,049	4	4	1,650	886	168	596	1,331
Wayne	430	256	253	1	2	174	121	0	53	217
Webster	203	130	126	4	0	73	26	9	38	109
Wilkinson	387	73	73	0	0	314	103	2	209	211
Winston	470	188	186	2	0	282	83	2	197	248
Yalobusha	313	105	103	2	0	208	68	64	76	161
Yazoo	978	280	267	10	3	698	105	12	581	501

Table 3 - Page 4
Live Births - Mississippi - 1958

Cities over 10,000 Population	TOTAL*	WHITE			NONWHITE			SEX		
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male
Biloxi	1,519	1,320	1,314	3	3	199	144	1	54	771
Clarksdale	599	228	227	1	0	371	168	124	79	306
Columbus	584	268	261	5	2	316	98	30	188	285
Greenville	1,305	582	582	0	0	723	373	67	283	672
Greenwood	574	212	212	0	0	362	80	0	282	305
Gulfport	868	608	604	0	4	260	128	17	115	455
Hattiesburg	851	480	479	1	0	371	253	2	116	419
Jackson	3,227	1,476	1,475	1	0	1,751	1,366	113	272	1,667
Laurel	802	437	435	2	0	365	346	1	18	407
McComb	309	190	190	0	0	119	102	5	12	162
Meridian	1,062	593	593	0	0	469	432	15	22	554
Natchez	811	347	347	0	0	464	448	1	15	397
Pascagoula	545	392	392	0	0	153	76	10	67	279
Tupelo	371	256	256	0	0	115	70	22	23	184
Vicksburg	820	357	357	0	1	463	326	1	136	417
										403

* Includes transcripts furnished by the National Office of Vital Statistics of births to residents of Mississippi occurring out-of-state.

TABLE 4
FETAL DEATHS
MISSISSIPPI
1958

By Place of Occurrence, Reallocated to Mother's Usual Residence
by Sex, Race and Attendant

County	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total in Hosp.	Phys. in Hosp.	Other	Total in Hosp.	Phys. in Hosp.	Other	Male	Female	Unde- termined
STATE	2,572	2,520	965	857	105	3	1,555	883	174	498	1,014
Adams	59	37	8	8	0	0	29	26	0	3	21
Alcorn	23	16	10	8	2	0	6	5	1	8	4
Amité	7	12	1	1	0	0	11	7	1	3	0
Attala	12	11	1	1	0	0	10	1	1	8	2
Benton	3	9	4	4	0	0	5	2	1	2	2
Bolivar	59	57	11	10	1	0	46	22	1	23	33
Calhoun	17	13	6	6	0	0	7	6	0	1	4
Carroll	13	18	2	2	0	0	16	3	4	9	10
Chickasaw	11	14	5	5	0	0	9	6	2	1	6
Choctaw	6	8	4	4	0	0	4	3	1	5	4
Claiborne	8	13	2	1	1	0	11	4	0	10	2
Clarke	3	10	3	3	0	0	7	5	0	2	3
Clay	9	9	2	2	0	0	7	6	1	0	5
Coahoma	83	79	8	8	0	0	71	31	15	25	36
Copiah	23	24	6	5	1	0	18	15	0	3	11
Covington	3	4	3	3	0	0	1	1	0	0	2
DeSoto	15	26	1	1	0	0	25	14	3	10	12
Forrest	49	48	23	23	0	0	25	18	3	4	16
Franklin	7	8	2	2	0	0	6	3	1	2	4
George	8	9	5	5	0	0	4	2	0	2	2
Greene	5	4	2	2	0	0	2	0	0	2	1
Grenada	21	3	3	3	0	0	18	8	4	6	11
Hancock	2	3	3	3	0	0	0	0	0	1	2
Harrison	85	79	48	48	1	0	31	20	1	10	45

Table 4 - Page 2
Fetal Deaths - Mississippi - 1958

County	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total	Phys. in Hosp.	Other	Total	Phys. in Hosp.	Other	Male	Female	Unde- termined
Hinds	387	348	209	137	72	0	139	91	24	88	42
Holmes	44	45	8	7	0	1	37	18	16	14	23
Humphreys	28	34	3	3	0	0	31	11	0	13	17
Issaquena	2	2	0	0	0	0	2	1	1	1	1
Itawamba	0	3	2	2	0	0	1	1	0	2	1
Jackson	52	59	44	41	3	0	15	12	2	1	26
Jasper	12	27	8	7	1	0	19	10	1	8	12
Jefferson	9	16	1	1	0	0	15	8	3	7	9
Jeff Davis	11	13	3	3	29	0	22	20	0	7	10
Jones	94	51	29	21	0	0	8	4	0	20	21
Kemper	7	9	1	1	6	2	12	1	2	4	3
Lafayette	20	20	8	6	2	0	0	0	9	12	5
Lamar	0	11	8	8	0	0	3	3	0	5	3
Lauderdale	98	65	22	22	3	1	6	6	6	6	27
Lawrence	5	9	3	1	1	1	11	6	1	6	22
Leake	6	12	1	1	11	27	11	9	2	5	1
Lee	27	22	27	27	0	0	67	32	5	40	37
Leflore	108	94	0	0	0	0	0	0	0	0	0
Lincoln	27	26	9	9	0	0	17	11	4	2	16
Lowndes	35	38	19	16	2	0	19	10	4	5	15
Madison	55	58	6	6	0	0	52	22	6	6	23
Marion	36	36	21	20	4	0	15	9	7	8	12
Marshall	45	30	4	4	14	0	26	11	2	3	7
Monroe	26	27	14	14	0	0	13	8	2	9	7
Montgomery	13	15	1	1	0	0	14	12	0	3	2
Neshoba	10	17	6	6	11	0	11	15	0	2	1
Newton	41	40	19	18	4	0	21	15	4	3	20
Noxubee	32	31	4	4	5	0	27	17	12	10	8
Oktibbeha	25	22	5	5	6	0	30	13	7	10	0
Panola	28	36	6	6	0	0	0	0	0	0	12

Table 4 - Page 3
Fetal Deaths - Mississippi - 1958

County	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total	Phys. in Hosp.	Other	Total	Phys. in Hosp.	Other	Male	Female	Undetermined
Pearl River	27	27	16	15	1	0	11	10	0	9	11
Perry	1	10	7	7	0	0	3	2	1	5	3
Pike	36	31	4	4	0	1	27	24	0	15	11
Pontotoc	15	14	9	8	0	0	5	5	0	6	4
Prentiss	9	8	8	8	0	0	0	0	0	7	0
Quitman	19	17	2	2	0	0	15	6	3	6	9
Rankin	12	27	21	13	8	0	6	5	0	1	5
Scott	16	20	6	5	0	0	1	8	0	6	5
Sharkey	3	9	1	1	0	0	0	6	2	4	0
Simpson	16	17	9	9	0	1	5	4	0	3	3
Smith	2	10	5	5	0	1	5	5	0	8	1
Stone	11	9	4	3	0	1	5	1	3	3	4
Sunflower	60	70	18	17	1	0	0	52	19	6	27
Tallahatchie	32	34	3	3	0	0	0	31	14	2	15
Tate	18	24	0	0	9	9	1	24	7	3	13
Tippah	11	10	9	9	0	0	0	0	0	0	8
Tishomingo	3	5	5	4	2	0	0	0	0	0	2
Tunica	19	19	2	2	0	0	0	17	4	2	11
Union	32	26	23	23	0	0	0	0	3	0	4
Walthall	31	19	6	6	0	0	0	13	11	0	2
Warren	77	45	12	12	0	0	0	33	26	1	6
Washington	204	203	110	108	2	0	0	93	66	1	13
Wayne	10	12	5	5	0	0	0	7	5	0	0
Webster	2	5	2	2	0	0	0	3	0	3	0
Wilkinson	21	23	0	0	0	0	0	23	9	0	14
Winston	20	24	8	5	4	1	0	19	10	2	7
Yalobusha	8	46	6	2	2	0	0	6	3	0	4
Yazoo	43	46	6	6	0	0	0	40	15	1	21

Table 4 - Page 4
Fetal Deaths - Mississippi - 1958

Cities over 10,000 Population	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total Phys. in Hosp.	Total Phys. in Home	Other	Total Phys. in Hosp.	Total Phys. in Home	Other	Male	Female	Unde- termined
Biloxi	41	33	23	22	1	0	10	7	3	22	10
Clarksdale	56	24	5	5	0	0	19	9	3	14	5
Columbus	31	24	12	9	2	1	12	4	4	10	5
Greenville	139	141	95	93	2	0	46	37	5	20	11
Greenwood	81	48	24	24	0	0	24	17	0	16	24
Gulfport	43	27	9	9	0	0	18	11	1	6	5
Hattiesburg	48	35	16	16	0	0	19	14	2	3	10
Jackson	345	231	138	86	52	0	93	67	19	7	57
Laurel	94	34	14	14	0	0	20	18	2	0	16
McComb	23	11	3	3	0	0	8	7	0	1	5
Meridian	94	49	17	17	0	0	32	25	5	21	16
Natchez	22	33	7	7	0	0	26	25	0	1	18
Pascagoula	5	19	14	13	1	0	5	4	1	0	8
Tupelo	26	11	5	5	0	0	6	5	1	0	5
Vicksburg	77	31	9	9	0	0	22	16	1	1	15

TABLE 5
DEATHS
MISSISSIPPI
1958

By Cause by Race on Basis of Place of Residence

International List Number	Cause of Death*	Total	White	Non-white
<u>001-999</u>	<u>TOTAL ALL CAUSES</u>	<u>21,541</u>	<u>10,811</u>	<u>10,730</u>
<u>001-019</u>	<u>Tuberculosis, all forms</u>	<u>172</u>	<u>82</u>	<u>90</u>
002	Pulmonary	145	73	72
003	Pleural	1	0	1
008	Unspecified site	8	2	6
010	Meninges and central nervous system	9	2	7
011	Intestines, peritoneum, and mesenteric glands	3	0	3
015	Lymphatic system	1	1	0
016	Genito-urinary system	3	2	1
019	Disseminated	2	2	0
<u>020-029</u>	<u>Syphilis and its sequelae</u>	<u>41</u>	<u>8</u>	<u>33</u>
020	Congenital	3	0	3
022	Aneurysm of aorta	15	5	10
023	Other cardiovascular	17	2	15
025	General paralysis of insane	1	1	0
026	Other central nervous system	3	0	3
027	Other forms of late syphilis	2	0	2
031	Chronic gonococcal infection of genito-urinary system	1	0	1
041	Paratyphoid fever	1	1	0
042	Other Salmonella infections	2	0	2
<u>045-048</u>	<u>Dysentery, all forms</u>	<u>33</u>	<u>7</u>	<u>26</u>
045	Bacillary	16	1	15
046	Amebiasis	12	6	6
048	Unspecified forms	5	0	5
049	Food poisoning	3	2	1
<u>050, 051</u>	<u>Scarlet fever and streptococcal sore throat</u>	<u>2</u>	<u>1</u>	<u>1</u>
051	Streptoccal sore throat	2	1	1
053	Septicemia and pyemia	19	7	12
055	Diphtheria	7	1	6
056	Whooping cough	12	0	12
057	Meningococcal infections	22	16	6
061	Tetanus	16	1	15
064	Other bacterial diseases	1	0	1
070	Vincent's infection	1	0	1
080	Acute poliomyelitis	1	1	0
081	Late effects of acute poliomyelitis	1	0	1
082	Acute infectious encephalitis	10	7	3
083	Late effects of acute infectious encephalitis	1	1	0
085	Measles	15	5	10
087	Chickenpox	7	2	5
088	Herpes zoster	1	1	0
089	Mumps	1	1	0
092	Infectious hepatitis	15	8	7
096	Other diseases attributable to viruses	7	3	4

Table 5 - Page 2
Deaths - Mississippi - 1958

International List Number	Cause of Death*	Total	White	Non-white
122	Other protozoal diseases	1	1	0
130	Infestation with worms of other, mixed, or unspecified type	5	0	5
132	Actinomycosis	2	2	0
134	Other fungus infections	7	3	4
138	Other infective and parasitic diseases	6	1	5
<u>140-205</u>	<u>Malignant neoplasm</u>	<u>2,525</u>	<u>1,524</u>	<u>1,001</u>
<u>140-148</u>	<u>Buccal cavity and pharynx</u>	<u>40</u>	<u>31</u>	<u>9</u>
140	Lip	2	2	0
141	Tongue	6	4	2
142	Salivary glands	7	7	0
144	Other parts of mouth and mouth, unspecified	6	4	2
145	Oral mesopharynx	3	3	0
146	Nasopharynx	4	3	1
148	Pharynx, unspecified	12	8	4
<u>150-159</u>	<u>Digestive organs and peritoneum</u>	<u>795</u>	<u>458</u>	<u>337</u>
150	Esophagus	33	18	15
151	Stomach	207	94	113
152	Small intestine, including duodenum	11	4	7
153	Large intestine except rectum	200	133	67
154	Rectum	69	41	28
155	Biliary passages and liver (stated to be primary site)	35	25	10
156	Liver (secondary and unspecified)	84	44	40
157	Pancreas	129	88	41
158	Peritoneum	6	5	1
159	Unspecified digestive organs	21	6	15
<u>160-165</u>	<u>Respiratory system</u>	<u>318</u>	<u>229</u>	<u>89</u>
160	Nose, nasal cavities, middle ear, and accessory sinuses	7	5	2
161	Larynx	22	12	10
162	Trachea, and bronchus and lung specified as primary	108	87	21
163	Lung and bronchus, unspecified as to whether primary or secondary	171	118	53
164	Mediastinum	2	2	0
165	Thoracic organs (secondary)	8	5	3
<u>170-181</u>	<u>Breast and genito-urinary organs</u>	<u>784</u>	<u>391</u>	<u>393</u>
170	Breast	178	102	76
171	Cervix uteri	144	63	81
172	Corpus uteri	9	4	5
174	Uterus, unspecified	89	35	54
175	Ovary, Fallopian tube, and broad ligament	53	29	24
176	Other and unspecified female genital organs	13	9	4
177	Prostate	203	104	99
178	Testis	3	3	0
179	Other and unspecified male genital organs	6	3	3
180	Kidney	22	9	13
181	Bladder and other urinary organs	64	30	34
<u>190-199</u>	<u>Other and unspecified sites</u>	<u>334</u>	<u>229</u>	<u>105</u>
190	Melanoma of skin	26	26	0

Table 5 - Page 3
Deaths - Mississippi - 1958

International List Number	Cause of Death*	Total	White	Non-white
191	Skin, other	34	26	8
192	Eye	9	7	2
193	Brain and other parts of nervous system	61	53	8
194	Thyroid gland	6	2	4
195	Other endocrine glands	4	2	2
196	Bone including jaw bone	25	21	4
197	Connective tissue	7	6	1
198	Lymph nodes, secondary and unspecified	7	6	1
199	Other and unspecified sites	155	80	75
<u>200-205</u>	<u>Lymphatic and hematopoietic tissues</u>	<u>254</u>	<u>186</u>	<u>68</u>
200	Lymphosarcoma and reticulosarcoma	50	39	11
201	Hodgkin's disease	33	26	7
202	Other forms of lymphoma (reticulosis)	5	4	1
203	Multiple myeloma (plasmacytoma)	39	19	20
204	Leukemia and aleukemia	127	98	29
<u>210-229</u>	<u>Benign neoplasms</u>	<u>31</u>	<u>12</u>	<u>19</u>
212	Respiratory system	2	0	2
214	Uterine fibromyoma	7	1	6
215	Uterus, other	1	0	1
216	Ovary	1	0	1
223	Brain and other parts of nervous system	12	8	4
224	Endocrine glands	3	2	1
227	Other muscular and connective tissues	2	0	2
228	Hemangioma and lymphangioma	3	1	2
<u>230-239</u>	<u>Neoplasms of unspecified nature</u>	<u>37</u>	<u>28</u>	<u>9</u>
230	Digestive organs	3	3	0
231	Respiratory organs	3	1	2
236	Other genito-urinary organs	3	1	2
237	Brain and other parts of nervous system	24	21	3
238	Skin and musculoskeletal system	1	1	0
239	Other and unspecified organs	3	1	2
241	Asthma	58	33	25
245	Other allergic disorders	7	4	3
251	Non-toxic nodular goiter	1	1	0
252	Thyrotoxicosis with or without goiter	7	1	6
253	Myxedema and cretinism	3	2	1
254	Other diseases of thyroid gland	2	1	1
260	Diabetes mellitus	317	163	154
270	Disorders of pancreatic internal secretion other than diabetes mellitus	1	1	0
272	Diseases of pituitary gland	4	3	1
274	Diseases of adrenal glands	2	2	0
277	Polyglandular dysfunction and other diseases of endocrine glands	1	0	1
281	Pellagra	2	0	2
286	Other avitaminoses and nutritional deficiency states	13	2	11
287	Obesity, not specified as of endocrine origin	6	1	5
289	Other metabolic diseases	6	2	4
<u>290-293</u>	<u>Anemias</u>	<u>61</u>	<u>25</u>	<u>36</u>
290	Pernicious and other hyperchromic	5	3	2
291	Iron deficiency (hyperchromic)	3	0	3
292	Other specified type	37	16	21

Table 5 - Page 4
Deaths - Mississippi - 1958

International List Number	Cause of Death*	Total	White	Non-white
293	Unspecified type	16	6	10
294	Polycythemia	5	3	2
295	Hemophilia	1	0	1
296	Purpura and other hemorrhagic conditions	9	5	4
297	Agranulocytosis	1	1	0
298	Diseases of spleen	2	2	0
299	Other diseases of blood and blood-forming organs	3	1	2
304	Senile psychosis	5	2	3
307	Alcoholic psychosis	2	2	0
309	Other and unspecified psychoses	1	0	1
316	Psychoneurosis with somatic symptoms affecting digestive system	1	0	1
322	Alcoholism	13	4	9
325	Mental deficiency	4	4	0
<u>330-334</u>	<u>Vascular lesions affecting central nervous system</u>	<u>2,867</u>	<u>1,426</u>	<u>1,441</u>
330	Subarachnoid hemorrhage	65	37	28
331	Cerebral hemorrhage	1,783	821	962
332	Cerebral embolism and thrombosis	601	374	227
334	Other and ill-defined	418	194	224
340	Meningitis, except meningococcal and tuberculous	40	14	26
342	Intracranial and intraspinal abscess	6	1	5
343	Encephalitis, myelitis, and encephalomylitis (except acute infectious)	7	4	3
344	Late effects of intracranial abscess or pyogenic infection	8	3	5
345	Multiple sclerosis	11	10	1
350	Paralysis agitans	16	13	3
351	Cerebral spastic infantile paralysis	4	2	2
352	Other cerebral paralysis	11	5	6
353	Epilepsy	32	8	24
355	Other diseases of brain	11	4	7
356	Motor neurone disease and muscular atrophy	11	5	6
357	Other diseases of spinal cord	2	0	2
364	Polyneuritis and polyradiculitis	2	2	0
391	Otitis media without mention of mastoiditis	2	1	1
<u>400-402</u>	<u>Rheumatic fever</u>	<u>13</u>	<u>5</u>	<u>8</u>
400	Without mention of heart involvement	1	1	0
401	With heart involvement	12	4	8
<u>410-443</u>	<u>Diseases of heart</u>	<u>6,515</u>	<u>3,924</u>	<u>2,591</u>
410	Diseases of mitral valve	53	28	25
411	Diseases of aortic valve	9	7	2
414	Other endocarditis specified as rheumatic	5	3	2
415	Other myocarditis specified as rheumatic	3	1	2
416	Other heart disease specified as rheumatic	59	33	26
420	Arteriosclerotic heart disease including coronary disease	3,825	2,750	1,075
421	Chronic endocarditis not specified as rheumatic	89	49	40

Table 5 - Page 5
Deaths - Mississippi - 1958

International List Number	Cause of Death*	Total	White	Non-white
422	Other myocardial degeneration	606	299	307
430	Acute and subacute endocarditis	7	2	5
431	Acute myocarditis not specified as rheumatic	7	3	4
433	Functional disease of heart	47	35	12
434	Other and unspecified diseases of heart	424	166	258
440	Essential benign hypertensive heart disease	1	0	1
441	Essential malignant hypertension with heart disease	20	4	16
442	Hypertensive heart disease with arteriolar nephrosclerosis	311	119	192
443	Other and unspecified hypertensive heart disease	1,049	425	624
<u>444-447</u>	<u>Hypertension without mention of heart</u>	<u>235</u>	<u>97</u>	<u>138</u>
444	Essential benign	59	21	38
445	Essential malignant	70	19	51
446	With arteriolar nephrosclerosis	98	56	42
447	Other hypertensive disease	8	1	7
450	General arteriosclerosis	268	178	90
<u>451-468</u>	<u>Other diseases of circulatory system</u>	<u>185</u>	<u>109</u>	<u>76</u>
451	Aortic aneurysm specified as nonsyphilitic, and dissecting aneurysm	66	46	20
452	Other aneurysm, except of heart and aorta	4	1	3
453	Peripheral vascular disease	6	5	1
454	Arterial embolism and thrombosis	5	4	1
455	Gangrene of unspecified cause	6	1	5
456	Other diseases of arteries	5	2	3
460	Varicose veins of lower extremities	1	1	0
461	Hemorrhoids	2	1	1
462	Varicose veins of other specified sites	4	4	0
463	Phlebitis and thrombophlebitis of lower extremities	16	10	6
464	Phlebitis and thrombophlebitis of other sites	10	8	2
465	Pulmonary embolism and infarction	40	14	26
466	Other venous embolism and thrombosis	7	6	1
467	Other diseases of circulatory system	12	5	7
468	Certain diseases of lymph nodes and lymph channels	1	1	0
470	Acute nasopharyngitis (common cold)	6	0	6
471	Acute sinusitis	1	0	1
472	Acute pharyngitis	1	0	1
473	Acute tonsilitis	5	0	5
474	Acute laryngitis and tracheitis	1	0	1
475	Acute upper respiratory infarction of multiple or unspecified site	6	2	4
<u>480-483</u>	<u>Influenza</u>	<u>196</u>	<u>62</u>	<u>134</u>
480	With pneumonia	125	36	89
481	With other respiratory manifestations, and influenza unqualified	59	21	38
482	With digestive manifestations, but without respiratory symptoms	6	3	3

Table 5 - Page 6
Deaths - Mississippi - 1958

International List Number	Cause of Death*	Total	White	Non-white
483	With nervous manifestations, but without digestive or respiratory symptoms	6	2	4
<u>490-493</u>	<u>Pneumonia</u>	<u>713</u>	<u>282</u>	<u>431</u>
490	Lobar	158	53	105
491	Broncho	302	131	171
492	Primary atypical	38	16	22
493	Other and unspecified	215	82	133
<u>500-502</u>	<u>Bronchitis</u>	<u>44</u>	<u>29</u>	<u>15</u>
500	Acute	13	7	6
501	Unqualified	13	8	5
502	Chronic	18	14	4
510	Hypertrophy of tonsils and adenoids	2	2	0
512	Chronic pharyngitis and nasopharyngitis	1	0	1
517	Other diseases of upper respiratory tract	5	4	1
518	Empyema	2	0	2
519	Pleurisy	3	1	2
520	Spontaneous pneumothorax	1	0	1
521	Abscess of lung	2	2	0
522	Pulmonary congestion and hypostasis	24	18	6
525	Other chronic interstitial pneumonia	23	20	3
526	Bronchiectasis	25	19	6
527	Other diseases of lung and pleural cavity	81	67	14
537	Diseases of salivary glands	2	1	1
538	Other diseases of buccal cavity	2	0	2
539	Diseases of esophagus	4	2	2
<u>540, 541</u>	<u>Ulcer of stomach and duodenum</u>	<u>102</u>	<u>73</u>	<u>29</u>
540	Stomach	54	36	18
541	Duodenum	48	37	11
543	Gastritis and duodenitis	3	1	2
544	Disorders of function of stomach	3	1	2
545	Other diseases of stomach and duodenum	10	6	4
<u>550-553</u>	<u>Appendicitis</u>	<u>12</u>	<u>2</u>	<u>10</u>
550	Acute	12	2	10
<u>560, 561, 570</u>	<u>Hernia and intestinal obstruction</u>	<u>127</u>	<u>64</u>	<u>63</u>
560	Hernia of abdominal cavity without mention of obstruction	8	4	4
561	Hernia of abdominal cavity with obstruction	23	11	12
570	Intestinal obstruction without mention of hernia	96	49	47
571	Gastro-enteritis and colitis, except ulcerative tive, age 4 weeks, and over	198	35	163
572	Chronic enteritis and ulcerative colitis	19	16	3
573	Functional disorders of intestines	2	2	0
575	Abscess of anal and rectal regions	1	1	0
576	Peritonitis	4	1	3
578	Other diseases of intestines and peritoneum	36	22	14
580	Acute and subacute yellow atrophy of liver	8	4	4
581	Cirrhosis of liver	95	64	31
582	Suppurative hepatitis and liver abscess	1	1	0
583	Other diseases of liver	13	8	5
584	Cholelithiasis	29	20	9
585	Cholecystitis without mention of calculi	23	12	11
586				

Table 5 - Page 7
 Deaths - Mississippi - 1958

International List Number	Cause of Death*	Total	White	Non-white
586-523	Other diseases of gall bladder and biliary ducts	15	10	5
587	Diseases of pancreas	38	30	8
590,591	<u>Acute nephritis and nephritis with edema</u>	58	11	47
590	Acute	28	6	22
591	With edema, including nephrosis	30	5	25
592-594	<u>Chronic and unspecified nephritis and other renal sclerosis</u>	264	89	175
592	Chronic nephritis	213	76	137
593	Nephritis not specified as acute or chronic	50	13	37
594	Other renal sclerosis	1	0	1
600	Infection of kidney	115	60	55
601	Hydronephrosis	3	2	1
602	Calculi of kidney and ureter	10	7	3
603	Other diseases of kidney and ureter	5	1	4
604	Calculi of other parts of urinary system	1	1	0
605	Cystitis	4	4	0
606	Other diseases of bladder	7	1	6
608	Stricture of urethra	7	0	7
609	Other diseases of urethra	3	0	3
610	Hyperplasia of prostate	85	36	49
611	Prostatitis	7	3	4
612	Other diseases of prostate	1	1	0
624	Salpingitis and oophoritis, unqualified	2	1	1
630	Infective disease of uterus, vagina, and vulva	4	2	2
633	Other diseases of uterus	3	0	3
640-689	<u>Diseases of pregnancy, childbirth, and puerperium</u>	53	10	43
640	Pyelitis and pyelonephritis of pregnancy	1	0	1
642	Toxemias of pregnancy	19	4	15
645	Ectopic pregnancy	3	0	3
648	Other complications arising from pregnancy	1	0	1
650	Abortion without mention of sepsis or toxemia	3	1	2
651	Abortion with sepsis	1	0	1
660	Delivery without stated complications	1	1	0
670	Delivery complicated by placenta previa or antepartum hemorrhage	3	0	3
672	Delivery complicated by other postpartum hemorrhage	7	1	6
674	Delivery complicated by disproportion or malposition of fetus	1	0	1
675	Delivery complicated by prolonged labor of other origin	2	0	2
677	Delivery with other trauma	2	0	2
681	Sepsis of childbirth and puerperium	4	2	2
682	Puerperal phlebitis and thrombosis	1	0	1
684	Puerperal pulmonary embolism	2	0	2
686	Other forms of puerperal toxemia	1	0	1

Table 5 - Page 8
 Deaths - Mississippi - 1958

International List Number	Cause of Death*	Total	White	Non-white
688	Other and unspecified complications of the puerperium	1	1	0
690	Boil and carbuncle	1	0	1
692	Other cellulitis and abscess without mention of lymphangitis	2	0	2
704	Pemphigus	1	0	1
705	Erythematous conditions	1	1	0
710	Other hypertrophic and atrophic conditions of skin	2	2	0
715	Chronic ulcer of skin	3	1	2
722	Rheumatoid arthritis and allied conditions	18	11	7
723	Osteo-arthritis (arthrosis) and allied conditions	6	6	0
725	Arthritis, unspecified	15	9	6
727	Rheumatism, unspecified	1	1	0
730	Osteomyelitis and periostitis	2	1	1
731	Osteitis deformans	1	1	0
733	Other diseases of bone	2	2	0
744	Other diseases of muscle, tendon, and fascia	8	6	2
<u>750-759</u>	<u>Congenital malformations</u>	<u>254</u>	<u>141</u>	<u>113</u>
750	Monstrosity	10	6	4
751	Spina bifida and meningocele	13	9	4
752	Congenital hydrocephalus	25	10	15
753	Other nervous system and sense organs	6	5	1
754	Circulatory system	126	74	52
756	Digestive system	31	13	18
757	Genito-urinary system	11	10	1
758	Bone and joint	5	0	5
759	Other and unspecified, not elsewhere classified	27	14	13
<u>760-762</u>	<u>Birth injuries, postnatal asphyxia and atelectasis</u>	<u>390</u>	<u>195</u>	<u>195</u>
760	Intracranial and spinal injury at birth	58	28	30
761	Other birth injury	70	34	36
762	Postnatal asphyxia and atelectasis	262	133	129
<u>763-768</u>	<u>Infections of newborn</u>	<u>120</u>	<u>40</u>	<u>80</u>
763	Pneumonia	84	35	49
764	Diarrhea	16	0	16
767	Umbilical sepsis	5	1	4
768	Other sepsis	15	4	11
<u>769-776</u>	<u>Other diseases peculiar to early infancy, and immaturity, unqualified</u>	<u>738</u>	<u>216</u>	<u>522</u>
769	Neonatal disorders arising from maternal toxemia	25	11	14
770	Hemolytic disease of newborn (erythroblastosis fetosa)	28	14	14
771	Hemorrhagic disease of newborn	11	3	8
772	Nutritional maladjustment	34	2	32
773	Ill-defined diseases peculiar to early infancy	95	37	58
774	Immaturity with mention of any other subsidiary condition	25	8	17
776	Immaturity, unqualified	520	141	379

Table 5 - Page 9
 Deaths - Mississippi - 1958

International List Number	Cause of Death*	Total	White	Non-white
<u>780-795</u>	<u>Symptoms, senility, and ill-defined conditions</u>	<u>1,930</u>	<u>362</u>	<u>1,568</u>
780	Certain symptoms referable to nervous system and special senses	11	3	8
781	Other symptoms referable to nervous system and special senses	1	1	0
782	Symptoms referable to cardiovascular and lymphatic system	45	17	28
783	Symptoms referable to respiratory system	8	4	4
784	Symptoms referable to upper gastro-intestinal tract	7	1	6
785	Symptoms referable to abdomen and lower gastro-intestinal tract	6	5	1
786	Symptoms referable to genito-urinary system	1	0	1
788	Other general symptoms	5	2	3
789	Abnormal urinary constituents of unspecified cause	1	1	0
792	Uremia, unqualified	12	7	5
794	Senility, without mention of psychosis	68	15	53
795	Ill-defined and unknown causes	1,765	306	1,459
<u>800-962</u>	<u>Accidents</u>	<u>1,379</u>	<u>729</u>	<u>650</u>
800-802	Railway	16	6	10
<u>810-835</u>	<u>Motor vehicle</u>	<u>554</u>	<u>381</u>	<u>173</u>
810	Collision with railway train, traffic	19	13	6
812	Traffic, to pedestrian	97	56	41
813	Traffic, to pedal cyclist	8	6	2
815	Traffic, to rider or passenger of motorcycle in collision with other motor vehicle	2	2	0
816	Other traffic, two or more motor vehicles	218	156	62
818	Traffic, collision with animal or animal-drawn vehicle	6	4	2
819	Traffic, collision with fixed or unspecified object	24	16	8
821	Noncollision traffic, to rider of motorcycle	1	1	0
822	Traffic, overturning in roadway	14	10	4
823	Traffic, running off roadway	117	82	35
824	Other noncollision traffic	8	4	4
825	Traffic, unspecified nature	24	18	6
830	Nontraffic, to pedestrian	14	11	3
834	Nontraffic, boarding and alighting	1	1	0
835	Nontraffic, other and unspecified nature	1	1	0
845	Other nonmotor road vehicle	7	4	3
850	Submersion of occupant of small boat	24	15	9
851	Other water transport injury by submersion	2	2	0
857	Other specified water transport	3	3	0
866	Other and unspecified aircraft	7	7	0
870	Poisoning by morphine and other opium derivatives	1	1	0
871	Poisoning by barbituric acid and derivatives	1	1	0
872	Poisoning by aspirin and salicylates	2	2	0
874	Poisoning by other analgesic and soporific drugs	1	1	0

Table 5 - Page 10
Deaths - Mississippi - 1958

International List Number	Cause of Death*	Total	White	Non-white
876	Poisoning by strychnine	1	0	1
880	Poisoning by alcohol	3	0	3
881	Poisoning by petroleum products	4	1	3
882	Poisoning by industrial solvents	1	0	1
886	Poisoning by arsenic and antimony, and their compounds	1	0	1
888	Poisoning by other and unspecified solid and liquid substances	4	0	4
890	Poisoning by utility (illuminating) gas	1	1	0
900	Fall on stairs	8	5	3
902	Other falls from one level to another	26	21	5
903	Fall on same level	16	13	3
904	Unspecified falls	73	50	23
910	Blow from falling object	26	13	13
912	Caused by machinery	28	16	12
913	Caused by cutting and piercing instrument	5	3	2
914	Caused by electrical current	25	16	9
916	Caused by fire and explosion of combustible material	217	44	173
917	Caused by hot substance, corrosive liquid, and steam	12	4	8
919	Caused by firearm	36	19	17
921	Inhalation and ingestion of food causing obstruction or suffocation	25	7	18
922	Inhalation and ingestion of other object causing obstruction or suffocation	6	1	5
923	Foreign body entering other orifice	1	0	1
924	Mechanical suffocation in bed and cradle	29	9	20
925	Mechanical suffocation in other and unspecified circumstances	10	2	8
926	Lack of care of infants under 1 year of age	2	0	2
927	Caused by bites and stings of venomous animals and insects	1	0	1
928	Other accidents caused by animals	4	1	3
929	Drowning and submersion	125	49	76
931	Excessive heat and insolation	3	2	1
932	Excessive cold	8	1	7
933	Hunger, thirst, and exposure	2	2	0
934	Cataclysm	10	6	4
935	Lightning	7	1	6
936	Other and unspecified accidents	22	9	13
950	Therapeutic misadventure in surgical treatment	4	1	3
951	Therapeutic misadventure in infusion or transfusion	2	1	1
954	Therapeutic misadventure in anesthesia	1	1	0
960	Late effects of motor vehicle accident	2	0	2
962	Late effects of other accidental injury	10	7	3
<u>963, 970-979</u>	<u>Suicide</u>	<u>149</u>	<u>124</u>	<u>25</u>
970	By analgesic and soporific substances	2	2	0
971	By other solid and liquid substances	3	3	0

Table 5 - Page 11
 Deaths - Mississippi - 1958

International List Number	Cause of Death*	Total	White	Non-white
973	By other gases	3	3	0
974	By hanging and strangulation	10	9	1
975	By submersion (drowning)	1	1	0
976	By firearms and explosives	129	106	23
979	By other and unspecified means	1	0	1
<u>964, 980-985</u>	<u>Homicide</u>	<u>207</u>	<u>48</u>	<u>159</u>
981	Assault by firearm and explosive	126	35	91
982	Assault by cutting and piercing instruments	52	8	44
983	Assault by other means	20	3	17
984	Injury by intervention of police	7	2	5
985	Execution	2	0	2

* When cause of death and International List Numbers are omitted, no deaths were reported.

TABLE 6
DEATHS
MISSISSIPPI
1958

By Place of Occurrence, Reallocated to Place of Residence,
by Race, with Rates

County	Place of Occurrence			Place of Residence					
	Number		Non-white	Number		Non-white	Rate*		Non-white
	Total	White		Total	White		Total	White	
STATE	21,372	10,706	10,666	21,541	10,811	10,730	9.9	8.4	11.9
Adams	435	146	289	385	132	253	11.7	6.6	19.5
Alcorn	262	224	38	261	222	39	9.7	10.6	6.5
Amite	83	38	45	134	78	56	6.7	9.6	4.7
Attala	227	126	101	249	143	106	12.4	12.4	12.5
Benton	38	21	17	71	39	32	6.4	7.2	5.7
Bolivar	627	137	490	647	178	469	16.6	11.9	19.5
Calhoun	130	101	29	143	113	30	8.4	10.3	5.0
Carroll	87	21	66	133	51	82	13.3	12.4	13.9
Chickasaw	152	85	67	166	97	69	9.8	12.0	7.8
Choctaw	91	62	29	105	71	34	9.5	9.5	9.7
Claiborne	100	24	76	136	37	99	13.6	11.9	14.3
Clarke	152	87	65	189	111	78	11.8	11.4	12.4
Clay	178	71	107	185	74	111	10.3	7.4	13.9
Coahoma	566	141	425	597	143	454	14.9	9.5	18.2
Copiah	278	124	154	315	151	164	11.2	10.8	11.7
Covington	88	60	28	116	83	33	6.4	7.9	4.4
DeSoto	164	37	127	225	66	159	9.8	6.9	11.8
Forrest	480	324	156	474	301	173	9.3	8.6	10.8
Franklin	81	41	40	102	54	48	11.9	11.2	12.6
George	75	63	12	84	72	12	5.2	5.1	6.0
Greene	68	54	14	83	64	19	10.6	10.5	11.2
Grenada	221	97	124	201	84	117	11.8	10.4	13.1
Hancock	99	75	24	133	103	30	11.1	11.2	10.7
Harrison	895	709	186	779	596	183	4.7	4.5	5.7
Hinds	2,138	1,109	1,029	1,502	706	796	8.5	6.6	11.5
Holmes	306	81	225	335	88	247	14.0	12.9	14.4
Humphreys	175	43	132	207	55	152	12.9	11.2	13.7
Issaquena	30	8	22	38	10	28	10.3	5.6	14.7
Itawamba	107	97	10	131	118	13	6.2	5.9	11.8
Jackson	364	257	107	383	275	108	6.5	6.8	5.8
Jasper	114	46	68	184	84	100	10.8	9.1	12.8
Jefferson	87	21	66	128	33	95	12.8	11.8	13.2
Jeff Davis	136	64	72	145	64	81	8.5	6.5	11.2
Jones	664	410	254	557	375	182	8.4	7.8	10.1
Kemper	104	47	57	128	57	71	10.7	10.8	10.6
Lafayette	158	85	73	173	98	75	6.2	5.2	8.3

Table 6 - Page 2
Deaths - Mississippi - 1958

County	Place of Occurrence			Place of Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Lamar	74	57	17	106	81	25	5.6	5.2	7.1
Lauderdale	880	530	350	724	420	304	12.3	11.4	13.8
Lawrence	67	28	39	118	66	52	9.8	7.8	14.4
Leake	160	95	65	188	112	76	8.5	8.6	8.4
Lee	364	257	107	360	252	108	8.2	7.9	9.0
Leflore	611	243	368	538	166	372	13.4	12.8	13.8
Lincoln	278	157	121	293	166	127	12.2	10.4	15.9
Lowndes	397	199	198	388	192	196	9.5	8.3	10.9
Madison	358	121	237	367	109	258	14.7	18.2	13.6
Marion	217	119	98	230	129	101	8.8	8.1	10.1
Marshall	250	64	186	238	68	170	10.3	10.6	10.2
Monroe	331	194	137	353	204	149	11.8	11.3	12.4
Montgomery	118	62	56	156	87	69	11.1	11.4	10.8
Neshoba	189	120	69	227	154	73	9.1	9.4	8.5
Newton	213	141	72	222	143	79	12.3	11.9	13.2
Noxubee	190	53	137	196	56	140	14.0	10.2	16.5
Oktibbeha	202	82	120	214	91	123	7.9	6.1	10.2
Panola	299	125	174	340	139	201	11.7	10.7	12.6
Pearl River	197	140	57	210	154	56	7.5	7.7	7.0
Perry	48	29	19	89	53	36	11.6	9.6	16.4
Pike	397	226	171	399	216	183	12.9	12.0	14.1
Pontotoc	165	125	40	191	148	43	9.1	9.2	8.6
Prentiss	164	148	16	148	133	15	7.8	8.3	5.0
Quitman	183	47	136	230	71	159	12.1	10.6	12.9
Rankin	388	176	212	226	128	98	6.3	5.8	7.0
Scott	216	116	100	239	130	109	12.0	10.8	13.6
Sharkey	102	26	76	116	32	84	12.6	10.7	13.5
Simpson	234	146	88	228	148	80	9.5	9.2	10.0
Smith	74	62	12	133	107	26	8.9	10.2	5.8
Stone	61	44	17	69	45	24	11.5	10.0	16.0
Sunflower	400	95	305	460	129	331	10.7	8.1	12.2
Tallahatchie	248	71	177	280	89	191	12.7	9.9	14.7
Tate	114	45	69	172	76	96	10.1	8.4	12.0
Tippah	153	120	33	168	134	34	9.9	10.7	7.6
Tishomingo	88	84	4	136	130	6	8.0	7.9	12.0
Tunica	192	28	164	206	30	176	12.9	15.0	12.6
Union	207	161	46	205	164	41	8.5	8.6	8.2
Walthall	131	87	44	117	67	50	7.8	7.4	8.3
Warren	689	251	438	513	210	303	13.9	11.0	16.8
Washington	784	234	550	798	209	589	12.7	8.0	15.9
Wayne	138	85	53	174	109	65	10.9	10.9	10.8
Webster	72	59	13	88	69	19	6.8	6.9	6.3

Table 6 - Page 3
Deaths - Mississippi - 1958

County	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Wilkinson	155	69	86	144	49	95	14.4	16.3	13.6
Winston	161	85	76	196	106	90	9.8	9.6	10.0
Yalobusha	131	66	65	155	87	68	11.9	12.4	11.3
Yazoo	322	98	224	369	127	242	13.2	11.5	14.2
Out-of-State	806	603	203	637	498	139	-	-	-

Cities over 10,000 Population**	Place of Occurrence			Place of Residence		
	Number			Number		
	Total	White	Non-white	Total	White	Non-white
Biloxi	276	220	56	278	223	55
Clarksdale	326	120	206	272	99	173
Columbus	265	145	120	204	101	103
Greenville	465	180	285	408	131	277
Greenwood	368	160	208	267	115	152
Gulfport	341	259	82	219	152	67
Hattiesburg	388	270	118	297	176	121
Jackson	1,841	986	855	1,084	530	554
Laurel	539	313	226	294	163	131
McComb	194	135	59	145	100	45
Meridian	757	466	291	505	296	209
Natchez	266	126	140	288	105	183
Pascagoula	73	48	25	139	102	37
Tupelo	200	139	61	132	87	45
Vicksburg	607	218	389	396	173	223
Total Urban#	10,337	5,722	4,615	7,408	3,890	3,518
Total Rural	11,035	4,984	6,051	14,133	6,921	7,212

* Rate per 1,000 population.

** Cities having 10,000 population and over according to 1950 census.

Includes all towns of 2,500 population or more according to 1950 census.

TABLE 7
DEATHS FROM SELECTED CAUSES
MISSISSIPPI
1958

By Age, Race, and Sex, on Basis of Place of Residence, with Rates

Inter-national List Number	Cause of Death	Total Rate*	Total by Race	Total by Sex and Race	Age in Years							Unknown Over 65								
					1-4	5-9	10-14	15-24	25-34	35-44	45-54									
001-999	<u>ALL CAUSES</u>	<u>21.541</u>	<u>2.2</u>	<u>W 10,811</u>	M 6,371	F 4,440	M 401	F 305	M 67	F 55	M 173	F 288	M 620	F 1,083	M 3,510	F 2,984	M 0	F 0		
001-138	<u>Infective and parasitic diseases</u>	<u>41.3</u>	<u>18.9</u>	<u>C 251</u>	M 162	F 102	M 54	F 39	M 7	F 4	M 2	F 2	M 1	F 1	M 15	F 18	M 5	F 7	M 1	F 1
001-019	<u>Tuberculosis (all forms)</u>	<u>17.2</u>	<u>7.9</u>	<u>W 82</u>	M 66	F 59	M 16	F 16	M 1	F 0	M 0	F 0	M 0	F 0	M 1	F 1	M 1	F 1	M 1	F 1
001-008	<u>Tuberculosis (respiratory system)</u>	<u>15.4</u>	<u>7.0</u>	<u>C 79</u>	M 75	F 53	M 13	F 26	M 0	F 0	M 0	F 1	M 0	F 0	M 1	F 1	M 2	F 2	M 0	F 0
020-029	<u>Syphilis & sequelae</u>	<u>4.1</u>	<u>1.9</u>	<u>W 8</u>	M 5	F 3	M 25	F 8	M 0	F 0	M 0	F 0	M 0	F 0	M 0	F 0	M 0	F 0	M 0	F 0
045-048	<u>Dysentery (all forms)</u>	<u>3.3</u>	<u>1.5</u>	<u>C 26</u>	M 7	F 3	M 4	F 3	M 16	F 12	M 10	F 5	M 2	F 0	M 0	F 0	M 1	F 1	M 0	F 0

Table 7 - Page 2
Deaths from Selected Causes - Mississippi - 1958

Table 7 - Page 3
Deaths from Selected Causes - Mississippi - 1958

Inter-national List Number	Cause of Death	Total	Rate*	Total by Race	Total by Sex and Race	Total by Age and Race	Age in Years								
							5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 & Over age	Unknown
210-239	<u>Benign and unspeci-fied neoplasms</u>	68	3.1	W 40 C 28	M 23 F 24	M 27 F 24	1 1/2	1 1/2	1 1/2	1 1/2	1 1/2	1 1/2	1 1/2	1 1/2	1 1/2
240-289	<u>Allergic, endocrine system, metabolic & nutritional diseases</u>	430	19.7	W 216 C 214	M 118 F 150	M 64	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0
260	Diabetes mellitus	317	14.5	W 163 C 154	M 97 F 115	M 66	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0
290-299	<u>Diseases of the blood and blood-forming organs</u>	82	3.8	W 37 C 45	M 27 F 28	M 19	1 1/2	1 1/2	1 1/2	1 1/2	1 1/2	1 1/2	1 1/2	1 1/2	1 1/2
290-293	Anemias	61	2.8	W 25 C 36	M 13 F 13	M 12	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0
300-326	<u>Mental, psycho-neurotic, & personality disorders</u>	26	1.2	W 14 C 14	M 12 F 12	M 9	1 1/2	1 1/2	1 1/2	1 1/2	1 1/2	1 1/2	1 1/2	1 1/2	1 1/2
330-334	<u>Vascular lesions affecting central nervous system</u>	2,867	131.3	W 1,426 C 1,441	M 711 F 758	M 715	683	711	711	711	711	711	711	711	711

Table 7 - Page 4
Deaths from Selected Causes - Mississippi - 1958

Table 7 - Page 4
Deaths from Selected Causes - Mississippi - 1958

International List Number	Cause of Death	Total	Rate*	Total by Race	Total by Sex and Race	Age in Years							
						5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 and over
340-398	Other diseases of the nervous system and sense organs	163	7.5	W 72 C 21	M 43 F 29	5 7 11	1 1 1	1 1 1	2 1 1	2 1 1	2 1 1	14 6 6	14 6 6
	Nonmeningococcal meningitis	72	3.3	W 32 C 40	M 16 F 11	4 7 10	0 1 0	0 0 0	1 0 0	1 0 0	1 0 0	2 3 0	2 3 0
	Diseases of the circulatory system	400-468	330.4	W 4,313 C 2,903	M 2,593 F 1,720	1 3 2	1 2 2	1 3 3	1 3 3	1 3 3	1 3 3	14 25 22	14 25 22
400-402	Rheumatic fever	13	0.6	W 5 C 8	M 3 F 2	0 0	0 0	1 0	1 0	1 0	1 0	0 0 0	0 0 0
	Chronic rheumatic heart disease	410-416	5.9	W 72 C 57	M 47 F 25	0 0	0 0	0 0	0 0	0 0	0 0	1 2 3	1 2 3
420-422	Arteriosclerotic and degenerative heart disease	4,520	207.0	C 1,422	M 1,942 F 1,156	1 0	0 0	0 0	0 0	0 0	0 0	10 12 16	10 12 16
	Other diseases of the heart	430-434	22.2	W 206 C 279	M 132 F 638	0 1	0 1	0 1	0 1	0 1	0 1	2 1 1	2 1 1

Table 7 - Page 5
Deaths from Selected Causes - Mississippi - 1958

50

International List Number	Cause of Death	Total Rate*	Race	Total by Race	Xes	Total by Sex and Race	Age in Years										Unknown Over 65
							1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	
440-443	Hypertension with heart disease	1,381	63.2	W C	548 833	M M	269 377	0 0	0 0	1 1	4 5	11 10	32 25	221 238	0 0	0 0	
444-447	Hypertension without mention of heart	235	10.8	C	97	M	49	0	0	1	1	1 1	3 12	39 13	0 37	0 30	0 0
470-527	Diseases of the respiratory system	1,142	52.3	C	634	M W	296 508	20 24	11 12	1 1	1 1	3 1	2 1	2 2	2 1	2 1	0 0
480-483	Influenza	196	9.0	C	134	M F	70 64	12 17	6 3	0 0	0 0	0 4	0 2	1 7	3 7	3 27	0 0
490-493	Pneumonia (excluding pneumonia of newborn)	713	32.6	C	431	M F	148 134	13 16	6 4	1 1	2 2	3 1	3 2	3 7	4 12	9 19	91 77
500-502	Bronchitis	44	2.0	C	15	M F	18 11	2 1	1 3	0 0	0 0	0 0	0 0	1 1	0 0	0 0	0 0
530-537	Diseases of the digestive system	747	24.2	C	371	M F	228 211	13 82	18 56	1 1	1 1	6 4	2 1	1 1	1 1	1 1	1 1

Table 7 - Page 6
Deaths from Selected Causes - Mississippi - 1958

Table 7 - Page 6
Deaths from Selected Causes - Mississippi - 1958

International List Number	Cause of Death	Total	Rate*	Race	Total by Race	Sex and Race	Total by Sex and Race	Age in Years		Over 65 & Unknown
								Under 1-4	10-14	
540,541	Ulcer of stomach and duodenum	102	4.7	C	29	M 24 F 25	M 18 F 24	1	0	23 0
550,553	Appendicitis	12	0.5	C	10	M 6 F 4	M 6 F 4	0	0	10 0
560,561, 570	Intestinal obstruction and hernia	127	5.8	C	63	M 35 F 31	M 29 F 31	2	1	17 0
571,572	Gastritis, duodenitis, enteritis & colitis, except diarrhea of the newborn	220	10.1	C	168	M 99 F 69	M 73 F 49	6	0	10 0
581	Cirrhosis of liver	95	4.3	C	31	M 46 F 18	M 17 F 14	1	0	10 0
590-637	Diseases of the genito-urinary system	579	26.5	C	360	M 139 F 80	M 123	1	0	10 0
590-594	Nephritis and nephrosis	322	14.7	C	222	M 100 F 136	M 47 F 86	1	0	10 0

Table 7 - Page 7
Deaths from Selected Causes - Mississippi - 1958

International List Number	Cause of Death	Total	Rate*	Race	Total by Race	Sex and Race	Age in Years	Unknown	
								Under 5	65 and over
610	Hyperplasia of prostate	85	3.9	W C	36 M 49 F	M F	0-4	0 0 0 0	34 0
	Complications of pregnancy, childbirth, and the puerperium	53	0.9	W C	10 M 43 F	M F	5-9	3 0 3 0	44 0
620-689*	Diseases of the skin and cellular tissue	10	0.4	W C	4 M 6 F	M F	10-14	2 1 2 0	9 2 5
690-716	Diseases of the bones & organs of movement	53	2.4	W C	37 M 16 F	M F	15-19	12 12 12 12	19 4 5
720-749	Congenital malformations	254	11.6	W C	141 M 113 F	M F	20-24	9 2 9 2	14 2 5
750-759	Certain diseases of early infancy	1,248	21.1	W C	451 M 797 F	M F	25-29	22 22 22 22	21 2 1
760-776*	Symptoms, senility, and ill-defined conditions	1,930	88.4	W C	362 M 1,568 F	M F	30-34	121 71 432 281	121 71 432 281
780-795									

Table 7 - Page 8
Deaths from Selected Causes - Mississippi - 1958

Table 7 - Page 8
Deaths from Selected Causes - Mississippi - 1958

Inter-national List Number	Cause of Death	Total	Rate*	Total by Race	Total by Sex and Race	Age in Years							
						5-9	10-14	15-24	25-34	35-44	45-54	55-64	65-74
800-999	Accidents, poisonings and violence	1,735	79.4	W 901 C 834	M 671 F 230	13	25	18	25	117	113	80	100
						13	17	7	27	26	25	23	16
						20	44	42	49	26	26	22	62
810-835	Motor vehicle accidents	554	25.4	C 173	M 304 F 130	3	11	10	14	79	59	34	25
						77	2	5	5	18	7	9	8
						130	3	4	8	27	18	24	13
						43	0	6	0	13	4	7	4
800-802, 840-962	All other accidents	825	37.8	C 477	M 228 F 120	10	14	8	10	30	36	22	32
						228	11	12	2	4	3	4	7
						120	11	36	19	23	23	26	30
						299	40	42	25	4	6	3	2
963, 970-979	Suicide	149	6.8	C 25	M 106 F 18	0	0	0	1	4	10	17	26
						106	0	0	0	1	1	1	5
						18	0	0	0	0	0	0	4
						21	0	0	0	0	0	0	2
						4	0	0	0	0	0	0	1
													0
980-985	Homicide	207	9.5	C 159	M 48 F 48	0	0	0	0	4	3	7	8
						48	0	0	0	1	31	35	23
						159	1	2	0	1	0	7	3
													2
													0

* Crude death rate per 1,000 population. Specific death rates per 100,000 population. Maternal death rates and rates for infant causes (760-776) per 1,000 live births.

TABLE 8
DEATHS UNDER 1 YEAR
MISSISSIPPI
1958

For Selected Causes, by Race, by Age, on Basis of Place of Residence, with Rates

Cause of Death and International List Number	Total	Rate*	Total by Race	DAYS						MONTHS									
				Under 1 day	1- 6	7- 13	14- 20	21- 27	Under 28 days	1	2	3	4	5	6	7	8		
ALL CAUSES	2,424	41.0	W C	<u>206</u> <u>1,718</u>	<u>293</u> <u>400</u>	<u>196</u> <u>324</u>	<u>29</u> <u>26</u>	<u>13</u> <u>22</u>	<u>9</u> <u>43</u>	<u>540</u> <u>935</u>	<u>36</u> <u>178</u>	<u>23</u> <u>128</u>	<u>26</u> <u>28</u>	<u>19</u> <u>23</u>	<u>12</u> <u>82</u>	<u>2</u> <u>54</u>	<u>7</u> <u>41</u>	<u>8</u> <u>18</u>	<u>2</u> <u>25</u>
Syphilis (020-029)	3	0.05	W C	0 3	0 0	0 1	0 0	0 0	0 0	0 0	0 0	0 1	0 2	0 0	0 0	0 0	0 0	0 0	
Diphtheria (055)	0	-	W C	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	
Whooping cough (056)	10	0.2	W C	0 10	0 0	0 0	0 0	0 0	0 1	0 1	0 0	0 0	0 2	0 3	0 1	0 0	0 0	0 0	
Measles (085)	8	0.1	W C	0 8	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 1	0 0	0 0	0 1	0 0	0 0	
Influenza (480-483)	31	0.5	W C	2 29	0 1	0 1	0 1	0 1	0 1	0 0	0 0	0 0	0 5	0 2	0 1	0 0	0 1	0 0	
**Pneumonia (490-493)	190	3.2	W C	29 161	0 0	0 1	0 0	0 0	0 0	0 1	29 27	4 14	5 27	2 14	1 21	3 11	4 11	8 6	2 3
Other respiratory diseases (470-475, 500-527)	30	0.5	W C	13 17	0 0	2 1	0 1	0 1	0 1	0 0	2 1	5 2	2 2	1 2	3 0	0 0	1 0	0 1	
**Diarrhea and enteritis (5710)	131	2.2	W C	11 120	0 0	0 0	0 0	0 0	0 0	0 0	23 17	1 17	2 17	1 15	1 15	6 10	5 7	4 2	

Table 8 - Page 2
Deaths Under 1 Year - Mississippi - 1958

Cause of Death and International List Number	Total	Rate*	Total by Race	DAYS												MONTHS						
				Under 1 day	1- 6	7- 13	14- 20	21- 27	28 days	Under 1 day	1	2	3	4	5	6	7	8	9	10	11	
Congenital malfor- mations (750-759)	191	3.2	W C	98 93	27	7	1	5	69	4	3	2	1	1	1	2	1	1	1	1	1	
Birth injuries (760-761)	128	2.2	W C	62 66	42	20	0	0	62	0	0	0	0	0	0	0	0	0	0	0	0	
Postnatal asphyxia & atelectasis (762)	262	4.4	W C	133 129	70	58	3	0	0	131	1	0	0	0	0	1	0	0	0	0	0	
**Infections of the newborn (763-768)	120	2.0	W C	40 80	4	17	14	3	40	0	0	0	0	0	0	0	0	0	0	0	0	
Hemolytic disease of the newborn (770)	28	0.5	W C	14 14	5	8	1	0	0	14	0	0	0	0	0	0	0	0	0	0	0	
All other defined dis. of early infancy (769,771,772)	70	1.2	W C	16 54	5	9	0	1	15	0	1	0	0	0	0	0	0	0	0	0	0	
Violent & accidental deaths (800-999)	100	1.7	W C	26 74	1	0	0	2	32	7	5	1	2	3	3	2	3	0	3	0	1	
Other specified causes	157	2.6	W C	49 108	6	1	2	5	0	14	10	6	4	2	2	1	3	3	1	2	5	
Ill-defined (773-776, 780-793,795)	965	16.3	W C	213 752	133	52	2	2	191	8	4	4	4	0	0	1	0	0	1	0	1	

* Rate per 1,000 live births.

** Pneumonia and diarrhea of the newborn (under 28 days) included in infections of the newborn (763-768).

TABLE 9
DEATHS FROM SELECTED CAUSES
MISSISSIPPI
1958

By Month of Occurrence on Basis of Place of Residence

Cause of Death and International List Number	Total	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
TOTAL ALL CAUSES	21,541	2,220	2,160	1,991	1,681	1,678	1,623	1,626	1,554	1,500	1,728	1,716	2,064
Dysentery (045-048)	33	3	4	1	2	0	3	6	5	2	2	3	2
Whooping cough (056)	12	0	0	0	0	3	0	4	0	2	1	1	1
Acute poliomyelitis (080)	1	0	0	0	0	0	0	0	0	1	0	0	0
Measles (085)	15	1	0	4	5	2	0	1	0	0	1	0	1
Influenza (480-483)	196	51	68	34	13	0	2	2	3	3	4	4	12
Lobar pneumonia (490)	158	25	33	25	13	10	2	2	8	4	7	8	21
Bronchopneumonia (491)	302	40	54	35	27	15	18	9	11	15	25	25	28
Other and unspecified pneumonia (492-493)	253	37	51	24	30	12	15	9	11	14	16	14	20
Diarrhea & enteritis (-2) (5710)	152	33	10	5	6	7	7	9	12	14	17	15	17
Diarrhea & enteritis (+2) (5711)	46	2	6	2	3	1	5	4	8	5	4	5	1
Accidents (800-962)	1,379	121	118	108	106	100	117	116	100	96	127	124	146
Motor vehicle accidents (810-835)	1,554	32	31	38	44	35	46	47	51	51	69	55	55

TABLE 10
MARRIAGES AND DIVORCES
MISSISSIPPI
1958

By County, by Race of Bride

County	Marriages						Divorces		
	Place of Licensure			Residence of Bride			Total	White	Non-white
	Total	White	Non-white	Total	White	Non-white			
<u>STATE</u>	<u>36,402</u>	<u>26,998</u>	<u>9,404</u>	<u>17,370</u>	<u>10,875</u>	<u>6,495</u>	<u>5,016</u>	<u>3,485</u>	<u>1,531</u>
Adams	525	367	158	308	184	124	166	107	59
Alcorn	3,682	3,413	269	214	190	24	88	72	16
Amite	216	162	54	114	63	51	29	13	16
Attala	124	84	40	140	95	45	47	32	15
Benton	107	39	68	61	35	26	7	3	4
Bolivar	440	184	256	420	173	247	90	52	38
Calhoun	116	93	23	126	99	27	30	22	8
Carroll	76	50	26	57	26	31	23	11	12
Chickasaw	175	117	58	147	96	51	26	17	9
Choctaw	79	50	29	64	34	30	4	2	2
Claiborne	108	54	54	68	24	44	33	9	24
Clarke	259	200	59	149	99	50	31	25	6
Clay	194	132	62	130	72	58	37	23	14
Coahoma	466	197	269	321	109	212	70	36	34
Copiah	217	119	98	208	115	93	59	27	32
Covington	128	92	36	116	82	34	23	19	4
DeSoto	3,921	2,701	1,220	169	73	96	35	22	13
Forrest	397	313	84	428	323	105	207	151	56
Franklin	77	58	19	60	37	23	29	19	10
George	891	838	53	106	98	8	13	11	2
Greene	276	252	24	66	55	11	17	13	4
Grenada	138	82	56	123	70	53	39	16	23
Hancock	681	632	49	90	79	11	48	45	3
Harrison	1,477	1,308	169	947	803	144	619	530	89
Hinds	1,462	916	546	1,601	1,037	564	424	313	111
Holmes	226	71	155	262	80	182	35	15	20
Humphreys	148	54	94	155	56	99	23	9	14
Issaquena	25	13	12	27	6	21	6	3	3
Itawamba	710	700	10	89	88	1	24	24	0
Jackson	1,485	1,108	377	507	396	111	137	111	26
Jasper	118	78	40	159	97	62	38	20	18
Jefferson	206	160	46	67	10	57	16	6	10
Jeff Davis	156	88	68	97	51	46	21	14	7
Jones	571	442	129	549	433	116	184	142	42
Kemper	74	40	34	78	40	38	16	8	8
Lafayette	131	86	45	136	83	53	13	7	6

County	Marriages						Divorces		
	Place of Licensure			Residence of Bride					
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Lamar	212	192	20	128	116	12	32	28	4
Lauderdale	1,109	754	355	634	462	172	170	137	33
Lawrence	73	48	25	115	72	43	26	18	8
Leake	170	117	53	156	105	51	36	27	9
Lee	389	319	70	344	273	71	62	52	10
Leflore	387	149	238	375	144	231	73	41	32
Lincoln	253	198	55	221	177	44	43	34	9
Lowndes	1,555	1,259	296	329	213	116	103	70	33
Madison	293	118	175	233	80	153	77	24	53
Marion	292	183	109	217	143	74	59	43	16
Marshall	564	388	176	167	54	113	34	15	19
Monroe	317	229	88	251	165	86	60	46	14
Montgomery	105	55	50	113	64	49	28	18	10
Neshoba	213	153	60	179	129	50	63	48	15
Newton	176	118	58	175	113	62	45	27	18
Noxubee	158	49	109	108	30	78	24	7	17
Oktibbeha	251	178	73	137	70	67	24	15	9
Panola	254	140	114	244	120	124	46	25	21
Pearl River	568	465	103	234	192	42	111	97	14
Perry	84	66	18	72	60	12	22	19	3
Pike	384	236	148	307	175	132	116	75	41
Pontotoc	154	136	18	126	108	18	26	20	6
Prentiss	317	298	19	155	141	14	36	35	1
Quitman	208	104	104	156	68	88	31	16	15
Rankin	390	279	111	203	120	83	50	38	12
Scott	239	168	71	213	136	77	57	39	18
Sharkey	98	33	65	79	37	42	18	8	10
Simpson	159	123	36	156	111	45	53	39	14
Smith	93	85	8	100	83	17	35	33	2
Stone	105	85	20	74	56	18	9	7	2
Sunflower	387	204	183	309	102	207	38	25	13
Tallahatchie*	183	53	130	139	56	83	37	17	20
Tate	556	475	81	131	57	74	34	16	18
Tippah	189	165	24	150	127	23	25	19	6
Tishomingo	1,380	1,282	98	140	139	1	27	25	2
Tunica	286	124	162	153	32	121	18	9	9
Union	210	176	34	154	123	31	42	30	12
Walthall	238	164	74	104	61	43	18	10	8
Warren	487	329	158	345	211	134	194	75	119
Washington	1,586	1,170	416	597	284	313	107	98	9
Wayne	275	184	91	138	98	40	38	25	13
Webster	69	57	12	70	57	13	12	12	0
Wilkinson	386	300	86	88	24	64	32	7	25
Winston	161	90	71	157	88	69	46	21	25
Yalobusha	123	80	43	112	61	51	19	16	3
Yazoo	234	127	107	223	127	96	53	30	23
Unstated				112	48	64			
Out-of-state residents				18,920	16,075	2,845			

* Records received from only one district, July-December, 1958.

TABLE 11
DIVORCES
MISSISSIPPI
1958

By Cause of Divorce

Cause	Total	White	Non-white
<u>TOTAL</u>	<u>5,016</u>	<u>3,485</u>	<u>1,531</u>
Cruel and inhuman treatment	2,420	2,032	388
Desertion	1,941	960	981
Adultery	241	146	95
Drunkenness	269	230	39
Penitentiary	43	36	7
Insanity	10	9	1
Annulment	32	21	11
Bigamy	20	18	2
Miscellaneous and other	15	11	4
<u>Not stated</u>	<u>25</u>	<u>22</u>	<u>3</u>

By Number of Minor Children

Minor Children	Total	White	Non-white
None	2,623	1,686	937
One	983	781	202
Two	659	515	144
Three or more	640	435	205
<u>Not stated</u>	<u>111</u>	<u>68</u>	<u>43</u>

By Number of Years Married

Number Years Married	Total	White	Non-white
<u>TOTAL</u>	<u>5,016</u>	<u>3,485</u>	<u>1,531</u>
Under one year	386	339	47
One year	609	548	61
Two years	450	374	76
Three years	384	305	79
Four years	292	229	63
5-9 years	956	649	307
10-19 years	1,061	675	386
Twenty years and over	778	298	480
<u>Not stated</u>	<u>100</u>	<u>68</u>	<u>32</u>

Table 12--Live Births, by Year, by Race, on Basis of Place of Occurrence
with Rates: Mississippi, 1913-1958

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1912**	6,311					
1913	41,103	21,874	19,229	21.9	26.6	18.2
1914	40,623	21,360	19,263	21.4	25.7	18.0
1915	39,484	20,999	18,485	20.5	23.8	17.1
1916	39,033	20,130	18,903	20.0	23.6	17.2
1917	37,141	19,056	18,085	18.8	22.0	16.3
1918	39,391	20,005	19,386	19.7	22.8	17.2
1919	42,491	22,028	20,463	21.0	24.9	18.0
1920	43,743	22,438	21,305	24.4	26.3	22.7
1921	46,482	23,703	22,779	25.6	27.3	24.1
1922	43,723	21,286	22,437	23.4	24.1	23.6
1923	43,266	21,140	22,126	23.3	23.6	23.1
1924	43,960	22,180	21,780	23.4	24.3	22.5
1925	45,563	22,265	23,298	24.0	24.1	23.9
1926	50,449	24,600	25,849	26.2	26.2	26.3
1927	49,204	24,163	25,041	25.3	25.3	25.3
1928	48,126	23,642	24,484	24.5	24.4	24.5
1929	45,570	22,105	23,465	22.9	22.5	23.3
1930	47,982	23,192	24,790	23.8	23.2	24.0
1931	45,123	21,637	23,486	22.2	21.4	22.9
1932	46,481	21,883	24,598	22.6	21.4	23.9
1933	44,184	20,407	23,777	21.3	19.7	22.9
1934	47,905	22,342	25,563	23.0	21.4	24.6
1935	48,291	21,972	26,319	23.0	20.8	25.1
1936	49,261	22,194	27,067	23.3	20.8	25.7
1937	52,170	23,285	28,885	24.4	21.6	27.3
1938	53,729	24,046	29,683	25.0	22.1	27.8
1939	52,288	23,505	28,783	24.1	21.4	26.8
1940	52,526	23,084	29,442	24.0	20.8	27.3
1941	54,561	24,167	30,394	24.7	21.6	28.0
1942	56,521	25,306	31,215	25.4	22.4	28.6
1943	59,408	27,611	31,797	28.9	26.4	31.6
1944#	56,586	25,647	30,939	28.3	25.2	31.6
1945#	53,540	23,426	30,114	26.9	23.1	30.9
1946#	61,430	29,715	31,715	26.8	25.4	28.2
1947#	66,291	32,423	33,868	28.6	27.5	29.9
1948#	65,291	29,666	35,625	28.0	24.9	31.2
1949#	66,128	28,786	37,342	28.1	24.0	32.5
1950#	64,427	27,188	37,239	29.6	22.9	37.6
1951#	65,512	27,798	37,714	30.1	23.4	38.1
1952#	64,509	28,073	36,436	29.6	23.6	36.8
1953#	64,061	28,312	35,749	29.4	23.8	36.1

Table 12 (continued)

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1954#	64,748	28,781	35,967	29.7	24.2	36.3
1955#	63,244	27,546	35,698	29.0	23.2	36.0
1956#	62,178	27,278	34,900	29.5	23.8	36.2
1957#	60,712	26,804	33,908	28.8	23.4	35.2
1958#	59,189	26,812	32,377	27.1	20.9	35.9

* Rate per 1,000 population. 1913-1920 population estimates based on U. S. Censuses of 1910 & 1920; 1921-1930 population estimates based on U. S. Censuses of 1920 & 1930; 1931-1940 population estimates based on U. S. Censuses of 1930 & 1940; 1941 & 1942 projected estimates based on U. S. Censuses of 1930 & 1940; 1943 & 1944 estimates of civilian population obtained from Ration Book Registration; 1945 estimate of civilian population, July 1, 1945, obtained from Bureau of Census Bulletin Series P-46, No. 3, dated February 12, 1946; 1946-1949 projected estimates as of July 1, each year based on U. S. Censuses of 1930 & 1940; 1950-1955 U. S. Census population of April 1, 1950; estimate for 1956-1957 as of July 1, 1956, prepared by Dr. Harald A. Pedersen, Sociology Department, Mississippi State University, with adjustments by Statistical Service, Mississippi State Board of Health; 1958 estimate as of January 1, prepared by Dr. George L. Wilber and Mr. Carl T. Robbins, Sociology Department, Mississippi State University.

** November-December, 1912.

On basis of place of residence.

Table 13-Deaths, by Year, by Race, on Basis of Place of Occurrence, with
Rates: Mississippi, 1913-1958

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1912**	4,341					
1913	21,305	7,643	13,662	11.4	9.3	13.0
1914	21,485	7,442	14,043	11.3	8.9	13.1
1915	24,001	8,088	15,913	12.5	9.6	14.7
1916	21,794	7,588	14,206	11.2	8.9	13.0
1917	23,579	8,048	15,531	11.9	9.3	14.0
1918	30,437	10,177	20,260	15.2	11.6	18.0
1919	22,667	8,142	14,525	12.7	9.5	15.5
1920	22,072	7,916	14,156	12.3	9.3	15.1
1921	19,991	7,308	12,683	11.0	8.4	13.4
1922	19,278	7,312	11,966	10.5	8.3	12.6
1923	20,363	7,688	12,675	11.0	8.6	13.2
1924	21,155	7,809	13,346	11.3	8.6	13.8
1925	22,229	8,143	14,086	11.7	8.8	14.4
1926	23,929	9,133	14,796	12.5	9.7	15.1
1927	23,257	8,765	14,492	12.0	9.2	14.6
1928	25,965	10,111	15,854	13.2	10.4	15.9
1929	25,868	10,048	15,820	13.0	10.2	15.7
1930	24,098	9,072	15,026	11.9	9.0	14.8
1931	22,007	8,641	13,366	10.8	8.5	13.0
1932	20,322	8,295	12,027	9.9	8.1	11.6
1933	21,617	8,898	12,719	10.4	8.6	12.2
1934	21,786	9,350	12,436	10.5	9.0	12.0
1935	21,285	9,186	12,099	10.1	8.7	11.6
1936	24,127	10,367	13,760	11.4	9.7	13.1
1937	24,005	10,055	13,950	11.2	9.3	13.2
1938	22,814	9,471	13,343	10.6	8.7	12.5
1939	22,701	9,464	13,237	10.5	8.6	12.3
1940	23,185	9,565	13,620	10.6	8.6	12.6
1941	22,573	9,442	13,131	10.2	8.4	12.1
1942	20,693	8,750	11,943	9.3	7.7	10.9
1943	20,850	9,139	11,711	10.2	8.7	11.6
1944#	20,199	8,830	11,369	10.1	8.7	11.6
1945#	19,024	8,452	10,572	9.6	8.3	10.8
1946#	19,321	8,995	10,326	8.4	7.7	9.2
1947#	20,007	9,238	10,769	8.6	7.8	9.5
1948#	20,183	9,216	10,967	8.6	7.7	9.6
1949#	20,489	9,233	11,256	8.7	7.7	9.8
1950#	20,567	9,466	11,101	9.4	8.0	11.2
1951#	21,007	9,593	11,414	9.6	8.1	11.5
1952#	20,657	9,387	11,270	9.5	7.9	11.4

Table 13 (continued)

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1953#	20,492	9,495	10,997	9.4	8.0	11.1
1954#	19,353	9,453	9,900	8.9	8.0	10.0
1955#	19,403	9,501	9,902	8.9	8.0	10.0
1956#	19,780	9,969	9,811	9.4	8.7	10.2
1957#	20,476	10,339	10,137	9.7	9.0	10.5
1958#	21,541	10,811	10,730	9.9	8.4	11.9

* Rate per 1,000 population. See footnote on Table 13 for source of population estimates.

** November-December, 1912.

On basis of place of residence.

Table 14--Maternal Deaths, by Year, by Race, on Basis of Place of Occurrence, with Rates: Mississippi, 1913-1958

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1913	447	162	285	10.9	7.4	14.8
1914	413	124	289	10.2	5.8	15.0
1915	289	88	201	7.3	4.2	10.9
1916	326	102	224	8.4	5.1	11.8
1917	297	105	192	8.0	5.5	10.6
1918	302	82	220	7.7	4.1	11.3
1919	407	173	234	9.6	7.8	11.4
1920	393	127	266	9.0	5.7	12.5
1921	391	145	246	8.4	6.1	10.8
1922	313	110	203	7.2	5.2	9.0
1923	321	108	213	7.4	5.1	9.6
1924	413	137	276	9.4	6.2	12.7
1925	435	143	292	9.5	6.4	12.5
1926	388	153	235	7.7	6.2	9.1
1927	409	134	275	8.3	5.5	11.0
1928	421	161	260	8.7	6.8	10.6
1929	410	152	258	9.0	6.9	11.0
1930	489	171	318	10.2	7.4	12.8
1931	352	131	221	7.8	6.1	9.4
1932	279	103	176	6.0	4.7	7.2
1933	315	116	199	7.1	5.7	8.4
1934	306	116	190	6.4	5.2	7.4
1935	334	137	197	6.9	6.2	7.5
1936	358	134	224	7.3	6.0	8.3
1937	370	150	220	7.1	6.4	7.6
1938	333	116	217	6.2	4.8	7.3
1939	311	97	214	5.9	4.1	7.4
1940	327	106	221	6.2	4.6	7.5
1941	293	93	200	5.4	3.8	6.6
1942	248	77	171	4.4	3.0	5.5
1943	233	66	167	3.9	2.4	5.3
1944**	225	72	153	4.0	2.8	4.9
1945**	216	64	152	4.0	2.7	5.0
1946**	199	67	132	3.2	2.2	4.2
1947**	175	50	125	2.6	1.5	3.7
1948**	177	38	139	2.7	1.3	3.9
1949**	147	32	115	2.2	1.1	3.1
1950**	165	30	135	2.6	1.1	3.6
1951**	143	28	115	2.2	1.0	3.0
1952**	119	15	104	1.8	0.5	2.8
1953**	119	24	95	1.8	0.8	2.6
1954**	104	11	93	1.6	0.4	2.6
1955**	76	12	64	1.2	0.4	1.8
1956**	74	12	62	1.2	0.4	1.8
1957**	75	14	61	1.2	0.5	1.8
1958**	53	10	43	0.9	0.4	1.3

* Rate per 1,000 live births.

** On basis of place of residence.

Table 15--Deaths Under 1 Year, by Year, by Race, on Basis of Place of Occurrence, with Rates: Mississippi, 1917-1958

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1917	3,689	1,296	2,393	99.3	68.0	132.3
1918						
1919	3,554			83.6		
1920	3,499	1,344	2,155	80.0	59.9	101.1
1921	3,157	1,243	1,914	67.9	52.4	84.0
1922	2,826	1,067	1,759	64.6	50.1	78.4
1923	2,908	1,104	1,804	67.2	52.2	81.5
1924	3,137	1,188	1,949	71.4	53.6	89.5
1925	3,176	1,179	1,997	69.7	53.0	85.7
1926	3,578	1,447	2,131	70.9	58.8	82.4
1927	3,291	1,303	1,988	66.9	54.0	79.4
1928	3,567	1,416	2,151	74.1	59.9	87.9
1929	3,275	1,268	2,007	71.9	57.4	85.5
1930	3,276	1,182	2,094	68.3	51.0	84.5
1931	2,547	961	1,586	56.4	44.4	67.5
1932	2,482	965	1,517	53.4	44.1	61.7
1933	2,800	1,035	1,765	63.4	50.7	74.2
1934	3,070	1,199	1,871	64.1	53.7	73.2
1935	2,587	1,044	1,543	53.6	47.5	58.6
1936	2,883	1,105	1,778	58.5	49.8	65.7
1937	3,098	1,174	1,924	59.4	50.4	66.6
1938	3,051	1,205	1,846	56.8	50.1	62.2
1939	2,908	1,135	1,773	55.6	48.3	61.6
1940	2,864	1,070	1,794	54.5	46.4	60.9
1941	2,969	1,113	1,856	54.4	46.1	61.1
1942	2,654	957	1,697	47.0	37.8	54.4
1943	2,774	1,137	1,637	46.7	41.2	51.5
1944**	2,489	974	1,515	44.0	38.0	49.0
1945**	2,192	846	1,346	40.9	36.1	44.7
1946**	2,298	1,005	1,293	37.4	33.8	40.8
1947**	2,428	1,009	1,419	36.6	31.1	41.9
1948**	2,470	988	1,482	37.8	33.3	41.6
1949**	2,612	900	1,712	39.5	31.3	45.8
1950**	2,348	763	1,585	36.4	28.1	42.6
1951**	2,589	814	1,775	39.5	29.3	47.1
1952**	2,592	760	1,832	40.2	27.1	50.3
1953**	2,534	807	1,727	39.6	28.5	48.3
1954**	2,360	744	1,616	36.4	25.8	44.9
1955**	2,338	680	1,658	37.0	24.7	46.4
1956**	2,353	734	1,619	37.8	26.9	46.4
1957**	2,257	694	1,563	37.2	25.9	46.1
1958**	2,424	706	1,718	41.0	26.3	53.1

* Rate per 1,000 live births.

** On basis of place of residence.

Table 16--Deaths Under 1 Month, by Year, by Race, on Basis of Place of Occurrence, with Rates: Mississippi, 1924-1958

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1924	1,598	730	868	36.4	32.9	39.8
1925	1,639	701	938	36.0	31.5	40.3
1926	1,731	823	908	34.3	33.4	35.1
1927	1,644	786	858	33.4	32.5	34.3
1928	1,788	804	984	37.2	34.0	40.2
1929	1,662	773	889	36.5	35.0	37.9
1930	1,692	715	977	35.3	30.8	39.4
1931	1,339	590	749	29.7	27.3	31.9
1932	1,361	572	789	29.3	26.1	32.1
1933	1,540	636	904	34.8	31.2	38.0
1934	1,505	644	861	31.4	28.8	33.7
1935	1,446	642	804	29.9	29.2	30.5
1936	1,494	676	818	30.3	30.4	30.2
1937	1,678	755	923	32.2	32.4	32.0
1938	1,688	747	941	31.4	31.1	31.7
1939	1,637	735	902	31.3	31.3	31.3
1940	1,613	680	933	30.7	29.4	31.7
1941	1,663	734	929	30.5	30.4	30.6
1942	1,544	638	906	27.3	25.2	29.0
1943	1,652	791	861	27.8	28.6	27.1
1944	1,436	661	775	25.2	25.4	25.1
1945**	1,351	624	727	25.2	26.6	24.1
1946**	1,561	794	767	25.4	26.7	24.2
1947**	1,595	750	845	24.1	23.1	24.9
1948**	1,674	773	901	25.6	26.0	25.3
1949**	1,594	661	933	24.1	23.0	25.0
1950**	1,446	560	886	22.4	20.6	23.8
1951**	1,679	626	1,053	25.6	22.5	27.9
1952**	1,658	581	1,077	25.7	20.7	29.6
1953**	1,630	635	995	25.4	22.4	27.8
1954**	1,521	601	920	23.5	20.9	25.6
1955**	1,440	527	913	22.8	19.1	25.6
1956**	1,451	581	870	23.3	21.3	24.9
1957**	1,425	540	885	23.5	20.1	26.1
1958**	1,475	540	935	24.9	20.1	28.9

* Rate per 1,000 live births.

** On basis of place of residence.

TABLE 17
DEATHS FROM HEART DISEASES, NEPHRITIS (ALL FORMS),
VASCULAR LESIONS, CANCER, AND TUBERCULOSIS (ALL FORMS)
MISSISSIPPI
1949-1958

Shown by Year and Race, on Basis of Place of Residence, with Rates*

YEAR	HEART DISEASES				NEPHRITIS (ALL FORMS)				VASCULAR LESIONS				CANCER				TUBERCULOSIS (ALL FORMS)			
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white		
1949 No.	5,496	3,060	2,436	729	236	493	2,093	976	1,117	2,039	1,205	834	581	190	391					
Rate	233.9	254.9	212.0	31.0	19.7	42.9	89.1	81.3	97.2	86.8	100.4	72.6	24.7	15.8	34.0					
1950 No.	5,546	3,158	2,388	733	246	487	2,200	971	1,229	2,143	1,333	810	558	179	379					
Rate	254.5	265.7	241.1	33.6	20.7	49.2	101.0	81.7	124.1	98.4	112.2	81.8	25.6	15.1	38.3					
1951 No.	5,626	3,134	2,492	681	210	471	2,344	1,089	1,255	2,186	1,326	860	497	158	339					
Rate	258.2	263.7	251.6	31.2	17.7	47.6	107.6	91.6	126.7	100.3	111.6	86.8	22.8	13.3	34.2					
1952 No.	5,876	3,256	2,620	658	214	444	2,386	1,098	1,288	2,189	1,285	904	418	133	285					
Rate	269.7	273.9	264.6	30.2	18.0	44.8	109.5	92.4	130.1	100.5	108.1	91.3	19.2	11.2	28.8					
1953 No.	5,758	3,183	2,575	474	164	310	2,404	1,125	1,279	2,230	1,348	882	308	85	223					
Rate	264.3	267.8	260.0	21.8	13.8	31.3	110.3	94.6	129.2	102.3	113.4	89.1	14.1	7.2	22.5					
1954 No.	5,708	3,325	2,383	448	169	279	2,422	1,176	1,246	2,332	1,420	912	226	80	146					
Rate	262.0	279.7	240.6	20.6	14.2	28.2	111.2	98.9	125.8	107.0	119.5	92.1	10.4	6.7	14.7					
1955 No.	5,969	3,517	2,452	401	144	257	2,447	1,169	1,278	2,317	1,402	915	211	66	145					
Rate	273.9	295.9	247.6	18.4	12.1	26.0	112.3	98.3	129.0	106.3	118.0	92.4	9.7	5.6	14.6					
1956 No.	6,104	3,703	2,401	410	161	249	2,550	1,229	1,321	2,432	1,439	993	199	80	119					
Rate	289.3	322.8	249.4	19.4	14.0	25.9	120.9	107.1	137.2	115.3	125.4	103.2	9.4	7.0	12.4					
1957 No.	6,367	3,706	2,661	362	143	219	2,800	1,420	1,380	2,496	1,482	1,014	161	61	100					
Rate	301.8	323.0	276.4	17.1	12.5	22.7	132.7	123.8	143.9	118.3	129.2	105.3	7.6	5.3	10.4					
1958 No.	6,515	3,924	2,591	322	100	222	2,867	1,426	1,441	2,525	1,524	1,001	172	82	90					
Rate	298.3	305.9	287.5	14.7	7.8	24.6	131.3	111.2	159.9	115.6	118.8	111.1	7.9	6.4	10.0					


* Rate per 100,000 population.

TABLE 18
BIRTHS, DEATHS, MARRIAGES
MISSISSIPPI
1958

By Month of Occurrence

Month	Births	Deaths	Marriages
<u>TOTAL</u>	<u>59,189</u>	<u>21,541</u>	<u>36,402</u>
January	5,160	2,220	3,808
February	4,454	2,160	3,482
March	4,733	1,991	4,179
April	4,230	1,681	4,005
May	4,307	1,678	5,005
June	4,491	1,623	6,163
July	5,365	1,626	1,256
August	5,544	1,554	1,887
September	5,460	1,500	1,458
October	5,211	1,728	1,494
November	5,005	1,716	1,641
<u>December</u>	<u>5,229</u>	<u>2,064</u>	<u>2,024</u>

COUNTY SUMMARIES



BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ADAMS COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,703	704	999	1,223	549	674	37.1	27.4	51.8
Fetal deaths	59	14	45	37	8	29	30.2	14.6	43.0
Deaths from all causes	435	146	289	385	132	253	11.7	6.6	19.5
Deaths under 1 year	58	14	44	43	10	33	35.2	18.2	49.0
Deaths under 1 month	33	8	25	22	6	16	18.0	10.9	23.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	1	3	5	-	4	15.2	5.0	30.8
Tbc., respiratory (001-008)	4	1	3	5	-	4	15.2	5.0	30.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	1	0	1	1	0	1	3.0	-	7.7
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	3.0	-	7.7
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	36	14	22	35	14	21	106.1	70.0	161.5
Diabetes (260)	5	2	3	4	3	1	12.1	15.0	7.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	65	16	49	55	14	41	166.7	70.0	315.4
Heart diseases (410-443)	147	54	93	139	50	89	421.2	250.0	684.6
Arteriosclerosis (450)	10	6	4	9	5	4	27.3	25.0	30.8
Influenza (480-483)	5	0	5	5	0	5	15.2	-	38.5
#Pneumonia (490-493)	15	3	12	14	3	11	42.4	15.0	84.6
#Diarrhea & enteritis (571)	7	0	7	6	0	6	18.2	-	46.2
Acute nephritis (590, 591)	0	0	0	1	1	0	3.0	5.0	-
Chronic nephritis, etc. (592-594)	2	0	2	1	0	1	3.0	-	7.7
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	9	4	5	10	5	5	30.3	25.0	38.5
Birth inj., atelect. (760-762)	5	2	3	5	2	3	4.1	3.6	4.4
#Other dis. early inf. (763-776)	25	5	20	13	2	11	10.6	3.6	16.3
Violent & acc. deaths (800-999)	34	18	16	27	13	14	81.8	65.0	107.7
Acc. deaths, total (800-962)	25	13	12	22	10	12	66.7	50.0	92.3
Motor vehicle acc. (810-835)	6	6	0	4	4	0	12.1	20.0	-
Ill-defined (780-795)	13	1	12	13	1	12	39.4	5.0	92.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ALCORN COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	599	481	118	513	403	110	19.0	19.2	18.3
Fetal deaths	23	17	6	16	10	6	31.2	24.8	54.5
Deaths from all causes	262	224	38	261	222	39	9.7	10.6	6.5
Deaths under 1 year	21	17	4	17	13	4	33.1	32.2	36.3
Deaths under 1 month	18	15	3	14	11	3	27.3	27.3	27.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	3.7	4.8	-
Tbc., respiratory (001-008)	0	0	0	1	1	-	3.7	4.8	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	42	37	5	43	37	6	159.2	176.2	100.0
Diabetes (260)	6	5	1	7	6	1	25.9	28.6	16.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	22	15	7	23	16	7	85.2	76.2	116.7
Heart diseases (410-443)	102	91	11	100	90	10	370.4	428.6	166.7
Arteriosclerosis (450)	3	3	0	4	4	0	14.8	19.0	-
Influenza (480-483)	5	2	3	5	2	3	18.5	9.5	50.0
#Pneumonia (490-493)	7	7	0	11	11	0	40.7	52.4	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	3	1	3	2	1	11.1	9.5	16.7
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	1	1	0	3.7	4.8	-
Birth inj., atelect. (760-762)	7	7	0	7	7	0	13.6	17.4	-
#Other dis. early inf. (763-776)	10	7	3	7	4	3	13.6	9.9	27.3
Violent & acc. deaths (800-999)	18	17	1	20	19	1	74.1	90.5	16.7
Acc. deaths, total (800-962)	13	12	1	15	14	1	55.6	66.7	16.7
Motor vehicle acc. (810-835)	3	3	0	7	7	0	25.9	33.3	-
Ill-defined (780-795)	9	8	1	9	8	1	33.3	38.1	16.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
AMITE COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	257	41	216	384	127	257	19.2	15.7	21.6
Fetal deaths	7	0	7	12	1	11	31.2	7.9	42.8
Deaths from all causes	83	38	45	134	78	56	6.7	9.6	4.7
Deaths under 1 year	7	0	7	10	2	8	26.0	15.7	31.1
Deaths under 1 month	4	0	4	6	2	4	15.6	15.7	15.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	5.0	-	8.4
Tbc., respiratory (001-008)	1	0	1	1	0	1	5.0	-	8.4
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	8	4	4	12	8	4	60.0	98.8	33.6
Diabetes (260)	1	0	1	1	0	1	5.0	-	8.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	19	6	13	31	18	13	155.0	222.2	109.2
Heart diseases (410-443)	30	20	10	43	29	14	215.0	358.0	117.6
Arteriosclerosis (450)	2	2	0	2	2	0	10.0	24.7	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	0	2	3	1	2	15.0	12.3	16.8
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	2	2	0	10.0	24.7	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	2	0	2	4	1	3	10.4	7.9	11.7
Violent & acc. deaths (800-999)	5	3	2	10	6	4	50.0	74.1	33.6
Acc. deaths, total (800-962)	3	1	2	8	4	4	40.0	49.4	33.6
Motor vehicle acc. (810-835)	0	0	0	2	1	1	10.0	12.3	8.4
Ill-defined (780-795)	6	1	5	7	2	5	35.0	24.7	42.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ATTALA COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	527	228	299	490	201	289	24.5	17.5	34.0
Fetal deaths	12	1	11	11	1	10	22.4	5.0	34.6
Deaths from all causes	227	126	101	249	143	106	12.4	12.4	12.5
Deaths under 1 year	26	4	22	28	7	21	57.1	34.8	72.7
Deaths under 1 month	13	4	9	15	6	9	30.6	29.8	31.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	2	1	1	10.0	8.7	11.8
Tbc., respiratory (001-008)	2	1	1	2	1	1	10.0	8.7	11.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	25	18	7	29	21	8	145.0	182.6	94.1
Diabetes (260)	5	2	3	3	1	2	15.0	8.7	23.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	36	22	14	38	24	14	190.0	208.7	164.7
Heart diseases (410-443)	74	49	25	81	54	27	405.0	469.6	317.6
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	5	2	3	5	2	3	25.0	17.4	35.3
#Pneumonia (490-493)	13	5	8	13	5	8	65.0	43.5	94.1
#Diarrhea & enteritis (571)	2	0	2	2	0	2	10.0	-	23.5
Acute nephritis (590, 591)	1	0	1	1	0	1	5.0	-	11.8
Chronic nephritis, etc. (592-594)	3	2	1	3	2	1	15.0	17.4	11.8
Maternal deaths (640-689)	3	1	2	2	1	1	4.1	5.0	3.5
Cong. malformations (750-759)	1	1	0	1	1	0	5.0	8.7	-
Birth inj., atelect. (760-762)	3	1	2	3	2	1	6.1	10.0	3.5
#Other dis. early inf. (763-776)	10	2	8	11	3	8	22.4	14.9	27.7
Violent & acc. deaths (800-999)	12	10	2	19	15	4	95.0	130.4	47.0
Acc. deaths, total (800-962)	11	9	2	16	13	3	80.0	113.0	35.3
Motor vehicle acc. (810-835)	3	3	0	8	7	1	40.0	60.9	11.8
Ill-defined (780-795)	15	3	12	14	2	12	70.0	17.4	141.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
BENTON COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	116	5	111	232	86	146	21.1	15.9	26.1
Fetal deaths	3	0	3	9	4	5	38.8	46.5	34.2
Deaths from all causes	38	21	17	71	39	32	6.4	7.2	5.7
Deaths under 1 year	3	0	3	13	5	8	56.0	58.1	54.8
Deaths under 1 month	0	0	0	6	3	3	25.9	34.9	20.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	1	0	9.1	18.5	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	2	2	0	4	3	1	36.4	55.6	17.8
Diabetes (260)	2	1	1	4	2	2	36.4	37.0	35.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	8	3	5	11	3	8	100.0	55.6	142.8
Heart diseases (410-443)	12	10	2	18	14	4	163.6	259.2	71.4
Arteriosclerosis (450)	1	0	1	1	0	1	9.1	-	17.8
Influenza (480-483)	1	0	1	1	0	1	9.1	-	17.8
#Pneumonia (490-493)	0	0	0	1	1	0	9.1	18.5	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	2	2	0	18.2	37.0	-
Maternal deaths (640-689)	0	0	0	1	1	0	4.3	11.6	-
Cong. malformations (750-759)	0	0	0	2	2	0	18.2	37.0	-
Birth inj., atelect. (760-762)	0	0	0	1	1	0	4.3	11.6	-
#Other dis. early inf. (763-776)	0	0	0	3	1	2	12.9	11.6	13.7
Violent & acc. deaths (800-999)	6	2	4	8	1	7	72.7	18.5	125.0
Acc. deaths, total (800-962)	6	2	4	8	1	7	72.7	18.5	125.0
Motor vehicle acc. (810-835)	5	1	4	4	0	4	36.4	-	71.4
Ill-defined (780-795)	4	2	2	4	1	3	36.4	18.5	53.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
BOLIVAR COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,895	404	1,491	1,775	425	1,350	45.5	28.3	56.2
Fetal deaths	59	10	49	57	11	46	32.1	25.9	34.1
Deaths from all causes	627	137	490	647	178	469	16.6	11.9	19.5
Deaths under 1 year	88	10	78	90	14	76	50.7	32.9	56.3
Deaths under 1 month	45	7	38	43	9	34	24.2	21.2	25.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	5	3	2	12	4	8	30.8	26.7	33.3
Tbc., respiratory (001-008)	5	3	2	10	4	6	25.6	26.7	25.0
Syphilis (020-029)	2	0	2	2	0	2	5.1	-	8.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	1	0	1	1	0	1	2.6	-	4.2
Whooping cough (056)	1	0	1	1	0	1	2.6	-	4.2
Meningococcal infections (057)	1	0	1	1	0	1	2.6	-	4.2
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	55	15	40	58	22	36	148.7	146.7	150.0
Diabetes (260)	15	0	15	12	0	12	30.8	-	50.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	82	16	66	81	21	60	207.7	140.0	250.0
Heart diseases (410-443)	174	63	111	175	72	103	448.7	480.0	429.2
Arteriosclerosis (450)	3	0	3	5	2	3	12.8	13.3	12.5
Influenza (480-483)	15	2	13	13	1	12	33.3	6.7	50.0
#Pneumonia (490-493)	21	1	20	23	2	21	59.0	13.3	87.5
#Diarrhea & enteritis (571)	12	1	11	12	2	10	30.8	13.3	41.7
Acute nephritis (590, 591)	0	0	0	2	0	2	5.1	-	8.3
Chronic nephritis, etc. (592-594)	8	3	5	13	4	9	33.3	26.7	37.5
Maternal deaths (640-689)	2	0	2	1	0	1	0.6	-	0.7
Cong. malformations (750-759)	7	3	4	9	5	4	23.1	33.3	16.7
Birth inj., atelect. (760-762)	13	4	9	11	3	8	6.2	7.0	5.9
#Other dis. early inf. (763-776)	21	0	21	21	3	18	11.8	7.0	13.3
Violent & acc. deaths (800-999)	42	15	27	41	17	24	105.1	113.3	100.0
Acc. deaths, total (800-962)	30	9	21	31	12	19	79.5	80.0	79.2
Motor vehicle acc. (810-835)	9	2	7	10	5	5	25.6	33.3	20.8
Ill-defined (780-795)	101	4	97	102	5	97	261.5	33.3	404.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CALHOUN COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	382	203	179	337	190	147	19.8	17.3	24.5
Fetal deaths	17	9	8	13	6	7	38.6	31.6	47.6
Deaths from all causes	130	101	29	143	113	30	8.4	10.3	5.0
Deaths under 1 year	14	4	10	12	4	8	35.6	21.0	54.4
Deaths under 1 month	9	3	6	7	3	4	20.8	15.8	27.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	2	2	0	11.8	18.2	-
Tbc., respiratory (001-008)	1	1	0	2	2	0	11.8	18.2	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	15	13	2	20	16	4	117.6	145.4	66.7
Diabetes (260)	1	1	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	17	14	3	21	17	4	123.5	154.5	66.7
Heart diseases (410-443)	43	38	5	47	42	5	276.5	381.8	83.3
Arteriosclerosis (450)	5	5	0	5	5	0	29.4	45.4	-
Influenza (480-483)	2	2	0	2	2	0	11.8	18.2	-
#Pneumonia (490-493)	3	2	1	3	2	1	17.6	18.2	16.7
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	2	2	0	11.8	18.2	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	1	1	0	5.9	9.1	-
Birth inj., atelect. (760-762)	2	0	2	1	0	1	3.0	-	6.8
#Other dis. early inf. (763-776)	5	2	3	4	2	2	11.9	10.5	13.6
Violent & acc. deaths (800-999)	14	10	4	13	10	3	76.5	90.9	50.0
Acc. deaths, total (800-962)	8	5	3	8	5	3	47.0	45.4	50.0
Motor vehicle acc. (810-835)	5	4	1	6	5	1	35.3	45.4	16.7
Ill-defined (780-795)	14	6	8	15	6	9	88.2	54.5	150.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CARROLL COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	238	40	198	284	77	207	28.4	18.8	35.1
Fetal deaths	13	0	13	18	2	16	63.4	26.0	77.3
Deaths from all causes	87	21	66	133	51	82	13.3	12.4	13.9
Deaths under 1 year	11	1	10	15	1	14	52.8	13.0	67.6
Deaths under 1 month	4	0	4	7	1	6	24.6	13.0	29.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	1	0	1	1	0	1	10.0	-	16.9
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	8	4	4	17	11	6	170.0	268.3	101.7
Diabetes (260)	0	0	0	1	1	0	10.0	24.4	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	6	1	5	15	6	9	150.0	146.3	152.5
Heart diseases (410-443)	22	5	17	34	14	20	340.0	341.5	339.0
Arteriosclerosis (450)	0	0	0	1	1	0	10.0	24.4	-
Influenza (480-483)	2	0	2	2	0	2	20.0	-	33.9
#Pneumonia (490-493)	2	0	2	3	0	3	30.0	-	50.8
#Diarrhea & enteritis (571)	2	0	2	3	0	3	30.0	-	50.8
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	10.0	-	16.9
Maternal deaths (640-689)	0	0	0	2	0	2	7.0	-	9.7
Cong. malformations (750-759)	1	0	1	1	0	1	10.0	-	16.9
Birth inj., atelect. (760-762)	1	0	1	2	1	1	7.0	13.0	4.8
#Other dis. early inf. (763-776)	3	0	3	4	0	4	14.1	-	19.3
Violent & acc. deaths (800-999)	11	4	7	12	5	7	120.0	122.0	118.6
Acc. deaths, total (800-962)	10	3	7	10	4	6	100.0	97.6	101.7
Motor vehicle acc. (810-835)	8	2	6	5	1	4	50.0	24.4	67.8
Ill-defined (780-795)	16	0	16	15	0	15	150.0	-	254.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CHICKASAW COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	421	199	222	467	219	248	27.5	27.0	27.9
Fetal deaths	11	3	8	14	5	9	30.0	22.8	36.3
Deaths from all causes	152	85	67	166	97	69	9.8	12.0	7.8
Deaths under 1 year	13	1	12	17	2	15	36.4	9.1	60.5
Deaths under 1 month	8	1	7	8	1	7	17.1	4.6	28.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	2	0	1	1	0	5.9	12.3	-
Tbc., respiratory (001-008)	2	2	0	1	1	0	5.9	12.3	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	1	0	5.9	12.3	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)/	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	13	12	1	13	13	0	76.5	160.5	-
Diabetes (260)	1	0	1	3	2	1	17.6	24.7	11.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	21	12	9	25	13	12	147.0	160.5	134.8
Heart diseases (410-443)	57	38	19	57	41	16	335.3	506.2	179.8
Arteriosclerosis (450)	1	1	0	1	1	0	5.9	12.3	-
Influenza (480-483)	1	0	1	1	0	1	5.9	-	11.2
#Pneumonia (490-493)	8	3	5	8	3	5	47.0	37.0	56.2
#Diarrhea & enteritis (571)	1	1	0	2	1	1	11.8	12.3	11.2
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	0	1	5.9	-	11.2
Maternal deaths (640-689)	2	0	2	1	0	1	2.1	-	4.0
Cong. malformations (750-759)	0	0	0	2	0	2	11.8	-	22.5
Birth inj., atelect. (760-762)	1	1	0	2	1	1	4.3	4.6	4.0
#Other dis. early inf. (763-776)	5	0	5	5	0	5	10.7	-	20.2
Violent & acc. deaths (800-999)	12	6	6	15	8	7	88.2	98.8	78.6
Acc. deaths, total (800-962)	4	1	3	7	3	4	41.2	37.0	44.9
Motor vehicle acc. (810-835)	1	1	0	3	3	0	17.6	37.0	-
Ill-defined (780-795)	14	0	14	11	0	11	64.7	-	123.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CHOCTAW COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	189	90	99	190	91	99	17.3	12.1	28.3
Fetal deaths	6	3	3	8	4	4	42.1	44.0	40.4
Deaths from all causes	91	62	29	105	71	34	9.5	9.5	9.7
Deaths under 1 year	5	1	4	7	3	4	36.8	33.0	40.4
Deaths under 1 month	3	1	2	5	3	2	26.3	33.0	20.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	9.1	13.3	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	9.1	13.3	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	1	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	13	11	2	13	10	3	118.2	133.3	85.7
Diabetes (260)	6	3	3	6	3	3	54.5	40.0	85.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	11	6	5	14	8	6	127.3	106.7	171.4
Heart diseases (410-443)	32	27	5	35	29	6	318.2	386.7	171.4
Arteriosclerosis (450)	2	0	2	3	1	2	27.3	13.3	57.1
Influenza (480-483)	1	1	0	1	1	0	9.1	13.3	-
#Pneumonia (490-493)	1	1	0	2	2	0	18.2	26.7	-
#Diarrhea & enteritis (571)	1	0	1	1	0	1	9.1	-	28.6
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	9.1	13.3	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	2	1	1	18.2	13.3	28.6
Birth inj., atelect. (760-762)	2	0	2	2	0	2	10.5	-	20.2
#Other dis. early inf. (763-776)	1	1	0	1	1	0	5.3	11.0	-
Violent & acc. deaths (800-999)	5	4	1	6	5	1	54.5	66.7	28.6
Acc. deaths, total (800-962)	4	4	0	5	5	0	45.4	66.7	-
Motor vehicle acc. (810-835)	0	0	0	2	2	0	18.2	26.7	-
Ill-defined (780-795)	4	1	3	3	0	3	27.3	-	85.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CLAIBORNE COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	236	31	205	288	39	249	28.8	12.6	36.1
Fetal deaths	8	1	7	13	2	11	45.1	51.3	44.2
Deaths from all causes	100	24	76	136	37	99	13.6	11.9	14.3
Deaths under 1 year	12	1	11	17	1	16	59.0	25.6	64.2
Deaths under 1 month	4	0	4	7	0	7	24.3	-	28.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	10.0	-	14.5
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	10.0	-	14.5
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	8	0	8	12	1	11	120.0	32.2	159.4
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	10	2	8	15	3	12	150.0	96.8	173.9
Heart diseases (410-443)	30	12	18	40	19	21	400.0	612.9	304.3
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	1	0	1	1	0	1	10.0	-	14.5
#Pneumonia (490-493)	7	0	7	7	0	7	70.0	-	101.4
#Diarrhea & enteritis (571)	1	0	1	2	0	2	20.0	-	29.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	2	0	2	20.0	-	29.0
Maternal deaths (640-689)	1	0	1	1	0	1	3.5	-	4.0
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	1	0	1	3.5	-	4.0
#Other dis. early inf. (763-776)	2	0	2	4	0	4	13.9	-	16.1
Violent & acc. deaths (800-999)	8	2	6	9	1	8	90.0	32.2	115.9
Acc. deaths, total (800-962)	6	2	4	6	1	5	60.0	32.2	72.5
Motor vehicle acc. (810-835)	1	1	0	1	1	0	10.0	32.2	-
Ill-defined (780-795)	21	2	19	23	3	20	230.0	96.8	289.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CLARKE COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	229	112	117	430	201	229	26.9	20.7	36.3
Fetal deaths	3	0	3	10	3	7	23.2	14.9	30.6
Deaths from all causes	152	87	65	189	111	78	11.8	11.4	12.4
Deaths under 1 year	11	3	8	10	1	9	23.2	5.0	39.3
Deaths under 1 month	7	2	5	7	1	6	16.3	5.0	26.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	6.2	10.3	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	6.2	10.3	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	21	13	8	28	17	11	175.0	175.2	174.6
Diabetes (260)	2	2	0	3	3	0	18.8	30.9	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	19	8	11	29	17	12	181.2	175.2	190.5
Heart diseases (410-443)	57	43	14	62	47	15	387.5	484.5	238.1
Arteriosclerosis (450)	3	0	3	2	0	2	12.5	-	31.7
Influenza (480-483)	1	0	1	1	0	1	6.2	-	15.9
#Pneumonia (490-493)	5	1	4	6	1	5	37.5	10.3	79.4
#Diarrhea & enteritis (571)	1	1	0	1	1	0	6.2	10.3	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	2	0	2	12.5	-	31.7
Maternal deaths (640-689)	0	0	0	1	0	1	2.3	-	4.4
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	2	1	1	2	0	2	4.6	-	8.7
#Other dis. early inf. (763-776)	4	1	3	4	1	3	9.3	5.0	13.1
Violent & acc. deaths (800-999)	16	12	4	16	12	4	100.0	123.7	63.5
Acc. deaths, total (800-962)	11	9	2	11	9	2	68.8	92.8	31.7
Motor vehicle acc. (810-835)	10	8	2	9	7	2	56.2	72.2	31.7
Ill-defined (780-795)	12	3	9	12	3	9	75.0	30.9	142.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CLAY COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	474	191	283	495	202	293	27.5	20.2	36.6
Fetal deaths	9	2	7	9	2	7	18.2	9.9	23.9
Deaths from all causes	178	71	107	185	74	111	10.3	7.4	13.9
Deaths under 1 year	20	5	15	22	6	16	44.4	29.7	54.6
Deaths under 1 month	14	5	9	16	6	10	32.3	29.7	34.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	2	0	2	11.1	-	25.0
Tbc., respiratory (001-008)	1	0	1	2	0	2	11.1	-	25.0
Syphilis (020-029)	1	0	1	1	0	1	5.6	-	12.5
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	0	1	1	0	1	5.6	-	12.5
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	19	8	11	20	9	11	111.1	90.0	137.5
Diabetes (260)	2	0	2	2	0	2	11.1	-	25.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	17	7	10	18	6	12	100.0	60.0	150.0
Heart diseases (410-443)	57	28	29	57	29	28	316.7	290.0	350.0
Arteriosclerosis (450)	2	2	0	2	2	0	11.1	20.0	-
Influenza (480-483)	1	0	1	1	0	1	5.6	-	12.5
#Pneumonia (490-493)	5	2	3	5	2	3	27.8	20.0	37.5
#Diarrhea & enteritis (571)	3	0	3	3	0	3	16.7	-	37.5
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	2	1	1	11.1	10.0	12.5
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	8	5	3	8	5	3	16.2	24.8	10.2
#Other dis. early inf. (763-776)	4	0	4	4	0	4	8.1	-	13.6
Violent & acc. deaths (800-999)	22	12	10	22	12	10	122.2	120.0	125.0
Acc. deaths, total (800-962)	16	7	9	16	7	9	88.9	70.0	112.5
Motor vehicle acc. (810-835)	6	4	2	7	4	3	38.9	40.0	37.5
Ill-defined (780-795)	24	3	21	25	3	22	138.9	30.0	275.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
COAHOMA COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,496	363	1,133	1,531	327	1,204	38.3	21.8	48.2
Fetal deaths	83	13	70	79	8	71	51.6	24.5	59.0
Deaths from all causes	566	141	425	597	143	454	14.9	9.5	18.2
Deaths under 1 year	82	8	74	78	6	72	50.9	18.3	59.8
Deaths under 1 month	40	4	36	37	3	34	24.2	9.2	28.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	11	2	9	12	2	10	30.0	13.3	40.0
Tbc., respiratory (001-008)	8	2	6	9	2	7	22.5	13.3	28.0
Syphilis (020-029)	2	0	2	4	0	4	10.0	-	16.0
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	2.5	-	4.0
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	64	25	39	72	24	48	180.0	160.0	192.0
Diabetes (260)	8	2	6	9	2	7	22.5	13.3	28.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	54	14	40	64	18	46	160.0	120.0	184.0
Heart diseases (410-443)	159	55	104	168	53	115	420.0	353.3	460.0
Arteriosclerosis (450)	4	2	2	5	3	2	12.5	20.0	8.0
Influenza (480-483)	11	0	11	10	0	10	25.0	-	40.0
#Pneumonia (490-493)	31	5	26	32	5	27	80.0	33.3	108.0
#Diarrhea & enteritis (571)	8	1	7	8	0	8	20.0	-	32.0
Acute nephritis (590, 591)	2	0	2	3	0	3	7.5	-	12.0
Chronic nephritis, etc. (592-594)	9	1	8	11	1	10	27.5	6.7	40.0
Maternal deaths (640-689)	1	1	0	2	1	1	1.3	3.0	0.8
Cong. malformations (750-759)	9	3	6	7	2	5	17.5	13.3	20.0
Birth inj., atelect. (760-762)	7	1	6	8	2	6	5.2	6.1	5.0
#Other dis. early inf. (763-776)	27	1	26	25	1	24	16.3	3.0	19.9
Violent & acc. deaths (800-999)	57	14	43	51	14	37	127.5	93.3	148.0
Acc. deaths, total (800-962)	49	12	37	44	12	32	110.0	80.0	128.0
Motor vehicle acc. (810-835)	19	8	11	12	7	5	30.0	46.7	20.0
Ill-defined (780-795)	53	0	53	54	1	53	135.0	6.7	212.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
COPIAH COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	877	248	629	716	221	495	25.6	15.8	35.4
Fetal deaths	23	3	20	24	6	18	33.5	27.1	36.4
Deaths from all causes	278	124	154	315	151	164	11.2	10.8	11.7
Deaths under 1 year	29	3	26	33	4	29	46.1	18.1	58.6
Deaths under 1 month	15	2	13	17	2	15	23.7	9.0	30.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	3.6	7.1	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	3.6	7.1	-
Syphilis (020-029)	1	1	0	1	1	0	3.6	7.1	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	3.6	-	7.1
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	2	1	1	2	1	1	7.1	7.1	7.1
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	39	21	18	44	26	18	157.1	185.7	128.6
Diabetes (260)	5	3	2	5	3	2	17.8	21.4	14.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	26	13	13	29	15	14	103.6	107.1	100.0
Heart diseases (410-443)	113	57	56	124	68	56	442.8	485.7	400.0
Arteriosclerosis (450)	2	1	1	3	2	1	10.7	14.3	7.1
Influenza (480-483)	4	1	3	4	1	3	14.3	7.1	21.4
#Pneumonia (490-493)	6	3	3	8	4	4	28.6	28.6	28.6
#Diarrhea & enteritis (571)	5	2	3	5	2	3	17.8	14.3	21.4
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	5	0	5	5	0	5	17.8	-	35.7
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	1	1	1	1	0	3.6	7.1	-
Birth inj., atelect. (760-762)	3	0	3	6	1	5	8.4	4.5	10.1
#Other dis. early inf. (763-776)	11	1	10	11	0	11	15.4	-	22.2
Violent & acc. deaths (800-999)	15	7	8	19	8	11	67.8	57.1	78.6
Acc. deaths, total (800-962)	12	5	7	16	7	9	57.1	50.0	64.3
Motor vehicle acc. (810-835)	5	3	2	5	2	3	17.8	14.3	21.4
Ill-defined (780-795)	11	2	9	12	2	10	42.8	14.3	71.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
COVINGTON COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	221	158	63	382	216	166	21.2	20.6	22.1
Fetal deaths	3	2	1	4	3	1	10.5	13.9	6.0
Deaths from all causes	88	60	28	116	83	33	6.4	7.9	4.4
Deaths under 1 year	2	1	1	4	2	2	10.5	9.2	12.0
Deaths under 1 month	0	0	0	2	1	1	5.2	4.6	6.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	5.6	9.5	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	5.6	9.5	-
Syphilis (020-029)	1	0	1	1	0	1	5.6	-	13.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	1	0	1	1	0	1	5.6	-	13.3
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	1	0	1	1	0	5.6	9.5	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	7	3	14	10	4	77.8	95.2	53.3
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	13	9	4	18	14	4	100.0	133.3	53.3
Heart diseases (410-443)	34	25	9	41	33	8	227.8	314.3	106.7
Arteriosclerosis (450)	0	0	0	2	1	1	11.1	9.5	13.3
Influenza (480-483)	2	2	0	2	2	0	11.1	19.0	-
#Pneumonia (490-493)	4	3	1	5	4	1	27.8	38.1	13.3
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	0	1	1	0	1	5.6	-	13.3
Chronic nephritis, etc. (592-594)	1	1	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	1	1	0	2.6	4.6	-
#Other dis. early inf. (763-776)	0	0	0	1	0	1	2.6	-	6.0
Violent & acc. deaths (800-999)	8	5	3	13	9	4	72.2	85.7	53.3
Acc. deaths, total (800-962)	6	4	2	11	8	3	61.1	76.2	40.0
Motor vehicle acc. (810-835)	3	2	1	7	5	2	38.9	47.6	26.7
Ill-defined (780-795)	4	2	2	5	2	3	27.8	19.0	40.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
DESOTO COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	643	60	583	703	170	533	30.6	17.9	39.5
Fetal deaths	15	0	15	26	1	25	37.0	5.9	46.9
Deaths from all causes	164	37	127	225	66	159	9.8	6.9	11.8
Deaths under 1 year	21	1	20	32	4	28	45.5	23.5	52.5
Deaths under 1 month	12	0	12	21	3	18	29.9	17.6	33.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	4.3	-	7.4
Tbc., respiratory (001-008)	1	0	1	1	0	1	4.3	-	7.4
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	4.3	-	7.4
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	4.3	-	7.4
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	5	12	31	13	18	134.8	136.8	133.3
Diabetes (260)	2	1	1	3	1	2	13.0	10.5	14.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	24	3	21	31	9	22	134.8	94.7	163.0
Heart diseases (410-443)	33	14	19	47	20	27	204.3	210.5	200.0
Arteriosclerosis (450)	4	1	3	3	1	2	13.0	10.5	14.8
Influenza (480-483)	1	0	1	2	1	1	8.7	10.5	7.4
#Pneumonia (490-493)	15	3	12	17	4	13	73.9	42.1	96.3
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	1	0	1	4.3	-	7.4
Chronic nephritis, etc. (592-594)	2	1	1	1	1	0	4.3	10.5	-
Maternal deaths (640-689)	1	0	1	1	0	1	1.4	-	1.9
Cong. malformations (750-759)	1	0	1	2	0	2	8.7	-	14.8
Birth inj., atelect.(760-762)	1	0	1	5	2	3	7.1	11.8	5.6
#Other dis. early inf. (763-776)	8	0	8	12	1	11	17.1	5.9	20.6
Violent & acc. deaths (800-999)	15	7	8	17	7	10	73.9	73.7	74.1
Acc. deaths, total (800-962)	11	7	4	12	6	6	52.2	63.2	44.4
Motor vehicle acc. (810-835)	5	2	3	6	3	3	26.1	31.6	22.2
Ill-defined (780-795)	24	1	23	27	1	26	117.4	10.5	192.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
FORREST COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,471	1,081	390	1,271	787	484	24.9	22.5	30.2
Fetal deaths	49	33	16	48	23	25	37.8	29.2	51.6
Deaths from all causes	480	324	156	474	301	173	9.3	8.6	10.8
Deaths under 1 year	46	32	14	44	29	15	34.6	36.8	31.0
Deaths under 1 month	28	20	8	28	19	9	22.0	24.1	18.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	2	2	8	6	2	15.7	17.1	12.5
Tbc., respiratory (001-008)	4	2	2	7	5	2	13.7	14.3	12.5
Syphilis (020-029)	0	0	0	1	0	1	2.0	-	6.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	1	0	1	0	1	2.0	-	6.2
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	59	46	13	56	40	16	109.8	114.3	100.0
Diabetes (260)	10	7	3	9	6	3	17.6	17.1	18.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	61	40	21	63	40	23	123.5	114.3	143.8
Heart diseases (410-443)	185	126	59	177	117	60	347.0	334.3	375.0
Arteriosclerosis (450)	7	6	1	5	4	1	9.8	11.4	6.2
Influenza (480-483)	1	1	0	1	1	0	2.0	2.8	-
#Pneumonia (490-493)	13	8	5	11	7	4	21.6	20.0	25.0
#Diarrhea & enteritis (571)	5	3	2	5	3	2	9.8	8.6	12.5
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	2	2	5	2	3	9.8	5.7	18.8
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	7	5	2	7	5	2	13.7	14.3	12.5
Birth inj., atelect. (760-762)	6	5	1	7	5	2	5.5	6.4	4.1
#Other dis. early inf. (763-776)	17	11	6	17	11	6	13.4	14.0	12.4
Violent & acc. deaths (800-999)	35	25	10	32	19	13	62.7	54.3	81.2
Acc. deaths, total (800-962)	25	21	4	20	13	7	39.2	37.1	43.8
Motor vehicle acc. (810-835)	16	15	1	5	4	1	9.8	11.4	6.2
Ill-defined (780-795)	20	5	15	22	5	17	43.1	14.3	106.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
FRANKLIN COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	133	60	73	232	96	136	27.0	20.0	35.8
Fetal deaths	7	1	6	8	2	6	34.5	20.8	44.1
Deaths from all causes	81	41	40	102	54	48	11.9	11.2	12.6
Deaths under 1 year	6	0	6	10	1	9	43.1	10.4	66.2
Deaths under 1 month	4	0	4	8	1	7	34.5	10.4	51.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	1	2	3	1	2	34.9	20.8	52.6
Tbc., respiratory (001-008)	3	1	2	3	1	2	34.9	20.8	52.6
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	8	4	4	9	6	3	104.6	125.0	78.9
Diabetes (260)	1	0	1	1	0	1	11.6	-	26.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	6	3	3	7	4	3	81.4	83.3	78.9
Heart diseases (410-443)	41	22	19	44	25	19	511.6	520.8	500.0
Arteriosclerosis (450)	6	4	2	7	5	2	81.4	104.2	52.6
Influenza (480-483)	1	0	1	1	0	1	11.6	-	26.3
#Pneumonia (490-493)	1	1	0	2	2	0	23.2	41.7	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	0	1	11.6	-	26.3
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	2	1	1	23.2	20.8	26.3
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	4	0	4	7	1	6	30.2	10.4	44.1
Violent & acc. deaths (800-999)	3	3	0	10	7	3	116.3	145.8	78.9
Acc. deaths, total (800-962)	3	3	0	10	7	3	116.3	145.8	78.9
Motor vehicle acc. (810-835)	0	0	0	3	3	0	34.9	62.5	-
Ill-defined (780-795)	4	1	3	4	1	3	46.5	20.8	78.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
GEORGE COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	337	282	55	345	288	57	21.6	20.6	28.5
Fetal deaths	8	4	4	9	5	4	26.1	17.4	70.2
Deaths from all causes	75	63	12	84	72	12	5.2	5.1	6.0
Deaths under 1 year	9	6	3	12	8	4	34.8	27.8	70.2
Deaths under 1 month	5	4	1	7	6	1	20.3	20.8	17.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	3	3	0	18.8	21.4	-
Tbc., respiratory (001-008)	0	0	0	3	3	0	18.8	21.4	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	1	0	1	1	0	6.2	7.1	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	9	0	9	8	1	56.2	57.1	50.0
Diabetes (260)	1	1	0	1	1	0	6.2	7.1	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	12	9	3	10	8	2	62.5	57.1	100.0
Heart diseases (410-443)	29	25	4	32	28	4	200.0	200.0	200.0
Arteriosclerosis (450)	0	0	0	1	1	0	6.2	7.1	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	2	0	1	1	0	6.2	7.1	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	6.2	7.1	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	3	2	1	18.8	14.3	50.0
Birth inj., atelect. (760-762)	3	3	0	4	4	0	11.6	13.9	-
#Other dis. early inf. (763-776)	2	1	1	3	2	1	8.7	6.9	17.5
Violent & acc. deaths (800-999)	10	8	2	10	8	2	62.5	57.1	100.0
Acc. deaths, total (800-962)	8	7	1	9	8	1	56.2	57.1	50.0
Motor vehicle acc. (810-835)	2	2	0	3	3	0	18.8	21.4	-
Ill-defined (780-795)	2	0	2	1	0	1	6.2	-	50.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
GREENE COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	112	62	50	189	120	69	24.2	19.7	40.6
Fetal deaths	5	3	2	4	2	2	21.2	16.7	29.0
Deaths from all causes	68	54	14	83	64	19	10.6	10.5	11.2
Deaths under 1 year	4	2	2	5	3	2	26.4	25.0	29.0
Deaths under 1 month	1	1	0	2	2	0	10.6	16.7	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	12.8	-	58.8
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	1	1	0	3	1	2	38.5	16.4	117.6
Diabetes (260)	0	0	0	1	0	1	12.8	-	58.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	10	8	2	12	10	2	153.8	163.9	117.6
Heart diseases (410-443)	28	23	5	35	29	6	448.7	475.4	352.9
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	1	0	1	1	0	1	12.8	-	58.8
#Pneumonia (490-493)	2	1	1	3	2	1	38.5	32.8	58.8
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	12.8	16.4	-
Birth inj., atelect. (760-762)	0	0	0	1	1	0	5.3	8.3	-
#Other dis. early inf. (763-776)	1	1	0	1	1	0	5.3	8.3	-
Violent & acc. deaths (800-999)	16	13	3	14	11	3	179.5	180.3	176.5
Acc. deaths, total (800-962)	14	13	1	12	11	1	153.8	180.3	58.8
Motor vehicle acc. (810-835)	7	7	0	7	7	0	89.7	114.8	-
Ill-defined (780-795)	4	3	1	5	4	1	64.1	65.6	58.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
GRENADA COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	555	230	325	516	194	322	30.4	24.0	36.2
Fetal deaths	21	1	20	21	3	18	40.7	15.5	55.9
Deaths from all causes	221	97	124	201	84	117	11.8	10.4	13.1
Deaths under 1 year	26	4	22	20	3	17	38.8	15.5	52.8
Deaths under 1 month	15	3	12	12	2	10	23.2	10.3	31.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	2	0	2	11.8	-	22.5
Tbc., respiratory (001-008)	2	0	2	2	0	2	11.8	-	22.5
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	25	17	8	30	19	11	176.5	234.6	123.6
Diabetes (260)	2	1	1	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	44	17	27	37	11	26	217.6	135.8	292.1
Heart diseases (410-443)	59	30	29	51	24	27	300.0	296.3	303.4
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	6	1	5	6	1	5	35.3	12.3	56.2
#Pneumonia (490-493)	7	2	5	6	2	4	35.3	24.7	44.9
#Diarrhea & enteritis (571)	3	1	2	2	0	2	11.8	-	22.5
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	5.9	12.3	-
Maternal deaths (640-689)	1	0	1	1	1	0	1.9	5.2	-
Cong. malformations (750-759)	2	1	1	2	2	0	11.8	24.7	-
Birth inj., atelect. (760-762)	3	3	0	2	2	0	3.9	10.3	-
#Other dis. early inf. (763-776)	8	0	8	8	0	8	15.5	-	24.8
Violent & acc. deaths (800-999)	25	10	15	23	8	15	135.3	98.8	168.5
Acc. deaths, total (800-962)	22	8	14	20	6	14	117.6	74.1	157.3
Motor vehicle acc. (810-835)	5	3	2	3	2	1	17.6	24.7	11.2
Ill-defined (780-795)	18	6	12	17	6	11	100.0	74.1	123.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HANCOCK COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	135	103	32	266	217	49	22.2	23.6	17.5
Fetal deaths	2	2	0	3	3	0	11.3	13.8	-
Deaths from all causes	99	75	24	133	103	30	11.1	11.2	10.7
Deaths under 1 year	5	5	0	7	4	3	26.3	18.4	61.2
Deaths under 1 month	1	1	0	3	1	2	11.3	4.6	40.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	2	1	1	16.7	10.9	35.7
Tbc., respiratory (001-008)	2	1	1	2	1	1	16.7	10.9	35.7
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	1	0	8.3	10.9	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	9	1	21	20	1	175.0	217.4	35.7
Diabetes (260)	2	2	0	3	3	0	25.0	32.6	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	17	12	5	23	17	6	191.7	184.8	214.3
Heart diseases (410-443)	27	23	4	42	36	6	350.0	391.3	214.3
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	1	0	2	2	0	16.7	21.7	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	0	1	8.3	-	35.7
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	2	0	2	7.5	-	40.8
#Other dis. early inf. (763-776)	1	1	0	1	1	0	3.8	4.6	-
Violent & acc. deaths (800-999)	21	10	11	15	6	9	125.0	65.2	321.4
Acc. deaths, total (800-962)	21	10	11	15	6	9	125.0	65.2	321.4
Motor vehicle acc. (810-835)	15	7	8	9	1	8	75.0	10.9	285.7
Ill-defined (780-795)	16	14	2	13	11	2	108.3	119.6	71.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HARRISON COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	3,691	2,991	700	3,394	2,722	672	20.6	20.5	21.0
Fetal deaths	85	53	32	79	48	31	23.3	17.6	46.1
Deaths from all causes	895	709	186	779	596	183	4.7	4.5	5.7
Deaths under 1 year	96	73	23	93	71	22	27.4	26.1	32.7
Deaths under 1 month	66	53	13	63	50	13	18.6	18.4	19.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	3	1	3	2	1	1.8	1.5	3.1
Tbc., respiratory (001-008)	2	2	0	2	2	0	1.2	1.5	-
Syphilis (020-029)	4	4	0	2	2	0	1.2	1.5	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	1	1	0	1	1	0	0.6	0.8	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	2	2	0	2	2	0	1.2	1.5	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	129	103	26	111	86	25	67.3	64.7	78.1
Diabetes (260)	9	7	2	8	6	2	4.8	4.5	6.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	98	74	24	84	61	23	50.9	45.9	71.9
Heart diseases (410-443)	283	240	43	243	199	44	147.3	149.6	137.5
Arteriosclerosis (450)	14	11	3	14	11	3	8.4	8.3	9.4
Influenza (480-483)	3	3	0	3	3	0	1.8	2.2	-
#Pneumonia (490-493)	27	23	4	21	16	5	12.7	12.0	15.6
#Diarrhea & enteritis (571)	5	4	1	4	4	0	2.4	3.0	-
Acute nephritis (590, 591)	1	1	0	1	1	0	0.6	0.8	-
Chronic nephritis, etc. (592-594)	5	3	2	5	3	2	3.0	2.2	6.2
Maternal deaths (640-689)	1	1	0	1	1	0	0.3	0.4	-
Cong. malformations (750-759)	14	12	2	15	13	2	9.1	9.8	6.2
Birth inj., atelect. (760-762)	25	19	6	25	19	6	7.4	7.0	8.9
#Other dis. early inf. (763-776)	29	26	3	26	23	3	7.7	8.4	4.5
Violent & acc. deaths (800-999)	78	55	23	75	50	25	45.4	37.6	78.1
Acc. deaths, total (800-962)	60	48	12	58	44	14	35.2	33.1	43.8
Motor vehicle acc. (810-835)	21	19	2	20	17	3	12.1	12.8	9.4
Ill-defined (780-795)	69	42	27	65	38	27	39.4	28.6	84.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HINDS COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
			Non-white	Number		Non-white	Rate*		Non-white
	Total	White		Total	White		Total	White	
Live births	5,546	2,926	2,620	5,146	2,568	2,578	29.2	24.0	37.4
Fetal deaths	387	233	154	348	209	139	67.6	81.4	53.9
Deaths from all causes	2,138	1,109	1,029	1,502	706	796	8.5	6.6	11.5
Deaths under 1 year	291	105	186	217	68	149	42.2	26.5	57.8
Deaths under 1 month	195	83	112	147	59	88	28.6	23.0	34.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	25	11	14	16	9	7	9.1	8.4	10.1
Tbc., respiratory (001-008)	20	8	12	14	7	7	8.0	6.5	10.1
Syphilis (020-029)	7	2	5	3	0	3	1.7	-	4.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	3	0	3	3	1	2	1.7	0.9	2.9
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	2	0	2	0	0	0	-	-	-
Whooping cough (056)	2	0	2	1	0	1	0.6	-	1.4
Meningococcal infections (057)	3	2	1	2	1	1	1.1	0.9	1.4
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	301	172	129	176	94	82	100.0	87.8	118.8
Diabetes (260)	36	19	17	29	13	16	16.5	12.1	23.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	189	101	88	151	69	82	85.8	64.5	118.8
Heart diseases (410-443)	534	334	200	397	237	160	225.6	221.5	231.9
Arteriosclerosis (450)	34	26	8	24	16	8	13.6	15.0	11.6
Influenza (480-483)	8	4	4	7	3	4	4.0	2.8	5.8
#Pneumonia (490-493)	74	33	41	52	18	34	29.5	16.8	49.3
#Diarrhea & enteritis (571)	14	1	13	13	0	13	7.4	-	18.8
Acute nephritis (590, 591)	9	2	7	6	0	6	3.4	-	8.7
Chronic nephritis, etc. (592-594)	32	9	23	26	6	20	14.8	5.6	29.0
Maternal deaths (640-689)	4	1	3	3	0	3	0.6	-	1.2
Cong. malformations (750-759)	43	29	14	26	16	10	14.8	15.0	14.5
Birth inj., atelect. (760-762)	60	29	31	42	20	22	8.2	7.8	8.5
#Other dis. early inf. (763-776)	100	38	62	80	30	50	15.5	11.7	19.4
Violent & acc. deaths (800-999)	167	92	75	116	59	57	65.9	55.1	82.6
Acc. deaths, total (800-962)	135	78	57	88	45	43	50.0	42.0	62.3
Motor vehicle acc. (810-835)	64	38	26	37	21	16	21.0	19.6	23.2
Ill-defined (780-795)	163	27	136	151	24	127	85.8	22.4	184.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HOLMES COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	804	135	669	786	120	666	32.8	17.6	38.7
Fetal deaths	44	8	36	45	8	37	57.2	66.7	55.6
Deaths from all causes	306	81	225	335	88	247	14.0	12.9	14.4
Deaths under 1 year	46	3	43	44	2	42	56.0	16.7	63.1
Deaths under 1 month	25	2	23	22	1	21	28.0	8.3	31.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	2	1	1	8.3	14.7	5.8
Tbc., respiratory (001-008)	0	0	0	2	1	1	8.3	14.7	5.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	4.2	-	5.8
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	0	1	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	43	15	28	50	16	34	208.3	235.3	197.7
Diabetes (260)	3	1	2	3	1	2	12.5	14.7	11.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	41	7	34	44	10	34	183.3	147.0	197.7
Heart diseases (410-443)	76	32	44	80	31	49	333.3	455.9	284.9
Arteriosclerosis (450)	5	2	3	5	2	3	20.8	29.4	17.4
Influenza (480-483)	3	2	1	3	2	1	12.5	29.4	5.8
#Pneumonia (490-493)	14	1	13	16	2	14	66.7	29.4	81.4
#Diarrhea & enteritis (571)	2	1	1	1	1	0	4.2	14.7	-
Acute nephritis (590, 591)	1	0	1	1	0	1	4.2	-	5.8
Chronic nephritis, etc. (592-594)	0	0	0	2	1	1	8.3	14.7	5.8
Maternal deaths (640-689)	1	0	1	1	0	1	1.3	-	1.5
Cong. malformations (750-759)	2	0	2	3	0	3	12.5	-	17.4
Birth inj., atelect. (760-762)	2	0	2	1	0	1	1.3	-	1.5
#Other dis. early inf. (763-776)	22	2	20	19	1	18	24.2	8.3	27.0
Violent & acc. deaths (800-999)	23	9	14	25	6	19	104.2	88.2	110.5
Acc. deaths, total (800-962)	22	9	13	22	6	16	91.7	88.2	93.0
Motor vehicle acc. (810-835)	10	7	3	7	3	4	29.2	44.1	23.2
Ill-defined (780-795)	40	2	38	41	3	38	170.8	44.1	220.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HUMPHREYS COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	662	109	553	664	115	549	41.5	23.5	49.4
Fetal deaths	28	3	25	34	3	31	51.2	26.1	56.5
Deaths from all causes	175	43	132	207	55	152	12.9	11.2	13.7
Deaths under 1 year	24	2	22	22	1	21	33.1	8.7	38.2
Deaths under 1 month	14	1	13	13	1	12	19.6	8.7	21.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	2	0	2	12.5	-	18.0
Tbc., respiratory (001-008)	1	0	1	1	0	1	6.2	-	9.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	0	1	1	0	1	6.2	-	9.0
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	15	5	10	21	8	13	131.2	163.3	117.1
Diabetes (260)	2	0	2	2	0	2	12.5	-	18.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	31	8	23	37	11	26	231.2	224.5	234.2
Heart diseases (410-443)	36	16	20	40	18	22	250.0	367.3	198.2
Arteriosclerosis (450)	3	1	2	4	2	2	25.0	40.8	18.0
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	0	1	3	2	1	18.8	40.8	9.0
#Diarrhea & enteritis (571)	5	0	5	5	0	5	31.2	-	45.0
Acute nephritis (590, 591)	2	0	2	3	0	3	18.8	-	27.0
Chronic nephritis, etc. (592-594)	5	0	5	6	0	6	37.5	-	54.0
Maternal deaths (640-689)	2	0	2	2	0	2	3.0	-	3.6
Cong. malformations (750-759)	3	2	1	2	1	1	12.5	20.4	9.0
Birth inj., atelect. (760-762)	2	0	2	2	0	2	3.0	-	3.6
#Other dis. early inf. (763-776)	12	1	11	11	1	10	16.6	8.7	18.2
Violent & acc. deaths (800-999)	8	4	4	10	4	6	62.5	81.6	54.0
Acc. deaths, total (800-962)	5	3	2	7	3	4	43.8	61.2	36.0
Motor vehicle acc. (810-835)	1	1	0	3	3	0	18.8	61.2	-
Ill-defined (780-795)	31	0	31	35	0	35	218.8	-	315.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ISSAQUNA COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	36	0	36	104	19	85	28.1	10.6	44.7
Fetal deaths	2	0	2	2	0	2	19.2	-	23.5
Deaths from all causes	30	8	22	38	10	28	10.3	5.6	14.7
Deaths under 1 year	3	0	3	6	0	6	57.7	-	70.6
Deaths under 1 month	1	0	1	4	0	4	38.5	-	47.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	2	1	1	6	2	4	162.2	111.1	210.5
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	4	1	3	5	2	3	135.1	111.1	157.9
Heart diseases (410-443)	7	1	6	8	2	6	216.2	111.1	315.8
Arteriosclerosis (450)	1	1	0	1	1	0	27.0	55.5	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	0	0	0	0	0	0	-	-	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	1	0	1	4	0	4	38.5	-	47.0
Violent & acc. deaths (800-999)	10	4	6	9	3	6	243.2	166.7	315.8
Acc. deaths, total (800-962)	9	4	5	8	3	5	216.2	166.7	263.2
Motor vehicle acc. (810-835)	2	1	1	2	1	1	54.0	55.5	52.6
Ill-defined (780-795)	3	0	3	3	0	3	81.1	-	157.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ITAWAMBA COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence				Rate*		
	Total	White	Non-white	Number			Total	White	Non-white	
				Total	White	Non-white				
Live births	62	53	9	241	224	17	11.5	11.2	15.4	
Fetal deaths	0	0	0	3	2	1	12.4	8.9	58.8	
Deaths from all causes	107	97	10	131	118	13	6.2	5.9	11.8	
Deaths under 1 year	2	1	1	4	1	3	16.6	4.5	176.5	
Deaths under 1 month	1	1	0	3	1	2	12.4	4.5	117.6	
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.										
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-	
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-	
Syphilis (020-029)	0	0	0	0	0	0	-	-	-	
Typhoid fever (040)	0	0	0	0	0	0	-	-	-	
Dysentery (045-048)	0	0	0	0	0	0	-	-	-	
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-	
Diphtheria (055)	0	0	0	0	0	0	-	-	-	
Whooping cough (056)	0	0	0	0	0	0	-	-	-	
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-	
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-	
Smallpox (084)	0	0	0	0	0	0	-	-	-	
Measles (085)	0	0	0	0	0	0	-	-	-	
Malaria (110-117)	0	0	0	0	0	0	-	-	-	
Cancer, incl. leukemia (140-205)	14	13	1	18	17	1	85.7	85.4	90.9	
Diabetes (260)	4	4	0	5	5	0	23.8	25.1	-	
Pellagra (281)	0	0	0	0	0	0	-	-	-	
Intracranial lesions (330-334)	21	20	1	24	23	1	114.3	115.6	90.9	
Heart diseases (410-443)	43	40	3	50	47	3	238.1	236.2	272.7	
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-	
Influenza (480-483)	0	0	0	0	0	0	-	-	-	
#Pneumonia (490-493)	1	1	0	2	2	0	9.5	10.0	-	
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-	
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-	
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	4.8	-	90.9	
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-	
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-	
Birth inj., atelect. (760-762)	0	0	0	2	0	2	8.3	-	117.6	
#Other dis. early inf. (763-776)	0	0	0	0	0	0	-	-	-	
Violent & acc. deaths (800-999)	9	6	3	12	9	3	57.1	45.2	272.7	
Acc. deaths, total (800-962)	5	3	2	6	4	2	28.6	20.1	181.8	
Motor vehicle acc. (810-835)	4	2	2	5	3	2	23.8	15.1	181.8	
Ill-defined (780-795)	8	7	1	8	7	1	38.1	35.2	90.9	

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JACKSON COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,277	889	388	1,532	1,120	412	26.0	27.6	22.3
Fetal deaths	52	38	14	59	44	15	38.5	39.3	36.4
Deaths from all causes	364	257	107	383	275	108	6.5	6.8	5.8
Deaths under 1 year	42	28	14	48	33	15	31.3	29.5	36.4
Deaths under 1 month	34	23	11	37	26	11	24.2	23.2	26.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	4	2	2	6.8	4.9	10.8
Tbc., respiratory (001-008)	2	0	2	4	2	2	6.8	4.9	10.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	1	1	0	1.7	2.5	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	1	0	1	1	0	1.7	2.5	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	42	30	12	51	39	12	86.4	96.3	64.9
Diabetes (260)	5	4	1	5	4	1	8.5	9.9	5.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	43	31	12	47	34	13	79.7	84.0	70.3
Heart diseases (410-443)	117	96	21	115	94	21	194.9	232.1	113.5
Arteriosclerosis (450)	4	4	0	3	3	0	5.1	7.4	-
Influenza (480-483)	8	6	2	7	5	2	11.9	12.3	10.8
#Pneumonia (490-493)	5	3	2	7	4	3	11.9	9.9	16.2
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	3.4	2.5	5.4
Maternal deaths (640-689)	2	1	1	1	1	0	0.7	0.9	-
Cong. malformations (750-759)	8	6	2	10	8	2	16.9	19.8	10.8
Birth inj., atelect. (760-762)	7	6	1	7	6	1	4.6	5.4	2.4
#Other dis. early inf. (763-776)	20	12	8	22	14	8	14.4	12.5	19.4
Violent & acc. deaths (800-999)	47	28	19	41	23	18	69.5	56.8	97.3
Acc. deaths, total (800-962)	36	24	12	31	19	12	52.5	46.9	64.9
Motor vehicle acc. (810-835)	14	13	1	12	10	2	20.3	24.7	10.8
Ill-defined (780-795)	22	12	10	17	7	10	28.8	17.3	54.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JASPER COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	124	45	79	438	152	286	25.8	16.5	36.7
Fetal deaths	12	2	10	27	8	19	61.6	52.6	66.4
Deaths from all causes	114	46	68	184	84	100	10.8	9.1	12.8
Deaths under 1 year	10	1	9	19	3	16	43.4	19.7	55.9
Deaths under 1 month	4	1	3	10	2	8	22.8	13.2	28.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	2	0	2	11.8	-	25.6
Tbc., respiratory (001-008)	1	0	1	2	0	2	11.8	-	25.6
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	7	3	18	14	4	105.9	152.2	51.3
Diabetes (260)	2	1	1	3	1	2	17.6	10.9	25.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	4	11	26	13	13	152.9	141.3	166.7
Heart diseases (410-443)	39	21	18	56	32	24	329.4	347.8	307.7
Arteriosclerosis (450)	1	1	0	3	1	2	17.6	10.9	25.6
Influenza (480-483)	3	0	3	3	0	3	17.6	-	38.5
#Pneumonia (490-493)	1	0	1	6	3	3	35.3	32.6	38.5
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	7	1	6	8	2	6	47.0	21.7	76.9
Chronic nephritis, etc. (592-594)	2	0	2	2	1	1	11.8	10.9	12.8
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	3	2	1	6	3	3	35.3	32.6	38.5
Birth inj., atelect. (760-762)	1	0	1	4	0	4	9.1	-	14.0
#Other dis. early inf. (763-776)	0	0	0	0	0	0	-	-	-
Violent & acc. deaths (800-999)	7	1	6	12	4	8	70.6	43.5	102.6
Acc. deaths, total (800-962)	5	1	4	10	4	6	58.8	43.5	76.9
Motor vehicle acc. (810-835)	3	0	3	4	2	2	23.5	21.7	25.6
Ill-defined (780-795)	14	3	11	16	3	13	94.1	32.6	166.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JEFFERSON COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	117	0	117	320	40	280	32.0	14.3	38.9
Fetal deaths	9	0	9	16	1	15	50.0	25.0	53.6
Deaths from all causes	87	21	66	128	33	95	12.8	11.8	13.2
Deaths under 1 year	5	0	5	13	1	12	40.6	25.0	42.8
Deaths under 1 month	3	0	3	8	1	7	25.0	25.0	25.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	7	3	4	10	4	6	100.0	142.8	83.3
Diabetes (260)	2	0	2	4	0	4	40.0	-	55.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	13	3	10	17	4	13	170.0	142.8	180.6
Heart diseases (410-443)	29	10	19	39	15	24	390.0	535.7	333.3
Arteriosclerosis (450)	2	2	0	2	2	0	20.0	71.4	-
Influenza (480-483)	2	0	2	2	0	2	20.0	-	27.8
#Pneumonia (490-493)	5	0	5	5	0	5	50.0	-	69.4
#Diarrhea & enteritis (571)	1	0	1	3	1	2	30.0	35.7	27.8
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	0	1	10.0	-	13.9
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	3	0	3	30.0	-	41.7
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	2	0	2	6	1	5	18.8	25.0	17.8
Violent & acc. deaths (800-999)	8	0	8	15	2	13	150.0	71.4	180.6
Acc. deaths, total (800-962)	7	0	7	14	2	12	140.0	71.4	166.7
Motor vehicle acc. (810-835)	0	0	0	3	1	2	30.0	35.7	27.8
Ill-defined (780-795)	7	0	7	7	0	7	70.0	-	97.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JEFFERSON DAVIS COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		Non-white
				Total	White	Non-white	Total	White	
Live births	346	147	199	390	129	261	22.9	13.2	36.2
Fetal deaths	11	3	8	13	3	10	33.3	23.2	38.3
Deaths from all causes	136	64	72	145	64	81	8.5	6.5	11.2
Deaths under 1 year	19	5	14	20	4	16	51.3	31.0	61.3
Deaths under 1 month	8	3	5	10	2	8	25.6	15.5	30.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	5.9	-	13.9
Tbc., respiratory (001-008)	0	0	0	1	0	1	5.9	-	13.9
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	4	5	13	7	6	76.5	71.4	83.3
Diabetes (260)	1	0	1	1	0	1	5.9	-	13.9
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	7	8	17	8	9	100.0	81.6	125.0
Heart diseases (410-443)	56	37	19	56	35	21	329.4	357.1	291.7
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	5	1	4	5	0	5	29.4	-	69.4
#Diarrhea & enteritis (571)	4	2	2	2	1	1	11.8	10.2	13.9
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	0	2	11.8	-	27.8
Maternal deaths (640-689)	1	0	1	1	0	1	2.6	-	3.8
Cong. malformations (750-759)	1	0	1	2	0	2	11.8	-	27.8
Birth inj., atelect. (760-762)	2	1	1	1	0	1	2.6	-	3.8
#Other dis. early inf. (763-776)	5	2	3	7	2	5	17.9	15.5	19.2
Violent & acc. deaths (800-999)	11	2	9	12	3	9	70.6	30.6	125.0
Acc. deaths, total (800-962)	10	1	9	11	2	9	64.7	20.4	125.0
Motor vehicle acc. (810-835)	5	1	4	4	2	2	23.5	20.4	27.8
Ill-defined (780-795)	12	1	11	10	1	9	58.8	10.2	125.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JONES COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	2,365	1,208	1,157	1,506	968	538	22.8	20.2	29.9
Fetal deaths	94	42	52	51	29	22	33.9	30.0	40.9
Deaths from all causes	664	410	254	557	375	182	8.4	7.8	10.1
Deaths under 1 year	99	48	51	70	40	30	46.5	41.3	55.8
Deaths under 1 month	72	39	33	47	33	14	31.2	34.1	26.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	1	2	3	1	2	4.5	2.1	11.1
Tbc., respiratory (001-008)	3	1	2	3	1	2	4.5	2.1	11.1
Syphilis (020-029)	2	1	1	1	1	0	1.5	2.1	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	1.5	-	5.6
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	2	1	1	1	1	0	1.5	2.1	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	80	63	17	70	56	14	106.1	116.7	77.8
Diabetes (260)	10	6	4	7	6	1	10.6	12.5	5.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	108	64	44	92	57	35	139.4	118.8	194.4
Heart diseases (410-443)	173	116	57	160	110	50	242.4	229.2	277.8
Arteriosclerosis (450)	3	3	0	2	2	0	3.0	4.2	-
Influenza (480-483)	6	4	2	6	4	2	9.1	8.3	11.1
#Pneumonia (490-493)	16	10	6	14	8	6	21.2	16.7	33.3
#Diarrhea & enteritis (571)	8	2	6	7	1	6	10.6	2.1	33.3
Acute nephritis (590, 591)	3	1	2	2	0	2	3.0	-	11.1
Chronic nephritis, etc. (592-594)	9	5	4	8	6	2	12.1	12.5	11.1
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	15	10	5	11	10	1	16.7	20.8	5.6
Birth inj., atelect. (760-762)	19	8	11	10	6	4	6.6	6.2	7.4
#Other dis. early inf. (763-776)	44	24	20	33	21	12	21.9	21.7	22.3
Violent & acc. deaths (800-999)	55	34	21	47	35	12	71.2	72.9	66.7
Acc. deaths, total (800-962)	46	28	18	38	30	8	57.6	62.5	44.4
Motor vehicle acc. (810-835)	15	12	3	15	14	1	22.7	29.2	5.6
Ill-defined (780-795)	32	15	17	29	13	16	43.9	27.1	88.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
KEMPER COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	206	37	169	294	60	234	24.5	11.3	34.9
Fetal deaths	7	0	7	9	1	8	30.6	16.7	34.2
Deaths from all causes	104	47	57	128	57	71	10.7	10.8	10.6
Deaths under 1 year	9	1	8	11	1	10	37.4	16.7	42.7
Deaths under 1 month	6	1	5	7	1	6	23.8	16.7	25.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	8.3	18.9	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	7	2	12	8	4	100.0	150.9	59.7
Diabetes (260)	2	1	1	3	1	2	25.0	18.9	29.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	9	3	6	12	5	7	100.0	94.3	104.5
Heart diseases (410-443)	30	11	19	36	15	21	300.0	283.0	313.4
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	11	7	4	12	7	5	100.0	132.1	74.6
#Diarrhea & enteritis (571)	2	0	2	2	0	2	16.7	-	29.8
Acute nephritis (590, 591)	1	1	0	1	1	0	8.3	18.9	-
Chronic nephritis, etc. (592-594)	2	2	0	2	2	0	16.7	37.7	-
Maternal deaths (640-689)	1	0	1	1	0	1	3.4	-	4.3
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	1	0	1	3.4	-	4.3
#Other dis. early inf. (763-776)	5	1	4	6	1	5	20.4	16.7	21.4
Violent & acc. deaths (800-999)	9	5	4	10	5	5	83.3	94.3	74.6
Acc. deaths, total (800-962)	6	3	3	8	4	4	66.7	75.5	59.7
Motor vehicle acc. (810-835)	3	2	1	5	3	2	41.7	56.6	29.8
Ill-defined (780-795)	14	2	12	15	2	13	125.0	37.7	194.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LAFAYETTE COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	505	259	246	524	261	263	18.7	13.7	29.2
Fetal deaths	20	7	13	20	8	12	38.2	30.6	45.6
Deaths from all causes	158	85	73	173	98	75	6.2	5.2	8.3
Deaths under 1 year	15	2	13	16	1	15	30.5	3.8	57.0
Deaths under 1 month	11	2	9	11	1	10	21.0	3.8	38.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	1	0	1	3.6	-	11.1
Tbc., respiratory (001-008)	1	0	1	1	0	1	3.6	-	11.1
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	13	9	4	17	12	5	60.7	63.2	55.6
Diabetes (260)	6	3	3	7	4	3	25.0	21.0	33.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	24	15	9	25	15	10	89.3	78.9	111.1
Heart diseases (410-443)	54	36	18	59	41	18	210.7	215.8	200.0
Arteriosclerosis (450)	5	3	2	5	3	2	17.8	15.8	22.2
Influenza (480-483)	3	2	1	3	2	1	10.7	10.5	11.1
#Pneumonia (490-493)	6	1	5	5	1	4	17.8	5.3	44.4
#Diarrhea & enteritis (571)	3	1	2	3	1	2	10.7	5.3	22.2
Acute nephritis (590, 591)	1	0	1	1	0	1	3.6	-	11.1
Chronic nephritis, etc. (592-594)	4	3	1	4	3	1	14.3	15.8	11.1
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	1	1	0	2	1	1	3.8	3.8	3.8
#Other dis. early inf. (763-776)	8	1	7	7	0	7	13.4	-	26.6
Violent & acc. deaths (800-999)	10	4	6	12	5	7	42.8	26.3	77.8
Acc. deaths, total (800-962)	10	4	6	12	5	7	42.8	26.3	77.8
Motor vehicle acc. (810-835)	3	3	0	4	3	1	14.3	15.8	11.1
Ill-defined (780-795)	8	1	7	9	2	7	32.1	10.5	77.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LAMAR COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	114	77	37	327	255	72	17.2	16.4	20.6
Fetal deaths	0	0	0	11	8	3	33.6	31.4	41.7
Deaths from all causes	74	57	17	106	81	25	5.6	5.2	7.1
Deaths under 1 year	9	6	3	16	10	6	48.9	39.2	83.3
Deaths under 1 month	7	5	2	10	6	4	30.6	23.5	55.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	5.3	6.4	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	1	0	5.3	6.4	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	7	5	2	14	11	3	73.7	71.0	85.7
Diabetes (260)	1	1	0	2	2	0	10.5	12.9	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	10	7	3	12	8	4	63.2	51.6	114.3
Heart diseases (410-443)	28	21	7	37	28	9	194.7	180.6	257.1
Arteriosclerosis (450)	0	0	0	1	1	0	5.3	6.4	-
Influenza (480-483)	3	3	0	3	3	0	15.8	19.4	-
#Pneumonia (490-493)	3	2	1	5	3	2	26.3	19.4	57.1
#Diarrhea & enteritis (571)	0	0	0	1	0	1	5.3	-	28.6
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	5.3	6.4	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	2	2	0	10.5	12.9	-
Birth inj., atelect. (760-762)	3	3	0	3	2	1	9.2	7.8	13.9
#Other dis. early inf. (763-776)	3	2	1	5	3	2	15.3	11.8	27.8
Violent & acc. deaths (800-999)	4	4	0	4	4	0	21.0	25.8	-
Acc. deaths, total (800-962)	4	4	0	4	4	0	21.0	25.8	-
Motor vehicle acc. (810-835)	4	4	0	2	2	0	10.5	12.9	-
Ill-defined (780-795)	4	1	3	4	1	3	21.0	6.4	85.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LAUDERDALE COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	2,088	1,061	1,027	1,591	851	740	27.0	23.0	33.6
Fetal deaths	98	34	64	65	22	43	40.8	25.8	58.1
Deaths from all causes	880	530	350	724	420	304	12.3	11.4	13.8
Deaths under 1 year	90	28	62	75	24	51	47.1	28.2	68.9
Deaths under 1 month	54	22	32	42	17	25	26.4	20.0	33.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	1	3	4	1	3	6.8	2.7	13.6
Tbc., respiratory (001-008)	4	1	3	4	1	3	6.8	2.7	13.6
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	2	0	2	2	0	3.4	5.4	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	1	0	2	2	0	3.4	5.4	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	2	1	1	2	1	1	3.4	2.7	4.5
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	89	60	29	76	50	26	128.8	135.1	118.2
Diabetes (260)	18	7	11	12	5	7	20.3	13.5	31.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	130	88	42	96	60	36	162.7	162.2	163.6
Heart diseases (410-443)	268	180	88	227	143	84	384.7	386.5	381.8
Arteriosclerosis (450)	8	8	0	7	7	0	11.9	18.9	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	30	13	17	28	12	16	47.4	32.4	72.7
#Diarrhea & enteritis (571)	11	2	9	8	1	7	13.6	2.7	31.8
Acute nephritis (590, 591)	1	1	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	8	6	2	7	5	2	11.9	13.5	9.1
Maternal deaths (640-689)	4	0	4	2	0	2	1.2	-	2.7
Cong. malformations (750-759)	12	10	2	10	7	3	16.9	18.9	13.6
Birth inj., atelect. (760-762)	19	6	13	13	5	8	8.2	5.9	10.8
#Other dis. early inf. (763-776)	29	11	18	24	9	15	15.1	10.6	20.3
Violent & acc. deaths (800-999)	69	45	24	58	40	18	98.3	108.1	81.8
Acc. deaths, total (800-962)	56	40	16	48	36	12	81.4	97.3	54.5
Motor vehicle acc. (810-835)	31	25	6	30	26	4	50.8	70.3	18.2
Ill-defined (780-795)	61	20	41	57	18	39	96.6	48.6	177.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LAWRENCE COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	180	23	157	256	124	132	21.3	14.8	36.7
Fetal deaths	5	0	5	9	3	6	35.2	24.2	45.4
Deaths from all causes	67	28	39	118	66	52	9.8	7.8	14.4
Deaths under 1 year	5	0	5	9	3	6	35.2	24.2	45.4
Deaths under 1 month	2	0	2	4	2	2	15.6	16.1	15.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	7	5	2	15	10	5	125.0	119.0	138.9
Diabetes (260)	1	0	1	2	1	1	16.7	11.9	27.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	12	3	9	18	9	9	150.0	107.1	250.0
Heart diseases (410-443)	19	13	6	37	26	11	308.3	309.5	305.6
Arteriosclerosis (450)	7	1	6	7	1	6	58.3	11.9	166.7
Influenza (480-483)	1	0	1	1	0	1	8.3	-	27.8
#Pneumonia (490-493)	4	1	3	4	1	3	33.3	11.9	83.3
#Diarrhea & enteritis (571)	0	0	0	1	0	1	8.3	-	27.8
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	8.3	11.9	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	2	2	0	7.8	16.1	-
#Other dis. early inf. (763-776)	1	0	1	1	0	1	3.9	-	7.6
Violent & acc. deaths (800-999)	5	2	3	10	7	3	83.3	83.3	83.3
Acc. deaths, total (800-962)	5	2	3	10	7	3	83.3	83.3	83.3
Motor vehicle acc. (810-835)	0	0	0	6	4	2	50.0	47.6	55.6
Ill-defined (780-795)	5	1	4	5	1	4	41.7	11.9	111.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LEAKE COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	345	147	198	411	168	243	18.7	12.9	27.0
Fetal deaths	6	0	6	12	1	11	29.2	6.0	45.3
Deaths from all causes	160	95	65	188	112	76	8.5	8.6	8.4
Deaths under 1 year	10	3	7	16	6	10	38.9	35.7	41.2
Deaths under 1 month	6	2	4	12	6	6	29.2	35.7	24.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	0	2	2	0	2	9.1	-	22.2
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	4.5	-	11.1
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	13	4	25	17	8	113.36	130.8	88.9
Diabetes (260)	3	3	0	5	4	1	22.7	30.8	11.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	24	13	11	28	17	11	127.3	130.8	122.2
Heart diseases (410-443)	58	39	19	62	43	19	281.8	330.8	211.1
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	2	0	2	2	0	2	9.1	-	22.2
#Pneumonia (490-493)	3	2	1	3	2	1	13.6	15.4	11.1
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	0	1	1	0	1	4.5	-	11.1
Chronic nephritis, etc. (592-594)	3	2	1	4	3	1	18.2	23.1	11.1
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	1	0	1	5	2	3	12.2	11.9	12.3
#Other dis. early inf. (763-776)	4	2	2	5	3	2	12.2	17.8	8.2
Violent & acc. deaths (800-999)	19	13	6	16	8	8	72.7	61.5	88.9
Acc. deaths, total (800-962)	14	10	4	11	5	6	50.0	38.5	66.7
Motor vehicle acc. (810-835)	7	7	0	4	2	2	18.2	15.4	22.2
Ill-defined (780-795)	6	0	6	7	0	7	31.8	-	77.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LEE COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,135	830	305	976	658	318	22.2	20.6	26.5
Fetal deaths	27	16	11	22	11	11	22.5	16.7	34.6
Deaths from all causes	364	257	107	360	252	108	8.2	7.9	9.0
Deaths under 1 year	36	24	12	30	19	11	30.7	28.9	34.6
Deaths under 1 month	24	17	7	20	13	7	20.5	19.8	22.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	2	1	1	4.5	3.1	8.3
Tbc., respiratory (001-008)	1	1	0	2	1	1	4.5	3.1	8.3
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	59	45	14	60	46	14	136.4	143.8	116.7
Diabetes (260)	2	2	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	54	29	25	54	32	22	122.7	100.0	183.3
Heart diseases (410-443)	121	92	29	120	91	29	272.7	284.4	241.7
Arteriosclerosis (450)	2	1	1	2	1	1	4.5	3.1	8.3
Influenza (480-483)	2	1	1	2	1	1	4.5	3.1	8.3
#Pneumonia (490-493)	16	10	6	17	10	7	38.6	31.2	58.3
#Diarrhea & enteritis (571)	1	0	1	1	0	1	2.3	-	8.3
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	2	0	1	1	0	2.3	3.1	-
Maternal deaths (640-689)	1	1	0	0	0	0	-	-	-
Cong. malformations (750-759)	6	3	3	4	1	3	9.1	3.1	25.0
Birth inj., atelect. (760-762)	15	10	5	13	8	5	13.3	12.2	15.7
#Other dis. early inf. (763-776)	7	4	3	6	3	3	6.1	4.6	9.4
Violent & acc. deaths (800-999)	31	25	6	28	22	6	63.6	68.8	50.0
Acc. deaths, total (800-962)	24	19	5	22	17	5	50.0	53.1	41.7
Motor vehicle acc. (810-835)	15	15	0	12	12	0	27.3	37.5	-
Ill-defined (780-795)	11	4	7	13	6	7	29.5	18.8	58.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LEFLORE COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,557	395	1,162	1,522	340	1,182	38.0	26.2	43.8
Fetal deaths	108	37	71	94	27	67	61.8	79.4	56.7
Deaths from all causes	611	243	368	538	166	372	13.4	12.8	13.8
Deaths under 1 year	79	8	71	78	8	70	51.2	23.5	59.2
Deaths under 1 month	46	8	38	45	8	37	29.6	23.5	31.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	2.5	7.7	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	2.5	7.7	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	3	0	3	3	0	3	7.5	-	11.1
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	2.5	-	3.7
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	62	32	30	48	21	27	120.0	161.5	100.0
Diabetes (260)	4	1	3	4	1	3	10.0	7.7	11.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	98	49	49	80	28	52	200.0	215.4	192.6
Heart diseases (410-443)	138	77	61	124	59	65	310.0	453.8	240.7
Arteriosclerosis (450)	6	4	2	4	1	3	10.0	7.7	11.1
Influenza (480-483)	5	1	4	5	1	4	12.5	7.7	14.8
#Pneumonia (490-493)	24	8	16	20	3	17	50.0	23.1	63.0
#Diarrhea & enteritis (571)	8	1	7	8	1	7	20.0	7.7	25.9
Acute nephritis (590, 591)	1	1	0	2	2	0	5.0	15.4	-
Chronic nephritis, etc. (592-594)	8	1	7	8	0	8	20.0	-	29.6
Maternal deaths (640-689)	2	0	2	1	0	1	0.6	-	0.8
Cong. malformations (750-759)	3	0	3	3	0	3	7.5	-	11.1
Birth inj., atelect. (760-762)	6	2	4	6	2	4	3.9	5.9	3.4
#Other dis. early inf. (763-776)	32	6	26	32	6	26	21.0	17.6	22.0
Violent & acc. deaths (800-999)	46	21	25	37	14	23	92.5	107.7	85.2
Acc. deaths, total (800-962)	33	15	18	25	9	16	62.5	69.2	59.2
Motor vehicle acc. (810-835)	11	6	5	8	4	4	20.0	30.8	14.8
Ill-defined (780-795)	95	8	87	92	5	87	230.0	38.5	322.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LINCOLN COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	697	421	276	671	369	302	28.0	23.1	37.8
Fetal deaths	27	13	14	26	9	17	38.7	24.4	56.3
Deaths from all causes	278	157	121	293	166	127	12.2	10.4	15.9
Deaths under 1 year	40	9	31	38	7	31	56.6	19.0	102.6
Deaths under 1 month	20	8	12	18	6	12	26.8	16.3	39.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	3	1	2	12.5	6.2	25.0
Tbc., respiratory (001-008)	0	0	0	3	1	2	12.5	6.2	25.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	29	23	6	35	25	10	145.8	156.2	125.0
Diabetes (260)	1	1	0	1	1	0	4.2	6.2	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	26	13	13	30	17	13	125.0	106.2	162.5
Heart diseases (410-443)	125	84	41	128	87	41	533.3	543.8	512.5
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	16	1	15	16	1	15	66.7	6.2	187.5
#Diarrhea & enteritis (571)	4	0	4	4	0	4	16.7	-	50.0
Acute nephritis (590, 591)	1	1	0	1	1	0	4.2	6.2	-
Chronic nephritis, etc. (592-594)	4	3	1	5	4	1	20.8	25.0	12.5
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	8	5	3	7	4	3	10.4	10.8	9.9
#Other dis. early inf. (763-776)	10	3	7	9	2	7	13.4	5.4	23.2
Violent & acc. deaths (800-999)	23	9	14	21	9	12	87.5	56.2	150.0
Acc. deaths, total (800-962)	20	8	12	18	8	10	75.0	50.0	125.0
Motor vehicle acc. (810-935)	13	7	6	9	6	3	37.5	37.5	37.5
Ill-defined (780-795)	18	4	14	16	4	12	66.7	25.0	150.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LOWNDES COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,176	581	595	1,158	558	600	28.2	24.3	33.3
Fetal deaths	35	16	19	38	19	19	32.8	34.0	31.7
Deaths from all causes	397	199	198	388	192	196	9.5	8.3	10.9
Deaths under 1 year	44	12	32	39	10	29	33.7	17.9	48.3
Deaths under 1 month	21	10	11	18	8	10	15.5	14.3	16.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	1	3	6	3	3	14.6	13.0	16.7
Tbc., respiratory (001-008)	4	1	3	6	3	3	14.6	13.0	16.7
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	43	26	17	47	31	16	114.6	134.8	88.9
Diabetes (260)	5	4	1	4	2	2	9.8	8.7	11.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	47	22	25	48	23	25	117.1	100.0	138.9
Heart diseases (410-443)	135	83	52	132	79	53	322.0	343.5	294.4
Arteriosclerosis (450)	7	4	3	7	4	3	17.1	17.4	16.7
Influenza (480-483)	2	0	2	2	0	2	4.9	-	11.1
#Pneumonia (490-493)	14	3	11	14	4	10	34.1	17.4	55.6
#Diarrhea & enteritis (571)	4	1	3	4	1	3	9.8	4.3	16.7
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	5	3	2	4	2	2	9.8	8.7	11.1
Maternal deaths (640-689)	1	0	1	0	0	0	-	-	-
Cong. malformations (750-759)	2	1	1	1	1	0	2.4	4.3	-
Birth inj., atelect. (760-762)	9	6	3	9	6	3	7.8	10.8	5.0
#Other dis. early inf. (763-776)	10	1	9	8	0	8	6.9	-	13.3
Violent & acc. deaths (800-999)	37	24	13	25	13	12	61.0	56.5	66.7
Acc. deaths, total (800-962)	36	23	13	23	12	11	56.1	52.2	61.1
Motor vehicle acc. (810-835)	17	15	2	5	4	1	12.2	17.4	5.6
Ill-defined (780-795)	47	7	40	48	9	39	117.1	39.1	216.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MADISON COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,048	181	867	1,033	184	849	41.3	30.7	44.7
Fetal deaths	55	5	50	58	6	52	56.1	32.6	61.2
Deaths from all causes	358	121	237	367	109	258	14.7	18.2	13.6
Deaths under 1 year	49	5	44	59	8	51	57.1	43.5	60.1
Deaths under 1 month	28	4	24	38	7	31	36.8	38.0	36.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	4	1	3	16.0	16.7	15.8
Tbc., respiratory (001-008)	1	0	1	4	1	3	16.0	16.7	15.8
Syphilis (020-029)	1	0	1	1	0	1	4.0	-	5.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	0	1	4.0	-	5.3
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	4.0	-	5.3
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	40	21	19	36	15	21	144.0	250.0	110.5
Diabetes (260)	6	2	4	5	1	4	20.0	16.7	21.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	53	22	31	49	19	30	196.0	316.7	157.9
Heart diseases (410-443)	95	33	62	97	31	66	388.0	516.7	347.4
Arteriosclerosis (450)	2	1	1	2	1	1	8.0	16.7	5.3
Influenza (480-483)	4	1	3	4	1	3	16.0	16.7	15.8
#Pneumonia (490-493)	12	2	10	13	2	11	52.0	33.3	57.9
#Diarrhea & enteritis (571)	5	0	5	5	0	5	20.0	-	26.3
Acute nephritis (590, 591)	1	0	1	1	0	1	4.0	-	5.3
Chronic nephritis, etc. (592-594)	6	3	3	6	3	3	24.0	50.0	15.8
Maternal deaths (640-689)	2	0	2	1	0	1	1.0	-	1.2
Cong. malformations (750-759)	3	0	3	2	0	2	8.0	-	10.5
Birth inj., atelect. (760-762)	3	2	1	5	3	2	4.8	16.3	2.4
#Other dis. early inf. (763-776)	20	2	18	27	4	23	26.1	21.7	27.1
Violent & acc. deaths (800-999)	28	11	17	29	8	21	116.0	133.3	110.5
Acc. deaths, total (800-962)	25	10	15	25	7	18	100.0	116.7	94.7
Motor vehicle acc. (810-835)	9	6	3	7	3	4	28.0	50.0	21.0
Ill-defined (780-795)	49	1	48	49	1	48	196.0	16.7	252.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MARION COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	525	310	215	593	354	239	22.8	22.1	23.9
Fetal deaths	36	20	16	36	21	15	60.7	59.3	62.8
Deaths from all causes	217	119	98	230	129	101	8.8	8.1	10.1
Deaths under 1 year	25	10	15	26	12	14	43.8	33.9	58.6
Deaths under 1 month	14	6	8	15	7	8	25.3	19.8	33.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	3.8	-	10.0
Tbc., respiratory (001-008)	1	0	1	1	0	1	3.8	-	10.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	1	1	0	3.8	6.2	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	19	8	11	19	9	10	73.1	56.2	100.0
Diabetes (260)	3	2	1	4	2	2	15.4	12.5	20.0
Pellagra (281)	1	0	1	1	0	1	3.8	-	10.0
Intracranial lesions (330-334)	46	27	19	40	21	19	153.8	131.2	190.0
Heart diseases (410-443)	59	37	22	64	42	22	246.2	262.5	220.0
Arteriosclerosis (450)	2	2	0	2	2	0	7.7	12.5	-
Influenza (480-483)	1	1	0	2	1	1	7.7	6.2	10.0
#Pneumonia (490-493)	5	3	2	5	3	2	19.2	18.8	20.0
#Diarrhea & enteritis (571)	3	1	2	3	2	1	11.5	12.5	10.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	2	2	5	2	3	19.2	12.5	30.0
Maternal deaths (640-689)	1	0	1	1	0	1	1.7	-	4.2
Cong. malformations (750-759)	2	1	1	3	2	1	11.5	12.5	10.0
Birth inj., atelect. (760-762)	2	2	0	2	2	0	3.4	5.6	-
#Other dis. early inf. (763-776)	10	4	6	11	5	6	18.5	14.1	25.1
Violent & acc. deaths (800-999)	13	7	6	21	11	10	80.8	68.8	100.0
Acc. deaths, total (800-962)	11	5	6	19	9	10	73.1	56.2	100.0
Motor vehicle acc. (810-835)	6	2	4	9	4	5	34.6	25.0	50.0
Ill-defined (780-795)	21	8	13	18	6	12	69.2	37.5	120.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MARSHALL COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	719	111	608	742	137	605	32.3	21.4	36.4
Fetal deaths	45	3	42	30	4	26	40.4	29.2	43.0
Deaths from all causes	250	64	186	238	68	170	10.3	10.6	10.2
Deaths under 1 year	26	4	22	17	3	14	22.9	21.9	23.1
Deaths under 1 month	21	4	17	13	3	10	17.5	21.9	16.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	3	1	2	13.0	15.6	12.0
Tbc., respiratory (001-008)	1	0	1	3	1	2	13.0	15.6	12.0
Syphilis (020-029)	1	0	1	1	0	1	4.3	-	6.0
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	1	0	1	1	0	4.3	15.6	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	3	6	14	4	10	60.9	62.5	60.2
Diabetes (260)	5	1	4	3	1	2	13.0	15.6	12.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	38	8	30	44	13	31	191.3	203.1	186.7
Heart diseases (410-443)	91	37	54	85	34	51	369.6	531.2	307.2
Arteriosclerosis (450)	3	1	2	3	1	2	13.0	15.6	12.0
Influenza (480-483)	4	1	3	3	0	3	13.0	-	18.1
#Pneumonia (490-493)	6	0	6	5	0	5	21.7	-	30.1
#Diarrhea & enteritis (571)	2	0	2	1	0	1	4.3	-	6.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	1	1	0	4.3	15.6	-
Maternal deaths (640-689)	3	0	3	3	0	3	4.0	-	5.0
Cong. malformations (750-759)	2	0	2	4	1	3	17.4	15.6	18.1
Birth inj., atelect. (760-762)	4	1	3	3	1	2	4.0	7.3	3.3
#Other dis. early inf. (763-776)	13	3	10	7	2	5	9.4	14.6	8.3
Violent & acc. deaths (800-999)	27	3	24	22	2	20	95.6	31.2	120.5
Acc. deaths, total (800-962)	23	3	20	16	2	14	69.6	31.2	84.3
Motor vehicle acc. (810-835)	13	2	11	9	1	8	39.1	15.6	48.2
Ill-defined (780-795)	21	2	19	21	3	18	91.3	46.9	108.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MONROE COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	849	435	414	795	408	387	26.5	22.7	32.2
Fetal deaths	26	12	14	27	14	13	34.0	34.3	33.6
Deaths from all causes	331	194	137	353	204	149	11.8	11.3	12.4
Deaths under 1 year	26	7	19	24	6	18	30.2	14.7	46.5
Deaths under 1 month	18	6	12	15	4	11	18.9	9.8	28.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	2	0	2	2	0	6.7	11.1	-
Tbc., respiratory (001-008)	2	2	0	2	2	0	6.7	11.1	-
Syphilis (020-029)	1	0	1	1	0	1	3.3	-	8.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	37	26	11	46	32	14	153.3	177.8	116.7
Diabetes (260)	4	3	1	4	3	1	13.3	16.7	8.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	58	28	30	60	28	32	200.0	155.6	266.7
Heart diseases (410-443)	93	74	19	99	77	22	330.0	427.8	183.3
Arteriosclerosis (450)	5	5	0	5	5	0	16.7	27.8	-
Influenza (480-483)	3	1	2	4	1	3	13.3	5.6	25.0
#Pneumonia (490-493)	14	6	8	14	6	8	46.7	33.3	66.7
#Diarrhea & enteritis (571)	2	0	2	2	0	2	6.7	-	16.7
Acute nephritis (590, 591)	2	0	2	2	0	2	6.7	-	16.7
Chronic nephritis, etc. (592-594)	9	1	8	12	3	9	40.0	16.7	75.0
Maternal deaths (640-689)	1	0	1	1	0	1	1.2	-	2.6
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	4	2	2	1	1	0	1.2	2.4	-
#Other dis. early inf. (763-776)	13	4	9	13	3	10	16.4	7.4	25.8
Violent & acc. deaths (800-999)	22	11	11	19	7	12	63.3	38.9	100.0
Acc. deaths, total (800-962)	14	7	7	13	5	8	43.3	27.8	66.7
Motor vehicle acc. (810-835)	5	5	0	3	3	0	10.0	16.7	-
Ill-defined (780-795)	33	10	23	33	10	23	110.0	55.6	191.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

'BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MONTGOMERY COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	311	136	175	305	124	181	21.8	16.3	28.3
Fetal deaths	13	2	11	15	1	14	49.2	8.1	77.3
Deaths from all causes	118	62	56	156	87	69	11.1	11.4	10.8
Deaths under 1 year	9	3	6	11	2	9	36.1	16.1	49.7
Deaths under 1 month	7	3	4	7	2	5	23.0	16.1	27.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	7.1	-	15.6
Tbc., respiratory (001-008)	1	0	1	1	0	1	7.1	-	15.6
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	12	5	20	14	6	142.8	184.2	93.8
Diabetes (260)	2	1	1	4	2	2	28.6	26.3	31.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	16	7	9	19	9	10	135.7	118.4	156.2
Heart diseases (410-443)	48	28	20	63	42	21	450.0	552.6	328.1
Arteriosclerosis (450)	1	0	1	1	0	1	7.1	-	15.6
Influenza (480-483)	1	0	1	1	0	1	7.1	-	15.6
#Pneumonia (490-493)	4	2	2	7	3	4	50.0	39.5	62.5
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	1	0	1	7.1	-	15.6
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	1	0	1	3.3	-	5.5
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	3	1	2	3	1	2	9.8	8.1	11.0
#Other dis. early inf. (763-776)	3	2	1	2	1	1	6.6	8.1	5.5
Violent & acc. deaths (800-999)	4	0	4	6	1	5	42.8	13.2	78.1
Acc. deaths, total (800-962)	3	0	3	5	1	4	35.7	13.2	62.5
Motor vehicle acc. (810-835)	1	0	1	3	1	2	21.4	13.2	31.2
Ill-defined (780-795)	6	0	6	10	1	9	71.4	13.2	140.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
NESHOBIA COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	457	254	203	482	277	205	19.3	16.9	23.8
Fetal deaths	10	4	6	17	6	11	35.3	21.7	53.6
Deaths from all causes	189	120	69	227	154	73	9.1	9.4	8.5
Deaths under 1 year	25	10	15	23	8	15	47.7	28.9	73.2
Deaths under 1 month	17	10	7	15	8	7	31.1	28.9	34.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	4.0	6.1	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	4.0	6.1	-
Syphilis (020-029)	1	0	1	1	0	1	4.0	-	11.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	22	17	5	27	20	7	108.0	122.0	81.4
Diabetes (260)	4	3	1	6	5	1	24.0	30.5	11.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	32	20	12	36	26	10	144.0	158.5	116.3
Heart diseases (410-443)	54	40	14	70	56	14	280.0	341.5	162.8
Arteriosclerosis (450)	0	0	0	1	1	0	4.0	6.1	-
Influenza (480-483)	1	1	0	2	2	0	8.0	12.2	-
#Pneumonia (490-493)	4	1	3	3	1	2	12.0	6.1	23.2
#Diarrhea & enteritis (571)	1	0	1	2	0	2	8.0	-	23.2
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	1	2	4	2	2	16.0	12.2	23.2
Maternal deaths (640-689)	1	0	1	1	0	1	2.1	-	4.9
Cong. malformations (750-759)	2	0	2	2	0	2	8.0	-	23.2
Birth inj., atelect. (760-762)	7	5	2	6	5	1	12.4	18.0	4.9
#Other dis. early inf. (763-776)	7	4	3	5	2	3	10.4	7.2	14.6
Violent & acc. deaths (800-999)	18	10	8	21	13	8	84.0	79.3	93.0
Acc. deaths, total (800-962)	16	9	7	19	12	7	76.0	73.2	81.4
Motor vehicle acc. (810-835)	8	7	1	8	8	0	32.0	48.8	-
Ill-defined (780-795)	11	3	8	14	4	10	56.0	24.4	116.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
NEWTON COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	388	245	143	424	204	220	23.6	17.0	36.7
Fetal deaths	41	22	19	40	19	21	94.3	93.1	95.4
Deaths from all causes	213	141	72	222	143	79	12.3	11.9	13.2
Deaths under 1 year	18	6	12	22	7	15	51.9	34.3	68.2
Deaths under 1 month	9	5	4	12	5	7	28.3	24.5	31.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	29	17	12	29	17	12	161.1	141.7	200.0
Diabetes (260)	4	2	2	5	2	3	27.8	16.7	50.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	27	22	5	29	22	7	161.1	183.3	116.7
Heart diseases (410-443)	68	53	15	62	48	14	344.4	400.0	233.3
Arteriosclerosis (450)	6	4	2	4	3	1	22.2	25.0	16.7
Influenza (480-483)	1	1	0	0	0	0	-	-	-
#Pneumonia (490-493)	6	4	2	3	1	2	16.7	8.3	33.3
#Diarrhea & enteritis (571)	1	0	1	3	1	2	16.7	8.3	33.3
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	11.1	8.3	16.7
Maternal deaths (640-689)	0	0	0	1	0	1	2.4	-	4.5
Cong. malformations (750-759)	3	2	1	4	3	1	22.2	25.0	16.7
Birth inj., atelect. (760-762)	2	2	0	4	2	2	9.4	9.8	9.1
#Other dis. early inf. (763-776)	4	0	4	5	0	5	11.8	-	22.7
Violent & acc. deaths (800-999)	12	8	4	18	13	5	100.0	108.3	83.3
Acc. deaths, total (800-962)	9	5	4	15	10	5	83.3	83.3	83.3
Motor vehicle acc. (810-835)	7	4	3	10	6	4	55.6	50.0	66.7
Ill-defined (780-795)	22	7	15	22	7	15	122.2	58.3	250.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
NOXUBEE COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	550	103	447	519	87	432	37.1	15.8	50.8
Fetal deaths	32	6	26	31	4	27	59.7	46.0	62.5
Deaths from all causes	190	53	137	196	56	140	14.0	10.2	16.5
Deaths under 1 year	21	3	18	23	3	20	44.3	34.5	46.3
Deaths under 1 month	10	2	8	11	2	9	21.2	23.0	20.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	7.1	-	11.8
Tbc., respiratory (001-008)	1	0	1	1	0	1	7.1	-	11.8
Syphilis (020-029)	1	0	1	1	0	1	7.1	-	11.8
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	21	8	13	23	10	13	164.3	181.8	152.9
Diabetes (260)	5	1	4	5	2	3	35.7	36.4	35.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	32	9	23	31	8	23	221.4	145.4	270.6
Heart diseases (410-443)	60	23	37	59	20	39	421.4	363.6	458.8
Arteriosclerosis (450)	2	0	2	3	1	2	21.4	18.2	23.5
Influenza (480-483)	1	0	1	1	0	1	7.1	-	11.8
#Pneumonia (490-493)	6	1	5	7	1	6	50.0	18.2	70.6
#Diarrhea & enteritis (571)	1	0	1	1	0	1	7.1	-	11.8
Acute nephritis (590, 591)	1	0	1	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	2	1	1	14.3	18.2	11.8
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	1	1	0	7.1	18.2	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	6	1	5	7	1	6	13.5	11.5	13.9
Violent & acc. deaths (800-999)	11	3	8	12	2	10	85.7	36.4	117.6
Acc. deaths, total (800-962)	9	3	6	10	2	8	71.4	36.4	94.1
Motor vehicle acc. (810-835)	2	1	1	3	1	2	21.4	18.2	23.5
Ill-defined (780-795)	29	1	28	29	1	28	207.1	18.2	329.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
OKTIBBEHA COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	824	362	462	759	312	447	28.1	20.8	37.2
Fetal deaths	25	8	17	22	5	17	29.0	16.0	38.0
Deaths from all causes	202	82	120	214	91	123	7.9	6.1	10.2
Deaths under 1 year	22	7	15	20	6	14	26.4	19.2	31.3
Deaths under 1 month	16	7	9	15	6	9	19.8	19.2	20.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	18	7	11	24	11	13	88.9	73.3	108.3
Diabetes (260)	6	4	2	7	5	2	25.9	33.3	16.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	46	17	29	47	21	26	174.1	140.0	216.7
Heart diseases (410-443)	63	31	32	62	28	34	229.6	186.7	283.3
Arteriosclerosis (450)	2	2	0	3	3	0	11.1	20.0	-
Influenza (480-483)	3	0	3	3	0	3	11.1	-	25.0
#Pneumonia (490-493)	4	2	2	4	2	2	14.8	13.3	16.7
#Diarrhea & enteritis (571)	4	0	4	3	0	3	11.1	-	25.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	0	2	3	0	3	11.1	-	25.0
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	5	0	5	5	0	5	18.5	-	11.2
Birth inj., atelect. (760-762)	1	1	0	1	1	0	1.3	3.2	-
#Other dis. early inf. (763-776)	10	5	5	10	5	5	13.2	16.0	11.2
Violent & acc. deaths (800-999)	11	5	6	11	5	6	40.7	33.3	50.0
Acc. deaths, total (800-962)	9	3	6	9	3	6	33.3	20.0	50.0
Motor vehicle acc. (810-835)	4	1	3	4	0	4	14.8	-	33.3
Ill-defined (780-795)	12	2	10	15	2	13	55.6	13.3	108.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PANOLA COUNTY, MISSISSIPPI, 1958

Shown By Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	798	254	544	831	264	567	28.6	20.3	35.4
Fetal deaths	28	4	24	36	6	30	43.3	22.7	52.9
Deaths from all causes	299	125	174	340	139	201	11.7	10.7	12.6
Deaths under 1 year	32	3	29	37	5	32	44.5	18.9	56.4
Deaths under 1 month	16	3	13	22	5	17	26.5	18.9	30.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	3.4	-	6.2
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	1	1	0	3.4	7.7	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	0	2	3	0	3	10.3	-	18.8
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	3.4	-	6.2
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	33	19	14	48	25	23	165.5	192.3	143.8
Diabetes (260)	1	0	1	1	0	1	3.4	-	6.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	45	19	26	50	22	28	172.4	169.2	175.0
Heart diseases (410-443)	103	56	47	108	56	52	372.4	430.8	325.0
Arteriosclerosis (450)	3	1	2	3	1	2	10.3	7.7	12.5
Influenza (480-483)	5	3	2	5	3	2	17.2	23.1	12.5
#Pneumonia (490-493)	8	2	6	12	4	8	41.4	30.8	50.0
#Diarrhea & enteritis (571)	2	0	2	1	0	1	3.4	-	6.2
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	5	3	2	6	3	3	20.7	23.1	18.8
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	0	2	4	1	3	13.8	7.7	18.8
Birth inj., atelect. (760-762)	8	2	6	9	2	7	10.8	7.6	12.3
#Other dis. early inf. (763-776)	7	1	6	9	2	7	10.8	7.6	12.3
Violent & acc. deaths (800-999)	24	11	13	20	8	12	69.0	61.5	75.0
Acc. deaths, total (800-962)	21	9	12	16	6	10	55.2	46.2	62.5
Motor vehicle acc. (810-835)	7	4	3	6	4	2	20.7	30.8	12.5
Ill-defined (780-795)	30	2	28	29	2	27	100.0	15.4	168.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PEARL RIVER COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	686	497	189	582	393	189	20.8	19.6	23.6
Fetal deaths	27	18	9	27	16	11	46.4	40.7	58.2
Deaths from all causes	197	140	57	210	154	56	7.5	7.7	7.0
Deaths under 1 year	22	11	11	22	12	10	37.8	30.5	52.9
Deaths under 1 month	16	9	7	19	11	8	32.6	28.0	42.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	3.6	5.0	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	3.6	5.0	-
Syphilis (020-029)	0	0	0	1	0	1	3.6	-	12.5
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	1	0	1	1	0	3.6	5.0	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	31	24	7	32	24	8	114.3	120.0	100.0
Diabetes (260)	2	1	1	2	1	1	7.1	5.0	12.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	10	5	5	10	7	3	35.7	35.0	37.5
Heart diseases (410-443)	58	49	9	57	48	9	203.6	240.0	112.5
Arteriosclerosis (450)	6	5	1	6	5	1	21.4	25.0	12.5
Influenza (480-483)	2	2	0	2	2	0	7.1	10.0	-
#Pneumonia (490-493)	4	2	2	4	2	2	14.3	10.0	25.0
#Diarrhea & enteritis (571)	1	0	1	1	1	0	3.6	5.0	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	1	2	2	1	1	7.1	5.0	12.5
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	1	1	0	3.6	5.0	-
Birth inj., atelect. (760-762)	8	5	3	9	6	3	15.5	15.3	15.9
#Other dis. early inf. (763-776)	5	3	2	7	4	3	12.0	10.2	15.9
Violent & acc. deaths (800-999)	34	21	13	41	26	15	146.4	130.0	187.5
Acc. deaths, total (800-962)	29	19	10	35	23	12	125.0	115.0	150.0
Motor vehicle acc. (810-835)	11	9	2	15	14	1	53.6	70.0	12.5
Ill-defined (780-795)	10	7	3	8	7	1	28.6	35.0	12.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PERRY COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number		Non-white	Total	White	Non-white
				Total	White				
Live births	79	58	21	220	158	62	28.6	28.7	28.2
Fetal deaths	1	0	1	10	7	3	45.4	44.3	48.4
Deaths from all causes	48	29	19	89	53	36	11.6	9.6	16.4
Deaths under 1 year	1	0	1	4	0	4	18.2	-	64.5
Deaths under 1 month	0	0	0	3	0	3	13.6	-	48.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	3	2	1	39.0	36.4	45.4
Tbc., respiratory (001-008)	1	0	1	3	2	1	39.0	36.4	45.4
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	8	6	2	11	9	2	142.8	163.6	90.9
Diabetes (260)	1	0	1	3	1	2	39.0	18.2	90.9
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	7	8	22	12	10	285.7	218.2	454.5
Heart diseases (410-443)	14	9	5	28	17	11	363.6	309.1	500.0
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	0	0	0	0	0	0	-	-	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	13.0	-	45.4
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	1	0	1	4.5	-	16.1
#Other dis. early inf. (763-776)	0	0	0	2	0	2	9.1	-	32.2
Violent & acc. deaths (800-999)	6	6	0	6	3	3	77.9	54.5	136.4
Acc. deaths, total (800-962)	6	6	0	6	3	3	77.9	54.5	136.4
Motor vehicle acc. (810-835)	2	2	0	1	0	1	13.0	-	45.4
Ill-defined (780-795)	0	0	0	1	1	0	13.0	18.2	-

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PIKE COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,021	537	484	895	407	488	28.9	22.6	37.5
Fetal deaths	36	6	30	31	4	27	34.6	9.8	55.3
Deaths from all causes	397	226	171	399	216	183	12.9	12.0	14.1
Deaths under 1 year	38	14	24	35	11	24	39.1	27.0	49.2
Deaths under 1 month	27	13	14	24	10	14	26.8	24.6	28.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	2	2	5	2	3	16.1	11.1	23.1
Tbc., respiratory (001-008)	3	1	2	4	1	3	12.9	5.6	23.1
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	3.2	-	7.7
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	3.2	-	7.7
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	43	32	11	45	30	15	145.2	166.7	115.4
Diabetes (260)	4	1	3	6	2	4	19.4	11.1	30.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	55	31	24	52	27	25	167.7	150.0	192.3
Heart diseases (410-443)	126	83	43	130	83	47	419.4	461.1	361.5
Arteriosclerosis (450)	7	5	2	8	6	2	25.8	33.3	15.4
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	17	10	7	15	8	7	48.4	44.4	53.8
#Diarrhea & enteritis (571)	2	0	2	2	0	2	6.4	-	15.4
Acute nephritis (590, 591)	2	0	2	3	1	2	9.7	5.6	15.4
Chronic nephritis, etc. (592-594)	6	0	6	6	0	6	19.4	-	46.2
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	8	6	2	5	3	2	16.1	16.7	15.4
Birth inj., atelect. (760-762)	3	1	2	3	1	2	3.4	2.4	4.1
#Other dis. early inf. (763-776)	19	8	11	18	7	11	20.1	17.2	22.5
Violent & acc. deaths (800-999)	26	11	15	24	8	16	77.4	44.4	123.1
Acc. deaths, total (800-962)	20	6	14	19	4	15	61.3	22.2	115.4
Motor vehicle acc. (810-835)	11	4	7	10	3	7	32.2	16.7	53.8
Ill-defined (780-795)	37	13	24	35	13	22	112.9	72.2	169.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PONTOTOC COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	292	216	76	301	229	72	14.3	14.3	14.4
Fetal deaths	15	9	6	14	9	5	46.5	39.3	69.4
Deaths from all causes	165	125	40	191	148	43	9.1	9.2	8.6
Deaths under 1 year	9	6	3	11	8	3	36.5	34.9	41.7
Deaths under 1 month	6	4	2	9	7	2	29.9	30.6	27.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	4.8	6.2	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	4.8	6.2	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	1	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	19	13	6	23	17	6	109.5	106.2	120.0
Diabetes (260)	4	4	0	3	3	0	14.3	18.8	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	21	16	5	22	16	6	104.8	100.0	120.0
Heart diseases (410-443)	66	55	11	72	61	11	342.8	381.2	220.0
Arteriosclerosis (450)	4	3	1	4	3	1	19.0	18.8	20.0
Influenza (480-483)	2	1	1	2	1	1	9.5	6.2	20.0
#Pneumonia (490-493)	9	8	1	10	9	1	47.6	56.2	20.0
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	4	3	1	4	4	0	19.0	25.0	-
Birth inj., atelect. (760-762)	2	1	1	5	3	2	16.6	13.1	27.8
#Other dis. early inf. (763-776)	2	1	1	2	1	1	6.6	4.4	13.9
Violent & acc. deaths (800-999)	12	8	4	20	14	6	95.2	87.5	120.0
Acc. deaths, total (800-962)	11	7	4	18	12	6	85.7	75.0	120.0
Motor vehicle acc. (810-835)	3	2	1	7	5	2	33.3	31.2	40.0
Ill-defined (780-795)	7	3	4	6	2	4	28.6	12.5	80.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar¹ to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PRENTISS COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	457	396	61	395	338	57	20.8	21.1	19.0
Fetal deaths	9	9	0	8	8	0	20.2	23.7	-
Deaths from all causes	164	148	16	148	133	15	7.8	8.3	5.0
Deaths under 1 year	5	4	1	5	4	1	12.6	11.8	17.5
Deaths under 1 month	5	4	1	5	4	1	12.6	11.8	17.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	5.3	6.2	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	5.3	6.2	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	24	24	0	21	21	0	110.5	131.2	-
Diabetes (260)	1	1	0	1	1	0	5.3	6.2	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	32	26	6	26	20	6	136.8	125.0	200.0
Heart diseases (410-443)	43	40	3	38	35	3	200.2	218.8	100.0
Arteriosclerosis (450)	3	3	0	2	2	0	10.5	12.5	-
Influenza (480-483)	1	0	1	1	0	1	5.3	-	33.3
#Pneumonia (490-493)	9	9	0	7	7	0	36.8	43.8	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	3	2	1	15.8	12.5	33.3
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	1	1	0	5.3	6.2	-
Birth inj., atelect. (760-762)	3	2	1	3	2	1	7.6	5.9	17.5
#Other dis. early inf. (763-776)	1	1	0	1	1	0	2.5	3.0	-
Violent & acc. deaths (800-999)	13	13	0	14	14	0	73.7	87.5	-
Acc. deaths, total (800-962)	6	6	0	7	7	0	36.8	43.8	-
Motor vehicle acc. (810-835)	5	5	0	5	5	0	26.3	31.2	-
Ill-defined (780-795)	4	4	0	6	6	0	31.6	37.5	-

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
QUITMAN COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	780	202	578	728	179	549	38.3	26.7	44.6
Fetal deaths	19	2	17	17	2	15	23.4	11.2	27.3
Deaths from all causes	183	47	136	230	71	159	12.1	10.6	12.9
Deaths under 1 year	22	3	19	28	4	24	38.5	22.3	43.7
Deaths under 1 month	16	3	13	20	4	16	27.5	22.3	29.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	3	1	2	15.8	14.9	16.3
Tbc., respiratory (001-008)	1	0	1	2	1	1	10.5	14.9	8.1
Syphilis (020-029)	0	0	0	1	0	1	5.3	-	8.1
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	33	12	21	38	17	21	200.0	253.7	170.7
Diabetes (260)	1	0	1	4	2	2	21.0	29.8	16.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	18	6	12	23	7	16	121.0	104.5	130.1
Heart diseases (410-443)	49	14	35	56	21	35	294.7	313.4	284.6
Arteriosclerosis (450)	2	1	1	3	2	1	15.8	29.8	8.1
Influenza (480-483)	4	0	4	4	0	4	21.0	-	32.5
#Pneumonia (490-493)	4	0	4	7	0	7	36.8	-	56.9
#Diarrhea & enteritis (571)	0	0	0	1	0	1	5.3	-	8.1
Acute nephritis (590, 591)	1	0	1	1	0	1	5.3	-	8.1
Chronic nephritis, etc. (592-594)	3	0	3	3	0	3	15.8	-	24.4
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	3	1	2	15.8	14.9	16.3
Birth inj., atelect. (760-762)	3	1	2	3	1	2	4.1	5.6	3.6
#Other dis. early inf. (763-776)	7	1	6	10	1	9	13.7	5.6	16.4
Violent & acc. deaths (800-999)	22	8	14	27	11	16	142.1	164.2	130.1
Acc. deaths, total (800-962)	18	7	11	23	10	13	121.0	149.2	105.7
Motor vehicle acc. (810-835)	4	2	2	5	3	2	26.3	44.8	16.3
Ill-defined (780-795)	30	2	28	31	3	28	163.2	44.8	227.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
RANKIN COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	248	44	204	699	385	314	19.4	17.5	22.4
Fetal deaths	12	10	2	27	21	6	38.6	54.5	19.1
Deaths from all causes	388	176	212	226	128	98	6.3	5.8	7.0
Deaths under 1 year	13	3	10	31	13	18	44.3	33.8	57.3
Deaths under 1 month	9	2	7	23	9	14	32.9	23.4	44.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	8	3	5	0	0	0	-	-	-
Tbc., respiratory (001-008)	8	3	5	0	0	0	-	-	-
Syphilis (020-029)	2	1	1	3	1	2	8.3	4.5	14.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	8	4	4	1	0	1	2.8	-	7.1
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	27	8	19	22	16	6	61.1	72.7	42.8
Diabetes (260)	2	0	2	3	2	1	8.3	9.1	7.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	64	39	25	26	18	8	72.2	81.8	57.1
Heart diseases (410-443)	103	64	39	61	43	18	169.4	195.4	128.6
Arteriosclerosis (450)	8	4	4	4	2	2	11.1	9.1	14.3
Influenza (480-483)	2	1	1	2	1	1	5.6	4.5	7.1
#Pneumonia (490-493)	16	3	13	13	5	8	36.1	22.7	57.1
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	0	1	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	0	0	0	-	-	-
Maternal deaths (640-689)	1	0	1	2	1	1	2.9	2.6	3.2
Cong. malformations (750-759)	3	2	1	8	6	2	22.2	27.3	14.3
Birth inj., atelect. (760-762)	2	1	1	6	3	3	8.6	7.8	9.6
#Other dis. early inf. (763-776)	6	1	5	13	4	9	18.6	10.4	28.7
Violent & acc. deaths (800-999)	23	10	13	16	8	8	44.4	36.4	57.1
Acc. deaths, total (800-962)	16	7	9	12	8	4	33.3	36.4	28.6
Motor vehicle acc. (810-835)	6	5	1	6	5	1	16.7	22.7	7.1
Ill-defined (780-795)	39	8	31	23	6	17	63.9	27.3	121.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SCOTT COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	555	319	236	558	288	270	27.9	24.0	33.8
Fetal deaths	16	5	11	20	6	14	35.8	20.8	51.8
Deaths from all causes	216	116	100	239	130	109	12.0	10.8	13.6
Deaths under 1 year	27	2	25	33	4	29	59.1	13.9	107.4
Deaths under 1 month	20	2	18	23	3	20	41.2	10.4	74.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	2	2	0	10.0	16.7	-
Tbc., respiratory (001-008)	1	1	0	2	2	0	10.0	16.7	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	1	0	1	1	0	1	5.0	-	12.5
Whooping cough (056)	2	0	2	3	0	3	15.0	-	37.5
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	22	14	8	25	16	9	125.0	133.3	112.5
Diabetes (260)	2	2	0	2	2	0	10.0	16.7	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	38	22	16	39	21	18	195.0	175.0	225.0
Heart diseases (410-443)	59	45	14	64	49	15	320.0	408.3	187.5
Arteriosclerosis (450)	1	0	1	1	0	1	5.0	-	12.5
Influenza (480-483)	2	0	2	2	0	2	10.0	-	25.0
#Pneumonia (490-493)	4	0	4	3	0	3	15.0	-	37.5
#Diarrhea & enteritis (571)	1	1	0	1	1	0	5.0	8.3	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	10.0	8.3	12.5
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	1	1	3	2	1	15.0	16.7	12.5
Birth inj., atelect. (760-762)	2	0	2	5	1	4	9.0	3.5	14.8
#Other dis. early inf. (763-776)	13	1	12	13	1	12	23.3	3.5	44.4
Violent & acc. deaths (800-999)	22	13	9	22	14	8	110.0	116.7	100.0
Acc. deaths, total (800-962)	18	13	5	18	14	4	90.0	116.7	50.0
Motor vehicle acc. (810-835)	7	6	1	8	7	1	40.0	58.3	12.5
Ill-defined (780-795)	23	3	20	22	3	19	110.0	25.0	237.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SHARKEY COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	233	39	194	355	65	290	38.6	21.7	46.8
Fetal deaths	3	1	2	9	1	8	25.4	15.4	27.6
Deaths from all causes	102	26	76	116	32	84	12.6	10.7	13.5
Deaths under 1 year	16	0	16	17	1	16	47.9	15.4	55.2
Deaths under 1 month	8	0	8	9	1	8	25.4	15.4	27.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	10.9	-	16.1
Tbc., respiratory (001-008)	0	0	0	1	0	1	10.9	-	16.1
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	0	2	2	0	2	21.7	-	32.2
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	15	3	12	13	3	10	141.3	100.0	161.3
Diabetes (260)	1	1	0	1	1	0	10.9	33.3	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	11	4	7	9	4	5	97.8	133.3	80.6
Heart diseases (410-443)	21	10	11	28	10	18	304.3	333.3	290.3
Arteriosclerosis (450)	2	0	2	2	0	2	21.7	-	32.2
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	0	1	2	0	2	21.7	-	32.2
#Diarrhea & enteritis (571)	2	0	2	2	0	2	21.7	-	32.2
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	0	4	3	0	3	32.6	-	48.4
Maternal deaths (640-689)	0	0	0	1	0	1	2.8	-	3.4
Cong. malformations (750-759)	0	0	0	1	1	0	10.9	33.3	-
Birth inj., atelect. (760-762)	1	0	1	2	0	2	5.6	-	6.9
#Other dis. early inf. (763-776)	5	0	5	6	1	5	16.9	15.4	17.2
Violent & acc. deaths (800-999)	7	1	6	9	3	6	97.8	100.0	96.8
Acc. deaths, total (800-962)	5	0	5	6	1	5	65.2	33.3	80.6
Motor vehicle acc. (810-835)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	19	3	16	19	3	16	206.5	100.0	258.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SIMPSON COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	527	263	264	486	262	224	20.2	16.4	28.0
Fetal deaths	16	6	10	17	9	8	35.0	34.4	35.7
Deaths from all causes	234	146	88	228	148	80	9.5	9.2	10.0
Deaths under 1 year	11	1	10	17	5	12	35.0	19.1	53.6
Deaths under 1 month	5	1	4	9	4	5	18.5	15.3	22.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	38	24	14	1	0	1	4.2	-	12.5
Tbc., respiratory (001-008)	36	23	13	1	0	1	4.2	-	12.5
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	2	0	2	8.3	-	25.0
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	4.2	-	12.5
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	26	19	7	33	24	9	137.5	150.0	112.5
Diabetes (260)	1	1	0	1	1	0	4.2	6.2	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	28	19	9	27	17	10	112.5	106.2	125.0
Heart diseases (410-443)	73	47	26	80	55	25	333.3	343.8	312.5
Arteriosclerosis (450)	1	1	0	3	3	0	12.5	18.8	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	6	3	3	9	7	2	37.5	43.8	25.0
#Diarrhea & enteritis (571)	0	0	0	1	0	1	4.2	-	12.5
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	0	3	2	0	2	8.3	-	25.0
Maternal deaths (640-689)	1	1	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	2	2	0	8.3	12.5	-
Birth inj., atelect. (760-762)	1	0	1	5	3	2	10.3	11.4	8.9
#Other dis. early inf. (763-776)	4	1	3	4	0	4	8.2	-	17.8
Violent & acc. deaths (800-999)	21	12	9	23	15	8	95.8	93.8	100.0
Acc. deaths, total (800-962)	18	11	7	20	14	6	83.3	87.5	75.0
Motor vehicle acc. (810-835)	11	9	2	10	9	1	41.7	56.2	12.5
Ill-defined (780-795)	13	4	9	14	4	10	58.3	25.0	125.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SMITH COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	127	90	37	342	207	135	22.8	19.7	30.0
Fetal deaths	2	2	0	10	5	5	29.2	24.2	37.0
Deaths from all causes	74	62	12	133	107	26	8.9	10.2	5.8
Deaths under 1 year	4	1	3	8	4	4	23.4	19.3	29.6
Deaths under 1 month	1	0	1	5	3	2	14.6	14.5	14.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	6.7	-	22.2
Tbc., respiratory (001-008)	0	0	0	1	0	1	6.7	-	22.2
Syphilis (020-029)	1	1	0	1	1	0	6.7	9.5	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	8	1	13	12	1	86.7	114.3	22.2
Diabetes (260)	1	1	0	3	2	1	20.0	19.0	22.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	14	11	3	19	14	5	126.7	133.3	111.1
Heart diseases (410-443)	37	34	3	53	49	4	353.3	466.7	88.9
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	0	0	0	2	2	0	13.3	19.0	-
#Diarrhea & enteritis (571)	0	0	0	1	1	0	6.7	9.5	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	1	1	0	2.9	4.8	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	1	0	1	4	2	2	11.7	9.7	14.8
Violent & acc. deaths (800-999)	7	5	2	19	13	6	126.7	123.8	133.3
Acc. deaths, total (800-962)	5	3	2	15	9	6	100.0	85.7	133.3
Motor vehicle acc. (810-835)	2	2	0	8	6	2	53.3	57.1	44.4
Ill-defined (780-795)	3	1	2	3	1	2	20.0	9.5	44.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
STONE COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	159	111	48	150	100	50	25.0	22.2	33.3
Fetal deaths	11	6	5	9	4	5	60.0	40.0	100.0
Deaths from all causes	61	44	17	69	45	24	11.5	10.0	16.0
Deaths under 1 year	1	1	0	3	2	1	20.0	20.0	20.0
Deaths under 1 month	1	1	0	2	2	0	13.3	20.0	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	7	6	1	9	8	1	150.0	177.8	66.7
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	10	6	4	10	4	6	166.7	88.9	400.0
Heart diseases (410-443)	30	24	6	28	21	7	466.7	466.7	466.7
Arteriosclerosis (450)	0	0	0	1	1	0	16.7	22.2	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	0	0	0	0	0	0	-	-	-
#Diarrhea & enteritis (571)	0	0	0	1	0	1	16.7	-	66.7
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	16.7	22.2	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	2	2	0	33.3	44.4	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	1	1	0	1	1	0	6.7	10.0	-
Violent & acc. deaths (800-999)	7	5	2	8	5	3	133.3	111.1	200.0
Acc. deaths, total (800-962)	6	4	2	6	4	2	100.0	88.9	133.3
Motor vehicle acc. (810-835)	5	3	2	4	2	2	66.7	44.4	133.3
Ill-defined (780-795)	2	1	1	2	1	1	33.3	22.2	66.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SUNFLOWER COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,354	263	1,091	1,459	311	1,148	33.9	19.4	42.5
Fetal deaths	60	9	51	70	18	52	48.0	57.9	45.3
Deaths from all causes	400	95	305	460	129	331	10.7	8.1	12.2
Deaths under 1 year	68	7	61	71	9	62	48.7	28.9	54.0
Deaths under 1 month	35	5	30	39	7	32	26.7	22.5	27.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	3	2	1	7.0	12.5	3.7
Tbc., Respiratory (001-008)	2	1	1	3	2	1	7.0	12.5	3.7
Syphilis (020-029)	2	0	2	2	0	2	4.6	-	7.4
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	2.3	-	3.7
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	2.3	-	3.7
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	35	13	22	51	22	29	118.6	137.5	107.4
Diabetes (260)	6	1	5	6	1	5	14.0	6.2	18.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	47	12	35	51	18	33	118.6	112.5	122.2
Heart diseases (410-443)	98	41	57	106	43	63	246.5	268.8	233.3
Arteriosclerosis (450)	5	1	4	6	2	4	14.0	12.5	14.8
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	18	3	15	20	5	15	46.5	31.2	55.6
#Diarrhea & enteritis (571)	7	1	6	8	1	7	18.6	6.2	25.9
Acute nephritis (590, 591)	4	1	3	2	1	1	4.6	6.2	3.7
Chronic nephritis, etc. (592-594)	7	0	7	7	0	7	16.3	-	25.9
Maternal deaths (640-689)	2	0	2	3	0	3	2.0	-	2.6
Cong. malformations (750-759)	5	0	5	7	1	6	16.3	6.2	22.2
Birth inj., atelect. (760-762)	5	2	3	5	3	2	3.4	9.6	1.7
#Other dis. early inf. (763-776)	22	2	20	26	3	23	17.8	9.6	20.0
Violent & acc. deaths (800-999)	38	7	31	43	12	31	100.0	75.0	114.8
Acc. deaths, total (800-962)	25	5	20	33	10	23	76.7	62.5	85.2
Motor vehicle acc. (810-835)	4	1	3	10	5	5	23.2	31.2	18.5
Ill-defined (780-795)	64	3	61	64	3	61	148.8	18.8	225.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TALLAHATCHIE COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	790	132	658	826	174	652	37.5	19.3	50.2
Fetal deaths	32	3	29	34	3	31	41.2	17.2	47.5
Deaths from all causes	248	71	177	280	89	191	12.7	9.9	14.7
Deaths under 1 year	36	6	30	40	9	31	48.4	51.7	47.5
Deaths under 1 month	22	5	17	23	5	18	27.8	28.7	27.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	3	1	2	13.6	11.1	15.4
Tbc., respiratory (001-008)	1	1	0	3	1	2	13.6	11.1	15.4
Syphilis (020-029)	1	0	1	1	0	1	4.5	-	7.7
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	4.5	-	7.7
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	1	0	1	1	0	4.5	11.1	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	30	11	19	33	12	21	150.0	133.3	161.5
Diabetes (260)	1	0	1	2	0	2	9.1	-	15.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	33	11	22	36	15	21	163.6	166.7	161.5
Heart diseases (410-443)	70	26	44	74	29	45	336.4	322.2	346.2
Arteriosclerosis (450)	6	4	2	7	5	2	31.8	55.5	15.4
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	9	0	9	9	0	9	40.9	-	69.2
#Diarrhea & enteritis (571)	2	0	2	4	1	3	18.2	11.1	23.1
Acute nephritis (590, 591)	2	0	2	2	0	2	9.1	-	15.4
Chronic nephritis, etc. (592-594)	4	0	4	3	0	3	13.6	-	23.1
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	4	1	3	4	1	3	18.2	11.1	23.1
Birth inj., atelect. (760-762)	7	3	4	8	3	5	9.7	17.2	7.7
#Other dis. early inf. (763-776)	11	1	10	11	1	10	13.3	5.7	15.3
Violent & acc. deaths (800-999)	19	6	13	25	10	15	113.6	111.1	115.4
Acc. deaths, total (800-962)	12	6	6	16	8	8	72.7	88.9	61.5
Motor vehicle acc. (810-835)	5	5	0	5	4	1	22.7	44.4	7.7
Ill-defined (780-795)	31	2	29	33	2	31	150.0	22.2	238.5

* Crude birth and death rate per 1,000 population: Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TATE COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	450	22	428	584	138	446	34.4	15.3	55.8
Fetal deaths	18	0	18	24	0	24	41.1	-	53.8
Deaths from all causes	114	45	69	172	76	96	10.1	8.4	12.0
Deaths under 1 year	13	0	13	21	2	19	36.0	14.5	42.6
Deaths under 1 month	11	0	11	17	2	15	29.1	14.5	33.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	2	2	0	11.8	22.2	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	11	7	4	17	12	5	100.0	133.3	62.5
Diabetes (260)	1	1	0	2	1	1	11.8	11.1	12.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	11	2	9	17	2	15	100.0	22.2	187.5
Heart diseases (410-443)	49	26	23	59	33	26	347.0	366.7	325.0
Arteriosclerosis (450)	3	1	2	6	3	3	35.3	33.3	37.5
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	3	0	3	5	1	4	29.4	11.1	50.0
#Diarrhea & enteritis (571)	1	0	1	2	0	2	11.8	-	25.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	3	1	2	17.6	11.1	25.0
Maternal deaths (640-689)	1	0	1	1	0	1	1.7	-	2.2
Cong. malformations (750-759)	1	0	1	3	2	1	17.6	22.2	12.5
Birth inj., atelect. (760-762)	0	0	0	1	0	1	1.7	-	2.2
#Other dis. early inf. (763-776)	9	0	9	12	1	11	20.5	7.2	24.7
Violent & acc. deaths (800-999)	10	4	6	13	4	9	76.5	44.4	112.5
Acc. deaths, total (800-962)	6	2	4	9	2	7	52.9	22.2	87.5
Motor vehicle acc. (810-835)	3	2	1	3	2	1	17.6	22.2	12.5
Ill-defined (780-795)	8	2	6	9	2	7	52.9	22.2	87.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TIPPAH COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	348	265	83	313	234	79	18.4	18.7	17.6
Fetal deaths	11	11	0	10	9	1	31.9	38.5	12.6
Deaths from all causes	153	120	33	168	134	34	9.9	10.7	7.6
Deaths under 1 year	16	11	5	18	13	5	57.5	55.6	63.3
Deaths under 1 month	9	8	1	11	10	1	35.1	42.7	12.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	2	2	0	11.8	16.0	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	5.9	8.0	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	1	0	5.9	8.0	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	15	2	17	15	2	100.0	120.0	44.4
Diabetes (260)	2	2	0	1	1	0	5.9	8.0	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	19	14	5	22	17	5	129.4	136.0	111.1
Heart diseases (410-443)	68	53	15	76	62	14	447.0	496.0	311.1
Arteriosclerosis (450)	0	0	0	1	1	0	5.9	8.0	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	5	4	1	6	5	1	35.3	40.0	22.2
#Diarrhea & enteritis (571)	2	1	1	2	1	1	11.8	8.0	22.2
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	0	0	0	-	-	-
Maternal deaths (640-689)	1	1	0	0	0	0	-	-	-
Cong. malformations (750-759)	6	1	5	5	0	5	29.4	-	111.1
Birth inj., atelect. (760-762)	4	4	0	5	5	0	16.0	21.4	-
#Other dis. early inf. (763-776)	3	3	0	5	5	0	16.0	21.4	-
Violent & acc. deaths (800-999)	12	8	4	11	5	6	64.7	40.0	133.3
Acc. deaths, total (800-962)	10	8	2	9	5	4	52.9	40.0	88.9
Motor vehicle acc. (810-835)	5	4	1	3	1	2	17.6	8.0	44.4
Ill-defined (780-795)	3	3	0	3	3	0	17.6	24.0	-

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TISHOMINGO COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	98	90	8	237	219	18	13.9	13.3	36.0
Fetal deaths	3	3	0	5	5	0	21.1	22.8	-
Deaths from all causes	88	84	4	136	130	6	8.0	7.9	12.0
Deaths under 1 year	3	3	0	6	6	0	25.3	27.4	-
Deaths under 1 month	2	2	0	3	3	0	12.6	13.7	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	3	3	0	17.6	18.2	-
Tbc., respiratory (001-008)	0	0	0	3	3	0	17.6	18.2	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	11	11	0	21	21	0	123.5	127.3	-
Diabetes (260)	4	4	0	6	6	0	35.3	36.4	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	23	20	3	29	26	3	170.6	157.6	600.0
Heart diseases (410-443)	27	27	0	39	37	2	229.4	224.2	400.0
Arteriosclerosis (450)	3	3	0	3	3	0	17.6	18.2	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	5	5	0	6	6	0	35.3	36.4	-
#Diarrhea & enteritis (571)	1	1	0	1	1	0	5.9	6.1	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	5.9	6.1	-
Maternal deaths (640-689)	0	0	0	1	1	0	4.2	4.6	-
Cong. malformations (750-759)	0	0	0	1	1	0	5.9	6.1	-
Birth inj., atelect. (760-762)	1	1	0	1	1	0	4.2	4.6	-
#Other dis. early inf. (763-776)	0	0	0	1	1	0	4.2	4.6	-
Violent & acc. deaths (800-999)	4	4	0	8	8	0	47.0	48.5	-
Acc. deaths, total (800-962)	3	3	0	6	6	0	35.3	36.4	-
Motor vehicle acc. (810-835)	2	2	0	3	3	0	17.6	18.2	-
Ill-defined (780-795)	3	2	1	3	2	1	17.6	12.1	200.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TUNICA COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	655	63	592	661	71	590	41.3	35.5	42.1
Fetal deaths	19	2	17	19	2	17	28.7	28.2	28.8
Deaths from all causes	192	28	164	206	30	176	12.9	15.0	12.6
Deaths under 1 year	32	2	30	30	1	29	45.4	14.1	49.2
Deaths under 1 month	12	2	10	10	1	9	15.1	14.1	15.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	6.2	-	7.1
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	6.2	-	7.1
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	2	0	2	2	0	2	12.5	-	14.3
Meningocosal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	14	2	12	16	1	15	100.0	50.0	107.1
Diabetes (260)	3	0	3	3	0	3	18.8	-	21.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	20	2	18	24	3	21	150.0	150.0	150.0
Heart diseases (410-443)	68	14	54	74	16	58	462.5	800.0	414.3
Arteriosclerosis (450)	0	0	0	1	1	0	6.2	50.0	-
Influenza (480-483)	5	0	5	5	0	5	31.2	-	35.7
#Pneumonia (490-493)	8	2	6	7	2	5	43.8	100.0	35.7
#Diarrhea & enteritis (571)	3	0	3	3	0	3	18.8	-	21.4
Acute nephritis (590, 591)	4	0	4	3	0	3	18.8	-	21.4
Chronic nephritis, etc. (592-594)	8	0	8	8	0	8	50.0	-	57.1
Maternal deaths (640-689)	1	0	1	1	0	1	1.5	-	1.7
Cong. malformations (750-759)	1	0	1	1	0	1	6.2	-	7.1
Birth inj., atelect. (760-762)	4	2	2	3	1	2	4.5	14.1	3.4
#Other dis. early inf. (763-776)	9	0	9	8	0	8	12.1	-	13.6
Violent & acc. deaths (800-999)	14	4	10	12	3	9	75.0	150.0	64.3
Acc. deaths, total (800-962)	11	3	8	9	2	7	56.2	100.0	50.0
Motor vehicle acc. (810-835)	0	0	0	1	1	0	6.2	50.0	-
Ill-defined (780-795)	20	1	19	22	1	21	137.5	50.0	150.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
UNION COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	461	365	96	402	306	96	16.8	16.1	19.2
Fetal deaths	32	28	4	26	23	3	64.7	75.2	31.2
Deaths from all causes	207	161	46	205	164	41	8.5	8.6	8.2
Deaths under 1 year	18	14	4	15	12	3	37.3	39.2	31.2
Deaths under 1 month	11	9	2	9	8	1	22.4	26.1	10.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	3	3	0	12.5	15.8	-
Tbc., respiratory (001-008)	1	1	0	3	3	0	12.5	15.8	-
Syphilis (020-029)	1	1	0	1	1	0	4.2	5.3	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	21	17	4	27	23	4	112.5	121.0	80.0
Diabetes (260)	3	2	1	3	2	1	12.5	10.5	20.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	30	22	8	28	23	5	116.7	121.0	100.0
Heart diseases (410-443)	86	67	19	81	62	19	337.5	326.3	380.0
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	2	0	2	2	0	8.3	10.5	-
#Diarrhea & enteritis (571)	4	2	2	4	2	2	16.7	10.5	40.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	1	1	0	1	1	0	2.5	3.3	-
Cong. malformations (750-759)	1	1	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	6	5	1	4	4	0	10.0	13.1	-
#Other dis. early inf. (763-776)	5	4	1	5	4	1	12.4	13.1	10.4
Violent & acc. deaths (800-999)	15	13	2	15	14	1	62.5	73.7	20.0
Acc. deaths, total (800-962)	12	10	2	13	12	1	54.2	63.2	20.0
Motor vehicle acc. (810-835)	4	3	1	6	5	1	25.0	26.3	20.0
Ill-defined (780-795)	13	9	4	14	10	4	58.3	52.6	80.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WALTHALL COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	619	362	257	350	137	213	23.3	15.2	35.5
Fetal deaths	31	18	13	19	6	13	54.3	43.8	61.0
Deaths from all causes	131	87	44	117	67	50	7.8	7.4	8.3
Deaths under 1 year	22	10	12	16	4	12	45.7	29.2	56.3
Deaths under 1 month	18	8	10	14	4	10	40.0	29.2	46.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	13	10	3	13	9	4	86.7	100.0	66.7
Diabetes (260)	1	1	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	23	15	8	21	13	8	140.0	144.4	133.3
Heart diseases (410-443)	45	35	10	33	23	10	220.0	255.6	166.7
Arteriosclerosis (450)	1	0	1	1	0	1	6.7	-	16.7
Influenza (480-483)	1	1	0	2	2	0	13.3	22.2	-
#Pneumonia (490-493)	3	3	0	1	1	0	6.7	11.1	-
#Diarrhea & enteritis (571)	1	0	1	1	0	1	6.7	-	16.7
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	6.7	-	16.7
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	2	2	0	13.3	22.2	-
Birth inj., atelect. (760-762)	4	2	2	3	1	2	8.6	7.3	9.4
#Other dis. early inf. (763-776)	11	4	7	8	1	7	22.8	7.3	32.9
Violent & acc. deaths (800-999)	7	4	3	9	5	4	60.0	55.6	66.7
Acc. deaths, total (800-962)	4	1	3	7	3	4	46.7	33.3	66.7
Motor vehicle acc. (810-835)	1	1	0	4	3	1	26.7	33.3	16.7
Ill-defined (780-795)	11	6	5	12	5	7	80.0	55.6	116.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WARREN COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,467	594	873	1,165	504	661	31.5	26.5	36.7
Fetal deaths	77	20	57	45	12	33	38.6	23.8	49.9
Deaths from all causes	689	251	438	513	210	303	13.9	11.0	16.8
Deaths under 1 year	84	18	66	59	16	43	50.6	31.7	65.0
Deaths under 1 month	54	13	41	37	10	27	31.8	19.8	40.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	1	0	1	2.7	-	5.6
Tbc., respiratory (001-008)	1	1	0	1	0	1	2.7	-	5.6
Syphilis (020-029)	3	0	3	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	1	1	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	1	0	1	1	0	2.7	5.3	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	94	43	51	65	29	36	175.7	152.6	200.0
Diabetes (260)	12	4	8	10	5	5	27.0	26.3	27.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	87	29	58	74	28	46	200.0	147.4	255.6
Heart diseases (410-443)	169	72	97	135	66	69	364.9	347.4	383.3
Arteriosclerosis (450)	8	7	1	8	7	1	21.6	36.8	5.6
Influenza (480-483)	8	3	5	7	2	5	18.9	10.5	27.8
#Pneumonia (490-493)	19	6	13	15	6	9	40.5	31.6	50.0
#Diarrhea & enteritis (571)	7	1	6	3	0	3	8.1	-	16.7
Acute nephritis (590, 591)	3	0	3	1	0	1	2.7	-	5.6
Chronic nephritis, etc. (592-594)	14	1	13	6	1	5	16.2	5.3	27.8
Maternal deaths (640-689)	4	0	4	0	0	0	-	-	-
Cong. malformations (750-759)	4	2	2	1	1	0	2.7	5.3	-
Birth inj., atelect. (760-762)	20	4	16	15	4	11	12.9	7.9	16.6
#Other dis. early inf. (763-776)	37	9	28	24	6	18	20.6	11.9	27.2
Violent & acc. deaths (800-999)	45	21	24	32	20	12	86.5	105.3	66.7
Acc. deaths, total (800-962)	39	19	20	28	19	9	75.7	100.0	50.0
Motor vehicle acc. (810-835)	11	8	3	8	7	1	21.6	36.8	5.6
Ill-defined (780-795)	63	13	50	60	11	49	162.2	57.9	272.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WASHINGTON COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	2,820	1,124	1,696	2,707	1,057	1,650	43.0	40.6	44.6
Fetal deaths	204	113	91	203	110	93	75.0	104.1	56.4
Deaths from all causes	784	234	550	798	209	589	12.7	8.0	15.9
Deaths under 1 year	118	34	84	116	30	86	42.8	28.4	52.1
Deaths under 1 month	72	28	44	71	25	46	26.2	23.6	27.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	1	3	2	1	1	3.2	3.8	2.7
Tbc., respiratory (001-008)	3	1	2	2	1	1	3.2	3.8	2.7
Syphilis (020-029)	3	0	3	3	0	3	4.8	-	8.1
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	0	2	2	0	2	3.2	-	5.4
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	2	1	1	2	1	1	3.2	3.8	2.7
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	92	37	55	86	28	58	136.5	107.7	156.8
Diabetes (260)	11	6	5	12	6	6	19.0	23.1	16.2
Pellagra (281)	0	0	0	1	0	1	1.6	-	2.7
Intracranial lesions (330-334)	78	18	60	89	15	74	141.3	57.7	200.0
Heart diseases (410-443)	210	63	147	210	56	154	333.3	215.4	416.2
Arteriosclerosis (450)	6	3	3	6	3	3	9.5	11.5	8.1
Influenza (480-483)	13	2	11	14	2	12	22.2	7.7	32.4
#Pneumonia (490-493)	28	15	13	28	13	15	44.4	50.0	40.5
#Diarrhea & enteritis (571)	11	1	10	11	1	10	17.5	3.8	27.0
Acute nephritis (590, 591)	2	0	2	2	0	2	3.2	-	5.4
Chronic nephritis, etc. (592-594)	10	2	8	13	1	12	20.6	3.8	32.4
Maternal deaths (640-689)	4	0	4	5	0	5	1.8	-	3.0
Cong. malformations (750-759)	8	4	4	8	3	5	12.7	11.5	13.5
Birth inj., atelect. (760-762)	12	6	6	14	6	8	5.2	5.7	4.8
#Other dis. early inf. (763-776)	57	19	38	55	16	39	20.3	15.1	23.6
Violent & acc. deaths (800-999)	56	25	31	53	20	33	84.1	76.9	89.2
Acc. deaths, total (800-962)	47	19	28	42	13	29	66.7	50.0	78.4
Motor vehicle acc. (810-835)	20	12	8	13	6	7	20.6	23.1	18.9
Ill-defined (780-795)	110	8	102	107	8	99	169.8	30.8	267.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WAYNE COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	452	314	138	430	256	174	26.9	25.6	29.0
Fetal deaths	10	5	5	12	5	7	27.9	19.5	40.2
Deaths from all causes	138	85	53	174	109	65	10.9	10.9	10.8
Deaths under 1 year	15	8	7	19	11	8	44.2	43.0	46.0
Deaths under 1 month	7	5	2	10	7	3	23.2	27.3	17.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	3	1	2	18.8	10.0	33.3
Tbc., respiratory (001-008)	0	0	0	3	1	2	18.8	10.0	33.3
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	1	0	2	1	1	12.5	10.0	16.7
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	7	10	19	8	11	118.8	80.0	183.3
Diabetes (260)	3	1	2	3	1	2	18.8	10.0	33.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	21	13	8	26	15	11	162.5	150.0	183.3
Heart diseases (410-443)	43	36	7	44	38	6	275.0	380.0	100.0
Arteriosclerosis (450)	2	1	1	4	3	1	25.0	30.0	16.7
Influenza (480-483)	1	0	1	1	0	1	6.2	-	16.7
#Pneumonia (490-493)	6	3	3	8	4	4	50.0	40.0	66.7
#Diarrhea & enteritis (571)	1	0	1	1	0	1	6.2	-	16.7
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	12.5	10.0	16.7
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	3	2	1	3	2	1	18.8	20.0	16.7
Birth inj., atelect. (760-762)	2	1	1	3	2	1	7.0	7.8	5.7
#Other dis. early inf. (763-776)	2	2	0	3	3	0	7.0	11.7	-
Violent & acc. deaths (800-999)	11	10	1	17	16	1	106.2	160.0	16.7
Acc. deaths, total (800-962)	9	9	0	16	15	1	100.0	150.0	16.7
Motor vehicle acc. (810-835)	3	3	0	7	7	0	43.8	70.0	-
Ill-defined (780-795)	14	2	12	16	3	13	100.0	30.0	216.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WEBSTER COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	125	67	58	203	130	73	15.6	13.0	24.3
Fetal deaths	2	0	2	5	2	3	24.6	15.4	41.1
Deaths from all causes	72	59	13	88	69	19	6.8	6.9	6.3
Deaths under 1 year	0	0	0	1	0	1	4.9	-	13.7
Deaths under 1 month	0	0	0	0	0	0	-	-	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	1	0	7.7	10.0	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	7	2	13	10	3	100.0	100.0	100.0
Diabetes (260)	1	1	0	1	1	0	7.7	10.0	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	20	18	2	20	18	2	153.8	180.0	66.7
Heart diseases (410-443)	26	23	3	29	26	3	223.1	260.0	100.0
Arteriosclerosis (450)	2	2	0	2	2	0	15.4	20.0	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	1	1	3	1	2	23.1	10.0	66.7
#Diarrhea & enteritis (571)	0	0	0	1	0	1	7.7	-	33.3
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	1	0	1	4.9	-	13.7
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	0	0	0	0	0	0	-	-	-
Violent & acc. deaths (800-999)	3	3	0	3	2	1	23.1	20.0	33.3
Acc. deaths, total (800-962)	3	3	0	2	2	0	15.4	20.0	-
Motor vehicle acc. (810-835)	3	3	0	2	2	0	15.4	20.0	-
Ill-defined (780-795)	6	2	4	7	2	5	53.8	20.0	166.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WILKINSON COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	441	188	253	387	73	314	38.7	24.3	44.8
Fetal deaths	21	0	21	23	0	23	59.4	-	73.2
Deaths from all causes	155	69	86	144	49	95	14.4	16.3	13.6
Deaths under 1 year	19	3	16	20	3	17	51.7	41.1	54.1
Deaths under 1 month	12	3	9	12	2	10	31.0	27.4	31.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	19	14	5	17	8	9	170.0	266.7	128.6
Diabetes (260)	1	0	1	1	0	1	10.0	-	14.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	24	10	14	26	9	17	260.0	300.0	242.8
Heart diseases (410-443)	50	22	28	43	14	29	430.0	466.7	414.3
Arteriosclerosis (450)	1	0	1	1	0	1	10.0	-	14.3
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	3	1	2	3	1	2	30.0	33.3	28.6
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	10.0	33.3	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	1	0	1	10.0	-	14.3
Birth inj., atelect. (760-762)	2	0	2	2	0	2	5.2	-	6.4
#Other dis. early inf. (763-776)	8	3	5	9	3	6	23.2	41.1	19.1
Violent & acc. deaths (800-999)	11	6	5	12	7	5	120.0	233.3	71.4
Acc. deaths, total (800-962)	9	6	3	10	7	3	100.0	233.3	42.8
Motor vehicle acc. (810-835)	5	5	0	7	7	0	70.0	233.3	-
Ill-defined (780-795)	19	2	17	19	2	17	190.0	66.7	242.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WINSTON COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	442	182	260	470	188	282	23.5	17.1	31.3
Fetal deaths	20	4	16	24	5	19	51.1	26.6	67.4
Deaths from all causes	161	85	76	196	106	90	9.8	9.6	10.0
Deaths under 1 year	23	7	16	26	11	15	55.3	58.5	53.2
Deaths under 1 month	13	6	7	18	9	9	38.3	47.9	31.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	5.0	9.1	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	5.0	9.1	-
Syphilis (020-029)	1	0	1	1	0	1	5.0	-	11.1
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (040, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	16	14	2	21	18	3	105.0	163.6	33.3
Diabetes (260)	3	1	2	4	1	3	20.0	9.1	33.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	19	10	9	23	12	11	115.0	109.1	122.2
Heart diseases (410-443)	42	28	14	50	31	19	250.0	281.8	211.1
Arteriosclerosis (450)	3	2	1	3	2	1	15.0	18.2	11.1
Influenza (480-483)	8	3	5	8	3	5	40.0	27.3	55.6
#Pneumonia (490-493)	5	1	4	5	1	4	25.0	9.1	44.4
#Diarrhea & enteritis (571)	2	0	2	1	0	1	5.0	-	11.1
Acute nephritis (590, 591)	0	0	0	2	0	2	10.0	-	22.2
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	5.0	9.1	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	4	2	2	5	3	2	25.0	27.3	22.2
Birth inj., atelect. (760-762)	4	2	2	6	2	4	12.8	10.6	14.2
#Other dis. early inf. (763-776)	7	2	5	10	5	5	21.3	26.6	17.7
Violent & acc. deaths (800-999)	9	5	4	12	8	4	60.0	72.7	44.4
Acc. deaths, total (800-962)	4	4	0	7	7	0	35.0	63.6	-
Motor vehicle acc. (810-835)	1	1	0	3	3	0	15.0	27.3	-
Ill-defined (780-795)	12	2	10	12	2	10	60.0	18.2	111.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
YALOBUSHA COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	305	86	219	313	105	208	24.1	15.0	34.7
Fetal deaths	8	2	6	8	2	6	25.6	19.0	28.8
Deaths from all causes	131	66	65	155	87	68	11.9	12.4	11.3
Deaths under 1 year	15	0	15	17	2	15	54.3	19.0	72.1
Deaths under 1 month	10	0	10	11	2	9	35.1	19.0	43.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	7.7	-	16.7
Tbc., respiratory (001-008)	1	0	1	1	0	1	7.7	-	16.7
Syphilis (020-029)	2	0	2	2	0	2	15.4	-	33.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	7.7	-	16.7
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	14	9	5	18	14	4	138.5	200.0	66.7
Diabetes (260)	1	0	1	2	1	1	15.4	14.3	16.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	21	12	9	28	18	10	215.4	257.1	166.7
Heart diseases (410-443)	47	29	18	51	32	19	392.3	457.1	316.7
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	4	1	3	4	1	3	30.8	14.3	50.0
#Pneumonia (490-493)	1	1	0	2	2	0	15.4	28.6	-
#Diarrhea & enteritis (571)	0	0	0	2	1	1	15.4	14.3	16.7
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	15.4	14.3	16.7
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	0	2	1	0	1	7.7	-	16.7
Birth inj., atelect. (760-762)	1	0	1	1	0	1	3.2	-	4.8
#Other dis. early inf. (763-776)	5	0	5	7	2	5	22.4	19.0	24.0
Violent & acc. deaths (800-999)	8	3	5	12	7	5	92.3	100.0	83.3
Acc. deaths, total (800-962)	5	2	3	8	5	3	61.5	71.4	50.0
Motor vehicle acc. (810-835)	1	1	0	4	4	0	30.8	57.1	-
Ill-defined (780-795)	12	2	10	12	1	11	92.3	14.3	183.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
YAZOO COUNTY, MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	934	241	693	978	280	698	34.9	25.4	41.0
Fetal deaths	43	5	38	46	6	40	47.0	21.4	57.3
Deaths from all causes	322	98	224	369	127	242	13.2	11.5	14.2
Deaths under 1 year	34	3	31	43	3	40	44.0	10.7	57.3
Deaths under 1 month	17	2	15	25	2	23	25.6	7.1	33.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	2	2	0	7.1	18.2	-
Tbc., respiratory (001-008)	1	1	0	2	2	0	7.1	18.2	-
Syphilis (020-029)	1	0	1	2	0	2	7.1	-	11.8
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	3.6	-	5.9
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	1	0	1	1	0	1	3.6	-	5.9
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	28	11	17	34	17	17	121.4	154.5	100.0
Diabetes (260)	4	2	2	3	2	1	10.7	18.2	5.9
Pellagra (281)	1	0	1	0	0	0	-	-	-
Intracranial lesions (330-334)	45	17	28	46	20	26	164.3	181.8	152.9
Heart diseases (410-443)	78	28	50	92	35	57	328.6	318.2	335.3
Arteriosclerosis (450)	4	4	0	6	6	0	21.4	54.5	-
Influenza (480-483)	3	1	2	3	2	1	10.7	18.2	5.9
#Pneumonia (490-493)	3	1	2	7	4	3	25.0	36.4	17.6
#Diarrhea & enteritis (571)	4	0	4	4	0	4	14.3	-	23.5
Acute nephritis (590, 591)	1	0	1	2	1	1	7.1	9.1	5.9
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	3.6	-	5.9
Maternal deaths (640-689)	0	0	0	1	0	1	1.0	-	1.4
Cong. malformations (750-759)	2	0	2	3	0	3	10.7	-	17.6
Birth inj., atelect. (760-762)	5	2	3	7	2	5	7.2	7.1	7.2
#Other dis. early inf. (763-776)	9	1	8	15	1	14	15.3	3.6	20.0
Violent & acc. deaths (800-999)	34	9	25	33	10	23	117.8	90.9	135.3
Acc. deaths, total (800-962)	27	5	22	26	6	20	92.8	54.5	117.6
Motor vehicle acc. (810-835)	8	3	5	9	4	5	32.1	36.4	29.4
Ill-defined (780-795)	67	7	60	66	7	59	235.7	63.6	347.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
STATE OF MISSISSIPPI, 1958

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	59673	27231	32442	59189	26812	32377	27.1	20.9	35.9
Fetal deaths	2572	1010	1562	2520	965	1555	42.6	36.0	48.0
Deaths from all causes	21372	10706	10666	21541	10811	10730	9.9	8.4	11.9
Deaths under 1 year	2431	712	1719	2424	706	1718	41.0	26.3	53.1
Deaths under 1 month	1478	547	931	1475	540	935	24.9	20.1	28.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	164	75	89	172	82	90	7.9	6.4	10.0
Tbc., respiratory (001-008)	146	68	78	154	75	79	7.0	5.8	8.8
Syphilis (020-029)	43	11	32	41	8	33	1.9	0.6	3.7
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	34	7	27	33	7	26	1.5	0.5	2.9
Scarlet fever & streptococcal sore throat (050, 051)	2	1	1	2	1	1	0.1	0.1	0.1
Diphtheria (055)	6	0	6	7	1	6	0.3	0.1	0.7
Whooping cough (056)	12	0	12	12	0	12	0.5	-	1.3
Meningococcal infections (057)	20	14	6	22	16	6	1.0	1.2	0.7
Acute poliomyelitis (080)	1	1	0	1	1	0	0.04	0.1	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	17	5	12	15	5	10	0.7	0.4	1.1
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	2444	1471	973	2525	1524	1001	115.6	118.8	111.1
Diabetes (260)	311	155	156	317	163	154	14.5	12.7	17.1
Pellagra (281)	2	0	2	2	0	2	0.1	-	0.2
Intracranial lesions (330-334)	2842	1414	1428	2867	1426	1441	131.3	111.2	159.9
Heart diseases (410-443)	6503	3933	2570	6515	3924	2591	298.3	305.9	287.5
Arteriosclerosis (450)	261	171	90	268	178	90	12.3	13.9	10.0
Influenza (480-483)	198	64	134	196	62	134	9.0	4.8	14.9
#Pneumonia (490-493)	706	280	426	713	282	431	32.6	22.0	47.8
#Diarrhea & enteritis (571)	196	34	162	198	35	163	9.1	2.7	18.1
Acute nephritis (590, 591)	57	10	47	58	11	47	2.6	0.8	5.2
Chronic nephritis, etc. (592-594)	255	85	170	264	89	175	12.1	6.9	19.4
Maternal deaths (640-689)	55	9	46	53	10	43	0.9	0.4	1.3
Cong. malformations (750-759)	242	131	111	254	141	113	11.6	11.0	12.5
Birth inj., atelect. (760-762)	383	192	191	390	195	195	6.6	7.3	6.0
#Other dis. early inf. (763-776)	867	264	603	858	256	602	14.5	9.5	18.6
Violent & acc. deaths (800-999)	1752	918	834	1735	901	834	79.4	70.2	92.5
Acc. deaths, total (800-962)	1390	742	648	1379	729	650	63.1	56.8	72.1
Motor vehicle acc. (810-835)	582	397	185	554	381	173	25.4	29.7	19.2
Ill-defined (780-795)	1956	376	1580	1930	362	1568	88.4	28.2	174.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.