


PUBLIC HEALTH STATISTICS
STATE OF
MISSISSIPPI
1959



VITAL STATISTICS
SUMMARIES

**Mississippi State Board of Health
Jackson, Mississippi**

A. L. Gray, M.D., Executive Officer

ANNUAL BULLETIN

OF

VITAL STATISTICS

1959

SUMMARIES

MISSISSIPPI STATE BOARD OF HEALTH


TABLE OF CONTENTS

	Page
General Summary	1
Table	
1. Estimated Population as of January 1, 1959, by County, by Race	19
2. Live Births, by County and City of Occurrence, Reallocated to Mother's Usual Residence, by Race, with Rates	21
3. Live Births, by County and City of Usual Residence of Mother, by Race and Attendant, and by Sex	24
4. Fetal Deaths, by County and City of Occurrence, Reallocated to Mother's Usual Residence, by Race and Attendant, and by Sex	28
5. Deaths, by Cause, by Race	32
6. Deaths, by County and City of Occurrence, Reallocated to Place of Residence, by Race, with Rates	43
7. Deaths from Selected Causes, by Race, Sex, and Age, with Rates	46
8. Deaths Under 1 Year, Shown for Selected Causes, by Race and Age, with Rates	54
9. Deaths from Selected Causes, by Month of Occurrence	56
10. Marriages, by County of Occurrence and of Residence of Bride, and Divorces, by County of Occurrence, by Race	57
11. Divorces, by Cause, by Number of Minor Children, by Number of Years Married, by Race	59
12. Live Births, 1913-1959, by Year, by Race, with Rates	60
13. Deaths, 1913-1959, by Year, by Race, with Rates	62
14. Maternal Deaths, 1913-1959, by Year, by Race, with Rates	64
15. Deaths Under 1 Year, 1917-1959, by Year, by Race, with Rates	65
16. Deaths under 1 Month, 1924-1959, by Year, by Race, with Rates	66
17. Deaths from Heart Diseases, Nephritis (all forms), Vascular Lesions, Cancer, and Tuberculosis (all forms), 1950-1959, by Year, by Race, with Rates	67
18. Births, Deaths, and Marriages, by Month of Occurrence	68

COUNTY TABLES AND STATE SUMMARY TABLE

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 1 MONTH, AND
DEATHS FROM CERTAIN SELECTED CAUSES, ON BASIS OF PLACE OF OCCURRENCE
AND PLACE OF RESIDENCE, BY RACE, WITH RATES:

	Page
For Each County	71
For the State of Mississippi	153



SUMMARY OF 1959 VITAL STATISTICS

INTRODUCTION

The 1959 Vital Statistics Bulletin is the seventeenth consecutive issue prepared from data obtained from records of live births, fetal deaths, deaths, marriages and divorces collected by the Division of Public Health Statistics.

Population and Rates

Population figures used in this bulletin are estimates as of January 1, 1959, prepared by Dr. George L. Wilber and Ellen Bryant, of the Sociology Department, Mississippi State University, according to the "vital rates" method.

No age or sex-specific rates (except for maternal and infant deaths) and no city rates were calculated for 1959 because current estimates by age, sex, and for cities are not available.

Occurrence and Residence Allocation

Births, fetal deaths, deaths, and marriages were classified by occurrence and residence. Occurrence data were obtained from certificates filed in Mississippi at the place of birth, fetal death, death, or issuance of marriage license. Residence data include transcripts of these events (except marriages) occurring to Mississippi residents outside the State, and exclude records filed in Mississippi for nonresidents. There were 943 births to residents of Mississippi which occurred outside the State in 1959 and were reported to the Division of Public Health Statistics; births occurring inside Mississippi to residents of other states numbered 1,538. Fetal death transcripts totaled 21 and the nonresident certificates 67. During the year, 836 death transcripts were received and 659 nonresident deaths were recorded. The total number of states sending transcripts was 38, including the District of Columbia. Birth records came from 31 states, fetal death records from 4 states, and death records from 28 states. Two reports on Mississippi residents came from foreign countries. In Table A the states furnishing the largest numbers of records are listed.

Some of the tables contain both residence and occurrence data; but, following recommended procedures, most of the tables and all of the rates are based on residence data only. Residence assignment for births and fetal deaths was determined on the basis of usual residence of mother; and for deaths, on the basis of usual residence of decedent. Deaths of inmates or employees of institutions were reallocated to previous residence regardless of length of stay in the institution.

Table A. Transcripts received, by place of occurrence

Place of Occurrence	Live Births	Deaths	Fetal Deaths
<u>Total</u>	<u>943</u>	<u>836</u>	<u>21</u>
Alabama	189	88	4
Arkansas	13	15	0
Colorado	29	0	0
Florida	9	18	0
Illinois	74	28	2
Kentucky	17	13	0
Louisiana	155	184	1
Missouri	20	10	0
Tennessee	342	396	14
Texas	11	28	0
Other	84	56	0

Urban-Rural Classification

Towns with populations of 2,500 or more according to the 1950 census were considered as urban areas. All other areas were classified as rural.

Classification of Causes of Death

Causes of death were assigned according to the Seventh Revision (1955) of the International Statistical Classification of Causes of Death. This revision differs only slightly from the Sixth Revision used from 1949 through 1957 and rates for the years prior to 1958 have not been adjusted for comparability with the Seventh Revision. Only underlying causes were coded.

LIVE BIRTHS

Resident Births

Live births to residents of Mississippi numbered 60,562 in 1959. This total was 1,373 greater than that for 1958, but less than the number in each year of the period 1946-1957. There was an increase of 1,167 for whites and 206 for nonwhites. The crude birth rate of 27.2 per 1,000 population was almost the same as the 1958 rate, which was the lowest since 1946. The State rate for whites in 1959 was 20.9 and county rates varied from 8.8 in Carroll County to 39.3 in Washington County. For nonwhites, the State rate was 36.6 and the range for the counties was from 14.8 in Tippah County to 57.5 in Bolivar County.

Characteristics

A breakdown of the 32,583 nonwhite births by specific race shows 32,410 were Negro, 116 were Indian, 27 were Chinese, 25 were Japanese, and 5 were other nonwhites. Births of males numbered 30,755 and there were 29,807 females. These totals yielded a sex ratio of 1,032 males to 1,000 females. August was the peak month of occurrence for live births. There were 5,682 in this month in comparison with only 4,347 in May when the fewest occurred.

During 1959, 21,212 children were born to urban residents and 39,350 to rural residents. Over half (54 per cent) of the infants born to urban residents were white,

but 58 per cent of the rural were nonwhite. Of the births to rural residents, 23,053 or 58.6 per cent were born to farm residents; 14,960 were born to nonfarm residents; and on 1,337 certificates information pertaining to farm residency was not stated. Comparison of the occurrence and residence figures indicates that a majority of the white mothers living in rural areas went to cities for delivery as compared with a much smaller percentage of the nonwhite mothers.

The number of illegitimate births during 1959 was 8,091 or 13.4 per cent of the total. This was 510 more than in 1958, with an increase in percentage of 0.6. White births classified as illegitimate totaled 370 or 1.3 per cent, and 7,721 or 23.7 per cent of the nonwhite births were illegitimate.

Liveborn infants classified as immature (birth weight 5 1/2 pounds or less) numbered 5,738 or 9.4 per cent of the total number of births registered. The incidence of immaturity was greater among the nonwhites than among the whites, 11.7 per cent in comparison with 6.7 per cent.

Classification by Attendant

In Table B, which gives the distribution of live births by attendant and race, it may be seen that, in 1959, about 72 per cent of all births occurred in hospitals or clinics (including physicians' offices or clinics which provide inpatient care). A small proportion, 3.5 per cent, was delivered by physicians in places other than hospitals or clinics; and 24.4 per cent were delivered at home by nonmedical attendants. Forty-five per cent of the nonwhite mothers, but less than one per cent of the white mothers, had no medical attendant.

Table B. Live births by attendant and race

Attendant	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
<u>Total</u>	<u>60,562</u>	<u>100.0</u>	<u>27,979</u>	<u>100.0</u>	<u>32,583</u>	<u>100.0</u>
Physician in hospital	43,664	72.1	27,666	98.9	15,998	49.1
Physician in home	2,102	3.5	187	0.7	1,915	5.9
Midwife	14,719	24.3	110	0.4	14,609	44.8
<u>Other</u>	<u>77</u>	<u>0.1</u>	<u>16</u>	<u>0.1</u>	<u>61</u>	<u>0.2</u>

There has been a steady increase in the proportion of live births occurring in hospitals in recent years (see Table C). The year 1951 was the first in which at least half of the deliveries took place in hospitals, and the proportion of hospital deliveries has increased each year since. The progress made by the nonwhites has been more rapid, but there is still a large difference between the percentages for the two racial groups.

Table C. Percentage of live births in hospitals or clinics, 1955-1959, by race

Year	Total	White	Nonwhite
1959	72.1	98.9	49.1
1958	70.4	98.6	47.0
1957	68.3	98.0	44.8
1956	65.7	97.2	41.0
1955	62.4	96.0	36.4

The proportions of live births attended by physicians during 1955-1959 are shown in Table D. The improvement shown in this table is less pronounced than that in the preceding table, but is also steady. This is the fourth year in which more than half the nonwhite mothers were attended by physicians.

Table D. Percentage of live births attended by physicians, 1955-1959, by race

Year	Total	White	Nonwhite
1959	75.6	99.5	55.0
1958	74.2	99.5	53.3
1957	73.6	99.5	53.2
1956	72.2	99.3	51.0
1955	70.4	99.1	48.3

Age of Mother

Table E gives the age distribution of mothers by race. Little variation in this distribution has occurred in recent years. Approximately half of the mothers were 15-24 years of age and more than one-third were 25-34 years old. In the periods 15-24 and 25-34 years the percentages of white mothers were slightly larger than those of nonwhites.

Table E. Live births by age of mother and race, with per cent of total births

Age of Mother	Total		White		Nonwhite	
	No.	%	No%	No%	No%	%
10-14 years	331	0.5	31	0.1	300	0.9
<u>15-44 years</u>	<u>60,009</u>	<u>99.1</u>	<u>27,895</u>	<u>99.7</u>	<u>32,114</u>	<u>98.6</u>
15-24 years	31,131	51.4	14,875	53.2	16,256	49.9
25-34 years	22,176	36.6	10,482	37.5	11,694	35.9
35-44 years	6,702	11.1	2,538	9.1	4,164	12.8
45-54 years	104	0.2	32	0.1	72	0.2
55-64 years	1	0.002	1	0.004	0	-
Unknown	117	0.2	20	0.1	97	0.3

FETAL DEATHS

In 1952 the registration of all fetal deaths, regardless of length of gestation, was initiated in Mississippi. Prior to that time, only stillbirths of 20 or more weeks of gestation were reported. Registration of early fetal deaths is still

far from complete, but there is no estimate of the degree of nonregistration. There were 2,649 fetal death certificates filed during 1959 and 2,603 certificates for residents. The resident total was 83 more than the comparable figure for 1958, the change being almost entirely among whites. For the State the fetal death ratio (fetal deaths per 1,000 live births) was 43.0. The ratio for whites was 37.3 with county ratios varying from zero in Leake and Webster Counties to 98.4 in Holmes County. The ratio for nonwhites was 47.8 with variations from zero in Itawamba, Prentiss, and Tishomingo Counties to 99.0 in Union County.

On 956 of the records the gestation period was stated to be less than 20 weeks; on 1,495 it was 20 weeks or more; and on 152 the length was not stated. In the period 1955-1959 there have been reported among the whites more fetal deaths after less than 20 weeks of gestation than after 20 weeks or more.

Using the certificates with sex stated, a sex ratio of 1,270 males to 1,000 females was determined. This ratio was the lowest since 1955. As usual, the sex ratio for fetal deaths was much higher than that for live births, but was not so reliable a figure because the sex was not stated on 773 or almost 30 per cent of the fetal death certificates.

The distribution by race and attendant is given in Table F. Approximately 60 per cent of the total were nonwhite. More than 73 per cent of the reported fetal deaths were attended by physicians in hospitals and about 18 per cent were attended by midwives. Although the proportion of white live births in hospitals was larger than the percentage of white fetal deaths in hospitals, the reverse was true for the nonwhites. This situation in the nonwhite group may be accounted for by the non-reporting of fetal deaths that occur at home and the tendency to transfer difficult and premature deliveries to a hospital.

Table F. Fetal deaths by attendant and race

Attendant	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
<u>Total</u>	<u>2,603</u>	<u>100.0</u>	<u>1,045</u>	<u>100.0</u>	<u>1,558</u>	<u>100.0</u>
Physician in hospital	1,914	73.5	986	94.4	928	59.6
Physician in home	198	7.6	53	5.1	145	9.3
Midwife	464	17.8	3	0.3	461	29.6
Other	27	1.0	3	0.3	24	1.5

Table G shows the percentage of fetal deaths delivered in hospitals in the last five years. Increases have occurred among both whites and nonwhites. The gap between the two race groups is a large one, but not so large as that for live births.

Table G. Percentage of fetal deaths delivered in hospitals, 1955-1959, by race

Year	Total	White	Nonwhite
1959	73.5	94.4	59.6
1958	69.0	88.8	56.8
1957	69.5	92.2	54.8
1956	67.2	91.0	52.7
1955	63.5	88.1	48.5

DEATHS

Resident Deaths

In 1959 the resident death total was 20,890, which is a decrease of 651 from the number for 1958. The crude death rate, based on the 1959 population estimate, was 9.4 per 1,000 population in comparison with a rate of 9.9 in 1958 based on the estimated population for that year. Slightly over one-third or 7,337 of the persons who died in 1959 were urban residents, but 49.5 per cent of the deaths occurred in urban areas.

Although the number of deaths of nonwhites was 594 less than that of the whites, the death rate of the nonwhites was 3.4 per 1,000 population greater than that of the whites. The total of 10,742 deaths of white persons was the second largest reported since 1913, the first complete year of death registration in Mississippi. The number of deaths of nonwhites in 1959 (10,148) was higher than in 1957, 1956, 1955 or 1954. The death rates of both whites and nonwhites were slightly lower than in 1958.

County rates for the white population ranged from 4.5 in Issaquena to 19.7 in Wilkinson, with a rate for the State of 8.0, while county rates for the nonwhites ranged from 3.5 in Prentiss to 24.0 in Tishomingo, with a State rate of 11.4.

The resident birth rate exceeded the death rate by 17.8 per 1,000 population. For whites the excess was 12.9 and for the nonwhites it was 25.2.

Age at Death

Table H shows the distribution of deaths by age and race. In comparison with the 1958 figures there were decreases in all the white groups except the 15-24, 45-54 and 75-84 year groups. The 15-24 year group showed an increase of almost 12 per cent over 1958 figures. Decreases occurred in all nonwhite groups except the 5-24, 35-44 and 55-64 year groups. For deaths of all races combined there were increases from 1958 in the 5-9, 15-24, 35-44 and 55-64 year groups, with the largest increase in the age group 15-24 years.

Table H. Age-race distribution of deaths

Age Group	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
All Ages	20,890	100.0	10,742	100.0	10,148	100.0
Under 1 year	2,355	11.3	677	6.3	1,678	16.5
1-4 years	391	1.9	112	1.0	279	2.7
5-9 years	168	0.8	61	0.6	107	1.0
10-14 years	141	0.7	64	0.6	77	0.8
15-24 years	505	2.4	228	2.1	277	2.7
25-34 years	496	2.4	198	1.8	298	2.9
35-44 years	955	4.6	438	4.1	517	5.1
45-54 years	1,891	9.0	974	9.1	917	9.0
55-64 years	3,003	14.4	1,523	14.2	1,480	14.6
65-74 years	4,528	21.7	2,518	23.4	2,010	19.8
75-84 years	4,449	21.3	2,805	26.1	1,644	16.2
85 years +	1,988	9.5	1,139	10.6	849	8.4
Unknown	20	0.1	5	0.05	15	0.1

Comparison of the age distribution of the two races reveals significant differences. The percentage of deaths of nonwhites which occurred among children under 10 years of age was more than double that for the whites, and in the groups from 10 through 44 years relatively more deaths of nonwhites than of whites were registered. "Retirement" age or 65 years had been reached by 60 per cent of the whites who died in 1959 and only 44 per cent of the nonwhites. The median age at death for the State was 66.2; for the whites 69.4; and for the nonwhites 61.2.

Leading Causes of Death

In Table I are listed the ten leading causes of death in the State in 1959. The first six of these causes are the same as in the six preceding years. Chronic and unspecified nephritis and other renal sclerosis, which was in seventh place in 1953 through 1957, and in ninth place in 1958, does not appear among the ten leading causes in 1959. Diabetes mellitus, which had been successively in tenth, eighth and seventh place, was again in eighth place in 1959. General arteriosclerosis, which was in tenth place in 1957, and in eighth place in 1958, moved to seventh place in 1959. Hypertension, in ninth place in 1959, for the first time appeared as one of the ten leading causes. Congenital malformations remained in tenth place.

Altogether the ten leading causes were responsible for 77 per cent of the deaths. For the whites the same ten causes were listed as for the total population with the exception that other diseases of the circulatory system shared ninth place with diabetes, and hypertension was not on the list. For nonwhites, homicide ranked in seventh place and gastritis in ninth place, while general arteriosclerosis and congenital malformations were not on the list.

Table I. Ten leading causes of death by race, with per cent of total deaths

Cause and International List Number	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
1. Diseases of heart (410-443)	6,219	29.8	3,772	35.1	2,447	24.1
2. Vascular lesions affecting central nervous system (330-334)	2,683	12.8	1,386	12.9	1,297	12.8
3. Malignant neoplasms (140-205)	2,600	12.4	1,580	14.7	1,020	10.0
4. Accidents (800-962)	1,501	7.2	837	7.8	664	6.5
5. Certain diseases of early infancy (760-776)	1,213	5.8	429	4.0	784	7.7
6. Pneumonia and influenza (480-493)	768	3.7	328	3.0	440	4.3
7. General arteriosclerosis (450)	294	1.4	193	1.8	101	1.0
8. Diabetes mellitus (260)	281	1.3	128	1.2	153	1.5
9. Hypertension without mention of heart (444-447)	255	1.2	96	0.9	159	1.6
10. Congenital malformations (750-759)	247	1.2	140	1.3	107	1.0

Heart disease, the leading cause of death, was responsible for 29.8 per cent of all deaths in 1959. During the year 6,219 deaths were attributed to some form of heart disease. In Table J it may be seen that a large number of these (4,365 deaths) were specified as arteriosclerotic and degenerative heart disease, and that the next largest group was hypertensive heart disease with 1,285 deaths. The total death rate was 279.2 per 100,000 population, a decrease from 298.3 for 1958. Decreases occurred among both whites and nonwhites.

Table J. Deaths from heart disease, by race by type of disease

Cause of Death and International List Number	Race		
	Total	White	Nonwhite
Diseases of the heart			
Chronic rheumatic heart disease (410-416)	<u>6,219</u> 127	<u>3,772</u> 63	<u>2,447</u> 64
Arteriosclerotic and degenerative heart disease (420-422)	4,365	3,029	1,336
Other diseases of heart (430-434)	442	199	243
Hypertension with heart disease (440-443)	1,285	481	804

For the second leading cause of death, vascular lesions affecting the central nervous system, there were 2,683 deaths representing 12.8 per cent of the total. The components of this category are listed in Table K showing that a majority of the deaths were due to cerebral hemorrhage (1,682) and 563 were due to cerebral embolism or thrombosis. There was a decrease in rate from that of the previous year; and, as in previous years, the rate for nonwhites exceeded that for whites.

Table K. Deaths from vascular lesions affecting the central nervous system, by race by type of lesion

Cause of Death and International List Number	Race		
	Total	White	Nonwhite
Vascular lesions affecting central nervous system (330-334)			
Subarachnoid hemorrhage (330)	<u>2,683</u> 66	<u>1,386</u> 29	<u>1,297</u> 37
Cerebral hemorrhage (331)	1,682	825	857
Cerebral embolism and thrombosis (332)	563	357	206
Other and ill-defined (334)	372	175	197

In 1959 the cancer death totals for both racial groups were the highest on record in this State. There were 2,600 deaths from this cause or 75 more than in 1958. The distribution of these deaths by age, sex, and race is given in Table L. In comparison with 1958, there were increases among each race-sex group except nonwhite females. There were changes from 1958 in almost every age group with the largest decreases occurring in deaths of white males 55 to 64 years, white females 25 to 54 years, and nonwhite females 35 to 54 years. The largest increases occurred in white males 45 to 54 years and 65 to 74 years, in white females 55 years and over, and in nonwhite males 55-64 years.

More deaths occurred among the whites than among nonwhites in all age groups, and the excess was largest in the age group 75-84, ~~at there were more males than females~~ in the age groups 55-84, and more females in the age groups 35-54 and 85 years and over.

Table L. Cancer deaths by age, sex, and race

Age Group	Total	Male		Female	
		White	Nonwhite	White	Nonwhite
<u>All Ages</u>	<u>2,600</u>	<u>849</u>	<u>516</u>	<u>731</u>	<u>504</u>
Under 10 years	43	15	8	12	8
10-14 years	9	1	3	4	1
15-24 years	30	13	5	5	7
25-34 years	35	14	4	5	12
35-44 years	131	36	24	36	35
45-54 years	356	112	56	98	90
55-64 years	579	171	131	154	123
65-74 years	754	259	160	191	144
75-84 years	522	190	101	171	60
85 years & over	140	38	24	54	24
Unknown	1	0	0	1	0

Table M lists cancer deaths by primary site in descending order according to the number in the total population. The number of deaths assigned to the leading site, digestive organs and peritoneum, is 47 more than in 1958. Cancer of the breast and genito-urinary system accounted for the second largest number of cancer deaths. The two leading sites accounted for 76 per cent of the cancer deaths among nonwhite females, more than 70 per cent among white females, more than 65 per cent among nonwhite males, but less than 46 per cent among white males. Cancer of the respiratory system showed an increase of 27 deaths, remaining in third rank. There was an increase of 40 in the number of deaths from cancer of other and unspecified sites.

Table M. Cancer deaths by primary site, sex, and race, with percentage distribution

Primary site and International List Number	Male				Female			
	White		Nonwhite		White		Nonwhite	
	No.	%	No.	%	No.	%	No.	%
<u>Total</u>	<u>849</u>	<u>100.0</u>	<u>516</u>	<u>100.0</u>	<u>731</u>	<u>100.0</u>	<u>504</u>	<u>100.0</u>
1. Digestive organs and peritoneum (150-159)	254	29.9	195	37.8	258	35.3	135	26.8
2. Breast & genito-urinary system (170-181)	134	15.8	143	27.7	257	35.2	247	49.0
3. Respiratory system (160-165)	228	26.8	58	11.2	35	4.8	24	4.8
4. Lymphatic & hematopoietic tissues (200-205)	96	11.3	45	8.7	79	10.8	20	4.0
5. Brain & nervous system (192, 193)	27	3.2	5	1.0	24	3.3	8	1.6
6. Skin (190, 191)	27	3.2	3	0.6	10	1.4	4	0.8
7. Buccal cavity & pharynx (140-148)	18	2.1	7	1.4	13	1.8	4	0.8
8. Bone & connective tissue (196, 197)	8	0.9	7	1.4	11	1.5	4	0.8
9. Other & unspecified sites (194, 195, 198, 199)	57	6.7	53	10.3	44	6.0	58	11.5

The number of deaths in 1959 which resulted from accidents was 1,501, an increase of 122 over the number in 1958. The greatest increases were in the number of deaths from firearms and from motor vehicles. Among both whites and nonwhites the greatest increase in accidental deaths was in deaths from firearms. Among whites there were no decreases in the number of deaths from any of the six leading causes of accidental deaths; among nonwhites the greatest decrease was in the number of deaths from drowning. The largest number of deaths from all accidents occurred in December in 1959, as in 1957 and 1958, and January was in second place. December was first in the number of motor vehicle accidents, and October was second. Table N shows the six leading causes of accidental death in 1959 by race. The first five of these causes appeared on the 1958 list in the same order except that in 1958 drownings caused a larger number of deaths than falls. Machinery was among the six leading causes of accidental death for the first time since 1954.

Table N. Six leading causes of accidental deaths among residents, by race

Type of Accident	Total	White	Nonwhite
1. Motor vehicle	588	401	187
2. Fire and explosion	225	58	167
3. Falls	117	91	26
4. Drowning except water transport	109	58	51
5. Firearm	94	46	48
6. Machinery	41	22	19

Leading causes of death for each age group except infants are given in Table O. None of these lists was exactly like any other, but there were some causes which appeared repeatedly. Accidents was listed in every group, and appeared as the chief cause in all groups under 45 years of age. Malignant neoplasms also appeared on every list and heart disease on all except 1-4 years. Vascular lesions affecting the central nervous system was listed in the three groups 25 and over. Pneumonia and influenza appeared in three groups, 1-4 years, 5-14 years, and 65 years and over. In both lists for children under 15, congenital malformations was included, and for the 5-14, 15-24, and 25-44 year groups homicide was listed. The gastritis group was listed only for the 1-4 year group. In the age group 65-74 the five leading causes were the same as for the age group 65 years and over, but not in the same order. In the 65-74 year group malignant neoplasms were in second place, vascular lesions in third place, and accidents in fourth place. In the age groups 75-84 and 85 and over, accidents does not appear on the list of five leading causes, and general arteriosclerosis is in fifth place.

Ill-defined Causes

Of the 20,713 certificates filed for deaths that occurred in Mississippi in 1959 those with no satisfactory statement of cause numbered 2,485 or 12 per cent of the total. Senility was the only cause given on 56 of these certificates; ill-defined diseases of early infancy and those due to immaturity on 628; and other ill-defined and unknown causes on 1,801. For white persons the percentage of certificates in this category was 6.0 and for nonwhites 18.3.

Table 0. Five leading causes of death by age group

<u>Age and Cause of Death</u>	<u>No. of Deaths</u>	<u>% of Total</u>
1-4 years		
All Causes	<u>391</u>	<u>100.0</u>
1. Accidents	141	36.1
2. Pneumonia and influenza	55	14.1
3. Congenital malformations	24	6.1
4. Malignant neoplasms, including leukemia	23	5.9
5. Gastritis, duodenitis, enteritis and colitis	15	3.8
5-14 years		
All Causes	<u>309</u>	<u>100.0</u>
1. Accidents	152	49.2
2. Malignant neoplasms, including leukemia	26	8.4
3. Congenital malformations	17	5.5
4. Pneumonia and influenza	13	4.2
5. Heart disease	9	2.9
5. Homicide	9	2.9
5. Rheumatic fever	9	2.9
15-24 years		
All Causes	<u>505</u>	<u>100.0</u>
1. Accidents	242	47.9
2. Homicide	56	11.1
3. Malignant neoplasms, including leukemia	30	5.9
4. Heart disease	26	5.1
5. Maternal deaths	15	3.0
25-44 years		
All Causes	<u>1,451</u>	<u>100.0</u>
1. Accidents	324	22.3
2. Heart disease	240	16.5
3. Malignant neoplasms, including leukemia	166	11.4
4. Homicide	100	6.9
5. Vascular lesions affecting central nervous system	79	5.4
45-64 years		
All Causes	<u>4,894</u>	<u>100.0</u>
1. Heart disease	1,725	35.2
2. Malignant neoplasms, including leukemia	935	19.1
3. Vascular lesions affecting central nervous system	571	11.7
4. Accidents	258	5.3
5. Diabetes mellitus	89	1.8
65 years and over		
All Causes	<u>10,965</u>	<u>100.0</u>
1. Heart disease	4,198	38.3
2. Vascular lesions affecting central nervous system	2,015	18.4
3. Malignant neoplasms, including leukemia	1,416	12.9
4. Pneumonia and influenza	378	3.4
5. Accidents	292	2.7

Deaths with unsatisfactory statement of cause were so numerous that if placed on the list of ten leading causes they would rank fourth and would appear as one of the five chief causes of death in every age group. If the true causes of death for these persons could be determined and properly assigned, the order of principal causes might not be the same and even the causes might be different.

If deaths from ill-defined diseases peculiar to early infancy and deaths due to immaturity are excluded from the group of deaths without definite cause, this group is 9 per cent of all deaths recorded in Mississippi. In Benton, Carroll, DeSoto, Issaquena and Madison Counties, they were more than 20 per cent of the total deaths, and in Hancock, Humphreys, Kemper, Sharkey, and Yazoo Counties 15 to 20 per cent had no definite cause.

Infant Mortality

An infant mortality rate of 38.9 per 1,000 live births was recorded in 1959, lower than the 1958 rate by 2.1 per 1,000 births. The white rate showed a slightly greater decrease than the nonwhite rate. Distribution of infant deaths by age and race is shown in Table P. In each age group, the rate for nonwhites exceeded that for whites, with the greatest difference between the two racial groups occurring in the period 28 days to one year.

Table P. Infant deaths by age and race, with rates

<u>Age at Death</u>	<u>Total</u>		<u>White</u>		<u>Nonwhite</u>	
	<u>No.</u>	<u>Rate*</u>	<u>No.</u>	<u>Rate*</u>	<u>No.</u>	<u>Rate*</u>
<u>Total under 1 year</u>	<u>2,355</u>	<u>38.9</u>	<u>677</u>	<u>24.2</u>	<u>1,678</u>	<u>51.5</u>
Under 1 day	696	11.5	277	9.9	419	12.8
One to 28 days	754	12.4	238	8.5	516	15.8
28 days to 1 year	905	14.9	162	5.8	743	22.8

* Deaths per 1,000 live births.


Table Q gives the five leading causes of infant deaths. Birth injuries, which was in fourth place in 1957 and in fifth place in 1958, was again in fourth place in 1959, with gastritis, duodenitis, enteritis and colitis again in fifth place. Immaturity, which is no longer considered a separate classification in ranking causes of infant deaths, was mentioned either in combination with another cause or as the only cause on 45 per cent of the certificates for white infants and more than 32 per cent of those for nonwhites.

Table Q. Leading causes of infant deaths, by race

<u>Cause of Death</u>	<u>Total</u>		<u>White</u>		<u>Nonwhite</u>	
	<u>No.</u>	<u>%</u>	<u>No.</u>	<u>%</u>	<u>No.</u>	<u>%</u>
<u>All Causes</u>	<u>2,355</u>	<u>100.0</u>	<u>677</u>	<u>100.0</u>	<u>1,678</u>	<u>100.0</u>
1. Postnatal asphyxia and atelectasis	218	9.2	117	17.3	101	6.0
2. Influenza and pneumonia, except pneumonia of newborn	207	8.8	32	4.7	175	10.4
3. Congenital malformations	187	7.9	99	14.6	88	5.2
4. Birth injuries	138	5.8	68	10.0	70	4.2
5. Gastritis, duodenitis, enteritis & colitis, except diarrhea of newborn	128	5.4	8	1.2	120	7.2

Infant death certificates which showed ill-defined causes or no cause at all numbered 984 or 42 per cent of the total in 1959. For 34 per cent of the white and 45 per cent of the nonwhite infants no definite cause of death was given. Determination of the true cause of death for this large group of infants might change the leading causes of death which are listed above.

Infant mortality rates by race for 1920-1959 are shown in Graph 1. In general, the trend has been downward for both races, but in recent years there has been some upward swing in the curve for the nonwhites.



Maternal Mortality

The maternal mortality rate for 1959 was 0.9 per 1,000 live births in the total population, 0.3 for white mothers and 1.4 for nonwhite. The total of 55 maternal deaths was two more than in 1958. There was a decrease of one death among the white mothers, and an increase of three among the nonwhite. There is still a large discrepancy between the rates by race, but the gap has narrowed considerably in recent years.

In 1959, as in each year from 1947 through 1958, the two leading causes were toxemia and hemorrhage. Toxemia has been the leading cause in each year of this period except 1957.

Table R. Maternal deaths by cause and race, with rates

Cause of Death	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
<u>All Causes</u>	<u>55</u>	<u>0.9</u>	<u>9</u>	<u>0.3</u>	<u>46</u>	<u>1.4</u>
Toxemia	22	0.4	2	0.1	20	0.6
Hemorrhage	12	0.2	2	0.1	10	0.3
Sepsis	5	0.1	3	0.1	2	0.1
Ectopic pregnancy	2	0.03	0	-	2	0.1
Abortion without sepsis or toxemia	2	0.03	0	-	2	0.1
Abortion with sepsis	1	0.02	1	0.04	0	-
Abortion with toxemia, without mention of sepsis	2	0.03	0	-	2	0.1
<u>Other complications</u>	<u>9</u>	<u>0.1</u>	<u>1</u>	<u>0.04</u>	<u>8</u>	<u>0.2</u>

* Deaths per 1,000 live births.

Table S gives the age and race-specific maternal death rates. There were no deaths in the 10-14 year group and only one death in the 45-54 year group. At all ages the rates for nonwhite mothers were higher than those for white mothers.


Table S. Maternal deaths by age group and race, with rates

Age Group	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
<u>All Ages</u>	<u>55</u>	<u>0.9</u>	<u>9</u>	<u>0.3</u>	<u>46</u>	<u>1.4</u>
15-24 years	15	0.5	4	0.3	11	0.7
25-34 years	15	0.7	0	-	15	1.3
35-44 years	24	3.6	5	2.0	19	4.6
45-54 years	1	9.6	0	-	1	13.9

* Deaths per 1,000 live births to mothers in specific age-race group.

Maternal death rates by race for 1913-1959 are shown in Graph 2. A downward trend began in the 1930's and has been rapid since 1940.

Graph 2. Maternal Mortality
1913-1959



MARRIAGES

A new marriage law was passed at a special session of the legislature in December, 1957, becoming effective on July 1, 1958. The law was in effect during the last six months of 1958, and for the whole year of 1959. The total number of marriages decreased from 62,495 in 1957 to 36,402 in 1958 and to 20,447 in 1959. Marriages among whites decreased from 46,868 in 1957 to 26,998 in 1958 and to 13,547 in 1959. Among nonwhites they decreased from 15,627 in 1957 to 9,404 in 1958 and to 6,900 in 1959.

The average daily number of marriages was 56, 44 less than the average for 1958 and 115 less than the average for 1957. In June, the months with the largest total, the daily average was 73, and in December the daily average was 70.

This was the first marriage for 14,398 or 70.4 per cent of the brides and 14,433 or 70.6 per cent of the grooms. For less than one fourth of each group there had been one previous marriage. Ninety-nine brides and 94 grooms did not state the number of previous marriages.

Table T shows the age distribution of brides and grooms. Of the brides, 9,165 or 44.8 per cent were under 20 years of age, and 71.1 per cent were under 25 years of age. There was a much smaller percentage of grooms in the teen-age group - 15.8 per cent - while 41.5 per cent of the grooms were in the 20-24 year age group. A cross tabulation of the ages of brides and grooms shows marriages of 16 couples with

both 75 years of age and over. While the youngest bride of a groom of 75 years of age or more was 17 years of age, most marriages were between persons of approximately the same age.

Table T. Age distribution of brides and grooms

Age	No.	Bride		Groom	
		No.	%	No.	%
<u>Total</u>	<u>20,447</u>	<u>100.0</u>		<u>20,447</u>	<u>100.0</u>
Under 15 years	110	0.5		1	0.005
15 years	868	4.2		6	0.03
16 years	1,485	7.3		47	0.2
17 years	1,863	9.1		480	2.3
18 years	2,782	13.6		1,091	5.3
19 years	2,057	10.1		1,604	7.8
20-24 years	5,373	26.3		8,484	41.5
25-34 years	2,701	13.2		4,437	21.7
35-44 years	1,529	7.5		1,709	8.4
45-54 years	959	4.7		1,173	5.7
55 years and over	709	3.5		1,412	6.9
Unknown	11	0.05		3	0.01

Among the marriages between nonresident brides and grooms there were 6 brides and no groom under 15 years of age. Nonresident brides in the 15-19 year age group numbered 1,435 or 15.8 per cent of all brides in this age group.

In Table U are shown data on the residence status of both brides and grooms. In 1959 both parties were residents of Mississippi in 72 per cent of the marriages. In 1958 both parties were residents in 43 per cent of the marriages, with a percentage of 34 during the first six months and of almost 67 during the last six months. In 1957 both parties were residents in 30 per cent of the marriages. There were 4,062 out-of-state brides or 20 per cent of the total in 1959. In 1958 there were 18,920 nonresident brides or 52 per cent of the total, and in 1957 there were 41,510 nonresident brides or 66.4 per cent of the total.

Table U. Marriages in Mississippi by residence status of bride and groom

Groom	Total	Bride		
		Resident	Nonresident	Not Stated
<u>Total</u>	<u>20,447</u>	<u>16,288</u>	<u>4,062</u>	<u>97</u>
Resident	15,368	14,806	544	18
Nonresident	4,986	1,462	3,516	8
Not stated	93	20	2	71

A comparison during the last twelve years of the resident status of brides is shown in Table V. In each year 1947 through 1956 the number of out-of-state brides increased and, with the exception of 1955, the number of resident brides decreased. The declining trend among resident brides continued in the years 1957, 1958 and 1959, but the increase in nonresidents ended abruptly when the new marriage law was passed.

Table V. Marriages in Mississippi by residence status of bride 1947-1959

Year	Total	Resident	Nonresident	Not Stated
1959	20,447	16,288	4,062	97
1958	36,402	17,370	18,920	112
1957	62,495	20,864	41,510	121
1956	66,223	23,011	43,212	*
1955	66,423	23,683	42,740	*
1954	61,381	23,555	37,826	*
1953	61,751	24,497	37,254	*
1952	58,116	24,868	33,248	*
1951	57,321	25,713	31,608	*
1950	56,738	26,859	29,879	*
1949	52,765	27,717	25,048	*
1948	53,645	30,921	22,724	*
1947	53,584	34,067	19,517	*

* Not classified.

With the exception of Alaska, Delaware, Montana, Nevada and Utah, brides came to Mississippi from every state in the union and from the District of Columbia, Hawaii and Puerto Rico, and there were 10 from foreign countries. Bordering states furnished 15.3 per cent of the brides married in Mississippi and 16.5 per cent of the grooms, with totals as follows: Alabama, 614 brides and 701 grooms; Arkansas, 233 brides and 239 grooms; Louisiana, 808 brides and 946 grooms; and Tennessee, 1,466 brides and 1,488 grooms. Illinois furnished 234 brides and 303 grooms. These five states furnished 16.4 per cent of the brides married in Mississippi and 18 per cent of the grooms.

Each county in the state except Carroll, Covington and Humphreys issued marriage licenses to one or more out-of-state brides in 1959. Counties adjacent to Tennessee and Arkansas and the coastal counties issued the largest numbers, with most of the remainder issued in other counties adjacent to Alabama and Louisiana. In DeSoto County almost 90 per cent of the licenses were for nonresident brides, in Alcorn County more than 70 per cent, and in Hancock and George Counties more than half. Counties bordering Tennessee issued licenses to 45 per cent of the out-of-state brides.

Table W. Number of out-of-state brides obtaining marriage license in certain counties

DeSoto	1,114	Pearl River	113
Alcorn	465	Tishomingo	92
Harrison	363	Wilkinson	82
Jackson	230	Adams	68
Washington	229	Lowndes	66
Hancock	166	Hinds	53
George	126	Prentiss	53
Marshall	126	Greene	45

DIVORCES

Divorces granted in Mississippi during 1959 numbered 5,108 - 92 more than in 1958. There was an increase of 72 among the whites and of 20 among the nonwhites. Cruel and inhuman treatment and desertion continued to rank as the two leading causes and together accounted for 88 per cent of the total granted. Of the divorces among the whites, cruel and inhuman treatment was given as the main cause for 59 per cent; among nonwhites 63 per cent were granted on grounds of desertion. The wife was listed as the complainant on 64 per cent of the records. Only 250 suits or 5 per cent were contested.

No minor children were involved in 49 per cent of the divorces, and 22 per cent of the couples had only one minor child. Nine of the families had 9 or more minor children. The total number of minor children affected was 5,015.

Forty-one per cent of the divorces in 1959 were granted to persons married less than 5 years; 6 per cent had been married less than one year; but 16 per cent had been married 20 years or longer. More divorces were granted to couples in their second year of marriage than in any other year. The distribution of divorces according to duration of marriage was different for the two races. Of the white couples, 49 per cent were granted after less than 5 years of marriage and only 10 per cent after 20 years or longer; for the nonwhites, corresponding percentages were 24 and 31 respectively.

TABLE 1
ESTIMATED POPULATION*, MISSISSIPPI
AS OF JANUARY 1, 1959

County	Total	White	Nonwhite
<u>TOTAL</u>	<u>2,227,000</u>	<u>1,336,000</u>	<u>891,000</u>
Adams	31,000	18,000	13,000
Alcorn	26,000	20,000	6,000
Amite	16,000	8,000	8,000
Attala	20,000	11,000	9,000
Benton	10,000	5,000	5,000
Bolivar	41,000	17,000	24,000
Calhoun	15,000	11,500	3,500
Carroll	11,000	6,000	5,000
Chickasaw	16,000	8,000	8,000
Choctaw	10,000	7,000	3,000
Claiborne	11,000	2,000	9,000
Clarke	16,000	10,000	6,000
Clay	16,000	8,000	8,000
Coahoma	43,000	17,000	26,000
Copiah	28,000	14,000	14,000
Covington	16,000	10,000	6,000
DeSoto	22,000	9,000	13,000
Forrest	52,000	38,000	14,000
Franklin	9,000	5,000	4,000
George	15,000	13,000	2,000
Greene	8,000	6,000	2,000
Grenada	18,000	9,000	9,000
Hancock	13,000	10,000	3,000
Harrison	166,000	136,000	30,000
Hinds	177,000	104,000	73,000
Holmes	23,000	6,000	17,000
Humphreys	16,000	6,000	10,000
Issaquena	4,000	2,000	2,000
Itawamba	21,000	19,500	1,500
Jackson	67,000	52,000	15,000
Jasper	18,000	10,000	8,000
Jefferson	12,000	3,000	9,000
Jeff Davis	19,000	9,000	10,000
Jones	72,000	55,000	17,000
Lafayette	26,000	18,000	8,000

<u>County</u>	<u>Total</u>	<u>White</u>	<u>Nonwhite</u>
Lamar	18,000	14,000	4,000
Lauderdale	60,000	37,000	23,000
Lawrence	12,000	8,000	4,000
Leake	20,000	11,000	9,000
Lee	42,000	31,000	11,000
Leflore	40,000	16,000	24,000
Lincoln	26,000	17,000	9,000
Lowndes	41,000	26,000	15,000
Madison	27,000	9,000	18,000
Marion	28,000	17,000	11,000
Marshall	24,000	7,000	17,000
Monroe	35,000	22,000	13,000
Montgomery	13,000	8,000	5,000
Neshoba	26,000	18,000	8,000
Newton	21,000	14,000	7,000
Noxubee	14,000	5,000	9,000
Oktibbeha	28,000	16,000	12,000
Panola	29,000	14,000	15,000
Pearl River	29,000	20,000	9,000
Perry	9,000	7,000	2,000
Pike	32,000	18,000	14,000
Pontotoc	22,000	18,000	4,000
Prentiss	17,000	13,000	4,000
Quitman	20,000	7,000	13,000
Rankin	43,000	28,000	15,000
Scott	22,000	14,000	8,000
Sharkey	9,000	4,000	5,000
Simpson	27,000	17,000	10,000
Smith	17,000	12,000	5,000
Stone	8,000	6,500	1,500
Sunflower	44,000	17,000	27,000
Tallahatchie	22,000	10,000	12,000
Tate	18,000	9,000	9,000
Tippah	21,000	15,000	6,000
Tishomingo	15,000	14,500	500
Tunica	16,000	3,000	13,000
Union	24,000	18,000	6,000
Walthall	14,000	8,000	6,000
Warren	36,000	18,000	18,000
Washington	61,000	30,000	31,000
Wayne	17,000	11,000	6,000
We s ster	11,000	8,000	3,000
Wilkinson	11,000	3,000	8,000
Winston	20,000	11,000	9,000
Yalobusha	13,000	7,000	6,000
Yazoo	29,000	11,000	18,000

* Prepared by Dr. George L. Wilber and Ellen Bryant, Sociology Department, Mississippi State University.

TABLE 2
LIVE BIRTHS
MISSISSIPPI
1959

By Place of Occurrence, Reallocated to Mother's Usual Residence,
by Race, with Rates

County	Place of Occurrence			Place of Residence			Rate*		
	Number		Non-white	Number		Non-white	Rate*		Non-white
	Total	White		Total	White		Total	White	
STATE	61,157	28,448	32,709	60,562	27,979	32,583	27.2	20.9	36.6
Adams	1,663	665	998	1,200	508	692	38.7	28.2	53.2
Alcorn	599	495	104	522	419	103	20.1	21.0	17.2
Amite	232	36	196	360	144	246	22.5	14.2	30.8
Attala	522	222	300	499	197	302	25.0	17.9	33.6
Benton	96	3	93	192	58	134	19.2	11.6	26.8
Bolivar	1,826	335	1,491	1,735	355	1,380	42.3	20.9	57.5
Calhoun	384	194	190	350	196	154	23.3	17.0	44.0
Carroll	211	28	183	248	53	195	22.5	8.8	39.0
Chickasaw	413	203	210	455	213	242	28.4	26.6	30.2
Choctaw	197	102	95	199	104	95	19.9	14.8	31.7
Claiborne	244	29	215	310	46	264	28.2	23.0	29.3
Clarke	186	111	75	392	193	199	24.5	19.3	33.2
Clay	487	210	277	501	223	278	31.3	27.9	34.8
Coahoma	1,489	363	1,126	1,539	345	1,194	35.8	20.3	45.9
Copiah	875	297	578	719	252	467	25.7	18.0	33.4
Covington	248	184	64	356	208	148	22.2	20.8	24.7
DeSoto	713	100	613	708	171	537	32.2	19.0	41.3
Forrest	1,482	1,099	383	1,318	846	472	25.3	22.3	33.7
Franklin	121	56	65	205	91	114	22.8	18.2	28.5
George	334	282	52	336	281	55	22.4	21.6	27.5
Greene	145	94	51	208	143	65	26.0	23.8	32.5
Grenada	490	167	323	470	147	323	26.1	16.3	35.9
Hancock	165	132	33	334	279	55	25.7	27.9	18.3
Harrison	4,337	3,562	775	4,001	3,258	743	24.1	24.0	24.8
Hinds	5,739	3,002	2,737	5,333	2,659	2,674	30.1	25.6	36.6
Holmes	783	127	656	779	122	657	33.9	20.3	38.6
Humphreys	607	91	516	615	100	515	38.4	16.7	51.5
Issaquena	44	1	43	120	22	98	30.0	11.0	49.0
Itawamba	79	62	17	283	257	26	13.5	13.2	17.3
Jackson	1,393	987	406	1,730	1,296	434	25.8	24.9	28.9
Jasper	106	25	81	427	149	278	23.7	14.9	34.8
Jefferson	122	0	122	296	36	260	24.7	12.0	28.9
Jeff Davis	332	154	178	372	138	234	19.6	15.3	23.4
Jones	2,480	1,264	1,216	1,518	979	539	21.1	17.8	31.7
Kemper	206	44	162	286	77	209	23.8	15.4	29.8
Lafayette	510	266	244	532	266	266	20.5	14.8	33.2

Table 2 - Page 2
Live Births - Mississippi - 1959

County	Place of Occurrence			Place of Residence			Rate*		
	Number		Non-white	Number		Non-white	Rate*		Non-white
	Total	White		Total	White		Total	White	
Lamar	90	52	38	303	227	76	16.8	16.2	19.0
Lauderdale	2,230	1,108	1,122	1,693	875	818	28.2	23.6	35.6
Lawrence	174	28	146	282	133	149	23.5	16.6	37.2
Leake	379	174	205	453	200	253	22.6	18.2	28.1
Lee	1,179	848	331	973	631	342	23.2	20.4	31.1
Leflore	1,572	397	1,175	1,527	341	1,186	38.2	21.3	49.4
Lincoln	689	415	274	663	367	296	25.5	21.6	32.9
Lowndes	1,509	886	623	1,423	793	630	34.7	30.5	42.0
Madison	1,084	196	888	1,063	190	873	39.4	21.1	48.5
Marion	504	283	221	586	329	257	20.9	19.4	23.4
Marshall	751	117	634	784	146	638	32.7	20.8	37.5
Monroe	822	408	414	832	436	396	23.8	19.8	30.5
Montgomery	333	127	206	334	123	211	25.7	15.4	42.2
Neshoba	492	280	212	499	279	220	19.2	15.5	27.5
Newton	376	257	119	435	220	215	20.7	15.7	30.7
Noxubee	554	96	458	540	90	450	38.6	18.0	50.0
Oktibbeha	822	347	475	786	328	458	28.1	20.5	38.2
Panola	785	223	562	814	243	571	28.1	17.4	38.1
Pearl River	656	480	176	534	356	178	18.4	17.8	19.8
Perry	101	76	25	232	169	63	25.8	24.1	31.5
Pike	990	446	544	875	332	543	27.3	18.4	38.8
Pontotoc	294	218	76	288	209	79	13.1	11.6	19.8
Prentiss	455	371	84	379	301	78	22.3	23.2	19.5
Quitman	734	210	524	661	168	493	33.0	24.0	37.9
Rankin	305	77	228	757	429	328	17.6	15.3	21.9
Scott	634	360	274	578	293	285	26.3	20.9	35.6
Sharkey	249	46	203	340	62	278	37.8	15.5	55.6
Simpson	456	234	222	461	260	201	17.1	15.3	20.1
Smith	86	54	32	334	189	145	19.6	15.8	29.0
Stone	160	112	48	153	95	58	19.1	14.6	38.7
Sunflower	1,357	251	1,106	1,444	277	1,167	32.8	16.3	43.2
Tallahatchie	730	131	599	773	175	598	35.1	17.5	49.8
Tate	450	29	421	580	126	454	32.2	14.0	50.4
Tippah	398	294	104	334	245	89	15.9	16.3	14.8
Tishomingo	110	98	12	252	236	16	16.8	16.3	32.0
Tunica	672	68	604	658	67	591	41.1	22.3	45.5
Union	434	335	99	405	304	101	16.9	16.9	16.8
Walthall	604	320	284	362	138	224	25.8	17.2	37.3
Warren	1,496	616	880	1,192	533	659	33.1	29.6	36.6
Washington	2,830	1,239	1,591	2,727	1,178	1,549	44.7	39.3	50.0
Wayne	509	339	170	486	280	206	28.6	25.4	34.3
Webster	167	94	73	224	145	79	20.4	18.1	26.3
Wilkinson	465	194	271	386	69	317	35.1	23.0	39.6
Winston	429	167	262	474	190	284	23.7	17.3	31.6
Yalobusha	311	104	207	313	111	202	24.1	15.8	33.7
Yazoo	870	251	619	922	290	632	31.8	26.4	35.1
Out-of-State	943	782	161	1,538	1,251	287	-	-	-

Cities over 10,000 <u>Population**</u>	Place of Occurrence			Place of Residence		
	Number		Non- white	Number		Non- white
	Total	White		Total	White	
Biloxi	2,675	2,409	266	1,771	1,551	220
Clarksdale	840	359	481	614	241	373
Columbus	985	590	395	756	410	346
Greenville	1,446	729	717	1,366	619	747
Greenwood	818	392	426	578	205	368
Gulfport	1,521	1,146	375	1,011	721	290
Hattiesburg	1,441	1,095	346	839	493	346
Jackson	5,327	2,999	2,328	3,328	1,469	1,859
Laurel	2,412	1,222	1,190	795	428	367
McComb	645	332	313	270	135	135
Meridian	2,145	1,099	1,046	1,160	624	536
Natchez	748	597	151	754	310	444
Pascagoula	78	2	76	593	417	176
Tupelo	919	724	195	375	247	128
Vicksburg	1,449	615	834	850	400	450
Total Urban#	34,711	20,909	13,802	21,212	11,440	9,772
Total Rural	26,446	7,539	18,907	39,350	16,539	22,811

* Rate per 1,000 population.

** Cities having 10,000 population and over according to 1950 census.

Includes all towns of 2,500 population or more according to 1950 census.

TABLE 3
LIVE BIRTHS
MISSISSIPPI
1959
By County and Cities over 10,000, by Race, Attendant, and Sex
on Basis of Usual Residence of Mother

County	TOTAL*	WHITE			NONWHITE			SEX			
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	
STATE	<u>60,562</u>	<u>27,979</u>	<u>27,666</u>	<u>187</u>	<u>126</u>	<u>32,583</u>	<u>15,998</u>	<u>1,915</u>	<u>14,670</u>	<u>30,755</u>	<u>29,807</u>
Adams	1,200	508	506	0	2	692	665	6	21	611	589
Alcorn	522	419	401	16	2	103	64	18	24	284	238
Amitte	360	114	113	1	0	246	83	0	163	187	173
Attala	499	197	195	2	0	302	87	28	187	261	238
Benton	192	58	54	3	1	134	44	6	84	87	105
Bolivar	1,735	355	343	8	4	1,380	574	82	724	881	854
Calhoun	350	196	192	4	0	154	99	7	48	186	164
Carroll	248	53	52	1	0	195	51	4	140	116	132
Chickasaw	455	213	208	5	0	242	110	30	102	229	226
Choctaw	199	104	103	1	0	95	49	2	44	94	105
Claiborne	310	46	46	0	0	264	69	1	194	150	160
Clarke	392	193	190	3	0	199	173	5	21	199	193
Clay	501	223	222	0	1	278	127	40	111	242	259
Coahoma	1,539	345	339	3	3	1,194	410	98	686	752	787
Copiah	719	252	252	0	0	467	337	10	120	363	356
Covington	356	208	206	1	1	148	113	3	32	172	184
DeSoto	708	171	170	0	0	537	215	11	311	367	341
Forrest	1,318	846	839	0	7	472	320	6	146	667	651
Franklin	205	91	90	1	0	114	83	2	29	96	109
George	336	281	277	2	2	55	38	3	14	177	159
Greene	208	143	138	4	1	65	26	2	37	120	88
Grenada	470	147	145	1	1	323	40	75	208	231	239
Hancock	334	279	276	1	2	55	23	0	32	150	184
Harrison	4,001	3,258	3,246	2	10	743	398	45	300	2,087	1,914

Table 3 - Page 2
Live Births - Mississippi - 1959

County	TOTAL*	WHITE			NONWHITE			SEX	
		Total	Phys. in Hosp.	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male
Hinds	5,333	2,659	2,657	2	0	2,674	2,034	123	517
Holmes	779	122	120	1	1	657	94	38	525
Humphreys	615	100	98	1	1	515	104	2	409
Issaquena	120	22	20	1	1	98	56	0	42
Itawamba	283	257	247	10	0	26	24	0	2
Jackson	1,730	1,296	1,286	1	9	434	298	59	117
Jasper	427	149	146	3	0	278	212	3	63
Jefferson	296	36	36	0	0	260	137	3	120
Jeff Davis	372	138	137	1	0	234	146	2	86
Jones	1,518	979	976	0	3	539	499	7	33
Kemper	286	77	77	0	0	209	90	4	115
Lafayette	532	266	261	2	3	266	60	15	191
Lamar	303	227	224	3	0	76	55	11	10
Lauderdale	1,693	875	872	2	1	818	677	37	104
Lawrence	282	133	125	4	4	149	69	2	78
Leake	453	200	197	1	2	253	114	1	138
Lee	973	631	624	7	0	342	191	92	59
Leflore	1,527	341	338	2	2	1,186	158	60	968
Lincoln	663	367	366	0	1	296	222	17	57
Lowndes	1,423	793	783	4	6	630	252	40	338
Madison	1,063	190	189	1	0	873	251	42	580
Marion	586	329	324	4	1	257	108	14	135
Marshall	784	146	138	2	6	638	119	90	429
Monroe	832	436	430	4	2	396	174	55	167
Montgomery	334	123	122	0	1	211	64	4	143
Neshoba	499	279	276	1	2	220	139	2	79
Newton	435	220	218	2	0	215	163	24	28
Noxubee	540	90	90	0	0	450	98	3	349
Oktibbeha	786	328	325	3	0	458	281	1	176
Panola	814	243	240	2	1	571	272	75	224

Table 3 - Page 3
Live Births - Mississippi - 1959

County	TOTAL*	WHITE			NONWHITE			SEX			
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female
Pearl River	534	356	355	0	1	178	170	5	3	283	251
Perry	232	169	168	1	0	63	52	2	9	121	111
Pike	875	332	331	1	0	543	388	5	150	437	438
Pontotoc	288	209	207	1	1	79	36	7	36	156	132
Prentiss	379	301	301	0	0	78	50	5	23	207	172
Quitman	661	168	164	3	1	493	280	35	178	330	331
Rankin	757	429	427	0	2	328	257	10	61	386	371
Scott	578	293	290	2	1	285	156	3	126	293	285
Sharkey	340	62	60	1	1	278	152	0	126	176	164
Simpson	461	260	259	1	0	201	136	4	61	236	225
Smith	334	189	183	5	1	145	113	13	19	175	159
Stone	153	95	94	1	0	58	22	0	36	72	81
Sunflower	1,444	277	268	3	6	1,167	341	163	663	683	761
Tallahatchie	773	175	165	6	4	598	111	55	432	382	391
Tate	580	126	119	7	0	454	54	34	366	301	279
Tippah	334	245	235	8	2	89	49	27	13	158	176
Tishomingo	252	236	229	7	0	16	11	5	0	132	120
Tunica	658	67	62	3	2	591	168	20	403	329	329
Union	405	304	294	1	9	101	29	7	65	208	197
Walthall	362	138	134	1	3	224	183	0	41	193	169
Warren	1,192	533	532	0	1	659	461	0	198	606	586
Washington	2,727	1,178	1,176	2	0	1,549	881	111	557	1,363	1,364
Wayne	486	280	279	0	1	206	148	1	57	232	254
Webster	224	145	138	7	0	79	36	10	33	103	121
Wilkinson	386	69	68	1	0	317	120	2	195	204	182
Winston	474	190	188	0	2	284	99	0	185	245	229
Yalobusha	313	111	110	1	0	202	65	70	67	155	158
Yazoo	922	290	285	1	4	632	111	11	510	451	471

Table 3 - Page 4
Live Births - Mississippi - 1959

Cities over 10,000 Population	TOTAL*	WHITE			NONWHITE			SEX			
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female
Biloxi	1,771	1,551	1,548	1	2	220	167	2	51	939	832
Clarksdale	614	241	241	0	0	373	222	49	102	307	307
Columbus	756	410	407	2	1	346	130	20	196	380	376
Greenville	1,366	619	617	2	0	747	391	72	284	685	681
Greenwood	573	205	204	0	1	368	79	1	288	299	274
Gulfport	1,011	721	718	0	3	290	144	37	109	510	501
Hattiesburg	839	493	489	0	4	346	236	5	105	413	426
Jackson	3,328	1,469	1,469	0	0	1,859	1,568	101	190	1,668	1,660
Laurel	795	428	427	0	1	367	350	5	12	396	399
McComb	270	135	134	1	0	135	121	2	12	137	133
Meridian	1,160	624	623	1	0	536	474	33	29	587	573
Natchez	754	310	308	0	2	444	431	4	9	377	377
Pascagoula	593	417	415	0	2	176	99	19	58	309	284
Tupelo	375	247	247	0	0	128	74	36	18	181	194
Vicksburg	850	400	400	0	0	450	311	0	139	420	430

* Includes transcripts furnished by the National Office of Vital Statistics of births to residents of Mississippi occurring out-of-state.

TABLE 4
FETAL DEATHS
MISSISSIPPI
1959

By Place of Occurrence, Reallocated to Mother's Usual Residence
by Sex, Race and Attendant

County	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX			Unde- termined	
			Total	Phys. in Hosp.	Other	Total	Phys. in Hosp.	Phys. in Home	Male	Female			
STATE	<u>2,649</u>	<u>2,603</u>	<u>1,045</u>	<u>986</u>	<u>53</u>	<u>6</u>	<u>1,558</u>	<u>928</u>	<u>145</u>	<u>485</u>	<u>1,024</u>	<u>806</u>	<u>773</u>
Adams	51	32	7	7	0	0	25	25	0	0	17	13	2
Alcorn	19	15	12	12	0	0	3	1	1	1	6	9	0
Amité	2	8	3	3	0	0	5	3	0	2	1	5	2
Attala	23	17	3	3	0	0	14	9	1	4	8	5	4
Benton	6	15	2	2	0	0	13	7	3	3	5	6	4
Bolivar	79	76	15	15	0	0	61	31	6	24	35	34	7
Calhoun	6	6	1	1	0	0	5	4	0	1	3	2	1
Carroll	9	13	1	1	0	0	12	6	2	4	7	3	3
Chickasaw	17	19	7	6	1	0	12	7	2	3	8	8	3
Choctaw	3	4	3	3	0	0	1	1	0	0	1	2	1
Claiborne	8	8	1	1	0	0	7	1	1	5	4	3	1
Clarke	7	18	5	4	1	0	13	13	0	0	12	4	2
Clay	9	12	2	2	0	0	10	9	1	0	5	7	0
Coahoma	69	67	8	8	0	0	59	30	11	18	32	25	10
Copiah	23	20	2	1	1	0	18	11	2	5	7	12	1
Covington	5	8	4	3	1	0	4	3	0	1	4	4	0
DeSoto	19	30	3	3	0	0	27	19	1	7	18	10	2
Forrest	53	47	22	21	1	0	25	21	1	3	16	16	15
Franklin	7	9	3	3	0	0	6	4	1	1	3	5	1
George	8	8	5	5	0	0	3	2	1	0	3	5	0
Greene	6	6	4	4	0	0	2	1	0	1	5	1	0
Grenada	26	28	9	9	0	0	19	6	6	7	18	7	3
Hancock	4	6	4	4	0	0	2	1	0	1	1	3	2
Harrison	102	94	56	55	0	1	38	20	7	11	48	37	9

Table 4 - Page 2
Fetal Deaths - Mississippi - 1959

County	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total	Phys. in Hosp.	Other	Total	Phys. in Hosp.	Other	Male	Female	Undetermined
Hinds	390	345	230	203	25	2	115	89	13	78	48
Holmes	59	59	12	10	2	0	47	16	8	23	21
Humphreys	25	27	1	1	0	0	26	5	2	19	11
Issaquena	1	4	1	1	0	0	3	2	0	1	1
Itawamba	1	7	7	7	0	0	0	0	0	0	3
Jackson	80	86	55	50	3	2	31	21	9	1	31
Jasper	3	20	7	7	0	0	13	10	1	2	5
Jefferson	8	18	2	2	0	0	16	8	0	8	7
Jeff Davis	13	14	1	1	0	0	13	5	5	3	2
Jones	83	46	25	24	1	0	21	15	3	3	11
Kemper	10	12	1	1	0	0	11	3	0	8	2
Lafayette	26	26	8	8	0	0	18	4	2	12	9
Lamar	1	8	6	6	0	0	2	2	0	0	5
Lauderdale	93	68	30	29	1	0	38	33	1	4	37
Lawrence	8	11	3	2	1	0	8	5	0	3	6
Leake	11	13	0	0	0	0	13	7	0	6	7
Lee	75	61	48	48	0	0	13	12	1	0	14
Leflore	103	94	25	25	0	0	69	31	7	31	36
Lincoln	34	28	11	11	0	0	17	13	2	2	14
Lowndes	46	43	18	17	1	0	25	13	2	10	17
Madison	41	40	6	6	0	0	34	11	7	16	20
Marion	13	20	9	9	0	0	11	7	0	4	11
Marshall	46	34	4	4	0	0	30	7	3	20	13
Monroe	30	29	9	9	0	0	20	13	2	5	10
Montgomery	11	12	3	3	0	0	9	4	0	5	9
Neshoba	14	14	4	4	0	0	10	7	3	0	10
Newton	37	33	17	17	0	0	16	13	2	1	4
Noxubee	30	28	2	2	0	0	26	13	0	13	12
Oktibbeha	23	19	4	4	0	0	15	11	0	4	6
Panola	21	29	4	4	0	0	25	15	2	8	15

Table 4 - Page 3
Fetal Deaths - Mississippi - 1959

County	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total	Phys. in Hosp.	Phys. in Home	Total	Phys. in Hosp.	Phys. in Home	Male	Female	Undetermined
Pearl River	119	17	10	8	2	0	7	7	0	8	5
Perry	5	13	8	7	1	0	5	4	1	6	1
Pike	30	27	5	5	0	0	22	18	1	3	10
Pontotoc	12	13	10	0	0	0	3	2	1	2	5
Prentiss	6	5	5	5	0	0	0	0	0	3	2
Quitman	18	19	3	3	0	0	16	9	4	3	10
Rankin	19	..	46	20	16	4	0	26	22	0	4
Scott	10	19	5	5	0	0	14	12	0	2	8
Sharkey	10	15	2	2	0	0	13	6	0	7	7
Simpson	9	10	8	7	1	0	2	2	0	6	2
Smith	2	8	5	5	0	0	3	2	0	1	2
Stone	13	12	7	7	0	0	5	3	0	2	6
Sunflower	60	65	11	11	0	0	54	22	5	27	31
Tallahatchie	37	38	11	8	3	0	27	14	1	12	22
Tate	10	17	2	1	0	1	15	6	0	9	5
Tippah	10	13	10	10	0	0	3	1	2	0	3
Tishomingo	3	10	10	9	1	0	0	0	0	0	7
Tunica	25	25	4	3	1	0	21	5	0	16	8
Union	50	35	25	25	0	0	10	7	1	2	8
Walthall	26	15	3	3	0	0	12	9	0	3	9
Warren	85	48	8	8	0	0	40	32	0	8	20
Washington	207	199	109	109	0	0	90	68	4	18	64
Wayne	22	32	14	13	1	0	0	18	12	2	4
Webster	3	5	0	0	0	0	5	4	0	1	4
Wilkinson	11	14	1	1	0	0	0	0	13	8	0
Winston	10	14	3	3	0	0	11	5	0	6	12
Yalobusha	4	8	5	5	0	0	3	1	1	3	4
Yazoo	36	47	6	6	0	0	41	17	2	22	19

Table 4 - Page 4
Fetal Deaths - Mississippi - 1959

Cities over 10,000 Population	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total	Phys. in Hosp.	Other	Total	Phys. in Hosp.	Other	Male	Female	Unde- termined
Biloxi	60	44	36	36	0	8	6	1	23	15	6
Clarksdale	45	25	7	7	0	18	14	3	11	7	7
Columbus	29	20	9	8	1	0	11	7	1	3	5
Greenville	126	140	93	93	0	0	47	40	3	4	100
Greenwood	75	44	15	15	0	0	29	18	2	9	20
Gulfport	39	33	9	9	0	0	24	9	6	9	2
Hattiesburg	53	37	17	16	1	0	20	16	1	3	11
Jackson	373	231	147	130	16	1	84	66	11	7	143
Laurel	79	25	13	13	0	0	12	9	2	1	5
McComb	21	11	3	3	0	0	8	6	1	3	3
Meridian	93	41	20	19	1	0	21	18	1	2	4
Natchez	23	25	5	5	0	0	20	20	0	0	2
Pascagoula	6	36	23	21	2	0	13	9	3	1	19
Tupelo	73	31	24	24	0	0	7	6	1	0	18
Vicksburg	82	34	6	6	0	0	28	24	0	4	2
									16	16	

TABLE 5
DEATHS
MISSISSIPPI
1959

By Cause by Race on Basis of Place of Residence

International List Number	Cause of Death*	Total	White	Non-white
<u>001-999</u>	<u>TOTAL ALL CAUSES</u>	<u>20,890</u>	<u>10,742</u>	<u>10,148</u>
<u>001-019</u>	<u>Tuberculosis, all forms</u>	<u>153</u>	<u>62</u>	<u>91</u>
002	Pulmonary	135	58	77
003	Pleural	3	1	2
008	Unspecified site	4	1	3
010	Meninges and central nervous system	8	0	8
012	Bones and joints, active or unspecified	1	0	1
013	Late effects of tuberculosis of bones and joints	1	1	0
017	Adrenal glands	1	1	0
<u>020-029</u>	<u>Syphilis and its sequelae</u>	<u>43</u>	<u>9</u>	<u>34</u>
020	Congenital	1	1	0
022	Aneurysm of aorta	12	2	10
023	Other cardiovascular	17	3	14
024	Tabes dorsalis	3	1	2
025	General paralysis of insane	1	0	1
026	Other central nervous system	3	1	2
027	Other forms of late syphilis	3	1	2
028	Latent syphilis	1	0	1
029	Unqualified	2	0	2
035	Late effects of gonococcal infection	2	0	2
038	Granuloma inguinale, venereal	1	0	1
041	Paratyphoid fever	1	0	1
<u>045-048</u>	<u>Dysentery, all forms</u>	<u>26</u>	<u>4</u>	<u>22</u>
045	Bacillary	11	1	10
046	Amebiasis	8	2	6
048	Unspecified forms	7	1	6
049	Food poisoning	2	0	2
<u>050,051</u>	<u>Scarlet fever and streptococcal sore throat</u>	<u>2</u>	<u>0</u>	<u>2</u>
051	Streptococcal sore throat	2	0	2
053	Septicemia and pyemia	27	12	15
055	Diphtheria	4	0	4
056	Whooping cough	18	1	17
057	Meningococcal infections	8	4	4
061	Tetanus	11	1	10
070	Vincent's infection	1	0	1
080	Acute poliomyelitis	9	4	5
082	Acute infectious encephalitis	9	3	6
083	Late effects of acute infectious encephalitis	2	1	1
085	Measles	8	2	6
087	Chickenpox	2	0	2
088	Herpes zoster	1	1	0
089	Mumps	2	1	1
092	Infectious hepatitis	19	12	7
096	Other diseases attributable to viruses	10	2	8

Table 5 - Page 2
Deaths - Mississippi - 1959

International List Number	Cause of Death*	Total	White	Non-white
122	Other protozoal diseases	2	1	1
134	Other fungus infections	5	4	1
138	Other infective and parasitic diseases	4	2	2
<u>140-205</u>	<u>Malignant neoplasms</u>	<u>2,600</u>	<u>1,580</u>	<u>1,020</u>
<u>140-148</u>	<u>Buccal cavity and pharynx</u>	<u>42</u>	<u>31</u>	<u>11</u>
140	Lip	4	3	1
141	Tongue	6	4	2
142	Salivary gland	4	4	0
143	Floor of mouth	1	1	0
144	Other parts of mouth, and mouth, unspecified	7	3	4
145	Oral mesopharynx	2	2	0
146	Nasopharynx	4	3	1
147	Hypopharynx	1	1	0
148	Pharynx, unspecified	13	10	3
<u>150-159</u>	<u>Digestive organs and peritoneum</u>	<u>842</u>	<u>512</u>	<u>330</u>
150	Esophagus	40	25	15
151	Stomach	238	108	130
152	Small intestine, including duodenum	7	3	4
153	Large intestine, except rectum	210	142	68
154	Rectum	81	51	30
155	Biliary passages and liver (stated to be primary site)	37	27	10
156	Liver (secondary and unspecified)	83	49	34
157	Pancreas	125	94	31
158	Peritoneum	7	5	2
159	Unspecified digestive organs	14	8	6
<u>160-165</u>	<u>Respiratory system</u>	<u>345</u>	<u>263</u>	<u>82</u>
160	Nose, nasal cavities, middle ear, and accessory sinuses	6	5	1
161	Larynx	22	16	6
162	Bronchus and trachea, and lung specified as primary	125	99	26
163	Lung, unspecified as to whether primary or secondary	183	138	45
164	Mediastinum	3	1	2
165	Thoracic organs (secondary)	6	4	2
<u>170-181</u>	<u>Breast and genito-urinary organs</u>	<u>781</u>	<u>391</u>	<u>390</u>
170	Breast	171	104	67
171	Cervix uteri	120	47	73
172	Corpus uteri	8	2	6
174	Uterus, unspecified	90	26	64
175	Ovary, Fallopian tube, and broad ligament	71	47	24
176	Other and unspecified female genital organs	8	3	5
177	Prostate	210	87	123
178	Testis	5	4	1
179	Other and unspecified male genital organs	5	0	5
180	Kidney	40	35	5
181	Bladder and other urinary organs	53	36	17
<u>190-199</u>	<u>Other and unspecified sites</u>	<u>350</u>	<u>208</u>	<u>142</u>
190	Melanoma of skin	15	14	1
191	Skin, other	29	23	6
192	Eye	4	4	0
193	Brain and other parts of nervous system	60	47	13

Table 5 - Page 3
 Deaths - Mississippi - 1959

International List Number	Cause of Death*	Total	White	Non-white
194	Thyroid gland	9	5	4
195	Other endocrine glands	2	2	0
196	Bone	24	16	8
197	Connective tissue	6	3	3
198	Lymphnodes (secondary and unspecified)	3	3	0
199	Other and unspecified sites	198	91	107
<u>200-205</u>	<u>Lymphatic and hematopoietic tissues</u>	<u>240</u>	<u>175</u>	<u>65</u>
200	Lymphosarcoma and reticulosarcoma	54	43	11
201	Hodgkin's disease	27	17	10
202	Other forms of lymphoma (reticulososis)	6	2	4
203	Multiple myeloma (plasmacytoma)	26	18	8
204	Leukemia and aleukemia	125	94	31
205	Mycosis fungoides	2	1	1
<u>210-229</u>	<u>Benign neoplasm</u>	<u>20</u>	<u>10</u>	<u>10</u>
211	Other parts of digestive system	1	0	1
214	Uterine fibromyoma	4	0	4
215	Uterus, other	1	0	1
216	Ovary	2	1	1
223	Brain and other parts of nervous system	4	2	2
224	Endocrine glands	5	5	0
226	Lipoma	1	0	1
228	Hemangioma and lymphangioma	2	2	0
<u>230-239</u>	<u>Neoplasms of unspecified nature</u>	<u>28</u>	<u>16</u>	<u>12</u>
230	Digestive organs	1	0	1
231	Respiratory organs	5	3	2
237	Brain and other parts of nervous system	19	12	7
238	Skin and musculoskeletal system	1	0	1
239	Other and unspecified organs	22	11	1
240	Hay fever	1	0	1
241	Asthma	52	26	26
245	Other allergic disorders	4	3	1
251	Non-toxic nodular goiter	1	0	1
252	Thyrotoxicosis with or without goiter	5	2	3
254	Other diseases of thyroid gland	2	0	2
260	Diabetes mellitus	281	128	153
274	Diseases of adrenal glands	5	3	2
277	Polyglandular dysfunction and other diseases of endocrine glands	2	1	1
280	Beriberi	1	1	0
281	Pellagra	33	0	3
282	Scurvy	1	0	1
283	Active rickets	2	0	2
286	Other avitaminoses and nutritional deficiency states	30	15	15
287	Obesity, not specified as of endocrine origin	3	1	2
288	Gout	2	2	0
289	Other metabolic diseases	11	10	1
<u>290-293</u>	<u>Anemias</u>	<u>48</u>	<u>21</u>	<u>27</u>
290	Pernicious and other hyperchromic	6	5	1
292	Other, specified type	23	9	14
293	Unspecified type	19	7	12
296	Purpura and other hemorrhagic conditions	7	6	1
297	Agranulocytosis	1	1	0

Table 5 - Page 4
Deaths - Mississippi - 1959

International List Number	Cause of Death*	Total	White	Non-white
299	Other diseases of blood and blood-forming organs	2	1	1
300	Schizophrenia disorders	1	1	0
304	Senile psychosis	7	5	2
305	Presenile psychosis	2	2	0
307	Alcoholic psychosis	5	5	0
311	Hysterical reaction without mention of anxiety reaction	1	1	0
312	Phobic reaction	1	1	0
322	Alcoholism	14	7	7
325	Mental deficiency	1	1	0
<u>330-334</u>	<u>Vascular lesions affecting central nervous system</u>	<u>2,683</u>	<u>1,386</u>	<u>1,297</u>
330	Subarachnoid hemorrhage	66	29	37
331	Cerebral hemorrhage	1,682	825	857
332	Cerebral embolism and thrombosis	563	357	206
334	Other and ill-defined	372	175	197
340	Meningitis, except meningococcal and tuberculous	46	16	30
342	Intracranial and intraspinal abscess	4	2	2
343	Encephalitis, myelitis, and encephalomyelitis (except acute infectious)	10	4	6
344	Late effects of intracranial abscess or pyogenic infection	13	8	5
345	Multiple sclerosis	10	5	5
350	Paralysis agitans	21	20	1
351	Cerebral spastic infantile paralysis	4	3	1
352	Other cerebral paralysis	12	7	5
353	Epilepsy	20	6	14
355	Other diseases of brain	13	9	4
356	Motor neurone disease and muscular atrophy	9	6	3
357	Other diseases of spinal cord	3	2	1
364	Polyneuritis and polyradiculitis	1	0	1
367	Other diseases of cranial nerves	1	0	1
389	Blindness	1	1	0
391	Otitis media without mention of mastoiditis	3	3	0
<u>400-402</u>	<u>Rheumatic fever</u>	<u>14</u>	<u>2</u>	<u>12</u>
400	Without mention of heart involvement	3	0	3
401	With heart involvement	10	1	9
402	Chorea	1	1	0
<u>410-443</u>	<u>Diseases of heart</u>	<u>6,219</u>	<u>3,772</u>	<u>2,447</u>
410	Diseases of mitral valve	51	22	29
411	Diseases of aortic valve specified as rheumatic	4	4	0
414	Other endocarditis specified as rheumatic	5	3	2
415	Other myocarditis specified as rheumatic	4	1	3
416	Other heart disease specified as rheumatic	63	33	30
420	Arteriosclerotic heart disease, including coronary disease	3,825	2,773	1,052
421	Chronic endocarditis not specified as rheumatic	75	34	41
422	Other myocardial degeneration	465	222	243
430	Acute and subacute endocarditis	9	4	5

Table 5 - Page 5
 Deaths - Mississippi - 1959

International List Number	Cause of Death*	Total	White	Non-white
431	Acute myocarditis not specified as rheumatic	4	3	1
433	Functional disease of heart	56	30	26
434	Other and unspecified diseases of heart	373	162	211
441	Essential malignant hypertension with heart disease	14	6	8
442	Hypertensive heart disease with arteriolar nephrosclerosis	250	99	151
443	Other and unspecified hypertensive heart disease	1,021	376	645
<u>444-447</u>	<u>Hypertension without mention of heart</u>	<u>255</u>	<u>96</u>	<u>159</u>
444	Essential benign	79	23	56
445	Essential malignant hypertension	74	13	61
446	With arteriolar nephrosclerosis	99	58	41
447	Other hypertensive disease	3	2	1
450	General arteriosclerosis	294	193	101
<u>451-468</u>	<u>Other diseases of circulatory system</u>	<u>216</u>	<u>128</u>	<u>88</u>
451	Aortic aneurysm, nonsyphilitic, and dissecting aneurysm	62	47	15
452	Other aneurysm, except of heart and aorta	10	7	3
453	Peripheral vascular disease	3	0	3
454	Arterial embolism and thrombosis	8	4	4
455	Gangrene of unspecified cause	5	1	4
456	Other diseases of arteries	9	6	3
460	Varicose veins of lower extremities	5	3	2
462	Varicose veins of other specified sites	3	2	1
463	Phlebitis and thrombophlebitis of lower extremities	10	7	3
464	Phlebitis and thrombophlebitis of other sites	20	12	8
465	Pulmonary embolism and infarction	58	26	32
466	Other venous embolism and thrombosis	11	9	2
467	Other diseases of circulatory system	12	4	8
470	Acute nasopharyngitis (common cold)	3	0	3
472	Acute pharyngitis	2	1	1
473	Acute tonsilitis	2	1	1
474	Acute laryngitis and tracheitis	3	3	0
475	Acute upper respiratory infection of multiple or unspecified sites	2	0	2
<u>480-483</u>	<u>Influenza</u>	<u>82</u>	<u>25</u>	<u>57</u>
480	With pneumonia	544	13	41
481	With other respiratory manifestations, and influenza unqualified	25	11	14
482	With digestive manifestations, but without respiratory symptoms	2	1	1
483	With nervous manifestations, but without digestive or respiratory symptoms	1	0	1
<u>490-493</u>	<u>Pneumonia</u>	<u>686</u>	<u>303</u>	<u>383</u>
490	Lobar	151	62	89
491	Bronchopneumonia	256	124	132
492	Primary atypical	47	16	31
493	Other and unspecified	232	101	131
<u>500-502</u>	<u>Bronchitis</u>	<u>44</u>	<u>25</u>	<u>19</u>

Table 5 - Page 6
Deaths - Mississippi - 1959

International List Number	Cause of Death*	Total	White	Non- white
500	Acute	10	5	5
501	Unqualified	16	5	11
502	Chronic	18	15	3
510	Hypertrophy of tonsils and adenoids	2	2	0
511	Peritonsillar abscess	1	0	1
517	Other diseases of upper respiratory tract	2	0	2
518	Empyema	4	1	3
520	Spontaneous pneumothorax	3	3	0
521	Abscess of lung	4	2	2
522	Pulmonary congestion and hypostasis	25	15	10
523	Pneumoconiosis due to silica and silicates (occupational)	1	1	0
525	Other chronic interstitial pneumonia	17	14	3
526	Bronchiectasis	21	18	3
527	Other diseases of lung and pleural cavity	100	88	12
535	Other diseases of teeth and supporting structures	1	0	1
537	Diseases of salivary glands	1	1	0
539	Diseases of esophagus	10	7	3
<u>540, 541</u>	<u>Ulcer of stomach and duodenum</u>	<u>80</u>	<u>44</u>	<u>36</u>
540	Stomach	33	15	18
541	Ulcer of duodenum	47	29	18
543	Gastritis and duodenitis	3	2	1
544	Disorders of function of stomach	2	0	2
545	Other diseases of stomach and duodenum	4	3	1
<u>550-553</u>	<u>Appendicitis</u>	<u>26</u>	<u>12</u>	<u>14</u>
550	Acute	25	11	14
551	Unqualified	1	1	0
<u>560, 561, 570</u>	<u>Hernia and intestinal obstruction</u>	<u>122</u>	<u>57</u>	<u>65</u>
560	Hernia of abdominal cavity without mention of obstruction	13	11	2
561	Hernia of abdominal cavity with obstruction	17	5	12
570	Intestinal obstruction without mention of hernia	92	41	51
571	Gastro-enteritis and colitis, except ulcerative, age 4 weeks and over	175	25	150
572	Chronic enteritis and ulcerative colitis	21	17	4
573	Functional disorders of intestines	1	0	1
574	Anal fissure and fistula	1	1	0
575	Abscess of anal and rectal regions	1	0	1
576	Peritonitis	13	4	9
577	Peritoneal adhesion	1	0	1
578	Other diseases of intestines and peritoneum	28	17	11
580	Acute and subacute yellow atrophy of liver	6	3	3
581	Cirrhosis of liver	100	71	29
582	Suppurative hepatitis and liver abscess	1	0	1
583	Other diseases of liver	7	5	2
584	Cholelithiasis	22	15	7
585	Cholecystitis and cholangitis without mention of calculi	13	10	3
586	Other diseases of gallbladder and biliary ducts	10	7	3
587	Diseases of pancreas	34	25	9
<u>590, 591</u>	<u>Acute nephritis and nephritis with edema</u>	<u>51</u>	<u>12</u>	<u>39</u>

Table 5 - Page 7
 Deaths - Mississippi - 1959

International List Number	Cause of Death*	Total	White	Non-white
590	Acute	27	7	20
591	With edema, including nephrosis	24	5	19
<u>592-594</u>	<u>Chronic and unspecified nephritis and other renal sclerosis</u>	<u>239</u>	<u>89</u>	<u>150</u>
592	Chronic nephritis	197	71	126
593	Nephritis not specified as acute or chronic	40	18	22
594	Other renal sclerosis	2	0	2
600	Infections of kidney	108	61	47
601	Hydronephrosis	2	1	1
602	Calculi of kidney and ureter	9	6	3
603	Other diseases of kidney and ureter	17	5	12
604	Calculi of other parts of urinary system	2	1	1
605	Cystitis	6	2	4
606	Other diseases of bladder	4	1	3
608	Stricture of urethra	5	0	5
609	Other diseases of urethra	5	4	1
610	Hyperplasia of prostate	74	30	44
611	Prostatitis	15	9	6
612	Other diseases of prostate	1	1	0
617	Other diseases of male genital organs	3	1	2
621	Other diseases of breast	1	0	1
623	Chronic salpingitis and oophoritis	1	0	1
624	Salpingitis and oophoritis, unqualified	4	1	3
625	Other diseases of ovary and Fallopian tube	1	1	0
626	Diseases of parametrium and pelvic peritoneum (female)	2	1	1
631	Uterovaginal prolapse	3	3	0
633	Other diseases of uterus	4	2	2
634	Disorders of menstruation	1	0	1
<u>640-689</u>	<u>Diseases of pregnancy, childbirth, and puerperium</u>	<u>55</u>	<u>9</u>	<u>46</u>
642	Toxemias of pregnancy	18	2	16
645	Ectopic pregnancy	2	0	2
650	Abortion without mention of sepsis or toxemia	2	0	2
651	Abortion with sepsis	1	1	0
652	Abortion with toxemia, without mention of sepsis	2	0	2
660	Delivery without mention of complication	3	0	3
670	Delivery complicated by placenta previa or antepartum hemorrhage	4	0	4
671	Delivery complicated by retained placenta	2	0	2
672	Delivery complicated by other postpartum hemorrhage	6	2	4
674	Delivery complicated by disproportion or malposition of fetus	2	1	1
675	Delivery complicated by prolonged labor of other origin	3	0	3
681	Sepsis of childbirth and the puerperium	2	0	2
684	Puerperal pulmonary embolism	3	3	0
685	Puerperal eclampsia	3	0	3
686	Other forms of puerperal toxemia	1	0	1

Table 5 - Page 8
Deaths - Mississippi - 1959

International List Number	Cause of Death*	Total	White	Non-white
688	Other and unspecified complications of the puerperium	1	0	1
690	Boil and carbuncle	2	2	0
692	Other cellulitis and abscess without mention of lymphangitis	2	0	2
703	Other dermatitis	1	0	1
704	Pemphigus	1	1	0
705	Erythematous conditions	7	3	4
708	Fruitus and related conditions	1	0	1
710	Other hypertrophic and atrophic conditions of skin	3	2	1
715	Chronic ulcer of skin	7	5	2
720	Acute arthritis due to pyogenic organisms	1	0	1
722	Rheumatoid arthritis and allied conditions	19	16	3
723	Osteo-arthritis (arthrosis) and allied conditions	3	2	1
725	Arthritis, unspecified	6	1	5
730	Osteomyelitis and periostitis	1	1	0
733	Other diseases of bone	1	1	0
744	Other diseases of muscle, tendon, and fascia	8	3	5
<u>750-759</u>	<u>Congenital malformations</u>	<u>247</u>	<u>140</u>	<u>107</u>
750	Monstrosity	5	5	0
751	Spina bifida and meningocele	9	5	4
752	Congenital hydrocephalus	23	13	10
753	Other nervous system and sense organs	9	5	4
754	Circulatory system	132	80	52
756	Digestive system	20	8	12
757	Genito-urinary system	10	7	3
758	Bone and joint	5	4	1
759	Other and unspecified, not elsewhere classified	34	13	21
<u>760-762</u>	<u>Birth injuries, postnatal asphyxia and atelectasis</u>	<u>356</u>	<u>185</u>	<u>171</u>
760	Intracranial and spinal injury at birth	58	27	31
761	Other birth injury	80	41	39
762	Postnatal asphyxia and atelectasis	218	117	101
<u>763-768</u>	<u>Infections of newborn</u>	<u>142</u>	<u>24</u>	<u>118</u>
763	Pneumonia	96	20	76
764	Diarrhea	25	1	24
767	Umbilical sepsis	4	0	4
768	Other sepsis	17	3	14
<u>769-776</u>	<u>Other diseases peculiar to early infancy, and immaturity, unqualified</u>	<u>715</u>	<u>220</u>	<u>495</u>
769	Neonatal disorders arising from maternal toxemia	14	6	8
770	Hemolytic disease of newborn (erythroblastosis)	19	12	7
771	Hemorrhagic disease of newborn	10	2	8
772	Nutritional maladjustment	38	2	36
773	Ill-defined diseases peculiar to early infancy	139	66	73
774	Immaturity with mention of any other subsidiary condition	20	4	16

Table 5 - Page 9
Deaths - Mississippi - 1959

International List Number	Cause of Death*	Total	White	Non-white
776	Immaturity, unqualified	475	128	347
<u>780-795</u>	<u>Symptoms, senility, and ill-defined conditions</u>	<u>1,837</u>	<u>432</u>	<u>1,405</u>
780	Certain symptoms referable to nervous system and special senses	9	2	7
781	Other symptoms referable to nervous system and special senses	2	2	0
782	Symptoms referable to cardiovascular and lymphatic system	56	15	41
783	Symptoms referable to respiratory system	1	1	0
784	Symptoms referable to upper gastro-intestinal tract	2	1	1
785	Symptoms referable to abdomen and lower gastro-intestinal tract	4	2	2
786	Symptoms referable to genito-urinary system	3	1	2
788	Other general symptoms	6	2	4
792	Uremia unqualified	23	11	12
794	Senility without mention of psychosis	54	16	38
795	Ill-defined and unknown causes	1,677	379	1,298
<u>800-962</u>	<u>Accidents</u>	<u>1,501</u>	<u>837</u>	<u>664</u>
<u>800-802</u>	<u>Railway</u>	<u>14</u>	<u>4</u>	<u>10</u>
<u>810-835</u>	<u>Motor vehicle</u>	<u>588</u>	<u>401</u>	<u>187</u>
810	Collision with railway train, traffic	21	16	5
812	Traffic, to pedestrian	106	52	54
813	Traffic, to pedal cyclist	8	8	0
815	Traffic, to rider or passenger of motorcycle in collision with other motor vehicle	10	10	0
816	Other traffic, two or more motor vehicles	226	161	65
818	Traffic, collision with animal or animal-drawn vehicle	2	1	1
819	Traffic, collision with fixed or unspecified object	12	11	1
820	Traffic, while boarding and alighting	1	1	0
821	Noncollision traffic, to rider of motorcycle	1	1	0
822	Traffic, overturning in roadway	18	14	4
823	Traffic, running off roadway	119	85	34
824	Other noncollision traffic	18	6	12
825	Traffic, unspecified nature	31	26	5
830	Nontraffic, to pedestrian	8	5	3
835	Nontraffic, other and unspecified nature	7	4	3
845	Other nonmotor road vehicle	3	1	2
850	Submersion of occupant of small boat	20	16	4
851	Other water transport injury by submersion	5	4	1
857	Other specified water transport	10	10	0
860	Accident to personnel in military aircraft	15	15	0
866	Other and unspecified aircraft	12	11	1
871	Poisoning by barbituric acid and derivatives	2	2	0
872	Poisoning by aspirin and salicylates	1	1	0
881	Poisoning by petroleum products	1	1	0
882	Poisoning by industrial solvents	1	1	0

Table 5 - Page 10
Deaths - Mississippi - 1959

International List Number	Cause of Death*	Total	White	Non-white
8833	Poisoning by corrosive aromatics, acids, and caustic alkalis	1	0	1
888	Poisoning by other and unspecified solid and liquid substances	5	0	5
890	Poisoning by utility (illuminating) gas	1	1	0
891	Poisoning by motor vehicle exhaust gas	1	1	0
894	Poisoning by other specified gases and vapors	1	1	0
900	Fall on stairs	6	4	2
902	Other falls from one level to another	31	26	5
903	Fall on same level	12	10	2
904	Unspecified falls	68	51	17
910	Blow from falling or projected object	35	13	22
911	Caused by vehicle not elsewhere classified	2	0	2
912	Caused by machinery	41	22	19
913	Caused by cutting and piercing instruments	3	0	3
914	Caused by electric current	18	10	8
916	Caused by fire and explosion of combustible material	225	58	167
917	Caused by hot substance, corrosive liquid, and steam	9	4	5
919	Caused by firearm	94	46	48
921	Inhalation and ingestion of food causing obstruction or suffocation	29	17	12
922	Inhalation and ingestion of other object causing obstruction or suffocation	5	3	2
923	Foreign body entering other orifice	3	1	2
924	Mechanical suffocation in bed and cradle	31	6	25
925	Mechanical suffocation in other and unspecified circumstances	15	6	9
927	Caused by bites and stings of venomous animals and insects	1	1	0
928	Other accidents caused by animals	3	2	1
929	Drowning and submersion	109	58	51
931	Excessive heat and insolation	6	1	5
932	Excessive cold	5	1	4
933	Hunger, thirst, and exposure	9	2	7
934	Cataclysm	1	0	1
935	Lightning	10	2	8
936	Other and unspecified accidents	23	9	14
946	Other complications due to non-therapeutic medical and surgical procedures	1	1	0
950	Therapeutic misadventure in surgical treatment	6	3	3
951	Therapeutic misadventure in infusion or transfusion	5	3	2
953	Therapeutic misadventure in administration of drugs or biologicals	2	1	1
954	Therapeutic misadventure in anesthesia	1	0	1
955	Other and unspecified therapeutic misadventure	1	0	1
956	Late complication of surgical operation	1	0	1
960	Late effect of motor vehicle accident	4	3	1
962	Late effect of other accidental injury	5	3	2

Table 5 - Page 11
 Deaths - Mississippi - 1959

International List Number	Cause of Death*	Total	White	Non-white
<u>963,970-979</u>	<u>Suicide</u>	<u>130</u>	<u>110</u>	<u>20</u>
970	By analgesic and soporific substances	1	1	0
971	By other solid and liquid substances	3	2	1
973	By other gases	1	1	0
974	By hanging and strangulation	12	9	3
975	By submersion (drowning)	3	2	1
976	By firearms and explosives	101	88	13
977	By cutting and piercing instruments	7	6	1
979	By other and unspecified means	2	1	1
<u>964,980-985</u>	<u>Homicide</u>	<u>231</u>	<u>44</u>	<u>187</u>
981	Assault by firearm and explosive	152	32	120
982	Assault by cutting and piercing instruments	48	4	44
983	Assault by other means	22	7	15
984	Injury by intervention of police	8	1	7
985	Execution	1	0	1

* When cause of death and International List Number are omitted, no deaths were reported.

TABLE 6
DEATHS
MISSISSIPPI
1959

By Place of Occurrence, Reallocated to Place of Residence,
by Race, with Rates

County	Place of Occurrence			Place of Residence			Rate*		
	Number		Non-white	Number		Non-white	Rate*		Non-white
	Total	White		Total	White		Total	White	
STATE	20,713	10,633	10,080	20,8800	10,742	10,148	9.4	8.0	11.4
Adams	390	156	234	340	135	205	11.0	7.5	15.8
Alcorn	266	217	49	271	222	49	10.4	11.1	8.2
Amite	96	31	65	158	66	92	9.9	8.2	11.5
Attala	215	111	104	235	125	110	11.8	11.4	12.2
Benton	246	27	19	67	38	29	6.7	7.6	5.8
Bolivar	643	112	531	645	156	489	15.7	9.2	20.4
Calhoun	118	81	37	137	97	40	9.1	8.4	11.4
Carroll	55	18	37	94	44	50	8.5	7.3	10.0
Chickasaw	173	100	73	177	102	75	11.1	12.8	9.4
Choctaw	80	53	27	97	67	30	9.7	9.6	10.0
Claiborne	84	20	64	99	25	74	9.0	12.5	8.2
Clarke	151	92	59	184	111	73	11.5	11.1	12.2
Clay	181	77	104	176	79	97	11.0	9.9	12.1
Coahoma	484	127	357	511	118	393	11.9	6.9	15.1
Copiah	247	105	142	306	147	159	10.9	10.5	11.4
Covington	105	69	36	135	86	49	8.4	8.6	8.2
DeSoto	156	45	111	209	70	139	9.5	7.8	10.7
Forrest	483	344	139	475	307	168	9.1	8.1	12.0
Franklin	80	51	29	106	71	35	11.8	14.2	8.8
George	77	65	12	97	83	14	6.5	6.4	7.0
Greene	66	48	18	83	57	26	10.4	9.5	13.0
Grenada	183	76	107	165	62	103	9.2	6.9	11.4
Hancock	107	90	17	139	112	27	10.7	11.2	9.0
Harrison	1,079	851	228	908	698	210	5.5	5.1	7.0
Hinds	2,102	1,125	977	1,498	768	730	8.5	7.4	10.0
Holmes	299	72	227	333	94	239	14.5	15.7	14.0
Humphreys	173	37	136	200	47	153	12.5	7.8	15.3
Issaquena	33	4	29	51	9	42	12.8	4.5	21.0
Itawamba	81	76	5	119	113	6	5.7	5.8	4.0
Jackson	301	211	90	334	244	90	5.0	4.7	6.0
Jasper	97	42	55	160	76	84	8.9	7.6	10.5
Jefferson	84	14	70	128	30	98	10.7	10.0	10.9
Jeff Davis	111	61	50	108	56	52	5.7	6.2	5.2
Jones	665	399	266	507	340	167	7.0	6.2	9.8
Kemper	99	47	52	124	61	63	10.3	12.2	9.0
Lafayette	163	93	70	186	110	76	7.12	6.1	9.5

Table 6 - Page 2
Deaths - Mississippi - 1959

County	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Lamar	77	64	13	118	95	23	6.6	6.8	5.8
Lauderdale	883	525	358	705	410	295	11.8	11.1	12.8
Lawrence	73	35	38	123	65	58	10.2	8.1	14.5
Leake	125	77	48	171	105	66	8.6	9.5	7.3
Lee	406	280	126	402	276	126	9.6	8.9	11.4
Leflore	563	204	359	542	173	369	13.6	10.8	15.4
Lincoln	237	138	99	252	144	108	9.7	8.5	12.0
Lowndes	4051	202	203	439	212	227	10.7	8.2	15.1
Madison	322	97	225	333	100	233	12.3	11.1	12.9
Marion	180	117	63	210	133	77	7.5	7.8	7.0
Marshall	231	72	159	216	72	144	9.0	10.3	8.5
Monroe	322	193	129	353	212	141	10.1	9.6	10.8
Montgomery	141	77	64	131	73	58	10.1	9.1	11.6
Neshoba	155	108	47	205	142	63	7.9	7.9	7.9
Newton	163	106	57	192	112	80	9.1	8.0	11.4
Noxubee	178	52	126	184	59	125	13.1	11.8	13.9
Oktibbeha	195	100	95	192	91	101	6.8	5.7	8.4
Panola	265	110	155	300	125	175	10.3	8.9	11.7
Pearl River	182	136	46	202	152	50	7.0	7.6	5.6
Perry	44	26	18	85	58	27	9.4	8.3	13.5
Pike	371	221	150	373	212	161	11.6	11.8	11.5
Pontotoc	169	141	28	178	148	30	8.1	8.2	7.5
Prentiss	199	185	14	175	161	14	10.3	12.4	3.5
Quitman	177	44	133	215	65	150	10.8	9.3	11.5
Rankin	378	209	169	234	143	91	5.4	5.1	6.1
Scott	174	93	81	200	108	92	9.1	7.7	11.5
Sharkey	99	28	71	119	31	88	13.2	7.8	17.6
Simpson	218	136	82	194	127	67	7.2	7.5	6.7
Smith	75	57	18	128	100	28	7.5	8.3	5.6
Stone	62	39	23	58	34	24	7.2	5.2	16.0
Sunflower	356	83	273	402	99	303	9.1	5.8	11.2
Tallahatchie	237	77	160	284	103	181	12.9	10.3	15.1
Tate	110	40	70	163	75	88	9.0	8.3	9.8
Tippah	117	96	21	144	119	25	6.8	7.9	4.2
Tishomingo	87	79	8	128	116	12	8.5	8.0	24.0
Tunica	199	25	174	206	21	185	12.9	7.0	14.2
Union	191	135	56	198	148	50	8.2	8.2	8.3
Walhall	132	74	58	130	73	57	9.3	9.1	9.5
Warren	677	250	427	495	218	277	13.8	12.1	15.4
Washington	759	238	521	819	226	593	13.4	7.5	19.1
Wayne	124	71	53	164	91	73	9.6	8.3	12.2
Webster	92	79	13	121	100	21	11.0	12.5	7.0

Table 6 - Page 3
Deaths - Mississippi -1959

County	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Wilkinson	176	82	94	156	59	97	14.2	19.7	12.1
Winston	135	85	50	164	97	67	10.0	8.8	7.4
Yalobusha	141	85	56	153	93	60	11.8	13.3	10.0
Yazoo	338	115	223	372	140	232	12.8	12.7	12.9
Out-of-State	836	637	199	659	528	131	-	-	-

Cities over 10,000 Population**	Place of Occurrence			Place of Residence		
	Number			Number		
	Total	White	Non-white	Total	White	Non-white
Biloxi	300	243	57	297	236	61
Clarksdale	277	94	183	220	73	147
Columbus	301	166	135	254	119	135
Greenville	442	176	266	406	137	269
Greenwood	344	146	198	259	109	150
Gulfport	448	331	117	296	202	94
Hattiesburg	423	302	121	339	205	134
Jackson	1,805	980	825	1,034	551	483
Laurel	534	290	244	287	159	128
McComb	190	132	58	140	97	43
Meridian	768	462	306	505	297	208
Natchez	233	122	111	242	101	141
Pascagoula	45	27	18	112	77	35
Tupelo	223	150	73	145	84	61
Vicksburg	580	212	368	361	169	192
Total Urban#	10,247	5,726	4,521	7,337	3,929	3,408
Total Rural	10,466	4,907	5,559	13,553	6,813	6,740

* Rate per 1,000 population.

** Cities having 10,000 population and over according to 1950 census.

Includes all towns of 2,500 population or more according to 1950 census.

TABLE 7
DEATHS FROM SELECTED CAUSES
MISSISSIPPI
1959

By Age, Race, and Sex, on Basis of Place of Residence, with Rates

Inter-national List Number	Cause of Death	Total	Rate*	Race	Total by Sex and Race	Total by Sex and Race	Age in Years											
							1-4	5-9	10-14	15-24	25-34	35-44	45-54					
001-999	<u>All causes</u>	<u>20,890</u>	<u>9.4</u>	W	<u>10,742</u>	M	<u>6,283</u>	<u>363</u>	<u>55</u>	<u>32</u>	<u>43</u>	<u>180</u>	<u>144</u>	<u>673</u>	<u>1,037</u>	<u>3,445</u>	<u>1</u>	
							F	<u>4,459</u>	<u>314</u>	<u>57</u>	<u>29</u>	<u>48</u>	<u>128</u>	<u>301</u>	<u>486</u>	<u>3,017</u>	<u>4</u>	
							M	<u>5,491</u>	<u>929</u>	<u>139</u>	<u>49</u>	<u>54</u>	<u>160</u>	<u>263</u>	<u>476</u>	<u>772</u>	<u>2,461</u>	<u>4</u>
							F	<u>4,657</u>	<u>749</u>	<u>140</u>	<u>58</u>	<u>54</u>	<u>184</u>	<u>254</u>	<u>441</u>	<u>708</u>	<u>2,042</u>	<u>11</u>
001-138	Infective and parasitic diseases	373	16.7	C	247	M	84	5	5	1	0	3	2	9	16	32	0	
							F	42	3	2	1	2	5	7	3	16	0	
							M	126	129	25	6	10	12	9	26	23	0	
							F	118	118	31	11	3	5	7	17	24	0	
001-119	Tuberculosis (all forms)	153	6.9	C	91	M	62	46	0	0	0	0	0	2	5	8	25	0
							F	48	16	0	0	0	0	0	1	1	9	0
001-008	Tuberculosis (respiratory system)	142	6.4	C	82	M	60	45	0	0	0	0	0	2	5	8	12	0
							F	36	15	0	1	0	0	0	1	2	6	0
020-029	Syphilis & sequelae	43	1.9	C	34	M	9	4	0	0	0	0	0	0	0	1	2	1
							F	11	23	0	0	0	0	0	0	0	7	0
														1	1	3	6	

Table 7 - Page 2
Deaths from Selected Causes - Mississippi - 1959

Inter-national List Number	Cause of Death	Total	Rate*	Race	Total by Race	Sex and Race	Total by Sex and Race	Age in Years							Unknown Over 65	
								1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	
045-048	Dysentery (all forms)	26	1.2	C	22	W 4	M 4	1	0	0	0	0	0	0	0	0
						M 0	F 0	0	1	0	0	0	0	0	0	0
050,051	Scarlet fever and streptococcal sore throat	2	0.1	C	2	W 0	M 0	0	0	0	0	0	0	0	0	0
						M 0	F 0	0	1	0	0	0	0	0	0	0
055	Diphtheria	4	0.2	C	4	W 1	M 1	0	0	0	0	0	0	0	0	0
						M 6	F 2	0	0	0	0	0	0	0	0	0
056	Whooping cough	18	0.8	C	17	W 11	M 10	1	1	0	0	0	0	0	0	0
						M 3	F 1	0	1	0	0	0	0	0	0	0
057	Meningococcal infections	8	0.4	C	4	W 4	M 4	2	2	0	1	0	0	0	0	0
						M 2	F 2	0	1	0	0	0	0	0	0	0
080	Acute poliomyelitis	9	0.4	C	5	W 2	M 2	1	0	0	0	0	0	0	0	0
						M 2	F 2	0	1	0	0	0	0	0	0	0
085	Measles	8	0.4	C	6	W 4	M 4	1	3	0	0	0	0	0	0	0

Table 7 - Page 3
Deaths from Selected Causes - Mississippi - 1959

Inter-national List Number	Cause of Death	Total	Rate*	Race	Total by Race	X Sex	Total by Sex and Race	Age in Years	Unknown	
									Over 65	Under 4
<u>140-205</u>	Malignant neoplasms, including neoplasms of lymphatic and hematopoietic tissues	<u>2,600</u>	<u>116.7</u>	W	<u>1,580</u>	M	<u>849</u>	0-14	<u>4.87</u>	<u>0.1</u>
				M	<u>731</u>	F	<u>516</u>	15-24	<u>4.16</u>	<u>0.1</u>
				F	<u>504</u>			25-34	<u>2.85</u>	<u>0.1</u>
								35-44	<u>2.28</u>	<u>0.1</u>
								45-54		
								55-64		
								65-74		
<u>210-239</u>	Benign and unspeci-fied neoplasms	<u>48</u>	<u>2.2</u>	C	<u>22</u>	M	<u>19</u>	0-14	<u>1.71</u>	<u>0.1</u>
				W	<u>26</u>	F	<u>7</u>	15-24	<u>1.54</u>	<u>0.1</u>
				M	<u>13</u>			25-34	<u>1.31</u>	<u>0.1</u>
				F	<u>13</u>			35-44	<u>1.23</u>	<u>0.1</u>
<u>240-289</u>	Allergic, endocrine system, metabolic & nutritional diseases	<u>406</u>	<u>18.2</u>	C	<u>192</u>	M	<u>87</u>	0-14	<u>4.9</u>	<u>0.1</u>
				W	<u>214</u>	F	<u>105</u>	15-24	<u>7.4</u>	<u>0.1</u>
				M	<u>94</u>			25-34	<u>4.3</u>	<u>0.1</u>
				F	<u>120</u>			35-44	<u>5.2</u>	<u>0.1</u>
<u>260</u>	Diabetes mellitus	<u>281</u>	<u>12.6</u>	C	<u>153</u>	M	<u>53</u>	0-14	<u>2.1</u>	<u>0.1</u>
				W	<u>128</u>	F	<u>75</u>	15-24	<u>1.9</u>	<u>0.1</u>
				M	<u>88</u>			25-34	<u>1.3</u>	<u>0.1</u>
				F	<u>10</u>			35-44	<u>1.1</u>	<u>0.1</u>
<u>290-299</u>	Diseases of the blood and blood-forming organs	<u>58</u>	<u>2.6</u>	C	<u>29</u>	M	<u>13</u>	0-14	<u>9</u>	<u>0.1</u>
				W	<u>29</u>	F	<u>16</u>	15-24	<u>12</u>	<u>0.1</u>
				M	<u>17</u>			25-34	<u>4</u>	<u>0.1</u>
				F	<u>12</u>			35-44	<u>3</u>	<u>0.1</u>
<u>290-293</u>	Anemias	<u>48</u>	<u>2.2</u>	C	<u>27</u>	M	<u>10</u>	0-14	<u>1.1</u>	<u>0.1</u>
				W	<u>21</u>	F	<u>11</u>	15-24	<u>1.0</u>	<u>0.1</u>
				M	<u>17</u>			25-34	<u>1.0</u>	<u>0.1</u>
				F	<u>10</u>			35-44	<u>0.8</u>	<u>0.1</u>
<u>300-326</u>	Mental, psycho-neurotic & personality disorders	<u>32</u>	<u>1.4</u>	C	<u>9</u>	M	<u>13</u>	0-14	<u>4</u>	<u>0.1</u>
				W	<u>23</u>	F	<u>8</u>	15-24	<u>2</u>	<u>0.1</u>
				M	<u>8</u>			25-34	<u>1.0</u>	<u>0.1</u>

Table 7 - Page 4
Deaths from Selected Causes - Mississippi - 1959

International List Number	Cause of Death	Total	Rate*	Race	Total by Sex and Race	Total by Sex and Race	Age in Years	Unknown			
								Under 1-4	5-9	10-14	15-24
<u>330-334</u>	<u>Vascular lesions affecting central nervous system</u>	<u>2,683</u>	<u>120.5</u>	W	<u>1,386</u>	M	<u>665</u>	<u>1</u>	<u>0</u>	<u>0</u>	<u>86</u>
				C	<u>1,297</u>	F	<u>607</u>	<u>0</u>	<u>1</u>	<u>2</u>	<u>59</u>
								<u>1</u>	<u>1</u>	<u>0</u>	<u>114</u>
								<u>0</u>	<u>1</u>	<u>0</u>	<u>121</u>
<u>340-398</u>	<u>Other diseases of the nervous system and sense organs</u>	<u>171</u>	<u>7.7</u>	C	<u>92</u>	M	<u>51</u>	<u>5</u>	<u>3</u>	<u>4</u>	<u>33</u>
				W	<u>16</u>	F	<u>41</u>	<u>7</u>	<u>3</u>	<u>1</u>	<u>19</u>
								<u>1</u>	<u>1</u>	<u>0</u>	<u>12</u>
								<u>0</u>	<u>0</u>	<u>1</u>	<u>1</u>
<u>340</u>	<u>Nonmeningococcal meningitis</u>	<u>46</u>	<u>2.1</u>	C	<u>30</u>	M	<u>14</u>	<u>8</u>	<u>2</u>	<u>0</u>	<u>0</u>
				W	<u>16</u>	F	<u>16</u>	<u>6</u>	<u>4</u>	<u>1</u>	<u>1</u>
								<u>0</u>	<u>0</u>	<u>1</u>	<u>0</u>
<u>400-468</u>	<u>Diseases of the circulatory system</u>	<u>6,998</u>	<u>314.2</u>	C	<u>2,807</u>	M	<u>2,552</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>284</u>
				W	<u>4,191</u>	F	<u>1,639</u>	<u>1</u>	<u>0</u>	<u>0</u>	<u>19</u>
								<u>1</u>	<u>0</u>	<u>1</u>	<u>19</u>
								<u>0</u>	<u>0</u>	<u>0</u>	<u>21</u>
<u>400-402</u>	<u>Rheumatic fever</u>	<u>14</u>	<u>0.6</u>	C	<u>12</u>	M	<u>7</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
				W	<u>2</u>	F	<u>5</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
								<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>410-416</u>	<u>Chronic rheumatic heart disease</u>	<u>127</u>	<u>5.7</u>	C	<u>64</u>	M	<u>40</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
				W	<u>63</u>	F	<u>33</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
								<u>0</u>	<u>1</u>	<u>1</u>	<u>1</u>
<u>420-422</u>	<u>Arteriosclerotic and degenerative heart disease</u>	<u>4,365</u>	<u>196.0</u>	C	<u>1,336</u>	M	<u>1,963</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>240</u>
				W	<u>3,029</u>	F	<u>1,066</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>404</u>
								<u>1</u>	<u>1</u>	<u>1</u>	<u>1</u>
								<u>2</u>	<u>4</u>	<u>4</u>	<u>474</u>
								<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>

Table 7 - Page 5 Deaths from Selected Causes - Mississippi - 1959

Inter-national List Number	Cause of Death	Total	Rate*	Race	Total by Race	Total by Sex and Race	Total by Sex and Race	Total by Sex and Race	Age in Years	Unknown Over 65										
										Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 and over	
430-434	Other diseases of the heart	442	19.8	C	243	M 129 F 114	W 199 M 129	M 117	0	0	0	1	0	3	2	6	12	25	69	0
440-443	Hypertension with heart disease	1,285	57.7	C	804	M 41 F 434	M 481 F 370	M 235	0	0	0	0	0	0	0	0	2	2	35	0
444-447	Hypertension without mention of heart	255	11.4	C	159	M 81 F 78	M 96	M 55	0	0	0	0	0	0	0	0	1	5	47	0
470-527	Diseases of the respiratory system	1,004	45.1	C	502	M 297 F 205	M 502	M 290 F 212	13 27 113 80	3 7 32 25	3 0 0 2	2 1 1 1	1 1 1 1	0 0 0 2	9 8 10 13	17 16 16 16	45 32 32 32	0 0 0 0		
480-483	Influenza	82	3.7	C	57	M 17 F 8	M 25 F 28	M 29	0 0 11 8	0 0 0 3	0 0 0 0	0 1 1 1	0 1 1 1	0 0 0 0	1 3 3 0	13 11 11 13	117 45 45 13	0 0 0 0		
490-493	Pneumonia (excluding pneumonia of newborn)	686	30.8	C	383	M 218 F 165	M 303 F 218	M 155	11 21 93 63	2 4 3 21	2 1 1 4	1 1 1 1	1 5 5 5	5 5 5 5	6 6 6 6	13 13 13 13	45 45 45 45	1 1 1 1		
500-502	Bronchitis	44	2.0	C	19	M 15 F 10	M 25 F 11	M 15	0 2 3	0 2 3	0 1 1	0 0 0	0 0 0	0 0 0	0 0 0	3 1 1	12 3 2	0 0 0		

Table 7 - Page 6
Deaths from Selected Causes - Mississippi - 1959

Table 7 - Page
Deaths from Selected Causes - Mississippi , 1959

Inter-national List Number	Cause of Death	Total	Rate*	Race	Total by Race	Sex and Race	Total by Sex and Race	Age in Years		Unknown Over 65 & Under 14
								Under 14	15-24	
590-594	Nephritis and nephrosis	290	13.0	W	101	M	49	1	0	0
610	Hyperplasia of prostate	74	3.3	C	44	F	52	0	0	0
640-689*	Complications of pregnancy, childbirth, and the puerperium	55	0.9	W	9	M	101	1	0	0
690-716	Diseases of the skin, & cellular tissue	24	1.1	C	46	F	88	5	4	0
720-749	Diseases of the bones & organs of movement	39	1.8	W	13	M	30	3	2	0
750-759	Congenital malformations	247	11.1	C	24	F	44	11	10	0
760-776	Certain diseases of early infancy	1,213	20.0	W	107	M	140	29	24	3
								784	739	339

Table 7 - Page 8
Deaths from Selected Causes - Mississippi, 1959

Inter-national List Number	Cause of Death	Rate*	Race†	Total by Race	Total by Sex and Race	Total by Age and Race	Age in Years						
							10-14	15-24	25-34	35-44	45-54	55-64	Over 65
780-795	Symptoms, senility, and ill-defined conditions	1,837	82.5	W	432	M	295	15	2	2	19	44	66
						F	137	1	0	0	15	14	144
						M	775	20	5	8	28	55	84
						F	630	16	4	0	20	47	297
													0
800-999	Accidents, poison- ings and violence	1,862	83.6	C	871	M	718	12	19	11	35	98	115
						F	273	12	22	13	25	20	75
						M	632	37	45	33	22	17	20
						F	239	30	55	29	124	59	48
													83
810-835	Motor vehicle accidents	588	26.4	C	187	M	401	1	6	3	16	92	45
						F	99	2	13	6	5	16	10
						M	152	0	9	6	5	29	20
						F	35	2	6	5	1	1	4
													4
800-802, 840-962	All other accidents	913	41.0	C	477	M	293	11	13	7	18	37	43
						F	143	10	9	6	2	5	7
						M	299	37	36	18	23	49	22
						F	178	28	49	24	7	11	8
													7
963, 970-979	Suicide	130	5.8	C	20	M	110	0	0	1	0	5	23
						F	25	0	0	0	0	1	2
						M	16	0	0	0	0	1	1
						F	4	0	0	0	0	0	0
													0
964, 980-985	Homicide	231	10.4	C	187	M	44	0	0	0	1	1	39
						F	6	0	0	1	0	0	12
						M	165	0	0	1	5	43	36
						F	22	0	0	0	0	5	6
													4

* Crude death rate per 1,000 population. Specific death rates per 100,000 population. Maternal death rates and rates for infant causes (760-776) per 1,000 live births.

TABLE 8
DEATHS UNDER 1 YEAR
MISSISSIPPI
1959

For Selected Causes, by Race, by Age, on Basis of Place of Residence, with Rates

Table 2 - Page 2
Deaths Under 1 Year - Mississippi - 1959

Cause of Death and International List Number	Total	Rate*	Race	Total by Race	DAYS						MONTHS							
					Under 1 day	1- 6	7- 13	14- 20	21- 27	Under 28 days	1	2	3	4	5	6	7	
Postnatal asphyxia & atelectasis (762)	218	3.6	W C	117 101	68	45	1	2	0	116	0	0	0	0	0	0	0	0
**Infections of the newborn (763-768)	142	2.3	W C	24 118	4	10	6	0	4	24	0	0	0	0	0	0	0	0
Hemolytic disease of the newborn (770)	19	0.3	W C	12 7	9	3	0	0	0	12	0	0	0	0	0	0	0	0
All other defined dis. of early infancy (769,771,772)	62	1.0	W C	10 52	5	2	0	0	0	7	0	0	0	0	0	0	0	0
Violent & accidental deaths (800-999)	91	1.5	W C	24 67	1	0	1	0	1	4	5	2	2	3	0	2	1	0
Other specified causes	138	2.3	W C	48 90	4	9	0	1	1	15	8	7	4	1	3	1	2	1
III-defined (773-776, 780-793,795)	984	16.2	W C	228 756	118	80	2	1	2	203	7	3	4	3	4	1	1	0
					270	187	34	20	17	528	54	45	25	24	16	13	10	14
																		4

* Rate per 1,000 live births.

** Pneumonia and diarrhea of the newborn (under 28 days) included in infections of the newborn (763-768).

TABLE 9
DEATHS FROM SELECTED CAUSES
MISSISSIPPI
1959

By Month of Occurrence on Basis of Place of Residence

Cause of Death and International List Number	Total	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
TOTAL ALL CAUSES	<u>20,890</u>	<u>2,010</u>	<u>1,738</u>	<u>1,882</u>	<u>1,651</u>	<u>1,663</u>	<u>1,634</u>	<u>1,601</u>	<u>1,594</u>	<u>1,560</u>	<u>1,695</u>	<u>1,906</u>	<u>1,956</u>
Dysentery (045-048)	26	0	1	1	2	2	1	2	1	2	7	1	4
Whooping cough (056)	18	0	1	1	0	1	5	1	0	0	0	2	2
Acute Poliomyelitis (080)	9	1	1	0	0	0	0	0	0	3	0	2	5
Measles (085)	8	1	0	2	0	3	0	0	0	1	0	1	0
Influenza (480-483)	82	16	9	15	7	3	4	2	2	2	4	11	7
Lobar pneumonia (490)	151	14	18	16	11	12	9	8	9	12	11	12	19
Bronchopneumonia (491)	256	34	37	25	19	13	10	14	16	14	20	30	24
Other & unspecified pneumonia (492-493)	279	35	38	22	28	13	21	16	23	13	18	24	28
Diarrhea & enteritis (-2) (5710)	138	13	7	1	5	5	17	11	18	15	14	14	18
Diarrhea & enteritis (+2) (5711)	37	1	3	4	1	5	2	8	0	2	7	2	2
Accidents (800-962)	1,501	159	117	155	91	84	112	135	122	99	127	129	171
Motor vehicle accidents (810-835)	588	53	49	56	34	34	37	51	55	48	59	43	69

TABLE 10
MARRIAGES AND DIVORCES
MISSISSIPPI
1959

By County, by Race of Bride

County	Marriages						Divorces		
	Place of Licensure			Residence of Bride			Total	White	Non-white
	Total	White	Non-white	Total	White	Non-white			
STATE	20,447	13,547	6,900	16,288	10,267	6,021	5,108	3,557	1,551
Adams	340	206	134	268	152	116	142	98	44
Alcorn	649	583	66	192	168	24	82	73	9
Amite	117	72	45	83	46	37	32	18	14
Attala	185	112	73	180	110	70	50	22	28
Benton	64	31	33	54	28	26	7	4	3
Bolivar	408	145	263	433	154	279	87	46	41
Calhoun	122	104	18	108	90	18	23	18	5
Carroll	64	38	26	63	36	27	17	11	6
Chickasaw	148	94	54	129	88	41	25	16	9
Choctaw	69	43	26	55	31	24	6	5	1
Claiborne	58	21	37	54	19	35	32	8	24
Clarke	167	113	54	136	83	53	31	23	8
Clay	121	65	56	117	62	55	26	14	12
Coahoma	342	133	209	271	102	169	70	46	24
Copiah	205	106	99	226	114	112	68	34	34
Covington	86	62	24	102	81	21	29	24	5
DeSoto	1,259	756	503	124	53	71	40	21	19
Forrest	422	343	79	436	349	87	224	169	55
Franklin	71	48	23	61	37	24	28	15	13
George	247	227	20	108	99	9	28	26	2
Greene	101	90	11	64	54	10	12	7	5
Grenada	160	73	87	161	72	89	50	29	21
Hancock	280	260	20	73	67	6	59	53	6
Harrison	1,262	1,116	146	939	805	134	561	482	79
Hinds	1,517	913	604	1,533	952	581	512	335	177
Holmes	181	55	126	191	53	138	32	16	16
Humphreys	129	47	82	129	46	83	24	13	11
Issaquena	26	14	12	19	6	13	1	0	1
Itawamba	146	143	3	81	81	0	31	29	2
Jackson	778	595	183	524	406	118	168	138	30
Jasper	101	55	46	133	71	62	15	7	8
Jefferson	75	32	43	55	16	39	20	10	10
Jeff Davis	105	71	34	84	52	32	17	7	10
Jones	573	451	122	526	423	103	150	115	35
Kemper	78	42	36	77	43	34	11	8	3
Lafayette	124	76	48	126	76	50	36	20	16

Table 10 - Page 2

Marriages and Divorces - Mississippi - 1959

County	Marriages						Divorces		
	Place of Licensure			Residence of Bride			Total	White	Non-white
	Total	White	Non-white	Total	White	Non-white			
Lamar	152	134	18	130	114	16	11	10	1
Lauderdale	556	392	164	544	390	154	177	139	38
Lawrence	63	40	23	86	56	30	31	21	10
Leake	183	120	63	167	98	69	28	20	8
Lee	299	239	60	352	279	73	95	86	9
Leflore	337	146	191	336	149	187	88	47	41
Lincoln	205	152	53	182	134	48	51	39	12
Lowndes	346	240	106	289	198	91	83	63	20
Madison	243	83	160	210	56	154	86	17	69
Marion	211	154	57	191	144	47	65	53	12
Marshall	275	146	129	136	49	87	32	17	15
Monroe	228	179	49	225	170	55	76	55	21
Montgomery	88	58	30	97	68	29	23	15	8
Neshoba	185	136	49	181	136	45	67	53	14
Newton	141	99	42	150	102	48	34	23	11
Noxubee	102	40	62	102	42	60	26	12	14
Oktibbeha	151	96	55	132	79	53	31	12	19
Panola	213	103	110	217	101	116	46	33	13
Pearl River	317	272	45	204	167	37	89	76	13
Perry	83	69	14	80	67	13	16	15	1
Pike	284	188	96	261	159	102	124	82	42
Pontotoc	151	125	26	128	104	24	41	29	12
Prentiss	234	219	15	163	151	12	39	35	4
Quitman	192	83	109	151	58	93	29	19	10
Rankin	264	183	81	178	123	55	36	27	9
Scott	143	90	53	166	108	58	62	44	18
Sharkey	65	26	39	57	24	33	15	7	8
Simpson	167	110	57	172	106	66	50	40	10
Smith	96	81	15	107	82	25	25	21	4
Stone	120	99	21	84	69	15	16	14	2
Sunflower	296	115	181	307	116	191	51	28	23
Tallahatchie	227	87	140	175	73	102	29	20	9
Tate	175	99	76	134	63	71	23	10	13
Tippah	144	125	19	128	114	14	15	12	3
Tishomingo	191	178	13	109	105	4	25	25	0
Tunica	170	47	123	153	36	117	22	8	14
Union	148	111	37	131	93	38	34	27	7
Walthall	150	95	55	84	49	35	26	21	5
Warren	388	224	164	346	193	153	143	79	64
Washington	877	534	343	599	303	296	205	125	80
Wayne	112	77	35	92	64	28	40	25	15
Webster	59	52	7	73	63	10	14	12	2
Wilkinson	167	102	65	83	25	58	25	13	12
Winston	168	104	64	146	86	60	40	20	20
Yalobusha	75	47	28	101	59	42	14	10	4
Yazoo	226	113	113	234	117	117	64	38	26
Unstated				97	36	61			
Out-of-state residents				4,062	3,244	818			

TABLE 11
DIVORCES
MISSISSIPPI
1959

By Cause of Divorce

Cause	Total	White	Non-white
TOTAL	5,108	3,557	1,551
Cruel and inhuman treatment	2,513	2,089	424
Desertion	1,980	1,007	973
Adultery	243	154	89
Drunkenness	236	207	29
Penitentiary	34	27	7
Insanity	10	7	3
Annulment	25	22	3
Bigamy	18	6	12
Miscellaneous and other	18	15	3
Not stated	31	23	8

By Number of Minor Children

Minor Children	Total	White	Non-white
None	2,524	1,655	869
One	1,118	852	266
Two	712	535	177
Three or more	635	457	178
Not stated	119	58	61

By Number of Years Married

Number of Years Married	Total	White	Non-white
TOTAL	5,108	3,557	1,551
Under one year	298	270	28
One year	543	467	76
Two years	516	430	86
Three years	427	331	96
Four years	323	234	89
5-9 years	1,041	744	297
10-19 years	1,047	682	365
Twenty years and over	828	348	480
Not stated	85	51	34

Table 12--Live Births, by Year, by Race, on Basis of Place of Occurrence
with Rates: Mississippi, 1913-1959

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1912**	6,311					
1913	41,103	21,874	19,229	26.9	26.6	18.2
1914	40,623	21,360	19,263	21.4	25.7	18.0
1915	39,484	20,999	18,485	20.5	23.8	17.1
1916	39,033	20,130	18,903	20.0	23.6	17.2
1917	37,141	19,056	18,085	18.8	22.0	16.3
1918	39,391	20,005	19,386	19.7	22.8	17.2
1919	42,491	22,028	20,463	21.0	24.9	18.0
1920	43,743	22,438	21,305	24.4	26.3	22.7
1921	46,482	23,703	22,779	25.6	27.3	24.1
1922	43,723	21,286	22,437	23.4	24.1	23.6
1923	43,266	21,140	22,126	23.3	23.6	23.1
1924	43,960	22,180	21,780	23.4	24.3	22.5
1925	45,563	22,265	23,298	24.0	24.1	23.9
1926	50,449	24,600	25,849	26.2	26.2	26.3
1927	49,204	24,163	25,041	25.3	25.3	25.3
1928	48,126	23,642	24,484	24.5	24.4	24.5
1929	45,570	22,105	23,465	22.9	22.5	23.3
1930	47,982	23,192	24,790	23.8	23.2	24.0
1931	45,123	21,637	23,486	22.2	21.4	22.9
1932	46,481	21,883	24,598	22.6	21.4	23.9
1933	44,184	20,407	23,777	21.3	19.7	22.9
1934	47,905	22,342	25,563	23.0	21.4	24.6
1935	48,291	21,972	26,319	23.0	20.8	25.1
1936	49,261	22,194	27,067	23.3	20.8	25.7
1937	52,170	23,285	28,885	24.4	21.6	27.3
1938	53,729	24,046	29,683	25.0	22.1	27.8
1939	52,288	23,505	28,783	24.1	21.4	26.8
1940	52,526	23,084	29,442	24.0	20.8	27.3
1941	54,561	24,167	30,394	24.7	21.6	28.0
1942	56,521	25,306	31,215	25.4	22.4	28.6
1943	59,408	27,611	31,797	28.9	26.4	31.6
1944#	56,586	25,647	30,939	28.3	25.2	31.6
1945#	53,540	23,426	30,114	26.9	23.1	30.9
1946#	61,430	29,715	31,715	26.8	25.4	28.2
1947#	66,291	32,423	33,868	28.6	27.5	29.9
1948#	65,291	29,666	35,625	28.0	24.9	31.2
1949#	66,128	28,786	37,342	28.1	24.0	32.5
1950#	64,427	27,188	37,239	29.6	22.9	37.6
1951#	65,512	27,798	37,714	30.1	23.4	38.1
1952#	64,509	28,073	36,436	29.6	23.6	36.8
1953#	64,061	28,312	35,749	29.4	23.8	36.1
1954#	64,748	28,781	35,967	29.7	24.2	36.3
1955#	63,244	27,546	35,698	29.0	23.2	36.0

Table 12 (continued)

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1956#	62,178	27,278	34,900	29.5	23.8	36.2
1957#	60,712	26,804	33,908	28.8	23.4	35.2
1958#	59,189	26,812	32,377	27.1	20.9	35.9
1959#	60,562	27,979	32,583	27.2	20.9	36.6

* Rate per 1,000 population. 1913-1920 population estimates based on U. S. Censuses of 1910 & 1920; 1921-1930 population estimates based on U. S. Censuses of 1920 & 1930; 1931-1940 population estimates based on U. S. Censuses of 1930 & 1940; 1941 & 1942 projected estimates based on U. S. Censuses of 1930 & 1940; 1943 & 1944 estimates of civilian population obtained from Ration Book Registration; 1945 estimate of civilian population, July 1, 1945, obtained from Bureau of Census Bulletin Series P-46, No. 3, dated February 12, 1946; 1946-1949 projected estimates as of July 1, each year based on U. S. Censuses of 1930 & 1940; 1950-1955 U. S. Census population of April 1, 1950; estimate for 1956-1957 as of July 1, 1956, prepared by Dr. Harald A Pedersen, Sociology Department, Mississippi State University, with adjustments by Statistical Service, Mississippi State Board of Health; 1958 estimate as of January 1, prepared by Dr. George L. Wilber and Mr. Carl T. Robbins, Sociology Department, Mississippi State University. 1959 estimate as of January 1, prepared by Dr. George L. Wilber and Ellen Bryant, Sociology Department, Mississippi State University.

** November-December, 1912.

On basis of place of residence.

Table 13--Deaths, by Year, by Race, on Basis of Place of Occurrence, with
Rates: Mississippi, 1913-1959

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1912**	4,341					
1913	21,305	7,643	13,662	11.4	9.3	13.0
1914	21,485	7,442	14,043	11.3	8.9	13.1
1915	24,001	8,088	15,913	12.5	9.6	14.7
1916	21,794	7,588	14,206	11.2	8.9	13.0
1917	23,579	8,048	15,531	11.9	9.3	14.0
1918	30,437	10,177	20,260	15.2	11.6	18.0
1919	22,667	8,142	14,525	12.7	9.5	15.5
1920	22,072	7,916	14,156	12.3	9.3	15.1
1921	19,991	7,308	12,683	11.0	8.4	13.4
1922	19,278	7,312	11,966	10.5	8.3	12.6
1923	20,363	7,688	12,675	11.0	8.6	13.2
1924	21,155	7,809	13,346	11.3	8.6	13.8
1925	22,229	8,143	14,086	11.7	8.8	14.4
1926	23,929	9,133	14,796	12.5	9.7	15.1
1927	23,257	8,765	14,492	12.0	9.2	14.6
1928	25,965	10,111	15,854	13.2	10.4	15.9
1929	25,868	10,048	15,820	13.0	10.2	15.7
1930	24,098	9,072	15,026	11.9	9.0	14.8
1931	22,007	8,641	13,366	10.8	8.5	13.0
1932	20,322	8,295	12,027	9.9	8.1	11.6
1933	21,617	8,898	12,719	10.4	8.6	12.2
1934	21,786	9,350	12,436	10.5	9.0	12.0
1935	21,285	9,186	12,099	10.1	8.7	11.6
1936	24,127	10,367	13,760	11.4	9.7	13.1
1937	24,005	10,055	13,950	11.2	9.3	13.2
1938	22,814	9,471	13,343	10.6	8.7	12.5
1939	22,701	9,464	13,237	10.5	8.6	12.3
1940	23,185	9,565	13,620	10.6	8.6	12.6
1941	22,573	9,442	13,131	10.2	8.4	12.1
1942	20,693	8,750	11,943	9.3	7.7	10.9
1943	20,850	9,139	11,711	10.2	8.7	11.6
1944#	20,199	8,830	11,369	10.1	8.7	11.6
1945#	19,024	8,452	10,572	9.6	8.3	10.8
1946#	19,321	8,995	10,326	8.4	7.7	9.2
1947#	20,007	9,238	10,769	8.6	7.8	9.5
1948#	20,183	9,216	10,967	8.6	7.7	9.6
1949#	20,489	9,233	11,256	8.7	7.7	9.8
1950#	20,567	9,466	11,101	9.4	8.0	11.2
1951#	21,007	9,593	11,414	9.6	8.1	11.5
1952#	20,657	9,387	11,270	9.5	7.9	11.4
1953#	20,492	9,495	10,997	9.4	8.0	11.1
1954#	19,353	9,453	9,900	8.9	8.0	10.0
1955#	19,403	9,501	9,902	8.9	8.0	10.0

Table 13 (continued)

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1956#	19,780	9,969	9,811	9.4	8.7	10.2
1957#	20,476	10,339	10,137	9.7	9.0	10.5
1958#	21,541	10,811	10,730	9.9	8.4	11.9
1959#	20,890	10,742	10,148	9.4	8.0	11.4

* Rate per 1,000 population. See footnote on Table 12 for source of population estimates.

** November-December, 1912.

On basis of place of residence.

Table 14--Maternal Deaths, by Year, by Race, on Basis of Place
of Occurrence, with Rates: Mississippi, 1913-1959

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1913	447	162	285	10.9	7.4	14.8
1914	413	124	289	10.2	5.8	15.0
1915	289	88	201	7.3	4.2	10.9
1916	326	102	224	8.4	5.1	11.8
1917	297	105	192	8.0	5.5	10.6
1918	302	82	220	7.7	4.1	11.3
1919	407	173	234	9.6	7.8	11.4
1920	393	127	266	9.0	5.7	12.5
1921	391	145	246	8.4	6.1	10.8
1922	313	110	203	7.2	5.2	9.0
1923	321	108	213	7.4	5.1	9.6
1924	413	137	276	9.4	6.2	12.7
1925	435	143	292	9.5	6.4	12.5
1926	388	153	235	7.7	6.2	9.1
1927	409	134	275	8.3	5.5	11.0
1928	421	161	260	8.7	6.8	10.6
1929	410	152	258	9.0	6.9	11.0
1930	489	171	318	10.2	7.4	12.8
1931	352	131	221	7.8	6.1	9.4
1932	279	103	176	6.0	4.7	7.2
1933	315	116	199	7.1	5.7	8.4
1934	306	116	190	6.4	5.2	7.4
1935	334	137	197	6.9	6.2	7.5
1936	358	134	224	7.3	6.0	8.3
1937	370	150	220	7.1	6.4	7.6
1938	333	116	217	6.2	4.8	7.3
1939	311	97	214	5.9	4.1	7.4
1940	327	106	221	6.2	4.6	7.5
1941	293	93	200	5.4	3.8	6.6
1942	248	77	171	4.4	3.0	5.5
1943	233	66	167	3.9	2.4	5.3
1944**	225	72	153	4.0	2.8	4.9
1945**	216	64	152	4.0	2.7	5.0
1946**	199	67	132	3.2	2.2	4.2
1947**	175	50	125	2.6	1.5	3.7
1948**	177	38	139	2.7	1.3	3.9
1949**	147	32	115	2.2	1.1	3.1
1950**	165	30	135	2.6	1.1	3.6
1951**	143	28	115	2.2	1.0	3.0
1952**	119	15	104	1.8	0.5	2.8
1953**	119	24	95	1.8	0.8	2.6
1954**	104	11	93	1.6	0.4	2.6
1955**	76	12	64	1.2	0.4	1.8
1956**	74	12	62	1.2	0.4	1.8
1957**	75	14	61	1.2	0.5	1.8
1958**	53	10	43	0.9	0.4	1.3
1959**	55	9	46	0.9	0.3	1.4

* Rate per 1,000 live births.

** On basis of place of residence.

Table 15--Deaths Under 1 Year, by Year, by Race, on Basis of Place of Occurrence, with Rates: Mississippi, 1917-1959

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1917	3,689	1,296	2,393	99.3	68.0	132.3
1918						
1919	3,554			83.6		
1920	3,499	1,344	2,155	80.0	59.9	101.1
1921	3,157	1,243	1,914	67.9	52.4	84.0
1922	2,826	1,067	1,759	64.6	50.1	78.4
1923	2,908	1,104	1,804	67.2	52.2	81.5
1924	3,137	1,188	1,949	71.4	53.6	89.5
1925	3,176	1,179	1,997	69.7	53.0	85.7
1926	3,578	1,447	2,131	70.9	58.8	82.4
1927	3,291	1,303	1,988	66.9	54.0	79.4
1928	3,567	1,416	2,151	74.1	59.9	87.9
1929	3,275	1,268	2,007	71.9	57.4	85.5
1930	3,276	1,182	2,094	68.3	51.0	84.5
1931	2,547	961	1,586	56.4	44.4	67.5
1932	2,482	965	1,517	53.4	44.1	61.7
1933	2,800	1,035	1,765	63.4	50.7	74.2
1934	3,070	1,199	1,871	64.1	53.7	73.2
1935	2,587	1,044	1,543	53.6	47.5	58.6
1936	2,883	1,105	1,778	58.5	49.8	65.7
1937	3,098	1,174	1,924	59.4	50.4	66.6
1938	3,051	1,205	1,846	56.8	50.1	62.2
1939	2,908	1,135	1,773	55.6	48.3	61.6
1940	2,864	1,070	1,794	54.5	46.4	60.9
1941	2,969	1,113	1,856	54.4	46.1	61.1
1942	2,654	957	1,697	47.0	37.8	54.4
1943	2,774	1,137	1,637	46.7	41.2	51.5
1944**	2,489	974	1,515	44.0	38.0	49.0
1945**	2,192	846	1,346	40.9	36.1	44.7
1946**	2,298	1,005	1,293	37.4	33.8	40.8
1947**	2,428	1,009	1,419	36.6	31.1	41.9
1948**	2,470	988	1,482	37.8	33.3	41.6
1949**	2,612	900	1,712	39.5	31.3	45.8
1950**	2,348	763	1,585	36.4	28.1	42.6
1951**	2,589	814	1,775	39.5	29.3	47.1
1952**	2,592	760	1,832	40.2	27.1	50.3
1953**	2,534	807	1,727	39.6	28.5	48.3
1954**	2,360	744	1,616	36.4	25.8	44.9
1955**	2,388	680	1,658	37.0	24.7	46.4
1956**	2,353	734	1,619	37.8	26.9	46.4
1957**	2,257	694	1,563	37.2	25.9	46.1
1958**	2,424	706	1,718	41.0	26.3	53.1
1959**	2,355	677	1,678	38.9	24.2	51.5

* Rate per 1,000 live births.

** On basis of place of residence.




TABLE 17
DEATHS FROM HEART DISEASES, NEPHRITIS (ALL FORMS),
VASCULAR LESIONS, CANCER, AND TUBERCULOSIS (ALL FORMS)
MISSISSIPPI
1950-1959

Shown by Year and Race, on Basis of Place of Residence, with Rates*

YEAR	HEART DISEASES				NEPHRITIS (ALL FORMS)				VASCULAR LESIONS				CANCER				TUBERCULOSIS (ALL FORMS)	
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
1950 No.	5,546	3,158	2,388	733	246	487	2,200	971	1,229	2,143	1,333	810	558	179	1,333	558	179	379
Rate	254.5	265.7	241.1	33.6	20.7	49.2	101.0	81.7	124.1	98.4	112.2	81.8	25.6	15.1	1,333	558	179	38.3
1951 No.	5,626	3,134	2,492	681	210	471	2,344	1,089	1,255	2,186	1,326	860	497	158	1,326	497	158	339
Rate	258.2	263.7	251.6	31.2	17.7	47.6	107.6	91.6	126.7	100.3	111.6	86.8	22.8	13.3	111.6	86.8	22.8	34.2
1952 No.	5,876	3,256	2,620	658	214	444	2,386	1,098	1,288	2,189	1,285	904	418	133	1,285	418	133	285
Rate	269.7	273.9	264.6	30.2	18.0	44.8	109.5	92.4	130.1	100.5	108.1	91.3	19.2	11.2	108.1	91.3	19.2	28.8
1953 No.	5,758	3,183	2,575	474	164	310	2,404	1,125	1,279	2,230	1,348	882	308	85	1,348	882	308	223
Rate	264.3	267.8	260.0	21.8	13.8	31.3	110.3	94.6	129.2	102.3	113.4	89.1	14.1	7.2	113.4	89.1	14.1	22.5
1954 No.	5,708	3,325	2,383	448	169	279	2,422	1,176	1,246	2,332	1,420	912	226	80	1,420	912	226	146
Rate	262.0	279.7	240.6	20.6	14.2	28.2	111.2	98.9	125.8	107.0	119.5	92.1	10.4	6.7	119.5	92.1	10.4	14.7
1955 No.	5,969	3,517	2,452	401	144	257	2,447	1,169	1,278	2,317	1,402	915	211	66	1,402	915	211	145
Rate	273.9	295.9	247.6	18.4	12.1	26.0	112.3	98.3	129.0	106.3	118.0	92.4	9.7	5.6	118.0	92.4	9.7	14.6
1956 No.	6,104	3,703	2,401	410	161	249	2,550	1,229	1,321	2,432	1,439	993	199	80	1,439	993	199	119
Rate	289.3	322.8	249.4	19.4	14.0	25.9	120.9	107.1	137.2	115.3	125.4	103.2	9.4	7.0	115.3	125.4	103.2	12.4
1957 No.	6,367	3,706	2,661	362	143	219	2,800	1,420	1,380	2,496	1,482	1,014	161	61	1,482	1,014	161	100
Rate	301.8	323.0	276.4	17.1	12.5	22.7	132.7	123.8	143.9	118.3	129.2	105.3	7.6	5.3	118.3	129.2	105.3	10.4
1958 No.	6,515	3,924	2,591	322	100	222	2,867	1,426	1,441	2,525	1,524	1,001	172	82	1,524	1,001	172	90
Rate	298.3	305.9	287.5	14.7	7.8	24.6	131.3	111.2	159.9	115.6	118.8	111.1	7.9	6.4	115.6	118.8	111.1	10.0
1959 No.	6,219	3,772	2,447	290	101	189	2,683	1,386	1,297	2,600	1,580	1,020	153	62	1,580	1,020	153	91
Rate	279.2	282.3	274.6	13.0	7.6	21.2	120.5	103.7	145.6	116.7	118.3	114.5	6.9	4.6	114.5	116.7	118.3	10.2


* Rate per 100,000 population.

TABLE 18
BIRTHS, DEATHS, MARRIAGES
MISSISSIPPI
1959

By Month of Occurrence

Month	Births	Deaths	Marriages
<u>TOTAL</u>	<u>60,562</u>	<u>20,890</u>	<u>20,447</u>
January	5,325	2,010	1,599
February	4,698	1,738	1,414
March	4,994	1,882	1,462
April	4,450	1,651	1,407
May	4,347	1,663	1,735
June	4,691	1,634	2,175
July	5,362	1,601	1,737
August	5,682	1,594	1,957
September	5,433	1,560	1,645
October	5,231	1,695	1,598
November	5,038	1,906	1,546
December	5,311	1,956	2,172

COUNTY SUMMARIES



BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ADAMS COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1663	665	998	1200	508	692	38.7	28.2	53.2
Fetal deaths	51	11	40	32	7	25	26.7	13.8	36.1
Deaths from all causes	390	156	234	340	135	205	11.0	7.5	15.8
Deaths under 1 year	56	21	35	45	16	29	37.5	31.5	41.9
Deaths under 1 month	33	14	19	28	13	15	23.3	25.6	21.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	1	2	3	1	2	9.7	5.6	15.4
Tbc., respiratory (001-008)	3	1	2	3	1	2	9.7	5.6	15.4
Syphilis (020-029)	1	0	1	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	3.2	-	7.7
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	47	26	21	44	22	22	141.9	122.2	169.2
Diabetes (260)	9	2	7	7	1	6	22.6	5.6	46.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	52	21	31	44	16	28	141.9	88.9	215.4
Heart diseases (410-443)	128	46	82	116	44	72	374.2	244.4	553.8
Arteriosclerosis (450)	6	3	3	5	2	3	16.1	11.1	23.1
Influenza (480-483)	3	0	3	3	0	3	9.7	-	23.1
#Pneumonia (490-493)	12	3	9	10	2	8	32.2	11.1	61.5
#Diarrhea & enteritis (571)	1	0	1	1	0	1	3.2	-	7.7
Acute nephritis (590, 591)	2	0	2	1	0	1	3.2	-	7.7
Chronic nephritis, etc. (592-594)	3	2	1	2	1	1	6.4	5.6	7.7
Maternal deaths (640-689)	1	0	1	1	0	1	0.8	-	1.4
Cong. malformations (750-759)	5	1	4	3	1	2	9.7	5.6	15.4
Birth inj., atelect. (760-762)	13	10	3	9	8	1	7.5	15.7	1.4
#Other dis. early inf. (763-776)	24	8	16	17	5	12	14.2	9.8	17.3
Violent & acc. deaths (800-999)	23	13	10	29	17	12	93.5	94.4	92.3
Acc. deaths, total (800-962)	21	13	8	27	17	10	87.1	94.4	76.9
Motor vehicle acc. (810-835)	5	4	1	9	7	2	29.0	38.9	15.4
Ill-defined (780-795)	10	2	8	10	2	8	32.2	11.1	61.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ALCORN COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	599	495	104	522	419	103	20.1	21.0	17.2
Fetal deaths	19	16	3	15	12	3	28.7	28.6	29.1
Deaths from all causes	266	217	49	271	222	49	10.4	11.1	8.2
Deaths under 1 year	20	9	11	16	6	10	30.6	14.3	97.1
Deaths under 1 month	7	3	4	8	4	4	15.3	9.5	38.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	2	0	3	2	1	11.5	10.0	16.7
Tbc., respiratory (001-008)	2	2	0	3	2	1	11.5	10.0	16.7
Syphilis (020-029)	1	1	0	1	1	0	3.8	5.0	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	33	31	2	31	29	2	119.2	145.0	33.3
Diabetes (260)	2	1	1	4	3	1	15.4	15.0	16.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	48	39	9	45	39	6	173.1	195.0	100.0
Heart diseases (410-443)	95	81	14	98	85	13	376.9	425.0	216.7
Arteriosclerosis (450)	1	1	0	1	1	0	3.8	5.0	-
Influenza (480-483)	1	1	0	1	1	0	3.8	5.0	-
#Pneumonia (490-493)	5	5	0	10	9	1	38.5	45.0	16.7
#Diarrhea & enteritis (571)	2	1	1	1	1	0	3.8	5.0	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	1	0	1	3.8	-	16.7
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	1	1	1	0	1	3.8	-	16.7
Birth inj., atelect. (760-762)	2	1	1	3	2	1	5.7	4.8	9.7
#Other dis. early inf. (763-776)	8	4	4	6	2	4	11.5	4.8	38.8
Violent & acc. deaths (800-999)	21	16	5	20	14	6	76.9	70.0	100.0
Acc. deaths, total (800-962)	18	14	4	15	11	4	57.7	55.0	66.7
Motor vehicle acc. (810-835)	11	9	2	9	7	2	34.6	35.0	33.3
Ill-defined (780-795)	19	12	7	22	15	7	84.6	75.0	116.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
AMITE COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	232	36	196	360	114	246	22.5	14.2	30.8
Fetal deaths	2	0	2	8	3	5	22.2	26.3	20.3
Deaths from all causes	96	31	65	158	66	92	9.9	8.2	11.5
Deaths under 1 year	9	1	8	20	4	16	55.6	35.1	65.0
Deaths under 1 month	2	0	2	9	2	7	25.0	17.5	28.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	1	0	1	6.2	-	12.5
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	1	0	1	6.2	-	12.5
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	13	6	7	21	12	9	131.2	150.0	112.5
Diabetes (260)	1	0	1	1	0	1	6.2	-	12.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	19	4	15	25	9	16	156.2	112.5	200.0
Heart diseases (410-443)	34	12	22	51	21	30	318.8	262.5	375.0
Arteriosclerosis (450)	4	3	1	4	3	1	25.0	37.5	12.5
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	5	1	4	8	4	4	50.0	50.0	50.0
#Diarrhea & enteritis (571)	1	1	0	3	1	2	18.8	12.5	25.0
Acute nephritis (590, 591)	0	0	0	1	0	1	6.2	-	12.5
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	12.5	12.5	12.5
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	2	1	1	12.5	12.5	12.5
Birth inj., atelect. (760-762)	0	0	0	3	1	2	8.3	8.8	8.1
#Other dis. early inf. (763-776)	2	0	2	6	1	5	16.7	8.8	20.3
Violent & acc. deaths (800-999)	5	1	4	12	5	7	75.0	62.5	87.5
Acc. deaths, total (800-962)	5	1	4	10	4	6	62.5	50.0	75.0
Motor vehicle acc. (810-835)	0	0	0	1	1	0	6.2	12.5	-
Ill-defined (780-795)	5	1	4	6	2	4	37.5	25.0	50.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ATTALA COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	522	222	300	499	197	302	25.0	17.9	33.6
Fetal deaths	23	7	16	17	3	14	34.1	15.2	46.4
Deaths from all causes	215	111	104	235	125	110	11.8	11.4	12.2
Deaths under 1 year	21	1	20	20	3	17	40.1	15.2	56.3
Deaths under 1 month	18	1	17	17	3	14	34.1	15.2	46.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	5.0	-	11.1
Tbc., respiratory (001-008)	1	0	1	1	0	1	5.0	-	11.1
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	14	10	4	22	14	8	110.0	127.3	88.9
Diabetes (260)	6	1	5	6	2	4	30.0	18.2	44.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	39	24	15	44	27	17	220.0	245.4	188.9
Heart diseases (410-443)	73	44	29	78	49	29	390.0	445.4	322.2
Arteriosclerosis (450)	1	1	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	4	1	3	4	1	3	20.0	9.1	33.3
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	1	1	3	1	2	15.0	9.1	22.2
Birth inj., atelect. (760-762)	5	1	4	4	1	3	8.0	5.1	9.9
#Other dis. early inf. (763-776)	12	0	12	12	2	10	24.0	10.2	33.1
Violent & acc. deaths (800-999)	24	14	10	25	14	11	125.0	127.3	122.2
Acc. deaths, total (800-962)	15	10	5	16	10	6	80.0	90.9	66.7
Motor vehicle acc. (810-835)	5	4	1	5	4	1	25.0	36.4	11.1
Ill-defined (780-795)	17	7	10	15	6	9	75.0	54.5	100.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
BENTON COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	96	3	93	192	58	134	19.2	11.6	26.8
Fetal deaths	6	0	6	15	2	13	78.1	34.5	97.0
Deaths from all causes	46	27	19	67	38	29	6.7	7.6	5.8
Deaths under 1 year	6	1	5	10	2	8	52.1	34.5	59.7
Deaths under 1 month	3	1	2	7	2	5	36.4	34.5	37.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	5	4	1	8	7	1	80.0	140.0	20.0
Diabetes (260)	1	1	0	2	2	0	20.0	40.0	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	6	4	2	7	5	2	70.0	100.0	40.0
Heart diseases (410-443)	13	9	4	19	13	6	190.0	260.0	120.0
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	0	1	2	1	1	20.0	20.0	20.0
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	1	0	1	5.2	-	7.5
Cong. malformations (750-759)	1	1	0	1	1	0	10.0	20.0	-
Birth inj., atelect. (760-762)	0	0	0	1	0	1	5.2	-	7.5
#Other dis. early inf. (763-776)	2	1	1	5	2	3	26.0	34.5	22.4
Violent & acc. deaths (800-999)	5	2	3	5	1	4	50.0	20.0	80.0
Acc. deaths, total (800-962)	4	2	2	5	1	4	50.0	20.0	80.0
Motor vehicle acc. (810-835)	3	2	1	1	0	1	10.0	-	20.0
Ill-defined (780-795)	10	3	7	11	2	9	110.0	40.0	180.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
BOLIVAR COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,826	335	1,491	1,735	355	1,380	42.3	20.9	57.5
Fetal deaths	79	11	68	76	15	61	43.8	42.2	44.2
Deaths from all causes	643	112	531	645	156	489	15.7	9.2	20.4
Deaths under 1 year	80	7	73	86	13	73	49.6	36.6	52.9
Deaths under 1 month	39	5	34	42	9	33	24.2	25.4	23.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	1	2	4	2	2	9.8	11.8	8.3
Tbc., respiratory (001-008)	3	1	2	4	2	2	9.8	11.8	8.3
Syphilis (020-029)	0	0	0	2	1	1	4.9	5.9	4.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	4	1	3	4	1	3	9.8	5.9	12.5
Scarlet fever & streptococcal sore throat (050, 051)	1	0	1	1	0	1	2.4	-	4.2
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	2	0	2	3	0	3	7.3	-	12.5
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	2	0	2	2	0	2	4.9	-	8.3
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	58	14	44	64	20	44	156.1	117.6	183.3
Diabetes (260)	15	2	13	16	4	12	39.0	23.5	50.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	93	16	77	82	18	64	200.0	105.9	266.7
Heart diseases (410-443)	180	43	137	163	51	112	397.6	300.0	466.7
Arteriosclerosis (450)	7	2	5	6	2	4	14.6	11.8	16.7
Influenza (480-483)	4	2	2	4	2	2	9.8	11.8	8.3
#Pneumonia (490-493)	18	4	14	20	7	13	48.8	41.2	54.2
#Diarrhea & enteritis (571)	13	1	12	16	2	14	39.0	11.8	58.3
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	8	0	8	8	2	6	19.5	11.8	25.0
Maternal deaths (640-689)	3	1	2	2	1	1	1.2	2.8	0.7
Cong. malformations (750-759)	6	2	4	6	3	3	14.6	17.6	12.5
Birth inj., atelect.* (760-762)	8	1	7	8	1	7	4.6	2.8	5.1
#Other dis. early inf. (763-776)	22	3	19	22	5	17	12.7	14.1	12.3
Violent & acc. deaths (800-999)	45	8	37	52	13	39	126.8	76.5	162.5
Acc. deaths, total (800-962)	39	7	32	45	11	34	109.8	64.7	141.7
Motor vehicle acc. (810-835)	12	2	10	14	4	10	34.1	23.5	41.7
III-defined (780-795)	96	5	91	98	4	94	239.0	23.5	391.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CALHOUN COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	384	194	190	350	196	154	23.3	17.0	44.0
Fetal deaths	6	2	4	6	1	5	17.1	5.1	32.5
Deaths from all causes	118	81	37	137	97	40	9.1	8.4	11.4
Deaths under 1 year	14	4	10	10	3	7	28.6	15.3	45.4
Deaths under 1 month	6	3	3	2	1	1	5.7	5.1	6.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	6.7	-	28.6
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	12	10	2	18	15	3	120.0	130.4	85.7
Diabetes (260)	2	1	1	2	1	1	13.3	8.7	28.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	14	11	3	17	14	3	113.3	121.7	85.7
Heart diseases (410-443)	33	25	8	36	28	8	240.0	243.5	228.6
Arteriosclerosis (450)	5	5	0	5	5	0	33.3	43.5	-
Influenza (480-483)	1	0	1	1	0	1	6.7	-	28.6
#Pneumonia (490-493)	13	10	3	14	10	4	93.3	87.0	114.3
#Diarrhea & enteritis (571)	3	0	3	3	0	3	20.0	-	85.7
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	6.7	8.7	-
Maternal deaths (640-689)	2	1	1	1	0	1	2.8	-	6.5
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	5	2	3	2	1	1	5.7	5.1	6.5
Violent & acc. deaths (800-999)	8	5	3	13	8	5	86.7	69.6	142.8
Acc. deaths, total (800-962)	5	4	1	10	7	3	66.7	60.9	85.7
Motor vehicle acc. (810-835)	2	2	0	5	5	0	33.3	43.5	-
Ill-defined (780-795)	11	5	6	10	4	6	66.7	34.8	171.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CARROLL COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	211	28	183	248	53	195	22.5	8.8	39.0
Fetal deaths	9	0	9	13	1	12	52.4	18.9	61.5
Deaths from all causes	55	18	37	94	44	50	8.5	7.3	10.0
Deaths under 1 year	7	0	7	9	0	9	36.3	-	46.2
Deaths under 1 month	5	0	5	7	0	7	28.2	-	35.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	7	0	7	14	6	8	127.3	100.0	160.0
Diabetes (260)	0	0	0	1	0	1	9.1	-	20.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	3	1	2	7	4	3	63.6	66.7	60.0
Heart diseases (410-443)	11	4	7	21	13	8	190.9	216.7	160.0
Arteriosclerosis (450)	1	0	1	1	0	1	9.1	-	20.0
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	0	1	1	0	1	9.1	-	20.0
#Diarrhea & enteritis (571)	0	0	0	1	0	1	9.1	-	20.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	0	1	9.1	-	20.0
Maternal deaths (640-689)	0	0	0	1	0	1	4.0	-	5.1
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	1	0	1	2	0	2	8.1	-	10.2
#Other dis. early inf. (763-776)	2	0	2	3	0	3	12.1	-	15.4
Violent & acc. deaths (800-999)	6	5	1	13	11	2	118.2	183.3	40.0
Acc. deaths, total (800-962)	4	4	0	10	9	1	90.9	150.0	20.0
Motor vehicle acc. (810-835)	1	1	0	5	5	0	45.4	83.3	-
Ill-defined (780-795)	18	4	14	18	4	14	163.6	66.7	280.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CHICKASAW COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	413	203	210	455	213	242	28.4	26.6	30.2
Fetal deaths	17	5	12	19	7	12	41.8	32.9	49.6
Deaths from all causes	173	100	73	177	102	75	11.1	12.8	9.4
Deaths under 1 year	19	5	14	22	6	16	48.4	28.2	66.1
Deaths under 1 month	13	5	8	16	6	10	35.2	28.2	41.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	1	0	6.2	12.5	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	23	17	6	22	15	7	137.5	187.5	87.5
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	30	16	14	30	17	13	187.5	212.5	162.5
Heart diseases (410-443)	54	39	15	51	38	13	318.8	475.0	162.5
Arteriosclerosis (450)	1	1	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	6	4	2	7	5	2	43.8	62.5	25.0
#Diarrhea & enteritis (571)	4	0	4	4	1	3	25.0	12.5	37.5
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	6.2	12.5	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	2	2	0	12.5	25.0	-
Birth inj., atelect. (760-762)	3	2	1	2	1	1	4.4	4.7	4.1
#Other dis. early inf. (763-776)	7	2	5	9	2	7	19.8	9.4	28.9
Violent & acc. deaths (800-999)	16	7	9	13	5	8	81.2	62.5	100.0
Acc. deaths, total (800-962)	13	4	9	13	5	8	81.2	62.5	100.0
Motor vehicle acc. (810-835)	3	2	1	2	1	1	12.5	12.5	12.5
Ill-defined (780-795)	16	2	14	15	2	13	93.8	25.0	162.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CHOCTAW COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	197	102	95	199	104	95	19.9	14.8	31.7
Fetal deaths	3	2	1	4	3	1	20.1	28.8	10.5
Deaths from all causes	80	53	27	97	67	30	9.7	9.6	10.0
Deaths under 1 year	5	2	3	3	0	3	15.1	31.6	31.6
Deaths under 1 month	4	2	2	2	0	2	10.0	-	21.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	10.0	-	33.3
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	12	10	2	14	11	3	140.0	157.1	100.0
Diabetes (260)	2	2	0	3	2	1	30.0	28.6	33.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	6	3	3	11	7	4	110.0	100.0	133.3
Heart diseases (410-443)	25	18	7	30	24	6	300.0	342.8	200.0
Arteriosclerosis (450)	5	4	1	5	4	1	50.0	57.1	33.3
Influenza (480-483)	1	0	1	1	0	1	10.0	-	33.3
#Pneumonia (490-493)	5	3	2	5	3	2	50.0	42.8	66.7
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	1	0	1	1	0	1	5.0	-	10.5
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	2	1	1	1	0	1	5.0	-	10.5
#Other dis. early inf. (763-776)	1	1	0	0	0	0	-	-	-
Violent & acc. deaths (800-999)	2	1	1	6	4	2	60.0	57.1	66.7
Acc. deaths, total (800-962)	0	0	0	5	3	2	50.0	42.8	66.7
Motor vehicle acc. (810-835)	0	0	0	1	0	1	10.0	-	33.3
Ill-defined (780-795)	7	3	4	8	4	4	80.0	57.1	133.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CLAIBORNE COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	244	29	215	310	46	264	28.2	23.0	29.3
Fetal deaths	8	1	7	8	1	7	25.8	21.7	26.5
Deaths from all causes	84	20	64	99	25	74	9.0	12.5	8.2
Deaths under 1 year	11	0	11	13	1	12	41.9	21.7	45.4
Deaths under 1 month	6	0	6	6	0	6	19.4	-	22.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	1	0	1	1	0	1	9.1	-	11.1
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	6	1	5	11	3	8	100.0	150.0	88.9
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	10	4	6	11	4	7	100.0	200.0	77.8
Heart diseases (410-443)	32	10	22	35	12	23	318.2	600.0	255.6
Arteriosclerosis (450)	1	1	0	1	1	0	9.1	50.0	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	5	0	5	5	0	5	45.4	-	55.6
#Diarrhea & enteritis (571)	1	1	0	2	1	1	18.2	50.0	11.1
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	0	2	3	0	3	27.3	-	33.3
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	9.1	50.0	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	4	0	4	4	0	4	12.9	-	15.2
Violent & acc. deaths (800-999)	8	2	6	7	0	7	63.6	-	77.8
Acc. deaths, total (800-962)	6	2	4	5	0	5	45.4	-	55.6
Motor vehicle acc. (810-835)	3	2	1	1	0	1	9.1	-	11.1
Ill-defined (780-795)	8	0	8	9	0	9	81.8	-	100.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CLARKE COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	186	111	75	392	193	199	24.5	19.3	33.2
Fetal deaths	7	6	1	18	5	13	45.9	25.9	65.3
Deaths from all causes	151	92	59	184	111	73	11.5	11.1	12.2
Deaths under 1 year	11	5	6	13	7	6	33.2	36.3	30.2
Deaths under 1 month	5	3	2	7	4	3	17.8	20.7	15.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	11	7	4	16	12	4	100.0	120.0	66.7
Diabetes (260)	2	2	0	2	2	0	12.5	20.0	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	24	11	13	29	14	15	181.2	140.0	250.0
Heart diseases (410-443)	59	49	10	62	50	12	387.5	500.0	200.0
Arteriosclerosis (450)	4	0	4	4	1	3	25.0	10.0	50.0
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	3	1	2	5	3	2	31.2	30.0	33.3
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	2	1	1	12.5	10.0	16.7
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	3	1	2	4	2	2	25.0	20.0	33.3
Birth inj., atelect. (760-762)	2	1	1	3	2	1	7.6	10.4	5.0
#Other dis. early inf. (763-776)	3	2	1	4	2	2	10.2	10.4	10.0
Violent & acc. deaths (800-999)	14	7	7	15	6	9	93.8	60.0	150.0
Acc. deaths, total (800-962)	10	5	5	11	4	7	68.8	40.0	116.7
Motor vehicle acc. (810-835)	7	5	2	6	3	3	37.5	30.0	50.0
Ill-defined (780-795)	17	4	13	19	4	15	118.8	40.0	250.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CLAY COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	487	210	277	501	223	278	31.3	27.9	34.8
Fetal deaths	9	1	8	12	2	10	24.0	9.0	36.0
Deaths from all causes	181	77	104	176	79	97	11.0	9.9	12.1
Deaths under 1 year	19	3	16	19	2	17	37.9	9.0	61.2
Deaths under 1 month	8	1	7	8	1	7	16.0	4.5	25.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	22	5	17	25	8	17	156.2	100.0	212.5
Diabetes (260)	1	0	1	1	0	1	6.2	-	12.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	28	15	13	27	15	12	168.8	187.5	150.0
Heart diseases (410-443)	67	33	34	64	32	32	400.0	400.0	400.0
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	6	3	3	6	3	3	37.5	37.5	37.5
#Diarrhea & enteritis (571)	3	0	3	3	0	3	18.8	-	37.5
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	0	2	2	0	2	12.5	-	25.0
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	1	1	2	1	1	12.5	12.5	12.5
Birth inj., atelect. (760-762)	4	0	4	4	0	4	8.0	-	14.4
#Other dis. early inf. (763-776)	3	1	2	2	0	2	44.0	-	7.2
Violent & acc. deaths (800-999)	13	8	5	11	8	3	68.8	100.0	37.5
Acc. deaths, total (800-962)	11	7	4	9	6	3	56.2	75.0	37.5
Motor vehicle acc. (810-835)	5	4	1	2	2	0	12.5	25.0	-
Ill-defined (780-795)	14	2	12	12	2	10	75.0	25.0	125.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
COAHOMA COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,489	363	1,126	1,539	345	1,194	35.8	20.3	45.9
Fetal deaths	69	13	56	67	8	59	43.5	23.2	49.4
Deaths from all causes	484	127	357	511	118	393	11.9	6.9	15.1
Deaths under 1 year	64	6	58	66	8	58	42.9	23.2	48.6
Deaths under 1 month	35	5	30	36	6	30	23.4	17.4	25.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	2.3	-	3.8
Tbc., respiratory (001-008)	1	0	1	1	0	1	2.3	-	3.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	3	0	3	3	0	3	7.0	-	11.5
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	2.3	-	3.8
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	55	20	35	55	18	37	127.9	105.9	142.3
Diabetes (260)	4	3	1	3	3	0	7.0	17.6	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	49	13	36	55	13	42	127.9	76.5	161.5
Heart diseases (410-443)	141	34	107	158	37	121	367.4	217.6	465.4
Arteriosclerosis (450)	7	3	4	8	2	6	18.6	11.8	23.1
Influenza (480-483)	4	0	4	3	0	3	7.0	-	11.5
#Pneumonia (490-493)	26	10	16	28	9	19	65.1	52.9	73.1
#Diarrhea & enteritis (571)	5	1	4	5	1	4	11.6	5.9	15.4
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	7	2	5	8	2	6	18.6	11.8	23.1
Maternal deaths (640-689)	3	1	2	3	0	3	1.9	-	2.5
Cong. malformations (750-759)	7	1	6	9	3	6	20.9	17.6	23.1
Birth inj., atelect. (760-762)	10	3	7	11	4	7	7.1	11.6	5.9
#Other dis. early inf. (763-776)	21	1	20	21	0	21	13.6	-	17.6
Violent & acc. deaths (800-999)	55	20	35	41	9	32	95.4	52.9	123.1
Acc. deaths, total (800-962)	43	19	24	30	8	22	69.8	47.0	84.6
Motor vehicle acc. (810-835)	13	5	8	8	2	6	18.6	11.8	23.1
Ill-defined (780-795)	38	2	36	42	3	39	97.7	17.6	150.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
COPIAH COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	875	297	578	719	252	467	25.7	18.0	33.4
Fetal deaths	23	6	17	20	2	18	27.8	7.9	38.5
Deaths from all causes	247	105	142	306	147	159	10.9	10.5	11.4
Deaths under 1 year	20	0	20	26	2	24	36.2	7.9	51.4
Deaths under 1 month	7	0	7	10	1	9	13.9	4.0	19.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	2	1	1	7.1	7.1	7.1
Tbc., respiratory (001-008)	0	0	0	2	1	1	7.1	7.1	7.1
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	1	0	1	1	0	1	3.6	-	7.1
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	34	22	12	37	25	12	132.1	178.6	85.7
Diabetes (260)	5	2	3	6	3	3	21.4	21.4	21.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	29	8	21	38	14	24	135.7	100.0	171.4
Heart diseases (410-443)	91	45	46	102	55	47	364.3	392.8	335.7
Arteriosclerosis (450)	3	2	1	4	3	1	14.3	21.4	7.1
Influenza (480-483)	1	0	1	1	0	1	3.6	-	7.1
#Pneumonia (490-493)	7	0	7	11	3	8	39.3	21.4	57.1
#Diarrhea & enteritis (571)	3	0	3	3	0	3	10.7	-	21.4
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	6	2	4	5	2	3	17.8	14.3	21.4
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	2	1	1	7.1	7.1	7.1
Birth inj., atelect. (760-762)	0	0	0	2	0	2	2.8	-	4.3
#Other dis. early inf. (763-776)	7	0	7	7	0	7	9.7	-	15.0
Violent & acc. deaths (800-999)	20	4	16	27	11	16	96.4	78.6	114.3
Acc. deaths, total (800-962)	17	3	14	24	10	14	85.7	71.4	100.0
Motor vehicle acc. (810-835)	8	3	5	12	7	5	42.8	50.0	35.7
Ill-defined (780-795)	8	4	4	10	5	5	35.7	35.7	35.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
COVINGTON COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	248	184	64	356	208	148	22.2	20.8	24.7
Fetal deaths	5	3	2	8	4	4	22.5	19.2	27.0
Deaths from all causes	105	69	36	135	86	49	8.4	8.6	8.2
Deaths under 1 year	8	4	4	13	5	8	36.5	24.0	54.0
Deaths under 1 month	5	3	2	10	4	6	28.1	19.2	40.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	6.2	-	16.7
Tbc., respiratory (001-008)	0	0	0	1	0	1	6.2	-	16.7
Syphilis (020-029)	0	0	0	1	1	0	6.2	10.0	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	1	0	6.2	10.0	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	1	0	1	1	0	1	6.2	-	16.7
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	13	8	5	14	9	5	87.5	90.0	83.3
Diabetes (260)	1	0	1	1	0	1	6.2	-	16.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	21	14	7	25	16	9	156.2	160.0	150.0
Heart diseases (410-443)	33	25	8	40	29	11	250.0	290.0	183.3
Arteriosclerosis (450)	6	4	2	7	5	2	43.8	50.0	33.3
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	8	6	2	8	6	2	50.0	60.0	33.3
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	0	1	1	0	1	6.2	-	16.7
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	6.2	-	16.7
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	6.2	10.0	-
Birth inj., atelect. (760-762)	1	1	0	1	1	0	2.8	4.8	-
#Other dis. early inf. (763-776)	3	2	1	8	3	5	22.5	14.4	33.8
Violent & acc. deaths (800-999)	8	5	3	9	5	4	56.2	50.0	66.7
Acc. deaths, total (800-962)	5	4	1	7	4	3	43.8	40.0	50.0
Motor vehicle acc. (810-835)	3	2	1	3	1	2	18.8	10.0	33.3
Ill-defined (780-795)	5	2	3	5	2	3	31.2	20.0	50.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

#Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
DESOTO COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	713	100	613	708	171	537	32.2	19.0	41.3
Fetal deaths	19	0	19	30	3	27	42.4	17.5	50.3
Deaths from all causes	156	45	111	209	70	139	9.5	7.8	10.7
Deaths under 1 year	14	0	14	24	2	22	33.9	11.7	41.0
Deaths under 1 month	8	0	8	16	1	15	22.6	5.8	27.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	4.5	-	7.7
Tbc., respiratory (001-008)	1	0	1	1	0	1	4.5	-	7.7
Syphilis (020-029)	0	0	0	1	0	1	4.5	-	7.7
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	0	2	3	0	3	13.6	-	23.1
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	2	7	14	4	10	63.6	44.4	76.9
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	25	6	19	29	7	22	131.8	77.8	169.2
Heart diseases (410-443)	36	20	16	55	34	21	250.0	377.8	161.5
Arteriosclerosis (450)	1	0	1	1	0	1	4.5	-	7.7
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	9	3	6	10	3	7	45.4	33.3	53.8
#Diarrhea & enteritis (571)	0	0	0	1	0	1	4.5	-	7.7
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	2	1	1	9.1	11.1	7.7
Maternal deaths (640-689)	0	0	0	1	0	1	1.4	-	1.9
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	2	0	2	2	0	2	2.8	-	3.7
#Other dis. early inf. (763-776)	5	0	5	14	2	12	19.8	11.7	22.3
Violent & acc. deaths (800-999)	19	7	12	22	7	15	100.0	77.8	115.4
Acc. deaths, total (800-962)	14	7	7	17	7	10	77.3	77.8	76.9
Motor vehicle acc. (810-835)	7	4	3	8	3	5	36.4	33.3	38.4
Ill-defined (780-795)	34	5	29	34	6	28	154.5	66.7	215.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
FORREST COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,482	1,099	383	1,318	846	472	25.3	22.3	33.7
Fetal deaths	53	34	19	47	22	25	35.7	26.0	53.0
Deaths from all causes	483	344	139	475	307	168	9.1	8.1	12.0
Deaths under 1 year	53	31	22	52	22	30	39.4	26.0	63.6
Deaths under 1 month	39	24	15	34	14	20	25.8	16.5	42.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	4	2	2	7.7	5.3	14.3
Tbc., respiratory (001-008)	1	0	1	4	2	2	7.7	5.3	14.3
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	2	0	1	1	0	1.9	2.6	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	1	1	0	1.9	2.6	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	65	52	13	62	49	13	119.2	128.9	92.8
Diabetes (260)	7	3	4	8	3	5	15.4	7.9	35.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	52	36	16	51	32	19	98.1	84.2	135.7
Heart diseases (410-443)	161	120	41	149	104	45	286.5	273.7	321.4
Arteriosclerosis (450)	11	9	2	9	8	1	17.3	21.0	7.1
Influenza (480-483)	5	4	1	5	4	1	9.6	10.5	7.1
#Pneumonia (490-493)	12	12	0	12	10	2	23.1	26.3	14.3
#Diarrhea & enteritis (571)	2	1	1	4	1	3	7.7	2.6	21.4
Acute nephritis (590, 591)	3	1	2	3	1	2	5.8	2.6	14.3
Chronic nephritis, etc. (592-594)	2	2	0	2	2	0	3.8	5.3	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	8	3	5	7	2	5	13.5	5.3	35.7
Birth inj., atelect. (760-762)	9	7	2	8	5	3	6.1	5.9	6.4
#Other dis. early inf. (763-776)	23	13	10	22	8	14	16.7	9.4	29.7
Violent & acc. deaths (800-999)	44	31	13	40	26	14	76.9	68.4	100.0
Acc. deaths, total (800-962)	39	29	10	32	22	10	61.5	57.9	71.4
Motor vehicle acc. (810-835)	19	15	4	15	13	2	28.8	34.2	14.3
Ill-defined (780-795)	28	8	20	30	8	22	57.7	21.0	157.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
FRANKLIN COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	121	56	65	205	91	114	22.8	18.2	28.5
Fetal deaths	7	1	6	9	3	6	43.9	33.0	52.6
Deaths from all causes	80	51	29	106	71	35	11.8	14.2	8.8
Deaths under 1 year	5	0	5	6	1	5	29.3	11.0	43.8
Deaths under 1 month	3	0	3	3	0	3	14.6	-	26.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	11	10	1	19	15	4	211.1	300.0	100.0
Diabetes (260)	6	3	3	6	3	3	66.7	60.0	75.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	5	2	3	10	7	3	111.1	140.0	75.0
Heart diseases (410-443)	34	25	9	41	30	11	455.6	600.0	275.0
Arteriosclerosis (450)	3	2	1	3	2	1	33.3	40.0	25.0
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	2	0	3	3	0	33.3	60.0	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	3	0	3	3	0	3	14.6	-	26.3
Violent & acc. deaths (800-999)	9	4	5	9	4	5	100.0	80.0	125.0
Acc. deaths, total (800-962)	7	3	4	7	3	4	77.8	60.0	100.0
Motor vehicle acc. (810-835)	2	1	1	1	0	1	11.1	-	25.0
Ill-defined (780-795)	3	0	3	3	0	3	33.3	-	75.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
GEORGE COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	334	282	52	336	281	55	22.4	21.6	27.5
Fetal deaths	8	5	3	8	5	3	23.8	17.8	54.5
Deaths from all causes	77	65	12	97	83	14	6.5	6.4	7.0
Deaths under 1 year	9	6	3	8	7	1	23.8	24.9	18.2
Deaths under 1 month	4	3	1	4	4	0	11.9	14.2	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	1	1	0	1	1	0	6.7	7.7	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	7	5	2	14	11	3	93.3	84.6	150.0
Diabetes (260)	2	1	1	2	1	1	13.3	7.7	50.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	1	1	0	3	2	1	20.0	15.4	50.0
Heart diseases (410-443)	39	37	2	41	38	3	273.3	292.3	150.0
Arteriosclerosis (450)	1	0	1	1	0	1	6.7	-	50.0
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	4	2	2	4	3	1	26.7	23.1	50.0
#Diarrhea & enteritis (571)	1	1	0	1	1	0	6.7	7.7	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	6.7	7.7	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	2	0	3	3	0	20.0	23.1	-
Birth inj., atelect. (760-762)	1	1	0	2	2	0	6.0	7.1	-
#Other dis. early inf. (763-776)	2	1	1	1	1	0	3.0	3.6	-
Violent & acc. deaths (800-999)	11	8	3	17	13	4	113.3	100.0	200.0
Acc. deaths, total (800-962)	10	7	3	14	11	3	93.3	84.6	150.0
Motor vehicle acc. (810-835)	5	4	1	8	7	1	53.3	53.8	50.0
Ill-defined (780-795)	0	0	0	1	1	0	6.7	7.7	-

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
GREENE COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	145	94	51	208	143	65	26.0	23.8	32.5
Fetal deaths	6	4	2	6	4	2	28.8	28.0	30.8
Deaths from all causes	66	48	18	83	57	26	10.4	9.5	13.0
Deaths under 1 year	5	3	2	10	3	7	48.1	21.0	107.7
Deaths under 1 month	2	2	0	4	2	2	19.2	14.0	30.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	1	0	1	1	0	1	12.5	-	50.0
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	12.5	-	50.0
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	7	4	3	13	8	5	162.5	133.3	250.0
Diabetes (260)	0	0	0	1	1	0	12.5	16.7	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	6	5	1	7	6	1	87.5	100.0	50.0
Heart diseases (410-443)	25	19	6	31	25	6	387.5	416.7	300.0
Arteriosclerosis (450)	1	1	0	1	1	0	12.5	16.7	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	0	1	2	0	2	25.0	-	100.0
#Diarrhea & enteritis (571)	1	1	0	1	0	1	12.5	-	50.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	12.5	16.7	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	2	0	2	2	0	25.0	33.3	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	0	0	0	2	0	2	9.6	-	30.8
Violent & acc. deaths (800-999)	12	8	4	7	5	2	87.5	83.3	100.0
Acc. deaths, total (800-962)	10	6	4	7	5	2	87.5	83.3	100.0
Motor vehicle acc. (810-835)	2	0	2	2	1	1	25.0	16.7	50.0
Ill-defined (780-795)	2	2	0	3	2	1	37.5	33.3	50.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
GRENADA COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	490	167	323	470	147	323	26.1	16.3	35.9
Fetal deaths	26	7	19	28	9	19	59.6	61.2	58.8
Deaths from all causes	183	76	107	165	62	103	9.2	6.9	11.4
Deaths under 1 year	25	3	22	26	4	22	55.3	27.2	68.1
Deaths under 1 month	16	2	14	16	3	13	34.0	20.4	40.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	5.6	-	11.1
Tbc., respiratory (001-008)	0	0	0	1	0	1	5.6	-	11.1
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	5.6	-	11.1
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	11	6	16	9	7	88.9	100.0	77.8
Diabetes (260)	1	0	1	1	0	1	5.6	-	11.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	31	11	20	27	7	20	150.0	77.8	222.2
Heart diseases (410-443)	47	21	26	46	20	26	255.6	222.2	288.9
Arteriosclerosis (450)	2	1	1	4	2	2	22.2	22.2	22.2
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	5	2	3	6	2	4	33.3	22.2	44.4
#Diarrhea & enteritis (571)	3	0	3	3	0	3	16.7	-	33.3
Acute nephritis (590, 591)	1	0	1	1	0	1	5.6	-	11.1
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	5.6	11.1	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	2	0	2	2	0	11.1	22.2	-
Birth inj., atelect. (760-762)	0	0	0	1	1	0	2.1	6.8	-
#Other dis. early inf. (763-776)	14	2	12	13	2	11	27.6	13.6	34.0
Violent & acc. deaths (800-999)	26	14	12	16	7	9	88.9	77.8	100.0
Acc. deaths, total (800-962)	20	10	10	11	4	7	61.1	44.4	77.8
Motor vehicle acc. (810-835)	9	6	3	5	2	3	27.8	22.2	33.3
Ill-defined (780-795)	15	2	13	15	2	13	83.3	22.2	144.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HANCOCK COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	165	132	33	334	279	55	25.7	27.9	18.3
Fetal deaths	4	3	1	6	4	2	18.0	14.3	36.4
Deaths from all causes	107	90	17	139	112	27	10.7	11.2	9.0
Deaths under 1 year	5	3	2	12	8	4	35.9	28.7	72.7
Deaths under 1 month	2	1	1	7	4	3	21.0	14.3	54.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	7.7	10.0	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	7.7	10.0	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	1	0	7.7	10.0	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	7	2	24	20	4	184.6	200.0	133.3
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	20	17	3	22	15	7	169.2	150.0	233.3
Heart diseases (410-443)	35	34	1	41	39	2	315.4	390.0	66.7
Arteriosclerosis (450)	0	0	0	1	0	1	7.7	-	33.3
Influenza (480-483)	1	1	0	1	1	0	7.7	10.0	-
#Pneumonia (490-493)	4	4	0	6	6	0	46.2	60.0	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	7.7	10.0	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	7.7	10.0	-
Birth inj., atelect. (760-762)	0	0	0	4	2	2	12.0	7.2	36.4
#Other dis. early inf. (763-776)	1	1	0	2	2	0	6.0	7.2	-
Violent & acc. deaths (800-999)	12	10	2	10	9	1	76.9	90.0	33.3
Acc. deaths, total (800-962)	10	8	2	8	7	1	61.5	70.0	33.3
Motor vehicle acc. (810-835)	7	5	2	4	4	0	30.8	40.0	-
Ill-defined (780-795)	16	9	7	11	5	6	84.6	50.0	200.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HARRISON COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	4,337	3,562	775	4,001	3,258	743	24.1	24.0	24.8
Fetal deaths	102	62	40	94	56	38	23.5	17.2	51.1
Deaths from all causes	1,079	851	228	908	698	210	5.5	5.1	7.0
Deaths under 1 year	114	78	36	107	74	33	26.7	22.7	44.4
Deaths under 1 month	88	61	27	79	55	24	19.7	16.9	32.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	9	5	4	9	5	4	5.4	3.7	13.3
Tbc., respiratory (001-008)	8	5	3	8	5	3	4.8	3.7	10.0
Syphilis (020-029)	2	0	2	3	0	3	1.8	-	10.0
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	3	3	0	1	1	0	0.6	0.7	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	133	107	26	113	90	23	68.1	66.2	76.7
Diabetes (260)	14	9	5	13	8	5	7.8	5.9	16.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	103	65	38	92	57	35	55.4	41.9	116.7
Heart diseases (410-443)	351	301	50	288	244	44	173.5	179.4	146.7
Arteriosclerosis (450)	14	12	2	13	12	1	7.8	8.8	3.3
Influenza (480-483)	2	2	0	2	2	0	1.2	1.5	-
#Pneumonia (490-493)	29	23	6	21	15	6	12.6	11.0	20.0
#Diarrhea & enteritis (571)	2	2	0	1	1	0	0.6	0.7	-
Acute nephritis (590, 591)	1	0	1	1	0	1	0.6	-	3.3
Chronic nephritis, etc. (592-594)	7	5	2	7	5	2	4.2	3.7	6.7
Maternal deaths (640-689)	1	1	0	1	1	0	0.2	0.3	-
Cong. malformations (750-759)	18	15	3	24	20	4	14.4	14.7	13.3
Birth inj., atelect. (760-762)	32	22	10	25	18	7	6.2	5.5	9.4
#Other dis. early inf. (763-776)	39	28	11	37	26	11	9.2	8.0	14.8
Violent & acc. deaths (800-999)	99	84	15	79	66	13	47.6	48.5	43.3
Acc. deaths, total (800-962)	80	68	12	66	56	10	39.8	41.2	33.3
Motor vehicle acc. (810-835)	24	23	1	25	23	2	15.1	16.9	6.7
Ill-defined (780-795)	84	56	28	72	46	26	43.4	33.8	86.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HINDS COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	5,739	3,002	2,737	5,333	2,659	2,674	30.1	25.6	36.6
Fetal deaths	390	256	134	345	230	115	64.7	86.5	43.0
Deaths from all causes	2,102	1,125	977	1,498	768	730	8.5	7.4	10.0
Deaths under 1 year	259	93	166	185	63	122	34.7	23.7	45.6
Deaths under 1 month	174	66	108	131	51	80	24.6	19.2	29.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	20	6	14	16	5	11	9.0	4.8	15.1
Tbc., respiratory (001-008)	15	6	9	12	5	7	6.8	4.8	9.6
Syphilis (020-029)	8	2	6	6	1	5	3.4	1.0	6.8
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	0.6	-	1.4
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	0.6	-	1.4
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	2	2	0	1	1	0	0.6	1.0	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	312	184	128	194	114	80	109.6	109.6	109.6
Diabetes (260)	34	14	20	21	6	15	11.9	5.8	20.5
Pellagra (281)	2	0	2	2	0	2	1.1	-	2.7
Intracranial lesions (330-334)	187	109	78	152	89	63	85.9	85.6	86.3
Heart diseases (410-443)	553	363	190	443	280	163	250.3	269.2	223.3
Arteriosclerosis (450)	24	16	8	18	11	7	10.2	10.6	9.6
Influenza (480-483)	1	1	0	1	1	0	0.6	1.0	-
#Pneumonia (490-493)	71	38	33	49	23	26	27.7	22.1	35.6
#Diarrhea & enteritis (571)	7	0	7	6	1	5	3.4	1.0	6.8
Acute nephritis (590, 591)	7	1	6	4	0	4	2.2	-	5.5
Chronic nephritis, etc. (592-594)	23	5	18	22	5	17	12.4	4.8	23.3
Maternal deaths (640-689)	2	0	2	1	0	1	0.2	-	0.4
Cong. malformations (750-759)	36	20	16	13	9	4	7.3	8.6	5.5
Birth inj., atelect. (760-762)	41	18	23	32	14	18	6.0	5.3	6.7
#Other dis. early inf. (763-776)	99	37	62	80	30	50	15.0	11.3	18.7
Violent & acc. deaths (800-999)	181	90	91	126	61	65	71.2	58.6	89.0
Acc. deaths, total (800-962)	130	69	61	82	44	38	46.3	42.3	52.0
Motor vehicle acc. (810-835)	58	39	19	37	26	11	20.9	25.0	15.1
Ill-defined (780-795)	156	40	116	144	32	112	81.4	30.8	153.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HOLMES COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	783	127	656	779	122	657	33.9	20.3	38.6
Fetal deaths	59	10	49	59	12	47	75.7	98.4	71.5
Deaths from all causes	299	72	227	333	94	239	14.5	15.7	14.0
Deaths under 1 year	32	2	30	40	4	36	51.3	32.8	54.8
Deaths under 1 month	21	1	20	27	2	25	34.6	16.4	38.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	0	3	4	0	4	17.4	-	23.5
Tbc., respiratory (001-008)	3	0	3	4	0	4	17.4	-	23.5
Syphilis (020-029)	3	0	3	3	0	3	13.0	-	17.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	41	15	26	47	19	28	204.3	316.7	164.7
Diabetes (260)	5	1	4	5	1	4	21.7	16.7	23.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	51	11	40	53	14	39	230.4	233.3	229.4
Heart diseases (410-443)	69	23	46	80	32	48	347.8	533.3	282.4
Arteriosclerosis (450)	4	0	4	3	0	3	13.0	-	17.6
Influenza (480-483)	1	1	0	1	1	0	4.3	16.7	-
#Pneumonia (490-493)	11	2	9	12	3	9	52.2	50.0	52.9
#Diarrhea & enteritis (571)	1	0	1	1	0	1	4.3	-	5.9
Acute nephritis (590, 591)	1	0	1	1	0	1	4.3	-	5.9
Chronic nephritis, etc. (592-594)	5	0	5	4	0	4	17.4	-	23.5
Maternal deaths (640-689)	1	0	1	1	0	1	1.3	-	1.5
Cong. malformations (750-759)	2	0	2	3	0	3	13.0	-	17.6
Birth inj., atelect. (760-762)	4	0	4	5	0	5	6.4	-	7.6
#Other dis. early inf. (763-776)	12	1	11	16	2	14	20.5	16.4	21.3
Violent & acc. deaths (800-999)	28	7	21	23	7	1	100.0	116.7	94.1
Acc. deaths, total (800-962)	24	6	18	18	5	13	78.3	83.3	76.5
Motor vehicle acc. (810-835)	13	4	9	3	1	2	13.0	16.7	11.8
Ill-defined (780-795)	35	3	32	38	4	34	165.2	66.7	200.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HUMPHREYS COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	607	91	516	615	100	515	38.4	16.7	51.5
Fetal deaths	25	0	25	27	1	26	43.9	10.0	50.5
Deaths from all causes	173	37	136	200	47	153	12.5	7.8	15.3
Deaths under 1 year	36	4	32	41	4	37	66.7	40.0	71.8
Deaths under 1 month	19	3	16	25	3	22	40.6	30.0	42.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	19	7	12	22	8	14	137.5	133.3	140.0
Diabetes (260)	2	0	2	3	0	3	18.8	-	30.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	20	1	19	24	4	20	150.0	66.7	200.0
Heart diseases (410-443)	30	16	14	35	19	16	218.8	316.7	160.0
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	1	0	1	1	0	1	6.2	-	10.0
#Pneumonia (490-493)	6	0	6	7	0	7	43.8	-	70.0
#Diarrhea & enteritis (571)	6	0	6	5	0	5	31.2	-	50.0
Acute nephritis (590, 591)	1	0	1	1	0	1	6.2	-	10.0
Chronic nephritis, etc. (592-594)	2	0	2	2	0	2	12.5	-	20.0
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	2	1	1	12.5	16.7	10.0
Birth inj., atelect. (760-762)	3	1	2	3	1	2	4.9	10.0	3.9
#Other dis. early inf. (763-776)	14	1	13	19	1	18	30.9	10.0	35.0
Violent & acc. deaths (800-999)	25	6	19	27	6	21	168.8	100.0	210.0
Acc. deaths, total (800-962)	17	4	13	19	4	15	118.8	66.7	150.0
Motor vehicle acc. (810-835)	7	2	5	8	2	6	50.0	33.3	60.0
III-defined (780-795)	30	1	29	28	1	27	175.0	16.7	270.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ISSAQUNA COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	44	1	43	120	22	98	30.0	11.0	49.0
Fetal deaths	1	0	1	4	1	3	33.3	45.4	30.6
Deaths from all causes	33	4	29	51	9	42	12.8	4.5	21.0
Deaths under 1 year	3	0	3	7	0	7	58.3	-	71.4
Deaths under 1 month	1	0	1	5	0	5	41.7	-	51.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	2	0	2	3	0	3	75.0	-	150.0
Diabetes (260)	1	0	1	3	0	3	75.0	-	150.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	3	0	3	4	1	3	100.0	50.0	150.0
Heart diseases (410-443)	9	1	8	18	7	11	450.0	350.0	550.0
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	0	2	2	0	2	50.0	-	100.0
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	0	1	1	0	1	25.0	-	50.0
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	0	1	25.0	-	50.0
Birth inj., atelect. (760-762)	0	0	0	1	0	1	8.3	-	10.2
#Other dis. early inf. (763-776)	1	0	1	4	0	4	33.3	-	40.8
Violent & acc. deaths (800-999)	5	2	3	3	0	3	75.0	-	150.0
Acc. deaths, total (800-962)	5	2	3	3	0	3	75.0	-	150.0
Motor vehicle acc. (810-835)	3	1	2	2	0	2	50.0	-	100.0
Ill-defined (780-795)	8	0	8	9	0	9	225.0	-	450.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ITAWAMBA COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	79	62	17	283	257	26	13.5	13.2	17.3
Fetal deaths	1	1	0	7	7	0	24.7	27.2	-
Deaths from all causes	81	76	5	119	113	6	5.7	5.8	4.0
Deaths under 1 year	1	1	0	11	10	1	38.9	38.9	38.5
Deaths under 1 month	1	1	0	10	9	1	35.3	35.0	38.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	4.8	5.1	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	4.8	5.1	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	1	1	0	4.8	5.1	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	21	20	1	24	23	1	114.3	117.9	66.7
Diabetes (260)	3	3	0	3	3	0	14.3	15.4	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	12	10	2	15	13	2	71.4	66.7	133.3
Heart diseases (410-443)	29	29	0	38	38	0	181.0	194.9	-
Arteriosclerosis (450)	2	2	0	4	4	0	19.0	20.5	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	1	1	4	3	1	19.0	15.4	66.7
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	1	1	0	3.5	3.9	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	5	4	1	17.7	15.6	38.5
#Other dis. early inf. (763-776)	1	1	0	5	5	0	17.7	19.4	-
Violent & acc. deaths (800-999)	3	2	1	6	5	1	28.6	25.6	66.7
Acc. deaths, total (800-962)	3	2	1	6	5	1	28.6	25.6	66.7
Motor vehicle acc. (810-835)	2	2	0	4	4	0	19.0	20.5	-
Ill-defined (780-795)	5	5	0	5	5	0	23.8	25.6	-

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JACKSON COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,393	987	406	1,730	1,296	434	25.8	24.9	28.9
Fetal deaths	80	52	28	86	55	31	49.7	42.4	71.4
Deaths from all causes	301	211	90	334	244	90	5.0	4.7	6.0
Deaths under 1 year	31	14	17	40	22	18	23.1	17.0	41.5
Deaths under 1 month	21	12	9	31	20	11	17.9	15.4	25.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	0	1	2	1	1	3.0	1.9	6.7
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	33	20	13	39	27	12	58.2	51.9	80.0
Diabetes (260)	3	2	1	3	2	1	4.5	3.8	6.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	51	33	18	46	30	16	68.6	57.7	106.7
Heart diseases (410-443)	72	62	10	81	69	12	120.9	132.7	80.0
Arteriosclerosis (450)	6	5	1	5	4	1	7.5	7.7	6.7
Influenza (480-483)	2	1	1	2	1	1	3.0	1.9	6.7
#Pneumonia (490-493)	5	3	2	6	4	2	9.0	7.7	13.3
#Diarrhea & enteritis (571)	2	0	2	3	1	2	4.5	1.9	13.3
Acute nephritis (590, 591)	2	2	0	2	2	0	3.0	3.8	-
Chronic nephritis, etc. (592-594)	3	2	1	3	2	1	4.5	3.8	6.7
Maternal deaths (640-689)	1	1	0	1	1	0	0.6	0.8	-
Cong. malformations (750-759)	6	5	1	7	6	1	10.4	11.5	6.7
Birth inj., atelect. (760-762)	8	6	2	11	8	3	6.4	6.2	6.9
#Other dis. early inf. (763-776)	10	3	7	14	7	7	8.1	5.4	16.1
Violent & acc. deaths (800-999)	50	35	15	53	39	14	79.1	75.0	93.3
Acc. deaths, total (800-962)	43	29	14	44	31	13	65.7	59.6	86.7
Motor vehicle acc. (810-835)	10	5	5	13	8	5	19.4	15.4	33.3
Ill-defined (780-795)	25	18	7	23	17	6	34.3	32.7	40.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JASPER COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence:

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	106	25	81	427	149	278	23.7	14.9	34.8
Fetal deaths	3	0	3	20	7	13	46.8	47.0	46.8
Deaths from all causes	97	42	55	160	76	84	8.9	7.6	10.5
Deaths under 1 year	4	0	4	15	1	14	35.1	6.7	50.4
Deaths under 1 month	0	0	0	10	1	9	23.4	6.7	32.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	5.6	10.0	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	5.6	10.0	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	4	5	14	7	7	77.8	70.0	87.5
Diabetes (260)	1	1	0	2	1	1	11.1	10.0	12.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	14	6	8	23	15	8	127.8	150.0	100.0
Heart diseases (410-443)	34	18	16	50	29	21	277.8	290.0	262.5
Arteriosclerosis (450)	0	0	0	1	0	1	5.6	-	12.5
Influenza (480-483)	1	0	1	1	0	1	5.6	-	12.5
#Pneumonia (490-493)	4	3	1	7	4	3	38.9	40.0	37.5
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	5	3	2	4	3	1	22.2	30.0	12.5
Chronic nephritis, etc. (592-594)	2	0	2	4	1	3	22.2	10.0	37.5
Maternal deaths (640-689)	0	0	0	1	0	1	2.3	-	3.6
Cong. malformations (750-759)	0	0	0	1	1	0	5.6	10.0	-
Birth inj., atelect. (760-762)	0	0	0	2	0	2	4.7	-	7.2
#Other dis. early inf. (763-776)	0	0	0	7	0	7	16.4	-	25.2
Violent & acc. deaths (800-999)	11	6	5	17	9	8	94.4	90.0	100.0
Acc. deaths, total (800-962)	8	5	3	13	8	5	72.2	80.0	62.5
Motor vehicle acc. (810-835)	3	3	0	6	6	0	33.3	60.0	-
Ill-defined (780-795)	13	0	13	15	1	14	83.3	10.0	175.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JEFFERSON COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	122	0	122	296	36	260	24.7	12.0	28.9
Fetal deaths	8	0	8	18	2	16	60.8	55.6	61.5
Deaths from all causes	84	14	70	128	30	98	10.7	10.0	10.9
Deaths under 1 year	9	0	9	13	0	13	43.9	-	50.0
Deaths under 1 month	0	0	0	2	0	2	6.8	-	7.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	1	0	1	8.3	-	11.1
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	15	4	11	23	6	17	191.7	200.0	188.9
Diabetes (260)	1	0	1	2	1	1	16.7	33.3	11.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	12	2	10	15	2	13	125.0	66.7	144.4
Heart diseases (410-443)	30	6	24	44	16	28	366.7	533.3	311.1
Arteriosclerosis (450)	0	0	0	1	1	0	8.3	33.3	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	0	2	4	0	4	33.3	-	44.4
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	1	0	1	8.3	-	11.1
Chronic nephritis, etc. (592-594)	3	0	3	3	0	3	25.0	-	33.3
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	0	2	2	0	2	16.7	-	22.2
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	3	0	3	5	0	5	16.9	-	19.2
Violent & acc. deaths (800-999)	6	1	5	10	2	8	83.3	66.7	88.9
Acc. deaths, total (800-962)	6	1	5	8	2	6	66.7	66.7	66.7
Motor vehicle acc. (810-835)	0	0	0	1	1	0	8.3	33.3	-
Ill-defined (780-795)	3	0	3	5	0	5	41.7	-	55.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JEFFERSON DAVIS COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	332	154	178	372	138	234	19.6	15.3	23.4
Fetal deaths	13	1	12	14	1	13	37.6	7.2	55.6
Deaths from all causes	111	61	50	108	56	52	5.7	6.2	5.2
Deaths under 1 year	15	1	14	14	1	13	37.6	7.2	55.6
Deaths under 1 month	9	1	8	9	1	8	24.2	7.2	34.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	19	12	7	22	12	10	115.8	133.3	100.0
Diabetes (260)	1	1	0	1	1	0	5.3	11.1	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	7	5	2	7	5	2	36.8	55.6	20.0
Heart diseases (410-443)	27	17	10	29	18	11	152.6	200.0	110.0
Arteriosclerosis (450)	1	1	0	2	1	1	10.5	11.1	10.0
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	3	0	3	3	0	3	15.8	-	30.0
#Diarrhea & enteritis (571)	1	0	1	1	0	1	5.3	-	10.0
Acute nephritis (590, 591)	2	1	1	2	1	1	10.5	11.1	10.0
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	1	0	1	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	1	0	1	5.3	-	10.0
Birth inj., atelect. (760-762)	1	0	1	1	0	1	2.7	-	4.3
#Other dis. early inf. (763-776)	2	0	2	2	0	2	5.4	-	8.5
Violent & acc. deaths (800-999)	17	9	8	14	7	7	73.7	77.8	70.0
Acc. deaths, total (800-962)	12	7	5	10	5	5	52.6	55.6	50.0
Motor vehicle acc. (810-835)	5	2	3	5	2	3	26.3	22.2	30.0
Ill-defined (780-795)	14	6	8	11	3	8	57.9	33.3	80.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JONES COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	2,480	1,264	1,216	1,518	979	539	21.1	17.8	31.7
Fetal deaths	83	35	48	46	25	21	30.3	25.5	39.0
Deaths from all causes	665	399	266	507	340	167	7.0	6.2	9.8
Deaths under 1 year	100	32	68	56	24	32	36.9	24.5	59.4
Deaths under 1 month	76	25	51	42	19	23	27.7	19.4	42.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	4	2	2	5.6	3.6	11.8
Tbc., respiratory (001-008)	1	1	0	3	2	1	4.2	3.6	5.9
Syphilis (020-029)	1	1	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	1.4	-	5.9
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	2	1	1	1	0	1	1.4	-	5.9
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	82	59	23	71	57	14	98.6	103.6	82.4
Diabetes (260)	14	6	8	10	5	5	13.9	9.1	29.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	100	57	43	80	48	32	111.1	87.3	188.2
Heart diseases (410-443)	176	133	43	144	108	36	200.0	196.4	211.8
Arteriosclerosis (450)	6	5	1	6	5	1	8.3	9.1	5.9
Influenza (480-483)	2	1	1	1	1	0	1.4	1.8	-
#Pneumonia (490-493)	20	10	10	16	9	7	22.2	16.4	41.2
#Diarrhea & enteritis (571)	11	3	8	3	2	1	4.2	3.6	5.9
Acute nephritis (590, 591)	3	1	2	3	1	2	4.2	1.8	11.8
Chronic nephritis, etc. (592-594)	5	3	2	3	2	1	4.2	3.6	5.9
Maternal deaths (640-689)	3	0	3	1	0	1	0.6	-	1.8
Cong. malformations (750-759)	8	6	2	4	3	1	5.6	5.4	5.9
Birth inj., atelect. (760-762)	18	10	8	12	8	4	7.9	8.2	7.4
#Other dis. early inf. (763-776)	53	12	41	26	8	18	17.1	8.2	33.4
Violent & acc. deaths (800-999)	51	32	19	39	25	14	54.2	45.4	82.4
Acc. deaths, total (800-962)	38	27	11	28	21	7	38.9	38.2	41.2
Motor vehicle acc. (810-835)	19	15	4	12	9	3	16.7	16.4	17.6
III-defined (780-795)	36	22	14	31	21	10	43.0	38.2	58.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
KEMPER COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	206	44	162	286	77	209	23.8	15.4	29.8
Fetal deaths	10	0	10	12	1	11	42.0	13.0	52.6
Deaths from all causes	99	47	52	124	61	63	10.3	12.2	9.0
Deaths under 1 year	6	2	4	6	2	4	21.0	26.0	19.1
Deaths under 1 month	2	0	2	2	0	2	7.0	-	9.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	12	4	8	16	6	10	133.3	120.0	142.8
Diabetes (260)	2	1	1	2	1	1	16.7	20.0	14.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	12	9	3	15	12	3	125.0	240.0	42.8
Heart diseases (410-443)	28	13	15	36	18	18	300.0	360.0	257.1
Arteriosclerosis (450)	3	1	2	3	1	2	25.0	20.0	28.6
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	6	5	1	7	6	1	58.3	120.0	14.3
#Diarrhea & enteritis (571)	1	0	1	1	0	1	8.3	-	14.3
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	3	0	3	25.0	-	42.8
Maternal deaths (640-689)	0	0	0	1	1	0	3.5	13.0	-
Cong. malformations (750-759)	2	2	0	2	2	0	16.7	40.0	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	1	0	1	1	0	1	3.5	-	4.8
Violent & acc. deaths (800-999)	9	4	5	11	4	7	91.7	80.0	100.0
Acc. deaths, total (800-962)	6	3	3	8	3	5	66.7	60.0	71.4
Motor vehicle acc. (810-835)	0	0	0	1	0	1	8.3	-	14.3
Ill-defined (780-795)	16	3	13	17	4	13	141.7	80.0	185.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LAFAYETTE COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	510	266	244	532	266	266	20.5	14.8	33.2
Fetal deaths	26	8	18	26	8	18	48.9	30.1	67.7
Deaths from all causes	163	93	70	186	110	76	7.2	6.1	9.5
Deaths under 1 year	15	4	11	18	5	13	33.8	18.8	48.9
Deaths under 1 month	11	3	8	13	4	9	24.4	15.0	33.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	3.8	-	12.5
Tbc., respiratory (001-008)	1	0	1	1	0	1	3.8	-	12.5
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	1	1	0	1	1	0	3.8	5.6	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	23	17	6	28	20	8	107.7	111.1	100.0
Diabetes (260)	2	1	1	2	1	1	7.7	5.6	12.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	22	14	8	23	16	7	88.5	88.9	87.5
Heart diseases (410-443)	47	34	13	53	39	14	203.8	216.7	175.0
Arteriosclerosis (#50)	5	5	0	5	5	0	19.2	27.8	-
Influenza (480-483)	2	0	2	1	0	1	3.8	-	12.5
#Pneumonia (490-493)	7	3	4	9	5	4	34.6	27.8	50.0
#Diarrhea & enteritis (571)	1	0	1	1	0	1	3.8	-	12.5
Acute nephritis (590, 591)	2	0	2	2	0	2	7.7	-	25.0
Chronic nephritis, etc. (592-594)	5	1	4	6	2	4	23.1	11.1	50.0
Maternal deaths (640-689)	1	0	1	1	0	1	1.9	-	3.8
Cong. malformations (750-759)	0	0	0	1	0	1	3.8	-	12.5
Birth inj., atelect. (760-762)	2	0	2	2	0	2	3.8	-	7.5
#Other dis. early inf. (763-776)	6	2	4	7	2	5	13.2	7.5	18.8
Violent & acc. deaths (800-999)	16	9	7	17	10	7	65.4	55.6	87.5
Acc. deaths, total (800-962)	8	6	2	10	7	3	38.5	38.9	37.5
Motor vehicle acc. (810-835)	3	2	1	3	2	1	11.5	11.1	12.5
Ill-defined (780-795)	12	3	9	13	4	9	50.0	22.2	112.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LAMAR COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	90	52	38	303	227	76	16.8	16.2	19.0
Fetal deaths	1	0	1	8	6	2	26.4	26.4	26.3
Deaths from all causes	77	64	13	118	95	23	6.6	6.8	5.8
Deaths under 1 year	1	0	1	9	8	1	29.7	35.2	13.2
Deaths under 1 month	0	0	0	8	8	0	26.4	35.2	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	2	1	1	11.1	7.1	25.0
Tbc., respiratory (001-008)	1	0	1	2	1	1	11.1	7.1	25.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	13	12	1	18	15	3	100.0	107.1	75.0
Diabetes (260)	2	2	0	1	1	0	5.6	7.1	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	9	8	1	14	9	5	77.8	64.3	125.0
Heart diseases (410-443)	25	22	3	41	37	4	227.8	264.3	100.0
Arteriosclerosis (450)	3	3	0	3	3	0	16.7	21.4	-
Influenza (480-483)	1	1	0	1	1	0	5.6	7.1	-
#Pneumonia (490-493)	4	3	1	4	3	1	22.2	21.4	25.0
#Diarrhea & enteritis (571)	1	0	1	1	0	1	5.6	-	25.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	1	0	1	1	0	1	3.3	-	13.2
Cong. malformations (750-759)	1	0	1	2	1	1	11.1	7.1	25.0
Birth inj., atelect. (760-762)	0	0	0	2	2	0	6.6	8.8	-
#Other dis. early inf. (763-776)	0	0	0	4	4	0	13.2	17.6	-
Violent & acc. deaths (800-999)	7	6	1	12	10	2	66.7	71.4	50.0
Acc. deaths, total (800-962)	7	6	1	11	10	1	61.1	71.4	25.0
Motor vehicle acc. (810-835)	4	3	1	7	7	0	38.9	50.0	-
Ill-defined (780-795)	3	2	1	4	3	1	22.2	21.4	25.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LAUDERDALE COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	2,230	1,108	1,122	1,693	875	818	28.2	23.6	35.6
Fetal deaths	93	36	57	68	30	38	40.2	34.3	46.4
Deaths from all causes	883	525	358	705	410	295	11.8	11.1	12.8
Deaths under 1 year	85	31	54	72	27	45	42.5	30.8	55.0
Deaths under 1 month	50	23	27	44	20	24	26.0	22.8	29.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	7	1	6	9	1	8	15.0	2.7	34.8
Tbc., respiratory (001-008)	7	1	6	9	1	8	15.0	2.7	34.8
Syphilis (020-029)	1	1	0	1	1	0	1.7	2.7	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	2	0	2	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	93	61	32	89	58	31	148.3	156.8	134.8
Diabetes (260)	13	3	10	11	3	8	18.3	8.1	34.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	102	75	27	77	53	24	128.3	143.2	104.3
Heart diseases (410-443)	289	192	97	225	144	81	375.0	389.2	352.2
Arteriosclerosis (450)	13	9	4	10	7	3	16.7	18.9	13.0
Influenza (480-483)	4	2	2	3	1	2	5.0	2.7	8.7
#Pneumonia (490-493)	36	19	17	26	11	15	43.3	29.7	65.2
#Diarrhea & enteritis (571)	5	3	2	3	1	2	5.0	2.7	8.7
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	8	1	7	5	0	5	8.3	-	21.7
Maternal deaths (640-689)	3	1	2	1	0	1	0.6	-	1.2
Cong. malformations (750-759)	9	3	6	6	1	5	10.0	2.7	21.7
Birth inj., atelect. (760-762)	15	11	4	12	9	3	7.1	10.3	3.7
#Other dis. early inf. (763-776)	24	10	14	21	10	11	12.4	11.4	13.4
Violent & acc. deaths (800-999)	78	45	33	62	37	25	103.3	100.0	108.7
Acc. deaths, total (800-962)	60	39	21	46	29	17	76.7	78.4	73.9
Motor vehicle acc. (810-835)	31	23	8	26	18	8	43.3	48.6	34.8
Ill-defined (780-795)	61	25	36	59	25	34	98.3	67.6	147.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LAWRENCE COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	174	28	146	282	133	149	23.5	16.6	37.2
Fetal deaths	8	2	6	11	3	8	39.0	22.6	53.7
Deaths from all causes	73	35	38	123	65	58	10.2	8.1	14.5
Deaths under 1 year	6	1	5	9	2	7	31.9	15.0	47.0
Deaths under 1 month	3	1	2	6	2	4	21.3	15.0	26.8
SELECTED CAUSES OF DEATH									
AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	2	1	1	16.7	12.5	25.0
Tbc., respiratory (001-008)	0	0	0	1	0	1	8.3	-	25.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	7	3	17	12	5	141.7	150.0	125.0
Diabetes (260)	4	0	4	5	0	5	41.7	-	125.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	8	4	4	11	7	4	91.7	87.5	100.0
Heart diseases (410-443)	20	9	11	33	19	14	275.0	237.5	350.0
Arteriosclerosis (450)	3	1	2	4	2	2	33.3	25.0	50.0
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	5	0	5	6	0	6	50.0	-	150.0
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	2	0	2	16.7	-	50.0
Maternal deaths (640-689)	1	0	1	2	0	2	7.1	-	13.4
Cong. malformations (750-759)	2	2	0	3	3	0	25.0	37.5	-
Birth inj., atelect. (760-762)	1	0	1	2	0	2	7.1	-	13.4
#Other dis. early inf. (763-776)	0	0	0	1	0	1	3.5	-	6.7
Violent & acc. deaths (800-999)	9	6	3	16	10	6	133.3	125.0	150.0
Acc. deaths, total (800-962)	7	5	2	13	9	4	108.3	112.5	100.0
Motor vehicle acc. (810-835)	1	1	0	4	4	0	33.3	50.0	-
Ill-defined (780-795)	4	2	2	7	4	3	58.3	50.0	75.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LEAKE COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	379	174	205	453	200	253	22.6	18.2	28.1
Fetal deaths	11	0	11	13	0	13	28.7	-	51.4
Deaths from all causes	125	77	48	171	105	66	8.6	9.5	7.3
Deaths under 1 year	13	3	10	20	5	15	44.2	25.0	59.3
Deaths under 1 month	8	2	6	14	4	10	30.9	20.0	39.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	1	0	1	5.0	-	11.1
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	1	0	1	1	0	5.0	9.1	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	14	3	19	15	4	95.0	136.4	44.4
Diabetes (260)	1	1	0	2	1	1	10.0	9.1	11.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	17	10	7	26	16	10	130.0	145.4	111.1
Heart diseases (410-443)	47	31	16	55	36	19	275.0	327.3	211.1
Arteriosclerosis (450)	2	2	0	3	3	0	15.0	27.3	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	3	1	2	7	4	3	35.0	36.4	33.3
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	0	1	1	0	1	5.0	-	11.1
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	10.0	9.1	11.1
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	2	1	1	4.4	5.0	4.0
#Other dis. early inf. (763-776)	6	2	4	10	3	7	22.1	15.0	27.7
Violent & acc. deaths (800-999)	14	7	7	18	10	8	90.0	90.9	88.9
Acc. deaths, total (800-962)	12	7	5	16	10	6	80.0	90.0	66.7
Motor vehicle acc. (810-835)	6	4	2	7	5	2	35.0	45.4	22.2
Ill-defined (780-795)	2	0	2	4	1	3	20.0	9.1	33.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LEE COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,179	848	331	973	631	342	23.2	20.4	31.1
Fetal deaths	75	62	13	61	48	13	62.7	76.1	38.0
Deaths from all causes	406	280	126	402	276	126	9.6	8.9	11.4
Deaths under 1 year	51	28	23	37	15	22	38.0	23.8	64.3
Deaths under 1 month	37	23	14	25	12	13	25.7	19.0	38.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	2.4	3.2	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	2.4	3.2	-
Syphilis (020-029)	1	0	1	1	0	1	2.4	-	9.1
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	2.4	-	9.1
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	1	0	1	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	1	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	51	39	12	56	42	14	133.3	135.5	127.3
Diabetes (260)	4	2	2	5	3	2	11.9	9.7	18.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	50	28	22	55	33	22	131.0	106.4	200.0
Heart diseases (410-443)	141	99	42	137	95	42	326.2	306.4	381.8
Arteriosclerosis (450)	3	3	0	2	2	0	4.8	6.4	-
Influenza (480-483)	1	1	0	2	2	0	4.8	6.4	-
#Pneumonia (490-493)	15	10	5	14	9	5	33.3	29.0	45.4
#Diarrhea & enteritis (571)	3	0	3	3	0	3	7.1	-	27.3
Acute nephritis (590, 591)	2	0	2	2	0	2	4.8	-	18.2
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	4.8	3.2	9.1
Maternal deaths (640-689)	1	1	0	0	0	0	-	-	-
Cong. malformations (750-759)	4	3	1	5	4	1	11.9	12.9	9.1
Birth inj., atelect. (760-762)	19	13	6	12	7	5	12.3	11.1	14.6
#Other dis. early inf. (763-776)	16	8	8	11	3	8	11.3	4.8	23.4
Violent & acc. deaths (800-999)	32	28	4	38	34	4	90.5	109.7	36.4
Acc. deaths, total (800-962)	25	24	1	30	29	1	71.4	93.5	9.1
Motor vehicle acc. (810-835)	13	12	1	16	15	1	38.1	48.4	9.1
Ill-defined (780-795)	15	6	9	14	5	9	33.3	16.1	81.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

**BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LEFLORE COUNTY, MISSISSIPPI, 1959**

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,572	397	1,175	1,527	341	1,186	38.2	21.3	49.4
Fetal deaths	103	30	73	94	25	69	61.6	73.3	58.2
Deaths from all causes	563	204	359	542	173	369	13.6	10.8	15.4
Deaths under 1 year	87	13	74	87	9	78	57.0	26.4	65.8
Deaths under 1 month	46	9	37	47	5	42	30.8	14.7	35.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	0	3	5	0	5	12.5	-	20.8
Tbc., respiratory (001-008)	3	0	3	5	0	5	12.5	-	20.8
Syphilis (020-029)	2	0	2	2	0	2	5.0	-	8.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	2.5	-	4.2
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	2.5	-	4.2
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	64	33	31	62	28	34	155.0	175.0	141.7
Diabetes (260)	5	4	1	4	4	0	10.0	25.0	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	79	32	47	73	22	51	182.5	137.5	212.5
Heart diseases (410-443)	138	73	65	127	60	67	317.5	375.0	279.2
Arteriosclerosis (450)	4	3	1	4	3	1	10.0	18.8	4.2
Influenza (480-483)	2	0	2	2	0	2	5.0	-	8.3
#Pneumonia (490-493)	23	6	17	19	3	16	47.5	18.8	66.7
#Diarrhea & enteritis (571)	8	0	8	6	0	6	15.0	-	25.0
Acute nephritis (590, 591)	1	0	1	1	0	1	2.5	-	4.2
Chronic nephritis, etc. (592-594)	6	3	3	5	3	2	12.5	18.8	8.3
Maternal deaths (640-689)	0	0	0	1	0	1	0.6	-	0.8
Cong. malformations (750-759)	9	6	3	9	5	4	22.5	31.2	16.7
Birth inj., atelect. (760-762)	8	3	5	7	2	5	4.6	5.9	4.2
#Other dis. early inf. (763-776)	36	5	31	39	3	36	25.5	8.8	30.4
Violent & acc. deaths (800-999)	52	17	35	53	21	32	132.5	131.2	133.3
Acc. deaths, total (800-962)	40	14	26	41	17	24	102.5	106.2	100.0
Motor vehicle acc. (810-835)	15	7	8	17	10	7	42.5	62.5	29.2
Ill-defined (780-795)	68	8	60	65	5	60	162.5	31.2	250.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LINCOLN COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	689	415	274	663	367	296	25.5	21.6	32.9
Fetal deaths	34	16	18	28	11	17	42.2	30.0	57.4
Deaths from all causes	237	138	99	252	144	108	9.7	8.5	12.0
Deaths under 1 year	25	6	19	26	7	19	39.2	19.1	64.2
Deaths under 1 month	15	5	10	15	6	9	22.6	16.3	30.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	2	1	3	2	1	11.5	11.8	11.1
Tbc., respiratory (001-008)	3	2	1	3	2	1	11.5	11.8	11.1
Syphilis (020-029)	0	0	0	1	0	1	3.8	-	11.1
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	3.8	-	11.1
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	33	23	10	33	22	11	126.9	129.4	122.2
Diabetes (260)	4	3	1	4	4	0	15.4	23.5	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	36	12	24	35	11	24	134.6	64.7	266.7
Heart diseases (410-443)	96	65	31	102	66	36	392.3	388.2	400.0
Arteriosclerosis (450)	3	3	0	4	4	0	15.4	23.5	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	0	2	2	0	2	7.7	-	22.2
#Diarrhea & enteritis (571)	7	2	5	6	1	5	23.1	5.9	55.6
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	2	1	2	2	0	7.7	11.8	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	1	1	2	1	1	7.7	5.9	11.1
Birth inj., atelect. (760-762)	6	3	3	7	4	3	10.6	10.9	10.1
#Other dis. early inf. (763-776)	5	1	4	5	2	3	7.5	5.4	10.1
Violent & acc. deaths (800-999)	16	12	4	16	9	7	61.5	52.9	77.8
Acc. deaths, total (800-962)	14	11	3	14	8	6	53.8	47.0	66.7
Motor vehicle acc. (810-835)	8	7	1	8	5	3	30.8	29.4	33.3
Ill-defined (780-795)	9	2	7	12	3	9	46.2	17.6	100.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LOWNDES COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,509	886	623	1,423	793	630	34.7	30.5	42.0
Fetal deaths	46	22	24	43	18	25	30.2	22.7	39.7
Deaths from all causes	405	202	203	439	212	227	10.7	8.2	15.1
Deaths under 1 year	55	22	33	56	20	36	39.4	25.2	57.1
Deaths under 1 month	34	18	16	34	17	17	23.9	21.4	27.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	3	1	2	7.3	3.8	13.3
Tbc., respiratory (001-008)	1	0	1	3	1	2	7.3	3.8	13.3
Syphilis (020-029)	2	1	1	1	1	0	2.4	3.8	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	0	1	2.4	-	6.7
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	2.4	-	6.7
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	41	28	13	45	30	15	109.8	115.4	100.0
Diabetes (260)	1	1	0	1	1	0	2.4	3.8	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	49	22	27	59	27	32	143.9	103.8	213.3
Heart diseases (410-443)	127	71	56	129	69	60	314.6	265.4	400.0
Arteriosclerosis (450)	11	8	3	11	7	4	26.8	26.9	26.7
Influenza (480-483)	4	0	4	4	0	4	9.8	-	26.7
#Pneumonia (490-493)	12	4	8	14	5	9	34.1	19.2	60.0
#Diarrhea & enteritis (571)	3	1	2	3	1	2	7.3	3.8	13.3
Acute nephritis (590, 591)	1	0	1	1	0	1	2.4	-	6.7
Chronic nephritis, etc. (592-594)	4	1	3	6	1	5	14.6	3.8	33.3
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	8	6	2	11	7	4	26.8	26.9	26.7
Birth inj., atelect. (760-762)	14	11	3	12	9	3	8.4	11.3	4.8
#Other dis. early inf. (763-776)	17	5	12	17	5	12	11.9	6.3	19.0
Violent & acc. deaths (800-999)	37	20	17	45	27	18	109.8	103.8	120.0
Acc. deaths, total (800-962)	31	17	14	41	26	15	100.0	100.0	100.0
Motor vehicle acc. (810-835)	10	7	3	12	9	3	29.3	34.6	20.0
Ill-defined (780-795)	32	4	28	32	3	29	78.0	11.5	193.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MADISON COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,084	196	888	1,063	190	873	39.4	21.1	48.5
Fetal deaths	41	7	34	40	6	34	37.6	31.6	38.9
Deaths from all causes	322	97	225	333	100	233	12.3	11.1	12.9
Deaths under 1 year	30	2	28	31	3	28	29.2	15.8	32.1
Deaths under 1 month	16	2	14	17	3	14	16.0	15.8	16.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	3.7	11.1	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	3.7	11.1	-
Syphilis (020-029)	1	0	1	1	0	1	3.7	-	5.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	1	0	1	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	27	9	18	33	10	23	122.2	111.1	127.8
Diabetes (260)	6	2	4	5	2	3	18.5	22.2	16.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	55	22	33	51	19	32	188.9	211.1	177.8
Heart diseases (410-443)	73	33	40	82	33	49	303.7	366.7	272.2
Arteriosclerosis (450)	0	0	0	1	1	0	3.7	11.1	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	11	0	11	12	1	11	44.4	11.1	61.1
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	6	0	6	6	0	6	22.2	-	33.3
Maternal deaths (640-689)	1	0	1	1	0	1	0.9	-	1.1
Cong. malformations (750-759)	1	0	1	1	0	1	3.7	-	5.6
Birth inj., atelect. (760-762)	1	1	0	1	1	0	0.9	5.3	-
#Other dis. early inf. (763-776)	10	1	9	11	2	9	10.3	10.5	10.3
Violent & acc. deaths (800-999)	32	12	20	32	13	19	118.5	144.4	105.6
Acc. deaths, total (800-962)	27	10	17	27	11	16	100.0	122.2	88.9
Motor vehicle acc. (810-835)	8	7	1	8	6	2	29.6	66.7	11.1
Ill-defined (780-795)	73	5	68	68	4	64	251.8	44.4	355.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MARION COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	504	283	221	586	329	257	20.9	19.4	23.4
Fetal deaths	13	6	7	20	9	11	34.1	27.4	42.8
Deaths from all causes	180	117	63	210	133	77	7.5	7.8	7.0
Deaths under 1 year	9	1	8	13	3	10	22.2	9.1	38.9
Deaths under 1 month	5	0	5	7	1	6	11.9	3.0	23.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	3.6	-	9.1
Tbc., respiratory (001-008)	1	0	1	1	0	1	3.6	-	9.1
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	16	12	4	25	16	9	89.3	94.1	81.8
Diabetes (260)	1	1	0	3	2	1	10.7	11.8	9.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	44	35	9	46	37	9	164.3	217.6	81.8
Heart diseases (410-443)	45	32	13	51	36	15	182.1	211.8	136.4
Arteriosclerosis (450)	3	2	1	2	1	1	7.1	5.9	9.1
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	5	3	2	8	6	2	28.6	35.3	18.2
#Diarrhea & enteritis (571)	2	0	2	2	0	2	7.1	-	18.2
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	7.1	5.9	9.1
Maternal deaths (640-689)	1	0	1	1	0	1	1.7	-	3.9
Cong. malformations (750-759)	1	1	0	2	1	1	7.1	5.9	9.1
Birth inj., atelect. (760-762)	1	0	1	1	0	1	1.7	-	3.9
#Other dis. early inf. (763-776)	4	0	4	4	0	4	6.8	-	15.6
Violent & acc. deaths (800-999)	23	14	9	22	12	10	78.6	70.6	90.9
Acc. deaths, total (800-962)	22	14	8	20	11	9	71.4	64.7	81.8
Motor vehicle acc. (810-835)	9	7	2	10	7	3	35.7	41.2	27.3
Ill-defined (780-795)	12	5	7	12	5	7	42.8	29.4	63.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MARSHALL COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	751	117	634	784	146	638	32.7	20.8	37.5
Fetal deaths	46	2	44	34	4	30	43.4	27.4	47.0
Deaths from all causes	231	72	159	216	72	144	9.0	10.3	8.5
Deaths under 1 year	36	5	31	29	6	23	37.0	41.1	36.0
Deaths under 1 month	19	2	17	15	4	11	19.1	27.4	17.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	2	0	2	1	0	1	4.2	-	5.9
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	2	0	2	2	0	2	8.3	-	11.8
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	23	10	13	26	12	14	108.3	171.4	82.4
Diabetes (260)	3	1	2	2	0	2	8.3	-	11.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	27	8	19	27	6	21	112.5	85.7	123.5
Heart diseases (410-443)	72	33	39	63	28	35	262.5	400.0	205.9
Arteriosclerosis (450)	4	1	3	3	1	2	12.5	14.3	11.8
Influenza (480-483)	2	0	2	2	0	2	8.3	-	11.8
#Pneumonia (490-493)	12	3	9	11	3	8	45.8	42.8	47.0
#Diarrhea & enteritis (571)	3	0	3	3	0	3	12.5	-	17.6
Acute nephritis (590, 591)	1	0	1	1	0	1	4.2	-	5.9
Chronic nephritis, etc. (592-594)	4	1	3	2	1	1	8.3	14.3	5.9
Maternal deaths (640-689)	2	0	2	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	2	1	1	8.3	14.3	5.9
Birth inj., atelect. (760-762)	3	0	3	3	0	3	3.8	-	4.7
#Other dis. early inf. (763-776)	16	3	13	10	3	7	12.8	20.5	11.0
Violent & acc. deaths (800-999)	16	5	11	20	6	14	83.3	85.7	82.4
Acc. deaths, total (800-962)	11	3	8	14	4	10	58.3	57.1	58.8
Motor vehicle acc. (810-835)	5	2	3	7	3	4	29.2	42.8	23.5
Ill-defined (780-795)	19	1	18	19	1	18	79.2	14.3	105.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MONROE COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	822	408	414	832	436	396	23.8	19.8	30.5
Fetal deaths	30	7	23	29	9	20	34.8	20.6	50.5
Deaths from all causes	322	193	129	353	212	141	10.1	9.6	10.8
Deaths under 1 year	23	6	17	22	5	17	26.4	11.5	42.9
Deaths under 1 month	13	3	10	13	3	10	15.6	6.9	25.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	2.8	4.5	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	2.8	4.5	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	0	2	2	0	2	5.7	-	15.4
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	1	0	1	2.8	-	7.7
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	42	30	12	47	35	12	134.3	159.1	92.3
Diabetes (260)	5	1	4	6	2	4	17.1	9.1	30.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	50	22	28	51	21	30	145.7	95.4	230.8
Heart diseases (410-443)	110	79	31	121	84	37	345.7	381.8	284.6
Arteriosclerosis (450)	10	9	1	10	9	1	28.6	40.9	7.7
Influenza (480-483)	1	0	1	1	0	1	2.8	-	7.7
#Pneumonia (490-493)	4	2	2	9	7	2	25.7	31.8	15.4
#Diarrhea & enteritis (571)	3	1	2	2	0	2	5.7	-	15.4
Acute nephritis (590, 591)	1	0	1	1	0	1	2.8	-	7.7
Chronic nephritis, etc. (592-594)	12	6	6	12	6	6	34.3	27.3	46.2
Maternal deaths (640-689)	1	0	1	1	0	1	1.2	-	2.5
Cong. malformations (750-759)	1	1	0	1	1	0	2.8	4.5	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	14	4	10	12	2	10	14.4	4.6	25.2
Violent & acc. deaths (800-999)	18	12	6	22	15	7	62.8	68.2	53.8
Acc. deaths, total (800-962)	15	9	6	18	12	6	51.4	54.5	46.2
Motor vehicle acc. (810-835)	6	5	1	5	4	1	14.3	18.2	7.7
Ill-defined (780-795)	20	6	14	22	7	15	62.8	31.8	115.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MONTGOMERY COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	333	127	206	334	123	211	25.7	15.4	42.2
Fetal deaths	11	1	10	12	3	9	35.9	24.4	42.6
Deaths from all causes	141	77	64	131	73	58	10.1	9.1	11.6
Deaths under 1 year	12	4	8	13	5	8	38.9	40.6	37.9
Deaths under 1 month	8	4	4	9	5	4	26.9	40.6	19.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	20	14	6	18	13	5	138.5	162.5	100.0
Diabetes (260)	1	1	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	25	11	14	24	11	13	184.6	137.5	260.0
Heart diseases (410-443)	47	31	16	41	25	16	315.4	312.5	320.0
Arteriosclerosis (450)	1	1	0	1	1	0	7.7	12.5	-
Influenza (480-483)	1	0	1	1	0	1	7.7	-	20.0
#Pneumonia (490-493)	4	2	2	5	3	2	38.5	37.5	40.0
#Diarrhea & enteritis (571)	1	0	1	1	0	1	7.7	-	20.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	3	2	1	23.1	25.0	20.0
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	7.7	12.5	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	8	4	4	8	4	4	24.0	32.5	19.0
Violent & acc. deaths (800-999)	19	10	9	13	8	5	100.0	100.0	100.0
Acc. deaths, total (800-962)	15	8	7	10	6	4	76.9	75.0	80.0
Motor vehicle acc. (810-835)	8	2	6	2	0	2	15.4	-	40.0
Ill-defined (780-795)	9	1	8	8	1	7	61.5	12.5	140.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
NESHOBIA COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	492	280	212	499	279	220	19.2	15.5	27.5
Fetal deaths	14	4	10	14	4	10	28.0	14.3	45.4
Deaths from all causes	155	108	47	205	142	63	7.9	7.9	7.9
Deaths under 1 year	19	7	12	23	9	14	46.1	32.2	63.6
Deaths under 1 month	9	5	4	10	5	5	20.0	17.9	22.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	2	1	3	2	1	11.5	11.1	12.5
Tbc., respiratory (001-008)	3	2	1	3	2	1	11.5	11.1	12.5
Syphilis (020-029)	1	0	1	1	0	1	3.8	-	12.5
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	1	0	1	1	0	1	3.8	-	12.5
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	22	17	5	33	25	8	126.9	138.9	100.0
Diabetes (260)	1	0	1	2	0	2	7.7	-	25.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	29	25	4	34	30	4	130.8	166.7	50.0
Heart diseases (410-443)	40	31	9	52	40	12	200.0	222.2	150.0
Arteriosclerosis (450)	2	1	1	2	1	1	7.7	5.6	12.5
Influenza (480-483)	1	0	1	1	0	1	3.8	-	12.5
#Pneumonia (490-493)	4	3	1	3	2	1	11.5	11.1	12.5
#Diarrhea & enteritis (571)	3	1	2	3	2	1	11.5	11.1	12.5
Acute nephritis (590, 591)	1	0	1	1	0	1	3.8	-	12.5
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	3.8	-	12.5
Maternal deaths (640-689)	0	0	0	1	0	1	2.0	-	4.5
Cong. malformations (750-759)	5	2	3	10	4	6	38.5	22.2	75.0
Birth inj., atelect. (760-762)	3	2	1	2	1	1	4.0	3.6	4.5
#Other dis. early inf. (763-776)	6	2	4	5	2	3	10.0	7.2	13.6
Violent & acc. deaths (800-999)	10	7	3	17	13	4	65.4	72.2	50.0
Acc. deaths, total (800-962)	9	6	3	13	12	1	50.0	66.7	12.5
Motor vehicle acc. (810-835)	6	4	2	6	6	0	23.1	33.3	-
Ill-defined (780-795)	11	5	6	13	7	6	50.0	38.9	75.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
NEWTON COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	376	257	119	435	220	215	20.7	15.7	30.7
Fetal deaths	37	19	18	33	17	16	75.9	77.3	74.4
Deaths from all causes	163	106	57	192	112	80	9.1	8.0	11.4
Deaths under 1 year	13	7	6	15	6	9	34.5	27.3	41.9
Deaths under 1 month	9	4	5	11	4	7	25.3	18.2	32.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	1	1	0	4.8	7.1	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	0	1	4.8	-	14.3
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	25	18	7	30	19	11	142.8	135.7	157.1
Diabetes (260)	2	0	2	3	0	3	14.3	-	42.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	30	21	9	31	20	11	147.6	142.8	157.1
Heart diseases (410-443)	49	37	12	57	39	18	271.4	278.6	257.1
Arteriosclerosis (450)	4	2	2	4	3	1	19.0	21.4	14.3
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	2	0	3	3	0	14.3	21.4	-
#Diarrhea & enteritis (571)	1	0	1	2	0	2	9.5	-	28.6
Acute nephritis (590, 591)	0	0	0	1	0	1	4.8	-	14.3
Chronic nephritis, etc. (592-594)	2	1	1	1	1	0	4.8	7.1	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	8	6	2	7	5	2	33.3	35.7	28.6
Birth inj., atelect. (760-762)	1	0	1	2	0	2	4.6	-	9.3
#Other dis. early inf. (763-776)	3	0	3	3	0	3	6.9	-	14.0
Violent & acc. deaths (800-999)	10	8	2	11	6	5	52.4	42.8	71.4
Acc. deaths, total (800-962)	6	6	0	6	4	2	28.6	28.6	28.6
Motor vehicle acc. (810-835)	5	5	0	3	2	1	14.3	14.3	14.3
Ill-defined (780-795)	14	2	12	13	1	12	61.9	7.1	171.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

**BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
NOXUBEE COUNTY, MISSISSIPPI, 1959**

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	554	96	458	540	90	450	38.6	18.0	50.0
Fetal deaths	30	2	28	28	2	26	51.8	22.2	57.8
Deaths from all causes	178	52	126	184	59	125	13.1	11.8	13.9
Deaths under 1 year	21	1	20	23	2	21	42.6	22.2	46.7
Deaths under 1 month	14	1	13	15	2	13	27.8	22.2	28.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	0	4	5	1	4	35.7	20.0	44.4
Tbc., respiratory (001-008)	4	0	4	5	1	4	35.7	20.0	44.4
Syphilis (020-029)	1	0	1	2	0	2	14.3	-	22.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	7.1	-	11.1
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	22	11	11	23	11	12	164.3	220.0	133.3
Diabetes (260)	1	0	1	1	0	1	7.1	-	11.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	32	8	24	31	12	19	221.4	240.0	211.1
Heart diseases (410-443)	55	23	32	50	21	29	357.1	420.0	322.2
Arteriosclerosis (450)	4	1	3	3	1	2	21.4	20.0	22.2
Influenza (480-483)	1	0	1	1	0	1	7.1	-	11.1
#Pneumonia (490-493)	3	0	3	5	2	3	35.7	40.0	33.3
#Diarrhea & enteritis (571)	1	0	1	1	0	1	7.1	-	11.1
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	2	0	2	2	0	2	3.7	-	4.4
Cong. malformations (750-759)	2	0	2	2	0	2	14.3	-	22.2
Birth inj., atelect. (760-762)	1	0	1	2	1	1	3.7	11.1	2.2
#Other dis. early inf. (763-776)	10	1	9	10	1	9	18.5	11.1	20.0
Violent & acc. deaths (800-999)	8	3	5	8	3	5	57.1	60.0	55.6
Acc. deaths, total (800-962)	8	3	5	7	3	4	50.0	60.0	44.4
Motor vehicle acc. (810-835)	2	1	1	1	1	0	7.1	20.0	-
Ill-defined (780-795)	22	1	21	23	1	22	164.3	20.0	244.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
OKTIBBEHA COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	822	347	475	786	328	458	28.1	20.5	38.2
Fetal deaths	23	4	19	19	4	15	24.2	12.2	32.8
Deaths from all causes	195	100	95	192	91	101	6.8	5.7	8.4
Deaths under 1 year	29	11	18	24	8	16	30.5	24.4	34.9
Deaths under 1 month	17	9	8	13	6	7	16.5	18.3	15.3
SELECTED CAUSES OF DEATH									
AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	3.6	6.2	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	3.6	6.2	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	21	11	10	20	9	11	71.4	56.2	91.7
Diabetes (260)	2	1	1	1	1	0	3.6	6.2	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	30	18	12	29	17	12	103.6	106.2	100.0
Heart diseases (410-443)	55	31	24	58	31	27	207.1	193.8	225.0
Arteriosclerosis (450)	4	2	2	3	1	2	10.7	6.2	16.7
Influenza (480-483)	2	0	2	2	0	2	7.1	-	16.7
#Pneumonia (490-493)	4	0	4	3	0	3	10.7	-	25.0
#Diarrhea & enteritis (571)	4	0	4	4	0	4	14.3	-	33.3
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	7.1	6.2	8.3
Maternal deaths (640-689)	2	0	2	2	0	2	2.5	-	4.4
Cong. malformations (750-759)	3	1	2	3	1	2	10.7	6.2	16.7
Birth inj., atelect. (760-762)	6	3	3	4	2	2	5.1	6.1	4.4
#Other dis. early inf. (763-776)	7	4	3	5	2	3	6.4	6.1	6.6
Violent & acc. deaths (800-999)	19	12	7	20	11	9	71.4	68.8	75.0
Acc. deaths, total (800-962)	14	9	5	14	8	6	50.0	50.0	50.0
Motor vehicle acc. (810-835)	5	2	3	5	2	3	17.8	12.5	25.0
Ill-defined (780-795)	15	3	12	15	3	12	53.6	18.8	100.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PANOLA COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	785	223	562	814	243	571	28.1	17.4	38.1
Fetal deaths	21	1	20	29	4	25	35.6	16.5	43.8
Deaths from all causes	265	110	155	300	125	175	10.3	8.9	11.7
Deaths under 1 year	28	6	22	30	7	23	36.8	28.8	40.3
Deaths under 1 month	12	4	8	14	5	9	17.2	20.6	15.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	2	0	2	6.9	-	13.3
Tbc., respiratory (001-008)	1	0	1	2	0	2	6.9	-	13.3
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	44	21	23	51	24	27	175.9	171.4	180.0
Diabetes (260)	2	0	2	2	0	2	6.9	-	13.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-384)	28	8	20	36	15	21	124.1	107.1	140.0
Heart diseases (410-443)	95	49	46	98	49	49	337.9	350.0	326.7
Arteriosclerosis (450)	3	1	2	3	1	2	10.3	7.1	13.3
Influenza (480-483)	2	0	2	2	0	2	6.9	-	13.3
#Pneumonia (490-493)	7	1	6	7	1	6	24.1	7.1	40.0
#Diarrhea & enteritis (571)	5	0	5	3	0	3	10.3	-	20.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	4	0	4	13.8	-	26.7
Maternal deaths (640-689)	0	0	0	1	1	0	1.2	4.1	-
Cong. malformations (750-759)	2	1	1	2	1	1	6.9	7.1	6.7
Birth inj., atelect. (760-762)	5	2	3	5	2	3	6.1	8.2	5.2
#Other dis. early inf. (763-776)	5	1	4	5	1	4	6.1	4.1	7.0
Violent & acc. deaths (800-999)	20	15	5	20	13	7	69.0	92.8	46.7
Acc. deaths, total (800-962)	20	15	5	19	13	6	65.5	92.8	40.0
Motor vehicle acc. (810-835)	11	8	3	9	5	4	31.0	35.7	26.7
111-defined (780-795)	29	5	24	33	8	25	113.8	57.1	166.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PEARL RIVER COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	656	480	176	534	356	178	18.4	17.8	19.8
Fetal deaths	19	12	7	17	10	7	31.8	28.1	39.3
Deaths from all causes	182	136	46	202	152	50	7.0	7.6	5.6
Deaths under 1 year	20	12	8	20	10	10	37.4	28.1	56.2
Deaths under 1 month	12	8	4	13	8	5	24.3	22.5	28.1
SELECTED CAUSES OF DEATH									
AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	3.4	5.0	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	3.4	5.0	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	23	20	3	25	22	3	86.2	110.0	33.3
Diabetes (260)	2	2	0	3	3	0	10.3	15.0	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	12	3	13	9	4	44.8	45.0	44.4
Heart diseases (410-443)	64	52	12	66	53	13	227.6	265.0	144.4
Arteriosclerosis (450)	4	3	1	3	2	1	10.3	10.0	11.1
Influenza (480-483)	1	1	0	1	1	0	3.4	5.0	-
#Pneumonia (490-493)	6	5	1	6	5	1	20.7	25.0	11.1
#Diarrhea & enteritis (571)	1	0	1	1	0	1	3.4	-	11.1
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	3.4	5.0	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	4	1	3	4	1	3	13.8	5.0	33.3
Birth inj., atelect. (760-762)	5	3	2	5	3	2	9.4	8.4	11.2
#Other dis. early inf. (763-776)	6	6	0	5	4	1	9.4	11.2	5.6
Violent & acc. deaths (800-999)	14	9	5	24	20	4	82.8	100.0	44.4
Acc. deaths, total (800-962)	8	7	1	20	19	1	69.0	95.0	11.1
Motor vehicle acc. (810-835)	6	5	1	9	8	1	31.0	40.0	11.1
Ill-defined (780-795)	21	9	12	22	10	12	75.9	50.0	133.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PERRY COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	101	76	25	232	169	63	25.8	24.1	31.5
Fetal deaths	5	3	2	13	8	5	56.0	47.3	79.4
Deaths from all causes	44	26	18	85	58	27	9.4	8.3	13.5
Deaths under 1 year	4	1	3	5	3	2	21.6	17.8	31.7
Deaths under 1 month	1	0	1	2	2	0	8.6	11.8	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	2	0	2	22.2	-	100.0
Tbc., respiratory (001-008)	1	0	1	2	0	2	22.2	-	100.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	8	5	3	15	11	4	166.7	157.1	200.0
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	5	2	3	10	5	5	111.1	71.4	250.0
Heart diseases (410-443)	14	11	3	22	18	4	244.4	257.1	200.0
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	0	1	3	2	1	33.3	28.6	50.0
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	11.1	14.3	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	11.1	14.3	-
Birth inj., atelect. (760-762)	0	0	0	1	1	0	4.3	5.9	-
#Other dis. early inf. (763-776)	1	0	1	1	1	0	4.3	5.9	-
Violent & acc. deaths (800-999)	8	5	3	15	10	5	166.7	142.8	250.0
Acc. deaths, total (800-962)	6	4	2	13	9	4	144.4	128.6	200.0
Motor vehicle acc. (810-835)	2	1	1	6	5	1	66.7	71.4	50.0
Ill-defined (780-795)	5	3	2	5	3	2	55.6	42.8	100.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PIKE COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence				Rate*		
	Total	White	Non-white	Number		Rate*		Total	White	Non-white
				Total	White	Non-white	Total			
Live births	990	446	544	875	332	543	27.3	18.4	38.8	
Fetal deaths	30	8	22	27	5	22	30.8	15.1	40.5	
Deaths from all causes	371	221	150	373	212	161	11.6	11.8	11.5	
Deaths under 1 year	33	8	25	31	7	24	35.4	21.1	44.2	
Deaths under 1 month	15	5	10	15	4	11	17.1	12.0	20.2	
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.										
Tbc., all forms (001-019)	4	1	3	5	1	4	15.6	5.6	28.6	
Tbc., respiratory (001-008)	4	1	3	5	1	4	15.6	5.6	28.6	
Syphilis (020-029)	0	0	0	0	0	0	-	-	-	
Typhoid fever (040)	0	0	0	0	0	0	-	-	-	
Dysentery (045-048)	0	0	0	0	0	0	-	-	-	
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-	
Diphtheria (055)	0	0	0	0	0	0	-	-	-	
Whooping cough (056)	2	0	2	1	0	1	3.1	-	7.1	
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-	
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-	
Smallpox (084)	0	0	0	0	0	0	-	-	-	
Measles (085)	0	0	0	0	0	0	-	-	-	
Malaria (110-117)	0	0	0	0	0	0	-	-	-	
Cancer, incl. leukemia (140-205)	53	39	14	58	45	13	181.2	250.0	92.8	
Diabetes (260)	4	3	1	3	2	1	9.4	11.1	7.1	
Pellagra (281)	0	0	0	0	0	0	-	-	-	
Intracranial lesions (330-334)	51	31	20	53	30	23	165.6	166.7	164.3	
Heart diseases (410-443)	110	68	42	115	66	49	359.4	366.7	350.0	
Arteriosclerosis (450)	11	7	4	9	6	3	28.1	33.3	21.4	
Influenza (480-483)	0	0	0	0	0	0	-	-	-	
#Pneumonia (490-493)	14	7	7	12	5	7	37.5	27.8	50.0	
#Diarrhea & enteritis (571)	2	1	1	2	1	1	6.2	5.6	7.1	
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-	
Chronic nephritis, etc. (592-594)	3	0	3	3	0	3	9.4	-	21.4	
Maternal deaths (640-689)	1	1	0	0	0	0	-	-	-	
Cong. malformations (750-759)	5	1	4	5	1	4	15.6	5.6	28.6	
Birth inj., atelect. (760-762)	2	2	0	1	1	0	1.1	3.0	-	
#Other dis. early inf. (763-776)	10	3	7	11	3	8	12.6	9.0	14.7	
Violent & acc. deaths (800-999)	32	19	13	25	13	12	78.1	72.2	85.7	
Acc. deaths, total (800-962)	26	17	9	20	11	9	62.5	61.1	64.3	
Motor vehicle acc. (810-835)	14	11	3	7	4	3	21.9	22.2	21.4	
Ill-defined (780-795)	32	11	21	28	8	20	87.5	44.4	142.8	

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PONTOTOC COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	294	218	76	288	209	79	13.1	11.6	19.8
Fetal deaths	12	9	3	13	10	3	45.1	47.8	38.0
Deaths from all causes	169	141	28	178	148	30	8.1	8.2	7.5
Deaths under 1 year	10	4	6	9	4	5	31.2	19.1	63.3
Deaths under 1 month	8	3	5	7	3	4	24.3	14.4	50.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	4.5	-	25.0
Tbc., respiratory (001-008)	1	0	1	1	0	1	4.5	-	25.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	1	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	22	18	4	25	21	4	113.6	116.7	100.0
Diabetes (260)	2	2	0	1	1	0	4.5	5.6	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	20	14	6	18	12	6	81.8	66.7	150.0
Heart diseases (410-443)	69	63	6	73	65	8	331.8	361.1	200.0
Arteriosclerosis (450)	5	5	0	3	3	0	13.6	16.7	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	9	8	1	9	8	1	40.9	44.4	25.0
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590; 591)	1	1	0	1	1	0	4.5	5.6	-
Chronic nephritis, etc. (592-594)	2	1	1	1	0	1	4.5	-	25.0
Maternal deaths (640-689)	0	0	0	1	1	0	3.5	4.8	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	3	1	2	3	1	2	10.4	4.8	25.3
#Other dis. early inf. (763-776)	6	3	3	5	3	2	17.4	14.4	25.3
Violent & acc. deaths (800-999)	12	9	3	19	15	4	86.4	83.3	100.0
Acc. deaths, total (800-962)	9	6	3	16	12	4	72.7	66.7	100.0
Motor vehicle acc. (810-835)	7	6	1	11	9	2	50.0	50.0	50.0
Ill-defined (780-795)	1	1	0	2	2	0	9.1	11.1	-

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PRENTISS COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	455	371	84	379	301	78	22.3	23.2	19.5
Fetal deaths	6	6	0	5	5	0	13.2	16.6	-
Deaths from all causes	199	185	14	175	161	14	10.3	12.4	3.5
Deaths under 1 year	18	14	4	19	15	4	50.1	49.8	51.3
Deaths under 1 month	14	12	2	15	13	2	39.6	43.2	25.6
SELECTED CAUSES OF DEATH									
AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	5.9	7.7	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	5.9	7.7	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	1	1	0	5.9	7.7	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	22	21	1	20	19	1	117.6	146.2	25.0
Diabetes (260)	8	8	0	6	6	0	35.3	46.2	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	33	30	3	24	21	3	141.2	161.5	75.0
Heart diseases (410-443)	54	52	2	46	45	1	270.6	346.2	25.0
Arteriosclerosis (450)	6	5	1	6	5	1	35.3	38.5	25.0
Influenza (480-483)	1	1	0	1	1	0	5.9	7.7	-
#Pneumonia (490-493)	9	7	2	7	4	3	41.2	30.8	75.0
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	4	0	4	4	0	23.5	30.8	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	6	5	1	6	5	1	15.8	16.6	12.8
#Other dis. early inf. (763-776)	7	6	1	8	7	1	21.1	23.2	12.8
Violent & acc. deaths (800-999)	13	12	1	15	14	1	88.2	107.7	25.0
Acc. deaths, total (800-962)	12	11	1	14	13	1	82.4	100.0	25.0
Motor vehicle acc. (810-835)	6	6	0	5	5	0	29.4	38.5	-
Ill-defined (780-795)	13	12	1	13	12	1	76.5	92.3	25.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
QUITMAN COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	734	210	524	661	168	493	33.0	24.0	37.9
Fetal deaths	18	3	15	19	3	16	28.7	17.8	32.4
Deaths from all causes	177	44	133	215	65	150	10.8	9.3	11.5
Deaths under 1 year	26	4	22	28	6	22	42.4	35.7	44.6
Deaths under 1 month	13	2	11	15	4	11	22.7	23.8	22.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	3	1	2	15.0	14.3	15.4
Tbc., respiratory (001-008)	2	1	1	3	1	2	15.0	14.3	15.4
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	5.0	-	7.7
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	28	11	17	38	15	23	190.0	214.3	176.9
Diabetes (260)	1	0	1	2	0	2	10.0	-	15.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	5	10	20	8	12	100.0	114.3	92.3
Heart diseases (410-443)	49	14	35	55	16	39	275.0	228.6	300.0
Arteriosclerosis (450)	1	0	1	1	0	1	5.0	-	7.7
Influenza (480-483)	1	0	1	2	0	2	10.0	-	15.4
#Pneumonia (490-493)	11	0	11	11	1	10	55.0	14.3	76.9
#Diarrhea & enteritis (571)	4	0	4	3	0	3	15.0	-	23.1
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	0	2	2	0	2	10.0	-	15.4
Maternal deaths (640-689)	0	0	0	1	1	0	1.5	6.0	-
Cong. malformations (750-759)	1	0	1	2	1	1	10.0	14.3	7.7
Birth inj., atelect. (760-762)	1	0	1	1	0	1	1.5	-	2.0
#Other dis. early inf. (763-776)	9	2	7	9	2	7	13.6	11.9	14.2
Violent & acc. deaths (800-999)	17	6	11	21	7	14	105.0	100.0	107.7
Acc. deaths, total (800-962)	15	5	10	19	6	13	95.0	85.7	100.0
Motor vehicle acc. (810-835)	7	4	3	8	4	4	40.0	57.1	30.8
Ill-defined (780-795)	17	2	15	16	3	13	80.0	42.8	100.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
RANKIN COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	305	77	228	757	429	328	17.6	15.3	21.9
Fetal deaths	19	5	14	46	20	26	60.8	46.6	79.3
Deaths from all causes	378	209	169	234	143	91	5.4	5.1	6.1
Deaths under 1 year	8	3	5	20	8	12	26.4	18.6	36.6
Deaths under 1 month	3	0	3	15	6	9	19.8	14.0	27.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	6	5	1	1	1	0	2.3	3.6	-
Tbc., respiratory (001-008)	5	4	1	1	1	0	2.3	3.6	-
Syphilis (020-029)	2	1	1	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	3	0	3	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	32	17	15	35	28	7	81.4	100.0	46.7
Diabetes (260)	3	1	2	3	2	1	7.0	7.1	6.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	84	55	29	27	10	17	62.8	35.7	113.3
Heart diseases (410-443)	94	54	40	71	49	22	165.1	175.0	146.7
Arteriosclerosis (450)	7	3	4	7	4	3	16.3	14.3	20.0
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	12	7	5	4	2	2	9.3	7.1	13.3
#Diarrhea & enteritis (571)	1	0	1	2	0	2	4.6	-	13.3
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	1	2	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	3	2	1	7.0	7.1	6.7
Birth inj., atelect. (760-762)	0	0	0	3	3	0	4.0	7.0	-
#Other dis. early inf. (763-776)	3	0	3	9	2	7	11.9	4.7	21.3
Violent & acc. deaths (800-999)	29	19	10	24	17	7	55.8	60.7	46.7
Acc. deaths, total (800-962)	25	17	8	21	15	6	48.8	53.6	40.0
Motor vehicle acc. (810-835)	6	5	1	8	5	3	18.6	17.8	20.0
Ill-defined (780-795)	31	6	25	16	5	11	37.2	17.8	73.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SCOTT COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	634	360	274	578	293	285	26.3	20.9	35.6
Fetal deaths	10	1	9	19	5	14	32.9	17.1	49.1
Deaths from all causes	174	93	81	200	108	92	9.1	7.7	11.5
Deaths under 1 year	25	7	18	30	11	19	51.9	37.5	66.7
Deaths under 1 month	18	6	12	21	9	12	36.3	30.7	42.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	2	1	1	9.1	7.1	12.5
Tbc., respiratory (001-008)	2	1	1	2	1	1	9.1	7.1	12.5
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	23	15	8	20	12	8	90.9	85.7	100.0
Diabetes (260)	1	0	1	1	0	1	4.5	-	12.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	34	19	15	39	22	17	177.3	157.1	212.5
Heart diseases (410-443)	46	29	17	50	33	17	227.3	235.7	212.5
Arteriosclerosis (450)	4	3	1	4	3	1	18.2	21.4	12.5
Influenza (480-483)	3	0	3	3	0	3	13.6	-	37.5
#Pneumonia (490-493)	4	1	3	4	2	2	18.2	14.3	25.0
#Diarrhea & enteritis (571)	0	0	0	1	0	1	4.5	-	12.5
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	2	0	2	2	0	9.1	14.3	-
Maternal deaths (640-689)	1	1	0	1	1	0	1.7	3.4	-
Cong. malformations (750-759)	2	1	1	3	2	1	13.6	14.3	12.5
Birth inj., atelect. (760-762)	3	2	1	4	3	1	6.9	10.2	3.5
#Other dis. early inf. (763-776)	10	4	6	9	3	6	15.6	10.2	21.0
Violent & acc. deaths (800-999)	7	4	3	12	6	6	54.5	42.8	75.0
Acc. deaths, total (800-962)	6	4	2	10	5	5	45.4	35.7	62.5
Motor vehicle acc. (810-835)	2	1	1	2	1	1	9.1	7.1	12.5
Ill-defined (780-795)	21	5	16	22	5	17	100.0	35.7	212.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SHARKEY COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	249	46	203	340	62	278	37.8	15.5	55.6
Fetal deaths	10	1	9	15	2	13	44.1	32.2	46.8
Deaths from all causes	99	28	71	119	31	88	13.2	7.8	17.6
Deaths under 1 year	21	0	21	22	1	21	64.7	16.1	75.5
Deaths under 1 month	9	0	9	9	1	8	26.5	16.1	28.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	2	2	4	2	2	44.4	50.0	40.0
Tbc., respiratory (001-008)	4	2	2	4	2	2	44.4	50.0	40.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	4	1	3	7	2	5	77.8	50.0	100.0
Diabetes (260)	5	3	2	3	2	1	33.3	50.0	20.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	9	1	8	14	2	12	155.6	50.0	240.0
Heart diseases (410-443)	29	15	14	28	12	16	311.1	300.0	320.0
Arteriosclerosis (450)	1	1	0	2	2	0	22.2	50.0	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	0	1	4	1	3	44.4	25.0	60.0
#Diarrhea & enteritis (571)	3	0	3	3	0	3	33.3	-	60.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	0	2	3	0	3	33.3	-	60.0
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	0	0	0	-	-	-
Birth inj., atelect. (760-762)	3	0	3	4	1	3	11.8	16.1	10.8
#Other dis. early inf. (763-776)	4	0	4	3	0	3	8.8	-	10.8
Violent & acc. deaths (800-999)	12	3	9	16	3	13	177.8	75.0	260.0
Acc. deaths, total (800-962)	11	2	9	15	2	13	166.7	50.0	260.0
Motor vehicle acc. (810-835)	3	2	1	2	1	1	22.2	25.0	20.0
Ill-defined (780-795)	15	1	14	15	1	14	166.7	25.0	280.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SIMPSON COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	456	234	222	461	260	201	17.1	15.3	20.1
Fetal deaths	9	7	2	10	8	2	21.7	30.8	10.0
Deaths from all causes	218	136	82	194	127	67	7.2	7.5	6.7
Deaths under 1 year	11	6	5	13	6	7	28.2	23.1	34.8
Deaths under 1 month	10	6	4	11	5	6	23.9	19.2	29.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	34	15	19	1	0	1	3.7	-	10.0
Tbc., respiratory (001-008)	33	15	18	1	0	1	3.7	-	10.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	25	16	9	25	16	9	92.6	94.1	90.0
Diabetes (260)	3	3	0	4	3	1	14.8	17.6	10.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	26	13	13	26	13	13	96.3	76.5	130.0
Heart diseases (410-443)	66	50	16	57	48	9	211.1	282.4	90.0
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	7	2	5	9	2	7	33.3	11.8	70.0
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	2	2	0	7.4	11.8	-
Birth inj., atelect. (760-762)	5	4	1	6	4	2	13.0	15.4	10.0
#Other dis. early inf. (763-776)	5	2	3	5	1	4	10.8	3.8	19.9
Violent & acc. deaths (800-999)	15	10	5	19	12	7	70.4	70.6	70.0
Acc. deaths, total (800-962)	12	7	5	14	9	5	51.8	52.9	50.0
Motor vehicle acc. (810-835)	5	4	1	6	6	0	22.2	35.3	-
Ill-defined (780-795)	15	8	7	18	10	8	66.7	58.8	80.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SMITH COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	86	54	32	334	189	145	19.6	15.8	29.0
Fetal deaths	2	1	1	8	5	3	24.0	26.4	20.7
Deaths from all causes	75	57	18	128	100	28	7.5	8.3	5.6
Deaths under 1 year	2	0	2	14	3	11	41.9	15.9	75.9
Deaths under 1 month	0	0	0	7	2	5	21.0	10.6	34.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	5.9	8.3	-
Tbc., respiratory (001-110)	1	1	0	1	1	0	5.9	8.3	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	1	1	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	15	12	3	24	21	3	141.2	175.0	60.0
Diabetes (260)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330334)	6	5	1	16	14	2	94.1	116.7	40.0
Heart diseases (410-443)	21	18	3	31	27	4	182.4	225.0	80.0
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	2	2	0	2	2	0	11.8	16.7	-
#Pneumonia (490-493)	2	1	1	4	3	1	23.5	25.0	20.0
#Diarrhea & enteritis (571)	0	0	0	2	0	2	11.8	-	40.0
Acute nephritis (590, 591)	2	0	2	2	0	2	11.8	-	40.0
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	5.9	8.3	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	2	0	2	11.8	-	40.0
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	0	0	0	5	2	3	15.0	10.6	20.7
Violent & acc. deaths (800-999)	11	7	4	17	12	5	100.0	100.0	100.0
Acc. deaths, total (800-962)	9	5	4	14	9	5	82.4	75.0	100.0
Motor vehicle acc. (810-835)	4	3	1	6	5	1	35.3	41.7	20.0
Ill-defined (780-795)	5	3	2	5	3	2	29.4	25.0	40.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
STONE COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	160	112	48	153	95	58	19.1	14.6	38.7
Fetal deaths	13	8	5	12	7	5	78.4	73.7	86.2
Deaths from all causes	62	39	23	58	34	24	7.2	5.2	16.0
Deaths under 1 year	9	6	3	8	4	4	52.3	42.1	69.0
Deaths under 1 month	6	5	1	6	4	2	39.2	42.1	34.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	2	1	1	25.0	15.4	66.7
Tbc., respiratory (001-008)	1	0	1	2	1	1	25.0	15.4	66.7
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	12.5	-	66.7
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	12	10	2	13	9	4	162.5	138.5	266.7
Diabetes (260)	0	0	0	1	1	0	12.5	15.4	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	7	4	3	5	3	2	62.5	46.2	133.3
Heart diseases (410-443)	14	9	5	12	8	4	150.0	123.1	266.7
Arteriosclerosis (450)	0	0	0	1	1	0	12.5	15.4	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	0	0	0	2	2	0	25.0	30.8	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	12.5	15.4	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	3	3	0	2	2	0	13.1	21.0	-
#Other dis. early inf. (763-776)	2	2	0	3	2	1	19.6	21.0	17.2
Violent & acc. deaths (800-999)	10	8	2	4	1	3	50.0	15.4	200.0
Acc. deaths, total (800-962)	9	8	1	3	1	2	37.5	15.4	133.3
Motor vehicle acc. (810-835)	5	5	0	2	1	1	25.0	15.4	66.7
III-defined (780-795)	8	1	7	8	1	7	100.0	15.4	466.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SUNFLOWER COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,357	251	1,106	1,444	277	1,167	32.8	16.3	43.2
Fetal deaths	60	11	49	65	11	54	45.0	39.7	46.3
Deaths from all causes	356	83	273	402	99	303	9.1	5.8	11.2
Deaths under 1 year	62	3	59	68	5	63	47.1	18.0	54.0
Deaths under 1 month	34	2	32	34	3	31	23.5	10.8	26.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	0	4	4	1	3	9.1	5.9	11.1
Tbc., respiratory (001-008)	4	0	4	4	1	3	9.1	5.9	11.1
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	2.3	-	3.7
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	34	9	25	38	13	25	86.4	76.5	92.6
Diabetes (260)	4	0	4	5	0	5	11.4	-	18.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	50	9	41	53	11	42	120.4	64.7	155.6
Heart diseases (410-443)	91	32	59	105	36	69	238.6	211.8	255.6
Arteriosclerosis (450)	2	1	1	3	2	1	6.8	11.8	3.7
Influenza (480-483)	2	0	2	3	0	3	6.8	-	11.1
#Pneumonia (490-493)	16	1	15	19	4	15	43.2	23.5	55.6
#Diarrhea & enteritis (571)	8	1	7	10	1	9	22.7	5.9	33.3
Acute nephritis (590, 591)	1	0	1	1	0	1	2.3	-	3.7
Chronic nephritis, etc. (592-594)	4	2	2	3	2	1	6.8	11.8	3.7
Maternal deaths (640-689)	2	0	2	2	0	2	1.4	-	1.7
Cong. malformations (750-759)	2	1	1	5	2	3	11.4	11.8	11.1
Birth inj., atelect. (760-762)	7	1	6	8	1	7	5.5	3.6	6.0
#Other dis. early inf. (763-776)	27	1	26	25	2	23	17.3	7.2	19.7
Violent & acc. deaths (800-999)	34	14	20	42	14	28	95.4	82.4	103.7
Acc. deaths, total (800-962)	28	11	17	36	12	24	81.8	70.6	88.9
Motor vehicle acc. (810-835)	8	3	5	9	3	6	20.4	17.6	22.2
Ill-defined (780-795)	40	2	38	40	1	39	90.9	5.9	144.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TALLAHATCHIE COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	730	131	599	773	175	598	35.1	17.5	49.8
Fetal deaths	37	11	26	38	11	27	49.2	62.8	45.2
Deaths from all causes	237	77	160	284	103	181	12.9	10.3	15.1
Deaths under 1 year	42	6	36	46	7	39	59.5	40.0	65.2
Deaths under 1 month	20	4	16	21	5	16	27.2	28.6	26.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	3	1	2	13.6	10.0	16.7
Tbc., respiratory (001-008)	0	0	0	2	1	1	9.1	10.0	8.3
Syphilis (020-029)	2	0	2	2	0	2	9.1	-	16.7
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	4.5	-	8.3
Scarlet fever & streptococcal sore throat (050, 051)	1	1	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	4	0	4	4	0	4	18.2	-	33.3
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	4.5	-	8.3
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	21	9	12	36	19	17	163.6	190.0	141.7
Diabetes (260)	2	0	2	3	1	2	13.6	10.0	16.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	27	14	13	30	17	13	136.4	170.0	108.3
Heart diseases (410-443)	70	27	43	74	29	45	336.4	290.0	375.0
Arteriosclerosis (450)	2	1	1	3	2	1	13.6	20.0	8.3
Influenza (480-483)	1	0	1	1	0	1	4.5	-	8.3
#Pneumonia (490-493)	5	0	5	6	0	6	27.3	-	50.0
#Diarrhea & enteritis (571)	3	0	3	7	0	7	31.8	-	58.3
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	1	3	6	1	5	27.3	10.0	41.7
Maternal deaths (640-689)	3	0	3	2	0	2	2.6	-	3.3
Cong. malformations (750-759)	3	1	2	3	1	2	13.6	10.0	16.7
Birth inj., atelect. (760-762)	3	1	2	3	1	2	3.9	5.7	3.3
#Other dis. early inf. (763-776)	12	2	10	13	3	10	16.8	17.1	16.7
Violent & acc. deaths (800-999)	18	10	8	24	12	12	109.1	120.0	100.0
Acc. deaths, total (800-962)	15	9	6	19	11	8	86.4	110.0	66.7
Motor vehicle acc. (810-835)	7	5	2	9	6	3	40.9	60.0	25.0
Ill-defined (780-795)	31	3	28	31	3	28	140.9	30.0	233.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

**BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TATE COUNTY, MISSISSIPPI, 1959**

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	450	29	421	580	126	454	32.2	14.0	50.4
Fetal deaths	10	1	9	17	2	15	29.3	15.9	33.0
Deaths from all causes	110	40	70	163	75	88	9.0	8.3	9.8
Deaths under 1 year	8	0	8	18	5	13	31.0	39.7	28.6
Deaths under 1 month	4	0	4	13	5	8	22.4	39.7	17.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	2	0	2	2	0	11.1	22.2	-
Tbc., respiratory (001-008)	2	2	0	2	2	0	11.1	22.2	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	15	5	10	24	13	11	133.3	144.4	122.2
Diabetes (260)	2	0	2	3	1	2	16.7	11.1	22.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	11	3	8	16	7	9	88.9	77.8	100.0
Heart diseases (410-443)	44	19	25	55	27	28	305.6	300.0	311.1
Arteriosclerosis (450)	1	0	1	1	0	1	5.6	-	11.1
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	0	2	5	1	4	27.8	11.1	44.4
#Diarrhea & enteritis (571)	2	0	2	2	0	2	11.1	-	22.2
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	0	3	4	0	4	22.2	-	44.4
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	1	1	0	5.6	11.1	-
Birth inj., atelect. (760-762)	0	0	0	3	3	0	5.2	23.8	-
#Other dis. early inf. (763-776)	2	0	2	8	2	6	13.8	15.9	13.2
Violent & acc. deaths (800-999)	9	7	2	11	8	3	61.1	88.9	33.3
Acc. deaths, total (800-962)	7	6	1	9	7	2	50.0	77.8	22.2
Motor vehicle acc. (810-835)	2	2	0	6	5	1	33.3	55.6	11.1
Ill-defined (780-795)	12	1	11	15	3	12	83.3	33.3	133.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TIPPAH COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	398	294	104	334	245	89	15.9	16.3	14.8
Fetal deaths	10	6	4	13	10	3	38.9	40.8	33.7
Deaths from all causes	117	96	21	144	119	25	6.8	7.9	4.2
Deaths under 1 year	6	1	5	9	5	4	26.9	20.4	44.9
Deaths under 1 month	3	1	2	3	2	1	9.0	8.2	11.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	16	1	19	18	1	90.5	120.0	16.7
Diabetes (260)	1	1	0	3	3	0	14.3	20.0	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	20	14	6	24	17	7	114.3	113.3	116.7
Heart diseases (410-443)	48	45	3	51	48	3	242.8	320.0	50.0
Arteriosclerosis (450)	0	0	0	1	1	0	4.8	6.7	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	3	1	2	5	3	2	23.8	20.0	33.3
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	1	0	1	0	0	0	-	-	-
Cong. malformations (750-759)	2	1	1	2	1	1	9.5	6.7	16.7
Birth inj., atelect. (760-762)	2	1	1	1	1	0	3.0	4.1	-
#Other dis. early inf. (763-776)	1	0	1	2	1	1	6.0	4.1	11.2
Violent & acc. deaths (800-999)	10	8	2	15	9	6	71.4	60.0	100.0
Acc. deaths, total (800-962)	7	6	1	10	7	3	47.6	46.7	50.0
Motor vehicle acc. (810-835)	3	2	1	5	3	2	23.8	20.0	33.3
Ill-defined (780-795)	5	2	3	5	2	3	23.8	13.3	50.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TISHOMINGO COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	110	98	12	252	236	16	16.8	16.3	32.0
Fetal deaths	3	3	0	10	10	0	39.7	42.4	-
Deaths from all causes	87	79	8	128	116	12	8.5	8.0	24.0
Deaths under 1 year	6	5	1	11	9	2	43.6	38.1	125.0
Deaths under 1 month	3	2	1	5	4	1	19.8	16.9	62.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	2	0	2	2	0	13.3	13.8	-
Tbc., respiratory (001-008)	2	2	0	2	2	0	13.3	13.8	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	6	6	0	9	8	1	60.0	55.2	200.0
Diabetes (260)	1	1	0	1	1	0	6.7	6.9	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	23	21	2	36	33	3	240.0	227.6	600.0
Heart diseases (410-443)	30	29	1	34	32	2	226.7	220.7	400.0
Arteriosclerosis (450)	4	4	0	5	5	0	33.3	34.5	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	3	3	0	3	3	0	20.0	20.7	-
#Diarrhea & enteritis (571)	0	0	0	1	0	1	6.7	-	200.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	0	2	4	2	2	26.7	13.8	400.0
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	4	3	1	5	4	1	19.8	16.9	62.5
#Other dis. early inf. (763-776)	0	0	0	1	1	0	4.0	4.2	-
Violent & acc. deaths (800-999)	3	2	1	6	6	0	40.0	41.4	-
Acc. deaths, total (800-962)	3	2	1	6	6	0	40.0	41.4	-
Motor vehicle acc. (810-835)	1	0	1	3	3	0	20.0	20.7	-
Ill-defined (780-795)	2	1	1	4	3	1	26.7	20.7	200.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TUNICA COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	672	68	604	658	67	591	41.1	22.3	45.5
Fetal deaths	25	3	22	25	4	21	38.0	59.7	35.5
Deaths from all causes	199	25	174	206	21	185	12.9	7.0	14.2
Deaths under 1 year	30	1	29	30	1	29	45.6	14.9	49.1
Deaths under 1 month	19	1	18	18	1	17	27.4	14.9	28.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	4	1	3	25.0	33.3	23.1
Tbc., respiratory (001-008)	2	0	2	4	1	3	25.0	33.3	23.1
Syphilis (020-029)	0	0	0	1	0	1	6.2	-	7.7
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	16	3	13	24	3	21	150.0	100.0	161.5
Diabetes (260)	1	1	0	2	2	0	12.5	66.7	-
Pellagra (281)	1	0	1	1	0	1	6.2	-	7.7
Intracranial lesions (330-334)	11	0	11	13	0	13	81.2	-	100.0
Heart diseases (410-443)	75	11	64	73	10	63	456.2	333.3	484.6
Arteriosclerosis (450)	1	0	1	1	0	1	6.2	-	7.7
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	6	1	5	6	0	6	37.5	-	46.2
#Diarrhea & enteritis (571)	0	0	0	1	0	1	6.2	-	7.7
Acute nephritis (590, 591)	3	0	3	3	0	3	18.8	-	23.1
Chronic nephritis, etc. (592-594)	8	2	6	8	2	6	50.0	66.7	46.2
Maternal deaths (640-689)	1	0	1	2	0	2	3.0	-	3.4
Cong. malformations (750-759)	1	0	1	1	0	1	6.2	-	7.7
Birth inj., atelect. (760-762)	5	0	5	5	0	5	7.6	-	8.5
#Other dis. early inf. (763-776)	13	1	12	12	1	11	18.2	14.9	18.6
Violent & acc. deaths (800-999)	24	5	19	15	0	15	93.8	-	115.4
Acc. deaths, total (800-962)	24	5	19	15	0	15	93.8	-	115.4
Motor vehicle acc. (810-835)	12	4	8	3	0	3	18.8	-	23.1
Ill-defined (780-795)	19	1	18	20	1	19	125.0	33.3	146.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
UNION COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	434	335	99	405	304	101	16.9	16.9	16.8
Fetal deaths	50	38	12	35	25	10	86.4	82.2	99.0
Deaths from all causes	191	135	56	198	148	50	8.2	8.2	8.3
Deaths under 1 year	19	10	9	19	12	7	46.9	39.5	69.3
Deaths under 1 month	12	7	5	14	10	4	34.6	33.0	39.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	2	2	0	8.3	11.1	-
Tbc., respiratory (001-008)	0	0	0	2	2	0	8.3	11.1	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	18	14	4	22	18	4	91.7	100.0	66.7
Diabetes (260)	2	2	0	3	3	0	12.5	16.7	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	22	18	4	21	18	3	87.5	100.0	50.0
Heart diseases (410-443)	78	61	17	80	62	18	333.3	344.4	300.0
Arteriosclerosis (450)	5	2	3	6	3	3	25.0	16.7	50.0
Influenza (480-483)	1	0	1	1	0	1	4.2	-	16.7
#Pneumonia (490-493)	1	0	1	2	1	1	8.3	5.6	16.7
#Diarrhea & enteritis (571)	2	0	2	1	0	1	4.2	-	16.7
Acute nephritis (590, 591)	1	1	0	1	1	0	4.2	5.6	-
Chronic nephritis, etc. (592-594)	2	1	1	3	2	1	12.5	11.1	16.7
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	4.2	5.6	-
Birth inj., atelect. (760-762)	5	4	1	6	5	1	14.8	16.4	9.9
#Other dis. early inf. (763-776)	6	3	3	6	4	2	14.8	13.2	19.8
Violent & acc. deaths (800-999)	21	12	9	15	10	5	62.5	55.6	83.3
Acc. deaths, total (800-962)	20	12	8	15	10	5	62.5	55.6	83.3
Motor vehicle acc. (810-835)	11	10	1	9	8	1	37.5	44.4	16.7
Ill-defined (780-795)	15	8	7	16	9	7	66.7	50.0	116.7

* Crude birth and death rate per 1,000 population. Rate for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WALTHALL COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	604	320	284	362	138	224	25.8	17.2	37.3
Fetal deaths	26	9	17	15	3	12	41.4	21.7	53.6
Deaths from all causes	132	74	58	130	73	57	9.3	9.1	9.5
Deaths under 1 year	17	1	16	18	2	16	49.7	14.5	71.4
Deaths under 1 month	10	1	9	12	2	10	33.1	14.5	44.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	7.1	12.5	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	7.1	12.5	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	7.1	-	16.7
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	20	13	7	14	7	7	100.0	87.5	116.7
Diabetes (260)	3	2	1	3	2	1	21.4	25.0	16.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	21	13	8	23	15	8	164.3	187.5	133.3
Heart diseases (410-443)	45	32	13	41	30	11	292.8	375.0	183.3
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	3	0	3	3	0	3	21.4	-	50.0
#Diarrhea & enteritis (571)	2	0	2	3	1	2	21.4	12.5	33.3
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	14.3	12.5	16.7
Maternal deaths (640-689)	1	0	1	1	0	1	2.8	-	4.5
Cong. malformations (750-759)	0	0	0	1	1	0	7.1	12.5	-
Birth inj., atelect. (760-762)	6	1	5	6	1	5	16.6	7.2	22.3
#Other dis. early inf. (763-776)	4	0	4	5	0	5	13.8	-	22.3
Violent & acc. deaths (800-999)	5	2	3	6	3	3	42.8	37.5	50.0
Acc. deaths, total (800-962)	5	2	3	6	3	3	42.8	37.5	50.0
Motor vehicle acc. (810-835)	2	2	0	2	2	0	14.3	25.0	-
Ill-defined (780-795)	9	1	8	9	1	8	64.3	12.5	133.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WARREN COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,496	616	880	1,192	533	659	33.1	29.6	36.6
Fetal deaths	85	12	73	48	8	40	40.3	15.0	60.7
Deaths from all causes	677	250	427	495	218	277	13.8	12.1	15.4
Deaths under 1 year	63	8	55	42	8	34	35.2	15.0	51.6
Deaths under 1 month	43	6	37	27	6	21	22.6	11.2	31.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	1	2	3	2	1	8.3	11.1	5.6
Tbc., respiratory (001-008)	2	0	2	2	1	1	5.6	5.6	5.6
Syphilis (020-029)	2	0	2	1	0	1	2.8	-	5.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	103	40	63	65	27	38	180.6	150.0	211.1
Diabetes (260)	5	1	4	2	1	1	5.6	5.6	5.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	65	30	35	58	30	28	161.1	166.7	155.6
Heart diseases (410-443)	187	85	102	150	79	71	416.7	438.9	394.4
Arteriosclerosis (450)	11	5	6	9	5	4	25.0	27.8	22.2
Influenza (480-483)	3	0	3	3	0	3	8.3	-	16.7
#Pneumonia (490-493)	15	4	11	11	4	7	30.6	22.2	38.9
#Diarrhea & enteritis (571)	9	0	9	4	0	4	11.1	-	22.2
Acute nephritis (590, 591)	1	1	0	1	1	0	2.8	5.6	-
Chronic nephritis, etc. (592-594)	9	4	5	4	3	1	11.1	16.7	5.6
Maternal deaths (640-689)	1	0	1	0	0	0	-	-	-
Cong. malformations (750-759)	5	3	2	3	3	0	8.3	16.7	-
Birth inj., atelect. (760-762)	3	1	2	1	1	0	0.8	1.9	-
#Other dis. early inf. (763-776)	31	3	28	21	3	18	17.6	5.6	27.3
Violent & acc. deaths (800-999)	68	26	42	48	23	25	133.3	127.8	138.9
Acc. deaths, total (800-962)	57	22	35	40	20	20	111.1	111.1	111.1
Motor vehicle acc. (810-835)	16	9	7	17	10	7	47.2	55.6	38.9
Ill-defined (780-795)	66	14	52	59	11	48	163.9	61.1	266.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WASHINGTON COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	2,830	1,239	1,591	2,727	1,178	1,549	44.7	39.3	50.0
Fetal deaths	207	121	86	199	109	90	73.0	92.5	58.1
Deaths from all causes	759	238	521	819	226	593	13.4	7.5	19.1
Deaths under 1 year	127	24	103	124	21	103	45.5	17.8	66.5
Deaths under 1 month	75	22	53	76	20	56	27.9	17.0	36.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	3	1	2	4.9	3.3	6.4
Tbc., respiratory (001-008)	1	1	0	3	1	2	4.9	3.3	6.4
Syphilis (020-029)	4	1	3	4	1	3	6.6	3.3	9.7
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	1.6	-	3.2
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	0	0	0	-	-	-
Meningococcal infections (057)	1	0	1	1	0	1	1.6	-	3.2
Acute poliomyelitis (080)	0	0	0	1	0	1	1.6	-	3.2
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	79	33	46	87	28	59	142.6	93.3	190.3
Diabetes (260)	4	1	3	6	1	5	9.8	3.3	16.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	76	22	54	80	19	61	131.1	63.3	196.8
Heart diseases (410-443)	220	85	135	238	81	157	390.2	270.0	506.4
Arteriosclerosis (450)	6	3	3	7	2	5	11.5	6.7	16.1
Influenza (480-483)	6	0	6	5	0	5	8.2	-	16.1
#Pneumonia (490-493)	25	9	16	23	7	16	37.7	23.3	51.6
#Diarrhea & enteritis (571)	10	1	9	10	0	10	16.4	-	32.2
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	13	6	7	12	4	8	19.7	13.3	25.8
Maternal deaths (640-689)	4	0	4	5	0	5	1.8	-	3.2
Cong. malformations (750-759)	10	5	5	13	7	6	21.3	23.3	19.4
Birth inj., atelect. (760-762)	16	8	8	15	7	8	5.5	5.9	5.2
#Other dis. early inf. (763-776)	48	10	38	50	10	40	18.3	8.5	25.8
Violent & acc. deaths (800-999)	71	20	51	88	29	59	144.3	96.7	190.3
Acc. deaths, total (800-962)	60	16	44	78	27	51	127.9	90.0	164.5
Motor vehicle acc. (810-835)	19	7	12	24	9	15	39.3	30.0	48.4
Ill-defined (780-795)	111	8	103	112	7	105	183.6	23.3	338.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WAYNE COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	509	339	170	486	280	206	28.6	25.4	34.3
Fetal deaths	22	10	12	32	14	18	65.8	50.0	87.4
Deaths from all causes	124	71	53	164	91	73	9.6	8.3	12.2
Deaths under 1 year	15	7	8	20	7	13	41.2	25.0	63.1
Deaths under 1 month	7	4	3	13	6	7	26.7	21.4	34.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	2	1	1	11.8	9.1	16.7
Tbc., respiratory (001-008)	1	0	1	2	1	1	11.8	9.1	16.7
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	1	0	1	1	0	1	5.9	-	16.7
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	13	11	2	16	12	4	94.1	109.1	66.7
Diabetes (260)	0	0	0	1	1	0	5.9	9.1	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	23	11	12	23	11	12	135.3	100.0	200.0
Heart diseases (410-443)	30	21	9	47	33	14	276.5	300.0	233.3
Arteriosclerosis (450)	5	1	4	6	2	4	35.3	18.2	66.7
Influenza (480-483)	1	0	1	1	0	1	5.9	-	16.7
#Pneumonia (490-493)	2	1	1	2	1	1	11.8	9.1	16.7
#Diarrhea & enteritis (571)	0	0	0	2	1	1	11.8	9.1	16.7
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	4	2	2	23.5	18.2	33.3
Maternal deaths (640-689)	0	0	0	1	0	1	2.0	-	4.8
Cong. malformations (750-759)	1	0	1	1	0	1	5.9	-	16.7
Birth inj., atelect. (760-762)	1	1	0	3	2	1	6.2	7.1	4.8
#Other dis. early inf. (763-776)	7	5	2	8	3	5	16.5	10.7	24.3
Violent & acc. deaths (800-999)	10	7	3	13	7	6	76.5	63.6	100.0
Acc. deaths, total (800-962)	9	6	3	12	6	6	70.6	54.5	100.0
Motor vehicle acc. (810-835)	4	4	0	5	5	0	29.4	45.4	-
Ill-defined (780-795)	12	4	8	13	4	9	76.5	36.4	150.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WEBSTER COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	167	94	73	224	145	79	20.4	18.1	26.3
Fetal Deaths	3	0	3	5	0	5	22.3	-	63.3
Deaths from all causes	92	79	13	121	100	21	11.0	12.5	7.0
Deaths under 1 year	4	2	2	16	10	6	71.4	69.0	75.9
Deaths under 1 month	4	2	2	14	9	5	62.5	62.1	63.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	2	0	2	2	0	18.2	25.0	-
Tbc., respiratory (001-008)	2	2	0	2	2	0	18.2	25.0	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	7	7	0	14	13	1	127.3	162.5	33.3
Diabetes (260)	0	0	0	1	1	0	9.1	12.5	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	26	21	5	26	20	6	236.4	250.0	200.0
Heart diseases (410-443)	32	30	2	35	33	2	318.2	412.5	66.7
Arteriosclerosis (450)	0	0	0	1	1	0	9.1	12.5	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	2	0	4	3	1	36.4	37.5	33.3
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	9.1	12.5	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	9.1	12.5	-
Birth inj., atelect. (760-762)	1	1	0	4	3	1	17.8	20.7	12.6
#Other dis. early inf. (763-776)	3	1	2	10	6	4	44.6	41.4	50.6
Violent & acc. deaths (800-999)	9	9	0	6	5	1	54.5	62.5	33.3
Acc. deaths, total (800-962)	9	9	0	5	4	1	45.4	50.0	33.3
Motor vehicle acc. (810-835)	7	7	0	4	3	1	36.4	37.5	33.3
Ill-defined (780-795)	4	1	3	6	2	4	54.5	25.0	133.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WILKINSON COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence				Rate*		
	Total	White	Non-white	Number			Total	White	Non-white	
				Total	White	Non-white				
Live births	465	194	271	386	69	317	35.1	23.0	39.6	
Fetal deaths	11	1	10	14	1	13	36.3	14.5	41.0	
Deaths from all causes	176	82	94	156	59	97	14.2	19.7	12.1	
Deaths under 1 year	24	9	15	20	6	14	51.8	87.0	44.2	
Deaths under 1 month	15	7	8	13	5	8	33.7	72.5	25.2	
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.										
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-	-
Measles (085)	1	0	1	1	0	1	9.1	-	12.5	
Malaria (110-117)	0	0	0	0	0	0	-	-	-	-
Cancer, incl. leukemia (140-205)	25	15	10	21	10	11	190.9	333.3	137.5	
Diabetes (260)	1	0	1	2	0	2	18.2	-	25.0	
Pellagra (281)	0	0	0	0	0	0	-	-	-	-
Intracranial lesions (330-334)	18	11	7	17	9	8	154.5	300.0	100.0	
Heart diseases (410-443)	48	27	21	37	18	19	336.4	600.0	237.5	
Arteriosclerosis (450)	1	1	0	1	1	0	9.1	33.3	-	
Influenza (480-483)	0	0	0	0	0	0	-	-	-	-
#Pneumonia (490-493)	8	1	7	7	0	7	63.6	-	87.5	
#Diarrhea & enteritis (571)	3	1	2	1	1	0	9.1	33.3	-	
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	9.1	-	12.5	
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-	-
Cong. malformations (750-759)	1	1	0	3	1	2	27.3	33.3	25.0	
Birth inj., atelect. (760-762)	6	3	3	6	2	4	15.5	29.0	12.6	
#Other dis. early inf. (763-776)	9	4	5	6	2	4	15.5	29.0	12.6	
Violent & acc. deaths (800-999)	9	3	6	7	1	6	63.6	33.3	75.0	
Acc. deaths, total (800-962)	6	3	3	5	1	4	45.4	33.3	50.0	
Motor vehicle acc. (810-835)	1	1	0	1	0	1	9.1	-	12.5	
Ill-defined (780-795)	18	3	15	17	3	14	154.5	100.0	175.0	

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WINSTON COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	429	167	262	474	190	284	23.7	17.3	31.6
Fetal deaths	10	2	8	14	3	11	29.5	15.8	38.7
Deaths from all causes	135	85	50	164	97	67	8.2	8.8	7.4
Deaths under 1 year	15	7	8	17	8	9	35.9	42.1	31.7
Deaths under 1 month	10	7	3	11	8	3	23.2	42.1	10.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	1	0	1	1	0	5.0	9.1	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	13	10	3	13	10	3	65.0	90.9	33.3
Diabetes (260)	5	1	4	7	1	6	35.0	9.1	66.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	25	15	10	26	15	11	130.0	136.4	122.2
Heart diseases (410-443)	43	32	11	53	37	16	265.0	336.4	177.8
Arteriosclerosis (450)	2	2	0	3	3	0	15.0	27.3	-
Influenza (480-483)	2	1	1	2	1	1	10.0	9.1	11.1
#Pneumonia (490-493)	4	3	1	5	3	2	25.0	27.3	22.2
#Diarrhea & enteritis (571)	2	1	1	2	1	1	10.0	9.1	11.1
Acute nephritis (590, 591)	2	1	1	2	1	1	10.0	9.1	11.1
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	5.0	9.1	-
Maternal deaths (640-689)	1	0	1	1	0	1	2.1	-	3.5
Cong. malformations (750-759)	3	2	1	3	2	1	15.0	18.2	11.1
Birth inj., atelect. (760-762)	3	2	1	4	3	1	8.4	15.8	3.5
#Other dis. early inf. (763-776)	6	4	2	6	4	2	12.6	21.0	7.0
Violent & acc. deaths (800-999)	6	2	4	12	4	8	60.0	36.4	88.9
Acc. deaths, total (800-962)	3	1	2	7	2	5	35.0	18.2	55.6
Motor vehicle acc. (810-835)	1	1	0	1	1	0	5.0	9.1	-
Ill-defined (780-795)	4	1	3	7	2	5	35.0	18.2	55.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
YALOBUSHUA COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	311	104	207	313	111	202	24.1	15.8	33.7
Fetal deaths	4	0	4	8	5	3	25.6	45.0	14.8
Deaths from all causes	141	85	56	153	93	60	11.8	13.3	10.0
Deaths under 1 year	10	0	10	12	1	11	38.3	9.0	54.4
Deaths under 1 month	7	0	7	9	1	8	28.8	9.0	39.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	1	0	1	1	0	1	7.7	-	16.7
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	14	10	4	18	14	4	138.5	200.0	66.7
Diabetes (260)	4	2	2	4	2	2	30.8	28.6	33.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	23	13	10	27	16	11	207.7	228.6	183.3
Heart diseases (410-443)	53	36	17	54	36	18	415.4	514.3	300.0
Arteriosclerosis (450)	2	1	1	2	2	0	15.4	28.6	-
Influenza (480-483)	1	1	0	2	1	1	15.4	14.3	16.7
#Pneumonia (490-493)	4	3	1	4	3	1	30.8	42.8	16.7
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	7	4	3	6	3	3	46.2	42.8	50.0
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	2	1	1	15.4	14.3	16.7
Birth inj., atelect. (760-762)	1	0	1	1	0	1	3.2	-	5.0
#Other dis. early inf. (763-776)	5	0	5	6	0	6	19.2	-	29.7
Violent & acc. deaths (800-999)	10	8	2	9	7	2	69.2	100.0	33.3
Acc. deaths, total (800-962)	10	8	2	9	7	2	69.2	100.0	33.3
Motor vehicle acc. (810-835)	4	3	1	5	4	1	38.5	57.1	16.7
Ill-defined (780-795)	6	2	4	5	1	4	38.5	14.3	66.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
YAZOO COUNTY, MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	870	251	619	922	290	632	31.8	26.4	35.1
Fetal deaths	36	5	31	47	6	41	51.0	20.7	64.9
Deaths from all causes	338	115	223	372	140	232	12.8	12.7	12.9
Deaths under 1 year	40	8	32	45	10	35	48.8	34.5	55.4
Deaths under 1 month	26	5	21	29	6	23	31.4	20.7	36.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	3	0	3	10.3	-	16.7
Tbc., respiratory (001-008)	1	0	1	2	0	2	6.9	-	11.1
Syphilis (020-029)	1	0	1	1	0	1	3.4	-	5.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	33	18	15	36	19	17	124.1	172.7	94.4
Diabetes (260)	4	1	3	6	2	4	20.7	18.2	22.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	42	18	24	40	19	21	137.9	172.7	116.7
Heart diseases (410-443)	69	34	35	77	41	36	265.5	372.7	200.0
Arteriosclerosis (450)	3	1	2	4	1	3	13.8	9.1	16.7
Influenza (480-483)	3	1	2	3	1	2	10.3	9.1	11.1
#Pneumonia (490-493)	11	4	7	14	5	9	48.3	45.4	50.0
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	0	1	2	0	2	6.9	-	11.1
Chronic nephritis, etc. (592-594)	1	0	1	2	0	2	6.9	-	11.1
Maternal deaths (640-689)	4	0	4	4	0	4	4.3	-	6.3
Cong. malformations (750-759)	3	0	3	3	1	2	10.3	9.1	11.1
Birth inj., atelect. (760-762)	2	1	1	3	2	1	3.2	6.9	1.6
#Other dis. early inf. (763-776)	16	4	12	16	4	12	17.4	13.8	19.0
Violent & acc. deaths (800-999)	42	16	26	45	22	23	155.2	200.0	127.8
Acc. deaths, total (800-962)	33	15	18	37	21	16	127.6	190.9	88.9
Motor vehicle acc. (810-835)	8	8	0	13	13	0	44.8	118.2	-
Ill-defined (780-795)	67	6	61	67	6	61	231.0	54.5	338.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
STATE OF MISSISSIPPI, 1959

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	61157	28448	32709	60562	27979	32583	27.2	20.9	36.6
Fetal deaths	2649	1081	1568	2603	1045	1558	43.0	37.3	47.8
Deaths from all causes	20713	10633	10080	20890	10742	10148	9.4	8.0	11.4
Deaths under 1 year	2329	657	1672	2355	677	1678	38.9	24.2	51.5
Deaths under 1 month	1409	484	925	1450	515	935	23.9	18.4	28.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	152	61	91	153	62	91	6.9	4.6	10.2
Tbc., respiratory (001-008)	141	59	82	142	60	82	6.4	4.5	9.2
Syphilis (020-029)	41	9	32	43	9	34	1.9	0.7	3.8
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	27	4	23	26	4	22	1.2	0.3	2.5
Scarlet fever & streptococcal sore throat (050, 051)	3	1	2	2	0	2	0.1	-	0.2
Diphtheria (055)	4	0	4	4	0	4	0.2	-	0.4
Whooping cough (056)	18	1	17	18	1	17	0.8	0.1	1.9
Meningococcal infections (057)	8	4	4	8	4	4	0.4	0.3	0.4
Acute poliomyelitis (080)	7	3	4	9	4	5	0.4	0.3	0.6
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	8	2	6	8	2	6	0.4	0.1	0.7
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	2496	1509	987	2600	1580	1020	116.7	118.3	114.5
Diabetes (260)	277	122	155	281	128	153	12.6	9.6	17.2
Pellagra (281)	3	0	3	3	0	3	0.1	-	0.3
Intracranial lesions (330-334)	2693	1397	1296	2683	1386	1297	120.5	103.7	145.6
Heart diseases (410-443)	6218	3791	2427	6219	3772	2447	279.2	282.3	274.6
Arteriosclerosis (450)	297	191	106	294	193	101	13.2	14.4	11.3
Influenza (480-483)	83	25	58	82	25	57	3.7	1.9	6.4
#Pneumonia (490-493)	672	294	378	686	303	383	30.8	22.7	43.0
#Diarrhea & enteritis (571)	177	26	151	175	25	150	7.8	1.9	16.8
Acute nephritis (590, 591)	52	13	39	51	12	39	2.3	0.9	4.4
Chronic nephritis, etc. (592-594)	235	86	149	239	89	150	10.7	6.7	16.8
Maternal deaths (640-689)	55	9	46	55	9	46	0.9	0.3	1.4
Cong. malformations (750-759)	227	121	106	247	140	107	11.1	10.5	12.0
Birth inj., atelect. (760-762)	352	183	169	356	185	171	5.9	6.6	5.2
#Other dis. early inf. (763-776)	855	246	609	857	244	613	14.2	8.7	18.8
Violent & acc. deaths (800-999)	1856	988	868	1862	991	871	83.6	74.2	97.8
Acc. deaths, total (800-962)	1493	830	663	1501	837	664	67.4	62.6	74.5
Motor vehicle acc. (810-835)	605	408	197	588	401	187	26.4	30.0	21.0
III-defined (780-795)	1857	444	1413	1837	432	1405	82.5	32.3	157.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, maternal deaths, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.