


PUBLIC HEALTH STATISTICS
STATE OF
MISSISSIPPI
1960



VITAL STATISTICS
SUMMARIES

Mississippi State Board of Health

Jackson, Mississippi

A. L. Gray, M.D., Executive Officer

ANNUAL BULLETIN

OF

VITAL STATISTICS

1960

SUMMARIES

MISSISSIPPI STATE BOARD OF HEALTH

TABLE OF CONTENTS

	Page
General Summary	1
Table	
1. Census Population as of April 1, 1960, by County, by Race	19
2. Live Births, by County and City of Occurrence, Reallocated to Mother's Usual Residence, by Race, with Rates	21
3. Live Births, by County and City of Usual Residence of Mother, by Race and Attendant, and by Sex	24
4. Fetal Deaths, by County and City of Occurrence, Reallocated to Mother's Usual Residence, by Race and Attendant, and by Sex	28
5. Deaths, by Cause, by Race	32
6. Deaths, by County and City of Occurrence, Reallocated to Place of Residence, by Race, with Rates	43
7. Deaths from Selected Causes, by Race, Sex, and Age, with Rates	46
8. Deaths Under 1 Year, Shown for Selected Causes, by Race and Age, with Rates	54
9. Deaths from Selected Causes, by Month of Occurrence	56
10. Marriages, by County of Occurrence and of Residence of Bride, and Divorces, by County of Occurrence, by Race	57
11. Divorces, by Cause, by Number of Minor Children, by Number of Years Married, by Race	59
12. Live Births, 1913-1960, by Year, by Race, with Rates	60
13. Deaths, 1913-1960, by Year, by Race, with Rates	62
14. Maternal Deaths, 1913-1960, by Year, by Race, with Rates	64
15. Deaths Under 1 Year, 1917-1960, by Year, by Race, with Rates	65
16. Deaths Under 1 Month, 1924-1960, by Year, by Race, with Rates	66
17. Deaths from Heart Diseases, Nephritis (all forms), Vascular Lesions, Cancer, and Tuberculosis (all forms), 1951-1960, by Year, by Race, with Rates	67
18. Births, Deaths, and Marriages, by Month of Occurrence	68

COUNTY TABLES AND STATE SUMMARY TABLE

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 1 MONTH, AND
DEATHS FROM CERTAIN SELECTED CAUSES, ON BASIS OF PLACE OF OCCURRENCE
AND PLACE OF RESIDENCE, BY RACE, WITH RATES:

	Page
For Each County	71
For the State of Mississippi	153

SUMMARY OF 1960 VITAL STATISTICS

INTRODUCTION

The 1960 Vital Statistics Bulletin is the eighteenth consecutive issue prepared from data obtained from records of live births, fetal deaths, deaths, marriages and divorces collected by the Division of Public Health Statistics.

Population

Population figures used in this bulletin are those of the census count of April 1, 1960, from final tabulations published by the U. S. Bureau of the Census.

Occurrence and Residence Allocation

Births, fetal deaths, deaths, and marriages were classified by occurrence and residence. Occurrence data were obtained from certificates filed in Mississippi at the place of birth, fetal death, death, or issuance of marriage license. Residence data include transcripts of these events (except marriages) occurring to Mississippi residents outside the State, and exclude records filed in Mississippi for nonresidents. There were 1,009 births to residents of Mississippi which occurred outside the State in 1960 and were reported to the Division of Public Health Statistics; births occurring inside Mississippi to residents of other states numbered 1,530. Fetal death transcripts totaled 23 and the nonresident certificates 60. During the year, 812 death transcripts were received and 682 nonresident deaths were recorded. The total number of states sending transcripts was 40, including the District of Columbia. Birth records came from 29 states, fetal death records from 6 states, and death records from 33 states. Three reports on Mississippi residents came from foreign countries. In Table A the states furnishing the largest numbers of records are listed.

Some of the tables contain both residence and occurrence data; but, following recommended procedures, most of the tables and all of the rates are based on residence data only. Residence assignment for births and fetal deaths was determined on the basis of usual residence of mother; and for deaths, on the basis of usual residence of decedent. Deaths of inmates or employees of institutions were reallocated to previous residence regardless of length of stay in the institution.

Table A. Transcripts received, by place of occurrence

Place of Occurrence	Live Births	Deaths	Fetal Deaths
<u>Total</u>	<u>1,009</u>	<u>812</u>	<u>23</u>
Alabama	224	76	5
Arkansas	19	12	0
Colorado	29	3	0
Florida	4	20	0
Illinois	62	23	3
Kentucky	33	4	0
Louisiana	135	169	2
Missouri	31	13	0
Tennessee	392	412	11
Texas	4	17	0
<u>Other</u>	<u>76</u>	<u>63</u>	<u>2</u>

Urban-Rural Classification

Towns with populations of 2,500 or more according to the 1960 census were considered as urban areas. All other areas were classified as rural.

Classification of Causes of Death

Causes of death were assigned according to the Seventh Revision (1955) of the International Statistical Classification of Causes of Death. This revision differs only slightly from the Sixth Revision used from 1949 through 1957 and rates for the years prior to 1958 have not been adjusted for comparability with the Seventh Revision. Only underlying causes were coded.

LIVE BIRTHS

Resident Births

Live births to residents of Mississippi numbered 59,210 in 1960. This total was 1,352 less than that for 1959, and was less than the number in each year for the period 1946-1957. There was a decrease of 781 for whites and 571 for nonwhites from 1959. The crude birth rate of 27.2 per 1,000 population was the same as the 1959 rate. Except for the 1958 rate of 27.1, this was the lowest since 1946. The State rate for whites in 1960 was 21.6 and county rates varied from 9.6 in Carroll County to 30.8 in Harrison County. For nonwhites, the State rate was 34.8 and the range for the counties was from 23.6 in Tishomingo County to 43.4 in Smith County.

Characteristics

A breakdown of the 32,012 nonwhite births by specific race shows 31,853 were Negro, 116 were Indian, 23 were Chinese, 15 were Japanese, and 5 were other nonwhites. Births of males numbered 29,952 and there were 29,258 females. These totals yielded a sex ratio of 1,024 males to 1,000 females. September was the peak month of occurrence for live births. There were 5,600 in this month in comparison with only 4,160 in May when the fewest occurred.

During 1960, 22,189 children were born to urban residents and 37,021 to rural residents. Over half (54 per cent) of the infants born to urban residents were white, but 59 per cent of the rural were nonwhite. Of the births to rural residents, 22,280 or 60.2 per cent were born to farm residents; 13,906 were born to nonfarm residents; and on 835 certificates information pertaining to farm residency was not stated. Comparison of the occurrence and residence figures indicates that a majority of the white mothers living in rural areas went to cities for delivery as compared with a much smaller percentage of the nonwhite mothers.

The number of illegitimate births during 1960 was 8,407 or 14.2 per cent of the total. This was 316 more than in 1959, with an increase in percentage of 0.8. White births classified as illegitimate totaled 388 or 1.4 per cent, and 8,019 or 25.0 per cent of the nonwhite births were illegitimate.

Liveborn infants classified as immature (birth weight 5 1/2 pounds or less) numbered 5,495 or 9.2 per cent of the total number of births registered. The incidence of immaturity was greater among the nonwhites than among the whites, 11.4 per cent in

comparison with 6.7 per cent.

Classification by Attendant

In Table B, which gives the distribution of live births by attendant and race, it may be seen that, in 1960, about 73 per cent of all births occurred in hospitals or clinics (including physicians' offices or clinics which provide inpatient care). A small proportion, 2.8 per cent, was delivered by physicians in places other than hospitals or clinics; and 24 per cent were delivered at home by nonmedical attendants. Almost 44 per cent of the nonwhite mothers, but less than one per cent of the white mothers, had no medical attendant.

Table B. Live births by attendant and race

Attendant	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
<u>Total</u>	<u>59,210</u>	<u>100.0</u>	<u>27,198</u>	<u>100.0</u>	<u>32,012</u>	<u>100.0</u>
Physician in hospital	43,349	73.2	26,931	99.0	16,418	51.3
Physician in home	1,687	2.8	135	0.5	1,552	4.8
Midwife	14,082	23.8	122	0.4	13,960	43.6
Other	92	0.2	10	0.04	82	0.2

There has been a steady increase in the proportion of live births occurring in hospitals in recent years (see Table C). The year 1951 was the first in which at least half of the deliveries took place in hospitals, and the proportion of hospital deliveries has increased each year since. The progress made by the nonwhites has been more rapid, and in 1960, for the first time, more than half of the nonwhite mothers were delivered in hospitals. There is still, however, a large difference between the percentages for the two racial groups.

Table C. Percentage of live births in hospitals or clinics, 1956-1960, by race

Year	Total	White	Nonwhite
1960	73.2	99.0	51.3
1959	72.1	98.9	49.1
1958	70.4	98.6	47.0
1957	68.3	98.0	44.8
1956	65.7	97.2	41.0

The proportions of live births attended by physicians during 1956-1960 are shown in Table D. The improvement shown in this table is less pronounced than that in the preceding table, but is also steady. This is the fifth year in which more than half the nonwhite mothers were attended by physicians.

Table D. Percentage of live births attended by physicians, 1956-1960, by race

Year	Total	White	Nonwhite
1960	76.1	99.5	56.1
1959	75.6	99.5	55.0
1958	74.2	99.5	53.3
1957	73.6	99.5	53.2
1956	72.2	99.3	51.0

Age of Mother

Table E gives the age distribution of mothers by race. Little variation in this distribution has occurred in recent years. Approximately half of the mothers were 15-24 years of age and more than one-third were 25-34 years old. In the periods 15-24 and 25-34 years the percentages of white mothers were slightly larger than those of nonwhites.

Table E. Live births by age of mother and race, with per cent of total births

Age of Mother	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
10-14 years	301	0.5	25	0.09	276	0.9
15-44 years	58,696	99.1	27,123	99.7	31,573	98.6
15-24 years	31,001	52.4	14,417	53.0	16,584	51.8
25-34 years	21,420	36.2	10,340	38.0	11,080	34.6
35-44 years	6,275	10.6	2,366	8.7	3,909	12.2
45-54 years	123	0.2	39	0.1	84	0.3
Unknown	90	0.2	11	0.04	79	0.2

FETAL DEATHS

In 1952 the registration of all fetal deaths, regardless of length of gestation, was initiated in Mississippi. Prior to that time, only stillbirths of 20 or more weeks gestation were reported. Registration of early fetal deaths is still far from complete, but there is no estimate of the degree of nonregistration. There were 2,421 fetal death certificates filed during 1960 and 2,384 certificates for residents. The resident total was 219 less than the comparable figure for 1959, with a decrease of 90 for whites and 129 for nonwhites. For the state the fetal death ratio (fetal deaths per 1,000 live births) was 40.3. The ratio for whites was 35.1, with county ratios varying from zero in Winston County to 142.8 in Issaquena County. The ratio for nonwhites was 44.6, with variations from zero in George, Itawamba, Lamar, Pontotoc, Tishomingo and Webster Counties to 96.8 in Perry County.

On 835 of the records the gestation period was stated to be less than 20 weeks; on 1,403 it was 20 weeks or more; and on 146 the length was not stated. In the period 1956-1960 there have been reported among the whites more fetal deaths after less than 20 weeks of gestation than after 20 weeks or more.

Using the certificates with sex stated, a sex ratio of 1,252 males to 1,000 females was determined. This ratio was the lowest recorded recently. As usual, the sex ratio for fetal deaths was much higher than that for live births, but was not so reliable a figure because the sex was not stated on 695 or 29 per cent of the fetal death certificates.

The distribution by race and attendant is given in Table F. Approximately 60 per cent of the total were nonwhite. About 75 per cent of the reported fetal deaths were attended by physicians in hospitals and about 18 per cent were attended by midwives. Although the proportion of white live births in hospitals was larger than the percentage of white fetal deaths in hospitals, the reverse was true for the nonwhites. This situation in the nonwhite group may be accounted for by the nonreporting of fetal deaths that occur at home and the tendency to transfer difficult and premature deliveries to a hospital.

Table F. Fetal deaths by attendant and race

Attendant	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
<u>Total</u>	<u>2,384</u>	<u>100.0</u>	<u>955</u>	<u>100.0</u>	<u>1,429</u>	<u>100.0</u>
Physician in hospital	1,768	74.2	912	95.5	856	59.9
Physician in home	171	7.2	41	4.3	130	9.1
Midwife	431	18.1	0	-	431	30.2
Other	14	0.6	2	0.2	12	0.8

Table G shows the percentage of fetal deaths delivered in hospitals in the last five years. Increases have occurred among both whites and nonwhites. The gap between the two race groups is a large one, but not so large as that for live births.

Table G. Percentage of fetal deaths delivered in hospitals, 1956-1960, by race

Year	Total	White	Nonwhite
1960	74.2	95.5	59.9
1959	73.5	94.4	59.6
1958	69.0	88.8	56.8
1957	69.5	92.2	54.8
<u>1956</u>	<u>67.2</u>	<u>91.0</u>	<u>52.7</u>

DEATHS

Resident Deaths

In 1960 the resident death total was 21,713, an increase of 823 over the number for 1959. The crude death rate, based on the 1960 census population, was 10.0 per 1,000 population in comparison with a rate of 9.4 in 1959 based on the estimated population for that year. There were 7,948 urban residents who died in 1960 (36.6 per cent of the total) but 51.8 per cent of the deaths occurred in urban areas.

Although the number of deaths of nonwhites was 695 less than that of the whites, the death rate of the nonwhites was 2.5 per 1,000 greater than that of the whites. The total of 11,204 deaths of white persons was the largest reported since 1913, the first complete year of death registration in Mississippi. The number of deaths of nonwhites was higher than in any of the six preceding years except 1958.

The mortality rate for whites was greater than in 1959, while the rate for nonwhites remained the same.

County rates for the white population ranged from 5.8 in Jackson to 16.5 in Yalobusha, with a rate for the State of 8.9, while county rates for the nonwhites ranged from 6.4 in Webster to 19.6 in Issaquena, with a State rate of 11.4.

The resident birth rate exceeded the death rate by 17.2 per 1,000 population. For whites the excess was 12.7 and for the nonwhites it was 23.4.

Age at Death

Table H shows the distribution of deaths by age and race. In comparison with the 1959 figures there were increases in all the white groups except the 10-24 year group and the group 85 years and over showed an increase of 15.6 per cent. Increases occurred in all nonwhite groups except the 1-4, 15-44, and 55-64 year groups. For deaths of all races combined there were decreases only in the 1-4, 10-24 and 55-64 year groups, the greatest decrease being in the 15-24 year group and the largest increase in the group 85 years and over.

Comparison of the age distribution of the two races reveals significant differences. The death rate for nonwhites under 55 was almost double that for whites. In all age groups except 85 years and over the death rate for nonwhites was higher than that for whites. "Retirement" age or 65 years had been reached by 60 per cent of the whites who died in 1960 and only 47 per cent of the nonwhites. The median age at death for the state was 65.8; for the whites 68.4; and for the nonwhites 61.5.

Table H. Age-race distribution of deaths

Age Group	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
All Ages	<u>21,713</u>	<u>10.0</u>	<u>11,204</u>	<u>8.9</u>	<u>10,509</u>	<u>11.4</u>
Under 1 year	2,457	41.5	714	26.2	1,743	54.4
1-4 years	378	1.7	116	1.1	262	2.3
5-9 years	185	0.7	64	0.5	121	0.9
10-14 years	139	0.6	51	0.4	88	0.8
15-24 years	465	1.4	222	1.2	243	1.7
25-34 years	514	2.2	228	1.5	286	3.6
35-44 years	966	4.0	460	2.9	506	6.3
45-54 years	1,951	8.5	982	6.7	969	11.7
55-64 years	2,965	17.2	1,597	14.8	1,368	21.4
65-74 years	4,769	38.3	2,621	34.9	2,148	43.5
75-84 years	4,615	85.6	2,831	83.2	1,784	89.7
85 years +	2,297	198.2	1,317	198.5	980	197.8
Unknown	12	-	1	-	11	-

* Rate per 1,000 population in the specified age and race group, except for the age group under 1 year. Rate for deaths under 1 year per 1,000 live births.

Leading Causes of Death

In Table I are listed the ten leading causes of death in the State in 1960. The first six of these causes are the same as in the seven preceding years. Congenital malformations, which was in eighth place in 1956, in ninth place in 1957, and in tenth place in 1958 and 1959, does not appear among the ten leading causes in 1960. Diabetes mellitus, which was in eighth place in 1959, moved to seventh place in 1960. General arteriosclerosis moved from seventh to eighth place. Hypertension remained in ninth place. Gastritis, duodenitis, enteritis and colitis, in tenth place in 1960, had not appeared as one of the ten leading causes since 1952.

Altogether the ten leading causes were responsible for 77 per cent of the deaths. For the whites the same ten causes were listed as for the total population with the exception that congenital malformations ranked in eighth place, and suicide in tenth place, while hypertension and gastritis, duodenitis, enteritis and colitis were not on the list. For nonwhites chronic and unspecified nephritis and other renal sclerosis ranked in ninth place, and general arteriosclerosis was not on the list.

Table I. Ten leading causes of death by race, with per cent of total deaths

Cause and International List Number	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
1. Diseases of heart (410-443)	6,312	29.1	3,879	34.6	2,433	23.2
2. Vascular lesions affecting central nervous system (330-334)	2,960	13.6	1,513	13.5	1,447	13.8
3. Malignant neoplasms (140-205)	2,635	12.1	1,644	14.7	991	9.4
4. Accidents (800-962)	1,504	6.9	827	7.4	677	6.4
5. Certain diseases of early infancy (760-776)	1,306	6.0	465	4.2	841	8.0
6. Pneumonia and influenza (480-493)	958	4.4	389	3.5	569	5.4
7. Diabetes mellitus (260)	309	1.4	139	1.2	170	1.6
8. General arteriosclerosis (450)	306	1.4	198	1.8	108	1.0
9. Hypertension without mention of heart (444-447)	250	1.2	98	0.9	152	1.4
10. Gastritis, duodenitis, enteritis & colitis, except diarrhea of newborn (543,571,572)	249	1.1	57	0.5	192	1.8

Heart disease, the leading cause of death, was responsible for 29.1 per cent of all deaths in 1960. During the year 6,312 deaths were attributed to some form of heart disease. In Table J it may be seen that a large number of these (4,572 deaths) were specified as arteriosclerotic and degenerative heart disease, and that the next largest group was hypertensive heart disease with 1,138 deaths. The total death rate was 289.8 per 100,000 population, an increase from 279.2 for 1959, due to an increase among whites.

Table J. Deaths from heart disease, by race by type of disease

Cause and International List Number	Race		
	Total	White	Nonwhite
Diseases of the heart			
Chronic rheumatic heart disease (410-416)	6,312	3,879	2,433
Arteriosclerotic and degenerative heart disease (420-422)	108	56	52
Other diseases of heart (430-434)	4,572	3,206	1,366
Hypertension with heart disease (440-443)	494	235	259
	1,138	382	756

For the second leading cause of death, vascular lesions affecting the central nervous system, there were 2,960 deaths representing 13.6 per cent of the total. The components of this category are listed in Table K showing that a majority of the deaths were due to cerebral hemorrhage (1,832) and 650 were due to cerebral embolism or thrombosis. There was an increase in rate over that of the previous year; and, as in previous years, the rate for nonwhites exceeded that for whites.

Table K. Deaths from vascular lesions affecting the central nervous system, by race by type of lesion

Cause and International List Number	Race		
	Total	White	Nonwhite
Vascular lesions affecting central nervous system (330-334)	2,960	1,513	1,447
Subarachnoid hemorrhage (330)	77	43	34
Cerebral hemorrhage (331)	1,832	892	940
Cerebral embolism and thrombosis (332)	650	395	255
Other and ill-defined (334)	401	183	218

In 1960 the cancer death totals for whites were the highest on record in this State. For nonwhites they were lower than in any of the preceding four years. There were 2,635 deaths from this cause or 35 more than in 1959. The distribution of these deaths by age, sex, and race is given in Table L. In comparison with 1959, for each sex group there was an increase among whites and a decrease among nonwhites. For particular age groups the greatest decreases occurred in deaths of white males 75 to 84 years, nonwhite males 55 to 64 years, white females 55 to 64 years, and nonwhite females 65 to 74 years. The largest increases occurred in white males 55 to 64 years, nonwhite males 65 to 74 years, and in white females 75 to 84 years.

More deaths occurred among whites than among nonwhites in all age groups, and the excess was largest in the age group 75-84. There were more deaths of males than females in the age group 55-74, and more of females than males in the age groups 25-54 and 75 years and over.

Table L. Cancer deaths by age, sex, and race, with rates

Age in Years	Number of Deaths					Rate*				
	Total	Male		Female		Total	Male		Female	
		White	Non-white	White	Non-white		White	Non-white	White	Non-white
All Ages	2,635	890	505	754	486	121.0	142.4	114.0	119.2	101.7
Under 10	40	17	7	10	6	7.4	12.8	5.0	7.9	4.3
10-14	12	4	2	3	3	5.0	6.2	3.4	4.9	5.2
15-24	21	9	6	2	4	6.3	9.2	8.6	2.2	5.5
25-34	52	13	7	15	17	22.0	17.0	20.3	18.8	37.1
35-44	149	40	19	51	39	62.4	51.9	54.3	63.0	85.3
45-54	368	114	57	98	99	160.4	157.6	147.4	132.2	223.6
55-64	588	241	105	134	108	341.8	462.2	348.5	239.2	320.0
65-74	746	258	180	188	120	599.1	736.2	757.4	469.0	468.2
75-84	515	154	98	197	66	955.2	1,029.9	1,011.8	1,032.6	647.1
85 +	144	40	24	56	24	1,242.6	1,471.1	1,044.4	1,429.7	904.0

* Rate per 100,000 population in the specified age, sex and race group.

Table M lists cancer deaths by primary site in descending order according to the number in the total population. The number of deaths assigned to the leading site, digestive organs and peritoneum, is 30 less than in 1959. Cancer of the breast and genito-urinary system accounted for the second largest number of cancer deaths. The two leading sites accounted for more than 75 per cent of the cancer deaths among non-white females, 68 per cent among white females, 64 per cent among nonwhite males, but less than 46 per cent among white males. Cancer of the respiratory system showed an increase of 12 deaths, remaining in third rank. The largest increase was that of 26 in the number of deaths from cancer of the buccal cavity and pharynx.

Table M. Cancer deaths by primary site, sex, and race, with rates

Primary Site and International List Number	Male				Female			
	White		Nonwhite		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*	No.	Rate*
Total	890	142.4	505	114.0	754	119.2	486	101.7
1. Digestive organs and peritoneum (150-159)	263	42.1	181	40.9	245	38.7	123	25.7
2. Breast & genito-urinary system (170-181)	142	22.7	142	32.0	268	42.4	244	51.1
3. Respiratory system (160-165)	228	36.5	69	15.6	39	6.2	21	4.4
4. Lymphatic & hematopoietic tissues (200-205)	94	15.0	42	9.5	73	11.5	23	4.8
5. Brain & nervous system (192,193)	38	6.1	9	2.0	24	3.8	9	1.9
6. Buccal cavity & pharynx (140-148)	31	5.0	9	2.0	18	2.8	10	2.1
7. Skin (190,191)	31	5.0	5	1.1	18	2.8	2	0.4
8. Bone & connective tissue (196,197)	14	2.2	8	1.8	8	1.3	3	0.6
9. Other & unspecified sites (194,195,198,199)	49	7.8	40	9.0	61	9.6	51	10.7

* Rate per 100,000 population in the specified sex and race group.

The number of deaths in 1960 which resulted from accidents was 1,504, three more than in 1959. The greatest increases were in the number of deaths from motor vehicles and from falls. The greatest increase among whites was in deaths from motor vehicles, and among nonwhites, from drowning. The greatest decrease for whites was in deaths from drowning. Among nonwhites there was a large decrease in deaths from machinery, which ranked sixth among leading causes in 1959. The largest number of deaths from all accidents occurred in December in 1960, as in the three preceding years, and July was in second place. The number of deaths from motor vehicle accidents was greatest in August and second in June. Table N shows the six leading causes of accidental death in 1960 by race. The first five of these causes appeared on the 1959 list in the same order. Machinery was in sixth place among leading causes of accidental death in 1959 while mechanical suffocation in bed or cradle was in sixth place in 1960.

Table N. Six leading causes of accidental deaths among residents, by race

Type of Accident	Total	White	Nonwhite
1. Motor vehicle	620	428	192
2. Fire and explosion	235	60	175
3. Falls	130	97	33
4. Drowning except water transport	115	44	71
5. Firearm	92	46	46
6. Mechanical suffocation in bed and cradle	35	9	26

Leading causes of death for each age group except infants are given in Table O. None of these lists was exactly like any other, but there were some causes which appeared repeatedly. Accidents was listed in every group, and appeared as the chief cause in all groups under 45 years of age. Malignant neoplasms also appeared on every list and heart disease on all except 1-4 and 15-24 years. Vascular lesions affecting the central nervous system was listed in the three groups 25 and over. Pneumonia and influenza appeared in all groups except 25-44 years. In both lists for children under 15, congenital malformations was included, and for the 15-24 year and 25-44 year groups homicide was listed. Gastritis, duodenitis, enteritis and colitis was listed only for the 1-4 year group. In the age group 65-74 (not shown here) the five leading causes were the same as for the total 65 years and over, and in the same order. In the age groups 75-84 and 85 and over, accidents does not appear on the list of five leading causes, and general arteriosclerosis is in fifth place.

Ill-defined Causes

Of the 21,583 certificates filed for deaths that occurred in Mississippi in 1960 those with no satisfactory statement of cause numbered 2,634 or 12 per cent of the total. For white persons the percentage of certificates in this category was 6.0 and for non-whites 18.8. Senility was the only cause given on 61 of these certificates; ill-defined diseases of early infancy and those due to immaturity unqualified on 753; and other ill-defined and unknown causes on 1,820.

Deaths with unsatisfactory statement of cause were so numerous that if placed on the list of ten leading causes they would rank fourth and would appear as one of the five chief causes of death in every age group. If the true causes of death for these persons could be determined and properly assigned, the order of principal causes might not be the same and even the causes might be different.

If deaths from ill-defined diseases of early infancy and deaths from immaturity unqualified are excluded from the group of deaths without definite cause, the remaining ill-defined deaths are 8.7 per cent of all deaths recorded in Mississippi. This group constitutes more than 20 per cent of the total deaths in Carroll and DeSoto Counties and 15 to 20 per cent of the total deaths in Amite, Holmes, Humphreys, Issaquena, Kemper, Madison and Noxubee Counties.

Table 0. Five leading causes of death by age group

<u>Age and Cause of Death</u>	<u>No. of Deaths</u>	<u>% of Total</u>
1-4 years		
All Causes	<u>378</u>	<u>100.0</u>
1. Accidents	106	28.0
2. Pneumonia and influenza	61	16.1
3. Congenital malformations	30	7.9
4. Gastritis, duodenitis, enteritis and colitis	23	6.1
5. Malignant neoplasms, including leukemia	21	5.6
5-14 years		
All Causes	<u>324</u>	<u>100.0</u>
1. Accidents	148	45.7
2. Malignant neoplasms, including leukemia	28	8.6
3. Pneumonia and influenza	15	4.6
4. Heart disease	11	3.4
5. Congenital malformations	9	2.8
15-24 years		
All Causes	<u>465</u>	<u>100.0</u>
1. Accidents	242	52.0
2. Homicide	39	8.4
3. Malignant neoplasms	21	4.5
3. Pneumonia and influenza	21	4.5
5. Maternal deaths	17	3.6
25-44 years		
All Causes	<u>1,480</u>	<u>100.0</u>
1. Accidents	347	23.4
2. Heart disease	228	15.4
3. Malignant neoplasms	201	13.6
4. Homicide	81	5.5
5. Vascular lesions affecting central nervous system	75	5.1
45-64 years		
All Causes	<u>4,916</u>	<u>100.0</u>
1. Heart disease	1,591	32.4
2. Malignant neoplasms	956	19.4
3. Vascular lesions affecting central nervous system	587	11.9
4. Accidents	297	6.0
5. Pneumonia and influenza	125	2.5
65 years and over		
All Causes	<u>11,681</u>	<u>100.0</u>
1. Heart disease	4,454	38.1
2. Vascular lesions affecting central nervous system	2,280	19.5
3. Malignant neoplasms	1,405	12.0
4. Pneumonia and influenza	475	4.1
5. Accidents	272	2.3

Infant Mortality

An infant mortality rate of 41.5 per 1,000 live births was recorded in 1960, higher than the 1959 rate by 2.6 per 1,000 births. The nonwhite rate showed a greater increase than the white rate. Distribution of infant deaths by age and race is shown in Table P. In each age group, the rate for nonwhites exceeded that for whites, with the greatest difference between the two racial groups occurring in the period 28 days to one year.

Table P. Infant deaths by age and race, with rates

Age at Death	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
<u>Total under 1 year</u>	<u>2,457</u>	<u>41.5</u>	<u>714</u>	<u>26.2</u>	<u>1,743</u>	<u>54.4</u>
Under 1 day	703	11.9	277	10.2	426	13.3
One to 28 days	794	13.4	273	10.0	521	16.3
28 days to one year	960	16.2	164	6.0	796	24.9

* Deaths per 1,000 live births.


Table Q gives the five leading causes of infant deaths. Influenza and pneumonia, except pneumonia of newborn, was in first place for the first time since 1952. Gastritis, duodenitis, enteritis and colitis, which was in fourth place in 1958 and in fifth place in 1959, was back in fourth place, with birth injuries which was in fifth place in 1958 and in fourth place in 1959, was back in fifth place. Immaturity, which is no longer considered a separate classification in ranking causes of infant deaths, was mentioned either in combination with another cause or as the only cause on 43 per cent of the certificates for white infants and 35 per cent of those for nonwhites.

Table Q. Leading causes of infant deaths, by race

Cause of Death	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
<u>All Causes</u>	<u>2,457</u>	<u>100.0</u>	<u>714</u>	<u>100.0</u>	<u>1,743</u>	<u>100.0</u>
1. Influenza & pneumonia, except pneumonia of newborn	227	9.2	29	4.1	198	11.4
2. Postnatal asphyxia & atelectasis	226	9.2	136	19.0	90	5.2
3. Congenital malformations	164	6.7	95	13.3	69	4.0
4. Gastritis, duodenitis, enteritis & colitis, except diarrhea of newborn	163	6.6	12	1.7	151	8.7
5. Birth injuries	106	4.3	50	7.0	56	3.2

Infant death certificates which showed ill-defined causes or no cause at all numbered 1,058 or 43 per cent of the total in 1960. For 33 per cent of the white and 47 per cent of the nonwhite infants no definite cause of death was given. Determination of the true cause of death for this large group of infants might change the leading causes of death which are listed above.

Infant mortality rates by race for 1920-1960 are shown in Graph 1. In general, the trend has been downward for both races, but in recent years there has been some upward swing in the curve for the nonwhites and, in 1960, an increase for whites.



Maternal Mortality

The maternal mortality rate for 1960 was 10 per 10,000 live births in the total population, 2.9 for white mothers and 15.9 for nonwhite. (Note that the population base has been changed from the previously used 1,000 live births to 10,000 live births.) The total of 59 maternal deaths was four more than in 1959. There was a decrease of one death among the white mothers and an increase of five among the nonwhite. The discrepancy between the rates by race narrowed considerably in recent years but the rate for nonwhites has increased slightly in the last two years.

In 1960, as in each year from 1947 through 1959, the two leading causes were toxemia and hemorrhage. Toxemia has been the leading cause in each year of this period except 1957.

Table R. Maternal deaths by cause and race, with rates

<u>Cause of Death</u>	<u>Total</u>		<u>White</u>		<u>Nonwhite</u>	
	<u>No.</u>	<u>Rate*</u>	<u>No.</u>	<u>Rate*</u>	<u>No.</u>	<u>Rate*</u>
<u>All Causes</u>	<u>59</u>	<u>10.0</u>	<u>8</u>	<u>2.9</u>	<u>51</u>	<u>15.9</u>
Toxemia	32	5.4	2	0.7	30	9.4
Hemorrhage	7	1.2	0	-	7	2.2
Sepsis	3	0.5	1	0.4	2	0.6
Abortion with sepsis	3	0.5	1	0.4	2	0.6
Abortion without sepsis or toxemia	1	0.2	0	-	1	0.3
<u>Other complications</u>	<u>13</u>	<u>2.2</u>	<u>4</u>	<u>1.5</u>	<u>9</u>	<u>2.8</u>

* Deaths per 10,000 live births.


Table S gives the age and race-specific maternal death rates. There were no deaths in the 10-14 year group and none in the 45-54 year group. At all ages the rates for nonwhite mothers were higher than those for white mothers.

Table S. Maternal deaths by age group and race, with rates

<u>Age Group</u>	<u>Total</u>		<u>White</u>		<u>Nonwhite</u>	
	<u>No.</u>	<u>Rate*</u>	<u>No.</u>	<u>Rate*</u>	<u>No.</u>	<u>Rate*</u>
<u>All Ages</u>	<u>59</u>	<u>10.0</u>	<u>8</u>	<u>2.9</u>	<u>51</u>	<u>15.9</u>
15-24 years	17	5.5	3	2.1	14	8.4
25-34 years	21	9.8	3	2.9	18	16.2
35-44 years	21	33.5	2	8.4	19	48.6

* Deaths per 10,000 live births to mothers in specific age-race group.

Maternal death rates by race for 1913-1960 are shown in Graph 2. A downward trend began in the 1930's and has been rapid since 1940, though in 1959 and 1960 there was an increase in the death rate among nonwhite mothers.



MARRIAGES

Marriages increased from 20,447 in 1959 to 21,233 in 1960. Among whites they increased from 13,547 to 14,029, and among nonwhites from 6,900 to 7,204.

The average daily number of marriages was 58, two more than the average for 1959. In June, the month with the largest total, the daily average was 78, and in December the daily average was 75.

This was the first marriage for 15,061 or 70.9 per cent of the brides and 14,976 or 70.5 per cent of the grooms. For less than one fourth of each group there had been one previous marriage. One hundred and three brides and 86 grooms did not state the number of previous marriages.

Table T shows the age distribution of brides and grooms. Of the brides, 9,570 or 45 per cent were under 20 years of age; and 71.2 per cent were under 25 years of age. There was a much smaller percentage of grooms in the teen-age group - 16.8 per cent - while 40.9 per cent of the grooms were in the 20-24 year age group. A cross tabulation of the ages of brides and grooms shows marriages of 16 couples with both 75 years of age and over. While the youngest bride of a groom of 75 years of age or more was 22 years of age, most marriages were between persons of approximately the same age.

Table T. Age distribution of brides and grooms

Age Groups	Bride		Groom	
	No.	%	No.	%
<u>Total</u>	<u>21,233</u>	100.0	<u>21,233</u>	100.0
Under 15 years	109	0.5	0	-
15 years	857	4.0	6	0.03
16 years	1,509	7.1	44	0.2
17 years	2,099	9.9	527	2.5
18 years	2,832	13.3	1,209	5.7
19 years	2,164	10.2	1,782	8.4
20-24 years	5,549	26.1	8,683	40.9
25-34 years	2,725	12.8	4,473	21.1
35-44 years	1,554	7.3	1,725	8.1
45-54 years	1,038	4.9	1,227	5.8
55 years and over	782	3.7	1,547	7.3
<u>Unknown</u>	<u>15</u>	<u>0.07</u>	<u>10</u>	<u>0.05</u>

Among the marriages between nonresident brides and grooms there were 5 brides and no groom under 15 years of age. Nonresident brides in the 15-19 year age group numbered 1,512 or 16 per cent of all brides in this age group.

In Table U are shown data on the residence status of both brides and grooms. In 1960 both parties were residents of Mississippi in 74 per cent of the marriages, compared with 72 per cent in 1959. Out-of-state brides were 19 per cent of the total in 1960 and 20 per cent in 1959.

Table U. Marriages in Mississippi by residence status of bride and groom

Groom	Bride			Not Stated
	Total	Resident	Nonresident	
<u>Total</u>	<u>21,233</u>	<u>17,198</u>	<u>3,979</u>	<u>56</u>
Resident	16,265	15,737	520	8
Nonresident	4,898	1,440	3,452	6
Not stated	70	21	7	42

A comparison during the last fourteen years of the resident status of brides is shown in Table V. In each year 1947 through 1956 the number of out-of-state brides increased and, with the exception of 1955, the number of resident brides decreased. The declining trend among resident brides continued through 1959, but the 1960 total shows an increase. The increase in nonresidents ended abruptly when the new marriage law became effective in July, 1958 and there were fewer nonresidents in 1960 than in 1959, the first full year in which the law was applicable.

:

Table V. Marriages in Mississippi by residence status of bride 1947-1960

Year	Total	Resident	Nonresident	Not Stated
1960	21,233	17,198	3,979	56
1959	20,447	16,288	4,062	97
1958	36,402	17,370	18,920	112
1957	62,495	20,864	41,510	121
1956	66,223	23,011	43,212	*
1955	66,423	23,683	42,740	*
1954	61,381	23,555	37,826	*
1953	61,751	24,497	37,254	*
1952	58,116	24,868	33,248	*
1951	57,321	25,713	31,608	*
1950	56,738	26,859	29,879	*
1949	52,765	27,717	25,048	*
1948	53,645	30,921	22,724	*
1947	53,584	34,067	19,517	*

* Not classified.

With the exception of Alaska, Connecticut, Hawaii, Idaho, Montana, Nevada, South Dakota and Wyoming, brides came to Mississippi from every state in the Union and from the District of Columbia, with 9 from foreign countries. Bordering states furnished 14.6 per cent of the brides married in Mississippi and 15.9 per cent of the grooms, with totals as follows: Alabama, 549 brides and 628 grooms; Arkansas, 235 brides and 236 grooms; Louisiana, 944 brides and 1,079 grooms; and Tennessee, 1,364 brides and 1,434 grooms. Illinois furnished 214 brides and 287 grooms. These five states furnished 15.6 per cent of the brides married in Mississippi and 17.2 per cent of the grooms.

Each county in the state issued marriage licenses to one or more out-of-state brides in 1960. Counties adjacent to Tennessee and Arkansas and the coastal counties issued the largest numbers, with most of the remainder issued in other counties adjacent to Alabama and Louisiana. In DeSoto County more than 80 per cent of the licenses were for nonresident brides; in Alcorn, Wilkinson and Hancock Counties about 70 per cent; and in George County more than half. Counties bordering Tennessee issued licenses to 43 per cent of the out-of-state brides.

DIVORCES

Divorces granted in Mississippi during 1960 numbered 5,260 - 152 more than in 1959. There was a decrease of 4 among the whites and an increase of 156 among the nonwhites. Cruel and inhuman treatment and desertion continued to rank as the two leading causes and together accounted for 89 per cent of the total granted. Of the divorces among whites, cruel and inhuman treatment was given as the main cause for 62 per cent; among nonwhites 65 per cent were granted on grounds of desertion. The wife was listed as complainant on 64 per cent of the records.

No minor children were involved in 47 per cent of the divorces, and 22 per cent of the couples had only one minor child. Twelve of the families had 9 or more minor children. The total number of minor children affected was 5,802.

Thirty-seven per cent of the divorces in 1960 were granted to persons married less than 5 years; 5 per cent had been married less than one year; but 18 per cent had been married 20 years or longer. More divorces were granted to couples in their second year of marriage than in any other year. The distribution of divorces according to duration of marriage was different for the two races. Of the white couples, 46 per cent were granted after less than 5 years of marriage and only 10 per cent after 20 years or longer; for the nonwhites, corresponding percentages were 19 and 35 respectively.

TABLE 1
CENSUS POPULATION, MISSISSIPPI
AS OF APRIL 1, 1960

County	Total	White	Nonwhite
<u>Total</u>	<u>2,178,141</u>	<u>1,257,546</u>	<u>920,595</u>
Adams	37,730	19,035	18,695
Alcorn	25,282	21,949	3,333
Amite	15,573	7,130	8,443
Attala	21,335	11,789	9,546
Benton	7,723	4,114	3,609
Bolivar	54,464	17,521	36,943
Calhoun	15,941	11,595	4,346
Carroll	11,177	4,677	6,500
Chickasaw	16,891	10,380	6,511
Choctaw	8,423	5,903	2,520
Claiborne	10,845	2,600	8,245
Clarke	16,493	10,001	6,492
Clay	18,933	9,214	9,719
Coahoma	46,212	14,630	31,582
Copiah	27,051	12,992	14,059
Covington	13,637	8,896	4,741
DeSoto	23,891	9,248	14,643
Forrest	52,722	37,970	14,752
Franklin	9,286	5,486	3,800
George	11,098	9,811	1,287
Greene	8,366	6,443	1,923
Grenada	18,409	9,352	9,057
Hancock	14,039	11,784	2,255
Harrison	119,489	100,233	19,256
Hinds	187,045	112,205	74,840
Holmes	27,096	7,595	19,501
Humphreys	19,093	5,758	13,335
Issaquena	3,576	1,176	2,400
Itawamba	15,080	14,206	874
Jackson	55,522	44,658	10,864
Jasper	16,909	8,402	8,507
Jefferson	10,142	2,489	7,653
Jeff Davis	13,540	6,126	7,414
Jones	59,542	44,095	15,447
Kemper	12,277	4,828	7,449
Lafayette	21,355	14,110	7,245

County	Total	White	Nonwhite
Lamar	13,675	11,443	2,232
Lauderdale	67,119	43,635	23,484
Lawrence	10,215	6,354	3,861
Leake	18,660	10,559	8,101
Lee	40,589	30,300	10,289
Leflore	47,142	16,699	30,443
Lincoln	26,759	18,407	8,352
Lowndes	46,639	28,871	17,768
Madison	32,904	9,267	23,637
Marion	23,293	15,408	7,885
Marshall	24,503	7,264	17,239
Monroe	33,953	21,932	12,021
Montgomery	13,320	7,349	5,971
Neshoba	20,927	15,026	5,901
Newton	19,517	12,950	6,567
Noxubee	16,826	4,724	12,102
Oktibbeha	26,175	14,727	11,448
Panola	28,791	12,565	16,226
Pearl River	22,411	17,221	5,190
Perry	8,745	6,333	2,412
Pike	35,063	19,655	15,408
Pontotoc	17,232	13,946	3,286
Prentiss	17,949	15,763	2,186
Quitman	21,019	7,715	13,304
Rankin	34,322	21,504	12,818
Scott	21,187	13,050	8,137
Sharkey	10,738	3,247	7,491
Simpson	20,454	13,254	7,200
Smith	14,303	11,056	3,247
Stone	7,013	5,302	1,711
Sunflower	45,750	14,730	31,020
Tallahatchie	24,081	8,580	15,501
Tate	18,138	7,696	10,442
Tippah	15,093	12,337	2,756
Tishomingo	13,889	13,210	679
Tunica	16,826	3,505	13,321
Union	18,904	15,592	3,312
Walthall	13,512	7,412	6,100
Warren	42,206	22,447	19,759
Washington	78,638	35,239	43,399
Wayne	16,258	10,449	5,809
Webster	10,580	7,938	2,642
Wilkinson	13,235	3,807	9,428
Winston	19,246	10,853	8,393
Yalobusha	12,502	6,962	5,540
Yazoo	31,653	12,862	18,791

TABLE 2
LIVE BIRTHS
MISSISSIPPI
1960

By Place of Occurrence, Reallocated to Mother's Usual Residence,
by Race, with Rates

County	Place of Occurrence			Place of Residence					
	Number		Non-white	Number			Rate*		
	Total	White		Total	White	Non-white	Total	White	Non-white
STATE	59,731	27,638	32,093	59,210	27,198	32,012	27.2	21.6	34.8
Adams	1,699	707	992	1,171	493	678	31.0	25.9	36.3
Alcorn	509	426	83	481	399	82	19.0	18.2	24.6
Amite	203	20	183	345	107	238	22.2	15.0	28.2
Attala	481	203	278	472	195	277	22.1	16.5	29.0
Benton	106	3	103	206	64	142	26.7	15.6	39.3
Bolivar	1,861	303	1,558	1,803	344	1,459	33.1	19.6	39.5
Calhoun	341	165	176	329	178	151	20.6	15.4	34.7
Carroll	193	17	176	233	45	188	20.8	9.6	28.9
Chickasaw	406	219	187	430	215	215	25.4	20.7	33.0
Choctaw	190	98	92	185	94	91	22.0	15.9	36.1
Claiborne	212	27	185	283	40	243	26.1	15.4	29.5
Clarke	167	97	70	377	171	206	22.8	17.1	31.7
Clay	457	195	262	478	214	264	25.2	23.2	27.2
Coahoma	1,514	339	1,175	1,533	313	1,220	33.2	21.4	38.6
Copiah	868	302	566	695	252	443	25.7	19.4	31.5
Covington	194	130	64	329	153	176	24.1	17.2	37.1
DeSoto	670	61	609	707	160	547	29.6	17.3	37.4
Forrest	1,502	1,123	379	1,271	803	468	24.1	21.1	31.7
Franklin	127	43	84	223	80	143	24.0	14.6	37.6
George	293	248	45	300	252	48	27.0	25.7	37.3
Greene	162	107	55	193	125	68	23.1	19.4	35.4
Grenada	522	196	326	509	185	324	27.6	19.8	35.8
Hancock	165	124	41	287	228	59	20.4	19.3	26.2
Harrison	4,121	3,370	751	3,798	3,083	715	31.8	30.8	37.1
Hinds	6,080	3,079	3,001	5,524	2,667	2,857	29.5	23.8	38.2
Holmes	761	137	624	768	129	639	28.3	17.0	32.8
Humphreys	630	109	521	632	120	512	33.1	20.8	38.4
Issaquena	57	0	57	114	14	100	31.9	11.9	41.7
Itawamba	82	64	18	300	278	22	19.9	19.6	25.2
Jackson	1,267	859	408	1,608	1,171	437	29.0	26.2	40.2
Jasper	98	18	80	427	150	277	25.2	17.8	32.6
Jefferson	96	0	96	274	38	236	27.0	15.3	30.8
Jeff Davis	295	145	150	322	110	212	23.8	18.0	28.6
Jones	2,513	1,275	1,238	1,469	967	502	24.7	21.9	32.5
Kemper	207	42	165	296	67	229	24.1	13.9	30.7
Lafayette	484	248	236	497	250	247	23.3	17.7	34.1

Table 2 - Page 2
Live Births - Mississippi - 1960

County	Place of Occurrence			Place of Residence					
	Number		Non-white	Number			Rate*		
	Total	White		Total	White	Non-white	Total	White	Non-white
Lamar	86	56	30	307	243	64	22.4	21.2	28.7
Lauderdale	2,154	1,063	1,091	1,603	842	761	23.9	19.3	32.4
Lawrence	159	39	120	268	146	122	26.2	23.0	31.6
Leake	329	149	180	423	192	231	22.7	18.2	28.5
Lee	1,240	943	297	999	693	306	24.6	22.9	29.7
Leflore	1,534	396	1,138	1,501	351	1,150	31.8	21.0	37.8
Lincoln	668	415	253	661	382	279	24.7	20.8	33.4
Lowndes	1,468	854	614	1,370	753	617	29.4	26.1	34.7
Madison	1,041	165	876	1,048	160	888	31.8	17.3	37.6
Marion	552	334	218	623	368	255	26.7	23.9	32.3
Marshall	762	123	639	808	151	657	33.0	20.8	38.1
Monroe	820	424	396	824	452	372	24.3	20.6	30.9
Montgomery	342	167	175	326	150	176	24.5	20.4	29.5
Neshoba	466	249	217	453	243	210	21.6	16.2	35.6
Newton	349	230	119	416	194	222	21.3	15.0	33.8
Noxubee	485	85	400	482	83	399	28.6	17.6	33.0
Oktibbeha	784	311	473	758	295	463	29.0	20.0	40.4
Panola	811	219	592	840	235	605	29.2	18.7	37.3
Pearl River	617	475	142	536	376	160	23.9	21.8	30.8
Perry	61	43	18	230	168	62	26.3	26.5	25.7
Pike	997	505	492	863	388	475	24.6	19.7	30.8
Pontotoc	331	249	82	344	254	90	20.0	18.2	27.4
Frentiss	427	361	66	372	310	62	20.7	19.7	28.4
Quitman	746	188	558	665	153	512	31.6	19.8	38.5
Rankin	272	73	199	837	505	332	24.4	23.5	25.9
Scott	560	338	222	526	280	246	24.8	21.4	30.2
Sharkey	252	51	201	340	65	275	31.7	20.0	36.7
Simpson	480	252	228	489	261	228	23.9	19.7	31.7
Smith	66	39	27	328	187	141	22.9	16.9	43.4
Stone	149	112	37	154	105	49	22.0	19.8	28.6
Sunflower	1,334	224	1,110	1,408	245	1,163	30.8	16.6	37.5
Tallahatchie	733	129	604	772	159	613	32.0	18.5	39.5
Tate	402	18	384	551	134	417	30.4	17.4	39.9
Tippah	393	291	102	332	242	90	22.0	19.6	32.6
Tishomingo	150	140	10	239	223	16	17.2	16.9	23.6
Tunica	616	70	546	605	67	538	36.0	19.1	40.4
Union	386	287	99	385	290	95	20.4	18.6	28.7
Walthall	523	291	232	326	128	198	24.1	17.3	32.4
Warren	1,402	596	806	1,111	508	603	26.3	22.6	30.5
Washington	2,734	1,084	1,650	2,641	1,042	1,599	33.6	29.6	36.8
Wayne	437	302	135	431	248	183	26.5	23.7	31.5
Webster	148	84	64	188	121	67	17.8	15.2	25.4
Wilkinson	432	194	238	352	75	277	26.6	19.7	29.4
Winston	377	159	218	421	177	244	21.9	16.3	29.1
Yalobusha	288	100	188	307	124	183	24.6	17.8	33.0
Yazoo	835	262	573	878	276	602	27.7	21.4	32.0
Out-of-State	1,009	842	167	1,530	1,282	248	-	-	-

Table 2 - Page 2
Live Births - Mississippi - 1960

Cities over 10,000 Population**	Place of Occurrence			Place of Residence			Rate*		
	Number		Number	Number		Total	White	Non-white	Rate*
	Total	White		Total	White				
Biloxi	2,519	2,272	247	1,593	1,393	200	36.2	36.3	35.0
Clarksdale	869	336	533	606	217	389	28.7	22.1	34.5
Cleveland	436	196	240	268	119	149	26.3	19.3	37.2
Columbus	909	513	396	769	418	351	31.0	28.2	35.3
Corinth	482	415	67	198	141	57	17.3	15.4	24.6
Greenville	1,372	686	686	1,395	693	702	33.6	32.9	34.4
Greenwood	812	392	420	583	223	360	28.5	22.7	33.9
Gulfport	1,463	1,086	377	915	643	272	30.3	27.0	42.7
Hattiesburg	1,474	1,122	352	857	493	364	24.5	20.8	32.4
Jackson	5,701	3,077	2,624	4,114	2,075	2,039	28.5	22.4	39.5
Laurel	2,468	1,251	1,217	744	387	357	26.7	21.6	35.8
McComb	685	405	280	298	179	119	24.8	20.8	34.8
Meridian	2,081	1,056	1,025	1,169	643	526	23.7	19.7	31.4
Natchez	871	649	222	766	309	457	32.2	27.0	37.0
Pascagoula	73	1	72	514	354	160	30.0	26.7	41.2
Tupelo	1,001	787	214	386	264	122	22.4	20.2	29.4
Vicksburg	1,378	596	782	759	338	421	26.0	21.8	30.9
Yazoo City	453	258	195	321	134	187	28.6	26.6	30.2
Total Urban#	36,551	21,425	15,126	22,189	12,048	10,141	28.2	24.1	35.3
Total Rural	23,180	6,213	16,967	37,021	15,150	21,871	26.6	20.0	34.5

* Rate per 1,000 population.

** Cities having 10,000 population and over according to 1960 census.

Includes all incorporated towns of 2,500 population or more according to 1960 census.

TABLE 3
LIVE BIRTHS
MISSISSIPPI
1960

By County and Cities over 10,000, by Race, Attendant, and Sex
on Basis of Usual Residence of Mother

County	TOTAL*	WHITE			NONWHITE			SEX			
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	
STATE	<u>59,210</u>	<u>27,198</u>	<u>26,931</u>	<u>135</u>	<u>132</u>	<u>32,012</u>	<u>16,418</u>	<u>1,552</u>	<u>14,042</u>	<u>29,952</u>	<u>29,258</u>
Adams	1,171	493	493	0	0	678	662	4	12	581	590
Alcorn	481	399	386	12	1	82	47	16	19	261	220
Anite	345	107	104	3	0	238	83	5	150	175	170
Attala	472	195	188	3	4	277	75	30	172	247	225
Benton	206	64	61	3	0	142	39	7	96	98	108
Bolivar	1,803	344	337	2	5	1,459	687	62	710	906	897
Calhoun	329	178	178	0	0	151	86	3	62	164	165
Carroll	233	45	42	1	2	188	45	5	138	119	114
Chickasaw	430	215	205	10	0	215	99	28	88	216	214
Choctaw	185	94	93	0	1	91	35	4	52	86	99
Claiborne	283	40	39	0	1	243	62	3	178	152	131
Clarke	377	171	168	2	1	206	175	7	24	194	183
Clay	478	214	214	0	0	264	152	31	81	235	243
Coahoma	1,533	313	310	0	3	1,220	524	76	620	774	759
Copiah	695	252	249	3	0	443	351	11	81	333	362
Covington	329	153	153	0	0	176	143	5	28	157	172
DeSoto	707	160	160	0	0	547	232	11	304	338	369
Forrest	1,271	803	803	0	0	468	351	0	117	639	632
Franklin	223	80	80	0	0	143	111	0	32	103	120
George	300	252	245	6	1	48	32	1	15	162	138
Greene	193	125	119	5	1	68	42	3	23	99	94
Grenada	509	185	183	0	2	324	43	61	220	270	239
Hancock	287	228	227	1	0	59	32	0	27	152	135
Harrison	3,798	3,083	3,065	2	16	715	389	59	267	1,962	1,836

Table 3 - Page 2
Live Births - Mississippi - 1960

County	TOTAL*	WHITE			NONWHITE			SEX		
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male
Hinds	5,524	2,667	2,667	0	0	2,857	2,299	84	474	2,878
Holmes	768	129	127	2	0	639	120	36	483	368
Humphreys	632	120	116	0	4	512	82	6	424	324
Issaquena	114	14	14	0	0	100	45	0	55	52
Itawamba	300	278	270	7	1	22	19	1	2	168
Jackson	1,608	1,171	1,167	2	2	437	249	63	125	852
Jasper	427	150	149	1	0	277	205	5	67	205
Jefferson	274	38	38	0	0	236	141	2	93	151
Jeff Davis	322	110	109	0	1	212	137	2	73	158
Jones	1,469	967	963	0	4	502	471	3	28	730
Kemper	296	67	66	0	1	229	96	0	133	154
Lafayette	497	250	246	1	3	247	47	10	190	248
Lamar	307	243	242	0	1	64	48	1	15	150
Lauderdale	1,603	842	839	2	1	761	656	27	78	818
Lawrence	268	146	141	2	3	122	66	3	53	140
Leake	423	192	188	3	1	231	119	6	106	217
Lee	999	693	691	2	0	306	195	56	55	500
Leflore	1,501	351	346	5	0	1,150	143	40	967	768
Lincoln	661	382	378	0	4	279	224	14	41	326
Lowndes	1,370	753	747	2	4	617	231	36	350	714
Madison	1,048	160	159	0	1	888	259	28	601	540
Marion	623	368	364	1	3	255	99	17	139	301
Marshall	808	151	146	0	5	657	167	67	423	415
Monroe	824	452	448	4	0	372	176	30	166	395
Montgomery	326	150	150	0	0	176	67	2	107	168
Neshoba	453	243	241	0	2	210	130	1	79	221
Newton	416	194	192	2	0	222	174	19	29	228
Noxubee	482	83	81	0	2	399	88	5	306	243
Oktibbeha	758	295	292	2	1	463	280	0	183	388
Panola	840	235	232	3	0	605	293	79	233	422

Table 3 - Page 3
Live Births - Mississippi - 1960

County	TOTAL*	WHITE			NONWHITE			SEX			
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female
Pearl River	536	376	376	0	0	160	154	1	5	271	265
Perry	230	168	166	2	0	62	48	4	10	122	108
Pike	863	388	386	1	1	475	316	2	157	430	433
Pontotoc	344	254	252	0	2	90	43	3	44	166	178
Prentiss	372	310	309	1	0	62	44	2	16	176	196
Quitman	665	153	150	1	2	512	302	18	192	317	348
Rankin	837	505	505	0	0	332	270	6	36	400	
Scott	526	280	278	1	1	246	125	1	120	257	269
Sharkey	340	65	64	0	1	275	150	0	125	152	188
Simpson	489	261	260	1	0	228	171	1	56	235	254
Smith	328	187	187	0	0	141	114	6	21	171	157
Stone	154	105	105	0	0	49	20	0	29	73	81
Sunflower	1,408	245	231	2	12	1,163	377	143	643	710	698
Tallahatchie	772	159	152	1	6	613	117	36	460	370	402
Tate	551	134	130	3	1	417	59	26	332	270	281
Tippah	332	242	231	9	2	90	45	24	21	178	154
Tishomingo	239	223	223	0	0	16	16	0	0	112	127
Tunica	605	67	64	1	2	538	162	26	350	297	308
Union	385	290	277	3	10	95	24	13	58	213	172
Walthall	326	128	127	0	1	198	135	0	63	156	170
Warren	1,111	508	508	0	0	603	443	5	155	573	538
Washington	2,641	1,042	1,039	1	2	1,599	948	77	574	1,312	1,329
Wayne	431	248	243	1	4	183	128	5	50	210	221
Webster	188	121	113	8	0	67	34	10	23	89	99
Wilkinson	352	75	75	0	0	277	97	0	180	184	168
Winston	421	177	177	0	0	244	81	2	161	212	209
Yalobusha	307	124	121	3	0	183	77	64	42	166	141
Yazoo	878	276	271	2	3	602	95	2	505	428	450

Table 3 - Page 4
Live Births - Mississippi - 1960

Cities Over 10,000 Population	TOTAL*	WHITE			NONWHITE			SEX	
		Total	Phys. in Hosp.	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male
Biloxi	1,593	1,393	1,390	1	200	146	10	44	818
Clarksdale	606	217	217	0	389	299	13	77	315
Cleveland	268	119	119	0	149	115	6	28	129
Columbus	769	418	415	1	351	121	31	199	400
Corinth	198	141	139	2	57	38	9	10	111
Greenville	1,395	693	691	1	702	423	46	233	698
Greenwood	583	223	223	0	360	64	1	295	302
Gulfport	915	643	639	0	272	136	37	99	494
Hattiesburg	857	493	493	0	364	278	0	86	436
Jackson	4,114	2,075	2,075	0	2,039	1,797	80	162	2,161
Laurel	744	387	387	0	357	346	2	9	378
McComb	298	179	179	0	119	103	0	16	156
Meridian	1,169	643	641	1	526	481	26	19	578
Natchez	766	309	309	0	457	450	2	5	372
Pascagoula	514	354	354	0	160	88	12	60	277
Tupelo	386	264	264	0	122	71	29	22	195
Vicksburg	759	338	338	0	421	306	4	111	392
Yazoo City	321	134	133	0	187	39	1	147	171

TABLE 4
FETAL DEATHS
MISSISSIPPI

**By Place of Occurrence, Relocated to Mother's Usual Residence
by Sex, Race and Attendant**

Table 4 - Page 2
Fetal Deaths - Mississippi - 1960

County	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total	Phys. in Hosp.	Other	Total	Phys. in Hosp.	Other	Male	Female	Undetermined
Hinds	384	350	219	196	22	1	131	103	18	10	79
Holmes	40	41	7	7	0	0	34	9	4	21	20
Humphreys	17	20	1	1	0	0	19	8	1	10	13
Issaquena	3	8	2	2	0	0	6	3	0	3	6
Itawamba	1	4	4	4	0	0	0	0	0	0	2
Jackson	44	54	33	33	0	0	21	20	1	0	18
Jasper	6	21	4	3	1	0	17	12	1	4	12
Jefferson	7	11	1	1	0	0	10	3	1	6	6
Jeff Davis	9	10	2	2	0	0	8	4	0	4	5
Jones	108	58	28	28	0	0	30	29	1	0	20
Kemper	10	15	1	1	0	0	14	11	1	2	4
Lafayette	20	20	2	2	0	0	18	3	0	15	8
Lamar	3	7	7	7	0	0	0	0	0	0	4
Lauderdale	71	46	11	11	0	0	35	28	4	3	20
Lawrence	5	9	3	3	0	0	6	6	0	0	3
Leake	6	12	3	3	0	0	9	4	1	4	9
Lee	50	44	32	32	0	0	12	12	0	0	11
Leflore	90	80	25	25	0	0	55	24	2	29	42
Lincoln	18	22	9	9	0	0	13	9	1	3	9
Lowndes	62	54	25	24	1	0	29	17	2	10	24
Madison	39	40	6	5	1	0	34	21	2	11	21
Marion	22	25	14	14	0	0	11	5	0	6	8
Marshall	48	37	6	6	0	0	31	16	7	8	20
Monroe	21	26	15	14	1	0	11	9	2	0	13
Montgomery	12	11	3	3	0	0	8	4	0	4	0
Neshoba	14	15	5	4	1	0	10	6	0	4	6
Newton	23	26	9	9	0	0	17	13	3	1	7
Noxubee	22	23	3	3	0	0	20	12	1	7	8
Oktibbeha	18	19	6	6	0	0	13	7	1	5	11
Panola	27	33	3	3	0	0	30	20	2	8	13
											20

Table 4 - Page 3
Fetal Deaths - Mississippi - 1960

County	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total	Phys. in Hosp.	Other	Total	Phys. in Hosp.	Other	Male	Female	Unde- termined
Pearl River	26	22	13	13	0	0	9	9	0	15	5
Perry	2	8	2	2	0	0	6	4	2	0	3
Pike	22	18	7	7	0	0	11	6	0	5	5
Pontotoc	6	5	5	5	0	0	0	0	0	8	0
Prentiss	15	12	11	10	0	1	1	1	0	0	2
Quitman	22	22	4	4	0	0	18	13	2	3	12
Rankin	13	37	24	23	1	0	13	9	0	4	7
Scott	15	15	3	3	0	0	12	7	0	5	9
Sharkey	14	13	1	1	0	0	12	6	1	5	6
Simpson	13	13	7	7	0	0	6	6	0	0	7
Smith	2	12	7	7	0	0	5	4	0	1	6
Stone	8	9	7	7	0	0	2	2	0	0	1
Sunflower	46	51	12	10	2	0	39	20	6	13	24
Tallahatchie	36	34	5	5	0	0	29	11	1	17	21
Tate	19	26	3	2	1	0	23	5	2	16	13
Tippah	22	19	11	11	0	0	8	8	0	0	5
Tishomingo	2	3	3	3	0	0	0	0	0	2	0
Tunica	29	29	2	2	0	0	27	9	3	15	13
Union	23	21	17	17	0	0	4	1	1	2	3
Walhall	26	15	6	6	0	0	9	8	0	1	4
Warren	63	39	19	19	0	0	20	15	1	4	17
Washington	216	213	112	111	1	0	101	72	9	20	62
Wayne	8	16	5	5	0	0	11	9	0	2	5
Webster	0	2	2	2	0	0	0	0	0	0	0
Wilkinson	30	26	3	3	0	0	23	10	1	12	17
Winston	10	13	0	0	0	0	13	5	3	5	7
Yalobusha	6	8	4	4	0	0	4	2	1	1	2
Yazoo	27	34	4	4	0	0	30	9	3	18	16

Table 4 - Page 4
Fetal Deaths - Mississippi - 1960

Cities Over 10,000 Population	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total Hosp.	Phys. in Hosp.	Other	Total Hosp.	Phys. in Hosp.	Other	Male	Female	Unde- termined
Biloxi	47	32	26	25	1	0	6	5	1	0	10
Clarksdale	38	18	5	5	0	0	13	7	2	4	7
Cleveland	10	10	3	3	0	0	7	5	0	2	9
Columbus	51	35	19	18	1	0	16	8	1	2	3
Corinth	5	3	1	1	0	0	2	1	0	1	1
Greenville	135	129	89	88	1	0	40	31	3	6	28
Greenwood	64	26	14	14	0	0	12	6	0	6	10
Gulfport	28	20	5	5	0	0	15	5	4	6	9
Hattiesburg	41	26	10	10	0	0	16	11	0	5	11
Jackson	375	293	191	173	17	1	102	80	17	5	60
Laurel	108	30	13	13	0	0	17	16	1	0	11
McComb	10	7	4	4	0	0	3	1	0	2	4
Meridian	66	25	8	8	0	0	17	13	3	1	10
Natchez	18	17	8	7	1	0	9	8	0	1	9
Pascagoula	1	21	13	13	0	0	8	7	1	0	8
Tupelo	43	21	15	15	0	0	6	6	0	0	6
Vicksburg	62	30	14	14	0	0	16	11	1	4	13
Yazoo City	17	15	3	3	0	0	12	3	2	7	9

TABLE 5
DEATHS
MISSISSIPPI
1960

By Cause by Race on Basis of Place of Residence

International List Number	Cause of Death*	Total	White	Non-white
<u>001-999</u>	<u>TOTAL ALL CAUSES</u>	<u>21,713</u>	<u>11,204</u>	<u>10,509</u>
<u>001-019</u>	<u>Tuberculosis, all forms</u>	<u>147</u>	<u>57</u>	<u>90</u>
002	Pulmonary	127	51	76
003	Pleural	1	0	1
005	Tracheobronchial glandular, with symptoms	1	1	0
008	Unspecified site	6	2	4
010	Meninges and central nervous system	4	1	3
011	Intestines, peritoneum, and mesenteric glands	3	1	2
012	Bones, and joints, active or unspecified	1	0	1
016	Genito-urinary system	1	1	0
019	Disseminated tuberculosis	3	0	3
<u>020-029</u>	<u>Syphilis and its sequelae</u>	<u>29</u>	<u>5</u>	<u>24</u>
020	Congenital	2	0	2
022	Aneurysm of aorta	11	3	8
023	Other cardiovascular	13	1	12
025	General paralysis of insane	1	0	1
026	Other central nervous system	1	1	0
029	Unqualified	1	0	1
030	Acute or unspecified gonorrhea	1	0	1
035	Late effects of gonococcal infection	1	0	1
041	Paratyphoid fever	1	0	1
<u>045-048</u>	<u>Dysentery, all forms</u>	<u>27</u>	<u>8</u>	<u>19</u>
045	Bacillary	10	5	5
046	Amebiasis	4	3	1
048	Unspecified forms	13	0	13
049	Food poisoning	3	1	2
<u>050,051</u>	<u>Scarlet fever and streptococcal sore throat</u>	<u>5</u>	<u>1</u>	<u>4</u>
051	Streptococcal sore throat	5	1	4
053	Septicemia and pyemia	35	20	15
055	Diphtheria	4	3	1
056	Whooping cough	7	0	7
057	Meningococcal infections	13	8	5
061	Tetanus	11	2	9
080	Acute poliomyelitis	4	2	2
082	Acute infectious encephalitis	12	9	3
085	Measles	9	4	5
087	Chickenpox	4	1	3
092	Infectious hepatitis	8	4	4
096	Other diseases attributable to viruses	4	2	2
130	Infestation with worms of other, mixed, and unspecified type	2	0	2
134	Other fungus infections	11	4	7
138	Other infective and parasitic diseases	4	3	1
<u>140-205</u>	<u>Malignant neoplasms</u>	<u>2,635</u>	<u>1,644</u>	<u>991</u>
<u>140-148</u>	<u>Buccal cavity and pharynx</u>	<u>68</u>	<u>49</u>	<u>19</u>
140	Lip	3	3	0

Table 5 - Page 2
Deaths - Mississippi - 1960

International List Number	Cause of Death*	Total	White	Non-white
141	Tongue	9	7	2
142	Salivary gland	9	6	3
143	Floor of mouth	2	1	1
144	Other parts of mouth, and mouth, unspecified	15	13	2
145	Oral mesopharynx	5	4	1
146	Nasopharynx	8	5	3
148	Pharynx, unspecified	17	10	7
<u>150-159</u>	<u>Digestive organs and peritoneum</u>	<u>812</u>	<u>508</u>	<u>304</u>
150	Esophagus	42	23	19
151	Stomach	226	99	127
152	Small intestine, including duodenum	6	5	1
153	Large intestine, except rectum	221	164	57
154	Rectum	60	34	26
155	Biliary passages and liver (stated to be primary site)	33	23	10
156	Liver (secondary and unspecified)	65	39	26
157	Pancreas	141	112	29
158	Peritoneum	4	3	1
159	Unspecified digestive organs	14	6	8
<u>160-165</u>	<u>Respiratory system</u>	<u>357</u>	<u>267</u>	<u>90</u>
160	Nose, nasal cavities, middle ear, and accessory sinuses	7	4	3
161	Larynx	20	13	7
162	Bronchus and trachea and lung specified as primary	136	112	24
163	Lung, unspecified as to whether primary or secondary	181	125	56
164	Mediastinum	3	3	0
165	Thoracic organs (secondary)	10	10	0
<u>170-181</u>	<u>Breast and genito-urinary organs</u>	<u>796</u>	<u>410</u>	<u>386</u>
170	Breast	169	112	57
171	Cervix uteri	158	54	104
172	Corpus uteri	5	3	2
173	Other parts of uterus	1	1	0
174	Uterus, unspecified	73	30	43
175	Ovary, Fallopian tube, and broad ligament	66	48	18
176	Other and unspecified female genital organs	9	5	4
177	Prostate	210	95	115
178	Testis	6	5	1
179	Other and unspecified male genital organs	3	1	2
180	Kidney	37	21	16
181	Bladder and other urinary organs	59	35	24
<u>190-199</u>	<u>Other and unspecified sites</u>	<u>370</u>	<u>243</u>	<u>127</u>
190	Melanoma of skin	23	20	3
191	Skin, other	33	29	4
192	Eye	8	6	2
193	Brain and other parts of nervous system	72	56	16
194	Thyroid gland	9	7	2
195	Other endocrine glands	1	1	0
196	Bone	25	17	8
197	Connective tissue	8	5	3

Table 5 - Page 3
 Deaths - Mississippi - 1960

International List Number	Cause of Death*	Total	White	Non-white
198	Lymph nodes (secondary and unspecified)	6	2	4
199	Other and unspecified sites	185	100	85
<u>200-205</u>	<u>Lymphatic and hematopoietic tissues</u>	<u>232</u>	<u>167</u>	<u>65</u>
200	Lymphosarcoma and reticulosarcoma	41	34	7
201	Hodgkin's disease	21	14	7
202	Other forms of lymphoma (reticulosis)	8	6	2
203	Multiple myeloma (plasmacytoma)	28	19	9
204	Leukemia and aleukemia	134	94	40
<u>210-229</u>	<u>Benign neoplasm</u>	<u>20</u>	<u>8</u>	<u>12</u>
210	Buccal cavity and pharynx	1	1	0
211	Other parts of digestive system	1	1	0
212	Respiratory system	1	0	1
214	Uterine fibromyoma	5	0	5
215	Uterus, other	1	1	0
223	Brain and other parts of nervous system	6	3	3
224	Endocrine glands	3	1	2
228	Hemangioma and lymphangioma	2	1	1
<u>230-239</u>	<u>Neoplasms of unspecified nature</u>	<u>39</u>	<u>20</u>	<u>19</u>
230	Digestive organs	2	1	1
231	Respiratory organs	3	3	0
236	Other	1	1	0
237	Brain and other parts of nervous system	24	12	12
238	Skin and musculoskeletal system	1	1	0
239	Other and unspecified organs	8	2	6
241	Asthma	74	40	34
242	Angioneurotic edema	1	1	0
245	Other allergic disorders	4	3	1
251	Non-toxic nodular goiter	1	0	1
252	Thyrotoxicosis with or without goiter	8	3	5
253	Myxedema and cretinism	2	2	0
254	Other diseases of thyroid gland	2	0	2
260	Diabetes mellitus	309	139	170
272	Diseases of pituitary gland	1	0	1
274	Diseases of adrenal glands	3	3	0
281	Pellagra	5	1	4
286	Other avitaminoses and nutritional deficiency states	29	9	20
287	Obesity, not specified as of endocrine origin	10	6	4
288	Gout	1	1	0
289	Other metabolic diseases	6	3	3
<u>290-293</u>	<u>Anemias</u>	<u>57</u>	<u>21</u>	<u>36</u>
290	Pernicious and other hyperchromic	3	2	1
292	Other specified type	38	17	21
293	Unspecified type	16	2	14
294	Polycythemia	3	2	1
296	Purpura and other hemorrhagic conditions	6	4	2
297	Agranulocytosis	1	1	0
298	Diseases of spleen	4	1	3
299	Other diseases of blood and blood-forming organs	2	0	2
300	Schizophrenic disorders	1	0	1
304	Senile psychosis	4	1	3

Table 5 - Page 4
 Deaths - Mississippi - 1960

International List Number	Cause of Death*	Total	White	Non-white
305	Presenile psychosis	2	2	0
307	Alcoholic psychosis	2	2	0
310	Anxiety reaction without mention of somatic symptoms	1	1	0
322	Alcoholism	12	7	5
323	Other drug addiction	1	1	0
325	Mental deficiency	5	1	4
<u>330-334</u>	<u>Vascular lesions affecting central nervous system</u>	<u>2,960</u>	<u>1,513</u>	<u>1,447</u>
330	Subarachnoid hemorrhage	77	43	34
331	Cerebral hemorrhage	1,832	892	940
332	Cerebral embolism and thrombosis	650	395	255
334	Other and ill-defined	401	183	218
340	Meningitis, except meningococcal and tuberculous	51	8	43
342	Intracranial and intraspinal abscess	5	3	2
343	Encephalitis, myelitis, and encephalo-myelitis, except acute infectious	10	6	4
344	Late effects of intracranial abscess or pyogenic infection	10	6	4
345	Multiple sclerosis	7	5	2
350	Paralysis agitans	21	16	5
351	Cerebral spastic infantile paralysis	15	10	5
352	Other cerebral paralysis	7	3	4
353	Epilepsy	16	3	13
355	Other diseases of brain	12	8	4
356	Motor neurone disease and muscular atrophy	13	9	4
357	Other diseases of spinal cord	1	0	1
364	Polyneuritis and polyradiculitis	4	2	2
366	Other and unspecified forms of neuralgia and neuritis	1	0	1
368	Other diseases of peripheral nerves except autonomic	1	0	1
377	Inflammation of optic nerve and retina	1	1	0
391	Otitis media without mention of mastoiditis	3	0	3
392	Otitis media with mastoiditis	1	1	0
394	Other inflammatory diseases of ear	1	0	1
<u>400-402</u>	<u>Rheumatic fever</u>	<u>9</u>	<u>3</u>	<u>6</u>
400	Without mention of heart involvement	2	0	2
401	With heart involvement	6	2	4
402	Chorea	1	1	0
<u>410-443</u>	<u>Diseases of heart</u>	<u>6,312</u>	<u>3,879</u>	<u>2,433</u>
410	Diseases of mitral valve	48	24	24
411	Diseases of aortic valve specified as rheumatic	4	4	0
414	Other endocarditis specified as rheumatic	3	1	2
415	Other myocarditis specified as rheumatic	7	2	5
416	Other heart disease specified as rheumatic	46	25	21
420	Arteriosclerotic heart disease, including coronary disease	3,973	2,870	1,103
421	Chronic endocarditis not specified as rheumatic	92	53	39

Table 5 - Page 5
 Deaths - Mississippi - 1960

International List Number	Cause of Death*	Total	White	Non-white
422	Other myocardial degeneration	507	283	224
430	Acute and subacute endocarditis	5	3	2
431	Acute myocarditis not specified as rheumatic	3	1	2
433	Functional disease of heart	61	36	25
434	Other and unspecified diseases of heart	425	195	230
440	Essential benign hypertensive heart disease	2	1	1
441	Essential malignant hypertension with heart disease	15	2	13
442	Hypertensive heart disease with arteriolar nephrosclerosis	260	96	164
443	Other and unspecified hypertensive heart disease	861	283	578
<u>444-447</u>	<u>Hypertension without mention of heart</u>	250	98	152
444	Essential benign	78	27	51
445	Essential malignant hypertension	69	13	56
446	With arteriolar nephrosclerosis	103	58	45
450	General arteriosclerosis	306	198	108
<u>451-468</u>	<u>Other diseases of circulatory system</u>	216	127	89
451	Aortic aneurysm, nonsyphilitic, and dissecting aneurysm	78	61	17
452	Other aneurysm, except of heart and aorta	7	4	3
453	Peripheral vascular disease	3	1	2
454	Arterial embolism and thrombosis	11	7	4
455	Gangrene of unspecified cause	9	0	9
456	Other diseases of arteries	7	5	2
460	Varicose veins of lower extremities	1	0	1
461	Hemorrhoids	2	2	0
462	Varicose veins of other specified sites	2	2	0
463	Phlebitis and thrombophlebitis of lower extremities	6	3	3
464	Phlebitis and thrombophlebitis of other sites	8	6	2
465	Pulmonary embolism and infarction	43	19	24
466	Other venous embolism and thrombosis	15	9	6
467	Other diseases of circulatory system	22	7	15
468	Certain diseases of lymph nodes and lymph channels	2	1	1
470	Acute nasopharyngitis (common cold)	9	3	6
471	Acute sinusitis	1	1	0
473	Acute tonsilitis	1	0	1
474	Acute laryngitis and tracheitis	1	0	1
475	Acute upper respiratory infection of multiple or unspecified sites	4	1	3
<u>480-483</u>	<u>Influenza</u>	238	71	167
480	With pneumonia	145	34	111
481	With other respiratory manifestations, and influenza unqualified	82	32	50
482	With digestive manifestations, but without respiratory symptoms	4	1	3
483	With nervous manifestations, but without digestive or respiratory symptoms	7	4	3
<u>490-493</u>	<u>Pneumonia</u>	720	318	402
490	Lobar	158	68	90

Table 5 - Page 6
Deaths - Mississippi - 1960

International List Number	Cause of Death*	Total	White	Non-white
491	Bronchopneumonia	280	134	146
492	Primary atypical	46	28	18
493	Other and unspecified	236	88	148
<u>500-502</u>	<u>Bronchitis</u>	<u>42</u>	<u>28</u>	<u>14</u>
500	Acute	12	5	7
501	Unqualified	15	9	6
502	Chronic	15	14	1
513	Chronic sinusitis	2	1	1
517	Other diseases of upper respiratory tract	4	2	2
518	Empyema	2	1	1
519	Pleurisy	3	2	1
521	Abscess of lung	3	2	1
522	Pulmonary congestion and hypostasis	26	12	14
525	Other chronic interstitial pneumonia	20	15	5
526	Bronchiectasis	25	22	3
527	Other diseases of lung and pleural cavity	115	99	16
536	Stomatitis	2	0	2
537	Diseases of salivary glands	1	1	0
538	Other diseases of buccal cavity	2	0	2
539	Diseases of esophagus	4	2	2
<u>540,541</u>	<u>Ulcer of stomach and duodenum</u>	<u>84</u>	<u>60</u>	<u>24</u>
540	Stomach	46	28	18
541	Ulcer of duodenum	38	32	6
542	Gastrojejunal ulcer	1	0	1
543	Gastritis and duodenitis	6	3	3
544	Disorders of function of stomach	3	1	2
545	Other diseases of stomach and duodenum	3	2	1
<u>550-553</u>	<u>Appendicitis</u>	<u>23</u>	<u>12</u>	<u>11</u>
550	Acute	19	11	8
551	Unqualified	3	0	3
553	Other diseases of appendix	1	1	0
<u>560,561,570</u>	<u>Hernia and intestinal obstruction</u>	<u>123</u>	<u>60</u>	<u>63</u>
560	Hernia of abdominal cavity without mention of obstruction	14	8	6
561	Hernia of abdominal cavity with obstruction	26	12	14
570	Intestinal obstruction without mention of hernia	83	40	43
571	Gastro-enteritis and colitis, except ulcerative, age 4 weeks and over	218	35	183
572	Chronic enteritis and ulcerative colitis	25	19	6
573	Functional disorders of intestines	1	1	0
575	Abscess of anal and rectal regions	2	0	2
576	Peritonitis	14	10	4
578	Other diseases of intestines and peritoneum	35	19	16
580	Acute and subacute yellow atrophy of liver	8	4	4
581	Cirrhosis of liver	96	70	26
582	Suppurative hepatitis and liver abscess	1	0	1
583	Other diseases of liver	12	7	5
584	Cholelithiasis	12	10	2
585	Cholecystitis and cholangitis, without mention of calculi	12	10	2
586	Other diseases of gallbladder and biliary ducts	13	9	4

Table 5 - Page 7
 Deaths - Mississippi - 1960

International List Number	Cause of Death*	Total	White	Non-white
587	Diseases of pancreas	34	22	12
<u>590,591</u>	Acute nephritis and nephritis with edema	<u>41</u>	<u>11</u>	<u>30</u>
590	Acute	21	7	14
591	With edema, including nephrosis	20	4	16
<u>592-594</u>	<u>Chronic and unspecified nephritis and other renal sclerosis</u>	<u>241</u>	<u>83</u>	<u>158</u>
592	Chronic nephritis	189	65	124
593	Nephritis not specified as acute or chronic	49	17	32
594	Other renal sclerosis	3	1	2
600	Infections of kidney	140	76	64
601	Hydronephrosis	3	2	1
602	Calculi of kidney and ureter	12	9	3
603	Other diseases of kidney and ureter	13	4	9
605	Cystitis	3	1	2
606	Other diseases of bladder	3	2	1
607	Urethritis (nonvenereal)	1	0	1
608	Stricture of urethra	13	2	11
609	Other diseases of urethra	9	5	4
610	Hyperplasia of prostate	76	27	49
611	Prostatitis	10	3	7
612	Other diseases of prostate	1	1	0
614	Orchitis and epididymitis	2	1	1
621	Other diseases of breast	2	1	1
626	Diseases of parametrium and pelvic peritoneum (female)	2	1	1
630	Infective disease of uterus, vagina, and vulva	2	0	2
633	Other diseases of uterus	2	0	2
<u>640-689</u>	<u>Diseases of pregnancy, childbirth, and puerperium</u>	<u>59</u>	<u>8</u>	<u>51</u>
642	Toxemias of pregnancy	31	2	29
648	Other complications arising from pregnancy	4	1	3
650	Abortion without mention of sepsis or toxemia	1	0	1
651	Abortion with sepsis	3	1	2
660	Delivery without mention of complication	2	1	1
670	Delivery complicated by placenta previa or antepartum hemorrhage	2	0	2
671	Delivery complicated by retained placenta	2	0	2
672	Delivery complicated by other postpartum hemorrhage	3	0	3
674	Delivery complicated by disproportion or mal-position of fetus	2	0	2
675	Delivery complicated by prolonged labor of other origin	1	0	1
677	Delivery with other trauma	2	2	0
678	Delivery with other complications of childbirth	1	0	1
681	Sepsis of childbirth and the puerperium	3	1	2
685	Puerperal eclampsia	1	0	1
688	Other and unspecified complications of the puerperium	1	0	1
690	Boil and carbuncle	2	0	2

Table 5 - Page 8
Deaths - Mississippi - 1960

International List Number	Cause of Death*	Total	White	Non-white
692	Other cellulitis and abscess without mention of lymphangitis	5	3	2
698	Other local infections of skin and subcutaneous tissue	2	2	0
703	Other dermatitis	1	0	1
704	Pemphigus	2	2	0
705	Erythematous conditions	5	3	2
706	Psoriasis and similar disorders	1	1	0
710	Other hypertrophic and atrophic conditions of skin	2	1	1
711	Other dermatoses	1	1	0
714	Diseases of sweat and sebaceous glands	1	0	1
715	Chronic ulcer of skin	4	1	3
720	Acute arthritis due to pyogenic organisms	1	1	0
722	Rheumatoid arthritis and allied conditions	16	14	2
723	Osteo-arthritis (arthrosis) and allied conditions	3	2	1
724	Other specified forms of arthritis	1	1	0
725	Arthritis, unspecified	5	2	3
730	Osteomyelitis and periostitis	1	0	1
733	Other diseases of bone	1	0	1
744	Other diseases of muscle, tendon, and fascia	10	9	1
745	Curvature of spine	1	0	1
<u>750-759</u>	<u>Congenital malformations</u>	<u>234</u>	<u>142</u>	<u>92</u>
750	Monstrosity	11	8	3
751	Spina bifida and meningocele	12	10	2
752	Congenital hydrocephalus	18	7	11
753	Other nervous system and sense organs	9	4	5
754	Circulatory system	119	72	47
756	Digestive system	17	10	7
757	Genito-urinary system	19	13	6
758	Bone and joint	5	4	1
759	Other and unspecified, not elsewhere classified	24	14	10
<u>760-762</u>	<u>Birth injuries, postnatal asphyxia and atelectasis</u>	<u>332</u>	<u>186</u>	<u>146</u>
760	Intracranial and spinal injury at birth	47	21	26
761	Other birth injury	59	29	30
762	Postnatal asphyxia and atelectasis	226	136	90
<u>763-768</u>	<u>Infections of newborn</u>	<u>127</u>	<u>36</u>	<u>91</u>
763	Pneumonia	79	24	55
764	Diarrhea	26	2	24
767	Umbilical sepsis	3	0	3
768	Other sepsis	19	10	9
<u>769-776</u>	<u>Other diseases peculiar to early infancy, and immaturity, unqualified</u>	<u>847</u>	<u>243</u>	<u>604</u>
769	Neonatal disorders arising from maternal toxemia	19	8	11
770	Hemolytic disease of newborn (erythroblastosis)	26	16	10
771	Hemorrhagic disease of newborn	14	6	8

Table 5 - Page 9
 Deaths - Mississippi - 1960

International List Number	Cause of Death*	Total	White	Non-white
772	Nutritional maladjustment	37	5	32
773	Ill-defined diseases peculiar to early infancy	159	66	93
774	Immaturity with mention of any other subsidiary condition	26	7	19
776	Immaturity, unqualified	566	135	431
<u>780-795</u>	<u>Symptoms, senility, and ill-defined conditions</u>	<u>1,852</u>	<u>428</u>	<u>1,424</u>
780	Certain symptoms referable to nervous system and special senses	4	1	3
782	Symptoms referable to cardiovascular and lymphatic system	35	8	27
783	Symptoms referable to respiratory system	3	2	1
784	Symptoms referable to upper gastro-intestinal tract	6	4	2
785	Symptoms referable to abdomen and lower gastro-intestinal tract	6	2	4
786	Symptoms referable to genito-urinary system	3	1	2
788	Other general symptoms	5	0	5
789	Abnormal urinary constituents of unspecified cause	1	0	1
790	Nervousness and debility	1	1	0
791	Headache	1	0	1
792	Uremia, unqualified	23	10	13
794	Senility without mention of psychosis	62	20	42
795	Ill-defined and unknown causes	1,702	379	1,323
<u>800-962</u>	<u>Accidents</u>	<u>1,504</u>	<u>827</u>	<u>677</u>
800-802	Railway	18	8	10
<u>810-835</u>	<u>Motor vehicle</u>	<u>620</u>	<u>428</u>	<u>192</u>
810	Collision with railway train, traffic	20	16	4
812	Traffic, to pedestrian	101	51	50
813	Traffic, to pedal cyclist	5	4	1
815	Traffic, to rider or passenger of motorcycle in collision with other motor vehicle	4	4	0
816	Other traffic, two or more motor vehicles	236	184	52
818	Traffic, collision with animal or animal-drawn vehicle	1	1	0
819	Traffic, collision with fixed or unspecified object	26	22	4
821	Noncollision traffic, to rider of motorcycle	1	0	1
822	Traffic, overturning in roadway	20	11	9
823	Traffic, running off roadway	140	98	42
824	Other noncollision traffic	15	5	10
825	Traffic, unspecified nature	33	22	11
830	Nontraffic, to pedestrian	6	4	2
835	Nontraffic, other and unspecified nature	12	6	6
845	Other nonmotor road vehicle	4	3	1
850	Submersion of occupant of small boat	32	18	14
856	Machinery accident in water transport	1	1	0
860	Accident to personnel in military aircraft	3	2	1
866	Other and unspecified aircraft	2	2	0

Table 5 - Page 10
Deaths - Mississippi - 1960

International List Number	Cause of Death*	Total	White	Non-white
32	Poisoning by aspirin and salicylates	2	1	1
93	Poisoning by other and unspecified drugs	3	3	0
880	Poisoning by alcohol	1	0	1
19	Poisoning by petroleum products	3	1	2
431	Poisoning by utility (illuminating) gas	9	4	5
424	Poisoning by motor vehicle exhaust gas	7	4	3
892	Poisoning by other carbon monoxide gas	1	0	1
3	Poisoning by unspecified gases and vapors	1	1	0
900	Fall on stairs	6	6	0
27	Fall from ladders	1	0	1
1	Other falls from one level to another	29	19	10
903	Fall on same level	15	12	3
2	Unspecified falls	79	60	19
910	Blow from falling or projected object	28	12	16
4	Caused by machinery	26	19	7
2	Caused by cutting and piercing instruments	6	1	5
5	Caused by electric current	16	13	3
915	Caused by explosion of pressure vessel	2	2	0
916	Caused by fire and explosion of combustible material	235	60	175
1	Caused by hot substance, corrosive liquid, and steam	6	1	5
13	Caused by firearm	92	46	46
42	Inhalation and ingestion of food causing obstruction or suffocation	22	11	11
323	Inhalation and ingestion of other object causing obstruction or suffocation	5	3	2
677	Foreign body entering other orifice	1	1	0
10	Mechanical suffocation in bed and cradle	35	9	26
192	Mechanical suffocation in other and unspecified circumstances	6	2	4
4	Lack of care of infants under 1 year of age	1	0	1
50	Caused by bites and stings of venomous animals and insects	1	1	0
1	Other accidents caused by animals	3	0	3
0	Drowning and submersion	115	44	71
52	Excessive heat and insolation	5	2	3
0	Excessive cold	8	2	6
9	Hunger, thirst, and exposure	9	3	6
42	Lightning	5	0	5
10	Other and unspecified accidents	18	10	8
11	Therapeutic misadventure in surgical treatment	5	3	2
2	Therapeutic misadventure in infusion or trans-fusion	3	0	3
6	Therapeutic misadventure in administration of drugs or biologicals	2	1	1
1	Therapeutic misadventure in anesthesia	2	2	0
14	Late complication of surgical operation	2	1	1
0				

Table 5 - Page 11
 Deaths - Mississippi - 1960

International List Number	Cause of Death*	Total	White	Non-white
960	Late effect of motor vehicle accident	3	2	1
962	Late effect of other accidental injury	5	3	2
<u>963,970-979</u>	<u>Suicide</u>	<u>153</u>	<u>131</u>	<u>22</u>
970	By analgesic and soporific substances	3	3	0
971	By other solid and liquid substances	6	5	1
974	By hanging and strangulation	16	14	2
975	By submersion (drowning)	6	5	1
976	By firearms and explosives	117	101	16
977	By cutting and piercing instruments	3	2	1
978	By jumping from high place	2	1	1
<u>964,980-985</u>	<u>Homicide</u>	<u>187</u>	<u>36</u>	<u>151</u>
981	Assault by firearm and explosive	121	27	94
982	Assault by cutting and piercing instruments	39	3	36
983	Assault by other means	21	5	16
984	Injury by intervention of police	5	1	4
985	Execution	1	0	1

* When cause of death and International List Number are omitted, no deaths were reported.

TABLE 6
DEATHS
MISSISSIPPI
1960

By Place of Occurrence, Reallocated to Place of Residence,
by Race, with Rates

County	Place of Occurrence			Place of Residence					
	Number		Non-white	Number		Non-white	Rate*		Non-white
	Total	White		Total	White		Total	White	
STATE	21,583	11,136	10,447	21,713	11,204	10,509	10.0	8.9	11.4
Adams	458	174	284	395	148	247	10.5	7.8	13.2
Alcorn	250	204	46	268	215	53	10.6	9.8	15.9
Amite	92	34	58	161	81	80	10.3	11.4	9.5
Attala	212	105	107	234	121	113	11.0	10.3	11.8
Benton	48	29	19	67	41	26	8.7	10.0	7.2
Bolivar	542	106	436	573	145	428	10.5	8.3	11.6
Calhoun	148	105	43	178	135	43	11.2	11.6	9.9
Carroll	65	26	39	103	43	60	9.2	9.2	9.2
Chickasaw	184	122	62	191	123	68	11.3	11.8	10.4
Choctaw	95	64	31	115	78	37	13.6	13.2	14.7
Claiborne	114	22	92	153	31	122	14.1	11.9	14.8
Clarke	155	90	65	198	116	82	12.0	11.6	12.6
Clay	180	92	88	192	96	96	10.1	10.4	9.9
Coahoma	511	126	385	531	118	413	11.5	8.1	13.1
Copiah	278	132	146	325	162	163	12.0	12.5	11.6
Covington	98	64	34	137	85	52	10.0	9.6	11.0
DeSoto	172	52	120	215	81	134	9.0	8.8	9.2
Forrest	538	368	170	525	312	213	10.0	8.2	14.4
Franklin	90	47	43	102	57	45	11.0	10.4	11.8
George	73	63	10	92	80	12	8.3	8.2	9.3
Greene	70	49	21	82	61	21	9.8	9.5	10.9
Grenada	217	110	107	214	97	117	11.6	10.4	12.9
Hancock	107	92	15	119	101	18	8.5	8.6	8.0
Harrison	1,025	775	250	902	667	235	7.5	6.6	12.2
Hinds	2,164	1,166	998	1,462	765	697	7.8	6.8	9.3
Holmes	299	83	216	327	93	234	12.1	12.2	12.0
Humphreys	175	53	122	202	60	142	10.6	10.4	10.6
Issaquena	42	13	29	62	15	47	17.3	12.8	19.6
Itawamba	88	82	6	110	101	9	7.3	7.1	10.3
Jackson	334	247	87	359	261	98	6.5	5.8	9.0
Jasper	92	40	52	161	78	83	9.5	9.3	9.8
Jefferson	86	13	73	115	24	91	11.3	9.6	11.9
Jeff Davis	113	51	62	134	61	73	9.9	10.0	9.8
Jones	717	447	270	557	391	166	9.4	8.9	10.7
Kemper	96	32	64	120	45	75	9.8	9.3	10.1
Lafayette	181	102	79	193	109	84	9.0	7.7	11.6

Table 6 - Page 2
Deaths - Mississippi - 1960

County	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Lamar	69	61	8	111	95	16	8.1	8.3	7.2
Lauderdale	961	586	375	777	442	335	11.6	10.1	14.3
Lawrence	102	52	50	135	79	56	13.2	12.4	14.5
Leake	130	81	49	176	108	68	9.4	10.2	8.4
Lee	419	301	118	404	288	116	10.0	9.5	11.3
Leflore	572	201	371	532	158	374	11.3	9.5	12.3
Lincoln	273	180	93	278	179	99	10.4	9.7	11.8
Lowndes	422	206	216	418	198	220	9.0	6.8	12.4
Madison	323	86	237	339	81	258	10.3	8.7	10.9
Marion	200	132	68	219	146	73	9.4	9.5	9.2
Marshall	271	96	175	260	83	177	10.6	11.4	10.3
Monroe	336	200	136	359	214	145	10.6	9.8	12.1
Montgomery	120	60	60	136	72	64	10.2	9.8	10.7
Neshoba	141	93	48	192	133	59	9.2	8.8	10.0
Newton	153	106	47	184	120	64	9.4	9.3	9.7
Noxubee	195	45	150	199	52	147	11.8	11.0	12.1
Oktibbeha	189	93	96	202	93	109	7.7	6.3	9.5
Panola	256	104	152	309	140	169	10.7	11.1	10.4
Pearl River	202	142	60	236	169	67	10.5	9.8	12.9
Perry	44	28	16	89	61	28	10.2	9.6	11.6
Pike	377	211	166	383	204	179	10.9	10.4	11.6
Pontotoc	179	148	31	195	162	33	11.3	11.6	10.0
Prentiss	200	169	31	186	158	28	10.4	10.0	12.8
Quitman	205	52	153	239	68	171	11.4	8.8	12.8
Rankin	395	215	180	261	164	97	7.6	7.6	7.6
Scott	191	111	80	218	127	91	10.3	9.7	11.2
Sharkey	111	22	89	131	32	99	12.2	9.8	13.2
Simpson	247	150	97	222	150	72	10.8	11.3	10.0
Smith	69	57	12	135	110	25	9.4	9.9	7.7
Stone	56	45	11	67	51	16	9.6	9.6	9.4
Sunflower	387	98	289	435	130	305	9.5	8.8	9.8
Tallahatchie	227	55	172	259	80	179	10.8	9.3	11.5
Tate	110	45	65	160	77	83	8.8	10.0	7.9
Tippah	142	119	23	148	128	20	9.8	10.4	7.2
Tishomingo	106	94	12	131	123	8	9.4	9.3	11.8
Tunica	209	32	177	223	31	192	13.2	8.8	14.4
Union	187	141	46	219	170	49	11.6	10.9	14.8
Walthall	115	71	44	107	63	44	7.9	8.5	7.2
Warren	722	260	462	541	216	325	12.8	9.6	16.4
Washington	801	246	555	841	226	615	10.7	6.4	14.2
Wayne	135	83	52	163	93	70	10.0	8.9	12.0
Webster	79	70	9	105	88	17	9.9	11.1	6.4
Wilkinson	172	77	95	157	56	101	11.9	14.7	10.7
Winston	196	130	66	216	146	70	11.2	13.4	8.3
Yalobusha	135	91	44	167	115	52	13.4	16.5	9.4
Yazoo	343	111	232	375	128	247	11.8	10.0	13.1
Out-of-State	812	613	199	682	545	137	-	-	-

Table 6 - Page 3
Deaths - Mississippi - 1960

Cities over 10,000 Population**	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Biloxi	302	237	65	299	228	71	6.8	5.9	12.4
Clarksdale	295	108	187	239	81	158	11.3	8.2	14.0
Cleveland	83	53	30	106	52	54	10.4	8.4	13.5
Columbus	290	154	136	233	110	123	9.4	7.4	12.4
Corinth	176	136	40	140	97	43	12.2	10.6	18.6
Greenville	479	190	289	436	142	294	10.5	6.7	14.4
Greenwood	343	137	206	242	96	146	11.8	9.8	13.8
Gulfport	385	260	125	266	168	98	8.8	7.0	15.4
Hattiesburg	456	312	144	345	184	161	9.9	7.7	14.3
Jackson	1,900	1,078	822	1,054	597	457	7.3	6.4	8.8
Laurel	599	353	246	274	158	116	9.8	8.8	11.6
McComb	197	137	60	141	94	47	11.7	10.9	13.8
Meridian	847	525	322	569	331	238	11.5	10.2	14.2
Natchez	274	143	131	284	111	173	11.9	9.7	14.0
Pascagoula	55	38	17	106	78	28	6.2	5.9	7.2
Tupelo	244	170	74	142	91	51	8.2	7.0	12.3
Vicksburg	617	226	391	395	168	227	13.6	10.8	16.6
Yazoo City	186	73	113	150	49	101	13.3	9.7	16.3
Total Urban#	11,175	6,382	4,793	7,948	4,222	3,726	10.1	8.4	13.0
Total Rural	10,408	4,754	5,654	13,765	6,982	6,783	9.9	9.2	10.7

* Rate per 1,000 population.

** Cities having 10,000 population and over according to 1960 census.

Includes all incorporated towns of 2,500 population or more according to 1960 census.

TABLE 7
DEATHS FROM SELECTED CAUSES
MISSISSIPPI
1960

By Age, Race, and Sex, on Basis of Place of Residence, with Rates

Inter-national List Number	Cause of Death	Total Rate*	Race by Race	Total by Sex and Race	Total by Sex and Race	Age in Years								
						1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	
001-999	All causes	21,713	10.0	W 11,204 M 10,509	M 6,606 F 4,598	417 297	65 51	39 45	31 20	160 62	302 158	708 274	1,104 493	3,621 3,148
001-138	Infective and parasitic diseases	342	15.7	W 134 M 208	M 85 F 49	10 5	5 4	3 1	1 1	3 5	1 3	10 13	11 12	30 23
001-019	Tuberculosis (all forms)	147	6.7	W 90 M 35	M 41 M 55	0 0	0 2	0 0	0 1	0 1	0 3	6 8	5 12	19 17
001-008	Tuberculosis (respiratory system)	135	6.2	W 54 M 81	M 40 M 51	0 0	0 0	0 0	0 1	0 1	1 3	6 8	5 12	18 17
020-029	Syphilis & sequelae	29	1.3	C 24 W 9	M 4 M 15	0 2	0 0	0 0	0 1	0 1	0 3	0 0	0 0	3 5

Table 7 - Page 2
Deaths from Selected Causes - Mississippi - 1960

Inter-national List Number	Cause of Death	Total	Rate*	Race	Total by Sex and Race	Total by Sex and Race	Age in Years	Unknown	
								65 Over	1 Under
045-048	Dysentery (all forms)	27	1.2	W	8	M	3	0	0
				C	19	M	5	1	1
						F	11	5	2
							7	1	0
050,051	Scarlet fever and streptococcal sore throat	5	0.2	W	1	M	1	0	0
				C	4	F	2	0	0
							0	1	0
							1	0	1
055	Diphtheria	4	0.2	W	3	M	1	0	0
				C	1	F	0	0	0
							0	0	0
056	Whooping cough	7	0.3	W	0	F	0	0	0
				C	7	M	4	3	2
						F	3	2	1
057	Meningococcal infections	13	0.6	W	8	M	4	2	0
				C	5	F	4	0	1
							1	0	0
080	Acute poliomyelitis	4	0.2	W	2	F	1	0	0
				C	2	M	0	1	0
085	Measles	9	0.4	W	4	F	3	1	2
				C	5	M	1	0	0
						F	4	1	1

Table 7 - Page 3
Deaths from Selected Causes - Mississippi - 1960

International List Number	Cause of Death	Total	Rate*	Race	Total by Race	Sex	Total by Sex and Race	Age in Years								
								5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 Over	Unknown
140-205	Malignant neoplasms, including neoplasms of lymphatic and hematopoietic tissues	2,635	121.0	W	1,644	M	890	6	4	9	13	40	114	241	452	0
				C	991	F	754	4	3	2	6	15	51	134	441	0
210-239	Benign and unspecified neoplasms	59	2.7	C	28	M	505	2	1	1	2	1	1	1	1	0
				W	31	F	486	0	0	0	0	0	0	0	0	0
240-289	Allergic, endocrine system, metabolic & nutritional diseases	456	20.9	C	245	M	22	6	12	19	15	106	97	17	65	0
				W	211	F	148	0	0	0	0	0	0	0	0	0
260	Diabetes mellitus	309	14.2	C	170	F	105	0	0	0	0	0	0	0	0	0
				W	139	M	106	0	0	0	0	0	0	0	0	0
290-299	Diseases of the blood and blood-forming organs	73	3.4	C	44	M	58	2	0	3	5	11	24	2	1	0
				W	29	F	148	0	0	0	0	0	0	0	0	0
290-293	Anemias	57	2.6	C	36	F	112	2	0	0	0	0	0	0	0	0
				W	21	M	8	0	0	0	0	0	0	0	0	0
300-326	Mental, psychoneurotic & personality disorders	28	1.3	C	13	M	15	11	4	5	3	1	1	1	3	0
				W	15	F	6	0	0	0	0	0	0	0	0	0

Table 7 - Page 4
Deaths from Selected Causes - Mississippi - 1960

International List Number	Cause of Death	Total	Rate*	Race	Total by Race	Total by Sex and Race	Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 and over	Unknown	Age in Years			
330-334	Vascular lesions affecting central nervous system	2,960	135.9	C	1,447	M 750 F 763	0	1	1	0	1	2	3	5	39	82	620	0	0	0	
340-398	Other diseases of the nervous system and sense organs	180	8.3	C	99	M 81 F 45	36	9	4	1	2	2	2	2	2	1	1	13	0	0	0
340	Nonmeningococcal meningitis	83	3.8	C	55	M 28 F 26	14	4	2	0	0	1	1	0	0	1	1	5	0	0	0
400-468	Diseases of the circulatory system	7,093	325.6	C	2,788	M 4,305 F 1,713	2,592	0	1	0	1	3	4	4	4	86	270	483	1,741	0	0
400-402	Rheumatic fever	9	0.4	C	6	M 3 F 2	1	0	0	0	0	1	0	0	0	0	0	174	1,435	0	0
410-416	Chronic rheumatic heart disease	108	5.0	C	52	M 56 F 28	3	0	0	0	0	2	1	0	0	0	0	138	266	1,030	3
420-422	Arteriosclerotic and degenerative heart disease	4,572	209.9	C	1,366	M 766 F 600	2,012 1,194	0	0	0	1	1	0	0	1	1	0	216	216	809	1

Table 7 - Page 5
Deaths from Selected Causes - Mississippi - 1960

Inter-national List Number	Cause of Death	Total	Rate*	Total by Race	Total by Sex and Race	M F	Age in Years															
								5-9	10-14	15-24	25-34	35-44	45-54	55-64	65-74	Over 75	Unknown					
430-434	Other diseases of the heart	494	22.7	C	259	M F	235	W	151	0	0	0	4	12	33	101	0					
									84	1	0	0	1	6	9	63	0					
									143	2	1	0	3	7	16	22	91	0				
									116	2	0	2	1	3	6	17	67	0				
440-443	Hypertension with heart disease	1,138	52.2	C	756	M F	382	W	172	0	0	0	0	3	5	24	140	0				
									210	0	0	0	0	0	1	5	14	190	0			
									394	0	0	0	0	1	13	34	77	269	0			
									362	0	0	0	0	1	3	21	40	64	233	0		
444-447	Hypertension without mention of heart	250	11.5	C	152	M F	98	W	53	0	0	0	0	1	2	5	7	38	0			
									83	0	0	0	0	1	4	15	11	7	41	0		
									69	0	0	0	0	1	3	11	15	16	23	0		
470-527	Diseases of the respiratory system	1,216	55.8	C	638	M F	578	W	348	27	7	3	0	4	7	5	21	217	0			
									230	11	11	2	1	3	4	16	168	0	129	0		
									353	120	38	1	1	6	5	20	29	110	0	110	0	
									285	95	18	1	1	3	4	5	14	19	110	0	110	0
480-483	Influenza	238	10.9	C	167	M F	71	W	33	0	0	0	0	1	1	1	1	7	23	0		
									74	17	4	0	2	2	2	4	1	4	20	0		
									93	22	7	2	0	2	3	2	6	8	41	0		
490-493	Pneumonia (excluding pneumonia of newborn)	720	33.0	C	402	M F	318	W	178	20	4	3	0	4	5	2	11	20	109	0		
									140	7	7	1	1	2	1	3	2	7	109	0		
									235	93	29	1	1	4	0	3	11	23	70	0		
									167	66	8	4	1	5	2	3	6	10	62	0		
500-502	Bronchitis	42	1.9	C	14	M F	28	W	20	3	2	0	0	0	1	1	5	8	0	0		
									9	2	1	0	0	0	0	0	0	5	0	0		
									5	3	2	0	0	0	0	0	0	4	0	0		
									14	3	2	0	0	0	0	0	0	0	0	0		

Age in Years
Total by

Inter-national List Number	Cause of Death	Total	Rate*	Race	Total by Race	Total by Sex and Race	$\frac{M}{F}$	$\frac{W}{M}$	Age in Years		Unknown over 65		
									6-5	10-14	15-24		
<u>530-587</u>	<u>Diseases of the digestive system</u>	<u>735</u>	<u>33.7</u>	<u>C</u>	<u>357</u>	<u>W</u>	<u>$\frac{14}{12}$</u>	<u>$\frac{202}{155}$</u>	<u>M</u>	<u>$\frac{1}{2}$</u>	<u>$\frac{3}{1}$</u>	<u>$\frac{38}{21}$</u>	<u>$\frac{94}{85}$</u>
540, 541	Ulcer of stomach and duodenum	84	3.8	C	378	W	$\frac{81}{87}$	$\frac{192}{186}$	M	1	0	$\frac{21}{20}$	$\frac{38}{34}$
550-553	Appendicitis	23	1.0	C	357	M	$\frac{11}{12}$	$\frac{202}{155}$	F	0	0	$\frac{16}{11}$	$\frac{34}{34}$
560, 561, 570	Intestinal obstruction and hernia	123	5.6	C	378	M	$\frac{7}{8}$	$\frac{192}{186}$	F	0	0	$\frac{21}{20}$	$\frac{38}{34}$
543, 571, 572	Gastritis, duodenitis, enteritis & colitis, except diarrhea of the newborn	249	11.4	C	357	M	$\frac{1}{1}$	$\frac{29}{29}$	F	0	0	$\frac{24}{24}$	$\frac{34}{34}$
581	Cirrhosis of liver	96	4.4	C	192	M	$\frac{1}{1}$	$\frac{86}{86}$	F	0	0	$\frac{16}{16}$	$\frac{34}{34}$
590-637	Diseases of the genito-urinary system	576	26.4	C	192	M	$\frac{1}{1}$	$\frac{106}{106}$	F	0	0	$\frac{14}{14}$	$\frac{34}{34}$
					229	W	$\frac{142}{87}$	$\frac{227}{120}$				$\frac{19}{19}$	$\frac{63}{63}$
					347	F	$\frac{87}{120}$	$\frac{227}{120}$				$\frac{12}{12}$	$\frac{147}{147}$

Table 7 - Page 7
Deaths from Selected Causes - Mississippi - 1960

International List Number	Cause of Death	Total	Rate*	Race	Total by Race	Sex	Total by Sex and Race	Age in Years									
								1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 & Over	Unknown
590-594	Nephritis and nephrosis	282	12.9	W	94	M	49	0	0	1	1	2	2	3	6	34	0
				C	188	F	45	0	0	1	1	2	3	5	30	0	
610	Hyperplasia of prostate	76	3.5	W	27	M	27	0	0	1	1	2	2	6	57	0	
	Complications of pregnancy, child-birth and the puerperium			C	49	F	86	0	0	0	0	0	0	48	0		
640-689*		59	10.0	W	51	M	49	0	0	0	0	0	0	0	0	0	
690-716	Diseases of the skin & cellular tissue	26	1.2	W	14	M	9	5	5	6	6	17	12	5	54	41	39
720-749	Diseases of the bones & organs of movement	39	1.8	W	29	M	12	5	5	5	5	10	60	42	278	187	492
				C	10	F	5	5	5	5	5	5	5	5	5	5	30
750-759	Congenital malformations	234	10.7	W	142	M	92	46	46	46	46	46	46	46	46	46	46
760-776*	Certain diseases of early infancy	1,306	22.0	W	465	M	841	841	841	841	841	841	841	841	841	841	841

Table 7 - Page 8
Deaths from Selected Causes

Table 7 - Page 8
Deaths from Selected Causes - Mississippi - 1960

International List Number	Cause of Death	Total Rate*	Total by Race	Total by Sex and Race	Age in Years										Unknown	
					1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84		
780-795	Symptoms, senility, and ill-defined conditions	1,852	85.0	W 428 C 1,424	M 298 F 130	13/14	2/1	1/1	2/1	8/0	14/6	52/11	60/18	144/78	1/0	
					M 790 F 634	158/122	24/19	10/3	5/1	10/9	1/5	73/48	79/59	393/342	1/0	
800-999	Accidents, poisonings and violence	1,844	84.6	W 994 C 850	M 721 F 273	11/12	21/10	20/11	13/6	105/33	125/33	74/24	114/85	0/0	0/0	
					M 572 F 278	36/39	39/37	37/27	27/11	84/21	73/22	45/22	54/22	56/22	0/0	
810-835	Motor vehicle accidents	620	28.5	C 192	M 428 F 120	2/2	7/7	1/1	1/1	90/44	28/44	17/42	14/21	13/41	0/0	
					M 143 F 49	2/3	10/3	5/5	5/5	30/15	18/15	20/17	10/17	13/13	0/0	
800-802, 840-962	All other accidents	884	40.6	C 485	M 276 F 187	9/10	12/10	15/11	12/11	30/27	28/30	46/46	29/29	49/49	0/0	
					M 399 F 34	3/34	4/34	4/34	4/34	9/10	9/10	6/7	7/7	69/69	0/0	
963, 970-979	Suicide	153	7.0	C 22	M 111 F 20	0/0	0/0	0/0	0/0	4/1	17/11	32/15	22/15	21/21	0/0	
					M 19 F 3	0/3	0/0	0/0	0/0	1/0	2/1	6/6	3/3	2/2	0/0	
964, 980-985	Homicide	187	8.6	C 151	M 112 F 39	0/1	1/1	0/0	0/0	1/0	30/14	22/14	24/6	7/5	3/0	0/0
					F 1	0/1	1/1	0/1	0/1	1/1	1/1	0/1	5/5	1/1	0/0	0/0

* Crude death rate per 1,000 population. Specific death rates per 10,000 live births. Rates for infant causes (760-776) per 1,000 live births. Maternal death rates per 100,000 population.

TABLE 8
DEATHS UNDER 1 YEAR
MISSISSIPPI
1960

For Selected Causes, by Race, by Age, on Basis of Place of Residence, with Rates

Cause of Death and International List Number	Total	Rate*	Total by Race	DAYS												MONTHS						
				Under 1 day	1- 6	7- 13	20	27	14- 21	28 days	Under 1 day	1	2	3	4	5	6	7	8	9	10	11
<u>ALL CAUSES</u>	<u>2,457</u>	<u>41.5</u>	W 714 C 1,743	277 426	212 321	35 71	16 58	10 58	550 947	30 200	27 132	24 104	17 45	14 47	8 45	19 47	5 33	9 28	3 31	3 31	8 12	
Syphilis (020-029)	2	0.03	W 0 C 2	0 1	0 0	0 0	0 0	0 0	0 1	0 1	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	
Diphtheria (055)	0	-	W 0 C 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	
Whooping cough (056)	5	0.08	W 5 C 5	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	
Measles (085)	2	0.03	W 1 C 1	0 0	0 0	0 0	0 0	1 1	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	
Influenza (480-483)	41	0.7	W 39 C 39	2 1	0 2	0 0	0 1	4 0	8 0	6 0	6 0	6 0	6 0	6 0	6 0	3 0	4 0	3 0	1 0	1 0	1 0	1 0
**Pneumonia (490-493)	186	3.1	W 159 C 159	27	0	0	0	0	0	0	5	5	6	1	2	1	2	1	1	1	1	2
Other respiratory diseases (470-475, 500-527)	26	0.4	W 9 C 17	0 1	0 1	1 0	0 0	0 0	1 0	0 3	1 5	3 5	3 1	1 0	2 3	1 3	3 1	0 3	0 1	2 0	0 0	0 0
**Diarrhea and enteritis (5710)	162	2.7	W 12 C 150	0	0	0	0	0	0	0	0	0	0	0	1	1	1	2	0	1	1	0
Congenital malfor- mations (750-759)	164	2.8	W 95 C 69	27 12	23 10	6 5	2 6	5 3	63 36	9 7	4 3	6 4	2 1	1 1	1 1	2 1	3 4	1 4	2 3	1 4	2 1	1 2
Birth injuries (760-761)	106	1.8	W 50 C 56	30 29	17 21	2 3	1 1	0 1	50 54	0 2	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	

Table 8 - Page 2
Deaths Under 1 Year - Mississippi - 1960

Cause of Death and International List Number	Total	Rate*	Total by Race	DAYS							MONTHS							
				Under 1 day	1-	7-	14-	21-	Under 28 days	1	2	3	4	5	6	7	8	9
Postnatal asphyxia & atelectasis (762)	226	3.8	W C	136 90	72 46	59 37	3 1	0 2	1 0	135 86	1 0	0 1	0 3	0 0	0 0	0 0	0 0	0 0
**Infections of the newborn (763-768)	127	2.1	W C	36 91	4 5	15 33	8 16	7 23	2 14	36 91	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0
Hemolytic disease of the newborn (770)	26	0.4	W C	16 10	8 4	7 5	1 0	0 0	0 0	16 9	0 1	0 0	0 0	0 0	0 0	0 0	0 0	0 0
All other defined dis- eases of early infancy (769,771,772)	70	1.2	W C	19 51	4 6	9 11	1 5	0 7	0 2	14 31	0 4	0 0	1 4	0 0	1 4	0 0	1 0	0 0
Violent & accidental deaths (800-999)	93	1.6	W C	23 70	0 0	2 4	0 1	1 4	0 3	3 12	1 11	2 8	1 6	2 4	2 8	2 2	1 5	0 5
Other specified causes	163	2.8	W C	53 110	1 3	6 6	5 6	2 2	1 6	15 23	5 18	3 10	8 9	5 7	3 10	2 8	0 5	3 2
III-defined (773-776, 780-793,795)	1,058	17.9	W C	235 823	131 319	73 191	9 33	3 24	1 24	217 590	3 80	5 48	1 30	4 25	2 17	1 7	1 10	0 5

* Rate per 1,000 live births.

** Pneumonia and diarrhea of the newborn (under 28 days) included in infections of the newborn (763-768).

TABLE 9
DEATHS FROM SELECTED CAUSES
MISSISSIPPI
1960

By Month of Occurrence on Basis of Place of Residence

Cause of Death and International List Number	Total	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
TOTAL ALL CAUSES	21,713	2,041	2,129	2,184	1,670	1,723	1,631	1,735	1,551	1,625	1,623	1,753	2,048
Dysentery (045-048)	27	2	6	1	0	1	2	3	4	3	2	1	2
Whooping cough (056)	7	1	0	0	2	0	0	1	1	1	0	0	0
Acute poliomyelitis (080)	4	0	0	0	0	0	1	2	1	0	0	0	0
Measles (085)	9	1	1	2	2	1	2	0	0	0	0	0	0
Influenza (480-483)	238	37	90	74	13	6	4	3	0	1	3	3	4
Lobar pneumonia (490)	158	20	33	12	13	8	9	9	9	1	15	9	20
Bronchopneumonia (491)	280	25	42	38	16	16	18	20	9	17	21	20	38
Other & unspecified pneumonia (492-493)	282	46	39	42	19	18	14	13	12	10	16	20	33
Diarrhea & enteritis (-2) (5710)	173	24	7	6	3	3	11	18	9	20	31	23	18
Diarrhea & enteritis (+2) (5711)	45	6	2	2	2	2	5	6	5	6	4	3	2
Accidents (800-962)	1,504	143	112	126	96	124	139	146	135	92	114	126	151
Motor vehicle accidents (810-835)	620	42	46	54	45	50	62	57	64	45	46	54	55

TABLE 10
MARRIAGES AND DIVORCES
MISSISSIPPI
1960

By County, by Race of Bride

County	Marriages						Divorces		
	Place of Licensure			Residence of Bride					
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
<u>TOTAL</u>	<u>21,233</u>	<u>14,029</u>	<u>7,204</u>	<u>17,198</u>	<u>10,765</u>	<u>6,433</u>	<u>5,260</u>	<u>3,553</u>	<u>1,707</u>
Adams	333	213	120	250	143	107	126	83	43
Alcorn	744	690	54	223	202	21	67	59	8
Amite	94	51	43	80	38	42	33	17	16
Attala	173	94	79	185	102	83	47	26	21
Benton	61	32	29	43	24	19	16	9	7
Bolivar	433	190	243	425	172	253	78	35	43
Calhoun	164	124	40	170	127	43	35	20	15
Carroll	69	38	31	74	34	40	19	5	14
Chickasaw	180	131	49	158	113	45	28	20	8
Choctaw	70	52	18	48	34	14	10	8	2
Claiborne	80	29	51	83	29	54	35	7	28
Clarke	144	89	55	133	78	55	29	17	12
Clay	144	54	60	113	54	59	34	18	16
Coahoma	355	141	214	299	115	184	84	39	45
Copiah	225	116	109	242	123	119	62	30	32
Covington	134	87	47	148	112	36	33	26	7
DeSoto	1,135	631	504	155	60	95	40	18	22
Forrest	517	395	122	508	384	124	202	148	54
Franklin	62	43	19	57	39	18	28	17	11
George	247	216	31	112	103	9	18	16	2
Greene	88	78	10	61	43	18	9	8	1
Grenada	155	86	69	161	87	74	49	30	19
Hancock	362	323	39	95	74	21	69	58	11
Harrison	1,330	1,168	162	1,026	866	160	557	477	80
Hinds	1,595	957	638	1,595	979	616	488	344	144
Holmes	207	63	144	235	69	166	37	15	22
Humphreys	147	60	87	145	59	86	42	18	24
Issaquena	15	5	10	17	2	15	4	1	3
Itawamba	148	142	6	93	86	7	32	29	3
Jackson	699	551	148	472	372	100	185	154	31
Jasper	102	62	40	139	78	61	18	12	6
Jefferson	80	27	53	71	15	56	25	4	21
Jeff Davis	101	48	53	82	41	41	20	11	9
Jones	574	452	122	513	409	104	167	137	30
Kosciusko	82	32	50	69	26	43	15	7	8
Lafayette	132	74	58	134	69	65	31	19	12

Table 10 - Page 2
 Marriages and Divorces - Mississippi - 1960

County	Marriages						Divorces		
	Place of Licensure			Residence of Bride					
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Lamar	117	108	9	110	104	6	25	21	4
Lauderdale	591	421	170	577	411	166	185	137	48
Lawrence	79	62	17	123	88	35	25	20	5
Leake	151	91	60	150	87	63	27	21	6
Lee	336	268	68	365	303	62	99	79	20
Leflore	374	157	217	362	161	201	98	49	49
Lincoln	210	150	60	175	133	42	68	49	19
Lowndes	323	223	100	295	192	103	83	62	21
Madison	280	109	171	233	88	145	96	35	61
Marion	258	181	77	213	138	75	63	43	20
Marshall	286	154	132	153	48	105	46	20	26
Monroe	250	193	57	238	177	61	68	53	15
Montgomery	132	85	47	124	81	43	33	18	15
Neshoba	199	141	58	185	133	52	52	34	18
Newton	140	96	44	149	103	46	31	21	10
Noxubee	94	31	63	97	36	61	17	3	14
Oktibbeha	169	91	78	148	78	70	24	13	11
Panola	225	110	115	224	119	105	58	35	23
Pearl River	312	260	52	204	168	36	87	74	13
Perry	65	52	13	66	54	12	17	16	1
Pike	262	169	93	242	151	91	103	68	35
Pontotoc	129	113	16	115	97	18	28	23	5
Prentiss	170	156	14	137	122	15	40	37	3
Quitman	193	83	110	189	74	115	34	20	14
Rankin	278	201	77	219	150	69	62	41	21
Scott	171	128	43	167	118	49	55	39	16
Sharkey	109	44	65	77	24	53	18	6	12
Simpson	138	104	34	146	102	44	39	26	13
Smith	119	92	27	131	89	42	36	34	2
Stone	98	84	14	72	64	8	18	15	3
Sunflower	329	108	221	331	107	224	77	40	37
Tallahatchie	233	99	134	202	78	124	46	23	23
Tate	170	98	72	127	61	66	15	10	5
Tippah	129	109	20	116	93	23	33	27	6
Tishomingo	167	158	9	131	124	7	29	25	4
Tunica	139	36	103	117	22	9	32	16	16
Union	184	151	33	160	129	31	35	25	10
Walthall	170	104	66	93	46	47	15	12	3
Warren	385	222	163	363	204	159	159	88	71
Washington	864	548	316	632	351	281	197	115	82
Wayne	139	94	45	105	73	32	44	29	15
Webster	82	74	8	96	87	9	13	12	1
Wilkinson	283	213	70	90	33	57	19	7	12
Winston	187	116	71	153	90	63	42	18	24
Yalobusha	105	70	35	122	81	41	19	13	6
Yazoo	258	128	130	260	132	128	78	39	39
Unstated				56	19	37			
<u>Out-of-state residents</u>				3,979	3,245	734			

TABLE 11
DIVORCES
MISSISSIPPI
1960

By Cause of Divorce

Cause	Total	White	Non-white
<u>TOTAL</u>	<u>5,260</u>	<u>3,553</u>	<u>1,707</u>
Cruel and inhuman treatment	2,631	2,195	436
Desertion	2,033	917	1,116
Adultery	249	150	99
Drunkenness	253	218	35
Penitentiary	34	27	7
Insanity	7	7	0
Annulment	12	10	2
Bigamy	11	5	6
Miscellaneous and other	16	11	5
<u>Not stated</u>	<u>14</u>	<u>13</u>	<u>1</u>

By Number of Minor Children

Minor Children	Total	White	Non-white
None	2,453	1,459	994
One	1,147	867	280
Two	780	627	153
Three or more	784	557	227
<u>Not stated</u>	<u>96</u>	<u>43</u>	<u>53</u>

By Number of Years Married

Number of Years Married	Total	White	Non-white
<u>TOTAL</u>	<u>5,260</u>	<u>3,553</u>	<u>1,707</u>
Under one year	272	247	25
One year	441	394	47
Two years	470	394	76
Three years	423	328	95
Four years	341	266	75
5-9 years	1,118	783	335
10-19 years	1,167	742	425
Twenty years and over	953	353	600
<u>Not stated</u>	<u>75</u>	<u>46</u>	<u>29</u>

Table 12--Live Births, by Year, by Race, on Basis of Place of Occurrence
with Rates: Mississippi, 1913-1960

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1912**	6,311					
1913	41,103	21,874	19,229	21.9	26.6	18.2
1914	40,623	21,360	19,263	21.4	25.7	18.0
1915	39,484	20,999	18,485	20.5	23.8	17.1
1916	39,033	20,130	18,903	20.0	23.6	17.2
1917	37,141	19,056	18,085	18.8	22.0	16.3
1918	39,391	20,005	19,386	19.7	22.8	17.2
1919	42,491	22,028	20,463	21.0	24.9	18.0
1920	43,743	22,438	21,305	24.4	26.3	22.7
1921	46,482	23,703	22,779	25.6	27.3	24.1
1922	43,723	21,286	22,437	23.4	24.1	23.6
1923	43,266	21,140	22,126	23.3	23.6	23.1
1924	43,960	22,180	21,780	23.4	24.3	22.5
1925	45,563	22,265	23,298	24.0	24.1	23.9
1926	50,449	24,600	25,849	26.2	26.2	26.3
1927	49,204	24,163	25,041	25.3	25.3	25.3
1928	48,126	23,642	24,484	24.5	24.4	24.5
1929	45,570	22,105	23,465	22.9	22.5	23.3
1930	47,982	23,192	24,790	23.8	23.2	24.0
1931	45,123	21,637	23,486	22.2	21.4	22.9
1932	46,481	21,883	24,598	22.6	21.4	23.9
1933	44,184	20,407	23,777	21.3	19.7	22.9
1934	47,905	22,342	25,563	23.0	21.4	24.6
1935	48,291	21,972	26,319	23.0	20.8	25.1
1936	49,261	22,194	27,067	23.3	20.8	25.7
1937	52,170	23,285	28,885	24.4	21.6	27.3
1938	53,729	24,046	29,683	25.0	22.1	27.8
1939	52,288	23,505	28,783	24.1	21.4	26.8
1940	52,526	23,084	29,442	24.0	20.8	27.3
1941	54,561	24,167	30,394	24.7	21.6	28.0
1942	56,521	25,306	31,215	25.4	22.4	28.6
1943	59,408	27,611	31,797	28.9	26.4	31.6
1944#	56,586	25,647	30,939	28.3	25.2	31.6
1945#	53,540	23,426	30,114	26.9	23.1	30.9
1946#	61,430	29,715	31,715	26.8	25.4	28.2
1947#	66,291	32,423	33,868	28.6	27.5	29.9
1948#	65,291	29,666	35,625	28.0	24.9	31.2
1949#	66,128	28,786	37,342	28.1	24.0	32.5
1950#	64,427	27,188	37,239	29.6	22.9	37.6
1951#	65,512	27,798	37,714	30.1	23.4	38.1
1952#	64,509	28,073	36,436	29.6	23.6	36.8
1953#	64,061	28,312	35,749	29.4	23.8	36.1
1954#	64,748	28,781	35,967	29.7	24.2	36.3
1955#	63,244	27,546	35,698	29.0	23.2	36.0

Table 12 (continued)

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1956#	62,178	28,958	34,900	29.5	23.8	36.2
1957#	60,712	26,804	33,908	28.8	23.4	35.2
1958#	59,189	26,812	32,377	27.1	20.9	35.9
1959#	60,562	27,979	32,583	27.2	20.9	36.6
1960#	59,210	27,198	32,012	27.2	21.6	34.8

* Rate per 1,000 population. 1913-1920 population estimates based on U. S. Censuses of 1910 & 1920; 1921-1930 population estimates based on U. S. Censuses of 1920 & 1930; 1931-1940 population estimates based on U. S. Censuses of 1930 & 1940; 1941 & 1942 projected estimates based on U. S. Censuses of 1930 & 1940; 1943 & 1944 estimates of civilian population obtained from Ration Book Registration; 1945 estimate of civilian population, July 1, 1945, obtained from Bureau of Census Bulletin Series P-46, No. 3, dated February 12, 1946; 1946-1949 projected estimates as of July 1, each year based on U. S. Censuses of 1930 & 1940; 1950-1955 U. S. Census population of April 1, 1950; estimate for 1956-1957 as of July 1, 1956, prepared by Dr. Harald A. Pedersen, Sociology Department, Mississippi State University, with adjustments by Statistical Service, Mississippi State Board of Health; 1958 estimate as of January 1, prepared by Dr. George L. Wilber and Mr. Carl T. Robbins, Sociology Department, Mississippi State University. 1959 estimate as of January 1, prepared by Dr. George L. Wilber and Ellen Bryant, Sociology Department, Mississippi State University; 1960 U. S. Census population of April 1, 1960.

** November-December, 1912.

On basis of place of residence.

3
3
3
3
6
9
2
2
5
6
1
8
1
3
0

Table 13--Deaths, by Year, by Race, on Basis of Place of Occurrence, with
Rates: Mississippi, 1913-1960

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1912**	4,341					
1913	21,305	7,643	13,662	11.4	9.3	13.0
1914	21,485	7,442	14,043	11.3	8.9	13.1
1915	24,001	8,088	15,913	12.5	9.6	14.7
1916	21,794	7,588	14,206	11.2	8.9	13.0
1917	23,579	8,048	15,531	11.9	9.3	14.0
1918	30,437	10,177	20,260	15.2	11.6	18.0
1919	22,667	8,142	14,525	12.7	9.5	15.5
1920	22,072	7,916	14,156	12.3	9.3	15.1
1921	19,991	7,308	12,683	11.0	8.4	13.4
1922	19,278	7,312	11,966	10.5	8.3	12.6
1923	20,363	7,688	12,675	11.0	8.6	13.2
1924	21,155	7,809	13,346	11.3	8.6	13.8
1925	22,229	8,143	14,086	11.7	8.8	14.4
1926	23,929	9,133	14,796	12.5	9.7	15.1
1927	23,257	8,765	14,492	12.0	9.2	14.6
1928	25,965	10,111	15,854	13.2	10.4	15.9
1929	25,868	10,048	15,820	13.0	10.2	15.7
1930	24,098	9,072	15,026	11.9	9.0	14.8
1931	22,007	8,641	13,366	10.8	8.5	13.0
1932	20,322	8,295	12,027	9.9	8.1	11.6
1933	21,617	8,898	12,719	10.4	8.6	12.2
1934	21,786	9,350	12,436	10.5	9.0	12.0
1935	21,285	9,186	12,099	10.1	8.7	11.6
1936	24,127	10,367	13,760	11.4	9.7	13.1
1937	24,005	10,055	13,950	11.2	9.3	13.2
1938	22,814	9,471	13,343	10.6	8.7	12.5
1939	22,701	9,464	13,237	10.5	8.6	12.3
1940	23,185	9,565	13,620	10.6	8.6	12.6
1941	22,573	9,442	13,131	10.2	8.4	12.1
1942	20,693	8,750	11,943	9.3	7.7	10.9
1943	20,850	9,139	11,711	10.2	8.7	11.6
1944#	20,199	8,830	11,369	10.1	8.7	11.6
1945#	19,024	8,452	10,572	9.6	8.3	10.8
1946#	19,321	8,995	10,326	8.4	7.7	9.2
1947#	20,007	9,238	10,769	8.6	7.8	9.5
1948#	20,183	9,216	10,967	8.6	7.7	9.6
1949#	20,489	9,233	11,256	8.7	7.7	9.8
1950#	20,567	9,466	11,101	9.4	8.0	11.2
1951#	21,007	9,593	11,414	9.6	8.1	11.5
1952#	20,657	9,387	11,270	9.5	7.9	11.4
1953#	20,492	9,495	10,997	9.4	8.0	11.1
1954#	19,353	9,453	9,900	8.9	8.0	10.0
1955#	19,403	9,501	9,902	8.9	8.0	10.0

Table 13 (continued)

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1956#	19,780	9,969	9,811	9.4	8.7	10.2
1957#	20,476	10,339	10,137	9.7	9.0	10.5
1958#	21,541	10,811	10,730	9.9	8.4	11.9
1959#	20,890	10,742	10,148	9.4	8.0	11.4
1960#	21,713	11,204	10,509	10.0	8.9	11.4

* Rate per 1,000 population. See footnote on Table 12 for source of population estimates.

** November-December, 1912.

On basis of place of residence.

Table 14--Maternal Deaths, by Year, by Race, on Basis of Place of Occurrence, with Rates: Mississippi, 1913-1960

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1913	447	162	285	108.8	74.1	148.2
1914	413	124	289	101.7	58.0	150.0
1915	289	88	201	73.2	41.9	108.7
1916	326	102	224	83.5	50.7	118.5
1917	297	105	192	80.0	55.1	106.2
1918	302	82	220	76.7	41.0	113.5
1919	407	173	234	95.8	78.5	114.4
1920	393	127	266	89.8	56.6	124.8
1921	391	145	246	84.1	61.2	108.0
1922	313	110	203	71.6	51.7	90.5
1923	321	108	213	74.2	51.1	96.3
1924	413	137	276	93.9	61.8	126.7
1925	435	143	292	95.5	64.2	125.3
1926	388	153	235	76.9	62.2	90.9
1927	409	134	275	83.1	55.4	109.8
1928	421	161	260	87.5	68.1	106.2
1929	410	152	258	90.0	68.8	110.0
1930	489	171	318	101.9	73.7	128.3
1931	352	131	221	78.0	60.5	94.1
1932	279	103	176	60.0	47.1	71.6
1933	315	116	199	71.3	56.8	83.7
1934	306	116	190	63.9	51.9	74.3
1935	334	137	197	69.2	62.4	74.8
1936	358	134	224	72.7	60.4	82.8
1937	370	150	220	70.9	64.4	76.2
1938	333	116	217	62.0	48.2	73.1
1939	311	97	214	59.5	41.3	74.3
1940	327	106	221	62.2	45.9	75.1
1941	293	93	200	53.7	38.5	65.8
1942	248	77	171	43.9	30.4	54.8
1943	233	66	167	39.2	23.9	52.5
1944**	225	72	153	39.8	28.1	49.4
1945**	216	64	152	40.3	27.3	50.5
1946**	199	67	132	32.4	22.5	41.6
1947**	175	50	125	26.4	15.4	36.9
1948**	177	38	139	27.1	12.8	39.0
1949**	147	32	115	22.2	11.1	30.8
1950**	165	30	135	25.6	11.0	36.2
1951**	143	28	115	21.8	10.1	30.5
1952**	119	15	104	18.4	5.3	28.5
1953**	119	24	95	18.6	8.5	26.6
1954**	104	11	93	16.1	3.8	25.8
1955**	76	12	64	12.0	4.4	17.9
1956**	74	12	62	11.9	4.4	17.8
1957**	75	14	61	12.4	5.2	18.0
1958**	53	10	43	9.0	3.7	13.3
1959**	55	9	46	9.1	3.2	14.1
1960**	59	8	51	10.0	2.9	15.9

* Rate per 10,000 live births.

** On basis of place of residence.

Table 15--Deaths Under 1 Year, by Year, by Race, on Basis of Place of Occurrence, with Rates: Mississippi, 1917-1960

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1917	3,689	1,296	2,393	99.3	68.0	132.3
1918						
1919	3,554			83.6		
1920	3,499	1,344	2,155	80.0	59.9	101.1
1921	3,157	1,243	1,914	67.9	52.4	84.0
1922	2,826	1,067	1,759	64.6	50.1	78.4
1923	2,908	1,104	1,804	67.2	52.2	81.5
1924	3,137	1,188	1,949	71.4	53.6	89.5
1925	3,176	1,179	1,997	69.7	53.0	85.7
1926	3,578	1,447	2,131	70.9	58.8	82.4
1927	3,291	1,303	1,988	66.9	54.0	79.4
1928	3,567	1,416	2,151	74.1	59.9	87.9
1929	3,275	1,268	2,007	71.9	57.4	85.5
1930	3,276	1,182	2,094	68.3	51.0	84.5
1931	2,547	961	1,586	56.4	44.4	67.5
1932	2,482	965	1,517	53.4	44.1	61.7
1933	2,800	1,035	1,765	63.4	50.7	74.2
1934	3,070	1,199	1,871	64.1	53.7	73.2
1935	2,587	1,044	1,543	53.6	47.5	58.6
1936	2,883	1,105	1,778	58.5	49.8	65.7
1937	3,098	1,174	1,924	59.4	50.4	66.6
1938	3,051	1,205	1,846	56.8	50.1	62.2
1939	2,908	1,135	1,773	55.6	48.3	61.6
1940	2,864	1,070	1,794	54.5	46.4	60.9
1941	2,969	1,113	1,856	54.4	46.1	61.1
1942	2,654	957	1,697	47.0	37.8	54.4
1943	2,774	1,137	1,637	46.7	41.2	51.5
1944**	2,489	974	1,515	44.0	38.0	49.0
1945**	2,192	846	1,346	40.9	36.1	44.7
1946**	2,298	1,005	1,293	37.4	33.8	40.8
1947**	2,428	1,009	1,419	36.6	31.1	41.9
1948**	2,470	988	1,482	37.8	33.3	41.6
1949**	2,612	900	1,712	39.5	31.3	45.8
1950**	2,348	763	1,585	36.4	28.1	42.6
1951**	2,589	814	1,775	39.5	29.3	47.1
1952**	2,592	760	1,832	40.2	27.1	50.3
1953**	2,534	807	1,727	39.6	28.5	48.3
1954**	2,360	744	1,616	36.4	25.8	44.9
1955**	2,338	680	1,658	37.0	24.7	46.4
1956**	2,353	734	1,619	37.8	26.9	46.4
1957**	2,257	694	1,563	37.2	25.9	46.1
1958**	2,424	706	1,718	41.0	26.3	53.1
1959**	2,355	677	1,678	38.9	24.2	51.5
1960**	2,457	714	1,743	41.5	26.2	54.4

* Rate per 1,000 live births.

** On basis of place of residence.

Table 16--Deaths Under 1 Month, by Year, by Race, on Basis of Place of Occurrence, with Rates: Mississippi, 1924-1960

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1924	1,598	730	868	36.4	32.9	39.8
1925	1,639	701	938	36.0	31.5	40.3
1926	1,731	823	908	34.3	33.4	35.1
1927	1,644	786	858	33.4	32.5	34.3
1928	1,788	804	984	37.2	34.0	40.2
1929	1,662	773	889	36.5	35.0	37.9
1930	1,692	715	977	35.3	30.8	39.4
1931	1,339	590	749	29.7	27.3	31.9
1932	1,361	572	789	29.3	26.1	32.1
1933	1,540	636	904	34.8	31.2	38.0
1934	1,505	644	861	31.4	28.8	33.7
1935	1,446	642	804	29.9	29.2	30.5
1936	1,494	676	818	30.3	30.4	30.2
1937	1,678	755	923	32.2	32.4	32.0
1938	1,688	747	941	31.4	31.1	31.7
1939	1,637	735	902	31.3	31.3	31.3
1940	1,613	680	933	30.7	29.4	31.7
1941	1,663	734	929	30.5	30.4	30.6
1942	1,544	638	906	27.3	25.2	29.0
1943	1,652	791	861	27.8	28.6	27.1
1944	1,436	661	775	25.2	25.4	25.1
1945**	1,351	624	727	25.2	26.6	24.1
1946**	1,561	794	767	25.4	26.7	24.2
1947**	1,595	750	845	24.1	23.1	24.9
1948**	1,674	773	901	25.6	26.0	25.3
1949**	1,594	661	933	24.1	23.0	25.0
1950**	1,446	560	886	22.4	20.6	23.8
1951**	1,679	626	1,053	25.6	22.5	27.9
1952**	1,658	581	1,077	25.7	20.7	29.6
1953**	1,630	635	995	25.4	22.4	27.8
1954**	1,521	601	920	23.5	20.9	25.6
1955**	1,440	527	913	22.8	19.1	25.6
1956**	1,451	581	870	23.3	21.3	24.9
1957**	1,425	540	885	23.5	20.1	26.1
1958**	1,475	540	935	24.9	20.1	28.9
1959**	1,450	515	935	23.9	18.4	28.7
1960**	1,497	550	947	25.3	20.2	29.6

* Rate per 1,000 live births.

** On basis of place of residence.

TABLE 17
DEATHS FROM HEART DISEASES, NEPHRITIS (ALL FORMS),
VASCULAR LESIONS, CANCER, AND TUBERCULOSIS (ALL FORMS)
MISSISSIPPI
1951-1960

Shown by Year and Race, on Basis of Place of Residence, with Rates*

YEAR	HEART DISEASES			NEPHRITIS (ALL FORMS)			VASCULAR LESIONS			CANCER			TUBERCULOSIS (ALL FORMS)			
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	
1951 No.	5,626	3,134	2,492	681	210	471	2,344	1,089	1,255	2,186	1,326	860	497	158	339	
Rate	258.2	263.7	251.6	31.2	17.7	47.6	107.6	91.6	126.7	100.3	111.6	86.8	22.8	13.3	34.2	
1952 No.	5,876	3,256	2,620	658	214	444	2,386	1,098	1,288	2,189	1,285	904	418	133	285	
Rate	269.7	273.9	264.6	30.2	18.0	44.8	109.5	92.4	130.1	100.5	108.1	91.3	19.2	11.2	28.8	
1953 No.	5,758	3,183	2,575	474	164	310	2,404	1,125	1,279	2,230	1,348	882	308	85	223	
Rate	264.3	267.8	260.0	21.8	13.8	31.3	110.3	94.6	129.2	102.3	113.4	89.1	14.1	7.2	22.5	
1954 No.	5,708	3,325	2,383	448	169	279	2,422	1,176	1,246	2,332	1,420	912	226	80	146	
Rate	262.0	279.7	240.6	20.6	14.2	28.2	111.2	98.9	125.8	107.0	119.5	92.1	10.4	6.7	14.7	
1955 No.	5,969	3,517	2,452	401	144	257	2,447	1,169	1,278	2,317	1,402	915	211	66	145	
Rate	273.9	295.9	247.6	18.4	12.1	26.0	112.3	98.3	129.0	106.3	118.0	92.4	9.7	5.6	14.6	
1956 No.	6,104	3,703	2,401	410	161	249	2,550	1,229	1,321	2,432	1,439	993	199	80	119	
Rate	289.3	322.8	249.4	19.4	14.0	25.9	120.9	107.1	137.2	115.3	125.4	103.2	9.4	7.0	12.4	
1957 No.	6,367	3,706	2,661	362	143	219	2,800	1,420	1,380	2,496	1,482	1,014	161	61	100	
Rate	301.8	323.0	276.4	17.1	12.5	22.7	132.7	123.8	143.9	118.3	129.2	105.3	..	7.6	5.3	10.4
1958 No.	6,515	3,924	2,591	322	100	222	2,867	1,426	1,441	2,525	1,524	1,001	172	82	90	
Rate	298.3	305.9	287.5	14.7	7.8	24.6	131.3	111.2	159.9	115.6	118.8	111.1	7.9	6.4	10.0	
1959 No.	6,219	3,772	2,447	290	101	189	2,683	1,386	1,297	2,600	1,580	1,020	153	62	91	
Rate	279.2	282.3	274.6	13.0	7.6	21.2	120.5	103.7	145.6	116.7	118.3	114.5	6.9	4.6	10.2	
1960 No.	6,312	3,879	2,433	282	94	188	2,960	1,513	1,447	2,635	1,644	991	147	57	90	
Rate	289.8	308.4	264.3	12.9	7.5	20.4	135.9	120.3	157.2	121.0	130.7	107.6	6.7	4.5	9.8	


* Rate per 100,000 population.

TABLE 18
BIRTHS, DEATHS, MARRIAGES
MISSISSIPPI
1960

By Month of Occurrence

Month	Births	Deaths	Marriages
<u>TOTAL</u>	<u>59,210</u>	<u>21,713</u>	<u>21,233</u>
January	5,064	2,041	1,573
February	4,677	2,129	1,387
March	5,031	2,184	1,364
April	4,489	1,670	1,591
May	4,160	1,723	1,703
June	4,529	1,631	2,341
July	5,216	1,735	2,116
August	5,510	1,551	1,908
September	5,600	1,625	1,697
October	5,014	1,623	1,652
November	4,842	1,753	1,579
<u>December</u>	<u>5,078</u>	<u>2,048</u>	<u>2,322</u>

COUNTY SUMMARIES



BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ADAMS COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,699	707	992	1,171	493	678	31.0	25.9	36.3
Fetal deaths	42	16	26	26	10	16	22.2	20.3	23.6
Deaths from all causes	458	174	284	395	148	247	10.5	7.8	13.2
Deaths under 1 year	56	17	39	37	13	24	31.6	26.4	35.4
Deaths under 1 month	32	9	23	26	9	17	22.2	18.2	25.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	4	1	3	10.6	5.2	16.0
Tbc., respiratory (001-008)	1	0	1	4	1	3	10.6	5.2	16.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	1	0	2.6	5.2	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	53	28	25	55	30	25	145.8	157.6	133.7
Diabetes (260)	16	5	11	9	2	7	23.8	10.5	37.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	61	27	34	48	19	29	127.2	99.8	155.1
Heart diseases (410-443)	142	53	89	128	45	83	339.2	236.4	444.0
Arteriosclerosis (450)	6	5	1	3	2	1	8.0	10.5	5.3
Influenza (480-483)	4	0	4	3	0	3	8.0	-	16.0
#Pneumonia (490-493)	11	2	9	7	1	6	18.6	5.2	32.1
#Diarrhea & enteritis (571)	5	0	5	1	0	1	2.6	-	5.3
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	5	0	5	5	0	5	13.2	-	26.7
Maternal deaths (640-689)	4	0	4	4	0	4	34.2	-	59.0
Cong. malformations (750-759)	6	4	2	5	4	1	13.2	21.0	5.3
Birth inj., atelect. (760-762)	7	5	2	7	5	2	6.0	10.1	2.9
#Other dis. early inf. (763-776)	25	6	19	16	3	13	13.7	6.1	19.2
Violent & acc. deaths (800-999)	48	18	30	49	18	31	129.9	94.6	165.8
Acc. deaths, total (800-962)	40	15	25	41	15	26	108.7	78.8	139.1
Motor vehicle acc. (810-835)	18	9	9	16	7	9	42.4	36.8	48.1
Ill-defined (780-795)	13	2	11	11	1	10	29.2	5.2	53.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ALCORN COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	509	426	83	481	399	82	19.0	18.2	24.6
Fetal deaths	5	3	2	6	4	2	12.5	10.0	24.4
Deaths from all causes	250	204	46	268	215	53	10.6	9.8	15.9
Deaths under 1 year	12	8	4	13	9	4	27.0	22.6	48.8
Deaths under 1 month	9	6	3	10	7	3	20.8	17.5	36.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	2	1	1	7.9	4.6	30.0
Tbc., respiratory (001-008)	1	1	0	2	1	1	7.9	4.6	30.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	1	0	1	1	0	4.0	4.6	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	1	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	31	27	4	38	33	5	150.3	150.3	150.0
Diabetes (260)	1	1	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	35	29	6	38	30	8	150.3	136.7	240.0
Heart diseases (410-443)	98	82	16	102	86	16	403.4	391.8	480.0
Arteriosclerosis (450)	5	5	0	8	8	0	31.6	36.4	-
Influenza (480-483)	2	0	2	2	0	2	7.9	-	60.0
#Pneumonia (490-493)	4	3	1	8	6	2	31.6	27.3	60.0
#Diarrhea & enteritis (571)	2	1	1	2	1	1	7.9	4.6	30.0
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	5	4	1	4	4	0	15.8	18.2	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	1	1	0	4.0	4.6	-
Birth inj., atelect. (760-762)	4	3	1	5	4	1	10.4	10.0	12.2
#Other dis. early inf. (763-776)	5	3	2	5	3	2	10.4	7.5	24.4
Violent & acc. deaths (800-999)	15	14	1	11	8	3	43.5	36.4	90.0
Acc. deaths, total (800-962)	14	13	1	11	8	3	43.5	36.4	90.0
Motor vehicle acc. (810-835)	8	8	0	1	1	0	4.0	4.6	-
Ill-defined (780-795)	11	8	3	10	7	3	40.0	31.9	90.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
AMITE COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

		Occurrence			Residence			Rate*		
		Total	White	Non-white	Number			Rate*		
					Total	White	Non-white	Total	White	Non-white
4.6	Live births	203	20	183	345	107	238	22.2	15.0	28.2
4.4	Fetal deaths	6	0	6	13	1	12	37.7	9.3	50.4
5.9	Deaths from all causes	92	34	58	161	81	80	10.3	11.4	9.5
8.8	Deaths under 1 year	10	0	10	19	1	18	55.1	9.3	75.6
6.6	Deaths under 1 month	5	0	5	12	1	11	34.8	9.3	46.2
	SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
30.0	Tbc., all forms (001-019)	1	0	1	1	0	1	6.4	-	11.8
30.0	Tbc., respiratory (001-008)	1	0	1	1	0	1	6.4	-	11.8
-	Syphilis (020-029)	0	0	0	0	0	0	-	-	-
-	Typhoid fever (040)	0	0	0	0	0	0	-	-	-
-	Dysentery (045-048)	0	0	0	0	0	0	-	-	-
-	Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
-	Diphtheria (055)	0	0	0	0	0	0	-	-	-
-	Whooping cough (056)	0	0	0	0	0	0	-	-	-
-	Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
-	Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
-	Smallpox (084)	0	0	0	0	0	0	-	-	-
-	Measles (085)	0	0	0	0	0	0	-	-	-
-	Malaria (110-117)	0	0	0	0	0	0	-	-	-
.50.0	Cancer, incl. leukemia (140-205)	6	4	2	15	10	5	96.3	140.2	59.2
-	Diabetes (260)	3	1	2	5	2	3	32.1	28.0	35.5
-	Pellagra (281)	0	0	0	0	0	0	-	-	-
240.0	Intracranial lesions (330-334)	12	4	8	23	13	10	147.7	182.3	118.4
480.0	Heart diseases (410-443)	30	15	15	51	33	18	327.5	462.8	213.2
-	Arteriosclerosis (450)	3	1	2	5	3	2	32.1	42.1	23.7
60.0	Influenza (480-483)	0	0	0	0	0	0	-	-	-
60.0	#Pneumonia (490-493)	3	2	1	6	3	3	38.5	42.1	35.5
30.0	#Diarrhea & enteritis (571)	1	0	1	1	0	1	6.4	-	11.8
-	Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
-	Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
-	Maternal deaths (640-689)	1	0	1	1	0	1	29.0	-	42.0
-	Cong. malformations (750-759)	1	1	0	2	2	0	12.8	28.0	-
12.2	Birth inj., atelect. (760-762)	0	0	0	1	0	1	2.9	-	4.2
24.4	#Other dis. early inf. (763-776)	1	0	1	4	1	3	11.6	9.3	12.6
90.0	Violent & acc. deaths (800-999)	5	1	4	9	3	6	57.8	42.1	71.1
90.0	Acc. deaths, total (800-962)	4	1	3	8	3	5	51.4	42.1	59.2
-	Motor vehicle acc. (810-835)	0	0	0	3	1	2	19.3	14.0	23.7
90.0	Ill-defined (780-795)	17	4	13	20	5	15	128.4	70.1	177.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (~28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ATTALA COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	481	203	278	472	195	277	22.1	16.5	29.0
Fetal deaths	17	4	13	17	4	13	36.0	20.5	46.9
Deaths from all causes	212	105	107	234	121	113	11.0	10.3	11.8
Deaths under 1 year	18	1	17	21	3	18	44.5	15.4	65.0
Deaths under 1 month	10	1	9	12	3	9	25.4	15.4	32.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	3	2	1	14.1	17.0	10.5
Tbc., respiratory (001-008)	0	0	0	2	1	1	9.4	8.5	10.5
Syphilis (020-029)	2	1	1	2	1	1	9.4	8.5	10.5
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcals infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	31	18	13	29	17	12	135.9	144.2	125.7
Diabetes (260)	6	3	3	5	3	2	23.4	25.4	21.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	35	18	17	29	17	12	135.9	144.2	125.7
Heart diseases (410-443)	65	31	34	78	41	37	365.6	347.8	387.6
Arteriosclerosis (450)	3	2	1	3	2	1	14.1	17.0	10.5
Influenza (480-483)	1	1	0	1	1	0	4.7	8.5	-
#Pneumonia (490-493)	8	2	6	8	2	6	37.5	17.0	62.8
#Diarrhea & enteritis (571)	1	0	1	1	0	1	4.7	-	10.5
Acute nephritis (590,591)	1	0	1	1	0	1	4.7	-	10.5
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	4.7	8.5	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	1	1	0	4.7	8.5	-
Birth inj., atelect. (760-762)	1	0	1	1	0	1	2.1	-	3.6
#Other dis. early inf. (763-776)	8	1	7	9	2	7	19.1	10.2	25.3
Violent & acc. deaths (800-999)	18	12	6	23	14	9	107.8	118.8	94.3
Acc. deaths, total (800-962)	16	12	4	20	14	6	93.7	118.8	62.8
Motor vehicle acc. (810-835)	12	10	2	12	11	1	56.2	93.3	10.5
Ill-defined (780-795)	14	8	6	16	8	8	75.0	67.8	83.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
BENTON COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	106	3	103	206	64	142	26.7	15.6	39.3
Fetal deaths	10	1	9	16	3	13	77.7	46.9	91.5
Deaths from all causes	48	29	19	67	41	26	8.7	10.0	7.2
Deaths under 1 year	1	0	1	4	0	4	19.4	-	28.2
Deaths under 1 month	0	0	0	2	0	2	9.7	-	14.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	7	5	2	10	6	4	129.5	145.8	110.8
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	17	11	6	23	17	6	297.8	413.2	166.2
Heart diseases (410-443)	8	6	2	12	9	3	155.4	218.8	83.1
Arteriosclerosis (450)	1	0	1	1	0	1	12.9	-	27.7
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	0	0	0	1	0	1	12.9	-	27.7
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	12.9	24.3	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	1	0	1	12.9	-	27.7
Birth inj., atelect. (760-762)	0	0	0	1	0	1	4.8	-	7.0
#Other dis. early inf. (763-776)	0	0	0	1	0	1	4.8	-	7.0
Violent & acc. deaths (800-999)	8	6	2	7	5	2	90.6	121.5	55.4
Acc. deaths, total (800-962)	8	6	2	7	5	2	90.6	121.5	55.4
Motor vehicle acc. (810-835)	5	5	0	5	5	0	64.7	121.5	-
Ill-defined (780-795)	5	0	5	5	0	5	64.7	-	138.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
BOLIVAR COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,861	303	1,558	1,803	344	1,459	33.1	19.6	39.5
Fetal deaths	84	13	71	89	19	70	49.4	55.2	48.0
Deaths from all causes	542	106	436	573	145	428	10.5	8.3	11.6
Deaths under 1 year	74	7	67	75	9	66	41.6	26.2	45.2
Deaths under 1 month	37	5	32	37	5	32	20.5	14.5	21.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	2	0	2	3.7	-	5.4
Tbc., respiratory (001-008)	2	0	2	2	0	2	3.7	-	5.4
Syphilis (020-029)	1	0	1	2	0	2	3.7	-	5.4
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	1.8	-	2.7
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	51	11	40	58	19	39	106.5	108.4	105.6
Diabetes (260)	11	2	9	9	2	7	16.5	11.4	18.9
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	96	11	85	87	14	73	159.7	79.9	197.6
Heart diseases (410-443)	164	51	113	180	65	115	330.5	371.0	311.3
Arteriosclerosis (450)	3	0	3	5	3	2	9.2	17.1	5.4
Influenza (480-483)	5	2	3	4	2	2	7.3	11.4	5.4
#Pneumonia (490-493)	8	0	8	13	4	9	23.9	22.8	24.4
#Diarrhea & enteritis (571)	14	0	14	14	0	14	25.7	-	37.9
Acute nephritis (590,591)	2	0	2	3	0	3	5.5	-	8.1
Chronic nephritis, etc. (592-594)	6	1	5	6	1	5	11.0	5.7	13.5
Maternal deaths (640-689)	1	0	1	3	0	3	16.6	-	20.6
Cong. malformations (750-759)	7	0	7	10	2	8	18.4	11.4	21.6
Birth inj., atelect. (760-762)	11	1	10	10	1	9	5.5	2.9	6.2
#Other dis. early inf. (763-776)	20	5	15	20	4	16	11.1	11.6	11.0
Violent & acc. deaths (800-999)	47	17	30	44	15	29	80.8	85.6	78.5
Acc. deaths, total (800-962)	40	16	24	38	14	24	69.8	79.9	65.0
Motor vehicle acc. (810-835)	12	8	4	7	4	3	12.8	22.8	8.1
Ill-defined (780-795)	60	2	58	59	3	56	108.3	17.1	151.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CALHOUN COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	341	165	176	329	178	151	20.6	15.4	34.7
Fetal deaths	12	3	9	9	4	5	27.4	22.5	33.1
Deaths from all causes	148	105	43	178	135	43	11.2	11.6	9.9
Deaths under 1 year	14	3	11	16	6	10	48.6	33.7	66.2
Deaths under 1 month	7	2	5	8	4	4	24.3	22.5	26.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	2	2	0	12.5	17.2	
Tbc., respiratory (001-008)	1	1	0	2	2	0	12.5	17.2	
Syphilis (020-029)	0	0	0	0	0	0	-	-	
Typhoid fever (040)	0	0	0	0	0	0	-	-	
Dysentery (045-048)	0	0	0	0	0	0	-	-	
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	
Diphtheria (055)	0	0	0	0	0	0	-	-	
Whooping cough (056)	0	0	0	0	0	0	-	-	
Meningococcal infections (057)	0	0	0	0	0	0	-	-	
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	
Smallpox (084)	0	0	0	0	0	0	-	-	
Measles (085)	0	0	0	0	0	0	-	-	
Malaria (110-117)	0	0	0	0	0	0	-	-	
Cancer, incl. leukemia (140-205)	17	11	6	25	18	7	156.8	155.2	161.1
Diabetes (260)	1	0	1	1	0	1	6.3	-	23.0
Pellagra (281)	0	0	0	0	0	0	-	-	
Intracranial lesions (330-334)	8	7	1	10	9	1	62.7	77.6	23.0
Heart diseases (410-443)	57	48	9	64	55	9	401.5	474.3	207.1
Arteriosclerosis (450)	7	4	3	8	5	3	50.2	43.1	69.0
Influenza (480-483)	2	1	1	3	2	1	18.8	17.2	23.0
#Pneumonia (490-493)	4	2	2	7	5	2	43.9	43.1	46.0
#Diarrhea & enteritis (571)	1	0	1	1	0	1	6.3	-	23.0
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	
Chronic nephritis, etc. (592-594)	1	1	0	3	2	1	18.8	17.2	23.0
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	
Birth inj., atelect. (760-762)	0	0	0	1	1	0	3.0	5.6	
#Other dis. early inf. (763-776)	6	1	5	6	2	4	18.2	11.2	26.5
Violent & acc. deaths (800-999)	15	12	3	15	13	2	94.1	112.1	46.0
Acc. deaths, total (800-962)	12	9	3	12	10	2	75.3	86.2	46.0
Motor vehicle acc. (810-835)	6	5	1	7	7	0	43.9	60.4	-
Ill-defined (780-795)	15	6	9	15	6	9	94.1	51.7	207.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CARROLL COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	193	17	176	233	45	188	20.8	9.6	28.9
Fetal deaths	8	1	7	14	3	11	60.1	66.7	58.5
Deaths from all causes	65	26	39	103	43	60	9.2	9.2	9.2
Deaths under 1 year	9	2	7	12	1	11	51.5	22.2	58.5
Deaths under 1 month	5	1	4	9	1	8	38.6	22.2	42.6
SELECTED CAUSES OF DEATH									
AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	8.9	21.4	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	8.9	21.4	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	5	3	2	6	3	3	53.7	64.1	46.2
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	4	0	4	10	5	5	89.5	106.9	76.9
Heart diseases (410-443)	13	8	5	28	18	10	250.5	384.9	153.8
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	1	0	1	1	0	1	8.9	-	15.4
#Pneumonia (490-493)	2	0	2	3	1	2	26.8	21.4	30.8
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	2	1	1	85.8	222.2	53.2
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	1	0	1	3	0	3	12.9	-	16.0
#Other dis. early inf. (763-776)	3	1	2	3	1	2	12.9	22.2	10.6
Violent & acc. deaths (800-999)	15	9	6	13	6	7	116.3	128.3	107.7
Acc. deaths, total (800-962)	15	9	6	12	6	6	107.4	128.3	92.3
Motor vehicle acc. (810-835)	11	7	4	8	4	4	71.6	85.5	61.5
Ill-defined (780-795)	14	2	12	18	3	15	161.0	64.1	230.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CHICKASAW COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	406	219	187	430	215	215	25.4	20.7	33.0
Fetal deaths	21	8	13	22	7	15	51.2	32.6	69.8
Deaths from all causes	184	122	62	191	123	68	11.3	11.8	10.4
Deaths under 1 year	14	9	5	12	6	6	27.9	27.9	27.9
Deaths under 1 month	9	5	4	8	3	5	18.6	14.0	23.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	1	0	1	1	0	1	5.9	-	15.4
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	1	0	1	1	0	5.9	9.6	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	18	15	3	20	14	6	118.4	134.9	92.2
Diabetes (260)	4	3	1	4	3	1	23.7	28.9	15.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	33	15	18	35	15	20	207.2	144.5	307.2
Heart diseases (410-443)	75	57	18	74	57	17	438.1	549.1	261.1
Arteriosclerosis (450)	3	3	0	2	2	0	11.8	19.3	-
Influenza (480-483)	2	1	1	1	1	0	5.9	9.6	-
#Pneumonia (490-493)	4	4	0	4	4	0	23.7	38.5	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	3	1	2	17.8	9.6	30.7
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	1	1	0	5.9	9.6	-
Birth inj., atelect. (760-762)	4	2	2	3	1	2	7.0	4.6	9.3
#Other dis. early inf. (763-776)	4	3	1	3	1	2	7.0	4.6	9.3
Violent & acc. deaths (800-999)	5	3	2	11	8	3	65.1	77.1	46.1
Acc. deaths, total (800-962)	3	2	1	9	7	2	53.3	67.4	30.7
Motor vehicle acc. (810-835)	1	1	0	4	4	0	23.7	38.5	-
Ill-defined (780-795)	7	4	3	7	4	3	41.4	38.5	46.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CHOCTAW COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	190	98	92	185	94	91	22.0	15.9	36.1
Fetal deaths	5	4	1	3	2	1	16.2	21.3	11.0
Deaths from all causes	95	64	31	115	78	37	13.6	13.2	14.7
Deaths under 1 year	5	1	4	6	2	4	32.4	21.3	44.0
Deaths under 1 month	2	1	1	2	1	1	10.8	10.6	11.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	11.9	-	39.7
Tbc., respiratory (001-008)	0	0	0	1	0	1	11.9	-	39.7
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever(040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	9	0	13	12	1	154.3	203.3	39.7
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	13	7	6	18	11	7	213.7	186.3	277.8
Heart diseases (410-443)	35	26	9	38	29	9	451.1	491.3	357.1
Arteriosclerosis (450)	2	1	1	2	1	1	23.7	16.9	39.7
Influenza (480-483)	2	2	0	2	2	0	23.7	33.9	-
#Pneumonia (490-493)	6	1	5	6	1	5	71.2	16.9	198.4
#Diarrhea & enteritis (571)	1	0	1	1	0	1	11.9	-	39.7
Acute nephritis (590,591)	0	0	0	1	1	0	11.9	16.9	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	11.9	16.9	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	2	1	1	2	1	1	10.8	10.6	11.0
Violent & acc. deaths (800-999)	6	5	1	9	6	3	106.8	101.6	119.0
Acc. deaths, total (800-962)	6	5	1	9	6	3	106.8	101.6	119.0
Motor vehicle acc. (810-835)	2	2	0	5	3	2	59.4	50.8	79.4
Ill-defined (780-795)	4	1	3	5	1	4	59.4	16.9	158.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CLAIBORNE COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	212	27	185	283	40	243	26.1	15.4	29.5
Fetal deaths	10	2	8	10	1	9	35.3	25.0	37.0
Deaths from all causes	114	22	92	153	31	122	14.1	11.9	14.8
Deaths under 1 year	17	0	17	19	0	19	67.1	-	78.2
Deaths under 1 month	6	0	6	8	0	8	28.3	-	32.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	3	6	15	5	10	138.3	192.3	121.3
Diabetes (260)	2	0	2	3	0	3	27.7	-	36.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	12	2	10	17	2	15	156.8	76.9	181.9
Heart diseases (410-443)	51	14	37	59	17	42	544.0	653.8	509.4
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	7	0	7	8	1	7	73.8	38.5	84.9
#Diarrhea & enteritis (571)	2	0	2	2	0	2	18.4	-	24.2
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	9.2	-	12.1
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	1	0	1	1	0	1	3.5	-	4.1
#Other dis. early inf. (763-776)	4	0	4	6	0	6	21.2	-	24.7
Violent & acc. deaths (800-999)	3	1	2	8	2	6	73.8	76.9	72.8
Acc. deaths, total (800-962)	2	0	2	7	1	6	64.5	38.5	72.8
Motor vehicle acc. (810-835)	0	0	0	1	0	1	9.2	-	12.1
Ill-defined (780-795)	12	0	12	16	1	15	147.5	38.5	181.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CLARKE COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	167	97	70	377	171	206	22.8	17.1	31.7
Fetal deaths	3	0	3	5	1	4	13.3	5.8	19.4
Deaths from all causes	155	90	65	198	116	82	12.0	11.6	12.6
Deaths under 1 year	12	3	9	21	5	16	55.7	29.2	77.7
Deaths under 1 month	10	3	7	17	5	12	45.1	29.2	58.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	3	1	2	18.2	10.0	30.8
Tbc., respiratory (001-008)	2	0	2	3	1	2	18.2	10.0	30.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	6.1	-	15.4
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	1	1	0	6.1	10.0	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	8	9	22	10	12	133.4	100.0	184.8
Diabetes (260)	4	2	2	5	3	2	30.3	30.0	30.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	23	13	10	32	18	14	194.0	180.0	215.6
Heart diseases (410-443)	58	42	16	64	49	15	388.0	490.0	231.0
Arteriosclerosis (450)	4	1	3	4	1	3	24.2	10.0	46.2
Influenza (480-483)	1	0	1	1	0	1	6.1	-	15.4
#Pneumonia (490-493)	2	0	2	5	1	4	30.3	10.0	61.6
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	6.1	10.0	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	1	1	0	6.1	10.0	-
Birth inj., atelect. (760-762)	3	1	2	4	2	2	10.6	11.7	9.7
#Other dis. early inf. (763-776)	5	2	3	10	3	7	26.5	17.5	34.0
Violent & acc. deaths (800-999)	9	4	5	12	6	6	72.8	60.0	92.4
Acc. deaths, total (800-962)	8	3	5	9	4	5	54.6	40.0	77.0
Motor vehicle acc. (810-835)	4	1	3	4	1	3	24.2	10.0	46.2
Ill-defined (780-795)	11	4	7	11	4	7	66.7	40.0	107.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
CLAY COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	457	195	262	478	214	264	25.2	23.2	27.2
Fetal deaths	11	4	7	11	4	7	23.0	18.7	26.5
Deaths from all causes	180	92	88	192	96	96	10.1	10.4	9.9
Deaths under 1 year	10	5	5	13	8	5	27.2	37.4	18.9
Deaths under 1 month	8	4	4	11	7	4	23.0	32.7	15.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	3	1	2	15.8	10.8	20.6
Tbc., respiratory (001-008)	1	0	1	3	1	2	15.8	10.8	20.6
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	1	0	1	5.3	-	10.3
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	8	9	19	11	8	100.4	119.4	82.3
Diabetes (260)	3	2	1	3	2	1	15.8	21.7	10.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	27	14	13	30	14	16	158.4	151.9	164.6
Heart diseases (410-443)	65	40	25	63	36	27	332.8	390.7	277.8
Arteriosclerosis (450)	1	1	0	1	1	0	5.3	10.8	-
Influenza (480-483)	2	1	1	3	1	2	15.8	10.8	20.6
#Pneumonia (490-493)	4	1	3	3	0	3	15.8	-	30.9
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	2	0	2	2	0	2	10.6	-	20.6
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	5.3	-	10.3
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	1	1	0	5.3	10.8	-
Birth inj., atelect. (760-762)	3	2	1	4	3	1	8.4	14.0	3.8
#Other dis. early inf. (763-776)	5	2	3	6	3	3	12.6	14.0	11.4
Violent & acc. deaths (800-999)	24	12	12	26	13	13	137.3	141.1	133.8
Acc. deaths, total (800-962)	19	11	8	20	12	8	105.6	130.2	82.3
Motor vehicle acc. (810-835)	8	7	1	8	7	1	42.2	76.0	10.3
Ill-defined (780-795)	11	1	10	12	1	11	63.4	10.8	113.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
COAHOMA COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,514	339	1,175	1,533	313	1,220	33.2	21.4	38.6
Fetal deaths	64	15	49	58	10	48	37.8	31.9	39.3
Deaths from all causes	511	126	385	531	118	413	11.5	8.1	13.1
Deaths under 1 year	82	5	77	81	5	76	52.8	16.0	62.3
Deaths under 1 month	53	3	50	52	3	49	33.9	9.6	40.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	0	3	4	0	4	8.6	-	12.7
Tbc., respiratory (001-008)	3	0	3	4	0	4	8.6	-	12.7
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	1	0	1	1	0	1	2.2	-	3.2
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	58	20	38	63	20	43	136.3	136.7	136.2
Diabetes (260)	1	1	0	3	1	2	6.5	6.8	6.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	63	20	43	64	17	47	138.5	116.2	148.8
Heart diseases (410-443)	149	47	102	158	43	115	341.9	293.9	364.1
Arteriosclerosis (450)	13	2	11	12	1	11	26.0	6.8	34.8
Influenza (480-483)	11	1	10	12	1	11	26.0	6.8	34.8
#Pneumonia (490-493)	22	9	13	19	6	13	41.1	41.0	41.2
#Diarrhea & enteritis (571)	8	1	7	9	1	8	19.5	6.8	25.3
Acute nephritis (590,591)	2	0	2	2	0	2	4.3	-	6.3
Chronic nephritis, etc. (592-594)	2	1	1	3	1	2	6.5	6.8	6.3
Maternal deaths (640-689)	2	0	2	2	0	2	13.0	-	16.4
Cong. malformations (750-759)	3	3	0	2	2	0	4.3	13.7	-
Birth inj., atelect. (760-762)	5	1	4	5	1	4	3.3	3.2	3.3
#Other dis. early inf. (763-776)	46	1	45	46	2	44	30.0	6.4	36.1
Violent & acc. deaths (800-999)	44	7	37	41	7	34	88.7	47.8	107.6
Acc. deaths, total (800-962)	29	5	24	29	6	23	62.8	41.0	72.8
Motor vehicle acc. (810-835)	4	2	2	6	3	3	13.0	20.5	9.5
Ill-defined (780-795)	33	1	32	36	3	33	77.9	20.5	104.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
COPIAH COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	868	302	566	695	252	443	25.7	19.4	31.5
Fetal deaths	13	2	11	14	2	12	20.1	7.9	27.1
Deaths from all causes	278	132	146	325	162	163	12.0	12.5	11.6
Deaths under 1 year	33	7	26	34	7	27	48.9	27.8	60.9
Deaths under 1 month	20	6	14	20	6	14	28.8	23.8	31.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	1	2	4	1	3	14.8	7.7	21.3
Tbc., respiratory (001-008)	2	1	1	3	1	2	11.1	7.7	14.2
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	34	22	12	43	28	15	159.0	215.5	106.7
Diabetes (260)	5	1	4	6	2	4	22.2	15.4	28.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	37	12	25	41	14	27	151.6	107.8	192.0
Heart diseases (410-443)	96	62	34	114	72	42	421.4	554.2	298.7
Arteriosclerosis (450)	2	0	2	4	2	2	14.8	15.4	14.2
Influenza (480-483)	6	0	6	7	1	6	25.9	7.7	42.7
#Pneumonia (490-493)	11	1	10	14	4	10	51.8	30.8	71.1
#Diarrhea & enteritis (571)	3	0	3	4	0	4	14.8	-	28.4
Acute nephritis (590,591)	3	2	1	2	1	1	7.4	7.7	7.1
Chronic nephritis, etc. (592-594)	5	2	3	6	2	4	22.2	15.4	28.4
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	4	2	2	5	3	2	18.5	23.1	14.2
Birth inj., atelect. (760-762)	3	1	2	2	0	2	2.9	-	4.5
#Other dis. early inf. (763-776)	13	4	9	12	4	8	17.3	15.9	18.0
Violent & acc. deaths (800-999)	14	5	9	15	6	9	55.4	46.2	64.0
Acc. deaths, total (800-962)	12	4	8	13	5	8	48.0	38.5	56.9
Motor vehicle acc. (810-835)	5	3	2	4	2	2	14.8	15.4	14.2
Ill-defined (780-795)	9	1	8	9	1	8	33.3	7.7	56.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
COVINGTON COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	194	130	64	329	153	176	24.1	17.2	37.1
Fetal deaths	5	4	1	10	6	4	30.4	39.2	22.7
Deaths from all causes	98	64	34	137	85	52	10.0	9.6	11.0
Deaths under 1 year	6	4	2	11	3	8	33.4	19.6	45.4
Deaths under 1 month	3	2	1	5	2	3	15.2	13.1	17.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	7.3	11.2	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	7.3	11.2	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	1	0	2	2	0	14.7	22.5	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	4	6	10	4	6	73.3	45.0	126.6
Diabetes (260)	2	2	0	1	1	0	7.3	11.2	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	14	8	6	17	11	6	124.7	123.6	126.6
Heart diseases (410-443)	30	20	10	40	27	13	293.3	303.5	274.2
Arteriosclerosis (450)	5	3	2	7	4	3	51.3	45.0	63.3
Influenza (480-483)	3	1	2	3	1	2	22.0	11.2	42.2
#Pneumonia (490-493)	5	5	0	10	6	4	73.3	67.4	84.4
#Diarrhea & enteritis (571)	2	1	1	3	1	2	22.0	11.2	42.2
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	7.3	-	21.1
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	4	3	1	4	2	2	12.2	13.1	11.4
#Other dis. early inf. (763-776)	0	0	0	1	0	1	3.0	-	5.7
Violent & acc. deaths (800-999)	9	6	3	11	7	4	80.7	78.7	84.4
Acc. deaths, total (800-962)	8	6	2	10	7	3	73.3	78.7	63.3
Motor vehicle acc. (810-835)	6	4	2	5	3	2	36.7	33.7	42.2
Ill-defined (780-795)	1	1	0	3	1	2	22.0	11.2	42.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
DESOTO COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	670	61	609	707	160	547	29.6	17.3	37.4
Fetal deaths	19	2	17	23	1	22	32.5	6.2	40.2
Deaths from all causes	172	52	120	215	81	134	9.0	8.8	9.2
Deaths under 1 year	24	1	23	32	7	25	45.3	43.8	45.7
Deaths under 1 month	16	0	16	21	4	17	29.7	25.0	31.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	3	0	3	12.6	-	20.5
Tbc., respiratory (001-008)	1	0	1	2	0	2	8.4	-	13.6
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	4.2	-	6.8
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	13	6	7	23	13	10	96.3	140.6	68.3
Diabetes (260)	0	0	0	1	1	0	4.2	10.8	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	20	7	13	24	9	15	100.4	97.3	102.4
Heart diseases (410-443)	32	15	17	46	25	21	192.5	270.3	143.4
Arteriosclerosis (450)	1	0	1	2	1	1	8.4	10.8	6.8
Influenza (480-483)	1	0	1	2	1	1	8.4	10.8	6.8
#Pneumonia (490-493)	7	1	6	9	3	6	37.7	32.4	41.0
#Diarrhea & enteritis (571)	2	0	2	2	0	2	8.4	-	13.6
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	1	2	5	2	3	20.9	21.6	20.5
Maternal deaths (640-689)	0	0	0	1	0	1	14.1	-	18.3
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	1	0	1	1	0	1	1.4	-	1.8
#Other dis. early inf. (763-776)	13	0	13	16	2	14	22.6	12.5	25.6
Violent & acc. deaths (800-999)	27	14	13	14	6	8	58.6	64.9	54.6
Acc. deaths, total (800-962)	22	12	10	9	4	5	37.7	43.2	34.1
Motor vehicle acc. (810-835)	11	5	6	2	1	1	8.4	10.8	6.8
Ill-defined (780-795)	35	5	30	40	6	34	167.4	64.9	232.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
FORREST COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,502	1,123	379	1,271	803	468	24.1	21.1	31.7
Fetal deaths	42	22	20	42	18	24	33.0	22.4	51.3
Deaths from all causes	538	368	170	525	312	213	10.0	8.2	14.4
Deaths under 1 year	48	27	21	52	20	32	40.9	24.9	68.4
Deaths under 1 month	29	19	10	30	14	16	23.6	17.4	34.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	3	1	2	5.7	2.6	13.6
Tbc., respiratory (001-008)	0	0	0	2	1	1	3.8	2.6	6.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	1	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	69	45	24	69	38	31	130.9	100.1	210.1
Diabetes (260)	14	6	8	14	6	8	26.6	15.8	54.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	68	45	23	63	34	29	119.5	89.5	196.6
Heart diseases (410-443)	175	135	40	161	117	44	305.4	308.1	298.3
Arteriosclerosis (450)	14	11	3	10	8	2	19.0	21.1	13.6
Influenza (480-483)	8	5	3	7	4	3	13.3	10.5	20.3
#Pneumonia (490-493)	17	12	5	23	14	9	43.6	36.9	61.0
#Diarrhea & enteritis (571)	7	3	4	8	2	6	15.2	5.3	40.7
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	5	5	0	4	4	0	7.6	10.5	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	11	7	4	10	6	4	19.0	15.8	27.1
Birth inj., atelect. (760-762)	10	5	5	9	4	5	7.1	5.0	10.7
#Other dis. early inf. (763-776)	12	8	4	16	5	11	12.6	6.2	23.5
Violent & acc. deaths (800-999)	51	43	8	49	35	14	92.9	92.2	94.9
Acc. deaths, total (800-962)	40	35	5	36	26	10	68.3	68.5	67.8
Motor vehicle acc. (810-835)	22	22	0	19	16	3	36.0	42.1	20.3
Ill-defined (780-795)	33	8	25	30	7	23	56.9	18.4	155.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
FRANKLIN COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	127	43	84	223	80	143	24.0	14.6	37.6
Fetal deaths	12	6	6	14	5	9	62.8	62.5	62.9
Deaths from all causes	90	47	43	102	57	45	11.0	10.4	11.8
Deaths under 1 year	9	2	7	10	2	8	44.8	25.0	55.9
Deaths under 1 month	7	2	5	7	2	5	31.4	25.0	35.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	13	9	4	13	7	6	140.0	127.6	157.9
Diabetes (260)	1	1	0	4	4	0	43.1	72.9	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	10	5	5	11	6	5	118.4	109.4	131.6
Heart diseases (410-443)	38	21	17	39	23	16	420.0	419.2	421.0
Arteriosclerosis (450)	1	1	0	2	2	0	21.5	36.4	-
Influenza (480-483)	1	0	1	1	0	1	10.8	-	26.3
#Pneumonia (490-493)	2	1	1	4	3	1	43.1	54.7	26.3
#Diarrhea & enteritis (571)	0	0	0	1	0	1	10.8	-	26.3
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	1	0	1	10.8	-	26.3
Birth inj., atelect. (760-762)	1	0	1	0	0	0	-	-	-
#Other dis. early inf. (763-776)	6	2	4	7	2	5	31.4	25.0	35.0
Violent & acc. deaths (800-999)	9	4	5	7	3	4	75.4	54.7	105.3
Acc. deaths, total (800-962)	8	3	5	6	2	4	64.6	36.4	105.3
Motor vehicle acc. (810-835)	5	1	4	4	1	3	43.1	18.2	78.9
Ill-defined (780-795)	3	1	2	3	1	2	32.3	18.2	52.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
GEORGE COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	293	248	45	300	252	48	27.0	25.7	37.3
Fetal deaths	5	5	0	8	8	0	26.7	31.7	-
Deaths from all causes	73	63	10	92	80	12	8.3	8.2	9.3
Deaths under 1 year	6	5	1	8	7	1	26.7	27.8	20.8
Deaths under 1 month	4	3	1	5	4	1	16.7	15.9	20.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	3	2	1	27.0	20.4	77.7
Tbc., respiratory (001-008)	1	1	0	3	2	1	27.0	20.4	77.7
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	16	14	2	19	17	2	171.2	173.3	155.4
Diabetes (260)	1	1	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	4	2	2	6	4	2	54.1	40.8	155.4
Heart diseases (410-443)	29	27	2	32	30	2	288.3	305.8	155.4
Arteriosclerosis (450)	2	2	0	2	2	0	18.0	20.4	-
Influenza (480-483)	2	2	0	2	2	0	18.0	20.4	-
#Pneumonia (490-493)	2	2	0	4	4	0	36.0	40.8	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	2	0	2	2	0	18.0	20.4	-
Birth inj., atelect. (760-762)	2	1	1	3	2	1	10.0	7.9	20.8
#Other dis. early inf. (763-776)	1	1	0	1	1	0	3.3	4.0	-
Violent & acc. deaths (800-999)	5	4	1	5	4	1	45.0	40.8	77.7
Acc. deaths, total (800-962)	5	4	1	5	4	1	45.0	40.8	77.7
Motor vehicle acc. (810-835)	2	2	0	2	2	0	18.0	20.4	-
Ill-defined (780-795)	0	0	0	0	0	0	-	-	-

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
GREENE COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	162	107	55	193	125	68	23.1	19.4	35.4
Fetal deaths	1	0	1	3	1	2	15.5	8.0	29.4
Deaths from all causes	70	49	21	82	61	21	9.8	9.5	10.9
Deaths under 1 year	11	8	3	14	10	4	72.5	80.0	58.8
Deaths under 1 month	4	3	1	8	7	1	41.4	56.0	14.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	1	1	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	1	0	12.0	15.5	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	7	2	12	9	3	143.4	139.7	156.0
Diabetes (260)	1	1	0	1	1	0	12.0	15.5	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	11	6	5	11	7	4	131.5	108.6	208.0
Heart diseases (410-443)	22	15	7	27	20	7	322.7	310.4	364.0
Arteriosclerosis (450)	2	2	0	2	2	0	23.9	31.0	-
Influenza (480-483)	1	1	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	1	0	2	2	0	23.9	31.0	-
#Diarrhea & enteritis (571)	2	1	1	2	1	1	23.9	15.5	52.0
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	3	3	0	35.8	46.6	-
Birth inj., atelect. (760-762)	4	3	1	4	3	1	20.7	24.0	14.7
#Other dis. early inf. (763-776)	2	2	0	2	2	0	10.4	16.0	-
Violent & acc. deaths (800-999)	6	3	3	7	4	3	83.7	62.1	156.0
Acc. deaths, total (800-962)	3	2	1	4	4	0	47.8	62.1	-
Motor vehicle acc. (810-835)	0	0	0	2	2	0	23.9	31.0	-
Ill-defined (780-795)	0	0	0	1	0	1	12.0	-	52.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
GRENADA COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	522	196	326	509	185	324	27.6	19.8	35.8
Fetal deaths	21	2	19	19	3	16	37.3	16.2	49.4
Deaths from all causes	217	110	107	214	97	117	11.6	10.4	12.9
Deaths under 1 year	24	5	19	25	5	20	49.1	27.0	61.7
Deaths under 1 month	16	3	13	17	4	13	33.4	21.6	40.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	3	1	2	16.3	10.7	22.1
Tbc., respiratory (001-008)	1	0	1	2	1	1	10.9	10.7	11.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	26	14	12	26	13	13	141.2	139.0	143.5
Diabetes (260)	2	1	1	2	2	0	10.9	21.4	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	52	20	32	48	15	33	260.7	160.4	364.4
Heart diseases (410-443)	54	36	18	46	25	21	249.9	267.3	231.9
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	4	1	3	4	1	3	21.7	10.7	33.1
#Pneumonia (490-493)	3	3	0	2	2	0	10.9	21.4	-
#Diarrhea & enteritis (571)	3	2	1	3	2	1	16.3	21.4	11.0
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	2	0	2	10.9	-	22.1
Maternal deaths (640-689)	1	0	1	1	0	1	19.6	-	30.9
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	1	1	0	1	1	0	2.0	5.4	-
#Other dis. early inf. (763-776)	12	2	10	14	3	11	27.5	16.2	34.0
Violent & acc. deaths (800-999)	17	9	8	19	9	10	103.2	96.2	110.4
Acc. deaths, total (800-962)	15	8	7	17	8	9	92.3	85.5	99.4
Motor vehicle acc. (810-835)	5	3	2	5	3	2	27.2	32.1	22.1
Ill-defined (780-795)	18	3	15	19	3	16	103.2	32.1	176.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HANCOCK COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	165	124	41	287	228	59	20.4	19.3	26.2
Fetal deaths	6	5	1	8	7	1	27.9	30.7	16.9
Deaths from all causes	107	92	15	119	101	18	8.5	8.6	8.0
Deaths under 1 year	3	3	0	3	3	0	10.4	13.2	-
Deaths under 1 month	3	3	0	3	3	0	10.4	13.2	-
SELECTED CAUSES OF DEATH									
AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	1	1	0	7.1	8.5	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	1	0	7.1	8.5	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	11	8	3	17	14	3	121.1	118.8	133.0
Diabetes (260)	0	0	0	2	1	1	14.2	8.5	44.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	18	17	1	19	16	3	135.3	135.8	133.0
Heart diseases (410-443)	34	31	3	38	35	3	270.7	297.0	133.0
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	1	1	0	1	1	0	7.1	8.5	-
#Pneumonia (490-493)	4	4	0	2	2	0	14.2	17.0	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	7.1	8.5	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	1	1	0	1	1	0	3.5	4.4	-
#Other dis. early inf. (763-776)	2	2	0	2	2	0	7.0	8.8	-
Violent & acc. deaths (800-999)	13	10	3	8	6	2	57.0	50.9	88.7
Acc. deaths, total (800-962)	10	9	1	5	5	0	35.6	42.4	-
Motor vehicle acc. (810-835)	8	8	0	4	4	0	28.5	33.9	-
Ill-defined (780-795)	15	12	3	13	10	3	92.6	84.9	133.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HARRISON COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	4,121	3,370	751	3,798	3,083	715	31.8	30.8	37.1
Fetal deaths	78	48	30	68	43	25	17.9	13.9	35.0
Deaths from all causes	1,025	775	250	902	667	235	7.5	6.6	12.2
Deaths under 1 year	110	76	34	109	76	33	28.7	24.6	46.2
Deaths under 1 month	89	69	20	87	67	20	22.9	21.7	28.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	2	1	7	4	3	5.8	4.0	15.6
Tbc., respiratory (001-008)	3	2	1	7	4	3	5.8	4.0	15.6
Syphilis (020-029)	3	1	2	3	2	1	2.5	2.0	5.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	1	0	1	1	0	0.8	1.0	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	127	93	34	117	88	29	97.9	87.8	150.6
Diabetes (260)	6	5	1	9	8	1	7.5	8.0	5.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	97	72	25	86	62	24	72.0	61.8	124.6
Heart diseases (410-443)	313	267	46	265	222	43	221.8	221.5	223.3
Arteriosclerosis (450)	12	12	0	9	9	0	7.5	9.0	-
Influenza (480-483)	5	5	0	4	4	0	3.3	4.0	-
#Pneumonia (490-493)	39	29	10	24	17	7	20.1	17.0	36.4
#Diarrhea & enteritis (571)	3	2	1	3	2	1	2.5	2.0	5.2
Acute nephritis (590,591)	1	1	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	11	7	4	9	6	3	7.5	6.0	15.6
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	11	9	2	12	10	2	10.0	10.0	10.4
Birth inj., atelect. (760-762)	45	34	11	44	33	11	11.6	10.7	15.4
#Other dis. early inf. (763-776)	34	27	7	33	26	7	8.7	8.4	9.8
Violent & acc. deaths (800-999)	99	70	29	90	59	31	75.3	58.9	161.0
Acc. deaths, total (800-962)	73	53	20	70	47	23	58.6	46.9	119.4
Motor vehicle acc. (810-835)	34	27	7	34	25	9	28.4	24.9	46.7
Ill-defined (780-795)	96	56	40	84	46	38	70.3	45.9	197.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HINDS COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	6,080	3,079	3,001	5,524	2,667	2,857	29.5	23.8	38.2
Fetal deaths	384	246	138	350	219	131	63.4	82.1	45.8
Deaths from all causes	2,164	1,166	998	1,462	765	697	7.8	6.8	9.3
Deaths under 1 year	262	93	169	191	65	126	34.6	24.4	44.1
Deaths under 1 month	173	75	98	134	53	81	24.2	19.9	28.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	18	3	15	13	2	11	7.0	1.8	14.7
Tbc., respiratory (001-008)	17	3	14	12	2	10	6.4	1.8	13.4
Syphilis (020-029)	7	1	6	4	1	3	2.1	0.9	4.0
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	1	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	1	0	1	1	0	1	0.5	-	1.3
Whooping cough (056)	1	0	1	1	0	1	0.5	-	1.3
Meningococcal infections (057)	3	2	1	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	333	210	123	194	126	68	103.7	112.3	90.9
Diabetes (260)	29	9	20	20	6	14	10.7	5.3	18.7
Pellagra (281)	3	0	3	3	0	3	1.6	-	4.0
Intracranial lesions (330-334)	226	112	114	158	71	87	84.5	63.3	116.2
Heart diseases (410-443)	502	317	185	379	238	141	202.6	212.1	188.4
Arteriosclerosis (450)	35	29	6	23	19	4	12.3	16.9	5.3
Influenza (480-483)	4	0	4	5	1	4	2.7	0.9	5.3
#Pneumonia (490-493)	62	33	29	39	19	20	20.8	16.9	26.7
#Diarrhea & enteritis (571)	7	2	5	6	0	6	3.2	-	8.0
Acute nephritis (590,591)	6	1	5	3	0	3	1.6	-	4.0
Chronic nephritis, etc. (592-594)	25	8	17	14	4	10	7.5	3.6	13.4
Maternal deaths (640-689)	4	1	3	2	0	2	3.6	-	7.0
Cong. malformations (750-759)	44	27	17	17	12	5	9.1	10.7	6.7
Birth inj., atelect. (760-762)	37	24	13	31	18	13	5.6	6.7	4.6
#Other dis. early inf. (763-776)	102	31	71	84	21	63	15.2	7.9	22.0
Violent & acc. deaths (800-999)	180	113	67	142	92	50	75.9	82.0	66.8
Acc. deaths, total (800-962)	147	96	51	113	77	36	60.4	68.6	48.1
Motor vehicle acc. (810-835)	72	49	23	55	42	13	29.4	37.4	17.4
Ill-defined (780-795)	180	39	141	153	30	123	81.8	26.7	164.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HOLMES COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	761	137	624	768	129	639	28.3	17.0	32.8
Fetal deaths	40	7	33	41	7	34	53.4	54.3	53.2
Deaths from all causes	299	83	216	327	93	234	12.1	12.2	12.0
Deaths under 1 year	33	4	29	35	3	32	45.6	23.2	50.1
Deaths under 1 month	19	1	18	20	1	19	26.0	7.8	29.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	2	0	2	7.4	-	10.2
Tbc., respiratory (001-008)	1	0	1	2	0	2	7.4	-	10.2
Syphilis (020-029)	1	0	1	1	0	1	3.7	-	5.1
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	3.7	-	5.1
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	2	2	0	1	1	0	3.7	13.2	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	36	15	21	44	20	24	162.4	263.3	123.1
Diabetes (260)	3	0	3	4	0	4	14.8	-	20.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	48	8	40	52	12	40	191.9	158.0	205.1
Heart diseases (410-443)	56	30	26	63	33	30	232.5	434.5	153.8
Arteriosclerosis (450)	10	1	9	10	1	9	36.9	13.2	46.2
Influenza (480-483)	2	1	1	2	1	1	7.4	13.2	5.1
#Pneumonia (490-493)	16	5	11	18	5	13	66.4	65.8	66.7
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	3.7	-	5.1
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	2	1	1	7.4	13.2	5.1
Birth inj., atelect. (760-762)	4	0	4	3	0	3	3.9	-	4.7
#Other dis. early inf. (763-776)	12	0	12	14	0	14	18.2	-	21.9
Violent & acc. deaths (800-999)	26	9	17	23	6	17	84.9	79.0	87.2
Acc. deaths, total (800-962)	19	7	12	15	3	12	55.4	39.5	61.5
Motor vehicle acc. (810-835)	7	4	3	4	2	2	14.8	26.3	10.2
Ill-defined (780-795)	57	5	52	56	6	50	206.7	79.0	256.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
HUMPHREYS COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	630	109	521	632	120	512	33.1	20.8	38.4
Fetal deaths	17	1	16	20	1	19	31.6	8.3	37.1
Deaths from all causes	175	53	122	202	60	142	10.6	10.4	10.6
Deaths under 1 year	35	7	28	38	7	31	60.1	58.3	60.5
Deaths under 1 month	17	5	12	18	5	13	28.5	41.7	25.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	5.2	-	7.5
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	11	4	7	16	6	10	83.8	104.2	75.0
Diabetes (260)	2	0	2	2	0	2	10.5	-	15.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	13	4	9	20	6	14	104.8	104.2	105.0
Heart diseases (410-443)	48	30	18	51	27	24	267.1	468.9	180.0
Arteriosclerosis (450)	1	0	1	1	0	1	5.2	-	7.5
Influenza (480-483)	0	0	0	1	0	1	5.2	-	7.5
#Pneumonia (490-493)	7	1	6	8	2	6	41.9	34.7	45.0
#Diarrhea & enteritis (571)	3	0	3	4	0	4	21.0	-	30.0
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	9	0	9	9	0	9	47.1	-	67.5
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	4	1	3	4	1	3	21.0	17.4	22.5
Birth inj., atelect. (760-762)	4	3	1	4	3	1	6.3	25.0	2.0
#Other dis. early inf. (763-776)	9	1	8	10	1	9	15.8	8.3	17.6
Violent & acc. deaths (800-999)	21	2	19	23	3	20	120.5	52.1	150.0
Acc. deaths, total (800-962)	18	1	17	20	2	18	104.8	34.7	135.0
Motor vehicle acc. (810-835)	3	1	2	3	2	1	15.7	34.7	7.5
Ill-defined (780-795)	28	1	27	29	2	27	151.9	34.7	202.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ISSAQUENA COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	57	0	57	114	14	100	31.9	11.9	41.7
Fetal deaths	3	0	3	8	2	6	70.2	142.8	60.0
Deaths from all causes	42	13	29	62	15	47	17.3	12.8	19.6
Deaths under 1 year	6	0	6	13	0	13	114.0	-	130.0
Deaths under 1 month	1	0	1	6	0	6	52.6	-	60.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	28.0	-	41.7
Tbc., respiratory (001-008)	1	0	1	1	0	1	28.0	-	41.7
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	3	1	2	4	1	3	111.8	85.0	125.0
Diabetes (260)	1	0	1	1	0	1	28.0	-	41.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	4	0	4	8	3	5	223.7	255.1	208.3
Heart diseases (410-443)	8	3	5	14	5	9	391.5	425.2	375.0
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	3	0	3	4	1	3	111.8	85.0	125.0
#Diarrhea & enteritis (571)	1	0	1	2	0	2	55.9	-	83.3
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	2	0	2	17.5	-	20.0
#Other dis. early inf. (763-776)	1	0	1	4	0	4	35.1	-	40.0
Violent & acc. deaths (800-999)	11	7	4	10	4	6	279.6	340.1	250.0
Acc. deaths, total (800-962)	9	5	4	9	3	6	251.7	255.1	250.0
Motor vehicle acc. (810-835)	3	2	1	5	2	3	139.8	170.1	125.0
Ill-defined (780-795)	7	2	5	7	1	6	195.7	85.0	250.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
ITAWAMBA COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	82	64	18	300	278	22	19.9	19.6	25.2
Fetal deaths	1	1	0	4	4	0	13.3	14.4	-
Deaths from all causes	88	82	6	110	101	9	7.3	7.1	10.3
Deaths under 1 year	4	2	2	7	4	3	23.3	14.4	136.4
Deaths under 1 month	2	2	0	5	4	1	16.7	14.4	45.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	2	0	2	2	0	13.3	14.1	-
Tbc., respiratory (001-008)	2	2	0	2	2	0	13.3	14.1	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	16	16	0	20	20	0	132.6	140.8	-
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	16	14	2	19	17	2	126.0	119.7	228.8
Heart diseases (410-443)	37	34	3	45	40	5	298.4	281.6	572.1
Arteriosclerosis (450)	1	1	0	1	1	0	6.6	7.0	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	2	0	3	3	0	19.9	21.1	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	1	1	0	6.6	7.0	-
Birth inj., atelect. (760-762)	1	1	0	2	2	0	6.7	7.2	-
#Other dis. early inf. (763-776)	0	0	0	2	1	1	6.7	3.6	45.4
Violent & acc. deaths (800-999)	5	5	0	4	4	0	26.5	28.2	-
Acc. deaths, total (800-962)	4	4	0	3	3	0	19.9	21.1	-
Motor vehicle acc. (810-835)	2	2	0	1	1	0	6.6	7.0	-
Ill-defined (780-795)	3	2	1	3	2	1	19.9	14.1	114.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

**BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JACKSON COUNTY, MISSISSIPPI, 1960**

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,267	859	408	1,608	1,171	437	29.0	26.2	40.2
Fetal deaths	44	28	16	54	33	21	33.6	28.2	48.0
Deaths from all causes	334	247	87	359	261	98	6.5	5.8	9.0
Deaths under 1 year	39	20	19	43	23	20	26.7	19.6	45.8
Deaths under 1 month	31	16	15	34	19	15	21.1	16.2	34.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	1.8	-	9.2
Tbc., respiratory (001-008)	0	0	0	1	0	1	1.8	-	9.2
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	0	1	1	0	1	1.8	-	9.2
Acute poliomyelitis (080)	0	0	0	1	1	0	1.8	2.2	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	1.8	-	9.2
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	36	26	10	45	32	13	81.0	71.6	119.7
Diabetes (260)	9	8	1	9	7	2	16.2	15.7	18.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	36	27	9	40	31	9	72.0	69.4	82.8
Heart diseases (410-443)	91	78	13	95	81	14	171.1	181.4	128.9
Arteriosclerosis (450)	4	4	0	4	4	0	7.2	9.0	-
Influenza (480-483)	1	1	0	1	1	0	1.8	2.2	-
#Pneumonia (490-493)	4	2	2	7	4	3	12.6	9.0	27.6
#Diarrhea & enteritis (571)	2	0	2	2	0	2	3.6	-	18.4
Acute nephritis (590,591)	1	1	0	1	1	0	1.8	2.2	-
Chronic nephritis, etc. (592-594)	3	2	1	3	3	0	5.4	6.7	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	4	2	2	4	2	2	7.2	4.5	18.4
Birth inj., atelect. (760-762)	5	2	3	5	2	3	3.1	1.7	6.9
#Other dis. early inf. (763-776)	24	13	11	25	14	11	15.5	12.0	25.2
Violent & acc. deaths (800-999)	53	39	14	48	35	13	86.4	78.4	119.7
Acc. deaths, total (800-962)	50	38	12	43	33	10	77.4	73.9	92.0
Motor vehicle acc. (810-835)	12	9	3	10	8	2	18.0	17.9	18.4
Ill-defined (780-795)	33	24	9	26	18	8	46.8	40.3	73.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JASPER COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	98	18	80	427	150	277	25.2	17.8	32.6
Fetal deaths	6	1	5	21	4	17	49.2	26.7	61.4
Deaths from all causes	92	40	52	161	78	83	9.5	9.3	9.8
Deaths under 1 year	7	1	6	14	3	11	32.8	20.0	39.7
Deaths under 1 month	1	0	1	6	1	5	14.0	6.7	18.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	7	3	19	14	5	112.4	166.6	58.8
Diabetes (260)	2	0	2	4	1	3	23.6	11.9	35.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	12	3	9	28	11	17	165.6	130.9	199.8
Heart diseases (410-443)	28	14	14	49	26	23	289.8	309.4	270.4
Arteriosclerosis (450)	0	0	0	1	1	0	5.9	11.9	-
Influenza (480-483)	2	1	1	2	1	1	11.8	11.9	11.8
#Pneumonia (490-493)	8	2	6	10	4	6	59.1	47.6	70.5
#Diarrhea & enteritis (571)	2	1	1	2	1	1	11.8	11.9	11.8
Acute nephritis (590,591)	1	0	1	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	5.9	-	11.8
Maternal deaths (640-689)	1	0	1	1	0	1	23.4	-	36.1
Cong. malformations (750-759)	1	1	0	3	1	2	17.7	11.9	23.5
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	1	0	1	4	1	3	9.4	6.7	10.8
Violent & acc. deaths (800-999)	7	5	2	10	6	4	59.1	71.4	47.0
Acc. deaths, total (800-962)	4	3	1	8	4	4	47.3	47.6	47.0
Motor vehicle acc. (810-835)	3	3	0	5	3	2	29.6	35.7	23.5
Ill-defined (780-795)	9	2	7	10	3	7	59.1	35.7	82.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JEFFERSON COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	96	0	96	274	38	236	27.0	15.3	30.8
Fetal deaths	7	0	7	11	1	10	40.1	26.3	42.4
Deaths from all causes	86	13	73	115	24	91	11.3	9.6	11.9
Deaths under 1 year	7	0	7	13	0	13	47.4	-	55.1
Deaths under 1 month	4	0	4	7	0	7	25.5	-	29.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	1	8	13	3	10	128.2	120.5	130.7
Diabetes (260)	1	0	1	1	0	1	9.8	-	13.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	20	3	17	22	5	17	216.9	200.9	222.1
Heart diseases (410-443)	29	8	21	38	12	26	374.7	482.1	339.7
Arteriosclerosis (450)	1	0	1	1	0	1	9.8	-	13.1
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	3	0	3	4	0	4	39.4	-	52.3
#Diarrhea & enteritis (571)	0	0	0	1	0	1	9.8	-	13.1
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	9.8	-	13.1
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	2	0	2	2	0	2	7.3	-	8.5
#Other dis. early inf. (763-776)	4	0	4	6	0	6	21.9	-	25.4
Violent & acc. deaths (800-999)	4	0	4	5	0	5	49.3	-	65.3
Acc. deaths, total (800-962)	3	0	3	4	0	4	39.4	-	52.3
Motor vehicle acc. (810-835)	1	0	1	1	0	1	9.8	-	13.1
Ill-defined (780-795)	8	1	7	10	2	8	98.6	80.4	104.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JEFFERSON DAVIS COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	295	145	150	322	110	212	23.8	18.0	28.6
Fetal deaths	9	2	7	10	2	8	31.0	18.2	37.7
Deaths from all causes	113	51	62	134	61	73	9.9	10.0	9.8
Deaths under 1 year	19	4	15	22	4	18	68.3	36.4	84.9
Deaths under 1 month	9	3	6	11	4	7	34.2	36.4	33.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	7.4	-	13.5
Tbc., respiratory (001-008)	0	0	0	1	0	1	7.4	-	13.5
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	1	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	5	2	3	8	4	4	59.1	65.3	54.0
Diabetes (260)	1	0	1	3	1	2	22.2	16.3	27.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	13	7	6	16	8	8	118.2	130.6	107.9
Heart diseases (410-443)	40	22	18	46	25	21	339.7	408.1	283.2
Arteriosclerosis (450)	2	1	1	3	2	1	22.2	32.6	13.5
Influenza (480-483)	1	0	1	2	1	1	14.8	16.3	13.5
#Pneumonia (490-493)	3	2	1	3	2	1	22.2	32.6	13.5
#Diarrhea & enteritis (571)	1	0	1	2	0	2	14.8	-	27.0
Acute nephritis (590,591)	1	1	0	1	1	0	7.4	16.3	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	7.4	16.3	-
Maternal deaths (640-689)	2	0	2	2	0	2	62.1	-	94.3
Cong. malformations (750-759)	2	1	1	3	2	1	22.2	32.6	13.5
Birth inj., atelect. (760-762)	1	1	0	2	2	0	6.2	18.2	-
#Other dis. early inf. (763-776)	6	2	4	5	2	3	15.5	18.2	14.2
Violent & acc. deaths (800-999)	15	7	8	9	2	7	66.5	32.6	94.4
Acc. deaths, total (800-962)	12	6	6	6	1	5	44.3	16.3	67.4
Motor vehicle acc. (810-835)	8	4	4	3	0	3	22.2	-	40.5
Ill-defined (780-795)	9	0	9	12	1	11	88.6	16.3	148.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
JONES COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	2,513	1,275	1,238	1,469	967	502	24.7	21.9	32.5
Fetal deaths	108	43	65	58	28	30	39.5	29.0	59.8
Deaths from all causes	717	447	270	557	391	166	9.4	8.9	10.7
Deaths under 1 year	116	39	77	64	33	31	43.6	34.1	61.8
Deaths under 1 month	73	32	41	42	27	15	28.6	27.9	29.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	3	0	3	5.0	-	19.4
Tbc., respiratory (001-008)	0	0	0	3	0	3	5.0	-	19.4
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	5	1	4	2	0	2	3.4	-	12.9
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	1	1	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	1.7	-	6.5
Meningococcal infections (057)	1	1	0	2	2	0	3.4	4.5	-
Acute poliomyelitis (080)	1	1	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	68	51	17	63	50	13	105.8	113.4	84.2
Diabetes (260)	11	7	4	5	4	1	8.4	9.1	6.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	99	66	33	79	56	23	132.7	127.0	148.9
Heart diseases (410-443)	184	133	51	154	114	40	258.6	258.5	258.9
Arteriosclerosis (450)	7	5	2	7	5	2	11.8	11.3	12.9
Influenza (480-483)	9	3	6	9	3	6	15.1	6.8	38.8
#Pneumonia (490-493)	41	26	15	31	24	7	52.1	54.4	45.3
#Diarrhea & enteritis (571)	14	3	11	11	4	7	18.5	9.1	45.3
Acute nephritis (590,591)	4	1	3	4	1	3	6.7	2.3	19.4
Chronic nephritis, etc. (592-594)	13	7	6	8	3	5	13.4	6.8	32.4
Maternal deaths (640-689)	2	1	1	2	1	1	13.6	10.3	19.9
Cong. malformations (750-759)	15	12	3	14	12	2	23.5	27.2	12.9
Birth inj., atelect. (760-762)	10	8	2	10	8	2	6.8	8.3	4.0
#Other dis. early inf. (763-776)	51	16	35	23	12	11	15.6	12.4	21.9
Violent & acc. deaths (800-999)	56	38	18	41	32	9	68.8	72.6	58.3
Acc. deaths, total (800-962)	49	34	15	36	29	7	60.5	65.8	45.3
Motor vehicle acc. (810-835)	26	20	6	16	15	1	26.9	34.0	6.5
Ill-defined (780-795)	27	15	12	27	15	12	45.3	34.0	77.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
KEMPER COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	207	42	165	296	67	229	24.1	13.9	30.7
Fetal deaths	10	1	9	15	1	14	50.7	14.9	61.1
Deaths from all causes	96	32	64	120	45	75	9.8	9.3	10.1
Deaths under 1 year	8	0	8	12	2	10	40.5	29.8	43.7
Deaths under 1 month	5	0	5	9	2	7	30.4	29.8	30.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	8.1	-	13.4
Tbc., respiratory (001-008)	1	0	1	1	0	1	8.1	-	13.4
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	4	5	9	5	4	73.3	103.6	53.7
Diabetes (260)	0	0	0	1	0	1	8.1	-	13.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	6	9	17	5	12	138.5	103.6	161.1
Heart diseases (410-443)	28	15	13	32	18	14	260.6	372.8	187.9
Arteriosclerosis (450)	1	0	1	1	0	1	8.1	-	13.4
Influenza (480-483)	3	0	3	3	0	3	24.4	-	40.3
#Pneumonia (490-493)	3	2	1	6	4	2	48.9	82.8	26.8
#Diarrhea & enteritis (571)	1	0	1	2	0	2	16.3	-	26.8
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	1	3	4	1	3	32.6	20.7	40.3
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	2	1	1	16.3	20.7	13.4
Birth inj., atelect. (760-762)	1	0	1	1	1	0	3.4	14.9	-
#Other dis. early inf. (763-776)	3	0	3	6	0	6	20.3	-	26.2
Violent & acc. deaths (800-999)	4	2	2	7	4	3	57.0	82.8	40.3
Acc. deaths, total (800-962)	2	0	2	5	2	3	40.7	41.4	40.3
Motor vehicle acc. (810-835)	2	0	2	3	1	2	24.4	20.7	26.8
III-defined (780-795)	19	1	18	21	2	19	171.0	41.4	255.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LAFAYETTE COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	484	248	236	497	250	247	23.3	17.7	34.1
Fetal deaths	20	2	18	20	2	18	40.2	8.0	72.9
Deaths from all causes	181	102	79	193	109	84	9.0	7.7	11.6
Deaths under 1 year	31	7	24	29	5	24	58.4	20.0	97.2
Deaths under 1 month	20	4	16	17	2	15	34.2	8.0	60.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	2	1	1	9.4	7.1	13.8
Tbc., respiratory (001-008)	0	0	0	1	1	0	4.7	7.1	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	13	11	2	18	14	4	84.3	99.2	55.2
Diabetes (260)	3	2	1	3	2	1	14.0	14.2	13.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	25	14	11	29	15	14	135.8	106.3	193.2
Heart diseases (410-443)	56	41	15	57	43	14	266.9	304.7	193.2
Arteriosclerosis (450)	4	4	0	4	4	0	18.7	28.3	-
Influenza (480-483)	2	0	2	2	0	2	9.4	-	27.6
#Pneumonia (490-493)	9	3	6	12	6	6	56.2	42.5	82.8
#Diarrhea & enteritis (571)	1	1	0	1	1	0	4.7	7.1	-
Acute nephritis (590,591)	1	1	0	1	1	0	4.7	7.1	-
Chronic nephritis, etc. (592-594)	4	2	2	4	2	2	18.7	14.2	27.6
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	1	1	0	4.7	7.1	-
Birth inj., atelect. (760-762)	6	2	4	5	1	4	10.1	4.0	16.2
#Other dis. early inf. (763-776)	14	1	13	13	0	13	26.2	-	52.6
Violent & acc. deaths (800-999)	17	12	5	14	9	5	65.6	63.8	69.0
Acc. deaths, total (800-962)	14	10	4	12	7	5	56.2	49.6	69.0
Motor vehicle acc. (810-835)	5	3	2	5	2	3	23.4	14.2	41.4
Ill-defined (780-795)	15	1	14	15	1	14	70.2	7.1	193.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LAMAR COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	86	56	30	307	243	64	22.4	21.2	28.7
Fetal deaths	3	3	0	7	7	0	22.8	28.8	-
Deaths from all causes	69	61	8	111	95	16	8.1	8.3	7.2
Deaths under 1 year	2	2	0	6	3	3	19.5	12.3	46.9
Deaths under 1 month	0	0	0	3	1	2	9.8	4.1	31.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	7.3	8.7	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	7.3	8.7	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	18	17	1	22	20	2	160.9	174.8	89.6
Diabetes (260)	3	3	0	4	4	0	29.2	35.0	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	7	6	1	15	14	1	109.7	122.3	44.8
Heart diseases (410-443)	19	16	3	32	27	5	234.0	236.0	224.0
Arteriosclerosis (450)	1	1	0	3	3	0	21.9	26.2	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	2	0	3	3	0	21.9	26.2	-
#Diarrhea & enteritis (571)	1	1	0	1	0	1	7.3	-	44.8
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	0	1	7.3	-	44.8
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	1	1	0	7.3	8.7	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	0	0	0	3	1	2	9.8	4.1	31.2
Violent & acc. deaths (800-999)	9	8	1	10	10	0	73.1	87.4	-
Acc. deaths, total (800-962)	9	8	1	10	10	0	73.1	87.4	-
Motor vehicle acc. (810-835)	6	5	1	4	4	0	29.2	35.0	-
Ill-defined (780-795)	2	0	2	2	0	2	14.6	-	89.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LAUDERDALE COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	2,154	1,063	1,091	1,603	842	761	23.9	19.3	32.4
Fetal deaths	71	14	57	46	11	35	28.7	13.1	46.0
Deaths from all causes	961	586	375	777	442	335	11.6	10.1	14.3
Deaths under 1 year	114	44	70	86	31	55	53.6	36.8	72.3
Deaths under 1 month	78	38	40	56	27	29	34.9	32.1	38.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	4	0	3	2	1	4.5	4.6	4.2
Tbc., respiratory (001-008)	4	4	0	3	2	1	4.5	4.6	4.2
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	1	1	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	1	0	1.5	2.3	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	100	78	22	85	66	19	126.6	151.2	80.9
Diabetes (260)	27	14	13	16	7	9	23.8	16.0	38.3
Pellagra (281)	1	1	0	1	1	0	1.5	2.3	-
Intracranial lesions (330-334)	143	101	42	101	62	39	150.5	142.1	166.1
Heart diseases (410-443)	252	178	74	227	154	73	338.2	352.9	310.8
Arteriosclerosis (450)	11	9	2	9	6	3	13.4	13.8	12.8
Influenza (480-483)	7	6	1	6	5	1	8.9	11.4	4.2
#Pneumonia (490-493)	35	15	20	29	9	20	43.2	20.6	85.2
#Diarrhea & enteritis (571)	10	1	9	5	0	5	7.4	-	21.3
Acute nephritis (590,591)	1	0	1	3	0	3	4.5	-	12.8
Chronic nephritis, etc. (592-594)	5	2	3	5	1	4	7.4	2.3	17.0
Maternal deaths (640-689)	1	1	0	0	0	0	-	-	-
Cong. malformations (750-759)	11	7	4	11	6	5	16.4	13.8	21.3
Birth inj., atelect. (760-762)	18	17	1	14	13	1	8.7	15.4	1.3
#Other dis. early inf. (763-776)	47	16	31	29	8	21	18.1	9.5	27.6
Violent & acc. deaths (800-999)	84	42	42	66	29	37	98.3	66.5	157.6
Acc. deaths, total (800-962)	68	32	36	51	20	31	76.0	45.8	132.0
Motor vehicle acc. (810-835)	30	17	13	18	8	10	26.8	18.3	42.6
Ill-defined (780-795)	106	33	73	94	28	66	140.0	64.2	281.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LAWRENCE COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	159	39	120	268	146	122	26.2	23.0	31.6
Fetal deaths	5	2	3	9	3	6	33.6	20.5	49.2
Deaths from all causes	102	52	50	135	79	56	13.2	12.4	14.5
Deaths under 1 year	17	4	13	18	6	12	67.2	41.1	98.4
Deaths under 1 month	10	3	7	11	4	7	41.0	27.4	57.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	9.8	-	25.9
Tbc., respiratory (001-008)	0	0	0	1	0	1	9.8	-	25.9
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	1	1	0	1	1	0	9.8	15.7	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	19	11	8	20	12	8	195.8	188.8	207.2
Diabetes (260)	2	1	1	2	1	1	19.6	15.7	25.9
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	10	1	9	16	8	8	156.6	125.9	207.2
Heart diseases (410-443)	32	20	12	41	29	12	401.4	456.4	310.8
Arteriosclerosis (450)	2	1	1	2	1	1	19.6	15.7	25.9
Influenza (480-483)	2	1	1	1	0	1	9.8	-	25.9
#Pneumonia (490-493)	6	3	3	4	2	2	39.2	31.5	51.8
#Diarrhea & enteritis (571)	2	1	1	2	1	1	19.6	15.7	25.9
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	0	2	4	1	3	39.2	15.7	77.7
Birth inj., atelect. (760-762)	1	1	0	1	1	0	3.7	6.8	-
#Other dis. early inf. (763-776)	8	2	6	9	3	6	33.6	20.5	49.2
Violent & acc. deaths (800-999)	6	4	2	12	8	4	117.5	125.9	103.6
Acc. deaths, total (800-962)	5	3	2	10	6	4	97.9	94.4	103.6
Motor vehicle acc. (810-835)	1	1	0	2	1	1	19.6	15.7	25.9
Ill-defined (780-795)	2	0	2	4	0	4	39.2	-	103.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LEAKE COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	329	149	180	423	192	231	22.7	18.2	28.5
Fetal deaths	6	0	6	12	3	9	28.4	15.6	39.0
Deaths from all causes	130	81	49	176	108	68	9.4	10.2	8.4
Deaths under 1 year	17	8	9	26	13	13	61.5	67.7	56.3
Deaths under 1 month	10	7	3	17	12	5	40.2	62.5	21.6
SELECTED CAUSES OF DEATH									
AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	5.4	9.5	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	5.4	9.5	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	15	10	5	25	18	7	134.0	170.5	86.4
Diabetes (260)	3	2	1	4	3	1	21.4	28.4	12.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	17	13	4	18	12	6	96.5	113.6	74.1
Heart diseases (410-443)	45	30	15	57	37	20	305.5	350.4	246.9
Arteriosclerosis (450)	4	4	0	5	5	0	26.8	47.4	-
Influenza (480-483)	1	1	0	3	2	1	16.1	18.9	12.3
#Pneumonia (490-493)	2	0	2	4	1	3	21.4	9.5	37.0
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	5.4	9.5	-
Maternal deaths (640-689)	1	0	1	1	0	1	23.6	-	43.3
Cong. malformations (750-759)	1	0	1	2	2	0	10.7	18.9	-
Birth inj., atelect. (760-762)	2	2	0	3	3	0	7.1	15.6	-
#Other dis. early inf. (763-776)	10	6	4	15	9	6	35.5	46.9	26.0
Violent & acc. deaths (800-999)	11	7	4	8	4	4	42.9	37.9	49.4
Acc. deaths, total (800-962)	9	6	3	6	3	3	32.2	28.4	37.0
Motor vehicle acc. (810-835)	6	6	0	2	2	0	10.7	18.9	-
Ill-defined (780-795)	8	0	8	8	2	6	42.9	18.9	74.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LEE COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,240	943	297	999	693	306	24.6	22.9	29.7
Fetal deaths	50	38	12	44	32	12	44.0	46.2	39.2
Deaths from all causes	419	301	118	404	288	116	10.0	9.5	11.3
Deaths under 1 year	43	25	18	34	17	17	34.0	24.5	55.6
Deaths under 1 month	30	17	13	23	11	12	23.0	15.9	39.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	2	0	3	3	0	7.4	9.9	-
Tbc., respiratory (001-008)	2	2	0	3	3	0	7.4	9.9	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	1	1	0	1	1	0	2.5	3.3	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	2	1	1	2	1	1	4.9	3.3	9.7
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	50	43	7	49	38	11	120.7	125.4	106.9
Diabetes (260)	5	1	4	5	1	4	12.3	3.3	38.9
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	61	41	20	61	42	19	150.3	138.6	184.7
Heart diseases (410-443)	139	103	36	140	103	37	344.9	339.9	359.6
Arteriosclerosis (450)	5	5	0	7	7	0	17.2	23.1	-
Influenza (480-483)	3	3	0	3	3	0	7.4	9.9	-
#Pneumonia (490-493)	16	10	6	16	10	6	39.4	33.0	58.3
#Diarrhea & enteritis (571)	2	1	1	2	1	1	4.9	3.3	9.7
Acute nephritis (590,591)	3	0	3	3	0	3	7.4	-	29.2
Chronic nephritis, etc. (592-594)	1	0	1	0	0	0	-	-	-
Maternal deaths (640-689)	1	0	1	1	0	1	10.0	-	32.7
Cong. malformations (750-759)	5	3	2	4	2	2	9.8	6.6	19.4
Birth inj., atelect. (760-762)	10	8	2	7	5	2	7.0	7.2	6.5
# Other dis. early inf. (763-776)	18	9	9	14	6	8	14.0	8.6	26.1
Violent & acc. deaths (800-999)	41	34	7	37	31	6	91.2	102.3	58.3
Acc. deaths, total (800-962)	36	31	5	32	28	4	78.8	92.4	38.9
Motor vehicle acc. (810-835)	18	17	1	18	17	1	44.3	56.1	9.7
Ill-defined (780-795)	8	4	4	9	5	4	22.2	16.5	38.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LEFLORE COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,534	396	1,138	1,501	351	1,150	31.8	21.0	37.8
Fetal deaths	90	32	58	80	25	55	53.3	71.2	47.8
Deaths from all causes	572	201	371	532	158	374	11.3	9.5	12.3
Deaths under 1 year	82	8	74	76	6	70	50.6	17.1	60.9
Deaths under 1 month	45	7	38	39	5	34	26.0	14.2	29.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	5	1	4	6	1	5	12.7	6.0	16.4
Tbc., respiratory (001-008)	4	1	3	5	1	4	10.6	6.0	13.1
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	2	0	2	4.2	-	6.6
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	0	1	2.1	-	3.3
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	63	29	34	62	27	35	131.5	161.7	115.0
Diabetes (260)	7	3	4	6	2	4	12.7	12.0	13.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	87	37	50	76	26	50	161.2	155.7	164.2
Heart diseases (410-443)	144	61	83	130	46	84	275.8	275.5	275.9
Arteriosclerosis (450)	5	0	5	6	1	5	12.7	6.0	16.4
Influenza (480-483)	2	0	2	3	0	3	6.4	-	9.8
#Pneumonia (490-493)	20	7	13	21	5	16	44.5	29.9	52.6
#Diarrhea & enteritis (571)	12	0	12	11	0	11	23.3	-	36.1
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	4.2	6.0	3.3
Maternal deaths (640-689)	4	1	3	3	0	3	20.0	-	26.1
Cong. malformations (750-759)	3	2	1	3	2	1	6.4	12.0	3.3
Birth inj., atelect. (760-762)	8	3	5	7	2	5	4.7	5.7	4.3
#Other dis. early inf. (763-776)	36	4	32	32	3	29	21.3	8.5	25.2
Violent & acc. deaths (800-999)	47	19	28	48	18	30	101.8	107.8	98.5
Acc. deaths, total (800-962)	39	15	24	40	14	26	84.8	83.8	85.4
Motor vehicle acc. (810-835)	18	9	9	20	10	10	42.4	59.9	32.8
Ill-defined (780-795)	78	13	65	69	7	62	146.4	41.9	203.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LINCOLN COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	668	415	253	661	382	279	24.7	20.8	33.4
Fetal deaths	18	9	9	22	9	13	33.3	23.6	46.6
Deaths from all causes	273	180	93	278	179	99	10.4	9.7	11.8
Deaths under 1 year	24	13	11	20	9	11	30.2	23.6	39.4
Deaths under 1 month	13	10	3	12	8	4	18.2	20.9	14.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	3.7	-	12.0
Tbc., respiratory (001-008)	1	0	1	1	0	1	3.7	-	12.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	23	11	12	31	17	14	115.8	92.4	167.6
Diabetes (260)	10	9	1	10	8	2	37.4	43.5	23.9
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	34	21	13	33	20	13	123.3	108.6	155.6
Heart diseases (410-443)	120	88	32	109	78	31	407.3	423.8	371.2
Arteriosclerosis (450)	8	7	1	8	7	1	29.9	38.0	12.0
Influenza (480-483)	2	1	1	1	0	1	3.7	-	12.0
#Pneumonia (490-493)	8	1	7	9	1	8	33.6	5.4	95.8
#Diarrhea & enteritis (571)	2	0	2	2	0	2	7.5	-	23.9
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	1	2	4	2	2	14.9	10.9	23.9
Maternal deaths (640-689)	1	1	0	1	1	0	15.1	26.2	-
Cong. malformations (750-759)	1	1	0	1	0	1	3.7	-	12.0
Birth inj., atelect. (760-762)	3	2	1	3	2	1	4.5	5.2	3.6
#Other dis. early inf. (763-776)	8	6	2	6	4	2	9.1	10.5	7.2
Violent & acc. deaths (800-999)	18	12	6	20	14	6	74.7	76.0	71.8
Acc. deaths, total (800-962)	15	10	5	17	12	5	63.5	65.2	59.9
Motor vehicle acc. (810-835)	9	8	1	9	7	2	33.6	38.0	23.9
Ill-defined (780-795)	13	4	9	13	4	9	48.6	21.7	107.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

**BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
LOWNDES COUNTY, MISSISSIPPI, 1960**

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence				Rate*		
	Total	White	Non-white	Number			Total	White	Non-white	
				Total	White	Non-white				
Live births	1,468	854	614	1,370	753	617	29.4	26.1	34.7	
Fetal deaths	62	32	30	54	25	29	39.4	33.2	47.0	
Deaths from all causes	422	206	216	418	198	220	9.0	6.8	12.4	
Deaths under 1 year	46	19	27	44	17	27	32.1	22.6	43.8	
Deaths under 1 month	27	15	12	25	13	12	18.2	17.3	19.4	
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.										
Tbc., all forms (001-019)	2	0	2	2	0	2	4.3	-	11.2	
Tbc., respiratory (001-008)	1	0	1	1	0	1	2.1	-	5.6	
Syphilis (020-029)	0	0	0	0	0	0	-	-	-	
Typhoid fever (040)	0	0	0	0	0	0	-	-	-	
Dysentery (045-048)	0	0	0	0	0	0	-	-	-	
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-	
Diphtheria (055)	0	0	0	0	0	0	-	-	-	
Whooping cough (056)	0	0	0	0	0	0	-	-	-	
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-	
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-	
Smallpox (084)	0	0	0	0	0	0	-	-	-	
Measles (085)	0	0	0	0	0	0	-	-	-	
Malaria (110-117)	0	0	0	0	0	0	-	-	-	
Cancer, incl. leukemia (140-205)	50	35	15	47	31	16	100.8	107.4	90.0	
Diabetes (260)	3	2	1	2	1	1	4.3	3.5	5.6	
Pellagra (281)	1	0	1	1	0	1	2.1	-	5.6	
Intracranial lesions (330-334)	50	24	26	54	25	29	115.8	86.6	163.2	
Heart diseases (410-443)	105	52	53	105	53	52	225.1	183.6	292.7	
Arteriosclerosis (450)	7	6	1	5	4	1	10.7	13.8	5.6	
Influenza (480-483)	17	1	16	15	1	14	32.2	3.5	78.8	
#Pneumonia (490-493)	13	3	10	15	4	11	32.2	13.8	61.9	
#Diarrhea & enteritis (571)	4	1	3	4	1	3	8.6	3.5	16.9	
Acute nephritis (590,591)	1	1	0	0	0	0	-	-	-	
Chronic nephritis, etc. (592-594)	5	3	2	7	3	4	15.0	10.4	22.5	
Maternal deaths (640-689)	4	2	2	3	1	2	21.9	13.3	32.4	
Cong. malformations (750-759)	6	4	2	6	4	2	12.9	13.8	11.2	
Birth inj., atelect. (760-762)	8	4	4	6	3	3	4.4	4.0	4.9	
#Other dis. early inf. (763-776)	14	8	6	13	7	6	9.5	9.3	9.7	
Violent & acc. deaths (800-999)	61	30	31	58	27	31	124.4	93.5	174.5	
Acc. deaths, total (800-962)	56	27	29	53	24	29	113.6	83.1	163.2	
Motor vehicle acc. (810-835)	23	15	8	18	11	7	38.6	38.1	39.4	
Ill-defined (780-795)	29	3	26	29	3	26	62.2	10.4	146.3	

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MADISON COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,041	165	876	1,048	160	888	31.8	17.3	37.6
Fetal deaths	39	5	34	40	6	34	38.2	37.5	38.3
Deaths from all causes	323	86	237	339	81	258	10.3	8.7	10.9
Deaths under 1 year	39	1	38	46	2	44	43.9	12.5	49.5
Deaths under 1 month	26	1	25	31	2	29	29.6	12.5	32.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	25	11	14	27	11	16	82.0	118.7	67.7
Diabetes (260)	4	0	4	5	0	5	15.2	-	21.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	44	15	29	40	10	30	121.6	107.9	126.9
Heart diseases (410-443)	87	32	55	82	28	54	249.2	302.1	228.4
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	2	1	1	2	1	1	6.1	10.8	4.2
#Pneumonia (490-493)	10	3	7	12	3	9	36.5	32.4	38.1
#Diarrhea & enteritis (571)	1	0	1	1	0	1	3.0	-	4.2
Acute nephritis (590,591)	1	0	1	1	0	1	3.0	-	4.2
Chronic nephritis, etc. (592-594)	5	2	3	7	2	5	21.3	21.6	21.2
Maternal deaths (640-689)	2	0	2	2	0	2	19.1	-	22.5
Cong. malformations (750-759)	2	0	2	5	1	4	15.2	10.8	16.9
Birth inj., atelect. (760-762)	2	0	2	3	0	3	2.9	-	3.4
#Other dis. early inf. (763-776)	20	1	19	22	1	21	21.0	6.2	23.6
Violent & acc. deaths (800-999)	26	9	17	25	8	17	76.0	86.3	71.9
Acc. deaths, total (800-962)	22	7	15	21	6	15	63.8	64.7	63.4
Motor vehicle acc. (810-835)	9	5	4	10	5	5	30.4	54.0	21.2
Ill-defined (780-795)	60	2	58	61	3	58	185.4	32.4	245.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MARION COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	552	334	218	623	368	255	26.7	23.9	32.3
Fetal deaths	22	13	9	25	14	11	40.1	38.0	43.1
Deaths from all causes	200	132	68	219	146	73	9.4	9.5	9.2
Deaths under 1 year	26	12	14	30	15	15	48.2	40.8	58.8
Deaths under 1 month	14	8	6	17	10	7	27.3	27.2	27.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	4.3	6.5	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	4.3	6.5	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	30	21	9	36	26	10	154.6	168.7	126.8
Diabetes (260)	2	1	1	2	1	1	8.6	6.5	12.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	43	31	12	44	32	12	188.9	207.7	152.2
Heart diseases (410-443)	52	34	18	53	33	20	227.5	214.2	253.6
Arteriosclerosis (450)	2	2	0	2	2	0	8.6	13.0	-
Influenza (480-483)	2	0	2	2	0	2	8.6	-	25.4
#Pneumonia (490-493)	10	3	7	10	4	6	42.9	26.0	76.1
#Diarrhea & enteritis (571)	3	1	2	2	0	2	8.6	-	25.4
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	2	2	0	8.6	13.0	-
Birth inj., atelect. (760-762)	5	2	3	6	3	3	9.6	8.2	11.8
#Other dis. early inf. (763-776)	6	4	2	7	4	3	11.2	10.9	11.8
Violent & acc. deaths (800-999)	12	8	4	13	9	4	55.8	58.4	50.7
Acc. deaths, total (800-962)	10	7	3	11	8	3	47.2	51.9	38.0
Motor vehicle acc. (810-835)	4	3	1	5	4	1	21.5	26.0	12.7
Ill-defined (780-795)	18	15	3	18	15	3	77.3	97.4	38.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MARSHALL COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	762	123	639	808	151	657	33.0	20.8	38.1
Fetal deaths	48	6	42	37	6	31	45.8	39.7	47.2
Deaths from all causes	271	96	175	260	83	177	10.6	11.4	10.3
Deaths under 1 year	39	5	34	31	6	25	38.4	39.7	38.0
Deaths under 1 month	28	5	23	25	5	20	30.9	33.1	30.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	2	0	2	8.2	-	11.6
Tbc., respiratory (001-008)	2	0	2	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	34	14	20	37	12	25	151.0	165.2	145.0
Diabetes (260)	8	5	3	6	3	3	24.5	41.3	17.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	43	13	30	39	13	26	159.2	179.0	150.8
Heart diseases (410-443)	85	42	43	78	33	45	318.3	454.3	261.0
Arteriosclerosis (450)	3	1	2	2	0	2	8.2	-	11.6
Influenza (480-483)	4	0	4	3	0	3	12.2	-	17.4
#Pneumonia (490-493)	5	0	5	4	0	4	16.3	-	23.2
#Diarrhea & enteritis (571)	3	0	3	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	1	2	3	1	2	12.2	13.8	11.6
Maternal deaths (640-689)	4	0	4	2	0	2	24.8	-	30.4
Cong. malformations (750-759)	2	0	2	3	1	2	12.2	13.8	11.6
Birth inj., atelect. (760-762)	5	2	3	4	2	2	5.0	13.2	3.0
#Other dis. early inf. (763-776)	23	3	20	17	3	14	21.0	19.9	21.3
Violent & acc. deaths (800-999)	15	8	7	17	5	12	69.4	68.8	69.6
Acc. deaths, total (800-962)	14	8	6	15	4	11	61.2	55.1	63.8
Motor vehicle acc. (810-835)	9	6	3	7	2	5	28.6	27.5	29.0
Ill-defined (780-795)	18	3	15	18	3	15	73.5	41.3	87.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MONROE COUNTY, MISSISSIPPI, 1960

Show by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	820	424	396	824	452	372	24.3	20.6	30.9
Fetal deaths	21	12	9	26	15	11	31.6	33.2	29.6
Deaths from all causes	336	200	136	359	214	145	10.6	9.8	12.1
Deaths under 1 year	23	8	15	24	9	15	29.1	19.9	40.3
Deaths under 1 month	13	6	7	14	7	7	17.0	15.5	18.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	2	1	1	5.9	4.6	8.3
Tbc., respiratory (001-008)	1	1	0	2	1	1	5.9	4.6	8.3
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	2.9	-	8.3
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	33	25	8	36	28	8	106.0	127.7	66.6
Diabetes (260)	4	0	4	4	0	4	11.8	-	33.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	67	40	27	68	41	27	200.3	186.9	224.6
Heart diseases (410-443)	100	73	27	109	79	30	321.0	360.2	249.6
Arteriosclerosis (450)	7	6	1	8	7	1	23.6	31.9	8.3
Influenza (480-483)	2	1	1	3	1	2	8.8	4.6	16.6
#Pneumonia (490-493)	.7	1	6	8	2	6	23.6	9.1	49.9
#Diarrhea & enteritis (571)	1	0	1	1	0	1	2.9	-	8.3
Acute nephritis (590,591)	2	2	0	2	2	0	5.9	9.1	-
Chronic nephritis, etc. (592-594)	13	2	11	13	2	11	38.3	9.1	91.5
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	3	2	1	3	2	1	8.8	9.1	8.3
Birth inj., atelect. (760-762)	1	0	1	1	0	1	1.2	-	2.7
#Other dis. early inf. (763-776)	11	5	6	12	6	6	14.6	13.3	16.1
Violent & acc. deaths (800-999)	30	19	11	29	17	12	85.4	77.5	99.8
Acc. deaths, total (800-962)	22	15	7	20	12	8	58.9	54.7	66.6
Motor vehicle acc. (810-835)	9	6	3	6	3	3	17.7	13.7	25.0
Ill-defined (780-795)	24	6	18	24	6	18	70.7	27.4	149.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
MONTGOMERY COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	342	167	175	326	150	176	24.5	20.4	29.5
Fetal deaths	12	2	10	11	3	8	33.7	20.0	45.4
Deaths from all causes	120	60	60	136	72	64	10.2	9.8	10.7
Deaths under 1 year	7	2	5	7	2	5	21.5	13.3	28.4
Deaths under 1 month	5	2	3	5	2	3	15.3	13.3	17.0
SELECTED CAUSES OF DEATH									
AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	1	0	1	1	0	1	7.5	-	16.7
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	13	8	5	15	10	5	112.6	136.1	83.7
Diabetes (260)	2	1	1	4	2	2	30.0	27.2	33.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	17	8	9	21	9	12	157.6	122.5	201.0
Heart diseases (410-443)	43	26	17	45	29	16	337.8	394.6	268.0
Arteriosclerosis (450)	2	0	2	2	0	2	15.0	-	33.5
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	4	2	2	4	2	2	30.0	27.2	33.5
#Diarrhea & enteritis (571)	2	0	2	1	0	1	7.5	-	16.7
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	7.5	13.6	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	2	1	1	15.0	13.6	16.7
Birth inj., atelect. (760-762)	2	1	1	1	1	0	3.1	6.7	-
#Other dis. early inf. (763-776)	2	0	2	2	0	2	6.1	-	11.4
Violent & acc. deaths (800-999)	8	3	5	10	4	6	75.1	54.4	100.5
Acc. deaths, total (800-962)	6	2	4	8	3	5	60.1	40.8	83.7
Motor vehicle acc. (810-835)	3	2	1	3	2	1	22.5	27.2	16.7
Ill-defined (780-795)	10	4	6	10	4	6	75.1	54.4	100.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR , AND
DEATHS FROM CERTAIN SELECTED CAUSES
NESHOBIA COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	466	249	217	453	243	210	21.6	16.2	35.6
Fetal deaths	14	4	10	15	5	10	33.1	20.6	47.6
Deaths from all causes	141	93	48	192	133	59	9.2	8.8	10.0
Deaths under 1 year	20	5	15	26	10	16	57.4	41.2	76.2
Deaths under 1 month	12	5	7	18	9	9	39.7	37.0	42.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	4.8	-	16.9
Tbc., respiratory (001-008)	0	0	0	1	0	1	4.8	-	16.9
Syphilis (020-029)	0	0	0	1	0	1	4.8	-	16.9
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	1	0	1	1	0	4.8	6.6	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	20	17	3	22	19	3	105.1	126.4	50.8
Diabetes (260)	2	1	1	2	1	1	9.6	6.6	16.9
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	23	17	6	27	21	6	129.0	139.8	101.7
Heart diseases (410-443)	43	31	12	62	46	16	296.3	306.1	271.1
Arteriosclerosis (450)	1	1	0	1	1	0	4.8	6.6	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	4	1	3	3	1	2	14.3	6.6	33.9
#Diarrhea & enteritis (571)	2	0	2	2	1	1	9.6	6.6	16.9
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	4.8	6.6	-
Maternal deaths (640-689)	0	0	0	1	1	0	22.1	41.2	-
Cong. malformations (750-759)	2	0	2	2	0	2	9.6	-	33.9
Birth inj., atelect. (760-762)	4	2	2	6	3	3	13.2	12.3	14.3
#Other dis. early inf. (763-776)	5	2	3	10	6	4	22.1	24.7	19.0
Violent & acc. deaths (800-999)	13	10	3	19	15	4	90.8	99.8	67.8
Acc. deaths, total (800-962)	11	8	3	16	13	3	76.4	86.5	50.8
Motor vehicle acc. (810-835)	8	7	1	11	10	1	52.6	66.6	16.9
Ill-defined (780-795)	7	1	6	8	1	7	38.2	6.6	118.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
NEWTON COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	349	230	119	416	194	222	21.3	15.0	33.8
Fetal deaths	23	11	12	26	9	17	62.5	46.4	76.6
Deaths from all causes	153	106	47	184	120	64	9.4	9.3	9.7
Deaths under 1 year	6	2	4	17	5	12	40.9	25.8	54.0
Deaths under 1 month	3	2	1	9	4	5	21.6	20.6	22.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	22	19	3	23	19	4	117.8	146.7	60.9
Diabetes (260)	3	2	1	3	3	0	15.4	23.2	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	33	24	9	42	29	13	215.2	223.9	198.0
Heart diseases (410-443)	47	35	12	51	39	12	261.3	301.2	182.7
Arteriosclerosis (450)	2	1	1	3	1	2	15.4	7.7	30.4
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	1	0	4	2	2	20.5	15.4	30.4
#Diarrhea & enteritis (571)	2	0	2	4	0	4	20.5	-	60.9
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	2	2	0	10.2	15.4	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	2	0	2	10.2	-	30.4
Birth inj., atelect. (760-762)	0	0	0	1	1	0	2.4	5.2	-
#Other dis. early inf. (763-776)	3	2	1	7	3	4	16.8	15.5	18.0
Violent & acc. deaths (800-999)	6	5	1	10	8	2	51.2	61.8	30.4
Acc. deaths, total (800-962)	5	4	1	9	7	2	46.1	54.0	30.4
Motor vehicle acc. (810-835)	0	0	0	3	3	0	15.4	23.2	-
Ill-defined (780-795)	14	5	9	17	6	11	87.1	46.3	167.5

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
NOXUBEE COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	485	85	400	482	83	399	28.6	17.6	33.0
Fetal deaths	22	2	20	23	3	20	47.7	36.1	50.1
Deaths from all causes	195	45	150	199	52	147	11.8	11.0	12.1
Deaths under 1 year	21	2	19	22	3	19	45.6	36.1	47.6
Deaths under 1 month	12	2	10	13	3	10	27.0	36.1	25.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	5.9	-	8.3
Tbc., respiratory (001-008)	1	0	1	1	0	1	5.9	-	8.3
Syphilis (020-029)	1	0	1	1	0	1	5.9	-	8.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	5.9	-	8.3
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	10	7	14	9	5	83.2	190.5	41.3
Diabetes (260)	2	0	2	2	0	2	11.9	-	16.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	31	6	25	31	7	24	184.2	148.2	198.3
Heart diseases (410-443)	48	13	35	48	15	33	285.3	317.5	272.7
Arteriosclerosis (450)	5	1	4	5	1	4	29.7	21.2	33.0
Influenza (480-483)	10	2	8	10	2	8	59.4	42.3	66.1
#Pneumonia (490-493)	5	0	5	6	1	5	35.6	21.2	41.3
#Diarrhea & enteritis (571)	2	1	1	2	1	1	11.9	21.2	8.3
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	0	3	4	0	4	23.8	-	33.0
Maternal deaths (640-689)	2	0	2	2	0	2	41.5	-	50.1
Cong. malformations (750-759)	2	1	1	2	1	1	11.9	21.2	8.3
Birth inj., atelect. (760-762)	1	0	1	1	0	1	2.1	-	2.5
#Other dis. early inf. (763-776)	9	1	8	10	2	8	20.7	24.1	20.0
Violent & acc. deaths (800-999)	13	4	9	12	3	9	71.3	63.5	74.4
Acc. deaths, total (800-962)	10	4	6	9	3	6	53.5	63.5	49.6
Motor vehicle acc. (810-835)	4	4	0	1	1	0	5.9	21.2	-
Ill-defined (780-795)	32	2	30	33	2	31	196.1	42.3	256.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
OKTIBBEHA COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	784	311	473	758	295	463	29.0	20.0	40.4
Fetal deaths	18	7	11	19	6	13	25.1	20.3	28.1
Deaths from all causes	189	93	96	202	93	109	7.7	6.3	9.5
Deaths under 1 year	24	7	17	25	6	19	33.0	20.3	41.0
Deaths under 1 month	17	7	10	15	5	10	19.8	16.9	21.6
SELECTED CAUSES OF DEATH									
AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	3.8	6.8	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	3.8	6.8	-
Syphilis (020-029)	0	0	0	1	0	1	3.8	-	8.7
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	3.8	-	8.7
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	29	17	12	30	18	12	114.6	122.2	104.8
Diabetes (260)	1	1	0	1	0	1	3.8	-	8.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	30	15	15	30	13	17	114.6	88.3	148.5
Heart diseases (410-443)	59	34	25	62	33	29	236.9	224.1	253.3
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	4	0	4	3	0	3	11.5	-	26.2
#Pneumonia (490-493)	6	1	5	6	1	5	22.9	6.8	43.7
#Diarrhea & enteritis (571)	3	0	3	4	0	4	15.3	-	34.9
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	3.8	-	8.7
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	0	2	3	1	2	11.5	6.8	17.5
Birth inj., atelect. (760-762)	2	1	1	2	1	1	2.6	3.4	2.2
#Other dis. early inf. (763-776)	11	6	5	8	4	4	10.6	13.6	8.6
Violent & acc. deaths (800-999)	13	8	5	16	10	6	61.1	67.9	52.4
Acc. deaths, total (800-962)	11	7	4	12	7	5	45.8	47.5	43.7
Motor vehicle acc. (810-835)	4	4	0	2	2	0	7.6	13.6	-
Ill-defined (780-795)	10	0	10	12	1	11	45.8	6.8	96.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PANOLA COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	811	219	592	840	235	605	29.2	18.7	37.3
Fetal deaths	27	2	25	33	3	30	39.3	12.8	49.6
Deaths from all causes	256	104	152	309	140	169	10.7	11.1	10.4
Deaths under 1 year	27	6	21	28	7	21	33.3	29.8	34.7
Deaths under 1 month	13	3	10	13	3	10	15.5	12.8	16.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	2	0	2	6.9	-	12.3
Tbc., respiratory (001-008)	2	0	2	2	0	2	6.9	-	12.3
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	2	0	2	6.9	-	12.3
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	3.5	-	6.2
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	31	15	16	40	22	18	138.9	175.1	110.9
Diabetes (260)	1	1	0	3	1	2	10.4	8.0	12.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	47	17	30	52	22	30	180.6	175.1	184.9
Heart diseases (410-443)	73	38	35	78	40	38	270.9	318.3	234.2
Arteriosclerosis (450)	1	1	0	3	3	0	10.4	23.9	-
Influenza (480-483)	5	0	5	5	1	4	17.4	8.0	24.6
#Pneumonia (490-493)	4	3	1	6	5	1	20.8	39.8	6.2
#Diarrhea & enteritis (571)	10	5	5	10	5	5	34.7	39.8	30.8
Acute nephritis (590,591)	1	0	1	1	0	1	3.5	-	6.2
Chronic nephritis, etc. (592-594)	4	2	2	4	2	2	13.9	15.9	12.3
Maternal deaths (640-689)	1	0	1	1	0	1	11.9	-	16.5
Cong. malformations (750-759)	1	0	1	4	3	1	13.9	23.9	6.2
Birth inj., atelect. (760-762)	3	2	1	3	2	1	3.6	8.5	1.6
#Other dis. early inf. (763-776)	9	1	8	9	1	8	10.7	4.2	13.2
Violent & acc. deaths (800-999)	22	8	14	29	12	17	100.7	95.5	104.8
Acc. deaths, total (800-962)	18	8	10	23	12	11	79.9	95.5	67.8
Motor vehicle acc. (810-835)	8	4	4	8	4	4	27.8	31.8	24.6
Ill-defined (780-795)	24	3	21	30	6	24	104.2	47.8	147.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PEARL RIVER COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		Non-white
				Total	White	Non-white	Total	White	
Live births	617	475	142	536	376	160	23.9	21.8	30.8
Fetal deaths	26	18	8	22	13	9	41.0	34.6	56.2
Deaths from all causes	202	142	60	236	169	67	10.5	9.8	12.9
Deaths under 1 year	24	12	12	31	17	14	57.8	45.2	87.5
Deaths under 1 month	16	10	6	21	13	8	39.2	34.6	50.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	2	1	1	8.9	5.8	19.3
Tbc., respiratory (001@008)	1	0	1	2	1	1	8.9	5.8	19.3
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	4.5	-	19.3
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	27	24	3	35	31	4	156.2	180.0	77.1
Diabetes (260)	4	3	1	3	2	1	13.4	11.6	19.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	22	17	5	22	18	4	98.2	104.5	77.1
Heart diseases (410-443)	64	47	17	68	50	18	303.4	290.3	346.8
Arteriosclerosis (450)	1	1	0	2	1	1	8.9	5.8	19.3
Influenza (480-483)	5	0	5	5	0	5	22.3	-	96.3
#Pneumonia (490-493)	4	2	2	7	5	2	31.2	29.0	38.5
#Diarrhea & enteritis (571)	0	0	0	1	1	0	4.5	5.8	-
Acute nephritis (590,591)	1	0	1	1	0	1	4.5	-	19.3
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	5	5	0	7	7	0	31.2	40.6	-
Birth inj., atelect. (760-762)	7	6	1	8	7	1	14.9	18.6	6.2
#Other dis. early inf. (763-776)	7	3	4	11	5	6	20.5	13.3	37.5
Violent & acc. deaths (800-999)	18	10	8	19	10	9	84.8	58.1	173.4
Acc. deaths, total (800-962)	12	8	4	14	9	5	62.5	52.3	96.3
Motor vehicle acc. (810-835)	6	6	0	7	7	0	31.2	40.6	-
Ill-defined (780-795)	9	6	3	10	7	3	44.6	40.6	57.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PERRY COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	61	43	18	230	168	62	26.3	26.5	25.7
Fetal deaths	2	0	2	8	2	6	34.8	11.9	96.8
Deaths from all causes	44	28	16	89	61	28	10.2	9.6	11.6
Deaths under 1 year	3	0	3	8	3	5	34.8	17.8	80.6
Deaths under 1 month	0	0	0	4	3	1	17.4	17.8	16.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	11.4	15.8	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	11.4	15.8	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	11.4	-	41.4
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	2	1	1	8	6	2	91.5	94.7	82.9
Diabetes (260)	1	1	0	3	2	1	34.3	31.6	41.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	9	7	2	19	14	5	217.3	221.1	207.3
Heart diseases (410-443)	13	9	4	25	18	7	285.9	284.2	290.2
Arteriosclerosis (450)	1	1	0	2	2	0	22.9	31.6	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	0	0	0	0	0	0	-	-	-
#Diarrhea & enteritis (571)	1	0	1	1	0	1	11.4	-	41.4
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	3	2	1	34.3	31.6	41.4
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	2	1	1	22.9	15.8	41.4
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	0	0	0	3	2	1	13.0	11.9	16.1
Violent & acc. deaths (800-999)	6	3	3	5	2	3	57.2	31.6	124.4
Acc. deaths, total (800-962)	4	2	2	5	2	3	57.2	31.6	124.4
Motor vehicle acc. (810-835)	1	1	0	1	1	0	11.4	15.8	-
Ill-defined (780-795)	2	1	1	2	1	1	22.9	15.8	41.4

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PIKE COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	997	505	492	863	388	475	24.6	19.7	30.8
Fetal deaths	22	8	14	18	7	11	20.8	18.0	23.2
Deaths from all causes	377	211	166	383	204	179	10.9	10.4	11.6
Deaths under 1 year	33	11	22	32	11	21	37.1	28.4	44.2
Deaths under 1 month	22	7	15	21	8	13	24.3	20.6	27.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	2.8	5.1	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	2.8	5.1	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	1	1	0	1	1	0	2.8	5.1	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	44	24	20	52	23	29	148.3	117.0	188.2
Diabetes (260)	1	1	0	1	1	0	2.8	5.1	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	57	32	25	64	40	24	182.5	203.5	155.8
Heart diseases (410-443)	133	85	48	122	72	50	347.9	366.3	324.5
Arteriosclerosis (450)	12	9	3	11	8	3	31.4	40.7	19.5
Influenza (480-483)	0	0	0	1	0	1	2.8	-	6.5
#Pneumonia (490-493)	10	3	7	10	3	7	28.5	15.3	45.4
#Diarrhea & enteritis (571)	1	0	1	1	0	1	2.8	-	6.5
Acute nephritis (590,591)	1	1	0	1	1	0	2.8	5.1	-
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	5.7	5.1	6.5
Maternal deaths (640-689)	1	0	1	1	0	1	11.6	-	21.0
Cong. malformations (750-759)	3	2	1	2	1	1	5.7	5.1	6.5
Birth inj., atelect. (760-762)	6	2	4	6	2	4	7.0	5.2	8.4
#Other dis. early inf. (763-776)	12	5	7	11	5	6	12.7	12.9	12.6
Violent & acc. deaths (800-999)	18	10	8	15	7	8	42.8	35.6	51.9
Acc. deaths, total (800-962)	13	8	5	11	6	5	31.4	30.5	32.4
Motor vehicle acc. (810-835)	9	4	5	8	3	5	22.8	15.3	32.4
Ill-defined (780-795)	41	12	29	37	9	28	105.5	45.8	181.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PONTOTOC COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	331	249	82	344	254	90	20.0	18.2	27.4
Fetal deaths	6	6	0	5	5	0	14.5	19.7	-
Deaths from all causes	179	148	31	195	162	33	11.3	11.6	10.0
Deaths under 1 year	9	5	4	11	7	4	32.0	27.6	44.4
Deaths under 1 month	6	4	2	8	6	2	23.2	23.6	22.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	5.8	7.2	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	5.8	7.2	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	16	1	21	20	1	121.9	143.4	30.4
Diabetes (260)	2	1	1	2	1	1	11.6	7.2	30.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	20	17	3	24	19	5	139.3	136.2	152.2
Heart diseases (410-443)	84	71	13	81	68	13	470.0	487.6	395.6
Arteriosclerosis (450)	5	5	0	4	4	0	23.2	28.7	-
Influenza (480-483)	4	4	0	4	4	0	23.2	28.7	-
#Pneumonia (490-493)	8	6	2	8	6	2	46.4	43.0	60.9
#Diarrhea & enteritis (571)	2	0	2	2	0	2	11.6	-	60.9
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	11.6	7.2	30.4
Maternal deaths (640-689)	2	1	1	2	1	1	58.1	39.4	111.1
Cong. malformations (750-759)	1	1	0	1	1	0	5.8	7.2	-
Birth inj., atelect. (760-762)	3	3	0	3	3	0	8.7	11.8	-
#Other dis. early inf. (763-776)	2	1	1	4	3	1	11.6	11.8	11.1
Violent & acc. deaths (800-999)	10	9	1	16	15	1	92.8	107.6	30.4
Acc. deaths, total (800-962)	3	2	1	8	7	1	46.4	50.2	30.4
Motor vehicle acc. (810-835)	1	0	1	3	2	1	17.4	14.3	30.4
Ill-defined (780-795)	2	0	2	2	0	2	11.6	-	60.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
PRENTISS COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	427	361	66	372	310	62	20.7	19.7	28.4
Fetal deaths	15	14	1	12	11	1	32.2	35.5	16.1
Deaths from all causes	200	169	31	186	158	28	10.4	10.0	12.8
Deaths under 1 year	7	3	4	6	2	4	16.1	6.4	64.5
Deaths under 1 month	3	3	0	2	2	0	5.4	6.4	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	4	0	3	3	0	16.7	19.0	-
Tbc., respiratory (001-008)	4	4	0	3	3	0	16.7	19.0	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	25	23	2	26	23	3	144.8	145.9	137.2
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	46	35	11	37	30	7	206.1	190.3	320.2
Heart diseases (410-443)	64	54	10	58	48	10	323.1	304.5	457.4
Arteriosclerosis (450)	2	1	1	2	1	1	11.1	6.3	45.7
Influenza (480-483)	1	1	0	2	2	0	11.1	12.7	-
#Pneumonia (490-493)	9	7	2	7	5	2	39.0	31.7	91.5
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	2	0	1	1	0	5.6	6.3	-
Birth inj., atelect. (760-762)	2	1	1	2	1	1	5.4	3.2	16.1
#Other dis. early inf. (763-776)	1	1	0	1	1	0	2.7	3.2	-
Violent & acc. deaths (800-999)	9	9	0	15	15	0	83.6	95.2	-
Acc. deaths, total (800-962)	6	6	0	12	12	0	66.8	76.1	-
Motor vehicle acc. (810-835)	1	1	0	5	5	0	27.8	31.7	-
Ill-defined (780-795)	13	10	3	11	8	3	61.3	50.8	137.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
QUITMAN COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	746	188	558	665	153	512	31.6	19.8	38.5
Fetal deaths	22	4	18	22	4	18	33.1	26.1	35.2
Deaths from all causes	205	52	153	239	68	171	11.4	8.8	12.8
Deaths under 1 year	42	4	38	45	6	39	67.7	39.2	76.2
Deaths under 1 month	21	1	20	23	1	22	34.6	6.5	43.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	1	3	5	1	4	23.8	13.0	30.1
Tbc., respiratory (001-008)	4	1	3	5	1	4	23.8	13.0	30.1
Syphilis (020-029)	1	0	1	1	0	1	4.8	-	7.5
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	0	2	1	0	1	4.8	-	7.5
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	4.8	-	7.5
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	15	7	8	23	10	13	109.4	129.6	97.7
Diabetes (260)	5	0	5	5	0	5	23.8	-	37.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	17	7	10	20	9	11	95.2	116.6	82.7
Heart diseases (410-443)	50	18	32	64	27	37	304.5	350.0	278.1
Arteriosclerosis (450)	1	0	1	0	0	0	-	-	-
Influenza (480-483)	11	0	11	10	0	10	47.6	-	75.2
#Pneumonia (490-493)	5	0	5	6	1	5	28.5	13.0	37.6
#Diarrhea & enteritis (571)	5	0	5	6	0	.6	28.5	-	45.1
Acute nephritis (590,591)	2	0	2	2	0	2	9.5	-	15.0
Chronic nephritis, etc. (592-594)	2	0	2	3	0	3	14.3	-	22.5
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	4.8	13.0	-
Birth inj., atelect. (760-762)	2	0	2	2	0	2	3.0	-	3.9
#Other dis. early inf. (763-776)	19	2	17	22	3	19	33.1	19.6	37.1
Violent & acc. deaths (800-999)	22	9	13	21	4	17	99.9	51.8	127.8
Acc. deaths, total (800-962)	21	9	12	19	4	15	90.4	51.8	112.7
Motor vehicle acc. (810-835)	8	7	1	5	3	2	23.8	38.9	15.0
Ill-defined (780-795)	22	3	19	21	3	18	99.9	38.9	135.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
RANKIN COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	272	73	199	837	505	332	24.4	23.5	25.9
Fetal deaths	13	3	10	37	24	13	44.2	47.5	39.2
Deaths from all causes	395	215	180	261	164	97	7.6	7.6	7.6
Deaths under 1 year	14	4	10	29	12	17	34.6	23.8	51.2
Deaths under 1 month	9	2	7	20	9	11	23.9	17.8	33.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	5	0	5	1	1	0	2.9	4.6	-
Tbc., respiratory (001-008)	4	0	4	1	1	0	2.9	4.6	-
Syphilis (020-029)	1	0	1	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	1	1	1	1	0	2.9	4.6	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	42	22	20	36	25	11	104.9	116.2	85.8
Diabetes (260)	2	2	0	4	3	1	11.6	14.0	7.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	74	34	40	34	21	13	99.1	97.6	101.4
Heart diseases (410-443)	97	55	42	61	44	17	177.7	204.6	132.6
Arteriosclerosis (450)	11	8	3	8	6	2	23.3	27.9	15.6
Influenza (480-483)	1	0	1	1	0	1	2.9	-	7.8
#Pneumonia (490-493)	24	18	6	7	4	3	20.4	18.6	23.4
#Diarrhea & enteritis (571)	1	0	1	1	0	1	2.9	-	7.8
Acute nephritis (590,591)	0	0	0	1	0	1	2.9	-	7.8
Chronic nephritis, etc. (592-594)	0	0	0	4	3	1	11.6	14.0	7.8
Maternal deaths (640-689)	0	0	0	1	1	0	11.9	19.8	-
Cong. malformations (750-759)	1	1	0	2	1	1	5.8	4.6	7.8
Birth inj., atelect. (760-762)	4	0	4	9	5	4	10.8	9.9	12.0
#Other dis. early inf. (763-776)	2	1	1	8	3	5	9.6	5.9	15.1
Violent & acc. deaths (800-999)	47	30	17	36	21	15	104.9	97.6	117.0
Acc. deaths, total (800-962)	39	24	15	32	19	13	93.2	88.4	101.4
Motor vehicle acc. (810-835)	16	13	3	16	11	5	46.6	51.2	39.0
Ill-defined (780-795)	22	12	10	12	5	7	35.0	23.2	54.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SCOTT COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	560	338	222	526	280	246	24.8	21.4	30.2
Fetal deaths	15	4	11	15	3	12	28.5	10.7	48.8
Deaths from all causes	191	111	80	218	127	91	10.3	9.7	11.2
Deaths under 1 year	23	7	16	28	8	20	53.2	28.6	81.3
Deaths under 1 month	10	4	6	12	4	8	22.8	14.3	32.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	2	2	0	9.4	15.3	-
Tbc., respiratory (001-008)	1	1	0	2	2	0	9.4	15.3	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	25	14	11	27	15	12	127.4	114.9	147.5
Diabetes (260)	4	2	2	3	1	2	14.2	7.7	24.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	31	22	9	31	21	10	146.3	160.9	122.9
Heart diseases (410-443)	62	45	17	63	45	18	297.4	344.8	221.2
Arteriosclerosis (450)	1	1	0	1	1	0	4.7	7.7	-
Influenza (480-483)	2	0	2	2	0	2	9.4	-	24.6
#Pneumonia (490-493)	4	2	2	4	3	1	18.9	23.0	12.3
#Diarrhea & enteritis (571)	1	0	1	3	1	2	14.2	7.7	24.6
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	4.7	-	12.3
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	3	2	1	2	1	1	9.4	7.7	12.3
Birth inj., atelect. (760-762)	4	2	2	5	3	2	9.5	10.7	8.1
#Other dis. early inf. (763-776)	9	3	6	10	2	8	19.0	7.1	32.5
Violent & acc. deaths (800-999)	16	8	8	28	19	9	132.2	145.6	110.6
Acc. deaths, total (800-962)	15	8	7	28	19	9	132.2	145.6	110.6
Motor vehicle acc. (810-835)	5	4	1	13	11	2	61.4	84.3	24.6
Ill-defined (780-795)	18	5	13	19	5	14	89.7	38.3	172.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SHARKEY COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	252	51	201	340	65	275	31.7	20.0	36.7
Fetal deaths	14	2	12	13	1	12	38.2	15.4	43.6
Deaths from all causes	111	22	89	131	32	99	12.2	9.8	13.2
Deaths under 1 year	22	1	21	22	2	20	64.7	30.8	72.7
Deaths under 1 month	10	1	9	9	1	8	26.5	15.4	29.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	1	0	9.3	30.8	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	11	3	8	16	6	10	149.0	184.8	133.5
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	3	12	21	6	15	195.6	184.8	200.2
Heart diseases (410-443)	30	11	19	24	7	17	223.5	215.6	226.9
Arteriosclerosis (450)	0	0	0	2	2	0	18.6	61.6	-
Influenza (480-483)	4	0	4	4	0	4	37.2	-	53.4
#Pneumonia (490-493)	6	0	6	5	0	5	46.6	-	66.7
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	1	0	1	1	0	1	9.3	-	13.3
Chronic nephritis, etc. (592-594)	5	0	5	5	0	5	46.6	-	66.7
Maternal deaths (640-689)	0	0	0	1	0	1	29.4	-	36.4
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	2	1	1	2	1	1	5.9	15.4	3.6
#Other dis. early inf. (763-776)	7	0	7	7	0	7	20.6	-	25.4
Violent & acc. deaths (800-999)	9	3	6	15	7	8	139.7	215.6	106.8
Acc. deaths, total (800-962)	9	3	6	12	6	6	111.8	184.8	80.1
Motor vehicle acc. (810-835)	1	0	1	4	3	1	37.2	92.4	13.3
Ill-defined (780-795)	13	0	13	15	1	14	139.7	30.8	186.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SIMPSON COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	480	252	228	489	261	228	23.9	19.7	31.7
Fetal deaths	13	4	9	13	7	6	26.6	26.8	26.3
Deaths from all causes	247	150	97	222	150	72	10.8	11.3	10.0
Deaths under 1 year	17	5	12	18	7	11	36.8	26.8	48.2
Deaths under 1 month	11	4	7	10	6	4	20.4	23.0	17.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	46	25	21	5	5	0	24.4	37.7	-
Tbc., respiratory (001-008)	44	24	20	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	4.9	-	13.9
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	36	20	16	34	21	13	166.2	158.4	180.6
Diabetes (260)	3	2	1	3	1	2	14.7	7.5	27.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	33	19	14	37	25	12	180.9	188.6	166.7
Heart diseases (410-443)	58	42	16	58	46	12	283.6	347.1	166.7
Arteriosclerosis (450)	2	0	2	5	3	2	24.4	22.6	27.8
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	11	9	2	10	7	3	48.9	52.8	41.7
#Diarrhea & enteritis (571)	1	0	1	2	1	1	9.8	7.5	13.9
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	2	1	3	2	1	14.7	15.1	13.9
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	1	1	2	1	1	9.8	7.5	13.9
Birth inj., atelect. (760-762)	2	1	1	1	1	0	2.0	3.8	-
#Other dis. early inf. (763-776)	6	1	5	7	3	4	14.3	11.5	17.5
Violent & acc. deaths (800-999)	9	6	3	16	13	3	78.2	98.1	41.7
Acc. deaths, total (800-962)	9	6	3	13	10	3	63.6	75.4	41.7
Motor vehicle acc. (810-835)	3	3	0	5	5	0	24.4	37.7	-
Ill-defined (780-795)	17	7	10	17	6	11	83.1	45.3	152.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SMITH COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	66	39	27	328	187	141	22.9	16.9	43.4
Fetal deaths	2	1	1	12	7	5	36.6	37.4	35.5
Deaths from all causes	69	57	12	135	110	25	9.4	9.9	7.7
Deaths under 1 year	2	0	2	11	5	6	33.5	26.7	42.6
Deaths under 1 month	1	0	1	4	2	2	12.2	10.7	14.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	1	1	0	7.0	9.0	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	19	15	4	28	22	6	195.8	199.0	184.8
Diabetes (260)	1	1	0	3	2	1	21.0	18.1	30.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	9	9	0	18	16	2	125.8	144.7	61.6
Heart diseases (410-443)	22	21	1	31	30	1	216.7	271.3	30.8
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	1	1	0	1	1	0	7.0	9.0	-
#Pneumonia (490-493)	1	0	1	5	3	2	35.0	27.1	61.6
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	3	2	1	21.0	18.1	30.8
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	5	4	1	35.0	36.2	30.8
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	0	0	0	3	2	1	9.1	10.7	7.1
Violent & acc. deaths (800-999)	6	4	2	13	11	2	90.9	99.5	61.6
Acc. deaths, total (800-962)	3	1	2	10	8	2	69.9	72.4	61.6
Motor vehicle acc. (810-835)	2	1	1	7	6	1	48.9	54.3	30.8
Ill-defined (780-795)	4	3	1	7	6	1	48.9	54.3	30.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
STONE COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	149	112	37	154	105	49	22.0	19.8	28.6
Fetal deaths	8	8	0	9	7	2	58.4	66.7	40.8
Deaths from all causes	56	45	11	67	51	16	9.6	9.6	9.4
Deaths under 1 year	4	3	1	5	3	2	32.5	28.6	40.8
Deaths under 1 month	1	1	0	2	1	1	13.0	9.5	20.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	1	0	1	1	0	14.2	18.9	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	3	2	1	8	7	1	114.1	132.0	58.4
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	5	5	0	7	7	0	99.8	132.0	-
Heart diseases (410-443)	28	23	5	28	22	6	399.2	414.9	350.7
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	1	0	3	3	0	42.8	56.6	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	0	1	14.2	-	58.4
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	1	1	0	1	1	0	6.5	9.5	-
#Other dis. early inf. (763-776)	0	0	0	1	0	1	6.5	-	20.4
Violent & acc. deaths (800-999)	10	8	2	9	7	2	128.3	132.0	116.9
Acc. deaths, total (800-962)	10	8	2	9	7	2	128.3	132.0	116.9
Motor vehicle acc. (810-835)	6	5	1	4	3	1	57.0	56.6	58.4
Ill-defined (780-795)	3	1	2	2	0	2	28.5	-	116.9

*Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
SUNFLOWER COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,334	224	1,110	1,408	245	1,163	30.8	16.6	37.5
Fetal deaths	46	10	36	51	12	39	36.2	49.0	33.5
Deaths from all causes	387	98	289	435	130	305	9.5	8.8	9.8
Deaths under 1 year	66	4	62	68	4	64	48.3	16.3	55.0
Deaths under 1 month	39	4	35	41	4	37	29.1	16.3	31.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	0	3	1	0	1	2.2	-	3.2
Tbc., respiratory (001-008)	3	0	3	1	0	1	2.2	-	3.2
Syphilis (020-029)	1	0	1	2	0	2	4.4	-	6.4
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	3	0	3	2	0	2	4.4	-	6.4
Scarlet fever & streptococcal sore throat (050,051)	1	0	1	1	0	1	2.2	-	3.2
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	47	20	27	58	26	32	126.8	176.5	103.2
Diabetes (260)	4	2	2	10	4	6	21.8	27.2	19.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	50	15	35	62	20	42	135.5	135.8	135.4
Heart diseases (410-443)	91	38	53	106	48	58	231.7	325.9	187.0
Arteriosclerosis (450)	4	2	2	5	2	3	10.9	13.6	9.7
Influenza (480-483)	5	0	5	4	0	4	8.7	-	12.9
#Pneumonia (490-493)	17	1	16	16	1	15	35.0	6.8	48.4
#Diarrhea & enteritis (571)	8	1	7	8	1	7	17.5	6.8	22.6
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	6	0	6	7	1	6	15.3	6.8	19.3
Maternal deaths (640-689)	4	0	4	1	0	1	7.1	-	8.6
Cong. malformations (750-759)	1	0	1	0	0	0	-	-	-
Birth inj., atelect. (760-762)	8	0	8	8	0	8	5.7	-	6.9
#Other dis. early inf. (763-776)	32	4	28	34	4	30	24.1	16.3	25.8
Violent & acc. deaths (800-999)	28	6	22	28	7	21	61.2	47.5	67.7
Acc. deaths, total (800-962)	22	6	16	23	7	16	50.3	47.5	51.6
Motor vehicle acc. (810-835)	6	3	3	8	5	3	17.5	33.9	9.7
Ill-defined (780-795)	41	0	41	41	2	39	89.6	13.6	125.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TALLAHATCHIE COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	733	129	604	772	159	613	32.0	18.5	39.5
Fetal deaths	36	7	29	34	5	29	44.0	31.4	47.3
Deaths from all causes	227	55	172	259	80	179	10.8	9.3	11.5
Deaths under 1 year	39	1	38	42	3	39	54.4	18.9	63.6
Deaths under 1 month	19	0	19	21	2	19	27.2	12.6	31.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	3	1	2	12.4	11.6	12.9
Tbc., respiratory (001-008)	0	0	0	1	0	1	4.2	-	6.4
Syphilis (020-029)	2	0	2	2	0	2	8.3	-	12.9
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	20	9	11	26	16	10	108.0	186.5	64.5
Diabetes (260)	1	0	1	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	22	3	19	26	4	22	108.0	46.6	141.9
Heart diseases (410-443)	70	29	41	77	35	42	319.8	407.9	271.0
Arteriosclerosis (450)	3	0	3	4	1	3	16.6	11.6	19.4
Influenza (480-483)	8	2	6	7	1	6	29.1	11.6	38.7
#Pneumonia (490-493)	6	1	5	8	3	5	33.2	35.0	32.2
#Diarrhea & enteritis (571)	2	0	2	3	0	3	12.4	-	19.4
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	5	0	5	5	0	5	20.8	-	32.2
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	1	0	1	4.2	-	6.4
Birth inj., atelect. (760-762)	2	0	2	3	1	2	3.9	6.3	3.3
#Other dis. early inf. (763-776)	14	0	14	15	1	14	19.4	6.3	22.8
Violent & acc. deaths (800-999)	21	4	17	28	10	18	116.3	116.6	116.1
Acc. deaths, total (800-962)	17	2	15	22	7	15	91.4	81.6	96.8
Motor vehicle acc. (810-835)	4	1	3	6	3	3	24.9	35.0	19.4
Ill-defined (780-795)	26	2	24	24	0	24	99.7	-	154.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TATE COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	402	18	384	551	134	417	30.4	17.4	39.9
Fetal deaths	19	1	18	26	3	23	47.2	22.4	55.2
Deaths from all causes	110	45	65	160	77	83	8.8	10.0	7.9
Deaths under 1 year	11	1	10	21	4	17	38.1	29.8	40.8
Deaths under 1 month	7	1	6	13	4	9	23.6	29.8	21.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	19	9	10	26	14	12	143.3	181.9	114.9
Diabetes (260)	0	0	0	1	1	0	5.5	13.0	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	13	6	7	22	13	9	121.3	168.9	86.2
Heart diseases (410-443)	39	18	21	51	26	25	281.2	337.8	239.4
Arteriosclerosis (450)	1	0	1	2	1	1	11.0	13.0	9.6
Influenza (480-483)	2	0	2	3	0	3	16.5	-	28.7
#Pneumonia (490-493)	6	0	6	6	0	6	33.1	-	57.5
#Diarrhea & enteritis (571)	0	0	0	3	0	3	16.5	-	28.7
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	11.0	13.0	9.6
Maternal deaths (640-689)	0	0	0	1	0	1	18.1	-	24.0
Cong. malformations (750-759)	2	1	1	2	1	1	11.0	13.0	9.6
Birth inj., atelect. (760-762)	1	0	1	2	1	1	3.6	7.5	2.4
#Other dis. early inf. (763-776)	3	0	3	10	2	8	18.1	14.9	19.2
Violent & acc. deaths (800-999)	6	4	2	11	9	2	60.6	116.9	19.2
Acc. deaths, total (800-962)	2	2	0	7	7	0	38.6	91.0	-
Motor vehicle acc. (810-835)	1	1	0	3	3	0	16.5	39.0	-
Ill-defined (780-795)	13	4	9	13	4	9	71.7	52.0	86.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TIPPAH COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	393	291	102	332	242	90	22.0	19.6	32.6
Fetal deaths	22	14	8	19	11	8	57.2	45.4	88.9
Deaths from all causes	142	119	23	148	128	20	9.8	10.4	7.2
Deaths under 1 year	12	5	7	10	6	4	30.1	24.8	44.4
Deaths under 1 month	5	2	3	5	2	3	15.1	8.3	33.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	2	0	2	2	0	13.2	16.2	-
Tbc., respiratory (001-008)	2	2	0	2	2	0	13.2	16.2	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	14	3	20	17	3	132.5	137.8	108.8
Diabetes (260)	1	1	0	1	1	0	6.6	8.1	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	19	16	3	17	15	2	112.6	121.6	72.6
Heart diseases (410-443)	56	51	5	59	55	4	390.9	445.8	145.1
Arteriosclerosis (450)	2	1	1	2	1	1	13.2	8.1	36.3
Influenza (480-483)	2	2	0	1	1	0	6.6	8.1	-
#Pneumonia (490-493)	6	5	1	7	6	1	46.4	48.6	36.3
#Diarrhea & enteritis (571)	2	1	1	1	1	0	6.6	8.1	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	3	0	3	3	0	19.9	24.3	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	2	0	2	2	0	13.2	16.2	-
Birth inj., atelect. (760-762)	3	1	2	0	0	0	-	-	-
#Other dis. early inf. (763-776)	4	1	3	5	2	3	15.1	8.3	33.3
Violent & acc. deaths (800-999)	12	12	0	9	9	0	59.6	73.0	-
Acc. deaths, total (800-962)	8	8	0	5	5	0	33.1	40.5	-
Motor vehicle acc. (810-835)	8	8	0	4	4	0	26.5	32.4	-
Ill-defined (780-795)	3	3	0	5	5	0	33.1	40.5	-

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TISHOMINGO COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	150	140	10	239	223	16	17.2	16.9	23.6
Fetal deaths	2	2	0	3	3	0	12.6	13.4	-
Deaths from all causes	106	94	12	131	123	8	9.4	9.3	11.8
Deaths under 1 year	2	2	0	5	5	0	20.9	22.4	-
Deaths under 1 month	1	1	0	5	5	0	20.9	22.4	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	13	13	0	16	16	0	115.2	121.1	-
Diabetes (260)	1	1	0	2	2	0	14.4	15.1	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	30	24	6	37	32	5	266.4	242.2	736.4
Heart diseases (410-443)	32	30	2	37	35	2	266.4	265.0	294.6
Arteriosclerosis (450)	1	1	0	2	2	0	14.4	15.1	-
Influenza (480-483)	1	1	0	1	1	0	7.2	7.6	-
#Pneumonia (490-493)	4	3	1	5	5	0	36.0	37.8	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc., (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	2	2	0	14.4	15.1	-
Birth inj., atelect. (760-762)	0	0	0	2	2	0	8.4	9.0	-
#Other dis. early inf. (763-776)	2	2	0	1	1	0	4.2	4.5	-
Violent & acc. deaths (800-999)	14	11	3	16	16	0	115.2	121.1	-
Acc. deaths, total (800-962)	12	9	3	14	14	0	100.8	106.0	-
Motor vehicle acc. (810-835)	8	8	0	12	12	0	86.4	90.8	-
Ill-defined (780-795)	3	3	0	5	4	1	36.0	30.3	147.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
TUNICA COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	616	70	546	605	67	538	36.0	19.1	40.4
Fetal deaths	29	2	27	29	2	27	47.9	29.8	50.2
Deaths from all causes	209	32	177	223	31	192	13.2	8.8	14.4
Deaths under 1 year	40	4	36	42	5	37	69.4	74.6	68.8
Deaths under 1 month	9	2	7	10	3	7	16.5	44.8	13.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	0	4	5	0	5	29.7	-	37.5
Tbc., respiratory (001-008)	4	0	4	5	0	5	29.7	-	37.5
Syphilis (020-029)	1	0	1	2	0	2	11.9	-	15.0
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	5.9	-	7.5
Scarlet fever & streptococcal sore throat (050,051)	1	0	1	1	0	1	5.9	-	7.5
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	15	4	11	23	7	16	136.7	199.7	120.1
Diabetes (260)	1	0	1	2	1	1	11.9	28.5	7.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	21	2	19	23	2	21	136.7	57.1	157.6
Heart diseases (410-443)	57	8	49	56	8	48	332.8	228.2	360.3
Arteriosclerosis (450)	0	0	0	1	0	1	5.9	-	7.5
Influenza (480-483)	4	0	4	4	0	4	23.8	-	30.0
#Pneumonia (490-493)	12	2	10	12	2	10	71.3	57.1	75.1
#Diarrhea & enteritis (571)	15	0	15	15	0	15	89.1	-	112.6
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	6	3	3	6	2	4	35.6	57.1	30.0
Maternal deaths (640-689)	3	0	3	3	0	3	49.6	-	55.8
Cong. malformations (750-759)	1	0	1	1	0	1	5.9	-	7.5
Birth inj., atelect. (760-762)	4	0	4	5	1	4	8.3	14.9	7.4
#Other dis. early inf. (763-776)	6	2	4	6	2	4	9.9	29.8	7.4
Violent & acc. deaths (800-999)	24	6	18	23	2	21	136.7	57.1	157.6
Acc. deaths, total (800-962)	18	5	13	18	2	16	107.0	57.1	120.1
Motor vehicle acc. (810-835)	6	3	3	6	2	4	35.6	57.1	30.0
Ill-defined (780-795)	23	2	21	21	0	21	124.8	-	157.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
UNION COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	386	287	99	385	290	95	20.4	18.6	28.7
Fetal deaths	23	19	4	21	17	4	54.5	58.6	42.1
Deaths from all causes	187	141	46	219	170	49	11.6	10.9	14.8
Deaths under 1 year	13	4	9	16	7	9	41.6	24.1	94.7
Deaths under 1 month	9	4	5	11	6	5	28.6	20.7	52.6
SELECTED CAUSES OF DEATH									
AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	1	0	1	1	0	5.3	6.4	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	23	21	2	33	31	2	174.6	198.8	60.4
Diabetes (260)	3	3	0	3	3	0	15.9	19.2	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	25	21	4	30	24	6	158.7	153.9	181.2
Heart diseases (410-443)	71	56	15	78	63	15	412.6	404.0	452.9
Arteriosclerosis (450)	2	2	0	2	2	0	10.6	12.8	-
Influenza (480-483)	5	3	2	5	3	2	26.4	19.2	60.4
#Pneumonia (490-493)	5	2	3	8	5	3	42.3	32.1	90.6
#Diarrhea & enteritis (571)	2	1	1	2	1	1	10.6	6.4	30.2
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	1	3	4	1	3	21.2	6.4	90.6
Maternal deaths (640-689)	1	0	1	1	0	1	26.0	-	105.3
Cong. malformations (750-759)	1	1	0	2	2	0	10.6	12.8	-
Birth inj., atelect. (760-762)	2	2	0	4	4	0	10.4	13.8	-
#Other dis. early inf. (763-776)	6	2	4	6	2	4	15.6	6.9	42.1
Violent & acc. deaths (800-999)	14	12	2	14	12	2	74.0	77.0	60.4
Acc. deaths, total (800-962)	13	11	2	13	11	2	68.8	70.5	60.4
Motor vehicle acc. (810-835)	5	5	0	5	5	0	26.4	32.1	-
Ill-defined (780-795)	11	5	6	11	5	6	58.2	32.1	181.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WALTHALL COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	523	291	232	326	128	198	24.1	17.3	32.4
Fetal deaths	26	14	12	15	6	9	46.0	46.9	45.4
Deaths from all causes	115	71	44	107	63	44	7.9	8.5	7.2
Deaths under 1 year	15	7	8	11	4	7	33.7	31.2	35.4
Deaths under 1 month	6	4	2	5	3	2	15.3	23.4	10.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	14	10	4	13	10	3	96.2	134.9	49.2
Diabetes (260)	5	2	3	5	2	3	37.0	27.0	49.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	12	9	3	8	4	4	59.2	54.0	65.6
Heart diseases (410-443)	31	21	10	32	22	10	236.8	296.8	163.9
Arteriosclerosis (450)	1	1	0	1	1	0	7.4	13.5	-
Influenza (480-483)	2	1	1	1	1	0	7.4	13.5	-
#Pneumonia (490-493)	6	3	3	6	3	3	44.4	40.5	49.2
#Diarrhea & enteritis (571)	2	0	2	2	0	2	14.8	-	32.8
Acute nephritis (590,591)	1	1	0	1	1	0	7.4	13.5	-
Chronic nephritis, etc. (592-594)	0	0	0	1	0	1	7.4	-	16.4
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	4	1	3	3	0	3	22.2	-	49.2
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	6	4	2	5	3	2	15.3	23.4	10.1
Violent & acc. deaths (800-999)	11	8	3	12	6	6	88.8	80.9	98.4
Acc. deaths, total (800-962)	10	7	3	12	6	6	88.8	80.9	98.4
Motor vehicle acc. (810-835)	2	2	0	2	1	1	14.8	13.5	16.4
Ill-defined (780-795)	13	7	6	12	7	5	88.8	94.4	82.0

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WARREN COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,402	596	806	1,111	508	603	26.3	22.6	30.5
Fetal deaths	63	21	42	39	19	20	35.1	37.4	33.2
Deaths from all causes	722	260	462	541	216	325	12.8	9.6	16.4
Deaths under 1 year	78	17	61	50	15	35	45.0	29.5	58.0
Deaths under 1 month	45	10	35	30	11	19	27.0	21.6	31.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	4	1	3	9.5	4.4	15.2
Tbc., respiratory (001-008)	1	0	1	3	0	3	7.1	-	15.2
Syphilis (020-029)	2	0	2	2	0	2	4.7	-	10.1
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	84	41	43	57	27	30	135.0	120.3	151.8
Diabetes (260)	13	3	10	6	2	4	14.2	8.9	20.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	99	37	62	83	30	53	196.6	133.6	268.2
Heart diseases (410-443)	184	85	99	158	81	77	374.4	360.8	389.7
Arteriosclerosis (450)	13	7	6	11	6	5	26.1	26.7	25.3
Influenza (480-483)	6	2	4	5	1	4	11.8	4.4	20.2
#Pneumonia (490-493)	16	7	9	16	8	8	37.9	35.6	40.5
#Diarrhea & enteritis (571)	8	0	8	4	0	4	9.5	-	20.2
Acute nephritis (590,591)	1	0	1	1	0	1	2.4	-	5.1
Chronic nephritis, etc. (592-594)	7	0	7	3	0	3	7.1	-	15.2
Maternal deaths (640-689)	1	0	1	0	0	0	-	-	-
Cong. malformations (750-759)	3	2	1	1	1	0	2.4	4.4	-
Birth inj., atelect. (760-762)	9	4	5	7	5	2	6.3	9.8	3.3
#Other dis. early inf. (763-776)	42	7	35	23	5	18	20.7	9.8	29.8
Violent & acc. deaths (800-999)	67	24	43	46	19	27	109.0	84.6	136.6
Acc. dsaths, total (800-962)	57	20	37	37	15	22	87.7	66.8	111.3
Motor vehicle acc. (810-835)	19	8	11	10	4	6	23.7	17.8	30.4
Ill-defined (780-795)	68	7	61	58	5	53	137.4	22.3	268.2

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WASHINGTON COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	2,734	1,084	1,650	2,641	1,042	1,599	33.6	29.6	36.8
Fetal deaths	216	118	98	213	112	101	80.6	107.5	63.2
Deaths from all causes	801	246	555	841	226	615	10.7	6.4	14.2
Deaths under 1 year	114	27	87	118	26	92	44.7	25.0	57.5
Deaths under 1 month	65	20	45	68	21	47	25.7	20.2	29.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	2	0	2	2.5	-	4.6
Tbc., respiratory (001-008)	0	0	0	2	0	2	2.5	-	4.6
Syphilis (020-029)	1	0	1	2	0	2	2.5	-	4.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	4	2	2	4	2	2	5.1	5.7	4.6
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	1.3	-	2.3
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	105	43	62	109	41	68	138.6	116.3	156.7
Diabetes (260)	7	0	7	11	0	11	14.0	-	25.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	86	21	65	94	19	75	119.5	53.9	172.8
Heart diseases (410-443)	189	64	125	192	60	132	244.2	170.3	304.2
Arteriosclerosis (450)	6	2	4	7	1	6	8.9	2.8	13.8
Influenza (480-483)	11	1	10	11	1	10	14.0	2.8	23.0
#Pneumonia (490-493)	33	12	21	30	8	22	38.1	22.7	50.7
#Diarrhea & enteritis (571)	11	1	10	10	1	9	12.7	2.8	20.7
Acute nephritis (590,591)	1	0	1	1	0	1	1.3	-	2.3
Chronic nephritis, etc. (592-594)	13	2	11	14	1	13	17.8	2.8	30.0
Maternal deaths (640-689)	4	0	4	4	0	4	15.1	-	25.0
Cong. malformations (750-759)	14	7	7	16	7	9	20.3	19.9	20.7
Birth inj., atelect. (760-762)	4	2	2	5	2	3	1.9	1.9	1.9
#Other dis. early inf. (763-776)	57	15	42	60	15	45	22.7	14.4	28.1
Violent & acc. deaths (800-999)	70	34	36	72	35	37	91.6	99.3	85.2
Acc. deaths, total (800-962)	53	27	26	56	28	28	71.2	79.4	64.5
Motor vehicle acc. (810-835)	25	18	7	27	18	9	34.3	51.1	20.7
Ill-defined (780-795)	116	15	101	119	14	105	151.3	39.7	241.9

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WAYNE COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	437	302	135	431	248	183	26.5	23.7	31.5
Fetal deaths	8	3	5	16	5	11	37.1	20.2	60.1
Deaths from all causes	135	83	52	163	93	70	10.0	8.9	12.0
Deaths under 1 year	15	10	5	19	8	11	44.1	32.2	60.1
Deaths under 1 month	7	5	2	12	5	7	27.8	20.2	38.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	6.2	-	17.2
Tbc., respiratory (001-008)	0	0	0	1	0	1	6.2	-	17.2
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	1	0	1	1	0	1	6.2	-	17.2
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	0	1	1	0	1	6.2	-	17.2
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	13	11	2	18	16	2	110.7	153.1	34.4
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	9	6	3	10	7	3	61.5	67.0	51.6
Heart diseases (410-443)	47	31	16	51	32	19	313.7	306.2	327.1
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	1	0	1	1	0	1	6.2	-	17.2
#Pneumonia (490-493)	4	2	2	5	3	2	30.8	28.7	34.4
#Diarrhea & enteritis (571)	1	1	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	12.3	9.6	17.2
Maternal deaths (640-689)	1	0	1	1	0	1	23.2	-	54.6
Cong. malformations (750-759)	1	1	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	2	1	1	2	1	1	4.6	4.0	5.5
#Other dis. early inf. (763-776)	5	4	1	10	4	6	23.2	16.1	32.8
Violent & acc. deaths (800-999)	17	8	9	22	12	10	135.3	114.8	172.1
Acc. deaths, total (800-962)	14	6	8	18	10	8	110.7	95.7	137.7
Motor vehicle acc. (810-835)	11	6	5	13	8	5	80.0	76.6	86.1
Ill-defined (780-795)	12	4	8	15	5	10	92.3	47.8	172.1

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WEBSTER COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	148	84	64	188	121	67	17.8	15.2	25.4
Fetal deaths	0	0	0	2	2	0	10.6	16.5	-
Deaths from all causes	79	70	9	105	88	17	9.9	11.1	6.4
Deaths under 1 year	6	4	2	10	5	5	53.2	41.3	74.6
Deaths under 1 month	3	2	1	8	4	4	42.6	33.0	59.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	9	1	15	13	2	141.8	163.8	75.7
Diabetes (260)	0	0	0	2	1	1	18.9	12.6	37.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	28	23	5	27	22	5	255.2	277.1	189.2
Heart diseases (410-443)	24	23	1	28	26	2	264.6	327.5	75.7
Arteriosclerosis (450)	2	2	0	2	2	0	18.9	25.2	-
Influenza (480-483)	2	2	0	1	1	0	9.4	12.6	-
#Pneumonia (490-493)	0	0	0	2	2	0	18.9	25.2	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	2	2	0	18.9	25.2	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	2	1	1	5	2	3	26.6	16.5	44.8
Violent & acc. deaths (800-999)	7	7	0	10	7	3	94.5	88.2	113.6
Acc. deaths, total (800-962)	6	6	0	9	6	3	85.1	75.6	113.6
Motor vehicle acc. (810-835)	3	3	0	5	3	2	47.2	37.8	75.7
Ill-defined (780-795)	2	1	1	4	3	1	37.8	37.8	37.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WILKINSON COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	432	194	238	352	75	277	26.6	19.7	29.4
Fetal deaths	30	8	22	26	3	23	73.9	40.0	83.0
Deaths from all causes	172	77	95	157	56	101	11.9	14.7	10.7
Deaths under 1 year	25	2	23	21	1	20	59.6	13.3	72.2
Deaths under 1 month	8	1	7	8	1	7	22.7	13.3	25.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	0	2	2	0	2	15.1	-	21.2
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	20	16	4	13	10	3	98.2	262.7	31.8
Diabetes (260)	3	3	0	5	3	2	37.8	78.8	21.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	25	11	14	24	7	17	181.3	183.9	180.3
Heart diseases (410-443)	34	18	16	34	16	18	256.9	420.3	190.9
Arteriosclerosis (450)	4	1	3	4	1	3	30.2	26.3	31.8
Influenza (480-483)	1	1	0	1	1	0	7.6	26.3	-
#Pneumonia (490-493)	10	0	10	8	1	7	60.4	26.3	74.2
#Diarrhea & enteritis (571)	1	0	1	1	0	1	7.6	-	10.5
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	2	2	3	1	2	22.7	26.3	21.2
Maternal deaths (640-689)	0	0	0	1	0	1	28.4	-	36.1
Cong. malformations (750-759)	1	0	1	1	0	1	7.6	-	10.6
Birth inj., atelect. (760-762)	2	0	2	1	0	1	2.8	-	3.6
#Other dis. early inf. (763-776)	9	2	7	9	1	8	25.6	13.3	28.9
Violent & acc. deaths (800-999)	13	6	7	12	4	8	90.7	105.1	84.8
Acc. deaths, total (800-962)	12	5	7	11	3	8	83.1	78.8	84.8
Motor vehicle acc. (810-835)	3	1	2	3	1	2	22.7	26.3	21.2
Ill-defined (780-795)	23	5	18	22	5	17	166.2	131.3	180.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
WINSTON COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	377	159	218	421	177	244	21.9	16.3	29.1
Fetal deaths	10	0	10	13	0	13	30.9	-	53.3
Deaths from all causes	196	130	66	216	146	70	11.2	13.4	8.3
Deaths under 1 year	20	5	15	21	6	15	49.9	33.9	61.5
Deaths under 1 month	13	4	9	14	5	9	33.2	28.2	36.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	5.2	9.2	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	5.2	9.2	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	0	1	1	0	1	5.2	-	11.9
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	23	20	3	29	25	4	150.7	230.4	47.6
Diabetes (260)	4	3	1	5	3	2	26.0	27.6	23.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	31	20	11	31	21	10	161.1	193.5	119.1
Heart diseases (410-443)	69	53	16	79	59	20	410.5	543.6	238.3
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	4	1	3	4	1	3	20.8	9.2	35.7
#Pneumonia (490-493)	5	3	2	5	3	2	26.0	27.6	23.8
#Diarrhea & enteritis (571)	3	1	2	3	1	2	15.6	9.2	23.8
Acute nephritis (590,591)	1	1	0	1	1	0	5.2	9.2	-
Chronic nephritis, etc. (592-594)	2	2	0	1	1	0	5.2	9.2	-
Maternal deaths (640-689)	1	1	0	1	1	0	23.8	56.5	-
Cong. malformations (750-759)	4	1	3	5	2	3	26.0	18.4	35.7
Birth inj., atelect. (760-762)	2	1	1	2	1	1	4.8	5.6	4.1
#Other dis. early inf. (763-776)	6	3	3	7	4	3	16.6	22.6	12.3
Violent & acc. deaths (800-999)	13	7	6	14	8	6	72.7	73.7	71.5
Acc. deaths, total (800-962)	9	5	4	10	6	4	52.0	55.3	47.6
Motor vehicle acc. (810-835)	1	1	0	2	2	0	10.4	18.4	-
Ill-defined (780-795)	6	1	5	7	3	4	36.4	27.6	47.6

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
YALOBUSHA COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	288	100	188	307	124	183	24.6	17.8	33.0
Fetal deaths	6	3	3	8	4	4	26.0	32.2	21.8
Deaths from all causes	135	91	44	167	115	52	13.4	16.5	9.4
Deaths under 1 year	8	0	8	11	1	10	35.8	8.1	54.6
Deaths under 1 month	7	0	7	9	1	8	29.3	8.1	43.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	2	1	1	16.0	14.4	18.0
Tbc., respiratory (001-008)	1	0	1	2	1	1	16.0	14.4	18.0
Syphilis (020-029)	1	0	1	1	0	1	8.0	-	18.0
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	20	17	3	21	18	3	168.0	258.5	54.2
Diabetes (260)	2	0	2	3	0	3	24.0	-	54.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	13	9	4	16	12	4	128.0	172.4	72.2
Heart diseases (410-443)	46	36	10	61	48	13	487.9	689.4	234.6
Arteriosclerosis (450)	1	1	0	1	1	0	8.0	14.4	-
Influenza (480-483)	4	2	2	4	2	2	32.0	28.7	36.1
#Pneumonia (490-493)	6	6	0	5	5	0	40.0	71.8	-
#Diarrhea & enteritis (571)	2	1	1	2	1	1	16.0	14.4	18.0
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	8.0	-	18.0
Maternal deaths (640-689)	1	0	1	1	0	1	32.6	-	54.6
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	1	1	0	3.2	8.1	-
#Other dis. early inf. (763-776)	4	0	4	5	0	5	16.3	-	27.3
Violent & acc. deaths (800-999)	8	5	3	4	2	2	32.0	28.7	36.1
Acc. deaths, total (800-962)	8	5	3	4	2	2	32.0	28.7	36.1
Motor vehicle acc. (810-835)	3	2	1	1	1	0	8.0	14.4	-
Ill-defined (780-795)	9	4	5	13	7	6	104.0	100.5	108.3

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
YAZOO COUNTY, MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	835	262	573	878	276	602	27.7	21.4	32.0
Fetal deaths	27	2	25	34	4	30	38.7	14.5	49.8
Deaths from all causes	343	111	232	375	128	247	11.8	10.0	13.1
Deaths under 1 year	39	6	33	52	7	45	59.2	25.4	74.8
Deaths under 1 month	22	5	17	29	6	23	33.0	21.7	38.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	3.2	-	5.3
Tbc., respiratory (001-008)	0	0	0	1	0	1	3.2	-	5.3
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	33	11	22	33	11	22	104.2	85.5	117.1
Diabetes (260)	6	0	6	5	0	5	15.8	-	26.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	64	23	41	70	30	40	221.1	233.2	212.9
Heart diseases (410-443)	88	40	48	91	45	46	287.5	349.9	244.8
Arteriosclerosis (450)	5	2	3	6	4	2	19.0	31.1	10.6
Influenza (480-483)	4	1	3	4	1	3	12.6	7.8	16.0
#Pneumonia (490-493)	6	1	5	8	1	7	25.3	7.8	37.2
#Diarrhea & enteritis (571)	2	0	2	2	0	2	6.3	-	10.6
Acute nephritis (590,591)	1	0	1	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	0	2	3	0	3	9.5	-	16.0
Maternal deaths (640-689)	2	0	2	2	0	2	22.8	-	33.2
Cong. malformations (750-759)	0	0	0	1	0	1	3.2	-	5.3
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	16	5	11	23	6	17	26.2	21.7	28.2
Violent & acc. deaths (800-999)	37	14	23	37	13	24	116.9	101.1	127.7
Acc. deaths, total (800-962)	28	11	17	28	10	18	88.4	77.7	95.8
Motor vehicle acc. (810-835)	10	7	3	9	5	4	28.4	38.9	21.3
Ill-defined (780-795)	51	6	45	56	7	49	176.9	54.4	260.8

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, DEATHS UNDER 1 YEAR, AND
DEATHS FROM CERTAIN SELECTED CAUSES
STATE OF MISSISSIPPI, 1960

Shown by Race by Place of Occurrence and Place of Residence

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	59731	27638	32093	59210	27198	32012	27.2	21.6	34.8
Fetal deaths	2421	992	1429	2384	955	1429	40.3	35.1	44.6
Deaths from all causes	21583	11136	10447	21713	11204	10509	10.0	8.9	11.4
Deaths under 1 year	2455	703	1752	2457	714	1743	41.5	26.2	54.4
Deaths under 1 month	1470	528	942	1497	550	947	25.3	20.2	29.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	145	57	88	147	57	90	6.7	4.5	9.8
Tbc., respiratory (001-008)	133	54	79	135	54	81	6.2	4.3	8.8
Syphilis (020-029)	26	3	23	29	5	24	1.3	0.4	2.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	27	8	19	27	8	19	1.2	0.6	2.1
Scarlet fever & streptococcal sore throat (050,051)	5	1	4	5	1	4	0.2	0.1	0.4
Diphtheria (055)	4	3	1	4	3	1	0.2	0.2	0.1
Whooping cough (056)	8	1	7	7	0	7	0.3	-	0.8
Meningococcal infections (057)	13	8	5	13	8	5	0.6	0.6	0.5
Acute poliomyelitis (080)	3	2	1	4	2	2	0.2	0.2	0.2
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	9	4	5	9	4	5	0.4	0.3	0.5
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	2540	1570	970	2635	1644	991	121.0	130.7	107.6
Diabetes (260)	309	142	167	309	139	170	14.2	11.0	18.5
Pellagra (281)	5	1	4	5	1	4	0.2	0.1	0.4
Intracranial lesions (330-334)	2969	1519	1450	2960	1513	1447	135.9	120.3	157.2
Heart diseases (410-443)	6266	3862	2404	6312	3879	2433	289.8	308.4	264.3
Arteriosclerosis (450)	303	195	108	306	198	108	14.0	15.7	11.7
Influenza (480-483)	243	73	170	238	71	167	10.9	5.6	18.1
#Pneumonia (490-493)	720	315	405	720	318	402	33.0	25.3	43.7
#Diarrhea & enteritis (571)	222	37	185	218	35	183	10.0	2.8	19.9
Acute nephritis (590,591)	44	14	30	41	11	30	1.9	0.9	3.2
Chronic nephritis, etc. (592-594)	235	82	153	241	83	158	11.1	6.6	17.2
Maternal deaths (640-689)	60	9	51	59	8	51	10.0	2.9	15.9
Cong. malformations (750-759)	227	134	93	234	142	92	10.7	11.3	10.0
Birth inj., atelect. (760-762)	330	180	150	332	186	146	5.6	6.8	4.6
#Other dis. early inf. (763-776)	975	283	692	974	279	695	16.4	10.2	21.7
Violent & acc. deaths (800-999)	1888	1043	845	1844	994	850	84.6	79.0	92.3
Acc. deaths, total (800-962)	1542	869	673	1504	827	677	69.0	65.8	73.5
Motor vehicle acc. (810-835)	676	478	198	620	428	192	28.5	34.0	20.8
Ill-defined (780-795)	1881	450	1431	1852	428	1424	85.0	34.0	154.7

* Crude birth and death rate per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 1 month, and deaths from infant causes (760-776) per 1,000 live births. Rates for maternal deaths per 10,000 live births. Specific death rates per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.