PUBLIC HEALTH STATISTICS

STATE OF

MISSISSIPPI

1961

VITAL STATISTICS

SUMMARIES

Mississippi State Board of Health
Jackson, Mississippi

A. L. Gray, M. D., Executive Officer

VITAL STATISTICS

MISSISSIPPI

1961

****

MISSISSIPPI STATE BOARD OF HEALTH
JACKSON, MISSISSIPPI

A. L. GRAY, M. D., EXECUTIVE OFFICER

TABLE OF CONTENTS

SYNOPSIS OF VITAL STATISTICS, MISSISSIPPI, 1961 — Page

Introduction I

Live Births II

Fetal Deaths V

Deaths VI

Marriages XVII

Divorces XX

POPULATION

Table 1. ~~Census Population,~~ Mississippi, April 1, 1960 1

LIVE BIRTHS

Table 2. By Place of Occurrence, and by Place of Residence of Mother, by Race, with Rates, Mississippi, 1961 4

Table 3. By Place of Residence of Mother, by Race and Attendant, and by Sex, Mississippi, 1961 7

FETAL DEATHS

Table 4. By Place of Occurrence, by Place of Residence of Mother, by Race and Attendant, and by Sex, Mississippi, 1961 11

DEATHS

Table 5. By Cause of Death, by Race, Resident Data, Mississippi, 1961 15

Table 6. By Place of Occurrence, and by Place of Residence, by Race, with Rates, Mississippi, 1961 26

Table 7. By Cause Groups and Certain Selected Causes, by Race, Sex, and Age, with Total Rates, Resident Data, Mississippi, 1961 29

~~Table 8. By Cause of Death, by Month, Resident Data, Mississippi, 1961~~ 37

DEATHS UNDER 1 YEAR

Table 9. By Selected Causes, by Race and Age, with Total Rates, Resident Data, Mississippi, 1961 38

MARRIAGES AND DIVORCES

Table ~~10~~ 9. Marriages by Place of Licensure, and by Place of Residence of Bride, by Race of Bride; Divorces by Place of Occurrence, by Race, Mississippi, 1961 42

Table ~~11~~ 10. Divorces by Cause of Divorce, by Race, Mississippi, 1961 44

Divorces by Number of Minor Children, by Race, Mississippi, 1961 44

Divorces by Number of Years Married, by Race, Mississippi, 1961 44

SUMMARY TABLES

Table 12. Live Births, Mississippi, 1913-1961, by Year, by Race, with Rates, Occurrence Data 1913-1943, Residence Data 1944-1961 45

Table 13. Deaths, Mississippi, 1913-1961, by Year, by Race, with Rates, Occurrence Data 1913-1943, Residence Data 1944-1961 47

Table 14. Maternal Deaths, Mississippi, 1913-1961, by Year, by Race, with Rates, Occurrence Data 1913-1943, Residence Data 1944-1961 49

Table 15. Deaths Under 1 Year, Mississippi, 1917-1961, by Year, by Race, with Rates, Occurrence Data 1917-1943, Residence Data 1944-1961 51

Table 16. Deaths Under 1 Month, Mississippi, 1924-1961, by Year, by Race, with Rates, Occurrence Data 1924-1944, Residence Data 1945-1961 53

Table 17. Deaths from Heart Diseases, Nephritis and Nephrosis, Vascular Lesions, Malignant Neoplasms, and Tuberculosis (all forms), Mississippi, 1952-1961, by Year and Race, with Rates, Resident Data 54

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES, by Place of Occurrence, and by Place of Residence, by Race, with Rates

County Tables 57

State Table 139

****

SYNOPSIS OF 1961 VITAL STATISTICS

INTRODUCTION

The 1961 Vital Statistics Bulletin is the nineteenth consecutive issue prepared from data obtained from records of live births, fetal deaths, deaths, marriages and divorces collected by the Division of Public Health Statistics.

Population

Population figures used in this bulletin are those of the census count of April 1, 1960, from final tabulations published by the U. S. Bureau of the Census.

Occurrence and Residence Allocation

Births, fetal deaths, and deaths were classified by place of occurrence and residence. Occurrence data were obtained from certificates filed in Mississippi at the place of the event. Residence data include transcripts of certificates filed in other states of events occurring to Mississippi residents and exclude records filed in Mississippi for nonresidents.

Some of the tables contain both residence and occurrence data; but, following recommended procedures, most of the tables and all of the rates are based on residence data only. Residence assignment for births and fetal deaths was determined on the basis of usual residence of mother, and for deaths, on the basis of usual residence of decedent. Deaths of inmates of institutions were reallocated to previous residence regardless of length of stay in the institution.

There were transcripts for 985 births to residents of Mississippi which occurred outside the State in 1961; births occurring inside Mississippi to residents of other states numbered 1,464. Fetal death transcripts totaled 21 and the nonresident certificates 81. During the year, 851 death transcripts were received and 651 nonresident deaths were recorded. The total number of states sending transcripts was 39, including the District of Columbia. Birth records came from 25 states, fetal death records from 6 states, and death records from 34 states. In Table A the states furnishing the largest numbers of records are listed.

Table A. Transcripts received, by place of occurrence

Place of Occurrence	Live Births	Deaths	Fetal Deaths
Total	985	851	21
Alabama	199	88	3
Arkansas	23	23	2
Colorado	32	5	0
Florida	6	17	0
Illinois	53	25	1
Kentucky	33	7	0
Louisiana	120	193	0
Missouri	20	10	0
Tennessee	435	391	13
Texas	8	23	0
Other	56	69	2

Marriages were classified according to residence of bride and place of issuance of marriage license. Out of state marriages of residents of Mississippi were not included in the tabulations. Divorces were classified only according to the place where the divorce was granted.

Urban-Rural Classification

Incorporated towns with population of 2,500 or more according to the 1960 census were considered as urban areas. All other areas were classified as rural.

Classification of Causes of Death

Causes of death were assigned according to the Seventh Revision (1955) of the International Statistical Classification of Diseases, Injuries, and Causes of Death. Since this revision differs only slightly from the Sixth Revision used from 1949 through 1957, rates for the years prior to 1958 have not been adjusted for comparability with the Seventh Revision. Only underlying causes were coded.

LIVE BIRTHS

Resident Births

Live births to residents of Mississippi numbered 59,523 in 1961. This total was 313 more than that for 1960, but was the third lowest figure in the period 1946-1961. There were 27,435 white births, an increase over 1960 of 237, and for nonwhites 32,088, an increase of 76. The crude birth rate of 27.3 per 1,000 population was nearly the same as that for 1960. Although this rate was the highest since 1957, it was still lower than the rates for the period 1947-1957. The State rate for whites in 1961 was 21.8 and county rates varied from 9.0 in Carroll County to 31.8 in Harrison County. For nonwhites, the State rate was 34.8 and the range for the counties was from 23.6 in Perry County to 43.1 in Smith County.

Characteristics

A count of the births by sex showed 30,505 males and 29,018 females. These totals yielded a sex ratio of 1,051 males to 1,000 females. The ratio of males to females was higher among white births than among nonwhites. Table B gives the live births by month. September was the peak month with April having the fewest.

Table B. Live births by month

Month	Births	Month	Births
January	5,192	July	5,407
February	4,698	August	5,466
March	4,897	September	5,482
April	4,208	October	5,214
May	4,375	November	4,866
June	4,616	December	5,102

Vital Statistics
Mississippi - 1961

During 1961, 22,302 children were born to urban residents and 37,221 to rural residents. As shown in Table C, the birth rate for whites was 24.3 for urban residents, compared with 20.2 for rural residents. For nonwhites the rates were 35.2 for urban and 34.7 for rural. The urban rate was higher for both racial groups, but there was a larger difference in the rates for whites. Table C also gives fertility rates per 1,000 females 15-44 years old. The rates for whites were 111.9 for urban residents and 106.5 for rural residents. These figures followed the same pattern as the crude birth rates, but for nonwhites it was reversed with the urban rate of 177.9 much lower than the rural rate of 205.7. Study of the fertility rates also shows that although there was a large difference between the rates for white and nonwhite urban residents, for rural residents the nonwhite rate was nearly double that of the whites.

Table C. Live birth rates and fertility rates for urban and rural residents, by race

Residence of Mother	Live Birth Rate*			Fertility Rate**		
	Total	White	Nonwhite	Total	White	Nonwhite
Total	27.3	21.8	34.8	143.2	108.8	196.0
Urban	28.3	24.3	35.2	134.6	111.9	177.9
Rural	26.8	20.2	34.7	148.9	106.5	205.7

* Live births per 1,000 population.
** Live births per 1,000 female population, 15-44 years of age.

The number of illegitimate births during 1961 was 8,554 or 14.4 per cent of the total. White births classified as illegitimate totaled 410 or 1.5 per cent, and 8,144 or 25.4 per cent of the nonwhite births were illegitimate. The percentage of illegitimate births among nonwhites has risen in recent years.

Among infants born alive in Mississippi in 1961 there were 5,407 with birth weight listed as 5 1/2 pounds or less and thus classified as immature. This figure was 9.0 per cent of the total. The number, 3,543, and percentage, 11.0, for nonwhites were higher than the corresponding figures of 1,864 and 6.7 per cent for the whites.

Classification by Attendant

In Table D, which gives the distribution of live births by attendant and race, it may be seen that in 1961 about 73 per cent of all births occurred in hospitals or clinics (including physicians' offices or clinics which provide inpatient care). A small proportion, 2.5 per cent, was delivered by physicians in places other than hospitals or clinics, and 24.2 per cent were delivered at home by nonmedical attendants. Almost 45 per cent of the nonwhite mothers, but less than one per cent of the white mothers, had no medical attendant.

Table D. Live births by attendant and race

Attendant	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
Total	59,523	100.0	27,435	100.0	32,088	100.0
Physician in hospital	43,589	73.2	27,219	99.2	16,370	51.0
Physician in home	1,512	2.5	96	0.3	1,416	4.4
Midwife	14,313	24.0	100	0.4	14,213	44.3
Other	109	0.2	20	0.1	89	0.3

In 1961, for the first time in many years, there was no increase in the proportion of total live births occurring in hospitals (see Table E). There was a small increment in deliveries of white infants in hospitals, but a balancing decline for the nonwhites. Even though this slight drop occurred, the percentage of 51.0 for nonwhites in 1961 is more than double that of 22.7 in 1951, ten years earlier, and over ten times the 4.2 per cent for 1941. However, there is still a large discrepancy between the current percentages for the two racial groups.

Table E. Percentage of live births in hospitals or clinics, by race, 1957-1961

Year	Total	White	Nonwhite
1957	68.3	98.0	44.8
1958	70.4	98.6	47.0
1959	72.1	98.9	49.1
1960	73.2	99.0	51.3
1961	73.2	99.2	51.0

Age of Mother

Table F gives the age distribution of mothers by race and also shows the birth rates for each group. Over half of the mothers were 15-24 years of age and more than one-third were 25-34 years old. Birth rates per 1,000 females were much higher for nonwhite mothers than those for white mothers at every age, and were more than three times as high in the groups 35 years and over. Comparison of these rates with the latest ones available (1959) for the entire United States shows that for whites the Mississippi figures are lower than those for the nation in all groups except 10-14 years (the same) and 15-19, 20-24 (higher) but that for nonwhites they are higher than the national rates for all ages except 15-19.

Table F. Live births by age of mother and race, with rates

Age of Mother	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
10-14	59,523 ~~309~~	2.6	27,435 ~~25~~	0.4	32,088 ~~284~~	4.9
15-19	11,597	121.6	4,670	91.2	6,927	156.9
20-24	19,464	284.9	10,179	252.4	9,285	331.6
25-29	13,060	214.1	6,538	170.2	6,522	288.8
30-34	8,410	130.6	3,635	88.2	4,775	205.8
35-39	4,915	76.1	1,827	43.9	3,088	134.4
40-44	1,567	25.2	522	13.3	1,045	45.9
45-49	103	1.6	30	0.8	73	3.1
50-54	3	0.05	1	0.03	2	0.1
Unknown	95		8		87	

* Live births per 1,000 female population in specific age-race group.

FETAL DEATHS

In 1952 the registration of all fetal deaths, regardless of length of gestation, was initiated in Mississippi. Prior to that time, only fetal deaths of 20 or more weeks gestation were reported. Registration of early fetal deaths is still far from complete, but there is no estimate of the degree of underregistration. There were 2,562 fetal death certificates filed during 1961 and 2,502 certificates for residents. The resident total was 118 more than the comparable figure for 1960, with an increase of 100 for white and 18 for nonwhites. For the State the fetal death ratio (fetal deaths per 1,000 live births) was 42.0. The ratio for whites was 38.4, the highest ever recorded, with county ratios varying from zero in Quitman and Sharkey Counties to 97.1 in Amite County. For nonwhites the ratio was 45.1, with variations from zero in Tishomingo County to 148.1 in Tippah County.

On 1,002 of the records the gestation period was stated to be less than 20 weeks; on 1,377 it was 20 weeks or more; and on 123 the length was not stated. In the period 1954-1961 there have been reported among the whites more fetal deaths after less than 20 weeks of gestation than after 20 weeks or more.

Using the certificates with sex stated, a sex ratio of 1,346 males to 1,000 females was determined. This ratio is the highest since 1957. As usual, the sex ratio for fetal deaths was much higher than for live births, but was not so reliable a figure because the sex was not stated on 841 or 34 per cent of the fetal death certificates.

The distribution by race and attendant is given in Table G. Approximately 58 per cent of the total were nonwhite. About 75 per cent of the reported fetal deaths were attended by physicians in hospitals and about 16 per cent were attended by midwives. Although the proportion of white live births in hospitals was larger than the percentage of white fetal deaths in hospitals, the reverse was true for the nonwhites. This situation in the nonwhite group may be accounted for by the under-reporting of fetal deaths that occur at home and the tendency to transfer difficult and premature deliveries to a hospital.

Table G. Fetal deaths by attendant and race

Attendant	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
Total	2,502	100.0	1,055	100.0	1,447	100.0
Physician in hospital	1,881	75.2	985	93.4	896	61.9
Physician in home	190	7.6	66	6.2	124	8.6
Midwife	410	16.4	2	0.2	408	28.2
Other	21	0.8	2	0.2	19	1.3

Table H shows the percentage of fetal deaths delivered in hospitals or clinics in the last five years. Percentages for the whites have shown slight variations but continued at a high level. There has been a steady increase among the nonwhites. The gap between the two race groups is a large one, but not so large as that for live births.

Table H. Percentage of fetal deaths delivered in hospitals or clinics, by race, 1957-1961

Year	Total	White	Nonwhite
1957	69.5	92.2	54.8
1958	69.0	88.8	56.8
1959	73.5	94.4	59.6
1960	74.2	95.5	59.9
1961	75.2	93.4	61.9

DEATHS

Total Deaths

In 1961 the resident death total was 21,232, a decrease of 481 from the number for 1960. The number of both white and nonwhite deaths declined slightly (white total 11,063 decreased 141 and the nonwhite total 10,169 decreased 340). The crude death rate was 9.7 per 1,000 population in comparison with a rate of 10.0 in 1960. Although the number of deaths of nonwhites was 894 less than that for whites, the death rate of the nonwhites (11.0) was greater than that of the whites (8.8).

About 37 per cent or 7,802 of the Mississippi residents who died in 1961 lived in urban locations, but 54 per cent of the deaths occurring in the state took place in urban areas. County rates for the white population ranged from 5.9 in Jackson County to 15.5 in Kemper, with a rate of 8.8 for the State; county rates for the nonwhites ranged from 5.3 in Benton to 15.4 in Warren, with a State rate of 11.0.

The resident birth rate exceeded the death rate by 17.6 per 1,000 population, or about the same amount as in the previous three years. For whites the excess was 13.0 and for nonwhites it was 23.8.

Vital Statistics
Mississippi - 1961

Age at Death

Table I shows the distribution of deaths by age and race. Comparison with the 1960 figures for total deaths shows only minor changes except in the age group under ten which decreased by 8.6 per cent. Among white persons there were relatively small differences from 1960 by age except for decreases of 10.7 per cent under 10 years and 10.2 per cent at 35-44 years and an increase of 19.6 per cent at 10-14 years. Among nonwhites, deaths of children under 10 years declined by 7.8 per cent with the other age groups having only small variations.

Comparison of the age-specific death rates for the two races reveals significant differences. The death rate for nonwhites under 55 (51.7) was double that for whites (25.8). Nonwhite rates for the next two decades, 55-74 years, were larger than those for whites, about the same at 75-84 years, and lower than the white rates at 85 years and over. Retirement age (65 years) had been reached by 61 per cent of the whites who died in 1961 but only 46 per cent of the nonwhites. The median age at death for the total population was 66.9, for the whites 69.8, and for the nonwhites 62.5.

Table I. Deaths by age and race, with rates

Age	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
All Ages	21,232	9.7	11,063	8.8	10,169	11.0
Under 1	2,250	37.8	645	23.5	1,605	50.0
1-4	357	1.6	101	1.0	256	2.2
5-9	152	0.6	52	0.4	100	0.7
10-14	143	0.6	61	0.5	82	0.7
15-24	446	1.3	211	1.1	235	1.6
25-34	513	2.2	225	1.4	288	3.6
35-44	920	3.8	413	2.6	507	6.3
45-54	1,934	8.4	962	6.6	972	11.7
55-64	3,029	17.6	1,657	15.3	1,372	21.5
65-74	4,611	37.0	2,513	33.4	2,098	42.5
75-84	4,554	84.5	2,870	84.3	1,684	84.7
85 & over	2,302	198.6	1,351	203.6	951	192.0
Unknown	21	-	2	-	19	-

* Deaths per 1,000 population in specific age-race group, except for the age group under 1 year. Deaths under 1 year per 1,000 live births.

Leading Causes of Death

In Table J are listed the ten leading causes of death in the State in 1961. These ten causes were responsible for 78 per cent of the deaths among Mississippi residents. Comparisons are made for 1949-1961 only because of the revisions in cause of death coding in 1949. Heart diseases, vascular lesions, and malignant neoplasms have been the first three leading causes of death in that order since 1949.

Accidents, certain diseases of early infancy, and pneumonia and influenza have been listed as fourth, fifth, and sixth since 1953 and were in these positions but in varying order in 1950-1952. General arteriosclerosis has ranked either seventh or eighth since 1958 and among the leading ten since 1955 and also for 1949 and 1950. Diabetes mellitus has appeared in either seventh or eighth place since 1957 and among the ten leading causes of death since 1949 except for 1955, 1952, and 1950. Congenital malformations has been among the first ten since 1953 except for 1954 and 1960. Except for 1959 and 1960, chronic and unspecified nephritis and other renal sclerosis has been among the ten leading causes of death each year since 1949.

The ten leading causes of death differed by race. For the whites malignant neoplasms was second, and vascular lesions affecting central nervous system was third, and other diseases of circulatory system replaced chronic and unspecified nephritis and other renal sclerosis in tenth place. For the nonwhite group certain diseases of early infancy was listed fourth and accidents fifth; diabetes and the nephritis group exchanged places and became tenth and eighth and homicide and gastritis, duodenitis, enteritis, and colitis, except diarrhea of newborn replaced the causes shown on the table in seventh and ninth position.

Table J. Ten leading causes of death by race, with per cent of total deaths

Cause and International List Number	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
1. Diseases of heart (400-402,410-443)	6,561	30.9	4,022	36.4	2,539	25.0
2. Vascular lesions affecting central nervous system (330-334)	2,891	13.6	1,464	13.2	1,427	14.0
3. Malignant neoplasms (140-205)	2,748	12.9	1,732	15.6	1,016	10.0
4. Accidents (800-962)	1,427	6.7	760	6.9	667	6.6
5. Certain diseases of early infancy (760-776)	1,201	5.6	416	3.8	785	7.7
6. Pneumonia and influenza (480-493)	721	3.4	307	2.8	414	4.1
7. General arteriosclerosis (450)	327	1.5	208	1.9	119	1.2
8. Diabetes mellitus (260)	291	1.4	149	1.3	142	1.4
9. Congenital malformations (750-759)	239	1.1	138	1.2	101	1.0
10. Chronic & unspecified nephritis & other renal sclerosis (592-594)	237	1.1	79	0.7	158	1.6

Diseases of the heart, the leading cause of death, was responsible for 30.9 per cent of all deaths in Mississippi in 1961. (Beginning with this year rheumatic fever is included with the diseases of the heart, following the practice of the National Vital Statistics Division.) As seen in Table K, 6,561 deaths were attributed to some form of heart disease and a large number (4,742) were specified as arteriosclerotic and degenerative heart disease. The second largest category was hypertensive heart disease. The death rate was 301.2 per 100,000 population, an increase from a rate of 290.2 for all types of heart disease in 1960. Both the white and nonwhite groups showed increases.

Table K. Deaths from diseases of the heart, by type, by race

Cause and International List Number	Total	White	Nonwhite
Diseases of the heart (400-402,410-443)	6,561	4,022	2,539
Rheumatic fever (400-402)	15	3	12
Chronic rheumatic heart disease (410-416)	123	72	51
Arteriosclerotic and degenerative heart disease (420-422)	4,742	3,314	1,428
Other diseases of heart (430-434)	554	248	306
Hypertensive heart disease (440-443)	1,127	385	742

The second leading cause of death, vascular lesions affecting the central nervous system, caused 2,891 deaths or 13.6 per cent of the total. There was a small decrease from 1960, and, as in previous years, the rate for nonwhites exceeded that for whites. The components of this category are listed in Table L showing that a majority of the deaths (1,795) were due to cerebral hemorrhage.

Table L. Deaths from vascular lesions affecting central nervous system, by type of lesion, by race

Cause and International List Number	Total	White	Nonwhite
Vascular lesions affecting central nervous system (330-334)	2,891	1,464	1,427
Subarachnoid hemorrhage (330)	94	50	44
Cerebral hemorrhage (331)	1,795	842	953
Cerebral embolism and thrombosis (332)	638	400	238
Other and ill-defined (333,334)	364	172	192

There were 2,748 deaths due to malignant neoplasms (third on the list of leading causes of death), or 113 more than in 1960. The malignant neoplasm death total for whites in 1961 was the highest on record in this State. For nonwhites this was the next to the highest year on record. The distribution of these deaths by age, sex, and race is given in Table M. White males had the largest number of deaths and the highest death rate; nonwhite females had the smallest. In comparison with 1960, each sex-race group showed an increase, the largest occurring among white males. The age group 85 and over was the one in which the greatest increase occurred and the only one in which an increase was shown for each sex-race group. Increases over 1960 in the totals for each age were noted except among those under 15 and those 35-44 years of age.

Comparison of the death rates for cancer by race shows that among males, the rates for whites were larger than the ones for nonwhites at all ages except 35-44 years, but among females the rates for whites were higher only in the group 10-14 years and 75 years and over. There were also differences by sex; among the whites, males had higher rates than females at all ages except under 10, but among nonwhites the rates for males were greater than those for females only in the 10-14 year group and those 55 and over.

Table M. Deaths from malignant neoplasms, by age, sex, and race, with rates

Age in Years	Number of Deaths					Rate*				
		Male		Female			Male		Female	
	Total	White	Non-white	White	Non-white	Total	White	Non-white	White	Non-white
All Ages	2,748#	945	515#	787	501	126.2	151.2	116.3	124.4	104.9
Under 10	36	9	7	9	11	6.7	6.8	5.0	7.1	7.9
10-14	9	4	3	2	0	3.7	6.2	5.0	3.3	-
15-24	29	14	4	6	5	8.7	14.3	5.7	6.6	6.9
25-34	54	17	7	17	13	22.9	22.3	20.3	21.4	28.4
35-44	139	39	21	39	40	58.2	50.6	60.1	48.2	87.5
45-54	381	126	56	107	92	166.1	174.2	144.8	144.3	207.8
55-64	607	219	112	172	104	352.8	420.0	371.7	307.1	308.2
65-74	785	282	161	203	139	630.4	804.6	677.4	506.4	542.4
75-84	522	183	102	168	69	968.2	1,223.8	1,053.1	880.6	676.5
85 & over	185	52	41	64	28	1,596.3	1,912.5	1,784.2	1,633.9	1,054.6

* Deaths per 100,000 population in specific age-sex-race group.
\# Includes one of unknown age.

Table N. lists deaths from malignant neoplasms by primary site. The leading site was digestive organs and peritoneum with 868 deaths, an increase of 56 from 1960. Cancer of the breast and genito-urinary system accounted for the second largest number of deaths with 827. The two leading sites were responsible for 77 per cent of the cancer deaths among nonwhite females, 69 per cent among white females, 65 per cent among nonwhite males, but only 46 per cent among white males. Cancer of the respiratory system showed an increase of 42 deaths and remained in third place. The death rate for cancer of this site among white males was much larger than the rates for the other three sex-race groups, and has increased from 5.4 in 1940 to 41.6 in 1961. The highest rate on the table was that for deaths from cancer of the breast and genito-urinary system among white females.

Vital Statistics
Mississippi - 1961

Table N. Deaths from malignant neoplasms, by primary site, sex, and race, with rates

Primary Site and International List Number	Male				Female			
	White		Nonwhite		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*	No.	Rate*
Total	945	151.2	515	116.3	787	124.4	501	104.9
Buccal cavity & pharynx (140-148)	26	4.2	10	2.2	16	2.5	5	1.0
Digestive organs & peritoneum (150-159)	284	45.4	207	46.7	224	35.4	153	32.0
Respiratory system (160-165)	260	41.6	71	16.0	46	7.3	22	4.6
Breast & genito-urinary organs (170-181)	146	23.4	129	29.1	319	50.4	233	48.8
Other & unspecified sites (190-199)	114	18.2	52	11.7	112	17.7	57	11.9
Skin (190,191)	19	3.0	4	0.9	22	3.5	5	1.0
Eye, brain & other parts of nervous (192,193)	29	4.6	6	1.4	22	3.5	7	1.5
Bone & connective tissue (196,197)	12	1.9	5	1.1	6	0.9	6	1.2
Other & unspecifies sites (194,195,198,199)	54	8.6	37	8.4	62	9.8	39	8.2
Lymphatic & hematopoietic tissues (200-205)	115	18.4	46	10.4	70	11.1	31	6.5

* Deaths per 100,000 population in the specific sex-race group.

The number of deaths in 1961 which resulted from accidents, the fourth-ranking cause, was 1,427, a drop of 77 from 1960. Most of the decline occurred among white persons. Deaths from motor vehicle accidents showed a large decrease, 48, among whites, but the total for nonwhites increased by 13. A decrease was noted in both racial groups for deaths caused by firearms and drowning. Also, there was a decline in the number of nonwhite deaths due to fire and explosion of combustible material. An increase in the number of deaths from accidents caused by machinery was recorded. Poisonings by solid and liquid substances caused more deaths than in 1960, but those by gases and vapors decreased. The largest number of deaths from all accidents occurred in December, as has been true for all but one year in the period 1950-1961. The average number of accidental deaths per day was 3.9.

Table O shows the five leading causes of accidental death in 1961 by race. These same causes have appeared on the list for a number of years.

Table O. Five leading causes of accidental deaths, by race

Type of Accident and International List Number	Total	White	Nonwhite
1. Motor vehicle (810-835)	585	380	205
2. Fire and explosion of combustible material (916)	215	61	154
3. Falls (900-904)	127	88	39
4. Drowning, except water transport (929)	98	37	61
5. Firearm (919)	67	36	31

Leading causes of death for each age group except infants are given in Table P. The leading causes of death differed in the various age groups, even though certain causes appeared more than once. Malignant neoplasms appeared in either the second or third place in each age group. Accidents were the leading cause of death in all age groups 1-44 years and was in fourth order at 45-64 years and sixth at 65 years and over. Diseases of the heart was among the five leading causes of death in all age groups 5 years and over and was first in those 45 years and over. Pneumonia and influenza appeared on the lists for the two youngest groups and also the oldest group. All groups 25 years and over had vascular lesions affecting the central nervous system on their list. A more detailed age breakdown of deaths among persons who have reached 65 years or above shows that for those 65-74 years old the five leading causes of death are the same as for 55-64 years, but for persons 75-84 and 85 and over, the leading causes of death are as listed for 65 years or above.

Ill-defined Causes

Of the 21,032 certificates filed for deaths that occurred in Mississippi in 1961, those with no satisfactory statement of cause numbered 2,407, or 11.4 per cent of the total. For white persons the percentage of certificates in this category was 6.0 and for nonwhites, 17.4. Senility was the only cause given on 74 of these certificates; ill-defined diseases of early infancy and immaturity unqualified appeared on 659; and other ill-defined and unknown causes on 1,674.

Deaths with unsatisfactory statement of cause were so numerous that if placed on the list of ten leading causes they would rank fourth and would appear as one of the five leading causes of death in every age group. If the true causes of death for these persons could be determined and properly assigned, the order of leading causes might not be the same and even the causes listed might be different.

If deaths from ill-defined diseases of early infancy and immaturity unqualified are excluded from the group of deaths without definite cause, the remaining ill-defined deaths comprised 8.3 per cent of all deaths recorded in Mississippi in 1961. This group constitutes 15 per cent or more of the total deaths in Amite, Carroll, Hancock, Holmes, Issaquena, Kemper, Madison, and Washington Counties.

Vital Statistics
Mississippi - 1961

Table P. Five leading causes of death by age group

Cause of Death	Age Group	No. of Deaths	% of Total
	1-4 years		
All Causes		357	100.0
1. Accidents		126	35.3
2. Pneumonia and influenza		46	12.9
3. Malignant neoplasms		21	5.9
4. Gastritis, duodenitis, enteritis, and colitis		17	4.8
4. Congenital malformations		17	4.8
	5-14 years		
All Causes		295	100.0
1. Accidents		146	49.5
2. Malignant neoplasms		19	6.4
3. Pneumonia and influenza		16	5.4
3. Congenital malformations		16	5.4
5. Diseases of heart		10	3.4
	15-24 years		
All Causes		446	100.0
1. Accidents		216	48.4
2. Homicide		52	11.6
3. Malignant neoplasms		29	6.5
4. Diseases of heart		16	3.6
4. Deliveries & complication of pregnancy, childbirth, & the puerperium		16	3.6
	25-44 years		
All Causes		1,433	100.0
1. Accidents		291	20.3
2. Diseases of heart		247	17.2
3. Malignant neoplasms		193	13.5
4. Vascular lesions affecting central nervous system		112	7.8
5. Homicide		88	6.1
	45-64 years		
All Causes		4,963	100.0
1. Diseases of heart		1,667	33.6
2. Malignant neoplasms		988	19.9
3. Vascular lesions affecting central nervous system		591	11.9
4. Accidents		282	5.7
5. Diabetes mellitus		86	1.7
	65 years and over		
All Causes		11,467	100.0
1. Diseases of heart		4,601	40.1
2. Vascular lesions affecting central nervous system		2,174	19.0
3. Malignant neoplasms		1,492	13.0
4. Pneumonia and influenza		347	3.0
5. General arteriosclerosis		301	2.6

Infant Mortality

In 1961, an infant mortality rate of 37.8 per 1,000 live births was recorded, a reduction of 3.7 from the 1960 rate. This is the lowest infant mortality rate since 1957. Distribution of infant deaths by age and race is shown in Table Q. In each age group, the rate for nonwhites exceeded that for whites, with the greatest difference between the two racial groups occurring in the period 28 days to one year.

Table Q. Infant deaths by age and race, with rates

Age	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
Total under 1 year	2,250	37.8	645	23.5	1,605	50.0
Under one day	667	11.2	282	10.3	385	12.0
One to 28 days	734	12.3	227	8.3	507	15.8
28 days to one year	849	14.3	136	5.0	713	22.2

* Deaths per 1,000 live births.

Table R gives the five leading causes of infant deaths. The causes were the same and in the same order as in 1960. Influenza and pneumonia, except pneumonia of newborn, was in first place. Although there was a decline from 1960 in the total number of infant deaths, there was an increase in the numbers dying from congenital malformations and from birth injuries. For the two racial groups, both the leading causes and their rank differed. The list for whites included accidents (28 deaths) and excluded the gastritis group; that for nonwhites included accidents (72 deaths), but not birth injuries. Immaturity, which is no longer considered a separate classification in ranking causes of infant deaths, was mentioned either in combination with another cause or as the only cause on 44 per cent of the certificates for white infants and 34 per cent of those for nonwhites.

Table R. Leading causes of infant deaths, by race

Cause of Death	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
All Causes	2,250	100.0	645	100.0	1,605	100.0
1. Influenza & pneumonia, except pneumonia of newborn	209	9.3	28	4.3	181	11.3
2. Postnatal asphyxia & atelectasis	191	8.5	113	17.5	78	4.8
3. Congenital malformations	184	8.2	107	16.6	77	4.8
4. Gastritis, duodenitis, enteritis & colitis, except diarrhea of newborn	134	6.0	9	1.4	125	7.8
5. Birth injuries	116	5.2	53	8.2	63	3.9

Infant death certificates which showed ill-defined causes or no cause at all numbered 950 or 42 per cent of the total in 1961. For 34 per cent of the white and 45 per cent of the nonwhite infants no definite cause of death was given. Determination of the true cause of death for this large group of infants might change the leading causes of death which are listed above.

Infant mortality rates by race for 1920-1961 are shown in Graph 1. In general, the trend has been downward for both races. In 1961 the curve for the nonwhites again turned down after swinging upward for several years.



Maternal Mortality

The maternal mortality rate for 1961 was 8.2 per 10,000 live births in the total population, 3.3 for white mothers, and 12.5 for nonwhites. The total of 49 maternal deaths was the lowest on record and ten less than in 1960. There was an increase of one death among the white mothers and a decrease of eleven among the nonwhite. The discrepancy between the rates by race has narrowed considerably in recent years.

In Table S maternal deaths are shown by cause. In 1961, as in each year from 1949 through 1960 (years with comparable data), the two leading causes were toxemia and hemorrhage. Toxemia has been the leading cause in each year of this period except 1957.

Table S. Maternal deaths by cause and race, with rates

Cause of Death	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
All Causes	49	8.2	9	3.3	40	12.5
Toxemia	18	3.0	2	0.7	16	5.0
Hemorrhage	14	2.4	1	0.4	13	4.0
Sepsis	6	1.0	2	0.7	4	1.2
Abortion without sepsis or toxemia	3	0.5	1	0.4	2	0.6
Abortion with sepsis	2	0.3	1	0.4	1	0.3
Other complications	6	1.0	2	0.7	4	1.2

* Deaths per 10,000 live births.

Table T gives the age and race-specific maternal death rates. There were three deaths in 10-14 year age group and one in the 45-54 year group. For all age groups except the youngest, in which the number of births is very small, the rates for nonwhite mothers were higher than those for white mothers. There was a decline from 1960 in the number of deaths occurring in the period 35-44 years of age.

Table T. Maternal deaths by age and race, with rates

Age	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
All Ages	49	8.2	9	3.3	40	12.5
10-14	3	97.1	1	#	2	70.4
15-24	16	5.2	4	2.7	12	7.4
25-34	17	7.9	4	3.9	13	11.5
35-44	12	18.5	0	-	12	29.0
45-54	1	94.3	0	-	1	#

* Deaths per 10,000 live births to mothers in specific age-race group.
\# Rate not shown for less than 100 live births.

Maternal death rates by race for 1913-1961 are shown in Graph 2. A downward trend began in the 1930's and has been rapid since 1940, though in 1959 and 1960 there was an increase in the death rate among nonwhite mothers.

Graph 2. Maternal Mortality
1913-1961



MARRIAGES

In 1961, the number of marriages occurring in Mississippi was 21,610 or 377 more than in 1960. The total for whites was 14,438 and for nonwhites was 7,172. Table U shows the number of marriages by month. The largest number occurred in June, but the total for December was nearly as large. The average daily number of marriages was 59.

Table U. Number of marriages and averages per day by month

Month	Number	Average per Day	Month	Number	Average per Day
Total	21,610	59			
January	1,515	49	July	2,003	65
February	1,377	49	August	2,020	65
March	1,447	47	September	1,861	62
April	1,613	54	October	1,646	53
May	1,565	50	November	1,625	54
June	2,484	83	December	2,454	79

Table V shows the age distribution of brides and grooms. Teen-aged brides numbered 9,768 or 45 per cent of the total. This figure included one bride age 11, three age 12, 18 who were 13 years old, and 104 who were only 14. A much smaller percentage of grooms were in the teen-age group - 17.3 per cent. There were 26 brides and 145 grooms 75 years of age or over. The oldest bride was 90 and two of the grooms were 99 years old. Most marriages were between persons of approximately the same age, but for some there was a wide range of differences in age of bride and groom. The greatest difference was in the marriage of a 15 year old girl and a man 86 years old.

Table V. Brides and grooms by age

Age	Bride		Groom	
	No.	%	No.	%
Total	21,610	100.0	21,610	100.0
Under 15	126	0.6	1	*
15	865	4.0	6	*
16	1,373	6.3	66	0.3
17	2,047	9.5	552	2.6
18	3,017	14.0	1,330	6.2
19	2,340	10.8	1,785	8.3
20-24	5,653	26.2	8,874	41.1
25-34	2,863	13.2	4,535	21.0
35-44	1,538	7.1	1,850	8.6
45-54	1,037	4.8	1,108	5.1
55 and over	719	3.3	1,481	6.8
Unknown	32	0.1	22	0.1

* Less than 0.06 per cent.

Ministers officiated at 71 per cent of the marriages in 1961. The number of previous marriages of brides and grooms is given in Table W. This was the first marriage for 72.0 per cent of the brides and 71.8 per cent of the grooms. It was the first for both bride and groom in 64.1 per cent of the marriages. Among those who had

been married before there was a much larger number of divorced than of widowed persons. Brides with more than one previous marriage numbered 1,376 or 6.4 per cent of the total married in the State, and 12 had 5 or more previous marriages. Corresponding figures for grooms were 1,452 or 6.7 per cent with multiple previous marriages and 14 with 5 or more prior to the latest.

Table W. Number of previous marriages of brides and grooms

Previous Marriages	Bride		Groom	
	No.	%	No.	%
Total	21,610	100.0	21,610	100.0
None	15,555	72.0	15,510	71.8
One	4,611	21.3	4,596	21.3
Two	1,124	5.2	1,148	5.3
Three or more	252	1.2	304	1.4
Not stated	68	0.3	52	0.2

The number of nonresident brides married in Mississippi in 1961 was 3,877 or 18 per cent of the total, and there were 3,355 nonresident couples. A comparison of the resident status of brides is shown in Table X for 1946-1961. In each year, 1948 through 1956, the number of out-of-state brides increased. The increase in nonresidents ended abruptly when a new and much stricter marriage law became effective in July, 1958, and there has been a decrease in the nonresident total each year that this law has been in effect. The number of resident brides declined each year in the period 1947-1959, with the exception of 1955, but an increase was noted in both 1960 and 1961.

Table X. Marriages by residence status of bride, 1946-1961

Year	Total	Resident		Nonresident		Not Stated
		No.	%	No.	%	
1946	64,204	41,882	65.2	22,058	34.4	264
1947	53,584	34,030	63.5	19,517	36.4	37
1948	53,645	30,921	57.6	22,724	42.4	*
1949	52,765	27,717	52.5	25,048	47.5	*
1950	56,738	26,859	47.3	29,879	52.7	*
1951	57,321	25,713	44.8	31,608	55.1	*
1952	58,116	24,868	42.8	33,248	57.2	*
1953	61,751	24,497	39.7	37,254	60.3	*
1954	61,381	23,555	38.4	37,826	61.6	*
1955	66,423	23,683	35.6	42,740	64.3	*
1956	66,223	23,011	34.7	43,212	65.2	*
1957	62,495	20,864	33.4	41,510	66.4	121
1958	36,402	17,370	47.7	18,920	52.0	112
1959	20,447	16,288	79.6	4,062	19.9	97
1960	21,233	17,198	81.0	3,979	18.7	56
1961	21,610	17,727	82.0	3,877	17.9	6

* Not classified.

Nonresident brides came to Mississippi from the District of Columbia and all but four states. There were 14 brides from foreign countries. Three-fourths of these out-of-state brides came from the bordering states of Alabama, Arkansas, Louisiana, and Tennessee and counties which bordered these states issued 80 per cent of the licenses to out-of-state brides. In DeSoto County which is adjacent to Memphis, Tennessee 83 per cent of the licenses issued were for nonresident brides.

In each year since July 1, 1956, when a new marriage record containing more detailed information was put into use in this state, the records have shown that some Mississippi licenses have been used for marriage ceremonies performed in other states. However, because these marriages did not occur in Mississippi and are of questionable legality, they have not been included in the annual tabulations. In 1961, 120 Mississippi licenses were known to have been used in other states. All but 8 of these licenses were issued in border counties and over half by either Adams or Wilkinson Counties. A majority (84) of the marriages took place in Louisiana; 30 took place in the other border states of Tennessee, Arkansas, and Alabama; and 6 occurred in other states. One or both of the parties was a resident of the state in which the marriage took place in 98 cases, but in 3 instances neither bride nor groom was a resident of Mississippi nor of the state where the ceremony was performed.

DIVORCES

Divorces granted in Mississippi during 1961 numbered 5,213 - 47 less than in 1960, but the second highest total since 1952. There were 3,545 granted to white couples and 1,668 to nonwhite couples. Cruel and inhuman treatment and desertion continued to rank as the two leading causes and together accounted for 88 per cent of the total divorces granted. Cruel and inhuman treatment was given as the main cause for 62 per cent of the divorces among whites; among nonwhites 66 per cent were granted on grounds of desertion. The wife was listed as complainant on 63 per cent of the records.

No minor children were involved in 47 per cent of the divorces, and 20 per cent of the couples had only one minor child. Ten of the families had 9 or more minor children. The total number of minor children affected was 5,977.

Thirty-five per cent of the divorces in 1961 were granted to persons married less than 5 years; 5 per cent had been married less than one year; but 17 per cent had been married 20 years or longer. More divorces were granted to couples in their second year of marriage than in any other year. The distribution of divorces according to duration of marriage was different for the two races. Of the white couples, 43 per cent were granted after less than 5 years of marriage and only 10 per cent after 20 years or longer; for the nonwhites, corresponding percentages were 19 and 32 respectively.

TABLE I
CENSUS POPULATION, MISSISSIPPI
APRIL 1, 1960

County	Total	White	Nonwhite
Total	2,178,141	1,257,546	920,595
Adams	37,730	19,035	18,695
Alcorn	25,282	21,949	3,333
Amite	15,573	7,130	8,443
Attala	21,335	11,789	9,546
Benton	7,723	4,114	3,609
Bolivar	54,464	17,521	36,943
Calhoun	15,941	11,595	4,346
Carroll	11,177	4,677	6,500
Chickasaw	16,891	10,380	6,511
Choctaw	8,423	5,903	2,520
Claiborne	10,845	2,600	8,245
Clarke	16,493	10,001	6,492
Clay	18,933	9,214	9,719
Coahoma	46,212	14,630	31,582
Copiah	27,051	12,992	14,059
Covington	13,637	8,896	4,741
DeSoto	23,891	9,248	14,643
Forrest	52,722	37,970	14,752
Franklin	9,286	5,486	3,800
George	11,098	9,811	1,287
Greene	8,366	6,443	1,923
Grenada	18,409	9,352	9,057
Hancock	14,039	11,784	2,255
Harrison	119,489	100,233	19,256
Hinds	187,045	112,205	74,840
Holmes	27,096	7,595	19,501
Humphreys	19,093	5,758	13,335
Issaquena	3,576	1,176	2,400
Itawamba	15,080	14,206	874
Jackson	55,522	44,658	10,864
Jasper	16,909	8,402	8,507
Jefferson	10,142	2,489	7,653
Jeff Davis	13,540	6,126	7,414
Jones	59,542	44,095	15,447
Kemper	12,277	4,828	7,449
Lafayette	21,355	14,110	7,245

Table 1 - Page 2
Census Population - Mississippi - April 1, 1960

County	Total	White	Nonwhite
Lamar	13,675	11,443	2,232
Lauderdale	67,119	43,635	23,484
Lawrence	10,215	6,354	3,861
Leake	18,660	10,559	8,101
Lee	40,589	30,300	10,289
Leflore	47,142	16,699	30,443
Lincoln	26,759	18,407	8,352
Lowndes	46,639	28,871	17,768
Madison	32,904	9,267	23,637
Marion	23,293	15,408	7,885
Marshall	24,503	7,264	17,239
Monroe	33,953	21,932	12,021
Montgomery	13,320	7,349	5,971
Neshoba	20,927	15,026	5,901
Newton	19,517	12,950	6,567
Noxubee	16,826	4,724	12,102
Oktibbeha	26,175	14,727	11,448
Panola	28,791	12,565	16,226
Pearl River	22,411	17,221	5,190
Perry	8,745	6,333	2,412
Pike	35,063	19,655	15,408
Pontotoc	17,232	13,946	3,286
Prentiss	17,949	15,763	2,186
Quitman	21,019	7,715	13,304
Rankin	34,322	21,504	12,818
Scott	21,187	13,050	8,137
Sharkey	10,738	3,247	7,491
Simpson	20,454	13,254	7,200
Smith	14,303	11,056	3,247
Stone	7,013	5,302	1,711
Sunflower	45,750	14,730	31,020
Tallahatchie	24,081	8,580	15,501
Tate	18,138	7,696	10,442
Tippah	15,093	12,337	2,756
Tishomingo	13,889	13,210	679
Tunica	16,826	3,505	13,321
Union	18,904	15,592	3,312
Walthall	13,512	7,412	6,100
Warren	42,206	22,447	19,759
Washington	78,638	35,239	43,399
Wayne	16,258	10,449	5,809
Webster	10,580	7,938	2,642
Wilkinson	13,235	3,807	9,428
Winston	19,246	10,853	8,393
Yalobusha	12,502	6,962	5,540
Yazoo	31,653	12,862	18,791

~~Table 1 - Page 3~~
~~Census Population - Mississippi - April 1, 1960~~
Census Population for Selected Mississippi Cities
April 1, 1960

City*	Total	White	Nonwhite
Biloxi	44,053	38,348	5,705
Clarksdale	21,105	9,818	11,287
Cleveland	10,172	6,172	4,000
Columbus	24,771	14,822	9,949
Corinth	11,453	9,136	2,317
Greenville	41,502	21,082	20,420
Greenwood	20,436	9,818	10,618
Gulfport	30,204	23,839	6,365
Hattiesburg	34,989	23,746	11,243
Jackson	144,422	92,793	51,629
Laurel	27,889	17,914	9,975
McComb	12,020	8,605	3,415
Meridian	49,374	32,596	16,778
Natchez	23,791	11,437	12,354
Pascagoula	17,155	13,269	3,886
Tupelo	17,221	13,073	4,148
Vicksburg	29,143	15,516	13,627
Yazoo City	11,236	5,046	6,190
Total Urban**	787,731	500,469	287,262
Total Rural	1,390,410	757,077	633,333

* Cities having 10,000 population and over according to 1960 census.

** Includes all incorporated towns of 2,500 population or more according to 1960 census.

TABLE 2
LIVE BIRTHS
MISSISSIPPI
1961

By Place of Occurrence, and by Place of Residence of Mother, by Race with Rates

County	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
STATE	60,002	27,797	32,205	59,523	27,435	32,088	27.3	21.8	34.8
Adams	1,588	649	939	1,062	443	619	28.1	23.3	33.1
Alcorn	534	429	105	504	400	104	19.9	18.2	31.2
Amite	178	10	168	340	103	237	21.8	14.4	28.1
Attala	482	190	292	482	183	299	22.6	15.5	31.3
Benton	123	2	121	221	68	153	28.6	16.5	42.4
Bolivar	1,994	348	1,646	1,892	373	1,519	34.7	21.3	41.1
Calhoun	371	171	200	377	200	177	23.6	17.2	40.7
Carroll	188	23	165	220	42	178	19.7	9.0	27.4
Chickasaw	412	226	186	439	225	214	26.0	21.7	32.9
Choctaw	173	98	75	188	105	83	22.3	17.8	32.9
Claiborne	203	24	179	275	40	235	25.4	15.4	28.5
Clarke	152	87	65	328	144	184	19.9	14.4	28.3
Clay	483	187	296	511	208	303	27.0	22.6	31.2
Coahoma	1,565	390	1,175	1,584	372	1,212	34.3	25.4	38.4
Copiah	859	307	552	713	266	447	26.4	20.5	31.8
Covington	192	141	51	330	170	160	24.2	19.1	33.7
DeSoto	687	74	613	748	180	568	31.3	19.5	38.8
Forrest	1,603	1,171	432	1,388	879	509	26.3	23.1	34.5
Franklin	143	67	76	214	93	121	23.0	17.0	31.8
George	314	264	50	317	266	51	28.6	27.1	39.6
Greene	146	96	50	198	133	65	23.7	20.6	33.8
Grenada	540	204	336	542	207	335	29.4	22.1	37.0
Hancock	203	137	66	321	249	72	22.9	21.1	31.9
Harrison	4,221	3,462	759	3,919	3,191	728	32.8	31.8	37.8
Hinds	6,195	3,168	3,027	5,491	2,670	2,821	29.4	23.8	37.7
Holmes	777	132	645	754	115	639	27.8	15.1	32.8
Humphreys	599	86	513	612	104	508	32.0	18.1	38.1
Issaquena	34	0	34	101	22	79	28.2	18.7	32.9
Itawamba	75	56	19	241	216	25	16.0	15.2	28.6
Jackson	1,341	937	404	1,652	1,223	429	29.8	27.4	39.5
Jasper	85	17	68	442	142	300	26.1	16.9	35.3
Jefferson	103	0	103	264	33	231	26.0	13.2	30.2
Jeff Davis	255	122	133	305	118	187	22.5	19.3	25.2
Jones	2,593	1,266	1,327	1,444	926	518	24.2	21.0	33.5
Kemper	198	34	164	274	61	213	22.3	12.6	28.6
Lafayette	521	286	235	553	302	251	25.9	21.4	34.6

Table 2 - Page 2
Live Births - Mississippi - 1961

County	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Lamar	95	72	23	311	237	74	22.7	20.7	33.2
Lauderdale	2,130	1,064	1,066	1,721	888	833	25.6	20.4	35.5
Lawrence	160	53	107	228	120	108	22.3	18.9	28.0
Leake	314	136	178	396	169	227	21.2	16.0	28.0
Lee	1,304	948	356	1,055	718	337	26.0	23.7	32.8
Leflore	1,570	452	1,118	1,533	403	1,130	32.5	24.1	37.1
Lincoln	608	368	240	625	352	273	23.4	19.1	32.7
Lowndes	1,492	860	632	1,411	773	638	30.2	26.8	35.9
Madison	966	155	811	1,010	180	830	30.7	19.4	35.1
Marion	483	275	208	564	336	228	24.2	21.8	28.9
Marshall	730	126	604	746	145	601	30.4	20.0	34.9
Monroe	748	379	369	786	422	364	23.1	19.2	30.3
Montgomery	359	146	213	349	132	217	26.2	18.0	36.3
Neshoba	434	238	196	439	250	189	21.0	16.6	32.0
Newton	404	242	162	434	221	213	22.2	17.1	32.4
Noxubee	516	110	406	501	106	395	29.8	22.4	32.6
Oktibbeha	787	340	447	745	310	435	28.5	21.0	38.0
Panola	821	227	594	848	245	603	29.4	19.5	37.2
Pearl River	589	447	142	514	343	171	22.9	19.9	32.9
Perry	45	39	6	211	154	57	24.1	24.3	23.6
Pike	959	457	502	857	369	488	24.4	18.8	31.7
Pontotoc	329	245	84	337	251	86	19.6	18.0	26.2
Prentiss	463	381	82	407	334	73	22.7	21.2	33.4
Quitman	779	212	567	697	170	527	33.2	22.0	39.6
Rankin	239	85	154	858	517	341	25.0	24.0	26.6
Scott	558	335	223	534	289	245	25.2	22.1	30.1
Sharkey	233	29	204	356	52	304	33.2	16.0	40.6
Simpson	440	231	209	503	253	250	24.6	19.1	34.7
Smith	75	42	33	330	190	140	23.1	17.2	43.1
Stone	149	107	42	160	105	55	22.8	19.8	32.1
Sunflower	1,287	241	1,046	1,398	271	1,127	30.6	18.4	36.3
Tallahatchie	735	125	610	782	159	623	32.5	18.5	40.2
Tate	426	23	403	590	147	443	32.5	19.1	42.4
Tippah	360	269	91	318	237	81	21.1	19.2	29.4
Tishomingo	179	160	19	251	229	22	18.1	17.3	32.4
Tunica	648	64	584	623	69	554	37.0	19.7	41.6
Union	397	294	103	388	285	103	20.5	18.3	31.1
Walthall	469	267	202	303	128	175	22.4	17.3	28.7
Warren	1,452	620	832	1,091	513	578	25.8	22.8	29.2
Washington	2,568	948	1,620	2,460	882	1,578	31.3	25.0	36.4
Wayne	438	313	125	433	259	174	26.6	24.8	30.0
Webster	168	90	78	204	123	81	19.3	15.5	30.6
Wilkinson	430	194	236	371	78	293	28.0	20.5	31.1
Winston	404	173	231	420	174	246	21.8	16.0	29.3
Yalobusha	288	117	171	279	112	167	22.3	16.1	30.1
Yazoo	844	237	607	900	263	637	28.4	20.4	33.9
Out-of-State	985	839	146	1,464	1,201	263	-	-	-

Table 2 - Page 3
Live Birth - Mississippi - 1961

City**	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Biloxi	2,691	2,415	276	1,651	1,418	233	37.5	37.0	40.8
Clarksdale	952	387	565	629	259	370	29.8	26.4	32.8
Cleveland	525	222	303	301	118	183	29.6	19.1	45.8
Columbus	895	509	386	751	390	361	30.3	26.3	36.3
Corinth	506	422	84	214	153	61	18.7	16.7	26.3
Greenville	1,449	727	722	1,277	554	723	30.8	26.3	35.4
Greenwood	846	451	395	631	281	350	30.9	28.6	33.0
Gulfport	1,376	1,030	346	809	555	254	26.8	23.3	39.9
Hattiesburg	1,562	1,170	392	891	521	370	25.5	21.9	32.9
Jackson	5,841	3,166	2,675	4,217	2,242	1,975	29.2	24.2	38.2
Laurel	2,556	1,248	1,308	735	377	358	26.4	21.0	35.9
McComb	641	376	265	294	161	133	24.4	18.7	38.9
Meridian	2,008	1,052	956	1,228	663	565	24.9	20.3	33.7
Natchez	1,585	649	936	687	282	405	28.9	24.6	32.8
Pascagoula	93	5	88	541	377	164	31.5	28.4	42.2
Tupelo	1,043	807	236	376	264	112	21.8	20.2	27.0
Vicksburg	1,420	620	800	763	354	409	26.2	22.8	30.0
Yazoo City	457	234	223	305	110	195	27.1	21.8	31.5
Total Urban#	37,998	21,863	16,135	22,302	12,177	10,125	28.3	24.3	35.2
Total Rural	22,004	5,934	16,070	37,221	15,258	21,963	26.8	20.2	34.7

* ~~Rate~~ per 1,000 population.

** Cities having 10,000 population and over according to 1960 census.

Includes all incorporated towns of 2,500 population or more according to 1960 census.

TABLE 3
LIVE BIRTHS
MISSISSIPPI
1961

By Place of Residence of Mother, by Race and Attendant, and by Sex

County	TOTAL	WHITE				NONWHITE				SEX	
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female
STATE	59,523	27,435	27,219	96	120	32,088	16,370	1,416	14,302	30,505	29,018
Adams	1,062	443	443	0	0	619	612	2	5	542	520
Alcorn	504	400	392	6	2	104	62	15	27	246	258
Amite	340	103	102	0	1	237	78	0	159	176	164
Attala	482	183	181	1	1	299	83	21	195	261	221
Benton	221	68	66	1	1	153	32	0	121	117	104
Bolivar	1,892	373	368	2	3	1,519	691	71	757	952	940
Calhoun	377	200	197	2	1	177	107	10	60	185	192
Carroll	220	42	39	1	2	178	43	1	134	119	101
Chickasaw	439	225	223	1	1	214	102	22	90	225	214
Choctaw	188	105	101	3	1	83	37	2	44	98	90
Claiborne	275	40	39	0	1	235	70	2	163	124	151
Clarke	328	144	141	2	1	184	140	6	38	157	171
Clay	511	208	208	0	0	303	181	18	104	261	250
Coahoma	1,584	372	369	1	2	1,212	511	33	668	833	751
Copiah	713	266	265	0	1	447	384	2	61	371	342
Covington	330	170	166	2	2	160	132	2	26	158	172
DeSoto	748	180	180	0	0	568	277	9	282	399	349
Forrest	1,388	879	878	1	0	509	373	6	130	693	695
Franklin	214	93	93	0	0	121	101	2	18	100	114
George	317	266	263	3	0	51	31	2	18	158	159
Greene	198	133	132	1	0	65	39	2	24	99	99
Grenada	542	207	205	0	2	335	49	78	208	292	250
Hancock	321	249	248	0	1	72	37	0	35	173	148
Harrison	3,919	3,191	3,165	2	24	728	399	54	275	2,015	1,904

Table 3 - Page 2
Live Births - Mississippi - 1961

County	TOTAL	WHITE				NONWHITE				SEX	
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female
Hinds	5,491	2,670	2,667	3	0	2,821	2,277	70	474	2,775	2,716
Holmes	754	115	111	3	1	639	98	32	509	351	403
Humphreys	612	104	103	1	0	508	88	2	418	331	281
Issaquena	101	22	22	0	0	79	48	0	31	52	49
Itawamba	241	216	212	4	0	25	20	3	2	131	110
Jackson	1,652	1,223	1,219	0	4	429	207	70	152	855	797
Jasper	442	142	142	0	0	300	241	2	57	214	228
Jefferson	264	33	33	0	0	231	128	1	102	140	124
Jeff Davis	305	118	117	1	0	187	102	2	83	143	162
Jones	1,444	926	924	1	1	518	488	2	28	728	716
Kemper	274	61	60	0	1	213	87	1	125	130	144
Lafayette	553	302	299	0	3	251	48	11	192	290	263
Lamar	311	237	237	0	0	74	66	0	8	152	159
Lauderdale	1,721	888	888	0	0	833	604	41	188	892	829
Lawrence	228	120	116	0	4	108	45	1	62	104	124
Leake	396	169	169	0	0	227	119	2	106	217	179
Lee	1,055	718	715	3	0	337	219	14	104	571	484
Leflore	1,533	403	400	0	3	1,130	134	50	946	802	731
Lincoln	625	352	352	0	0	273	230	0	43	328	297
Lowndes	1,411	773	766	3	4	638	220	20	398	735	676
Madison	1,010	180	180	0	0	830	237	10	583	508	502
Marion	564	336	326	5	5	228	91	22	115	304	260
Marshall	746	145	145	0	0	601	149	72	380	385	361
Monroe	786	422	417	3	2	364	162	28	174	425	361
Montgomery	349	132	131	0	1	217	74	7	136	171	178
Neshoba	439	250	246	0	4	189	134	2	53	220	219
Newton	434	221	220	1	0	213	153	42	18	216	218
Noxubee	501	106	105	1	0	395	96	4	295	245	256
Oktibbeha	745	310	308	2	0	435	242	1	192	370	375
Panola	848	245	242	2	1	603	317	73	213	437	411

Table 3 - Page 3
Live Births - Mississippi - 1961

County	TOTAL	WHITE				NONWHITE				SEX	
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female
Pearl River	514	343	342	0	1	171	156	3	12	263	251
Perry	211	154	153	1	0	57	50	2	5	106	105
Pike	857	369	369	0	0	488	304	4	180	442	415
Pontotoc	337	251	249	1	1	86	38	5	43	172	165
Prentiss	407	334	333	1	0	73	55	3	15	213	194
Quitman	697	170	165	3	2	527	317	18	192	359	338
Rankin	858	517	515	2	0	341	284	4	53	436	422
Scott	534	289	286	1	2	245	123	2	120	250	284
Sharkey	356	52	52	0	0	304	164	2	138	179	177
Simpson	503	253	253	0	0	250	186	0	64	251	252
Smith	330	190	188	2	0	140	107	11	22	177	153
Stone	160	105	105	0	0	55	27	0	28	84	76
Sunflower	1,398	271	263	1	7	1,127	373	154	600	733	665
Tallahatchie	782	159	155	1	3	623	98	35	490	416	366
Tate	590	147	145	2	0	443	72	36	335	302	288
Tippah	318	237	233	4	0	81	41	22	18	174	144
Tishomingo	251	229	226	3	0	22	22	0	0	124	127
Tunica	623	69	68	1	0	554	165	19	370	317	306
Union	388	285	274	1	10	103	31	6	66	203	185
Walthall	303	128	121	0	7	175	98	0	77	160	143
Warren	1,091	513	513	0	0	578	437	6	135	567	524
Washington	2,460	882	878	2	2	1,578	1,003	48	527	1,259	1,201
Wayne	433	259	257	0	2	174	126	4	44	216	217
Webster	204	123	117	6	0	81	32	16	33	103	101
Wilkinson	371	78	78	0	0	293	135	1	157	186	185
Winston	420	174	174	0	0	246	70	1	175	211	209
Yalobusha	279	112	112	0	0	167	66	66	35	145	134
Yazoo	900	263	259	2	2	637	95	3	539	481	419

Table 3 - Page 4
Live Births - Mississippi - 1961

City*	TOTAL	WHITE				NONWHITE				SEX	
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female
Biloxi	1,651	1,418	1,411	1	6	233	170	10	53	831	820
Clarksdale	629	259	259	0	0	370	275	4	91	330	299
Cleveland	301	118	118	0	0	183	126	13	44	153	148
Columbus	751	390	387	2	1	361	115	15	231	393	358
Corinth	214	153	153	0	0	61	41	8	12	99	115
Greenville	1,277	554	551	1	2	723	486	34	203	651	626
Greenwood	631	281	279	0	2	350	70	8	272	338	293
Gulfport	809	555	552	0	3	254	136	31	87	409	400
Hattiesburg	891	521	521	0	0	370	279	5	86	435	456
Jackson	4,217	2,242	2,240	2	0	1,975	1,732	60	183	2,129	2,088
Laurel	735	377	377	0	0	358	347	0	11	381	354
McComb.	294	161	161	0	0	133	116	2	15	165	129
Meridian	1,228	663	663	0	0	565	448	37	80	643	585
Natchez	687	282	282	0	0	405	401	2	2	333	354
Pascagoula	541	377	377	0	0	164	77	19	68	283	258
Tupelo	376	264	263	1	0	112	78	1	33	208	168
Vicksburg	763	354	354	0	0	409	314	6	89	386	377
Yazoo City	305	110	110	0	0	195	32	1	162	169	136

* Cities having 10,000 population and over according to 1960 census.

TABLE 4
FETAL DEATHS
MISSISSIPPI
1961

By Place of Occurrence, by Place of Residence of Mother, by Race and Attendant, and by Sex

County	Total by Occurrence	Total by Residence	WHITE				NONWHITE				SEX		
			Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female	Unde-termined
STATE	2,562	2,502	1,055	985	66	4	1,447	896	124	427	953	708	841
Adams	77	42	21	20	1	0	21	20	0	1	18	13	11
Alcorn	10	10	7	7	0	0	3	1	0	2	3	5	2
Amite	15	34	10	7	3	0	24	13	6	5	11	4	19
Attala	20	23	10	10	0	0	13	6	1	6	7	5	11
Benton	2	9	5	5	0	0	4	2	1	1	5	1	3
Bolivar	63	65	9	9	0	0	56	29	7	20	36	22	7
Calhoun	18	11	4	4	0	0	7	7	0	0	1	4	6
Carroll	8	13	1	1	0	0	12	4	2	6	7	5	1
Chickasaw	15	22	7	7	0	0	15	7	2	6	14	4	4
Choctaw	4	4	3	2	1	0	1	1	0	0	2	2	0
Claiborne	4	12	2	1	1	0	10	7	2	1	5	4	3
Clarke	1	8	1	0	1	0	7	7	0	0	4	2	2
Clay	9	9	2	2	0	0	7	6	0	1	5	3	1
Coahoma	60	58	8	7	1	0	50	23	6	21	29	16	13
Copiah	25	24	8	8	0	0	16	13	1	2	14	8	2
Covington	2	8	3	3	0	0	5	2	2	1	3	3	2
DeSoto	13	18	3	3	0	0	15	4	0	11	13	5	0
Forrest	59	50	25	25	0	0	25	24	0	1	18	11	21
Franklin	3	8	5	5	0	0	3	2	0	1	4	3	1
George	4	6	5	5	0	0	1	0	0	1	5	1	0
Greene	9	11	3	2	0	1	8	2	1	5	9	2	0
Grenada	34	36	8	8	0	0	28	8	11	9	13	12	11
Hancock	5	5	2	2	0	0	3	1	0	2	3	1	1
Harrison	112	99	74	73	0	1	25	17	1	7	39	27	33

Table 4 - Page 2
Fetal Deaths - Mississippi - 1961

County	Total by Occurrence	Total by Residence	WHITE				NONWHITE				SEX		
			Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female	Unde-termined
Hinds	414	340	221	179	41	1	119	91	14	14	57	63	220
Holmes	39	42	7	7	0	0	35	13	5	17	22	14	6
Humphreys	17	17	1	0	1	0	16	4	0	12	4	12	1
Issaquena	0	2	1	1	0	0	1	0	1	0	1	0	1
Itawamba	1	6	4	4	0	0	2	2	0	0	4	2	0
Jackson	71	78	59	56	3	0	19	10	8	1	26	20	32
Jasper	3	12	2	2	0	0	10	7	0	3	7	4	1
Jefferson	7	11	1	1	0	0	10	4	2	4	5	4	2
Jeff Davis	15	19	9	9	0	0	10	7	0	3	11	4	4
Jones	82	43	21	20	1	0	22	21	0	1	13	14	16
Kemper	6	7	3	3	0	0	4	4	0	0	2	3	2
Lafayette	18	15	5	5	0	0	10	4	0	6	10	4	1
Lamar	2	10	8	8	0	0	2	2	0	0	5	2	3
Lauderdale	85	62	21	21	0	0	41	34	5	2	34	18	10
Lawrence	4	10	5	5	0	0	5	3	0	2	1	7	2
Leake	9	12	3	3	0	0	9	4	0	5	5	7	0
Lee	51	46	33	33	0	0	13	11	0	2	14	11	21
Leflore	106	95	26	26	0	0	69	34	2	33	26	28	41
Lincoln	21	24	4	4	0	0	20	17	0	3	9	11	4
Lowndes	54	49	22	22	0	0	27	18	0	9	27	12	10
Madison	55	59	15	13	2	0	44	25	0	19	25	14	20
Marion	11	14	8	8	0	0	6	3	0	3	9	2	3
Marshall	39	36	7	7	0	0	29	12	1	16	20	10	6
Monroe	22	24	14	14	0	0	10	8	0	2	8	8	8
Montgomery	12	12	4	4	0	0	8	4	1	3	4	6	2
Neshoba	12	11	7	7	0	0	4	3	0	1	4	4	3
Newton	35	36	16	15	1	0	20	14	5	1	12	9	15
Noxubee	30	28	1	1	0	0	27	14	1	12	15	9	4
Oktibbeha	22	25	7	7	0	0	18	12	0	6	14	9	2
Panola	25	29	2	2	0	0	27	18	6	3	14	14	1

Table 4 - Page 3
Fetal Deaths - Mississippi - 1961

County	Total by Occurrence	Total by Residence	WHITE				NONWHITE				SEX		
			Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female	Unde-termined
Pearl River	11	7	6	6	0	0	1	1	0	0	4	3	0
Perry	0	8	2	2	0	0	6	6	0	0	4	4	0
Pike	35	27	12	12	0	0	15	13	0	2	13	10	4
Pontotoc	11	15	11	10	1	0	4	4	0	0	5	7	3
Prentiss	7	4	3	3	0	0	1	0	0	1	2	2	0
Quitman	18	23	0	0	0	0	23	17	3	3	12	10	1
Rankin	8	40	27	24	3	0	13	12	0	1	9	7	24
Scott	21	25	6	6	0	0	19	12	1	6	10	10	5
Sharkey	6	9	0	0	0	0	9	6	0	3	5	4	0
Simpson	9	15	8	7	1	0	7	7	0	0	8	3	4
Smith	2	8	4	4	0	0	4	2	1	1	4	2	2
Stone	5	7	4	4	0	0	3	2	0	1	2	3	2
Sunflower	41	46	9	9	0	0	37	19	4	14	21	18	7
Tallahatchie	37	40	12	12	0	0	28	12	3	13	16	13	11
Tate	17	24	2	2	0	0	22	6	3	13	13	10	1
Tippah	35	31	19	19	0	0	12	10	1	1	6	6	19
Tishomingo	2	4	4	4	0	0	0	0	0	0	2	2	0
Tunica	33	34	3	3	0	0	31	12	5	14	17	17	0
Union	32	26	24	24	0	0	2	1	0	1	5	5	16
Walthall	29	20	5	5	0	0	15	11	0	4	3	6	11
Warren	74	47	18	17	1	0	29	25	0	4	18	11	18
Washington	189	177	82	81	1	0	95	75	4	16	45	29	103
Wayne	8	13	8	7	1	0	5	4	0	1	9	3	1
Webster	3	4	1	1	0	0	3	3	0	0	1	2	1
Wilkinson	38	24	4	3	0	1	20	11	0	9	10	8	6
Winston	12	18	6	6	0	0	12	5	4	3	9	7	2
Yalobusha	7	5	2	1	1	0	3	2	0	1	1	3	1
Yazoo	32	42	10	10	0	0	32	9	1	22	23	15	4

Table 4 - Page 4
Fetal Deaths - Mississippi - 1961

City*	Total by Occurrence	Total by Residence	WHITE				NONWHITE				SEX		
			Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female	Unde-termined
Biloxi	76	50	44	44	0	0	6	6	0	0	20	8	22
Clarksdale	36	17	5	5	0	0	12	9	2	1	9	4	4
Cleveland	10	9	5	5	0	0	4	2	1	1	6	2	1
Columbus	42	29	15	15	0	0	14	8	0	6	15	9	5
Corinth	8	3	2	2	0	0	1	1	0	0	1	1	1
Greenville	145	125	67	66	1	0	58	50	1	7	25	11	89
Greenwood	76	39	20	20	0	0	19	13	1	5	10	5	24
Gulfport	32	24	10	10	0	0	14	10	0	4	10	10	4
Hattiesburg	59	42	19	19	0	0	23	22	0	1	16	10	16
Jackson	401	297	197	158	38	1	100	78	14	8	47	51	199
Laurel	82	23	6	5	1	0	17	16	0	1	8	6	9
McComb	21	6	1	1	0	0	5	5	0	0	2	3	1
Meridian	83	39	14	14	0	0	25	19	5	1	17	13	9
Natchez	76	34	18	17	1	0	16	16	0	0	13	11	10
Pascagoula	5	39	30	30	0	0	9	6	3	0	9	8	22
Tupelo	47	19	13	13	0	0	6	6	0	0	7	2	10
Vicksburg	73	32	10	9	1	0	22	19	0	3	12	8	12
Yazoo City	14	13	6	6	0	0	7	2	1	4	7	4	2

* Cities having 10,000 population and over according to 1960 census.

TABLE 5
DEATHS
MISSISSIPPI
1961

By Cause of Death, by Race, Resident Data

International List Number*	Cause of Death**	Total	White	Non-white
001-999	TOTAL ALL CAUSES	21,232	11,063	10,169
001-019	Tuberculosis, all forms	113	54	59
002	Pulmonary	100	49	51
008	Unspecified site	3	1	2
010	Meninges and central nervous system	2	0	2
011	Intestines, peritoneum, and mesenteric glands	4	2	2
012	Bones, and joints, active or unspecified	1	1	0
018	Other organs	2	1	1
019	Disseminated	1	0	1
020-029	Syphilis and its sequelae	31	4	27
022	Aneurysm of aorta	13	4	9
023	Other cardiovascular	17	0	17
025	General paralysis of insane	1	0	1
040	Typhoid fever	2	0	2
041	Paratyphoid fever	1	0	1
045-048	Dysentery, all forms	17	4	13
045	Bacillary	3	1	2
046	Amebiasis	11	3	8
048	Unspecified forms	3	0	3
049	Food poisoning	1	0	1
050,051	Scarlet fever and streptococcal sore throat	2	0	2
051	Streptococcal sore throat	2	0	2
053	Septicemia and pyemia	36	14	22
055	Diphtheria	3	0	3
056	Whooping cough	5	0	5
057	Meningococcal infections	3	2	1
061	Tetanus	15	2	13
063	Gas gangrene	1	0	1
064	Other bacterial diseases	1	1	0
080	Acute poliomyelitis	4	2	2
081	Late effects of acute poliomyelitis	1	0	1
082	Acute infectious encephalitis	6	2	4
085	Measles	9	4	5
086	Rubella (German measles)	1	0	1
087	Chickenpox	1	0	1
088	Herpes zoster	2	2	0
089	Mumps	1	0	1
092	Infectious hepatitis	13	8	5
096	Other diseases attributable to viruses	4	2	2
130	Infestation with worms of other, mixed, and unspecified type	2	0	2
134	Other fungus infections	7	3	4
138	Other infective and parasitic diseases	7	1	6
140-205	Malignant neoplasms	2,748	1,732	1,016
140-148	Buccal cavity and pharynx	57	42	15
140	Lip	3	3	0

Table 5 - Page 2
Deaths - Mississippi - 1961

International List Number*	Cause of Death**	Total	White	Non-white
141	Tongue	7	5	2
142	Salivary gland	8	5	3
143	Floor of mouth	1	1	0
144	Other parts of mouth and mouth unspecified	10	10	0
145	Oral mesopharynx	5	3	2
146	Nasopharynx	4	4	0
147	Hypopharynx	1	0	1
148	Pharynx, unspecified	18	11	7
150-159	Digestive organs and peritoneum	868	508	360
150	Esophagus	52	31	21
151	Stomach	241	117	124
152	Small intestine, including duodenum	5	5	0
153	Large intestine, except rectum	232	143	89
154	Rectum	69	46	23
155	Biliary passages and liver (stated to be primary site)	41	30	11
156	Liver (secondary and unspecified)	67	32	35
157	Pancreas	148	97	51
158	Peritoneum	2	2	0
159	Unspecified digestive organs	11	5	6
160-165	Respiratory system	399	306	93
160	Nose, nasal cavities, middle ear, and accessory sinuses	7	5	2
161	Larynx	17	14	3
162	Bronchus and trachea, and lung specified as primary	149	107	42
163	Lung, unspecified as to whether primary or secondary	213	172	41
164	Mediastinum	3	1	2
165	Thoracic organs (secondary)	10	7	3
170-181	Breast and genito-urinary organs	827	465	362
170	Breast	213	141	72
171	Cervix uteri	145	67	78
172	Corpus uteri	10	2	8
173	Other parts of uterus	1	0	1
174	Uterus, unspecified	67	25	42
175	Ovary, Fallopian tube, and broad ligament	60	43	17
176	Other and unspecified female genital organs	11	9	2
177	Prostate	213	109	104
178	Testis	3	2	1
179	Other and unspecified male genital organs	7	2	5
180	Kidney	42	30	12
181	Bladder and other urinary organs	55	35	20
190-199	Other and unspecified sites	335	226	109
190	Melanoma of skin	25	21	4
191	Skin, other	25	20	5
192	Eye	3	3	0
193	Brain and other parts of nervous system	61	48	13
194	Thyroid gland	3	2	1
195	Other endocrine glands	1	1	0
196	Bone	21	13	8
197	Connective tissue	8	5	3
198	Lymph nodes (secondary and unspecified)	3	1	2

Table 5 - Page 3
Deaths - Mississippi - 1961

International List Number*	Cause of Death**	Total	White	Non-white
199	Other and unspecified sites	185	112	73
200-205	Lymphatic and hematopoietic tissues	262	185	77
200	Lymphosarcoma and reticulosarcoma	51	38	13
201	Hodgkin's disease	32	23	9
202	Other forms of lymphoma (reticulosis)	13	9	4
203	Multiple myeloma (plasmocytoma)	31	18	13
204	Leukemia and aleukemia	134	97	37
205	Mycosis fungoides	1	0	1
210-229	Benign neoplasms	31	16	15
211	Other parts of digestive system	1	0	1
212	Respiratory system	4	2	2
214	Uterine fibromyoma	7	2	5
216	Ovary	2	1	1
219	Kidney and other urinary organs	1	1	0
223	Brain and other parts of nervous system	8	5	3
224	Endocrine glands	3	2	1
226	Lipoma	1	1	0
227	Muscular and connective tissue	1	1	0
228	Hemangioma and lymphangioma	2	1	1
229	Other and unspecified organs and tissues	1	0	1
230-239	Neoplasms of unspecified nature	25	14	11
230	Digestive organs	1	0	1
231	Respiratory organs	3	2	1
234	Ovary	1	1	0
236	Other genito-urinary organs	1	0	1
237	Brain and other parts of nervous system	16	10	6
238	Skin and musculoskeletal system	1	0	1
239	Other and unspecified organs	2	1	1
241	Asthma	52	17	35
242	Angioneurotic edema	1	0	1
245	Other allergic disorders	2	0	2
250	Simple goiter	2	0	2
252	Thyrotoxicosis with or without goiter	4	2	2
253	Myxedema and cretinism	4	4	0
260	Diabetes mellitus	291	149	142
271	Diseases of parathyroid gland	1	1	0
272	Diseases of pituitary gland	1	0	1
274	Diseases of adrenal glands	10	9	1
277	Polyglandular dysfunction and other diseases of endocrine glands	1	1	0
286	Other avitaminoses and nutritional deficiency states	30	12	18
287	Obesity, not specified as of endocrine origin	16	6	10
289	Other metabolic diseases	4	3	1
290-293	Anemias	55	22	33
290	Pernicious and other hyperchromic	6	3	3
292	Other specified type	31	10	21
293	Unspecified type	18	9	9
294	Polycythemia	9	7	2
295	Hemophelia	1	1	0
296	Purpura and other hemorrhagic conditions	7	2	5
298	Diseases of spleen	2	1	1
304	Senile psychosis	8	5	3
305	Presenile psychosis	2	0	2

Table 5 - Page 4
Deaths - Mississippi - 1961

International List Number*	Cause of Death**	Total	White	Non-white
307	Alcoholic psychosis	1	1	0
309	Other and unspecified psychoses	5	3	2
320	Pathological personality	1	1	0
322	Alcoholism	16	8	8
323	Other drug addiction	1	1	0
325	Mental deficiency	5	4	1
330-334	Vascular lesions affecting central nervous system	2,891	1,464	1,427
330	Subarachnoid hemorrhage	94	50	44
331	Cerebral hemorrhage	1,795	842	953
332	Cerebral embolism and thrombosis	638	400	238
334	Other and ill-defined	364	172	192
340	Meningitis, except meningococcal and tuberculous	47	10	37
341	Phlebitis and thrombophlebitis of intracranial venous sinuses	1	1	0
342	Intracranial and intraspinal abscess	2	2	0
343	Encephalitis, myelitis, and encephalomyelitis (except acute infectious)	11	4	7
344	Late effects of intracranial abscess or pyogenic infection	6	3	3
345	Multiple sclerosis	11	9	2
350	Paralysis agitans	10	9	1
351	Cerebral spastic infantile paralysis	11	5	6
352	Other cerebral paralysis	8	3	5
353	Epilepsy	28	9	19
355	Other diseases of brain	12	9	3
356	Motor neurone disease and muscular atrophy	8	7	1
357	Other diseases of spinal cord	4	3	1
361	Trigeminal neuralgia	1	1	0
364	Polyneuritis and polyradiculitis	3	1	2
366	Other and unspecified forms of neuralgia and neuritis	1	1	0
377	Inflammation of optic nerve and retina	1	1	0
391	Otitis media without mention of mastoiditis	1	0	1
392	Otitis media with mastoiditis	2	1	1
400-402, 410-443	Diseases of the heart	6,561	4,022	2,539
400	Rheumatic fever without mention of heart involvement	4	1	3
401	Rheumatic fever with heart involvement	11	2	9
410	Diseases of mitral valve	46	31	15
411	Diseases of aortic valve specified as rheumatic	10	8	2
414	Other endocarditis specified as rheumatic	5	5	0
415	Other myocarditis specified as rheumatic	2	1	1
416	Other heart disease specified as rheumatic	60	27	33
420	Arteriosclerotic heart disease, including coronary disease	4,230	3,029	1,201
421	Chronic endocarditis not specified as rheumatic	74	42	32
422	Other myocardial degeneration	438	243	195
430	Acute and subacute endocarditis	12	4	8
431	Acute myocarditis not specified as rheumatic	10	5	5
433	Functional disease of heart	72	43	29

Table 5 - Page 5
Deaths - Mississippi - 1961

International List Number*	Cause of Death**	Total	White	Non-white
434	Other and unspecified diseases of heart	460	196	264
440	Essential benign hypertensive heart disease	2	1	1
441	Essential malignant hypertensive heart disease	15	3	12
442	Hypertensive heart disease with arteriolar nephrosclerosis	227	85	142
443	Other and unspecified hypertensive heart disease	883	296	587
444-447	Hypertension without mention of heart	200	64	136
444	Essential benign	51	18	33
445	Essential malignant	54	12	42
446	With arteriolar nephrosclerosis	94	34	60
447	Other	1	0	1
450	General arteriosclerosis	327	208	119
451-468	Other diseases of circulatory system	210	133	77
451	Aortic aneurysm, nonsyphilitic, and dissecting aneurysm	74	49	25
452	Other aneurysm, except of heart and aorta	10	4	6
453	Peripheral vascular disease	4	1	3
454	Arterial embolism and thrombosis	6	5	1
455	Gangrene of unspecified cause	5	2	3
456	Other diseases of arteries	15	12	3
460	Varicose veins of lower extremities	2	2	0
461	Hemorrhoids	1	0	1
462	Varicose veins of other specified sites	4	3	1
463	Phlebitis and thrombophlebitis of lower extremities	12	11	1
464	Phlebitis and thrombophlebitis of other sites	8	7	1
465	Pulmonary embolism and infarction	37	19	18
466	Other venous embolism and thrombosis	11	5	6
467	Other diseases of circulatory system	20	13	7
468	Certain diseases of lymph nodes and lymph channels	1	0	1
470	Acute nasopharyngitis (common cold)	3	0	3
471	Acute sinusitis	1	0	1
472	Acute pharyngitis	1	1	0
473	Acute tonsillitis	4	3	1
475	Acute upper respiratory infection of multiple or unspecified sites	11	2	9
480-483	Influenza	59	19	40
480	With pneumonia	31	9	22
481	With other respiratory manifestations, and influenza unqualified	23	9	14
482	With digestive manifestations, but without respiratory symptoms	4	1	3
483	With nervous manifestations, but without digestive or respiratory symptoms	1	0	1
490-493	Pneumonia	662	288	374
490	Lobar	136	58	78
491	Broncho	280	137	143
492	Primary atypical	45	18	27
493	Other and unspecified	201	75	126

Table 5 - Page 6
Deaths - Mississippi - 1961

International List Number*	Cause of Death**	Total	White	Non-white
500-502	Bronchitis	45	30	15
500	Acute	10	7	3
501	Unqualified	12	4	8
502	Chronic	23	19	4
510	Hypertrophy of tonsils and adenoids	2	1	1
511	Peritonsillar abscess (quinsy)	1	0	1
513	Chronic sinusitis	1	1	0
514	Deflected nasal septum	1	1	0
518	Empyema	5	2	3
519	Pleurisy	5	1	4
521	Abscess of lung	10	4	6
522	Pulmonary congestion and hypostasis	11	5	6
525	Other chronic interstitial pneumonia	16	10	6
526	Bronchiectasis	18	17	1
527	Other diseases of lung and pleural cavity	97	69	28
535	Other diseases of teeth and supporting structures	1	0	1
536	Stomatitis	1	0	1
537	Diseases of salivary glands	1	1	0
538	Other diseases of buccal cavity	3	1	2
539	Diseases of esophagus	7	6	1
540,541	Ulcer of stomach and duodenum	80	53	27
540	Stomach	35	21	14
541	Duodenum	45	32	13
542	Gastrojejunal ulcer	1	0	1
543	Gastritis and duodenitis	4	1	3
544	Disorders of function of stomach	2	0	2
545	Other diseases of stomach and duodenum	7	5	2
550	Acute appendicitis	14	5	9
560,561,570	Hernia and intestinal obstruction	101	50	51
560	Hernia of abdominal cavity without mention of obstruction	7	4	3
561	Hernia of abdominal cavity with obstruction	19	8	11
570	Intestinal obstruction without mention of hernia	75	38	37
571	Gastro-enteritis and colitis, except ulcerative, age 4 weeks and over	173	23	150
572	Chronic enteritis and ulcerative colitis	14	12	2
576	Peritonitis	7	3	4
578	Other diseases of intestines and peritoneum	40	22	18
580	Acute and subacute yellow atrophy of liver	8	4	4
581	Cirrhosis of liver	125	84	41
582	Suppurative hepatitis and liver abscess	2	0	2
583	Other diseases of liver	10	7	3
584	Cholelithiasis	16	12	4
585	Cholecystitis and cholangitis, without mention of calculi	12	11	1
586	Other diseases of gallbladder and biliary ducts	11	10	1
587	Diseases of pancreas	28	17	11
590,591	Acute nephritis and nephritis with edema	47	12	35
590	Acute	22	8	14
591	With edema, including nephrosis	25	4	21
592-594	Chronic and unspecified nephritis and other renal sclerosis	237	79	158

Table 5 - Page 7
Deaths - Mississippi - 1961

International List Number*	Cause of Death**	Total	White	Non-white
592	Chronic nephritis	194	61	133
593	Nephritis not specified as acute or chronic	38	15	23
594	Other renal sclerosis	5	3	2
600	Infections of kidney	132	68	64
601	Hydronephrosis	4	4	0
602	Calculi of kidney and ureter	11	10	1
603	Other diseases of kidney and ureter	10	3	7
604	Calculi of other parts of urinary system	1	0	1
605	Cystitis	6	5	1
606	Other diseases of bladder	6	4	2
608	Stricture of urethra	5	0	5
609	Other diseases of urethra	4	0	4
610	Hyperplasia of prostate	61	22	39
611	Prostatitis	6	2	4
612	Other diseases of prostate	1	0	1
617	Other diseases of male genital organs	1	0	1
624	Salpingitis and oophoritis, unqualified	2	0	2
626	Diseases of parametrium and pelvic peritoneum (female)	1	1	0
630	Infective disease of uterus, vagina, and vulva	1	1	0
634	Disorders of menstruation	1	1	0
640-689	Deliveries and complications of pregnancy, childbirth, and the puerperium	49	9	40
640	Pyelitis and pyelonephritis of pregnancy	1	0	1
642	Toxemias of pregnancy	16	2	14
648	Other complications arising from pregnancy	2	0	2
650	Abortion without mention of sepsis or toxemia	3	1	2
651	Abortion with sepsis	2	1	1
660	Delivery without mention of complication	2	1	1
670	Delivery complicated by placenta previa or antepartum hemorrhage	7	1	6
671	Delivery complicated by retained placenta	3	0	3
672	Delivery complicated by other postpartum hemorrhage	4	0	4
677	Delivery with other trauma	1	0	1
681	Sepsis of childbirth and the puerperium	3	1	2
684	Puerperal pulmonary embolism	2	1	1
685	Puerperal eclampsia	2	0	2
687	Cerebral hemorrhage in the puerperium	1	1	0
692	Other cellulitis and abscess without mention of lymphangitis	6	3	3
698	Other local infections of skin and subcutaneous tissue	3	2	1
703	Other dermatitis	1	0	1
705	Erythematous conditions	5	1	4
710	Other hypertrophic and atrophic conditions of skin	6	3	3
715	Chronic ulcer of skin	5	2	3
722	Rheumatoid arthritis and allied conditions	24	16	8
723	Osteo-arthritis (arthrosis) and allied conditions	4	2	2
724	Other specified forms of arthritis	1	1	0

Table 5 - Page 8
Deaths - Mississippi - 1961

International List Number*	Cause of Death**	Total	White	Non-white
725	Arthritis, unspecified	7	3	4
730	Osteomyelitis and periostitis	2	1	1
744	Other diseases of muscle, tendon, and fascia	4	2	2
745	Curvature of spine	1	1	0
750-759	Congenital malformations	239	138	101
750	Monstrosity	5	4	1
751	Spina bifida and meningocele	19	15	4
752	Congenital hydrocephalus	18	9	9
753	Other nervous system and sense organs	5	4	1
754	Circulatory system	119	69	50
756	Digestive system	26	10	16
757	Genito-urinary system	11	10	1
758	Bone and joint	2	1	1
759	Other and unspecified, not elsewhere classified	34	16	18
760-762	Birth injuries, postnatal asphyxia and atelectasis	307	166	141
760	Intracranial and spinal injury at birth	44	26	18
761	Other birth injury	72	27	45
762	Postnatal asphyxia and atelectasis	191	113	78
763-768	Infections of newborn	141	29	112
763	Pneumonia	86	20	66
764	Diarrhea	40	4	36
766	Pemphigus neonatorum	1	0	1
767	Umbilical sepsis	4	1	3
768	Other sepsis	10	4	6
769-776	Other diseases peculiar to early infancy, and immaturity, unqualified	753	221	532
769	Neonatal disorders arising from certain diseases of the mother during pregnancy	14	5	9
770	Hemolytic disease of newborn (erythroblastosis)	14	11	3
771	Hemorrhagic disease of newborn	16	8	8
772	Nutritional maladjustment	43	2	41
773	Ill-defined diseases peculiar to early infancy	145	68	77
774	Immaturity with mention of any other subsidiary condition	15	7	8
776	Immaturity, unqualified	506	120	386
780-795	Symptoms, senility, and ill-defined conditions	1,730	444	1,286
780	Certain symptoms referable to nervous system and special senses	8	1	7
782	Symptoms referable to cardiovascular and lymphatic system	55	23	32
783	Symptoms referable to respiratory system	6	3	3
784	Symptoms referable to upper gastro-intestinal tract	2	1	1
785	Symptoms referable to abdomen and lower gastro-intestinal tract	4	2	2
788	Other general symptoms	3	2	1
789	Abnormal urinary constituents of unspecified cause	1	1	0
790	Nervousness and debility	1	1	0
791	Headache	1	0	1
792	Uremia	30	10	20

Table 5 - Page 9
Deaths - Mississippi - 1961

International List Number*	Cause of Death**	Total	White	Non-white
794	Senility without mention of psychosis	75	17	58
795	Ill-defined and unknown causes	1,544	383	1,161
800-962	Accidents	1,427	760	667
800-802	Railway	12	3	9
810-835	Motor vehicle	585	380	205
810	Traffic, collision with railway train	33	17	16
812	Traffic, to pedestrian	103	52	51
813	Traffic, to pedal cyclist	6	4	2
815	Traffic, to rider or passenger of motorcycle in collision with other motor vehicle	2	2	0
816	Other traffic, two or more motor vehicles	209	146	63
818	Traffic, collision with animal or animal-drawn vehicle	2	2	0
819	Traffic, collision with fixed or unspecified object	27	23	4
821	Noncollision traffic, to rider of motorcycle	2	2	0
822	Traffic, overturning in roadway	8	4	4
823	Traffic, running off roadway	136	91	45
824	Other noncollision traffic	9	8	1
825	Traffic, unspecified nature	33	19	14
830	Nontraffic, to pedestrian	13	9	4
835	Nontraffic, other and unspecified nature	2	1	1
845	Other nonmotor road vehicle	1	1	0
850	Submersion of occupant of small boat	24	16	8
851	Other water transport injury by submersion	4	3	1
856	Machinery accident in water transport	1	0	1
857	Other specified accidents in water transport	3	3	0
860	Accident to personnel in military aircraft	1	1	0
863	Injury to occupant by accident to other specified aircraft	1	1	0
866	Other and unspecified aircraft	4	4	0
870	Poisoning by morphine and other opium derivatives	1	1	0
871	Poisoning by barbituric acid and derivatives	1	1	0
872	Poisoning by aspirin and salicylates	2	2	0
874	Poisoning by other analgesic and soporific drugs	1	1	0
880	Poisoning by alcohol	6	0	6
881	Poisoning by petroleum products	6	3	3
883	Poisoning by corrosive aromatics, acids, and caustic alkalis	2	1	1
885	Poisoning by lead and its compounds	2	0	2
886	Poisoning by arsenic and antimony, and their compounds	2	1	1
888	Poisoning by other and unspecified solid and liquid substances	8	2	6
890	Poisoning by utility (illuminating) gas	2	2	0
891	Poisoning by motor vehicle exhaust gas	5	5	0
894	Poisoning by other specified gases and vapors	2	1	1
900	Fall on stairs	6	3	3
901	Fall from ladders	3	3	0

Table 5 - Page 10
Deaths - Mississippi - 1961

International List Number*	Cause of Death**	Total	White	Non-white
902	Other falls from one level to another	32	21	11
903	Fall on same level	11	8	3
904	Unspecified falls	75	53	22
910	Blow from falling or projected object	23	10	13
912	Caused by machinery	40	22	18
913	Caused by cutting and piercing instruments	4	1	3
914	Caused by electric current	17	9	8
916	Caused by fire and explosion of combustible material	215	61	154
917	Caused by hot substance, corrosive liquid, and steam	8	1	7
919	Caused by firearm	67	36	31
921	Inhalation and ingestion of food causing obstruction or suffocation	29	12	17
922	Inhalation and ingestion of other object causing obstruction or suffocation	6	4	2
923	Foreign body entering other orifice	1	1	0
924	Mechanical suffocation in bed and cradle	28	8	20
925	Mechanical suffocation in other and unspecified circumstances	10	4	6
926	Lack of care of infants under 1 year of age	2	0	2
927	Caused by bites and stings of venomous animals and insects	1	0	1
928	Other accidents caused by animals	1	1	0
929	Drowning and submersion	98	37	61
932	Excessive cold	5	2	3
933	Hunger, thirst, and exposure	13	3	10
934	Cataclysm	2	2	0
935	Lightning	6	3	3
936	Other and unspecified accidents	25	8	17
946	Other complications due to non-therapeutic medical and surgical procedures	1	0	1
950	Therapeutic misadventure in surgical treatment	6	5	1
951	Therapeutic misadventure in infusion or transfusion	1	1	0
954	Therapeutic misadventure in anesthesia	1	0	1
960	Late effect of motor vehicle accident	3	1	2
961	Late effect of accidental poisoning	1	0	1
962	Late effect of other accidental injury	10	8	2
963,970-979	Suicide	150	121	29
970	By analgesic and soporific substances	2	2	0
971	By other solid and liquid substances	8	6	2
973	By other gases	3	3	0
974	By hanging and strangulation	9	9	0
975	By submersion (drowning)	3	2	1
976	By firearms and explosives	118	92	26
977	By cutting and piercing instruments	3	3	0
978	By jumping from high place	1	1	0
979	By other and unspecified means	3	3	0

Table 5 - Page 11
Deaths - Mississippi - 1961

International List Number*	Cause of Death**	Total	White	Non-white
964,980-985	Homicide	226	47	179
964	Late effect of injury purposely inflicted by another person (not in war)	1	0	1
981	Assault by firearm and explosive	153	32	121
982	Assault by cutting and piercing instruments	41	4	37
983	Assault by other means	25	9	16
984	Injury by intervention of police	3	2	1
985	Execution	3	0	3
965,990-999	Operations of war	1	1	0
965	Late effects of injuries due to war operations	1	1	0

* International List Numbers are those of the International Statistical Classification of Diseases, Injuries, and Cause of Death, Seventh Revision, 1955.

**When cause of death and International List Number are omitted, no deaths were reported.

TABLE 6
DEATHS
MISSISSIPPI
1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

County	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
STATE	21,032	10,961	10,071	21,232	11,063	10,169	9.7	8.8	11.0
Adams	461	171	290	379	144	235	10.0	7.6	12.6
Alcorn	267	225	42	278	234	44	11.0	10.7	13.2
Amite	70	25	45	137	63	74	8.8	8.8	8.8
Attala	219	132	87	247	149	98	11.6	12.6	10.3
Benton	48	35	13	63	44	19	8.2	10.7	5.3
Bolivar	599	99	500	613	130	483	11.2	7.4	13.1
Calhoun	141	104	37	158	122	36	9.9	10.5	8.3
Carroll	65	22	43	118	57	61	10.6	12.2	9.4
Chickasaw	173	97	76	173	97	76	10.2	9.3	11.7
Choctaw	84	51	33	99	66	33	11.8	11.2	13.1
Claiborne	90	19	71	121	30	91	11.2	11.5	11.0
Clarke	135	79	56	192	120	72	11.6	12.0	11.1
Clay	170	77	93	174	79	95	9.2	8.6	9.8
Coahoma	546	134	412	567	132	435	12.3	9.0	13.8
Copiah	251	124	127	304	156	148	11.2	12.0	10.5
Covington	90	63	27	134	86	48	9.8	9.7	10.1
DeSoto	148	41	107	207	75	132	8.7	8.1	9.0
Forrest	552	386	166	535	342	193	10.1	9.0	13.1
Franklin	80	48	32	102	59	43	11.0	10.8	11.3
George	77	68	9	99	88	11	8.9	9.0	8.5
Greene	65	49	16	74	54	20	8.8	8.4	10.4
Grenada	219	107	112	195	78	117	10.6	8.3	12.9
Hancock	150	132	18	158	137	21	11.2	11.6	9.3
Harrison	991	780	211	825	633	192	6.9	6.3	10.0
Hinds	2,165	1,167	998	1,441	711	730	7.7	6.3	9.8
Holmes	329	76	253	363	95	268	13.4	12.5	13.7
Humphreys	172	45	127	208	55	153	10.9	9.6	11.5
Issaquena	24	5	19	40	13	27	11.2	11.0	11.2
Itawamba	86	80	6	120	113	7	8.0	8.0	8.0
Jackson	310	219	91	363	264	99	6.5	5.9	9.1
Jasper	82	37	45	158	84	74	9.3	10.0	8.7
Jefferson	88	19	69	143	38	105	14.1	15.3	13.7
Jeff Davis	99	54	45	113	53	60	8.3	8.6	8.1
Jones	704	422	282	548	378	170	9.2	8.6	11.0
Kemper	117	59	58	145	75	70	11.8	15.5	9.4
Lafayette	179	113	66	196	124	72	9.2	8.8	9.9

Table 6 - Page 2
Deaths - Mississippi - 1961

County	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Lamar	77	64	13	126	103	23	9.2	9.0	10.3
Lauderdale	946	574	372	749	434	315	11.2	9.9	13.4
Lawrence	86	49	37	123	84	39	12.0	13.2	10.1
Leake	130	89	41	162	106	56	8.7	10.0	6.9
Lee	393	290	103	380	277	103	9.4	9.1	10.0
Leflore	606	235	371	542	176	366	11.5	10.5	12.0
Lincoln	252	160	92	275	166	109	10.3	9.0	13.0
Lowndes	404	197	207	430	215	215	9.2	7.4	12.1
Madison	304	84	220	327	89	238	9.9	9.6	10.1
Marion	229	149	80	249	166	83	10.7	10.8	10.5
Marshall	204	55	149	232	70	162	9.5	9.6	9.4
Monroe	271	149	122	296	165	131	8.7	7.5	10.9
Montgomery	163	90	73	176	101	75	13.2	13.7	12.6
Neshoba	144	94	50	183	134	49	8.7	8.9	8.3
Newton	186	133	53	216	152	64	11.1	11.7	9.7
Noxubee	164	44	120	181	51	130	10.8	10.8	10.7
Oktibbeha	195	102	93	215	111	104	8.2	7.5	9.1
Panola	245	100	145	293	123	170	10.2	9.8	10.5
Pearl River	190	147	43	213	163	50	9.5	9.5	9.6
Perry	34	26	8	72	55	17	8.2	8.7	7.0
Pike	375	212	163	401	224	177	11.4	11.4	11.5
Pontotoc	178	151	27	192	165	27	11.1	11.8	8.2
Prentiss	178	152	26	171	144	27	9.5	9.1	12.4
Quitman	172	42	130	218	62	156	10.4	8.0	11.7
Rankin	335	178	157	238	150	88	6.9	7.0	6.9
Scott	169	118	51	196	133	63	9.2	10.2	7.7
Sharkey	105	31	74	133	39	94	12.4	12.0	12.5
Simpson	193	121	72	174	104	70	8.5	7.8	9.7
Smith	66	52	14	123	97	26	8.6	8.8	8.0
Stone	55	41	14	61	43	18	8.7	8.1	10.5
Sunflower	379	100	279	456	140	316	10.0	9.5	10.2
Tallahatchie	234	77	157	263	88	175	10.9	10.2	11.3
Tate	108	50	58	142	67	75	7.8	8.7	7.2
Tippah	141	114	27	147	118	29	9.7	9.6	10.5
Tishomingo	122	114	8	151	143	8	10.9	10.8	11.8
Tunica	201	38	163	209	39	170	12.4	11.1	12.8
Union	171	138	33	174	142	32	9.2	9.1	9.7
Walthall	132	76	56	123	67	56	9.1	9.0	9.2
Warren	694	250	444	505	200	305	12.0	8.9	15.4
Washington	818	236	582	854	232	622	10.8	6.6	14.3
Wayne	141	85	56	162	95	67	10.0	9.1	11.5
Webster	102	77	25	126	96	30	11.9	12.1	11.4
Wilkinson	140	52	88	145	43	102	11.0	11.3	10.8
Winston	143	86	57	165	99	66	8.6	9.1	7.9
Yalobusha	128	73	55	159	97	62	12.7	13.9	11.2
Yazoo	283	102	181	314	117	197	9.9	9.1	10.5
Out-of-State	851	618	233	651	516	135	-	-	-

Table 6 - Page 3
Deaths - Mississippi - 1961

City**	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Biloxi	300	250	50	269	220	49	6.1	5.7	8.6
Clarksdale	315	114	201	251	94	157	11.9	9.6	13.9
Cleveland	78	42	36	75	34	41	7.4	5.5	10.2
Columbus	287	146	141	255	127	128	10.3	8.6	12.9
Corinth	198	160	38	146	111	35	12.7	12.1	15.1
Greenville	522	189	333	460	146	314	11.1	6.9	15.4
Greenwood	371	158	213	265	106	159	13.0	10.8	15.0
Gulfport	393	291	102	236	166	70	7.8	7.0	11.0
Hattiesburg	463	332	131	359	206	153	10.3	8.7	13.6
Jackson	1,902	1,080	822	1,057	590	467	7.3	6.4	9.0
Laurel	568	313	255	268	150	118	9.6	8.4	11.8
McComb	198	130	68	142	93	49	11.8	10.8	14.3
Meridian	822	503	319	552	321	231	11.2	9.8	13.8
Natchez	414	159	255	272	105	167	11.4	9.2	13.5
Pascagoula	40	22	18	118	79	39	6.9	6.0	10.0
Tupelo	252	197	55	150	109	41	8.7	8.3	9.9
Vicksburg	606	224	382	355	148	207	12.2	9.5	15.2
Yazoo City	165	69	96	128	47	81	11.4	9.3	13.1
Total Urban#	11,262	6,328	4,934	7,802	4,199	3,603	9.9	8.4	12.5
Total Rural	9,770	4,633	5,137	13,430	6,864	6,566	9.6	9.1	10.4

* Rate per 1,000 population.

** Cities having 10,000 population and over according to 1960 census.

Includes all incorporated towns of 2,500 population or more according to 1960 census.

TABLE 7
DEATHS
MISSISSIPPI
1961

By Cause Groups and Certain Selected Causes, by Race, Sex, and Age
With Total Rates, Resident Data

International List Number#	Cause of Death	Total	Rate*	Race	Total by Race	Sex	Total by Sex and Race	Age in Years: Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 & Over	Unknown
001-999	ALL CAUSES					M	6,486	384	57	26	40	160	141	274	702	1,138	3,563	1
				W	11,063	F	4,577	261	44	26	21	51	84	139	260	519	3,171	1
						M	5,448	884	132	49	55	156	150	244	522	717	2,529	10
		21,232	9.7	C	10,169	F	4,721	721	124	51	27	79	138	263	450	655	2,204	9
001-138	Infective and parasitic diseases					M	56	1	3	3	0	2	3	3	8	8	25	0
				W	105	F	49	2	2	2	0	1	2	5	5	8	22	0
						M	102	14	9	3	1	0	2	6	26	18	22	1
		289	13.3	C	184	F	82	13	12	2	4	4	13	9	6	8	11	0
001-019	Tuberculosis (all forms)					M	32	0	0	0	0	1	0	3	4	6	18	0
				W	54	F	22	0	0	0	0	0	1	4	2	4	11	0
						M	35	0	2	0	0	0	0	3	13	4	12	1
		113	5.2	C	59	F	24	0	1	0	0	4	8	2	2	3	4	0
001-008	Tuberculosis (respiratory system)					M	32	0	0	0	0	1	0	3	4	6	18	0
				W	50	F	18	0	0	0	0	0	1	4	2	2	9	0
						M	33	0	1	0	0	0	0	3	12	4	12	1
		103	4.7	C	53	F	20	0	0	0	0	3	7	2	2	3	3	0
020-029	Syphilis & sequelae					M	4	0	0	0	0	0	0	0	0	2	2	0
				W	4	F	0	0	0	0	0	0	0	0	0	0	0	0
						M	20	0	0	0	0	0	0	1	5	10	4	0
		31	1.4	C	27	F	7	0	0	0	0	0	0	1	2	3	1	0

Table 7 - Page 2
Deaths - Mississippi - 1961

International List Number#	Cause of Death	Total	Rate*	Race	Total by Race	Sex	Total by Sex and Race	Age in Years: Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 & Over	Unknown
045-048	Dysentery (all forms)					M	1	0	0	0	0	0	0	0	0	0	1	0
				W	4	F	3	0	0	0	0	0	0	0	0	1	2	0
						M	9	3	0	0	0	0	0	0	1	2	3	0
		17	0.8	C	13	F	4	1	0	0	0	0	0	0	0	1	2	0
050,051	Scarlet fever and streptococcal sore throat					M	0	0	0	0	0	0	0	0	0	0	0	0
				W	0	F	0	0	0	0	0	0	0	0	0	0	0	0
						M	2	0	2	0	0	0	0	0	0	0	0	0
		2	0.1	C	2	F	0	0	0	0	0	0	0	0	0	0	0	0
055	Diphtheria					M	0	0	0	0	0	0	0	0	0	0	0	0
				W	0	F	0	0	0	0	0	0	0	0	0	0	0	0
						M	2	0	1	1	0	0	0	0	0	0	0	0
		3	0.1	C	3	F	1	0	0	1	0	0	0	0	0	0	0	0
056	Whooping cough					M	0	0	0	0	0	0	0	0	0	0	0	0
				W	0	F	0	0	0	0	0	0	0	0	0	0	0	0
						M	1	1	0	0	0	0	0	0	0	0	0	0
		5	0.2	C	5	F	4	3	1	0	0	0	0	0	0	0	0	0
057	Meningococcal infections					M	1	0	0	1	0	0	0	0	0	0	0	0
				W	2	F	1	0	1	0	0	0	0	0	0	0	0	0
						M	0	0	0	0	0	0	0	0	0	0	0	0
		3	0.1	C	1	F	1	0	1	0	0	0	0	0	0	0	0	0
080	Acute poliomyelitis					M	2	0	0	1	0	1	0	0	0	0	0	0
				W	2	F	0	0	0	0	0	0	0	0	0	0	0	0
						M	2	0	1	1	0	0	0	0	0	0	0	0
		4	0.2	C	2	F	0	0	0	0	0	0	0	0	0	0	0	0
085	Measles					M	3	0	3	0	0	0	0	0	0	0	0	0
				W	4	F	1	0	0	1	0	0	0	0	0	0	0	0
						M	2	0	2	0	0	0	0	0	0	0	0	0
		9	0.4	C	5	F	3	0	2	1	0	0	0	0	0	0	0	0

Table 7 - Page 3
Deaths - Mississippi - 1961

International List Number#	Cause of Death	Total	Rate*	Race	Total by Race	Sex	Total by Sex and Race	Age in Years: Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 & Over	Unknown
140-205	Malignant neoplasms, including neoplasms of lymphatic and hematopoietic tissues	2,748	126.2	W	1,732	M	945	2	5	2	4	14	17	39	126	219	517	0
						F	787	0	3	6	2	6	17	39	107	172	435	0
				C	1,016	M	515	2	5	0	3	4	7	21	56	112	304	1
						F	501	1	8	2	0	5	13	40	92	104	236	0
210-239	Benign and unspecified neoplasms	56	2.6	W	30	M	13	0	1	0	0	0	2	0	4	4	2	0
						F	17	0	1	0	0	1	1	5	0	3	6	0
				C	26	M	7	0	0	0	0	0	1	0	2	2	2	0
						F	19	0	1	1	0	1	1	3	3	8	1	0
240-289	Allergic, endocrine system, metabolic, & nutritional diseases	419	19.2	W	204	M	88	0	0	0	0	2	2	4	13	15	52	0
						F	116	1	0	0	0	1	3	5	3	21	82	0
				C	215	M	79	3	2	3	1	1	5	5	7	6	46	0
						F	136	3	2	2	0	1	4	4	24	31	65	0
260	Diabetes mellitus	291	13.4	W	149	M	59	0	0	0	0	1	1	3	7	8	39	0
						F	90	0	0	0	0	1	1	4	1	15	68	0
				C	142	M	44	0	0	1	1	1	2	2	4	3	30	0
						F	98	0	0	0	0	1	1	2	24	24	46	0
290-299	Diseases of the blood and blood-forming organs	74	3.4	W	33	M	19	0	2	0	0	0	1	1	1	4	10	0
						F	14	0	0	0	0	0	0	1	4	0	9	0
				C	41	M	20	1	2	1	1	3	1	1	0	4	6	0
						F	21	1	2	1	1	3	5	3	2	1	2	0
~~290-293~~	~~Anemias~~	55	2.5	W	22	M	11	0	2	0	0	0	0	0	1	2	6	0
						F	11	0	0	0	0	0	0	0	3	0	8	0
				C	33	M	16	1	2	1	1	3	1	0	0	3	4	0
						F	17	1	2	1	1	3	5	1	1	0	2	0
300-326	Mental, psychoneurotic & personality disorders	39	1.8	W	23	M	18	1	0	0	1	1	0	2	6	2	5	0
						F	5	0	1	0	0	0	1	0	0	1	2	0
				C	16	M	10	0	0	0	1	0	0	3	2	2	2	0
						F	6	0	0	0	0	0	1	1	1	1	2	0

Table 7 - Page 4
Deaths - Mississippi - 1961

International List Number#	Cause of Death	Total	Rate*	Race	Total by Race	Sex	Total by Sex and Race	Age in Years: Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 & Over	Unknown
330-334	Vascular lesions affecting central nervous system					M	737	0	0	0	0	1	5	14	26	99	592	0
				W	1,464	F	727	0	0	0	0	4	4	13	19	62	625	0
						M	653	0	1	0	1	1	7	27	64	116	435	1
		2,891	132.7	C	1,427	F	774	1	0	0	1	2	10	32	83	122	522	1
340-398	Other diseases of the nervous system and sense organs					M	40	2	3	3	1	3	3	4	7	5	9	0
				W	79	F	39	4	5	0	1	3	3	2	3	5	13	0
						M	55	14	8	3	4	7	5	2	2	3	7	0
		168	7.7	C	89	F	34	13	6	1	1	0	2	2	3	1	5	0
340	Nonmeningococcal meningitis					M	3	1	1	0	0	0	0	0	1	0	0	0
				W	10	F	7	3	3	0	0	0	0	0	0	0	1	0
						M	19	8	6	0	1	1	0	1	0	1	1	0
		47	2.2	C	37	F	18	10	4	0	1	0	0	0	1	0	2	0
400-468	Diseases of the circulatory system					M	2,635	1	3	0	0	4	13	85	281	522	1,726	0
				W	4,427	F	1,792	0	0	0	0	3	8	17	58	170	1,535	1
						M	1,506	5	3	0	1	8	15	47	158	251	1,014	4
		7,298	335.0	C	2,871	F	1,365	1	2	4	5	5	22	79	130	236	877	4
400-402	Rheumatic fever					M	3	0	0	0	0	1	1	0	0	1	0	0
				W	3	F	0	0	0	0	0	0	0	0	0	0	0	0
						M	3	0	1	0	1	0	1	0	0	0	0	0
		15	0.7	C	12	F	9	0	1	2	4	0	0	2	0	0	0	0
410-416	Chronic rheumatic heart disease					M	41	0	0	0	0	0	3	8	7	8	15	0
				W	72	F	31	0	0	0	0	1	3	3	2	3	19	0
						M	21	0	0	0	0	1	2	1	5	3	9	0
		123	5.6	C	51	F	30	0	0	1	1	1	4	5	0	7	11	0
420-422	Arteriosclerotic and degenerative heart disease					M	2,057	0	0	0	0	1	8	60	238	429	1,321	0
				W	3,314	F	1,257	0	0	0	0	0	1	9	38	124	1,084	1
						M	773	0	0	0	0	0	8	19	72	132	540	2
		4,742	217.7	C	1,428	F	655	0	0	1	0	1	7	32	48	119	444	3

Table 7 - Page 5
Deaths - Mississippi - 1961

International List Number#	Cause of Death	Total	Rate*	Race	Total by Race	Sex	Total by Sex and Race	Age in Years: Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 & Over	Unknown
430-434	Other diseases of the heart					M	162	1	3	0	0	2	0	12	15	30	99	0
				W	248	F	86	0	0	0	0	1	1	1	2	9	72	0
						M	172	4	2	0	0	5	2	4	18	34	103	0
		554	25.4	C	306	F	134	0	1	0	0	2	4	10	21	24	72	0
440-443	Hypertension with heart disease					M	170	0	0	0	0	0	0	2	9	26	133	0
				W	385	F	215	0	0	0	0	0	1	2	6	20	186	0
						M	356	0	0	0	0	0	0	13	42	60	241	0
		1,127	51.7	C	742	F	386	0	0	0	0	0	3	15	46	69	252	1
444-447	Hypertension without mention of heart					M	25	0	0	0	0	0	0	1	3	5	16	0
				W	64	F	39	0	0	0	0	0	0	0	0	5	34	0
						M	77	0	0	0	0	1	2	6	12	11	45	0
		200	9.2	C	136	F	59	0	0	0	0	0	3	12	7	6	31	0
470-527	Diseases of the respiratory system					M	257	18	5	2	1	7	2	4	17	42	159	0
				W	454	F	197	19	5	4	1	3	2	4	3	12	144	0
						M	277	106	18	3	5	1	2	6	17	21	97	1
		953	43.8	C	499	F	222	95	29	4	2	2	2	9	8	13	58	0
480-483	Influenza					M	8	0	2	0	0	0	0	1	0	1	4	0
				W	19	F	11	0	0	0	0	0	0	1	0	0	10	0
						M	21	8	2	0	0	0	0	0	0	4	7	0
		59	2.7	C	40	F	19	6	4	0	0	0	0	0	0	3	6	0
490-493	Pneumonia (excluding pneumonia of newborn)					M	145	16	1	1	0	7	0	0	8	19	93	0
				W	288	F	143	12	4	2	1	1	2	1	0	9	111	0
						M	192	84	11	2	5	0	1	2	11	7	68	1
		662	30.4	C	374	F	182	83	22	3	2	1	1	8	6	8	48	0
500-502	Bronchitis					M	19	1	1	0	0	0	0	1	1	3	12	0
				W	30	F	11	2	0	0	0	0	0	0	0	1	8	0
						M	10	1	4	0	0	0	0	0	1	2	2	0
		45	2.1	C	15	F	5	2	2	0	0	0	0	0	0	0	1	0

Table 7 - Page 6
Deaths - Mississippi - 1961

International List Number#	Cause of Death	Total	Rate*	Race	Total by Race	Sex	Total by Sex and Race	Age in Years: Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 & Over	Unknown
530-587	Diseases of the digestive system					M	212	9	4	1	1	2	3	10	35	46	101	0
				W	327	F	115	6	3	0	1	0	2	9	15	16	63	0
						M	184	73	9	0	1	4	10	3	19	16	49	0
		668	30.7	C	341	F	157	64	6	3	1	2	5	8	11	20	37	0
540,541	Ulcer of stomach and duodenum					M	44	0	0	0	0	0	1	1	9	13	20	0
				W	53	F	9	0	0	0	0	0	0	1	1	2	5	0
						M	21	0	0	0	0	1	1	0	6	6	7	0
		80	3.7	C	27	F	6	1	0	0	0	0	0	1	0	0	4	0
~~550-553~~	~~Appendicitis~~					M	3	0	1	1	0	0	0	0	1	0	0	0
				W	5	F	2	0	1	0	0	0	0	0	0	1	0	0
						M	5	0	0	0	0	0	2	0	1	0	2	0
		14	0.6	C	9	F	4	0	0	2	0	0	0	1	0	0	1	0
560,561 570	Intestinal obstruction and hernia					M	34	2	0	0	0	0	0	1	1	9	21	0
				W	50	F	16	2	0	0	0	0	1	1	2	1	9	0
						M	26	2	0	0	0	1	0	2	1	4	16	0
		101	4.6	C	51	F	25	1	0	0	1	0	2	2	5	5	9	0
543,571, 572	Gastritis, duodenitis, enteritis & colitis, except diarrhea of the newborn					M	25	6	3	0	0	0	0	1	1	3	11	0
				W	36	F	11	3	0	0	0	0	0	1	0	0	7	0
						M	84	67	8	0	0	0	4	0	1	1	3	0
		191	8.8	C	155	F	71	58	6	0	0	0	0	0	0	1	6	0
581	Cirrhosis of liver					M	56	0	0	0	0	0	0	4	16	13	23	0
				W	84	F	28	0	0	0	0	0	0	4	8	7	9	0
						M	20	2	1	0	1	0	0	0	6	3	7	0
		125	5.7	C	41	F	21	0	0	0	0	1	0	4	4	6	6	0
590-637	Diseases of the genito-urinary system					M	122	2	0	0	0	4	2	7	11	13	83	0
				W	212	F	90	0	0	0	0	1	4	8	5	8	64	0
						M	205	2	4	3	1	3	3	11	14	32	132	0
		537	24.6	C	325	F	120	1	0	2	0	1	6	19	16	24	50	1

Table 7 - Page 7
Deaths - Mississippi - 1961

International List Number#	Cause of Death	Total	Rate*	Race	Total by Race	Sex	Total by Sex and Race	Age in Years: Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 & Over	Unknown
590-594	Nephritis and nephrosis					M	49	0	0	0	0	4	2	4	7	6	26	0
				W	91	F	42	0	0	0	0	1	2	4	0	5	30	0
						M	103	1	4	3	1	3	3	4	6	15	63	0
		284	13.0	C	193	F	90	1	0	2	0	1	4	9	9	19	44	1
610	Hyperplasia of prostate					M	22	0	0	0	0	0	0	0	0	1	21	0
				W	22	F	0	0	0	0	0	0	0	0	0	0	0	0
						M	39	0	0	0	0	0	0	0	0	3	36	0
		61	2.8	C	39	F	0	0	0	0	0	0	0	0	0	0	0	0
640-689	Complications of pregnancy, child-birth and the puerperium					M	0	0	0	0	0	0	0	0	0	0	0	0
				W	9	F	9	0	0	0	1	4	4	0	0	0	0	0
						M	0	0	0	0	0	0	0	0	0	0	0	0
		49	8.2*	C	40	F	40	0	0	0	2	12	13	12	1	0	0	0
690-716	Diseases of the skin and cellular tissue					M	4	0	0	0	0	1	0	0	2	0	1	0
				W	11	F	7	0	0	0	0	0	1	1	1	1	3	0
						M	4	0	1	0	0	0	0	0	1	2	0	0
		26	1.2	C	15	F	11	0	0	0	1	0	2	2	1	1	4	0
720-749	Diseases of the bones and organs of movement					M	15	0	0	0	0	0	1	0	2	2	10	0
				W	26	F	11	0	0	0	1	0	0	0	1	0	9	0
						M	6	0	0	0	0	1	0	0	2	0	3	0
		43	2.0	C	17	F	11	0	0	0	0	0	0	0	1	0	10	0
750-759	Congenital malformations					M	86	66	6	3	2	2	1	1	3	1	1	0
				W	138	F	52	41	2	1	1	2	1	2	1	1	0	0
						M	62	47	6	4	0	2	1	1	1	0	0	0
		239	11.0	C	101	F	39	30	3	3	2	1	0	0	0	0	0	0
760-776	Certain diseases of early infancy					M	250	250	0	0	0	0	0	0	0	0	0	0
				W	416	F	166	165	1	0	0	0	0	0	0	0	0	0
						M	429	429	0	0	0	0	0	0	0	0	0	0
		1,201	20.2*	C	785	F	356	356	0	0	0	0	0	0	0	0	0	0

Table 7 - Page 8
Deaths - Mississippi - 1961

International List Number#	Cause of Death	Total	Rate*	Race	Total by Race	Sex	Total by Sex and Race	Age in Years: Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 & Over	Unknown
780-795	Symptoms, senility and ill-defined conditions					M	316	16	1	0	2	1	3	16	46	80	151	0
				W	444	F	128	11	1	0	1	1	2	7	12	16	77	0
						M	715	150	20	2	2	7	17	24	61	88	343	1
		1,730	79.4	C	1,286	F	571	107	12	4	1	9	18	21	43	70	283	3
800-999	Accidents, poisonings and violence					M	673	16	24	12	28	116	83	84	114	76	119	1
				W	929	F	256	12	20	13	12	21	29	21	23	23	82	0
						M	619	38	44	27	33	114	74	87	90	44	67	1
		1,804	82.8	C	875	F	256	35	41	22	6	31	21	19	25	15	41	0
810-835	Motor vehicle accidents					M	292	2	9	6	9	74	42	40	45	27	37	1
				W	380	F	88	3	12	5	4	10	13	9	5	15	12	0
						M	153	1	8	8	7	28	24	29	24	10	14	0
		585	26.8	C	205	F	52	0	5	7	1	12	4	8	6	4	5	0
800-802, 840-962	All other accidents					M	256	14	15	6	18	35	27	27	34	24	56	0
				W	380	F	124	9	8	7	7	7	6	5	6	3	66	0
						M	303	37	33	18	24	41	22	27	33	26	41	1
		842	38.6	C	462	F	159	34	36	15	4	9	7	1	13	7	33	0
963, 970-979	Suicide					M	90	0	0	0	1	5	10	10	24	20	20	0
				W	121	F	31	0	0	0	0	4	5	5	10	3	4	0
						M	25	0	0	0	0	5	1	5	8	0	6	0
		150	6.9	C	29	F	4	0	0	0	0	0	2	1	0	1	0	0
964, 980-985	Homicide					M	34	0	0	0	0	2	4	7	11	4	6	0
				W	47	F	13	0	0	1	1	0	5	2	2	2	0	0
						M	138	0	3	1	2	40	27	26	25	8	6	0
		226	10.4	C	179	F	41	1	0	0	1	10	8	9	6	3	3	0

International List Numbers are those of the International Statistical Classification of Diseases, Injuries, and Causes of Death, Seventh Revision, 1955.

* Crude death rate per 1,000 population. Specific death rates per 100,000 population. Rate for maternal deaths (640-689) per 10,000 live births. Rate for infant causes (760-776) per 1,000 live births.

TABLE 8
DEATHS
MISSISSIPPI
1961

By Cause of Death, by Month, Resident Data

Cause of Death and International List Number*	Total	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
TOTAL ALL CAUSES	21,232	2,063	1,770	1,747	1,753	1,647	1,664	1,640	1,597	1,661	1,838	1,801	2,051
Dysentery (045-048)	17	0	1	1	0	2	2	2	2	3	1	1	2
Whooping cough (056)	5	1	0	1	1	0	0	0	0	1	0	0	1
Acute poliomyelitis (080)	4	0	0	0	0	0	1	1	1	1	0	0	0
Measles (085)	9	0	0	2	0	1	2	0	1	0	0	0	3
Influenza (480-483)	59	21	7	6	4	0	1	2	1	0	5	7	5
Lobar pneumonia (490)	136	19	11	12	9	11	9	4	7	8	7	16	23
Bronchopneumonia (491)	280	42	29	24	22	16	15	15	16	17	23	29	32
Other & unspecified pneumonia (492,493)	246	38	20	15	23	8	19	11	11	20	19	27	35
Diarrhea & enteritis (-2 yrs.) (5710)	144	13	11	7	5	5	15	22	9	13	19	12	13
Diarrhea & enteritis (+2 yrs.) (5711)	29	1	2	1	3	1	1	3	5	5	2	1	4
Accidents (800-962)	1,427	132	109	122	108	102	98	118	129	106	122	132	149
Motor vehicle accidents (810-835)	585	48	36	41	59	48	39	48	51	48	60	50	57

* International List Numbers are those of the International Statistical Classification of Diseases, Injuries, and Causes of Death, Seventh Revision, 1955.

TABLE 9
DEATHS UNDER 1 YEAR
MISSISSIPPI
1961

By Selected Causes, by Race and Age, with Total Rates, Residence Data

Cause of Death and International List Number#	Total	Rate*	Race	Total by Race	DAYS Under 1 day	1-6	7-13	14-20	21-27	MONTHS Under 28 days	1	2	3	4	5	6	7	8	9	10	11
ALL CAUSES	2,250	37.8	W	645	282	186	21	15	5	509	23	26	19	14	15	3	16	6	3	6	5
			C	1,605	385	293	102	73	39	892	169	118	112	70	65	57	39	30	20	16	17
Dysentery, all forms (045-048)	4	0.07	W	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
			C	4	0	0	0	0	1	1	0	0	1	1	0	1	0	0	0	0	0
Septicemia and pyemia (053)	11	0.2	W	2	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0
			C	9	0	0	0	0	0	0	3	1	0	4	0	1	0	0	0	0	0
Whooping cough (056)	4	0.07	W	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
			C	4	0	0	0	0	0	0	0	2	1	0	0	0	0	0	0	1	0
Meningococcal infections (057)	0	-	W	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
			C	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Tetanus (061)	7	0.1	W	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
			C	7	0	3	2	2	0	7	0	0	0	0	0	0	0	0	0	0	0
Other infective and parasitic diseases	4	0.07	W	1	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0
			C	3	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	1	0
Malignant neoplasms (140-205)	5	0.08	W	2	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0
			C	3	0	0	0	0	0	0	1	1	0	1	0	0	0	0	0	0	0
Benign and unspecified neoplasms (210-239)	0	-	W	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
			C	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Diseases of thymus gland (273)	0	-	W	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
			C	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Meningitis, except meningococcal and tuberculous (340)	22	0.4	W	4	0	0	0	0	0	0	0	0	0	0	1	0	1	0	1	1	0
			C	18	0	0	2	0	0	2	3	1	1	1	5	2	2	0	0	1	0

Table 9 - Page 2
Deaths Under 1 Year - Mississippi - 1961

Cause of Death and International List Number#	Total	Rate*	Race	Total by Race	DAYS					MONTHS											
					Under 1 day	1-6	7-13	14-20	21-27	Under 28 days	1	2	3	4	5	6	7	8	9	10	11
Other diseases of nervous system and sense organs			W	2	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0
(330-334, 341-398)	12	0.2	C	10	0	0	0	1	0	1	2	0	2	1	0	2	0	1	1	0	0
Acute upper respiratory infections			W	2	0	1	0	0	0	1	1	0	0	0	0	0	0	0	0	0	0
(470-475)	11	0.2	C	9	0	0	1	1	1	3	2	0	1	1	0	1	0	1	0	0	0
Influenza & pneumonia, except pneumonia of			W	28	0	0	0	0	0	0	4	5	3	4	6	1	1	1	0	1	2
newborn (480-493)	209	3.5	C	181	0	1	0	1	0	2	33	20	29	16	18	19	16	8	9	3	8
			W	3	0	0	0	0	0	0	1	0	1	0	1	0	0	0	0	0	0
Bronchitis (500-502)	6	0.1	C	3	0	0	0	0	0	0	1	0	0	1	0	1	0	0	0	0	0
Other diseases of respiratory system			W	4	0	0	0	0	0	0	1	0	2	0	0	0	1	0	0	0	0
(510-527)	12	0.2	C	8	0	1	0	1	0	2	2	1	3	0	0	0	0	0	0	0	0
Hernia & intestinal obstruction			W	4	2	0	0	1	0	3	0	0	1	0	0	0	0	0	0	0	0
(560,561,570)	7	0.1	C	3	0	1	0	0	0	1	0	1	0	1	0	0	0	0	0	0	0
Gastritis, duodenitis, enteritis, & colitis, except diarrhea of new-			W	9	0	0	0	0	0	0	1	0	3	1	2	0	0	0	0	2	0
born (543,571,572)	134	2.2	C	125	0	0	0	0	1	1	30	19	18	13	12	12	9	3	3	5	0
Other diseases of digestive system (530-542,544-553,			W	2	0	1	1	0	0	2	0	0	0	0	0	0	0	0	0	0	0
573-587)	11	0.2	C	9	0	2	0	0	0	2	1	2	2	0	0	1	0	1	0	0	0
Congenital malformations (750-759)			W	107	35	29	5	4	1	74	6	10	3	3	2	1	2	3	1	1	1
	184	3.1	C	77	12	17	6	9	3	47	4	10	5	2	3	1	2	2	1	0	0

Table 9 - Page 3
Deaths Under 1 Year - Mississippi - 1961

Cause of Death and International List Number#	Total	Rate*	Race	Total by Race	DAYS Under 1 day	1-6	7-13	14-20	21-27	MONTHS Under 28 days	1	2	3	4	5	6	7	8	9	10	11
Birth injuries			W	53	32	19	2	0	0	53	0	0	0	0	0	0	0	0	0	0	0
(760,761)	116	1.9	C	63	47	16	0	0	0	63	0	0	0	0	0	0	0	0	0	0	0
Postnatal asphyxia			W	113	59	47	4	0	1	111	1	0	0	0	1	0	0	0	0	0	0
and atelectasis (762)	191	3.2	C	78	38	32	2	2	1	75	2	1	0	0	0	0	0	0	0	0	0
Pneumonia of newborn			W	20	3	13	1	2	1	20	0	0	0	0	0	0	0	0	0	0	0
(763)	86	1.4	C	66	5	25	18	10	8	66	0	0	0	0	0	0	0	0	0	0	0
Diarrhea of newborn			W	4	0	0	2	2	0	4	0	0	0	0	0	0	0	0	0	0	0
(764)	40	0.7	C	36	1	1	17	7	10	36	0	0	0	0	0	0	0	0	0	0	0
Other infections of			W	5	1	3	0	1	0	5	0	0	0	0	0	0	0	0	0	0	0
newborn (765-768)	15	0.2	C	10	2	4	2	1	0	9	0	0	0	0	0	1	0	0	0	0	0
Neonatal disorders arising from certain diseases of mother during			W	5	3	1	0	0	0	4	1	0	0	0	0	0	0	0	0	0	0
pregnancy (769)	14	0.2	C	9	4	4	1	0	0	9	0	0	0	0	0	0	0	0	0	0	0
Hemolytic diseases of newborn (erythroblastosis)			W	10	8	2	0	0	0	10	0	0	0	0	0	0	0	0	0	0	0
(770)	13	0.2	C	3	1	1	1	0	0	3	0	0	0	0	0	0	0	0	0	0	0
Hemorrhagic disease of			W	8	2	5	0	1	0	8	0	0	0	0	0	0	0	0	0	0	0
newborn (771)	16	0.3	C	8	2	2	1	2	1	8	0	0	0	0	0	0	0	0	0	0	0
Nutritional maladjustment			W	2	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0
(772)	43	0.7	C	41	0	1	3	1	1	6	8	5	10	5	3	1	0	1	0	0	2
Ill-defined diseases peculiar to early			W	68	29	37	2	0	0	68	0	0	0	0	0	0	0	0	0	0	0
infancy (773)	145	2.4	C	77	31	33	4	0	0	68	4	1	1	0	1	0	1	0	0	0	1
Immaturity with mention of any other subsidiary			W	7	7	0	0	0	0	7	0	0	0	0	0	0	0	0	0	0	0
condition (774)	15	0.2	C	8	1	1	1	1	0	4	2	2	0	0	0	0	0	0	0	0	0

Table 9 - Page 4
Deaths Under 1 Year - Mississippi - 1961

Cause of Death and International List Number#	Total	Rate*	Race	Total by Race	DAYS					MONTHS											
					Under 1 day	1-6	7-13	14-20	21-27	Under 28 days	1	2	3	4	5	6	7	8	9	10	11
Immaturity, unqualified			W	120	96	21	1	0	0	118	0	1	0	0	1	0	0	0	0	0	0
(776)	506	8.5	C	386	210	105	20	19	4	358	12	5	3	3	3	1	0	0	0	1	0
Symptoms & ill-defined			W	27	4	5	1	1	0	11	2	5	1	2	0	0	6	0	0	0	0
conditions (780-793,795)	284	4.8	C	257	29	38	15	13	7	102	36	35	27	11	16	10	5	10	3	0	2
			W	5	1	0	0	0	1	2	1	0	0	0	0	0	1	0	0	0	1
All other diseases	22	0.4	C	17	1	0	2	0	0	3	3	1	2	1	1	0	0	2	1	2	1
			W	28	0	2	2	2	1	7	2	3	5	2	0	1	3	2	1	1	1
Accidents (800-962)	100	1.7	C	72	1	5	4	2	1	13	19	9	6	8	3	2	4	1	2	2	3
			W	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Homicide (964,980-984)	1	0.02	C	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0

International List Numbers are those of the International Statistical Classification of Diseases, Injuries, and Causes of Death, Seventh Revision, 1955.

* ~~Rate~~ per 1,000 live births.

TABLE 10
MARRIAGES AND DIVORCES
MISSISSIPPI
1961

Marriages by Place of Licensure, and by Place of Residence of Bride, by Race of Bride
Divorces by Place of Occurrence, by Race

County	Marriages						Divorces		
	Place of Licensure			Residence of Bride					
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
STATE	21,610	14,438	7,172	17,727	11,223	6,504	5,213	3,545	1,668
Adams	365	253	112	286	182	104	133	88	45
Alcorn	717	657	60	269	236	33	73	59	14
Amite	125	77	48	98	54	44	17	11	6
Attala	190	108	82	181	101	80	39	19	20
Benton	57	28	29	48	32	16	14	5	9
Bolivar	479	198	281	450	174	276	98	59	39
Calhoun	131	103	28	116	92	24	31	24	7
Carroll	56	35	21	61	39	22	17	11	6
Chickasaw	187	124	63	176	111	65	21	15	6
Choctaw	81	53	28	71	44	27	7	4	3
Claiborne	58	16	42	55	13	42	26	8	18
Clarke	149	95	54	108	67	41	28	17	11
Clay	159	79	80	147	77	70	39	11	28
Coahoma	402	154	248	365	139	226	93	46	47
Copiah	168	85	83	186	95	91	70	37	33
Covington	132	93	39	131	94	37	24	14	10
DeSoto	1,129	656	473	166	61	105	41	18	23
Forrest	519	405	114	505	392	113	220	147	73
Franklin	62	39	23	70	47	23	16	13	3
George	206	191	15	112	105	7	21	21	0
Greene	76	61	15	48	33	15	15	11	4
Grenada	160	95	65	172	99	73	40	20	20
Hancock	380	347	33	100	82	18	45	38	7
Harrison	1,468	1,244	224	1,106	899	207	589	504	85
Hinds	1,558	1,016	542	1,581	1,058	523	580	412	168
Holmes	192	67	125	237	79	158	42	14	28
Humphreys	138	53	85	146	52	94	29	9	20
Issaquena	14	6	8	24	3	21	4	3	1
Itawamba	165	157	8	83	80	3	30	29	1
Jackson	683	544	139	489	398	91	153	121	32
Jasper	119	79	40	152	93	59	24	13	11
Jefferson	81	37	44	67	21	46	21	4	17
Jeff Davis	109	58	51	103	52	51	11	9	2
Jones	570	450	120	508	405	103	150	112	38
Kemper	86	38	48	72	32	40	16	10	6
Lafayette	121	77	44	112	64	48	32	22	10

Table 10 - Page 2
Marriages and Divorces - Mississippi - 1961

County	Marriages						Divorces		
	Place of Licensure			Residence of Bride					
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Lamar	161	146	15	156	141	15	20	17	3
Lauderdale	607	420	187	621	433	188	180	135	45
Lawrence	90	64	26	121	86	35	28	24	4
Leake	150	100	50	139	90	49	40	22	18
Lee	364	281	83	390	305	85	98	84	14
Leflore	422	171	251	394	167	227	79	41	38
Lincoln	225	183	42	188	147	41	52	41	11
Lowndes	363	243	120	330	211	119	106	74	32
Madison	256	99	157	218	75	143	97	28	69
Marion	234	173	61	209	154	55	59	43	16
Marshall	252	130	122	153	52	101	47	24	23
Monroe	281	184	97	276	173	103	84	68	16
Montgomery	116	79	37	128	86	42	23	16	7
Neshoba	177	134	43	183	131	52	66	48	18
Newton	164	106	58	159	101	58	32	23	9
Noxubee	109	38	71	111	37	74	14	4	10
Oktibbeha	158	87	71	133	72	61	26	17	9
Panola	232	122	110	246	128	118	58	34	24
Pearl River	306	269	37	214	176	38	81	68	13
Perry	76	64	12	70	59	11	14	11	3
Pike	274	181	93	266	177	89	91	59	32
Pontotoc	162	129	33	137	110	27	41	33	8
Prentiss	225	208	17	188	175	13	50	42	8
Quitman	190	89	101	149	67	82	31	12	19
Rankin	312	220	92	249	170	79	60	50	10
Scott	217	145	72	214	143	71	49	29	20
Sharkey	104	36	68	68	25	43	14	6	8
Simpson	223	170	53	188	139	49	42	34	8
Smith	106	82	24	110	81	29	30	28	2
Stone	92	67	25	66	45	21	18	17	1
Sunflower	312	120	192	351	135	216	52	25	27
Tallahatchie	205	77	128	182	72	110	27	14	13
Tate	192	96	96	162	56	106	26	10	16
Tippah	186	151	35	163	127	36	17	14	3
Tishomingo	159	152	7	112	110	2	30	29	1
Tunica	115	34	81	109	29	80	21	2	19
Union	172	140	32	151	122	29	33	24	9
Walthall	132	91	41	82	44	38	20	17	3
Warren	399	241	158	363	217	146	149	90	59
Washington	799	494	305	613	323	290	205	123	82
Wayne	123	74	49	105	64	41	47	33	14
Webster	70	55	15	73	57	16	11	7	4
Wilkinson	291	237	54	74	28	46	29	6	23
Winston	141	82	59	130	76	54	32	15	17
Yalobusha	106	64	42	117	73	44	15	12	3
Yazoo	268	132	136	265	129	136	60	34	26
Not stated				6	2	4			
Out-of-State				3,877	3,213	664			

TABLE 11
DIVORCES
MISSISSIPPI
1961

By Cause of Divorce, by Race

Cause	Total	White	Non-white
TOTAL	5,213	3,545	1,668
Cruel and inhuman treatment	2,666	2,208	458
Desertion	1,941	844	1,097
Adultery	219	156	63
Drunkenness	268	237	31
Penitentiary	40	32	8
Insanity	9	9	0
Annulment	24	24	0
Bigamy	19	13	6
Miscellaneous and other	13	10	3
Not stated	14	12	2

By Number of Minor Children, by Race

Minor Children	Total	White	Non-white
None	2,427	1,477	950
One	1,030	781	249
Two	825	646	179
Three or more	843	590	253
Not stated	88	51	37

By Number of Years Married, by Race

Number of Years Married	Total	White	Non-white
TOTAL	5,213	3,545	1,668
Under five	1,848	1,532	316
Under one	281	266	15
One	456	400	56
Two	405	328	77
Three	361	271	90
Four	345	267	78
5-9	1,166	812	354
10-19	1,228	793	435
Twenty and over	903	366	537
Not stated	68	42	26

TABLE 12
LIVE BIRTHS
MISSISSIPPI
1913-1961

By Year, by Race, with Rates, Occurrence Data 1913-1943,
Residence Data 1944-1961

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1913	41,103	21,874	19,229	21.9	26.6	18.2
1914	40,623	21,360	19,263	21.4	25.7	18.0
1915	39,484	20,999	18,485	20.5	23.8	17.1
1916	39,033	20,130	18,903	20.0	23.6	17.2
1917	37,141	19,056	18,085	18.8	22.0	16.3
1918	39,391	20,005	19,386	19.7	22.8	17.2
1919	42,491	22,028	20,463	21.0	24.9	18.0
1920	43,743	22,438	21,305	24.4	26.3	22.7
1921	46,482	23,703	22,779	25.6	27.3	24.1
1922	43,723	21,286	22,437	23.4	24.1	23.6
1923	43,266	21,140	22,126	23.3	23.6	23.1
1924	43,960	22,180	21,780	23.4	24.3	22.5
1925	45,563	22,265	23,298	24.0	24.1	23.9
1926	50,449	24,600	25,849	26.2	26.2	26.3
1927	49,204	24,163	25,041	25.3	25.3	25.3
1928	48,126	23,642	24,484	24.5	24.4	24.5
1929	45,570	22,105	23,465	22.9	22.5	23.3
1930	47,982	23,192	24,790	23.8	23.2	24.0
1931	45,123	21,637	23,486	22.2	21.4	22.9
1932	46,481	21,883	24,598	22.6	21.4	23.9
1933	44,184	20,407	23,777	21.3	19.7	22.9
1934	47,905	22,342	25,563	23.0	21.4	24.6
1935	48,291	21,972	26,319	23.0	20.8	25.1
1936	49,261	22,194	27,067	23.3	20.8	25.7
1937	52,170	23,285	28,885	24.4	21.6	27.3
1938	53,729	24,046	29,683	25.0	22.1	27.8
1939	52,288	23,505	28,783	24.1	21.4	26.8
1940	52,526	23,084	29,442	24.0	20.9	27.3
1941	54,561	24,167	30,394	24.7	21.6	28.0
1942	56,521	25,306	31,215	25.4	22.4	28.6
1943	59,408	27,611	31,797	28.9	26.4	31.6
1944#	56,586	25,647	30,939	28.3	25.2	31.6
1945#	53,540	23,426	30,114	26.9	23.1	30.9
1946#	61,430	29,715	31,715	26.8	25.4	28.2
1947#	66,291	32,423	33,868	28.6	27.5	29.9
1948#	65,291	29,666	35,625	28.0	24.9	31.2
1949#	66,128	28,786	37,342	28.1	24.0	32.5

Table 12 - Page 2
Live Births - Mississippi - 1913-1961

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1950#	64,427	27,188	37,239	29.6	22.9	37.6
1951#	65,512	27,798	37,714	30.1	23.4	38.1
1952#	64,509	28,073	36,436	29.6	23.6	36.8
1953#	64,061	28,312	35,749	29.4	23.8	36.1
1954#	64,748	28,781	35,967	29.7	24.2	36.3
1955#	63,244	27,546	35,698	29.0	23.2	36.0
1956#	62,178	27,278	34,900	29.5	23.8	36.2
1957#	60,712	26,804	33,908	28.8	23.4	35.2
1958#	59,189	26,812	32,377	27.1	20.9	35.9
1959#	60,562	27,979	32,583	27.2	20.9	36.6
1960#	59,210	27,198	32,012	27.2	21.6	34.8
1961#	59,523	27,435	32,088	27.3	21.8	34.8

* Rate per 1,000 population.
1913-1920, population estimates based on U. S. Censuses of 1910 and 1920.
1921-1930, estimates based on U. S. Censuses of 1920 and 1930.
1931-1940, estimates based on U. S. Censuses of 1930 and 1940.
1941 and 1942, projected estimates based on U. S. Censuses of 1930 and 1940.
1943 and 1944, estimates of civilian population obtained from Ration Book Registration.
1945, estimate of civilian population, July 1, 1945, obtained from Bureau of Census Bulletin, Series P-46, No. 3, dated February 12, 1946.
1946-1949, projected estimates as of July 1, each year based on U. S. Censuses of 1930 and 1940.
1950-1955, U. S. Census population of April 1, 1950.
1956 and 1957, estimate as of July 1, 1956, prepared by Dr. Harald A. Pedersen, Sociology Department, Mississippi State University, with adjustments by Statistical Service, Mississippi State Board of Health.
1958, estimate as of January 1, prepared by Dr. George L. Wilber and Mr. Carl T. Robbins, Sociology Department, Mississippi State University.
1959, estimate as of January 1, prepared by Dr. George L. Wilber and Ellen Bryant, Sociology Department, Mississippi State University.
1960 and 1961, U. S. Census population of April 1, 1960.

\# On basis of place of residence.

TABLE 13
DEATHS
MISSISSIPPI
1913-1961

By Year, by Race, with Rates, Occurrence Data 1913-1943,
Residence Data 1944-1961

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1913	21,305	7,643	13,662	11.4	9.3	13.0
1914	21,485 + 180	7,442	14,043	11.3	8.9	13.1
1915	24,001 + 2,516	8,088	15,913	12.5	9.6	14.7
1916	21,794 − 2,207	7,588	14,206	11.2	8.9	13.0
1917	23,579 + 1,785	8,048	15,531	11.9	9.3	14.0
1918	30,437 + 6,858	10,177	20,260	15.2	11.6	18.0
1919	22,667 − 7,770	8,142	14,525	12.7	9.5	15.5
1920	22,072 − 595	7,916	14,156	12.3	9.3	15.1
1921	19,991 − 2,081	7,308	12,683	11.0	8.4	13.4
1922	19,278 − 713	7,312	11,966	10.5	8.3	12.6
1923	20,363 + 1,085	7,688	12,675	11.0	8.6	13.2
1924	21,155 + 792	7,809	13,346	11.3	8.6	13.8
1925	22,229 + 1,074	8,143	14,086	11.7	8.8	14.4
1926	23,929 + 1,700	9,133	14,796	12.5	9.7	15.1
1927	23,257 − 672	8,765	14,492	12.0	9.2	14.6
1928	25,965 + 2,708	10,111	15,854	13.2	10.4	15.9
1929	25,868 − 97	10,048	15,820	13.0	10.2	15.7
1930	24,098 − 1,770	9,072	15,026	11.9	9.0	14.8
1931	22,007 − 2,091	8,641	13,366	10.8	8.5	13.0
1932	20,322 − 1,685	8,295	12,027	9.9	8.1	11.6
1933	21,617 + 1,295	8,898	12,719	10.4	8.6	12.2
1934	21,786 + 169	9,350	12,436	10.5	9.0	12.0
1935	21,285 − 501	9,186	12,099	10.1	8.7	11.6
1936	24,127 + 2,842	10,367	13,760	11.4	9.7	13.1
1937	24,005 − 122	10,055	13,950	11.2	9.3	13.2
1938	22,814 − 1,191	9,471	13,343	10.6	8.7	12.5
1939	22,701 − 113	9,464	13,237	10.5	8.6	12.3
1940	23,185 + 484	9,565	13,620	10.6	8.6	12.6
1941	22,573 − 612	9,442	13,131	10.2	8.4	12.1
1942	20,693 − 1,880	8,750	11,943	9.3	7.7	10.9
1943	20,850 + 157	9,139	11,711	10.2	8.7	11.6
1944#	20,199 − 651	8,830	11,369	10.1	8.7	11.6
1945#	19,024 − 1,175	8,452	10,572	9.6	8.3	10.8
1946#	19,321 + 297	8,995	10,326	8.4	7.7	9.2
1947#	20,007 + 686	9,238	10,769	8.6	7.8	9.5
1948#	20,183 + 176	9,216	10,967	8.6	7.7	9.6
1949#	20,489 + 306	9,233	11,256	8.7	7.7	9.8

Table 13 - Page 2
Deaths - Mississippi - 1913-1961

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1950#	20,567+ 72	9,466	11,101	9.4	8.0	11.2
1951#	21,007+440	9,593	11,414	9.6	8.1	11.5
1952#	20,657-350	9,387	11,270	9.5	7.9	11.4
1953#	20,492-165	9,495	10,997	9.4	8.0	11.1
1954#	19,353-1,139	9,453	9,900	8.9	8.0	10.0
1955#	19,403+ 50	9,501	9,902	8.9	8.0	10.0
1956#	19,780+377	9,969	9,811	9.4	8.7	10.2
1957#	20,476+696	10,339	10,137	9.7	9.0	10.5
1958#	21,541+1,065	10,811	10,730	9.9	8.4	11.9
1959#	20,890-651	10,742	10,148	9.4	8.0	11.4
1960#	21,713+823	11,204	10,509	10.0	8.9	11.4
1961#	21,232-481	11,063	10,169	9.7	8.8	11.0

* ~~Rate~~ per 1,000 population. See footnote on Table 12 for source of population estimates.

\# On basis of place of residence.

TABLE 14
MATERNAL DEATHS
MISSISSIPPI
1913-1961

By Year, by Race, with Rates, Occurrence Data 1913-1943,
Residence Data 1944-1961

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1913	447	162	285	108.8	74.1	148.2
1914	413	124	289	101.7	58.0	150.0
1915	289	88	201	73.2	41.9	108.7
1916	326	102	224	83.5	50.7	118.5
1917	297	105	192	80.0	55.1	106.2
1918	302	82	220	76.7	41.0	113.5
1919	407	173	234	95.8	78.5	114.4
1920	393	127	266	89.8	56.6	124.8
1921	391	145	246	84.1	61.2	108.0
1922	313	110	203	71.6	51.7	90.5
1923	321	108	213	74.2	51.1	96.3
1924	413	137	276	93.9	61.8	126.7
1925	435	143	292	95.5	64.2	125.3
1926	388	153	235	76.9	62.2	90.9
1927	409	134	275	83.1	55.4	109.8
1928	421	161	260	87.5	68.1	106.2
1929	410	152	258	90.0	68.8	110.0
1930	489	171	318	101.9	73.7	128.3
1931	352	131	221	78.0	60.5	94.1
1932	279	103	176	60.0	47.1	71.6
1933	315	116	199	71.3	56.8	83.7
1934	306	116	190	63.9	51.9	74.3
1935	334	137	197	69.2	62.4	74.8
1936	358	134	224	72.7	60.4	82.8
1937	370	150	220	70.9	64.4	76.2
1938	333	116	217	62.0	48.2	73.1
1939	311	97	214	59.5	41.3	74.3
1940	327	106	221	62.2	45.9	75.1
1941	293	93	200	53.7	38.5	65.8
1942	248	77	171	43.9	30.4	54.8
1943	233	66	167	39.2	23.9	52.5
1944#	225	72	153	39.8	28.1	49.4
1945#	216	64	152	40.3	27.3	50.5
1946#	199	67	132	32.4	22.5	41.6
1947#	175	50	125	26.4	15.4	36.9
1948#	177	38	139	27.1	12.8	39.0
1949#	147	32	115	22.2	11.1	30.8

Table 14 - Page 2
Maternal Deaths - Mississippi - 1913-1961

Year	Number*			Rate*		
	Total	White	Non-white	Total	White	Non-white
1950#	165	30	135	25.6	11.0	36.2
1951#	143	28	115	21.8	10.1	30.5
1952#	119	15	104	18.4	5.3	28.5
1953#	119	24	95	18.6	8.5	26.6
1954#	104	11	93	16.1	3.8	25.8
1955#	76	12	64	12.0	4.4	17.9
1956#	74	12	62	11.9	4.4	17.8
1957#	75	14	61	12.4	5.2	18.0
1958#	53	10	43	9.0	3.7	13.3
1959#	55	9	46	9.1	3.2	14.1
1960#	59	8	51	10.0	2.9	15.9
1961#	49	9	40	8.2	3.3	12.5

* Rate per 10,000 live births.

On basis of place of residence.

TABLE 15
DEATHS UNDER 1 YEAR
MISSISSIPPI
1917-1961

By Year, by Race, with Rates, Occurrence Data 1917-1943,
Residence Data 1944-1961

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1917	3,689	1,296	2,393	99.3	68.0	132.3
1918	**	**	**	**	**	**
1919	3,554	**	**	83.6	**	**
1920	3,499	1,344	2,155	80.0	59.9	101.1
1921	3,157	1,243	1,914	67.9	52.4	84.0
1922	2,826	1,067	1,759	64.6	50.1	78.4
1923	2,908	1,104	1,804	67.2	52.2	81.5
1924	3,137	1,188	1,949	71.4	53.6	89.5
1925	3,176	1,179	1,997	69.7	53.0	85.7
1926	3,578	1,447	2,131	70.9	58.8	82.4
1927	3,291	1,303	1,988	66.9	54.0	79.4
1928	3,567	1,416	2,151	74.1	59.9	87.9
1929	3,275	1,268	2,007	71.9	57.4	85.5
1930	3,276	1,182	2,094	68.3	51.0	84.5
1931	2,547	961	1,586	56.4	44.4	67.5
1932	2,482	965	1,517	53.4	44.1	61.7
1933	2,800	1,035	1,765	63.4	50.7	74.2
1934	3,070	1,199	1,871	64.1	53.7	73.2
1935	2,587	1,044	1,543	53.6	47.5	58.6
1936	2,883	1,105	1,778	58.5	49.8	65.7
1937	3,098	1,174	1,924	59.4	50.4	66.6
1938	3,051	1,205	1,846	56.8	50.1	62.2
1939	2,908	1,135	1,773	55.6	48.3	61.6
1940	2,864	1,070	1,794	54.5	46.4	60.9
1941	2,969	1,113	1,856	54.4	46.1	61.1
1942	2,654	957	1,697	47.0	37.8	54.4
1943	2,774	1,137	1,637	46.7	41.2	51.5
1944#	2,489	974	1,515	44.0	38.0	49.0
1945#	2,192	846	1,346	40.9	36.1	44.7
1946#	2,298	1,005	1,293	37.4	33.8	40.8
1947#	2,428	1,009	1,419	36.6	31.1	41.9
1948#	2,470	988	1,482	37.8	33.3	41.6
1949#	2,612	900	1,712	39.5	31.3	45.8

Table 15 - Page 2
Deaths Under 1 Year - Mississippi - 1917-1961

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1950#	2,348	763	1,585	36.4	28.1	42.6
1951#	2,589	814	1,775	39.5	29.3	47.1
1952#	2,592	760	1,832	40.2	27.1	50.3
1953#	2,534	807	1,727	39.6	28.5	48.3
1954#	2,360	744	1,616	36.4	25.8	44.9
1955#	2,338	680	1,658	37.0	24.7	46.4
1956#	2,353	734	1,619	37.8	26.9	46.4
1957#	2,257	694	1,563	37.2	25.9	46.1
1958#	2,424	706	1,718	41.0	26.3	53.1
1959#	2,355	677	1,678	38.9	24.2	51.5
1960#	2,457	714	1,743	41.5	26.2	54.4
1961#	2,250	645	1,605	37.8	23.5	50.0

Deaths

* ~~Rate~~ per 1,000 live births.

\# On basis of place of residence.

** Not available.

TABLE 16
DEATHS UNDER 1 MONTH*
MISSISSIPPI
1924-1961

By Year, by Race, with Rates, Occurrence Data 1924-1944,
Residence Data 1945-1961

Year	Number			Rate**		
	Total	White	Non-white	Total	White	Non-white
1924	1,598	730	868	36.4	32.9	39.8
1925	1,639	701	938	36.0	31.5	40.3
1926	1,731	823	908	34.3	33.4	35.1
1927	1,644	786	858	33.4	32.5	34.3
1928	1,788	804	984	37.2	34.0	40.2
1929	1,662	773	889	36.5	35.0	37.9
1930	1,692	715	977	35.3	30.8	39.4
1931	1,339	590	749	29.7	27.3	31.9
1932	1,361	572	789	29.3	26.1	32.1
1933	1,540	636	904	34.8	31.2	38.0
1934	1,505	644	861	31.4	28.8	33.7
1935	1,446	642	804	29.9	29.2	30.5
1936	1,494	676	818	30.3	30.4	30.2
1937	1,678	755	923	32.2	32.4	32.0
1938	1,688	747	941	31.4	31.1	31.7
1939	1,637	735	902	31.3	31.3	31.3
1940	1,613	680	933	30.7	29.4	31.7
1941	1,663	734	929	30.5	30.4	30.6
1942	1,544	638	906	27.3	25.2	29.0
1943	1,652	791	861	27.8	28.6	27.1
1944	1,436	661	775	25.2	25.4	25.1
1945#	1,351	624	727	25.2	26.6	24.1
1946#	1,561	794	767	25.4	26.7	24.2
1947#	1,595	750	845	24.1	23.1	24.9
1948#	1,674	773	901	25.6	26.0	25.3
1949#	1,594	661	933	24.1	23.0	25.0
1950#	1,446	560	886	22.4	20.6	23.8
1951#	1,679	626	1,053	25.6	22.5	27.9
1952#	1,658	581	1,077	25.7	20.7	29.6
1953#	1,630	635	995	25.4	22.4	27.8
1954#	1,521	601	920	23.5	20.9	25.6
1955#	1,440	527	913	22.8	19.1	25.6
1956#	1,451	581	870	23.3	21.3	24.9
1957#	1,425	540	885	23.5	20.1	26.1
1958#	1,475	540	935	24.9	20.1	28.9
1959#	1,450	515	935	23.9	18.4	28.7
1960#	1,497	550	947	25.3	20.2	29.6
1961#	1,401	509	892	23.5	18.6	27.8

* Figures for 1949-1961 include only the deaths under 28 days.

** ~~Rate~~ Death per 1,000 live births.

\# On basis of place of residence.

TABLE 17
DEATHS FROM HEART DISEASES, NEPHRITIS AND NEPHROSIS,
VASCULAR LESIONS, MALIGNANT NEOPLASMS, AND TUBERCULOSIS (ALL FORMS)
MISSISSIPPI
1952-1961

By Year and Race, with Rates, Resident Data

YEAR	HEART DISEASES (400-402,410-443)			NEPHRITIS & NEPHROSIS (590-594)			VASCULAR LESIONS (330-334)			MALIGNANT NEOPLASMS (140-205)			TUBERCULOSIS (ALL FORMS) (001-019)		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
1952 No.	5,905	3,264	2,641	658	214	444	2,386	1,098	1,288	2,189	1,285	904	418	133	285
Rate*	271.0	274.6	266.7	30.2	18.0	44.8	109.5	92.4	130.1	100.5	108.1	91.3	19.2	11.2	28.8
1953 No.	5,790	3,196	2,594	474	164	310	2,404	1,125	1,279	2,230	1,348	882	308	85	223
Rate*	265.7	268.9	261.9	21.8	13.8	31.3	110.3	94.6	129.2	102.3	113.4	89.1	14.1	7.2	22.5
1954 No.	5,738	3,333	2,405	448	169	279	2,422	1,176	1,246	2,332	1,420	912	226	80	146
Rate*	263.3	280.4	242.9	20.6	14.2	28.2	111.2	98.9	125.8	107.0	119.5	92.1	10.4	6.7	14.7
1955 No.	5,994	3,525	2,469	401	144	257	2,447	1,169	1,278	2,317	1,402	915	211	66	145
Rate*	275.1	296.6	249.3	18.4	12.1	26.0	112.3	98.3	129.0	106.3	118.0	92.4	9.7	5.6	14.6
1956 No.	6,117	3,709	2,408	410	161	249	2,550	1,229	1,321	2,432	1,439	993	199	80	119
Rate*	289.9	323.3	250.2	19.4	14.0	25.9	120.9	107.1	137.2	115.3	125.4	103.2	9.4	7.0	12.4
1957 No.	6,378	3,711	2,667	362	143	219	2,800	1,420	1,380	2,496	1,482	1,014	161	61	100
Rate*	302.3	323.5	277.1	17.1	12.5	22.7	132.7	123.8	143.9	118.3	129.2	105.3	7.6	5.3	10.4
1958 No.	6,528	3,929	2,599	322	100	222	2,867	1,426	1,441	2,525	1,524	1,001	172	82	90
Rate*	298.9	306.3	288.4	14.7	7.8	24.6	131.3	111.2	159.9	115.6	118.8	111.1	7.9	6.4	10.0
1959 No.	6,233	3,774	2,459	290	101	189	2,683	1,386	1,297	2,600	1,580	1,020	153	62	91
Rate*	279.9	282.5	276.0	13.0	7.6	21.2	120.5	103.7	145.6	116.7	118.3	114.5	6.9	4.6	10.2
1960 No.	6,321	3,882	2,439	282	94	188	2,960	1,513	1,447	2,635	1,644	991	147	57	90
Rate*	290.2	308.7	264.9	12.9	7.5	20.4	135.9	120.3	157.2	121.0	130.7	107.6	6.7	4.5	9.8
1961 No.	6,561	4,022	2,539	284	91	193	2,891	1,464	1,427	2,748	1,732	1,016	113	54	59
Rate*	301.2	319.8	275.8	13.0	7.2	21.0	132.7	116.4	155.0	126.2	137.7	110.4	5.2	4.3	6.4

* Rate per 100,000 population.

BIRTHS, FETAL DEATHS, DEATHS

THE COUNTIES AND THE STATE OF MISSISSIPPI, 1961

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES ADAMS COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,588	649	939	1,062	443	619	28.1	23.3	33.1
Fetal deaths	77	41	36	42	21	21	39.5	47.4	33.9
Deaths from all causes	461	171	290	379	144	235	10.0	7.6	12.6
Deaths under 1 year	59	12	47	39	9	30	36.7	20.3	48.5
Deaths under 28 days	36	9	27	22	7	15	20.7	15.8	24.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	2	0	2	5.3	-	10.7
Tbc., respiratory (001-008)	1	0	1	2	0	2	5.3	-	10.7
Syphilis (020-029)	1	0	1	1	0	1	2.6	-	5.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	2	0	2	1	0	1	2.6	-	5.3
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	61	30	31	57	27	30	151.1	141.8	160.5
Diabetes (260)	10	2	8	7	1	6	18.6	5.2	32.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	87	20	67	72	17	55	190.8	89.3	294.2
Heart diseases (400-402,410-443)	143	70	73	120	56	64	318.0	294.2	342.3
Hypertension (444-447)	4	0	4	4	0	4	10.6	-	21.4
Arteriosclerosis (450)	9	4	5	8	3	5	21.2	15.8	26.7
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	15	4	11	14	4	10	37.1	21.0	53.5
#Diarrhea & enteritis (571)	5	0	5	2	0	2	5.3	-	10.7
Acute nephritis (590,591)	1	0	1	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	0	2	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	3	2	1	2	1	1	5.3	5.2	5.3
Birth inj., atelect. (760-762)	11	5	6	6	4	2	5.6	9.0	3.2
#Other dis. early inf. (763-776)	24	3	21	17	3	14	16.0	6.8	22.6
Violent & acc. deaths (800-999)	27	14	13	22	14	8	58.3	73.5	42.8
Acc. deaths, total (800-962)	25	13	12	21	13	8	55.6	68.3	42.8
Motor vehicle acc. (810-835)	15	11	4	14	11	3	37.1	57.8	16.0
Ill-defined (780-795)	15	3	12	14	2	12	37.1	10.5	64.2

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Specific cause death rates per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES ALCORN COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	534	429	105	504	400	104	19.9	18.2	31.2
Fetal deaths	10	7	3	10	7	3	19.8	17.5	28.8
Deaths from all causes	267	225	42	278	234	44	11.0	10.7	13.2
Deaths under 1 year	17	9	8	16	8	8	31.7	20.0	76.9
Deaths under 28 days	13	9	4	12	8	4	23.8	20.0	38.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	2	1	3	2	1	11.9	9.1	30.0
Tbc., respiratory (001-008)	2	1	1	2	1	1	7.9	4.6	30.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	35	31	4	42	37	5	166.1	168.6	150.0
Diabetes (260)	4	3	1	4	3	1	15.8	13.7	30.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	35	28	7	35	27	8	138.4	123.0	240.0
Heart diseases (400-402,410-443)	110	99	11	110	99	11	435.1	451.0	330.0
Hypertension (444-447)	4	1	3	5	2	3	19.8	9.1	90.0
Arteriosclerosis (450)	4	3	1	5	4	1	19.8	18.2	30.0
Influenza (480-483)	2	2	0	2	2	0	7.9	9.1	-
#Pneumonia (490-493)	6	5	1	8	7	1	31.6	31.9	30.0
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	4.0	4.6	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	4	2	2	5	3	2	9.9	7.5	19.2
#Other dis. early inf. (763-776)	6	5	1	5	4	1	9.9	10.0	9.6
Violent & acc. deaths (800-999)	16	16	0	14	14	0	55.4	63.8	-
Acc. deaths, total (800-962)	15	15	0	14	14	0	55.4	63.8	-
Motor vehicle acc. (810-835)	6	6	0	2	2	0	7.9	9.1	-
Ill-defined (780-795)	14	8	6	15	9	6	59.3	41.0	180.0

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES AMITE COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	178	10	168	340	103	237	21.8	14.4	28.1
Fetal deaths	15	3	12	34	10	24	100.0	97.1	101.3
Deaths from all causes	70	25	45	137	63	74	8.8	8.8	8.8
Deaths under 1 year	4	1	3	8	2	6	23.5	19.4	25.3
Deaths under 28 days	2	0	2	3	0	3	8.8	-	12.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	2	1	1	12.8	14.0	11.8
Tbc., respiratory (001-008)	1	0	1	2	1	1	12.8	14.0	11.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	1	0	1	6.4	-	11.8
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	6.4	-	11.8
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	7	4	3	13	6	7	83.5	84.2	82.9
Diabetes (260)	1	1	0	4	2	2	25.7	28.0	23.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	8	3	5	20	9	11	128.4	126.2	130.3
Heart diseases (400-402,410-443)	22	8	14	41	20	21	263.3	280.5	248.7
Hypertension (444-447)	2	1	1	2	1	1	12.8	14.0	11.8
Arteriosclerosis (450)	2	1	1	3	1	2	19.3	14.0	23.7
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	1	0	6	4	2	38.5	56.1	23.7
#Diarrhea & enteritis (571)	0	0	0	2	0	2	12.8	-	23.7
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	6.4	14.0	-
Birth inj., atelect. (760-762)	0	0	0	1	0	1	2.9	-	4.2
#Other dis. early inf. (763-776)	1	0	1	1	0	1	2.9	-	4.2
Violent & acc. deaths (800-999)	8	2	6	13	5	8	83.5	70.1	94.8
Acc. deaths, total (800-962)	5	1	4	10	4	6	64.2	56.1	71.1
Motor vehicle acc. (810-835)	2	1	1	4	3	1	25.7	42.1	11.8
Ill-defined (780-795)	14	3	11	17	5	12	109.2	70.1	142.1

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
ATTALA COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	482	190	292	482	183	299	22.6	15.5	31.3
Fetal deaths	20	8	12	23	10	13	47.7	54.6	43.5
Deaths from all causes	219	132	87	247	149	98	11.6	12.6	10.3
Deaths under 1 year	19	8	11	19	7	12	39.4	38.2	40.1
Deaths under 28 days	14	6	8	13	5	8	27.0	27.3	26.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	2	1	1	9.4	8.5	10.5
Tbc., respiratory (001-008)	1	0	1	2	1	1	9.4	8.5	10.5
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	33	24	9	38	26	12	178.1	220.5	125.7
Diabetes (260)	4	2	2	4	2	2	18.7	17.0	21.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	33	21	12	34	21	13	159.4	178.1	136.2
Heart diseases (400-402,410-443)	77	51	26	83	56	27	389.0	475.0	282.8
Hypertension (444-447)	3	2	1	3	2	1	14.1	17.0	10.5
Arteriosclerosis (450)	3	2	1	4	3	1	18.7	25.4	10.5
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	8	4	4	9	6	3	42.2	50.9	31.4
#Diarrhea & enteritis (571)	2	1	1	3	2	1	14.1	17.0	10.5
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	1	1	0	4.7	8.5	-
Birth inj., ateleot. (760-762)	9	5	4	9	5	4	18.7	27.3	13.4
#Other dis. early inf. (763-776)	3	1	2	2	0	2	4.1	-	6.7
Violent & acc. deaths (800-999)	14	6	8	21	11	10	98.4	93.3	104.8
Acc. deaths, total (800-962)	8	5	3	12	8	4	56.2	67.8	41.9
Motor vehicle acc. (810-835)	2	1	1	6	4	2	28.1	33.9	21.0
Ill-defined (780-795)	11	5	6	12	5	7	56.2	42.4	73.3

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES BENTON COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	123	2	121	221	68	153	28.6	16.5	42.4
Fetal deaths	2	0	2	9	5	4	40.7	73.5	26.1
Deaths from all causes	48	35	13	63	44	19	8.2	10.7	5.3
Deaths under 1 year	2	0	2	3	1	2	13.6	14.7	13.1
Deaths under 28 days	2	0	2	3	1	2	13.6	14.7	13.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	6	5	1	9	7	2	116.5	170.2	55.4
Diabetes (260)	1	0	1	1	0	1	12.9	-	27.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	6	3	3	12	6	6	155.4	145.8	166.2
Heart diseases (400-402,410-443)	18	16	2	22	20	2	284.9	486.1	55.4
Hypertension (444-447)	0	0	0	1	0	1	12.9	-	27.7
Arteriosclerosis (450)	1	0	1	2	1	1	25.9	24.3	27.7
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	2	0	2	2	0	25.9	48.6	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	1	1	0	4.5	14.7	-
#Other dis. early inf. (763-776)	0	0	0	0	0	0	-	-	-
Violent & acc. deaths (800-999)	5	5	0	4	3	1	51.8	72.9	27.7
Acc. deaths, total (800-962)	5	5	0	4	3	1	51.8	72.9	27.7
Motor vehicle acc. (810-835)	4	4	0	1	1	0	12.9	24.3	-
Ill-defined (780-795)	7	2	5	6	1	5	77.7	24.3	138.5

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
BOLIVAR COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,994	348	1,646	1,892	373	1,519	34.7	21.3	41.1
Fetal deaths	63	5	58	65	9	56	34.4	24.1	36.9
Deaths from all causes	599	99	500	613	130	483	11.2	7.4	13.1
Deaths under 1 year	96	8	88	103	11	92	54.4	29.5	60.6
Deaths under 28 days	57	6	51	59	9	50	31.2	24.1	32.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	2	0	2	3.7	-	5.4
Tbc., respiratory (001-008)	2	0	2	2	0	2	3.7	-	5.4
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	59	10	49	62	15	47	113.8	85.6	127.2
Diabetes (260)	5	2	3	5	2	3	9.2	11.4	8.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	93	12	81	77	11	66	141.4	62.8	178.6
Heart diseases (400-402,410-443)	193	45	148	200	57	143	367.2	325.3	387.1
Hypertension (444-447)	7	1	6	7	1	6	12.8	5.7	16.2
Arteriosclerosis (450)	9	1	8	10	2	8	18.4	11.4	21.6
Influenza (480-483)	2	0	2	2	0	2	3.7	-	5.4
#Pneumonia (490-493)	15	1	14	17	3	14	31.2	17.1	37.9
#Diarrhea & enteritis (571)	7	0	7	7	0	7	12.8	-	18.9
Acute nephritis (590,591)	0	0	0	1	0	1	1.8	-	2.7
Chronic nephritis, etc. (592-594)	12	0	12	9	0	9	16.5	-	24.4
Maternal deaths (640-689)	1	0	1	1	0	1	5.3	-	6.6
Cong. malformations (750-759)	3	0	3	5	2	3	9.2	11.4	8.1
Birth inj., atelect. (760-762)	13	3	10	15	6	9	7.9	16.1	5.9
#Other dis. early inf. (763-776)	39	2	37	40	1	39	21.1	2.7	25.7
Violent & acc. deaths (800-999)	32	8	24	43	12	31	79.0	68.5	83.9
Acc. deaths, total (800-962)	28	6	22	38	10	28	69.8	57.1	75.8
Motor vehicle acc. (810-835)	7	1	6	8	2	6	14.7	11.4	16.2
Ill-defined (780-795)	55	5	50	59	7	52	108.3	40.0	140.8

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES CALHOUN COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	371	171	200	377	200	177	23.6	17.2	40.7
Fetal deaths	18	7	11	11	4	7	29.2	20.0	39.5
Deaths from all causes	141	104	37	158	122	36	9.9	10.5	8.3
Deaths under 1 year	20	6	14	19	5	14	50.4	25.0	79.1
Deaths under 28 days	12	4	8	12	3	9	31.8	15.0	50.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	2	2	0	12.5	17.2	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	6.3	8.6	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	14	14	0	16	16	0	100.4	138.0	-
Diabetes (260)	1	1	0	2	2	0	12.5	17.2	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	10	7	3	12	9	3	75.3	77.6	69.0
Heart diseases (400-402,410-443)	50	40	10	57	46	11	357.6	396.7	253.1
Hypertension (444-447)	0	0	0	0	0	0	-	-	-
Arteriosclerosis (450)	5	4	1	6	5	1	37.6	43.1	23.0
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	7	6	1	6	5	1	37.6	43.1	23.0
#Diarrhea & enteritis (571)	3	0	3	3	0	3	18.8	-	69.0
Acute nephritis (590,591)	1	1	0	1	1	0	6.3	8.6	-
Chronic nephritis, etc. (592-594)	3	2	1	2	2	0	12.5	17.2	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	1	0	1	6.3	-	23.0
Birth inj., atelect. (760-762)	1	0	1	1	0	1	2.6	-	5.6
#Other dis. early inf. (763-776)	10	4	6	10	3	7	26.5	15.0	39.5
Violent & acc. deaths (800-999)	9	7	2	14	11	3	87.8	94.9	69.0
Acc. deaths, total (800-962)	7	5	2	12	9	3	75.3	77.6	69.0
Motor vehicle acc. (810-835)	4	3	1	6	4	2	37.6	34.5	46.0
Ill-defined (780-795)	11	9	2	10	9	1	62.7	77.6	23.0

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES CARROLL COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	188	23	165	220	42	178	19.7	9.0	27.4
Fetal deaths	8	1	7	13	1	12	59.1	23.8	67.4
Deaths from all causes	65	22	43	118	57	61	10.6	12.2	9.4
Deaths under 1 year	5	0	5	7	0	7	31.8	-	39.3
Deaths under 28 days	5	0	5	7	0	7	31.8	-	39.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	8.9	21.4	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	8.9	21.4	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	7	1	6	16	7	9	143.2	149.7	138.5
Diabetes (260)	1	0	1	1	0	1	8.9	-	15.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	4	2	2	22	12	10	196.8	256.6	153.8
Heart diseases (400-402,410-443)	19	13	6	28	22	6	250.5	470.4	92.3
Hypertension (444-447)	7	1	6	8	1	7	71.6	21.4	107.7
Arteriosclerosis (450)	1	0	1	2	1	1	17.9	21.4	15.4
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	1	1	3	1	2	26.8	21.4	30.8
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	0	1	8.9	-	15.4
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	4	0	4	6	0	6	27.3	-	33.7
Violent & acc. deaths (800-999)	3	2	1	5	3	2	44.7	64.1	30.8
Acc. deaths, total (800-962)	2	1	1	4	2	2	35.8	42.8	30.8
Motor vehicle acc. (810-835)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	13	2	11	13	2	11	116.3	42.8	169.2

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES CHICKASAW COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	412	226	186	439	225	214	26.0	21.7	32.9
Fetal deaths	15	3	12	22	7	15	50.1	31.1	70.1
Deaths from all causes	173	97	76	173	97	76	10.2	9.3	11.7
Deaths under 1 year	11	3	8	14	5	9	31.9	22.2	42.0
Deaths under 28 days	4	2	2	5	4	1	11.4	17.8	4.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	16	10	6	19	13	6	112.5	125.2	92.2
Diabetes (260)	2	1	1	1	0	1	5.9	-	15.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	37	15	22	35	14	21	207.2	134.9	322.5
Heart diseases (400-402,410-443)	66	42	24	67	43	24	396.7	414.2	368.6
Hypertension (444-447)	1	0	1	1	0	1	5.9	-	15.4
Arteriosclerosis (450)	2	1	1	2	1	1	11.8	9.6	15.4
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	10	8	2	11	9	2	65.1	86.7	30.7
#Diarrhea & enteritis (571)	1	0	1	1	0	1	5.9	-	15.4
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	2	0	1	1	0	5.9	9.6	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	5.9	9.6	-
Birth inj., atelect. (760-762)	1	1	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	2	0	2	3	2	1	6.8	8.9	4.7
Violent & acc. deaths (800-999)	16	11	5	14	9	5	82.9	86.7	76.8
Acc. deaths, total (800-962)	15	10	5	12	7	5	71.0	67.4	76.8
Motor vehicle acc. (810-835)	7	6	1	8	6	2	47.4	57.8	30.7
Ill-defined (780-795)	5	1	4	6	1	5	35.5	9.6	76.8

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
CHOCTAW COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	173	98	75	188	105	83	22.3	17.8	32.9
Fetal deaths	4	3	1	4	3	1	21.3	28.6	12.0
Deaths from all causes	84	51	33	99	66	33	11.8	11.2	13.1
Deaths under 1 year	4	2	2	4	2	2	21.3	19.0	24.1
Deaths under 28 days	4	2	2	4	2	2	21.3	19.0	24.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	7	3	11	8	3	130.6	135.5	119.0
Diabetes (260)	1	1	0	1	1	0	11.9	16.9	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	10	5	5	15	9	6	178.1	152.5	238.1
Heart diseases (400-402,410-443)	27	18	9	32	22	10	379.9	372.7	396.8
Hypertension (444-447)	0	0	0	0	0	0	-	-	-
Arteriosclerosis (450)	4	1	3	5	2	3	59.4	33.9	119.0
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	1	1	3	2	1	35.6	33.9	39.7
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	11.9	16.9	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	11.9	16.9	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	3	2	1	3	2	1	16.0	19.0	12.0
Violent & acc. deaths (800-999)	9	4	5	9	4	5	106.8	67.8	198.4
Acc. deaths, total (800-962)	7	3	4	7	3	4	83.1	50.8	158.7
Motor vehicle acc. (810-835)	0	0	0	4	3	1	47.5	50.8	39.7
Ill-defined (780-795)	3	1	2	3	2	1	35.6	33.9	39.7

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES CLAIBORNE COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	203	24	179	275	40	235	25.4	15.4	28.5
Fetal deaths	4	2	2	12	2	10	43.6	50.0	42.6
Deaths from all causes	90	19	71	121	30	91	11.2	11.5	11.0
Deaths under 1 year	8	0	8	13	0	13	47.3	-	55.3
Deaths under 28 days	5	0	5	9	0	9	32.7	-	38.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	9.2	38.5	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	9.2	38.5	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	11	4	7	17	7	10	156.8	269.2	121.3
Diabetes (260)	0	0	0	1	0	1	9.2	-	12.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	8	0	8	13	2	11	119.9	76.9	133.4
Heart diseases (400-402,410-443)	38	13	25	41	13	28	378.0	500.0	339.6
Hypertension (444-447)	2	0	2	3	0	3	27.7	-	36.4
Arteriosclerosis (450)	1	0	1	1	0	1	9.2	-	12.1
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	0	1	1	0	1	9.2	-	12.1
#Diarrhea & enteritis (571)	0	0	0	2	1	1	18.4	38.5	12.1
Acute nephritis (590,591)	1	0	1	1	0	1	9.2	-	12.1
Chronic nephritis, etc. (592-594)	3	1	2	4	1	3	36.9	38.5	36.4
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	2	1	1	18.4	38.5	12.1
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	5	0	5	8	0	8	29.1	-	34.0
Violent & acc. deaths (800-999)	11	1	10	12	0	12	110.6	-	145.5
Acc. deaths, total (800-962)	10	1	9	11	0	11	101.4	-	133.4
Motor vehicle acc. (810-835)	2	1	1	3	0	3	27.7	-	36.4
Ill-defined (780-795)	7	0	7	8	1	7	73.8	38.5	84.9

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
CLARKE COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	152	87	65	328	144	184	19.9	14.4	28.3
Fetal deaths	1	0	1	8	1	7	24.4	6.9	38.0
Deaths from all causes	135	79	56	192	120	72	11.6	12.0	11.1
Deaths under 1 year	7	2	5	12	5	7	36.6	34.7	38.0
Deaths under 28 days	3	1	2	8	3	5	24.4	20.8	27.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	15	11	4	23	19	4	139.4	190.0	61.6
Diabetes (260)	2	1	1	2	0	2	12.1	-	30.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	19	12	7	30	21	9	181.9	210.0	138.6
Heart diseases (400-402,410-443)	52	41	11	65	53	12	394.1	529.9	184.8
Hypertension (444-447)	1	1	0	1	1	0	6.1	10.0	-
Arteriosclerosis (450)	2	0	2	3	1	2	18.2	10.0	30.8
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	0	1	1	1	0	6.1	10.0	-
#Diarrhea & enteritis (571)	0	0	0	1	1	0	6.1	10.0	-
Acute nephritis (590,591)	0	0	0	2	1	1	12.1	10.0	15.4
Chronic nephritis, etc. (592-594)	7	0	7	8	0	8	48.5	-	123.2
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	6.1	10.0	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	3	1	2	7	2	5	21.3	13.9	27.2
Violent & acc. deaths (800-999)	11	5	6	17	10	7	103.1	100.0	107.8
Acc. deaths, total (800-962)	8	3	5	13	8	5	78.8	80.0	77.0
Motor vehicle acc. (810-835)	5	3	2	6	5	1	36.4	50.0	15.4
Ill-defined (780-795)	14	4	10	16	3	13	97.0	30.0	200.2

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES CLAY COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	483	187	296	511	208	303	27.0	22.6	31.2
Fetal deaths	9	2	7	9	2	7	17.6	9.6	23.1
Deaths from all causes	170	77	93	174	79	95	9.2	8.6	9.8
Deaths under 1 year	12	2	10	10	2	8	19.6	9.6	26.4
Deaths under 28 days	5	2	3	5	2	3	9.8	9.6	9.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	1	0	5.3	10.8	-
Tbc., respiratory (001-008)	1	0	1	1	1	0	5.3	10.8	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	1	0	1	1	0	5.3	10.8	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	12	3	9	19	9	10	100.4	97.7	102.9
Diabetes (260)	1	0	1	1	0	1	5.3	-	10.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	31	19	12	31	19	12	163.7	206.2	123.5
Heart diseases (400-402,410-443)	68	38	30	69	35	34	364.4	379.8	349.8
Hypertension (444-447)	0	0	0	0	0	0	-	-	-
Arteriosclerosis (450)	2	1	1	2	1	1	10.6	10.8	10.3
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	3	1	2	1	0	1	5.3	-	10.3
#Diarrhea & enteritis (571)	4	0	4	3	0	3	15.8	-	30.9
Acute nephritis (590,591)	1	0	1	1	0	1	5.3	-	10.3
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	5.3	-	10.3
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	1	0	1	5.3	-	10.3
Birth inj., atelect. (760-762)	2	1	1	2	1	1	3.9	4.8	3.3
#Other dis. early inf. (763-776)	4	1	3	4	1	3	7.8	4.8	9.9
Violent & acc. deaths (800-999)	13	8	5	15	8	7	79.2	86.8	72.0
Acc. deaths, total (800-962)	8	5	3	11	6	5	58.1	65.1	51.4
Motor vehicle acc. (810-835)	2	0	2	3	1	2	15.8	10.8	20.6
Ill-defined (780-795)	9	1	8	9	1	8	47.5	10.8	82.3

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
COAHOMA COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,565	390	1,175	1,584	372	1,212	34.3	25.4	38.4
Fetal deaths	60	7	53	58	8	50	36.6	21.5	41.2
Deaths from all causes	546	134	412	567	132	435	12.3	9.0	13.8
Deaths under 1 year	103	8	95	102	8	94	64.4	21.5	77.6
Deaths under 28 days	66	7	59	65	6	59	41.0	16.1	48.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	2	1	1	4.3	6.8	3.2
Tbc., respiratory (001-008)	2	1	1	2	1	1	4.3	6.8	3.2
Syphilis (020-029)	1	0	1	1	0	1	2.2	-	3.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	0	2	3	1	2	6.5	6.8	6.3
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	2.2	-	3.2
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	2.2	-	3.2
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	62	22	40	68	21	47	147.1	143.5	148.8
Diabetes (260)	7	2	5	8	2	6	17.3	13.7	19.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	52	17	35	56	16	40	121.2	109.4	126.6
Heart diseases (400-402,410-443)	155	49	106	152	47	105	328.9	321.2	332.5
Hypertension (444-447)	2	0	2	2	0	2	4.3	-	6.3
Arteriosclerosis (450)	1	1	0	2	1	1	4.3	6.8	3.2
Influenza (480-483)	8	2	6	9	2	7	19.5	13.7	22.2
#Pneumonia (490-493)	24	7	17	23	4	19	49.8	27.3	60.2
#Diarrhea & enteritis (571)	4	0	4	4	0	4	8.6	-	12.7
Acute nephritis (590,591)	3	0	3	2	0	2	4.3	-	6.3
Chronic nephritis, etc. (592-594)	6	1	5	7	1	6	15.1	6.8	19.0
Maternal deaths (640-689)	1	0	1	1	0	1	6.3	-	8.2
Cong. malformations (750-759)	4	2	2	5	3	2	10.8	20.5	6.3
Birth inj., atelect. (760-762)	10	1	9	10	1	9	6.3	2.7	7.4
#Other dis. early inf. (763-776)	47	3	44	46	2	44	29.0	5.4	36.3
Violent & acc. deaths (800-999)	41	9	32	44	14	30	95.2	95.7	95.0
Acc. deaths, total (800-962)	31	7	24	35	12	23	75.7	82.0	72.8
Motor vehicle acc. (810-835)	10	3	7	14	5	9	30.3	34.2	28.5
Ill-defined (780-795)	50	2	48	53	3	50	114.7	20.5	158.3

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES COPIAH COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	859	307	552	713	266	447	26.4	20.5	31.8
Fetal deaths	25	7	18	24	8	16	33.7	30.1	35.8
Deaths from all causes	251	124	127	304	156	148	11.2	12.0	10.5
Deaths under 1 year	20	3	17	23	2	21	32.2	7.5	47.0
Deaths under 28 days	11	2	9	12	1	11	16.8	3.8	24.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	2	0	2	2	0	7.4	15.4	-
Tbc., respiratory (001-008)	2	2	0	2	2	0	7.4	15.4	-
Syphilis (020-029)	1	0	1	1	0	1	3.7	-	7.1
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	3.7	-	7.1
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	3.7	-	7.1
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	2	2	0	7.4	15.4	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	36	21	15	44	26	18	162.6	200.1	128.0
Diabetes (260)	5	2	3	6	3	3	22.2	23.1	21.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	18	9	9	24	12	12	88.7	92.4	85.4
Heart diseases (400-402,410-443)	109	63	46	118	71	47	436.2	546.5	334.3
Hypertension (444-447)	2	1	1	3	1	2	11.1	7.7	14.2
Arteriosclerosis (450)	3	2	1	4	3	1	14.8	23.1	7.1
Influenza (480-483)	1	0	1	2	1	1	7.4	7.7	7.1
#Pneumonia (490-493)	7	2	5	9	4	5	33.3	30.8	35.6
#Diarrhea & enteritis (571)	1	0	1	1	0	1	3.7	-	7.1
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	6	2	4	5	2	3	18.5	15.4	21.3
Maternal deaths (640-689)	0	0	0	1	0	1	14.0	-	22.4
Cong. malformations (750-759)	2	1	1	3	1	2	11.1	7.7	14.2
Birth inj., atelect. (760-762)	4	1	3	4	1	3	5.6	3.8	6.7
#Other dis. early inf. (763-776)	5	1	4	7	0	7	9.8	-	15.6
Violent & acc. deaths (800-999)	13	6	7	21	10	11	77.6	77.0	78.2
Acc. deaths, total (800-962)	9	5	4	17	9	8	62.8	69.3	56.9
Motor vehicle acc. (810-835)	3	2	1	6	3	3	22.2	23.1	21.3
Ill-defined (780-795)	9	1	8	10	2	8	37.0	15.4	56.9

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES COVINGTON COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	192	141	51	330	170	160	24.2	19.1	33.7
Fetal deaths	2	0	2	8	3	5	24.2	17.6	31.2
Deaths from all causes	90	63	27	134	86	48	9.8	9.7	10.1
Deaths under 1 year	6	4	2	16	5	11	48.5	29.4	68.8
Deaths under 28 days	4	4	0	7	4	3	21.2	23.5	18.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	15	12	3	17	13	4	124.7	146.1	84.4
Diabetes (260)	1	1	0	2	2	0	14.7	22.5	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	14	10	4	20	13	7	146.6	146.1	147.6
Heart diseases (400-402,410-443)	29	21	8	36	27	9	264.0	303.5	189.8
Hypertension (444-447)	2	2	0	2	1	1	14.7	11.2	21.1
Arteriosclerosis (450)	1	0	1	1	0	1	7.3	-	21.1
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	1	1	3	2	1	22.0	22.5	21.1
#Diarrhea & enteritis (571)	0	0	0	4	0	4	29.3	-	84.4
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	3	1	2	22.0	11.2	42.2
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	2	0	2	2	0	14.7	22.5	-
Birth inj., atelect. (760-762)	1	1	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	2	2	0	6	3	3	18.2	17.6	18.8
Violent & acc. deaths (800-999)	6	3	3	16	11	5	117.3	123.6	105.5
Acc. deaths, total (800-962)	5	3	2	15	11	4	110.0	123.6	84.4
Motor vehicle acc. (810-835)	2	2	0	6	6	0	44.0	67.4	-
Ill-defined (780-795)	6	2	4	8	2	6	58.7	22.5	126.6

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES DESOTO COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	687	74	613	748	180	568	31.3	19.5	38.8
Fetal deaths	13	0	13	18	3	15	24.1	16.7	26.4
Deaths from all causes	148	41	107	207	75	132	8.7	8.1	9.0
Deaths under 1 year	22	1	21	39	6	33	52.1	33.3	58.1
Deaths under 28 days	11	1	10	22	5	17	29.4	27.8	29.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	1	0	1	3	1	2	12.6	10.8	13.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	4.2	-	6.8
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	1	0	1	4.2	-	6.8
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	15	7	8	22	14	8	92.1	151.4	54.6
Diabetes (260)	0	0	0	1	1	0	4.2	10.8	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	26	7	19	36	12	24	150.7	129.8	163.9
Heart diseases (400-402,410-443)	40	14	26	52	25	27	217.6	270.3	184.4
Hypertension (444-447)	0	0	0	1	0	1	4.2	-	6.8
Arteriosclerosis (450)	1	1	0	1	1	0	4.2	10.8	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	5	1	4	8	2	6	33.5	21.6	41.0
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	0	1	4.2	-	6.8
Maternal deaths (640-689)	0	0	0	1	0	1	13.4	-	17.6
Cong. malformations (750-759)	6	1	5	6	1	5	25.1	10.8	34.1
Birth inj., atelect. (760-762)	2	0	2	4	2	2	5.3	11.1	3.5
#Other dis. early inf. (763-776)	7	0	7	14	2	12	18.7	11.1	21.1
Violent & acc. deaths (800-999)	24	7	17	25	7	18	104.6	75.7	122.9
Acc. deaths, total (800-962)	19	5	14	20	5	15	83.7	54.1	102.4
Motor vehicle acc. (810-835)	4	3	1	5	5	0	20.9	54.1	-
Ill-defined (780-795)	17	1	16	19	1	18	79.5	10.8	122.9

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES FORREST COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,603	1,171	432	1,388	879	509	26.3	23.1	34.5
Fetal deaths	59	36	23	50	25	25	36.0	28.4	49.1
Deaths from all causes	552	386	166	535	342	193	10.1	9.0	13.1
Deaths under 1 year	51	30	21	47	23	24	33.9	26.2	47.2
Deaths under 28 days	32	21	11	28	14	14	20.2	15.9	27.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	4	3	1	7.6	7.9	6.8
Tbc., respiratory (001-008)	1	1	0	4	3	1	7.6	7.9	6.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	1	0	1	1.9	-	6.8
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	79	62	17	73	52	21	138.5	136.9	142.4
Diabetes (260)	12	9	3	11	8	3	20.9	21.1	20.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	53	42	11	53	38	15	100.5	100.1	101.7
Heart diseases (400-402,410-443)	193	144	49	189	134	55	358.5	352.9	372.8
Hypertension (444-447)	2	1	1	3	0	3	5.7	-	20.3
Arteriosclerosis (450)	9	9	0	9	9	0	17.1	23.7	-
Influenza (480-483)	4	4	0	4	4	0	7.6	10.5	-
#Pneumonia (490-493)	18	12	6	18	9	9	34.1	23.7	61.0
#Diarrhea & enteritis (571)	3	2	1	2	1	1	3.8	2.6	6.8
Acute nephritis (590,591)	2	1	1	2	1	1	3.8	2.6	6.8
Chronic nephritis, etc. (592-594)	4	3	1	6	3	3	11.4	7.9	20.3
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	9	7	2	7	6	1	13.3	15.8	6.8
Birth inj., atelect. (760-762)	8	7	1	5	4	1	3.6	4.6	2.0
#Other dis. early inf. (763-776)	20	8	12	19	5	14	13.7	5.7	27.5
Violent & acc. deaths (800-999)	69	35	34	54	23	31	102.4	60.6	210.1
Acc. deaths, total (800-962)	55	30	25	41	19	22	77.8	50.0	149.1
Motor vehicle acc. (810-835)	33	18	15	22	9	13	41.7	23.7	88.1
Ill-defined (780-795)	19	6	13	20	7	13	37.9	18.4	88.1

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES FRANKLIN COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	143	67	76	214	93	121	23.0	17.0	31.8
Fetal deaths	3	2	1	8	5	3	37.4	53.8	24.8
Deaths from all causes	80	48	32	102	59	43	11.0	10.8	11.3
Deaths under 1 year	5	3	2	9	4	5	42.0	43.0	41.3
Deaths under 28 days	4	3	1	7	4	3	32.7	43.0	24.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	14	10	4	20	13	7	215.4	237.0	184.2
Diabetes (260)	2	0	2	2	0	2	21.5	-	52.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	10	7	3	11	6	5	118.4	109.4	131.6
Heart diseases (400-402,410-443)	33	14	19	39	18	21	420.0	328.1	552.6
Hypertension (444-447)	0	0	0	0	0	0	-	-	-
Arteriosclerosis (450)	1	1	0	1	1	0	10.8	18.2	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	1	0	1	1	0	10.8	18.2	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	10.8	18.2	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	2	1	1	21.5	18.2	26.3
Birth inj., atelect. (760-762)	0	0	0	1	0	1	4.7	-	8.3
#Other dis. early inf. (763-776)	3	3	0	4	3	1	18.7	32.2	8.3
Violent & acc. deaths (800-999)	6	5	1	5	3	2	53.8	54.7	52.6
Acc. deaths, total (800-962)	6	5	1	5	3	2	53.8	54.7	52.6
Motor vehicle acc. (810-835)	3	3	0	3	3	0	32.3	54.7	-
Ill-defined (780-795)	1	1	0	3	2	1	32.3	36.4	26.3

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
GEORGE COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	314	264	50	317	266	51	28.6	27.1	39.6
Fetal deaths	4	3	1	6	5	1	18.9	18.8	19.6
Deaths from all causes	77	68	9	99	88	11	8.9	9.0	8.5
Deaths under 1 year	4	2	2	8	5	3	25.2	18.8	58.8
Deaths under 28 days	2	2	0	5	5	0	15.8	18.8	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	0	1	1	0	1	9.0	-	77.7
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	9	1	13	12	1	117.1	122.3	77.7
Diabetes (260)	1	1	0	1	1	0	9.0	10.2	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	8	8	0	10	10	0	90.1	101.9	-
Heart diseases (400-402,410-443)	27	25	2	37	34	3	333.4	346.5	233.1
Hypertension (444-447)	0	0	0	0	0	0	-	-	-
Arteriosclerosis (450)	1	1	0	2	2	0	18.0	20.4	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	4	4	0	5	4	1	45.0	40.8	77.7
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	2	0	2	2	0	18.0	20.4	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	3	2	1	27.0	20.4	77.7
Birth inj., atelect. (760-762)	0	0	0	2	2	0	6.3	7.5	-
#Other dis. early inf. (763-776)	2	2	0	2	2	0	6.3	7.5	-
Violent & acc. deaths (800-999)	10	8	2	9	7	2	81.1	71.3	155.4
Acc. deaths, total (800-962)	9	7	2	8	6	2	72.1	61.2	155.4
Motor vehicle acc. (810-835)	6	4	2	5	3	2	45.0	30.6	155.4
Ill-defined (780-795)	1	0	1	1	0	1	9.0	-	77.7

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES GREENE COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	146	96	50	198	133	65	23.7	20.6	33.8
Fetal deaths	9	1	8	11	3	8	55.6	22.6	123.1
Deaths from all causes	65	49	16	74	54	20	8.8	8.4	10.4
Deaths under 1 year	5	0	5	8	1	7	40.4	7.5	107.7
Deaths under 28 days	2	0	2	4	1	3	20.2	7.5	46.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	12.0	15.5	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	12.0	15.5	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	9	0	10	10	0	119.5	155.2	-
Diabetes (260)	3	2	1	3	2	1	35.8	31.0	52.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	6	4	2	7	4	3	83.7	62.1	156.0
Heart diseases (400-402,410-443)	23	19	4	25	20	5	298.8	310.4	260.0
Hypertension (444-447)	0	0	0	0	0	0	-	-	-
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	0	1	4	3	1	47.8	46.6	52.0
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	2	0	2	23.9	-	104.0
Birth inj., atelect. (760-762)	1	0	1	2	0	2	10.1	-	30.8
#Other dis. early inf. (763-776)	1	0	1	2	1	1	10.1	7.5	15.4
Violent & acc. deaths (800-999)	13	11	2	12	10	2	143.4	155.2	104.0
Acc. deaths, total (800-962)	13	11	2	11	9	2	131.5	139.7	104.0
Motor vehicle acc. (810-835)	6	6	0	5	5	0	59.8	77.6	-
Ill-defined (780-795)	3	0	3	3	0	3	35.8	-	156.0

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
GRENADA COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	540	204	336	542	207	335	29.4	22.1	37.0
Fetal deaths	34	7	27	36	8	28	66.4	38.6	83.6
Deaths from all causes	219	107	112	195	78	117	10.6	8.3	12.9
Deaths under 1 year	23	4	19	23	3	20	42.4	14.5	59.7
Deaths under 28 days	17	4	13	17	3	14	31.4	14.5	41.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	24	15	9	26	14	12	141.2	149.7	132.5
Diabetes (260)	1	0	1	1	0	1	5.4	-	11.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	61	22	39	50	9	41	271.6	96.2	452.7
Heart diseases (400-402,410-443)	66	48	18	53	36	17	287.9	384.9	187.7
Hypertension (444-447)	1	0	1	1	0	1	5.4	-	11.0
Arteriosclerosis (450)	2	0	2	2	1	1	10.9	10.7	11.0
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	6	2	4	4	0	4	21.7	-	44.2
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	2	0	2	10.9	-	22.1
Maternal deaths (640-689)	1	0	1	1	0	1	18.4	-	29.8
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	2	2	0	2	2	0	3.7	9.7	-
#Other dis. early inf. (763-776)	15	2	13	15	1	14	27.7	4.8	41.8
Violent & acc. deaths (800-999)	12	2	10	9	2	7	48.9	21.4	77.3
Acc. deaths, total (800-962)	10	2	8	7	2	5	38.0	21.4	55.2
Motor vehicle acc. (810-835)	2	0	2	2	1	1	10.9	10.7	11.0
Ill-defined (780-795)	10	4	6	9	4	5	48.9	42.8	55.2

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES HANCOCK COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	203	137	66	321	249	72	22.9	21.1	31.9
Fetal deaths	5	2	3	5	2	3	15.6	8.0	41.7
Deaths from all causes	150	132	18	158	137	21	11.2	11.6	9.3
Deaths under 1 year	8	3	5	9	4	5	28.0	16.1	69.4
Deaths under 28 days	7	3	4	8	4	4	24.9	16.1	55.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	4	3	1	28.5	25.4	44.3
Tbc., respiratory (001-008)	2	1	1	4	3	1	28.5	25.4	44.3
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	8	2	16	13	3	114.0	110.3	133.0
Diabetes (260)	4	4	0	2	2	0	14.2	17.0	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	28	27	1	28	26	2	199.4	220.6	88.7
Heart diseases (400-402,410-443)	43	42	1	45	44	1	320.5	373.4	44.3
Hypertension (444-447)	0	0	0	1	0	1	7.1	-	44.3
Arteriosclerosis (450)	3	3	0	3	3	0	21.4	25.4	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	2	0	3	3	0	21.4	25.4	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	1	0	1	1	0	1	31.2	-	138.9
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	1	1	0	2	2	0	6.2	8.0	-
#Other dis. early inf. (763-776)	4	2	2	4	2	2	12.5	8.0	27.8
Violent & acc. deaths (800-999)	20	18	2	16	15	1	114.0	127.3	44.3
Acc. deaths, total (800-962)	16	14	2	13	12	1	92.6	101.8	44.3
Motor vehicle acc. (810-835)	7	7	0	5	5	0	35.6	42.4	-
Ill-defined (780-795)	29	21	8	24	16	8	171.0	135.8	354.8

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
HARRISON COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	4,221	3,462	759	3,919	3,191	728	32.8	31.8	37.8
Fetal deaths	112	87	25	99	74	25	25.3	23.2	34.3
Deaths from all causes	991	780	211	825	633	192	6.9	6.3	10.0
Deaths under 1 year	83	62	21	79	59	20	20.2	18.5	27.5
Deaths under 28 days	64	52	12	60	49	11	15.3	15.4	15.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	3	1	2	1	1	1.7	1.0	5.2
Tbc., respiratory (001-008)	3	3	0	1	1	0	0.8	1.0	-
Syphilis (020-029)	1	0	1	1	1	0	0.8	1.0	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	1	0	1	0	1	0.8	-	5.2
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	1	0	1	1	0	1	0.8	-	5.2
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	126	104	22	125	104	21	104.6	103.8	109.0
Diabetes (260)	14	9	5	16	11	5	13.4	11.0	26.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	104	83	21	88	67	21	73.6	66.8	109.0
Heart diseases (400-402,410-443)	333	282	51	258	213	45	215.9	212.5	233.7
Hypertension (444-447)	9	5	4	4	2	2	3.3	2.0	10.4
Arteriosclerosis (450)	15	12	3	14	11	3	11.7	11.0	15.6
Influenza (480-483)	2	2	0	1	1	0	0.8	1.0	-
#Pneumonia (490-493)	39	33	6	23	17	6	19.2	17.0	31.2
#Diarrhea & enteritis (571)	1	0	1	1	0	1	0.8	-	5.2
Acute nephritis (590,591)	2	1	1	2	1	1	1.7	1.0	5.2
Chronic nephritis, etc. (592-594)	5	3	2	4	2	2	3.3	2.0	10.4
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	11	9	2	12	10	2	10.0	10.0	10.4
Birth inj., atelect. (760-762)	20	17	3	18	16	2	4.6	5.0	2.7
#Other dis. early inf. (763-776)	32	24	8	31	23	8	7.9	7.2	11.0
Violent & acc. deaths (800-999)	75	53	22	66	46	20	55.2	45.9	103.9
Acc. deaths, total (800-962)	57	43	14	51	39	12	42.7	38.9	62.3
Motor vehicle acc. (810-835)	28	21	7	24	18	6	20.1	18.0	31.2
Ill-defined (780-795)	90	57	33	82	51	31	68.6	50.9	161.0

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES HINDS COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	6,195	3,168	3,027	5,491	2,670	2,821	29.4	23.8	37.7
Fetal deaths	414	262	152	340	221	119	61.9	82.8	42.2
Deaths from all causes	2,165	1,167	998	1,441	711	730	7.7	6.3	9.8
Deaths under 1 year	259	95	164	174	51	123	31.7	19.1	43.6
Deaths under 28 days	174	67	107	123	39	84	22.4	14.6	29.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	9	3	6	10	3	7	5.3	2.7	9.4
Tbc., respiratory (001-008)	7	3	4	8	3	5	4.3	2.7	6.7
Syphilis (020-029)	5	0	5	4	0	4	2.1	-	5.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050-051)	1	0	1	1	0	1	0.5	-	1.3
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	2	2	0	2	2	0	1.1	1.8	-
Acute poliomyelitis (080)	1	1	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	2	2	0	2	2	0	1.1	1.8	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	359	225	134	208	127	81	111.2	113.2	108.2
Diabetes (260)	31	14	17	19	7	12	10.2	6.2	16.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	220	118	102	155	80	75	82.9	71.3	100.2
Heart diseases (400-402,410-443)	536	328	208	385	225	160	205.8	200.5	213.8
Hypertension (444-447)	17	5	12	13	4	9	7.0	3.6	12.0
Arteriosclerosis (450)	34	25	9	29	20	9	15.5	17.8	12.0
Influenza (480-483)	1	1	0	1	0	1	0.5	-	1.3
#Pneumonia (490-493)	54	32	22	29	13	16	15.5	11.6	21.4
#Diarrhea & enteritis (571)	9	3	6	6	2	4	3.2	1.8	5.3
Acute nephritis (590,591)	6	1	5	5	0	5	2.7	-	6.7
Chronic nephritis, etc. (592-594)	25	9	16	13	5	8	7.0	4.4	10.7
Maternal deaths (640-689)	2	0	2	4	1	3	7.3	3.7	10.6
Cong. malformations (750-759)	59	36	23	21	11	10	11.2	9.8	13.4
Birth inj., atelect. (760-762)	39	20	19	29	11	18	5.3	4.1	6.4
#Other dis. early inf. (763-776)	94	25	69	69	17	52	12.6	6.4	18.4
Violent & acc. deaths (800-999)	175	92	83	130	62	68	69.5	55.2	90.9
Acc. deaths, total (800-962)	135	72	63	91	44	47	48.6	39.2	62.8
Motor vehicle acc. (810-835)	52	36	16	28	16	12	15.0	14.2	16.0
Ill-defined (780-795)	200	63	137	176	51	125	94.1	45.4	167.0

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
HOLMES COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	777	132	645	754	115	639	27.8	15.1	32.8
Fetal deaths	39	7	32	42	7	35	55.7	60.9	54.8
Deaths from all causes	329	76	253	363	95	268	13.4	12.5	13.7
Deaths under 1 year	42	2	40	43	2	41	57.0	17.4	64.2
Deaths under 28 days	29	2	27	30	2	28	39.8	17.4	43.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	3.7	-	5.1
Tbc., respiratory (001-008)	1	0	1	1	0	1	3.7	-	5.1
Syphilis (020-029)	2	0	2	2	0	2	7.4	-	10.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	3.7	-	5.1
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	30	11	19	38	15	23	140.2	197.5	117.9
Diabetes (260)	2	1	1	4	2	2	14.8	26.3	10.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	44	6	38	42	7	35	155.0	92.2	179.5
Heart diseases (400-402,410-443)	101	36	65	109	40	69	402.3	526.7	353.8
Hypertension (444-447)	7	2	5	10	2	8	36.9	26.3	41.0
Arteriosclerosis (450)	3	1	2	3	1	2	11.1	13.2	10.2
Influenza (480-483)	0	0	0	1	0	1	3.7	-	5.1
#Pneumonia (490-493)	20	1	19	24	5	19	88.6	65.8	97.4
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	5	0	5	7	0	7	25.8	-	35.9
Maternal deaths (640-689)	1	0	1	1	0	1	13.3	-	15.6
Cong. malformations (750-759)	1	0	1	2	0	2	7.4	-	10.2
Birth inj., atelect. (760-762)	2	0	2	2	0	2	2.6	-	3.1
#Other dis. early inf. (763-776)	18	1	17	19	1	18	25.2	8.7	28.2
Violent & acc. deaths (800-999)	30	9	21	26	8	18	96.0	105.3	92.3
Acc. deaths, total (800-962)	19	4	15	17	4	13	62.7	52.7	66.7
Motor vehicle acc. (810-835)	6	1	5	6	1	5	22.1	13.2	25.6
Ill-defined (780-795)	53	5	48	51	5	46	188.2	65.8	235.9

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
HUMPHREYS COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	599	86	513	612	104	508	32.0	18.1	38.1
Fetal deaths	17	0	17	17	1	16	27.8	9.6	31.5
Deaths from all causes	172	45	127	208	55	153	10.9	9.6	11.5
Deaths under 1 year	30	2	28	36	1	35	58.8	9.6	68.9
Deaths under 28 days	12	2	10	15	1	14	24.5	9.6	27.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	5.2	-	7.5
Tbc., respiratory (001-008)	0	0	0	1	0	1	5.2	-	7.5
Syphilis (020-029)	1	0	1	1	0	1	5.2	-	7.5
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	14	6	8	24	11	13	125.7	191.0	97.5
Diabetes (260)	1	0	1	2	1	1	10.5	17.4	7.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	24	4	20	27	4	23	141.4	69.5	172.5
Heart diseases (400-402,410-443)	47	23	24	51	24	27	267.1	416.8	202.5
Hypertension (444-447)	3	2	1	3	2	1	15.7	34.7	7.5
Arteriosclerosis (450)	4	2	2	3	1	2	15.7	17.4	15.0
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	11	1	10	10	1	9	52.4	17.4	67.5
#Diarrhea & enteritis (571)	6	0	6	7	0	7	36.7	-	52.5
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	8	0	8	9	0	9	47.1	-	67.5
Maternal deaths (640-689)	2	0	2	1	0	1	16.3	-	19.7
Cong. malformations (750-759)	1	0	1	2	0	2	10.5	-	15.0
Birth inj., atelect. (760-762)	1	0	1	1	0	1	1.6	-	2.0
#Other dis. early inf. (763-776)	10	2	8	14	1	13	22.9	9.6	25.6
Violent & acc. deaths (800-999)	15	4	11	20	8	12	104.8	138.9	90.0
Acc. deaths, total (800-962)	10	2	8	15	6	9	78.6	104.2	67.5
Motor vehicle acc. (810-835)	0	0	0	3	3	0	15.7	52.1	-
Ill-defined (780-795)	16	0	16	19	0	19	99.5	-	142.5

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES ISSAQUENA COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	34	0	34	101	22	79	28.2	18.7	32.9
Fetal deaths	0	0	0	2	1	1	19.8	45.4	12.6
Deaths from all causes	24	5	19	40	13	27	11.2	11.0	11.2
Deaths under 1 year	5	0	5	8	1	7	79.2	45.4	88.6
Deaths under 28 days	1	0	1	4	1	3	39.6	45.4	38.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	1	0	1	7	3	4	195.7	255.1	166.7
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	1	0	1	3	1	2	83.9	85.0	83.3
Heart diseases (400-402,410-443)	7	3	4	10	4	6	279.6	340.1	250.0
Hypertension (444-447)	0	0	0	1	0	1	28.0	-	41.7
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	0	2	2	0	2	55.9	-	83.3
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	1	0	1	9.9	-	12.6
#Other dis. early inf. (763-776)	2	0	2	3	0	3	29.7	-	38.0
Violent & acc. deaths (800-999)	5	2	3	5	3	2	139.8	255.1	83.3
Acc. deaths, total (800-962)	5	2	3	4	2	2	111.8	170.1	83.3
Motor vehicle acc. (810-835)	0	0	0	2	2	0	55.9	170.1	-
Ill-defined (780-795)	5	0	5	6	1	5	167.8	85.0	208.3

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES ITAWAMBA COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	75	56	19	241	216	25	16.0	15.2	28.6
Fetal deaths	1	1	0	6	4	2	24.9	18.5	80.0
Deaths from all causes	86	80	6	120	113	7	8.0	8.0	8.0
Deaths under 1 year	2	1	1	3	2	1	12.4	9.2	40.0
Deaths under 28 days	2	1	1	2	1	1	8.3	4.6	40.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	6.6	7.0	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	6.6	7.0	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	21	20	1	30	28	2	198.9	197.1	228.8
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	13	2	19	17	2	126.0	119.7	228.8
Heart diseases (400-402,410-443)	30	28	2	38	36	2	252.0	253.4	228.8
Hypertension (444-447)	2	2	0	2	2	0	13.3	14.1	-
Arteriosclerosis (450)	3	3	0	3	3	0	19.9	21.1	-
Influenza (480-483)	1	1	0	1	1	0	6.6	7.0	-
#Pneumonia (490-493)	3	3	0	4	4	0	26.5	28.2	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	2	1	1	13.3	7.0	114.4
Birth inj., atelect. (760-762)	1	1	0	1	1	0	4.1	4.6	-
#Other dis. early inf. (763-776)	0	0	0	0	0	0	-	-	-
Violent & acc. deaths (800-999)	5	5	0	8	8	0	53.0	56.3	-
Acc. deaths, total (800-962)	4	4	0	7	7	0	46.4	49.3	-
Motor vehicle acc. (810-835)	0	0	0	2	2	0	13.3	14.1	-
Ill-defined (780-795)	1	1	0	2	2	0	13.3	14.1	-

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
JACKSON COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,341	937	404	1,652	1,223	429	29.8	27.4	39.5
Fetal deaths	71	52	19	78	59	19	47.2	48.2	44.3
Deaths from all causes	310	219	91	363	264	99	6.5	5.9	9.1
Deaths under 1 year	42	24	18	43	24	19	26.0	19.6	44.3
Deaths under 28 days	29	19	10	29	18	11	17.6	14.7	25.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	2	0	2	2	0	3.6	4.5	-
Tbc., respiratory (001-008)	2	2	0	2	2	0	3.6	4.5	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	1	0	1.8	2.2	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	45	29	16	57	40	17	102.7	89.6	156.5
Diabetes (260)	11	8	3	11	8	3	19.8	17.9	27.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	32	19	13	42	28	14	75.6	62.7	128.9
Heart diseases (400-402,410-443)	84	70	14	95	81	14	171.1	181.4	128.9
Hypertension (444-447)	6	4	2	7	4	3	12.6	9.0	27.6
Arteriosclerosis (450)	3	2	1	4	3	1	7.2	6.7	9.2
Influenza (480-483)	1	1	0	1	1	0	1.8	2.2	-
#Pneumonia (490-493)	6	3	3	10	7	3	18.0	15.7	27.6
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	1	1	0	1	1	0	1.8	2.2	-
Chronic nephritis, etc. (592-594)	2	1	1	2	2	0	3.6	4.5	-
Maternal deaths (640-689)	1	1	0	1	1	0	6.0	8.2	-
Cong. malformations (750-759)	7	5	2	9	7	2	16.2	15.7	18.4
Birth inj., atelect. (760-762)	11	8	3	13	9	4	7.9	7.4	9.3
#Other dis. early inf. (763-776)	14	8	6	13	7	6	7.9	5.7	14.0
Violent & acc. deaths (800-999)	26	15	11	33	22	11	59.4	49.3	101.2
Acc. deaths, total (800-962)	22	12	10	29	19	10	52.2	42.5	92.0
Motor vehicle acc. (810-835)	7	6	1	13	12	1	23.4	26.9	9.2
Ill-defined (780-795)	31	22	9	31	19	12	55.8	42.5	110.4

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES JASPER COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	85	17	68	442	142	300	26.1	16.9	35.3
Fetal deaths	3	0	3	12	2	10	27.1	14.1	33.3
Deaths from all causes	82	37	45	158	84	74	9.3	10.0	8.7
Deaths under 1 year	3	1	2	12	3	9	27.1	21.1	30.0
Deaths under 28 days	0	0	0	5	1	4	11.3	7.0	13.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	2	1	1	11.8	11.9	11.8
Tbc., respiratory (001-008)	1	1	0	2	1	1	11.8	11.9	11.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	13	5	8	24	13	11	141.9	154.7	129.3
Diabetes (260)	7	3	4	11	6	5	65.0	71.4	58.8
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	4	2	2	16	9	7	94.6	107.1	82.3
Heart diseases (400-402,410-443)	32	16	16	42	25	17	248.4	297.5	199.8
Hypertension (444-447)	0	0	0	2	0	2	11.8	-	23.5
Arteriosclerosis (450)	2	1	1	3	2	1	17.7	23.8	11.8
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	0	2	8	4	4	47.3	47.6	47.0
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	1	0	1	1	0	1	5.9	-	11.8
Chronic nephritis, etc. (592-594)	1	0	1	2	1	1	11.8	11.9	11.8
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	2	0	2	11.8	-	23.5
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	0	0	0	5	1	4	11.3	7.0	13.3
Violent & acc. deaths (800-999)	7	5	2	14	9	5	82.8	107.1	58.8
Acc. deaths, total (800-962)	6	4	2	13	8	5	76.9	95.2	58.8
Motor vehicle acc. (810-835)	1	1	0	5	2	3	29.6	23.8	35.3
Ill-defined (780-795)	10	3	7	12	4	8	71.0	47.6	94.0

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES JEFFERSON COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	103	0	103	264	33	231	26.0	13.2	30.2
Fetal deaths	7	0	7	11	1	10	41.7	30.3	43.3
Deaths from all causes	88	19	69	143	38	105	14.1	15.3	13.7
Deaths under 1 year	5	0	5	7	0	7	26.5	-	30.3
Deaths under 28 days	3	0	3	6	0	6	22.7	-	26.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	9.8	-	13.1
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	9.8	-	13.1
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	9	3	6	17	6	11	167.6	241.1	143.7
Diabetes (260)	2	0	2	2	0	2	19.7	-	26.1
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	13	4	9	24	7	17	236.6	281.2	222.1
Heart diseases (400-402,410-443)	25	7	18	38	14	24	374.7	562.5	313.6
Hypertension (444-447)	1	1	0	2	1	1	19.7	40.2	13.1
Arteriosclerosis (450)	4	1	3	5	2	3	49.3	80.4	39.2
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	0	0	0	3	2	1	29.6	80.4	13.1
#Diarrhea & enteritis (571)	0	0	0	1	0	1	9.8	-	13.1
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	5	0	5	6	0	6	59.2	-	78.4
Maternal deaths (640-689)	1	0	1	1	0	1	37.9	-	43.3
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	1	0	1	3	0	3	11.4	-	13.0
Violent & acc. deaths (800-999)	11	0	11	17	3	14	167.6	120.5	182.9
Acc. deaths, total (800-962)	11	0	11	17	3	14	167.6	120.5	182.9
Motor vehicle acc. (810-835)	1	0	1	4	1	3	39.4	40.2	39.2
Ill-defined (780-795)	9	2	7	11	2	9	108.4	80.4	117.6

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES JEFFERSON DAVIS COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	255	122	133	305	118	187	22.5	19.3	25.2
Fetal deaths	15	7	8	19	9	10	62.3	76.3	53.5
Deaths from all causes	99	54	45	113	53	60	8.3	8.6	8.1
Deaths under 1 year	7	2	5	10	3	7	32.8	25.4	37.4
Deaths under 28 days	5	2	3	8	3	5	26.2	25.4	26.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	13	4	17	13	4	125.6	212.2	54.0
Diabetes (260)	3	2	1	3	2	1	22.2	32.6	13.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	21	11	10	19	9	10	140.3	146.9	134.9
Heart diseases (400-402,410-443)	26	12	14	32	13	19	236.3	212.2	256.3
Hypertension (444-447)	0	0	0	0	0	0	-	-	-
Arteriosclerosis (450)	1	1	0	1	1	0	7.4	16.3	-
Influenza (480-483)	1	0	1	1	0	1	7.4	-	13.5
#Pneumonia (490-493)	0	0	0	0	0	0	-	-	-
#Diarrhea & enteritis (571)	2	0	2	2	0	2	14.8	-	27.0
Acute nephritis (590,591)	1	1	0	1	1	0	7.4	16.3	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	1	0	1	1	0	1	32.8	-	53.5
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	2	1	1	2	1	1	6.6	8.5	5.3
#Other dis. early inf. (763-776)	3	1	2	6	2	4	19.7	16.9	21.4
Violent & acc. deaths (800-999)	7	5	2	9	2	7	66.5	32.6	94.4
Acc. deaths, total (800-962)	6	5	1	7	2	5	51.7	32.6	67.4
Motor vehicle acc. (810-835)	4	3	1	6	2	4	44.3	32.6	54.0
Ill-defined (780-795)	4	2	2	4	2	2	29.5	32.6	27.0

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
JONES COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	2,593	1,266	1,327	1,444	926	518	24.2	21.0	33.5
Fetal deaths	82	31	51	43	21	22	29.8	22.7	42.5
Deaths from all causes	704	422	282	548	378	170	9.2	8.6	11.0
Deaths under 1 year	92	29	63	54	22	32	37.4	23.8	61.8
Deaths under 28 days	59	24	35	32	15	17	22.2	16.2	32.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	2	2	0	3.4	4.5	-
Tbc., respiratory (001-008)	2	1	1	2	2	0	3.4	4.5	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	2	0	2	1	0	1	1.7	-	6.5
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	1	0	1	1	0	1	1.7	-	6.5
Whooping cough (056)	1	0	1	1	0	1	1.7	-	6.5
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	1	0	1	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	1.7	-	6.5
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	84	63	21	79	65	14	132.7	147.4	90.6
Diabetes (260)	14	9	5	13	9	4	21.8	20.4	25.9
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	113	68	45	81	53	28	136.0	120.2	181.3
Heart diseases (400-402,410-443)	169	129	40	147	119	28	246.9	269.9	181.3
Hypertension (444-447)	11	2	9	4	2	2	6.7	4.5	12.9
Arteriosclerosis (450)	2	2	0	3	3	0	5.0	6.8	-
Influenza (480-483)	1	0	1	1	0	1	1.7	-	6.5
#Pneumonia (490-493)	41	21	20	28	15	13	47.0	34.0	84.2
#Diarrhea & enteritis (571)	5	0	5	2	0	2	3.4	-	12.9
Acute nephritis (590,591)	7	2	5	6	2	4	10.1	4.5	25.9
Chronic nephritis, etc. (592-594)	13	3	10	10	2	8	16.8	4.5	51.8
Maternal deaths (640-689)	3	1	2	1	1	0	6.9	10.8	-
Cong. malformations (750-759)	11	7	4	8	6	2	13.4	13.6	12.9
Birth inj., atelect. (760-762)	9	5	4	6	5	1	4.2	5.4	1.9
#Other dis. early inf. (763-776)	47	17	30	23	9	14	15.9	9.7	27.0
Violent & acc. deaths (800-999)	62	36	26	53	34	19	89.0	77.1	123.0
Acc. deaths, total (800-962)	48	30	18	39	28	11	65.5	63.5	71.2
Motor vehicle acc. (810-835)	25	17	8	22	19	3	36.9	43.1	19.4
Ill-defined (780-795)	41	22	19	34	20	14	57.1	45.4	90.6

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
KEMPER COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	198	34	164	274	61	213	22.3	12.6	28.6
Fetal deaths	6	4	2	7	3	4	25.5	49.2	18.8
Deaths from all causes	117	59	58	145	75	70	11.8	15.5	9.4
Deaths under 1 year	11	2	9	13	2	11	47.4	32.8	51.6
Deaths under 28 days	7	2	5	9	2	7	32.8	32.8	32.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	7	3	14	10	4	114.0	207.1	53.7
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	5	10	23	9	14	187.3	186.4	187.9
Heart diseases (400-402,410-443)	42	30	12	51	37	14	415.4	766.4	187.9
Hypertension (444-447)	0	0	0	0	0	0	-	-	-
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	2	0	2	2	0	2	16.3	-	26.8
#Pneumonia (490-493)	4	3	1	5	4	1	40.7	82.8	13.4
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	16.3	20.7	13.4
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	5	2	3	7	2	5	25.5	32.8	23.5
Violent & acc. deaths (800-999)	12	4	8	12	4	8	97.7	82.8	107.4
Acc. deaths, total (800-962)	10	3	7	9	2	7	73.3	41.4	94.0
Motor vehicle acc. (810-835)	5	1	4	4	1	3	32.6	20.7	40.3
Ill-defined (780-795)	20	4	16	21	4	17	171.0	82.8	228.2

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
LAFAYETTE COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	521	286	235	553	302	251	25.9	21.4	34.6
Fetal deaths	18	6	12	15	5	10	27.1	16.6	39.8
Deaths from all causes	179	113	66	196	124	72	9.2	8.8	9.9
Deaths under 1 year	18	6	12	15	3	12	27.1	9.9	47.8
Deaths under 28 days	13	6	7	10	3	7	18.1	9.9	27.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	4.7	7.1	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	4.7	7.1	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	18	15	3	19	15	4	89.0	106.3	55.2
Diabetes (260)	0	0	0	1	1	0	4.7	7.1	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	22	8	14	25	10	15	117.1	70.9	207.0
Heart diseases (400-402,410-443)	76	54	22	82	59	23	384.0	418.1	317.5
Hypertension (444-447)	0	0	0	1	1	0	4.7	7.1	-
Arteriosclerosis (450)	5	4	1	4	3	1	18.7	21.3	13.8
Influenza (480-483)	1	1	0	1	1	0	4.7	7.1	-
#Pneumonia (490-493)	4	2	2	5	3	2	23.4	21.3	27.6
#Diarrhea & enteritis (571)	1	0	1	2	1	1	9.4	7.1	13.8
Acute nephritis (590,591)	2	1	1	2	1	1	9.4	7.1	13.8
Chronic nephritis, etc. (592-594)	1	0	1	4	3	1	18.7	21.3	13.8
Maternal deaths (640-689)	2	0	2	1	0	1	18.1	-	39.8
Cong. malformations (750-759)	1	1	0	1	1	0	4.7	7.1	-
Birth inj., atelect. (760-762)	4	1	3	4	1	3	7.2	3.3	12.0
#Other dis. early inf. (763-776)	9	4	5	6	1	5	10.8	3.3	19.9
Violent & acc. deaths (800-999)	16	11	5	17	10	7	79.6	70.9	96.6
Acc. deaths, total (800-962)	12	7	5	13	6	7	60.9	42.5	96.6
Motor vehicle acc. (810-835)	6	4	2	6	4	2	28.1	28.3	27.6
Ill-defined (780-795)	2	1	1	2	1	1	9.4	7.1	13.8

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES LAMAR COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	95	72	23	311	237	74	22.7	20.7	33.2
Fetal deaths	2	1	1	10	8	2	32.2	33.8	27.0
Deaths from all causes	77	64	13	126	103	23	9.2	9.0	10.3
Deaths under 1 year	3	2	1	6	3	3	19.3	12.6	40.5
Deaths under 28 days	2	1	1	4	1	3	12.9	4.2	40.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	12	9	3	16	13	3	117.0	113.6	134.4
Diabetes (260)	3	1	2	3	1	2	21.9	8.7	89.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	12	8	4	23	17	6	168.2	148.6	268.8
Heart diseases (400-402,410-443)	32	31	1	53	48	5	387.6	419.5	224.0
Hypertension (444-447)	0	0	0	2	1	1	14.6	8.7	44.8
Arteriosclerosis (450)	2	2	0	2	2	0	14.6	17.5	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	0	0	0	2	2	0	14.6	17.5	-
#Diarrhea & enteritis (571)	0	0	0	1	1	0	7.3	8.7	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	2	0	2	2	0	14.6	17.5	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	2	2	0	14.6	17.5	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	2	1	1	4	1	3	12.9	4.2	40.5
Violent & acc. deaths (800-999)	5	3	2	6	4	2	43.9	35.0	89.6
Acc. deaths, total (800-962)	4	3	1	5	4	1	36.6	35.0	44.8
Motor vehicle acc. (810-835)	1	1	0	0	0	0	-	-	-
Ill-defined (780-795)	2	2	0	4	3	1	29.2	26.2	44.8

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES LAUDERDALE COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	2,130	1,064	1,066	1,721	888	833	25.6	20.4	35.5
Fetal deaths	85	25	60	62	21	41	36.0	23.6	49.2
Deaths from all causes	946	574	372	749	434	315	11.2	9.9	13.4
Deaths under 1 year	85	26	59	65	17	48	37.8	19.1	57.6
Deaths under 28 days	61	23	38	47	16	31	27.3	18.0	37.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	4	1	3	6.0	2.3	12.8
Tbc., respiratory (001-008)	2	1	1	4	1	3	6.0	2.3	12.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	103	69	34	85	58	27	126.6	132.9	115.0
Diabetes (260)	18	9	9	15	9	6	22.3	20.6	25.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	158	94	64	117	62	55	174.3	142.1	234.2
Heart diseases (400-402,410-443)	285	210	75	231	160	71	344.2	366.7	302.3
Hypertension (444-447)	5	1	4	5	0	5	7.4	-	21.3
Arteriosclerosis (450)	20	18	2	13	11	2	19.4	25.2	8.5
Influenza (480-483)	1	1	0	1	1	0	1.5	2.3	-
#Pneumonia (490-493)	30	13	17	22	9	13	32.8	20.6	55.4
#Diarrhea & enteritis (571)	4	2	2	3	1	2	4.5	2.3	8.5
Acute nephritis (590,591)	2	1	1	1	0	1	1.5	-	4.2
Chronic nephritis, etc. (592-594)	5	0	5	5	1	4	7.4	2.3	17.0
Maternal deaths (640-689)	3	2	1	3	2	1	17.4	22.5	12.0
Cong. malformations (750-759)	4	3	1	6	2	4	8.9	4.6	17.0
Birth inj., atelect. (760-762)	13	8	5	11	7	4	6.4	7.9	4.8
#Other dis. early inf. (763-776)	40	12	28	28	8	20	16.3	9.0	24.0
Violent & acc. deaths (800-999)	75	45	30	46	25	21	68.5	57.3	89.4
Acc. deaths, total (800-962)	59	36	23	33	16	17	49.2	36.7	72.4
Motor vehicle acc. (810-835)	30	21	9	15	10	5	22.3	22.9	21.3
Ill-defined (780-795)	91	26	65	77	21	56	114.7	48.1	238.5

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES LAWRENCE COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	160	53	107	228	120	108	22.3	18.9	28.0
Fetal deaths	4	1	3	10	5	5	43.8	41.7	46.3
Deaths from all causes	86	49	37	123	84	39	12.0	13.2	10.1
Deaths under 1 year	6	1	5	7	3	4	30.7	25.0	37.0
Deaths under 28 days	2	0	2	3	2	1	13.2	16.7	9.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	16	10	6	18	13	5	176.2	204.6	129.5
Diabetes (260)	2	1	1	2	1	1	19.6	15.7	25.9
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	9	6	25	18	7	244.7	283.3	181.3
Heart diseases (400-402,410-443)	25	19	6	36	29	7	352.4	456.4	181.3
Hypertension (444-447)	0	0	0	0	0	0	-	-	-
Arteriosclerosis (450)	4	0	4	3	0	3	29.4	-	77.7
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	4	1	3	5	1	4	48.9	15.7	103.6
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	3	0	3	29.4	-	77.7
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	1	1	0	4.4	8.3	-
#Other dis. early inf. (763-776)	2	0	2	3	1	2	13.2	8.3	18.5
Violent & acc. deaths (800-999)	8	7	1	16	15	1	156.6	236.1	25.9
Acc. deaths, total (800-962)	6	5	1	13	12	1	127.3	188.8	25.9
Motor vehicle acc. (810-835)	2	1	1	3	2	1	29.4	31.5	25.9
Ill-defined (780-795)	5	1	4	4	1	3	39.2	15.7	77.7

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
LEAKE COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	314	136	178	396	169	227	21.2	16.0	28.0
Fetal deaths	9	3	6	12	3	9	30.3	17.8	39.6
Deaths from all causes	130	89	41	162	106	56	8.7	10.0	6.9
Deaths under 1 year	8	4	4	15	6	9	37.9	35.5	39.6
Deaths under 28 days	4	2	2	9	4	5	22.7	23.7	22.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	14	10	4	20	13	7	107.2	123.1	86.4
Diabetes (260)	2	0	2	3	1	2	16.1	9.5	24.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	22	17	5	23	19	4	123.2	179.9	49.4
Heart diseases (400-402,410-443)	57	43	14	67	49	18	359.0	464.0	222.2
Hypertension (444-447)	1	1	0	3	1	2	16.1	9.5	24.7
Arteriosclerosis (450)	3	1	2	4	1	3	21.4	9.5	37.0
Influenza (480-483)	1	0	1	0	0	0	-	-	-
#Pneumonia (490-493)	3	2	1	4	2	2	21.4	18.9	24.7
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	2	2	0	10.7	18.9	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	0	1	5.4	-	12.3
Birth inj., atelect. (760-762)	2	1	1	5	2	3	12.6	11.8	13.2
#Other dis. early inf. (763-776)	2	1	1	3	1	2	7.6	5.9	8.8
Violent & acc. deaths (800-999)	18	11	7	20	12	8	107.2	113.6	98.8
Acc. deaths, total (800-962)	15	10	5	17	11	6	91.1	104.2	74.1
Motor vehicle acc. (810-835)	7	6	1	7	6	1	37.5	56.8	12.3
Ill-defined (780-795)	1	1	0	3	3	0	16.1	28.4	-

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES LEE COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,304	948	356	1,055	718	337	26.0	23.7	32.8
Fetal deaths	51	35	16	46	33	13	43.6	46.0	38.6
Deaths from all causes	393	290	103	380	277	103	9.4	9.1	10.0
Deaths under 1 year	46	30	16	42	27	15	39.8	37.6	44.5
Deaths under 28 days	32	24	8	28	20	8	26.5	27.8	23.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	2	1	3	1	2	7.4	3.3	19.4
Tbc., respiratory (001-008)	3	2	1	3	1	2	7.4	3.3	19.4
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	56	45	11	54	44	10	133.0	145.2	97.2
Diabetes (260)	4	2	2	6	4	2	14.8	13.2	19.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	43	33	10	51	40	11	125.6	132.0	106.9
Heart diseases (400-402,410-443)	150	114	36	139	103	36	342.4	339.9	349.9
Hypertension (444-447)	1	0	1	0	0	0	-	-	-
Arteriosclerosis (450)	5	4	1	5	4	1	12.3	13.2	9.7
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	16	9	7	17	10	7	41.9	33.0	68.0
#Diarrhea & enteritis (571)	2	1	1	2	1	1	4.9	3.3	9.7
Acute nephritis (590,591)	1	1	0	1	1	0	2.5	3.3	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	1	0	1	1	0	1	9.5	-	29.7
Cong. malformations (750-759)	5	5	0	5	5	0	12.3	16.5	-
Birth inj., atelect. (760-762)	12	9	3	10	7	3	9.5	9.7	8.9
#Other dis. early inf. (763-776)	16	11	5	14	9	5	13.3	12.5	14.8
Violent & acc. deaths (800-999)	39	27	12	35	24	11	86.2	79.2	106.9
Acc. deaths, total (800-962)	31	22	9	29	20	9	71.4	66.0	87.5
Motor vehicle acc. (810-835)	18	16	2	15	13	2	37.0	42.9	19.4
Ill-defined (780-795)	4	1	3	4	1	3	9.8	3.3	29.2

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
LEFLORE COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,570	452	1,118	1,533	403	1,130	32.5	24.1	37.1
Fetal deaths	106	34	72	95	26	69	62.0	64.5	61.1
Deaths from all causes	606	235	371	542	176	366	11.5	10.5	12.0
Deaths under 1 year	82	10	72	81	7	74	52.8	17.4	65.5
Deaths under 28 days	47	8	39	43	4	39	28.0	9.9	34.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	0	3	3	0	3	6.4	-	9.8
Tbc., respiratory (001-008)	3	0	3	3	0	3	6.4	-	9.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	60	27	33	53	23	30	112.4	137.7	98.5
Diabetes (260)	7	1	6	8	1	7	17.0	6.0	23.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	100	50	50	72	28	44	152.7	167.7	144.5
Heart diseases (400-402,410-443)	158	76	82	146	60	86	309.7	359.3	282.5
Hypertension (444-447)	10	3	7	7	3	4	14.8	18.0	13.1
Arteriosclerosis (450)	4	3	1	1	0	1	2.1	-	3.3
Influenza (480-483)	2	0	2	3	0	3	6.4	-	9.8
#Pneumonia (490-493)	25	7	18	23	5	18	48.8	29.9	59.1
#Diarrhea & enteritis (571)	8	0	8	9	0	9	19.1	-	29.6
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	6	1	5	6	1	5	12.7	6.0	16.4
Maternal deaths (640-689)	3	1	2	2	1	1	13.0	24.8	8.8
Cong. malformations (750-759)	1	1	0	1	1	0	2.1	6.0	-
Birth inj., atelect. (760-762)	5	2	3	4	1	3	2.6	2.5	2.6
#Other dis. early inf. (763-776)	39	5	34	36	2	34	23.5	5.0	30.1
Violent & acc. deaths (800-999)	53	28	25	55	28	27	116.7	167.7	88.7
Acc. deaths, total (800-962)	46	23	23	49	23	26	103.9	137.7	85.4
Motor vehicle acc. (810-835)	13	8	5	14	8	6	29.7	47.9	19.7
Ill-defined (780-795)	73	14	59	70	11	59	148.5	65.9	193.8

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES LINCOLN COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	608	368	240	625	352	273	23.4	19.1	32.7
Fetal deaths	21	7	14	24	4	20	38.4	11.4	73.3
Deaths from all causes	252	160	92	275	166	109	10.3	9.0	13.0
Deaths under 1 year	17	7	10	24	9	15	38.4	25.6	54.9
Deaths under 28 days	10	6	4	15	8	7	24.0	22.7	25.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	3.7	5.4	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	3.7	5.4	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	32	26	6	33	25	8	123.3	135.8	95.8
Diabetes (260)	3	2	1	4	2	2	14.9	10.9	23.9
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	31	19	12	36	20	16	134.5	108.6	191.6
Heart diseases (400-402,410-443)	106	72	34	110	73	37	411.1	396.6	443.0
Hypertension (444-447)	1	1	0	2	1	1	7.5	5.4	12.0
Arteriosclerosis (450)	4	4	0	4	4	0	14.9	21.7	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	6	3	3	7	4	3	26.2	21.7	35.9
#Diarrhea & enteritis (571)	3	1	2	4	1	3	14.9	5.4	35.9
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	5	3	2	5	3	2	18.7	16.3	23.9
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	3	3	0	5	5	0	18.7	27.2	-
Birth inj., atelect. (760-762)	1	1	0	2	2	0	3.2	5.7	-
#Other dis. early inf. (763-776)	7	3	4	10	2	8	16.0	5.7	29.3
Violent & acc. deaths (800-999)	25	12	13	23	12	11	86.0	65.2	131.7
Acc. deaths, total (800-962)	22	12	10	18	10	8	67.3	54.3	95.8
Motor vehicle acc. (810-835)	12	10	2	8	7	1	29.9	38.0	12.0
Ill-defined (780-795)	13	3	10	12	2	10	44.8	10.9	119.7

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES LOWNDES COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,492	860	632	1,411	773	638	30.2	26.8	35.9
Fetal deaths	54	25	29	49	22	27	34.7	28.5	42.3
Deaths from all causes	404	197	207	430	215	215	9.2	7.4	12.1
Deaths under 1 year	51	24	27	51	24	27	36.1	31.0	42.3
Deaths under 28 days	34	20	14	33	20	13	23.4	25.9	20.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	4	0	4	8.6	-	22.5
Tbc., respiratory (001-008)	2	0	2	4	0	4	8.6	-	22.5
Syphilis (020-029)	0	0	0	1	0	1	2.1	-	5.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	45	28	17	49	33	16	105.1	114.3	90.0
Diabetes (260)	1	1	0	1	1	0	2.1	3.5	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	48	19	29	57	25	32	122.2	86.6	180.1
Heart diseases (400-402,410-443)	112	60	52	113	62	51	242.3	214.7	287.0
Hypertension (444-447)	9	4	5	8	3	5	17.2	10.4	28.1
Arteriosclerosis (450)	6	5	1	6	5	1	12.9	17.3	5.6
Influenza (480-483)	6	0	6	6	0	6	12.9	-	33.8
#Pneumonia (490-493)	14	5	9	15	4	11	32.2	13.8	61.9
#Diarrhea & enteritis (571)	2	0	2	2	0	2	4.3	-	11.2
Acute nephritis (590,591)	1	0	1	1	0	1	2.1	-	5.6
Chronic nephritis, etc. (592-594)	2	1	1	3	2	1	6.4	6.9	5.6
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	3	1	2	4	2	2	8.6	6.9	11.2
Birth inj., atelect. (760-762)	11	10	1	11	10	1	7.8	12.9	1.6
#Other dis. early inf. (763-776)	16	9	7	17	10	7	12.0	12.9	11.0
Violent & acc. deaths (800-999)	45	28	17	44	29	15	94.3	100.4	84.4
Acc. deaths, total (800-962)	31	19	12	30	19	11	64.3	65.8	61.9
Motor vehicle acc. (810-835)	12	10	2	10	8	2	21.4	27.7	11.2
Ill-defined (780-795)	50	9	41	47	9	38	100.8	31.2	213.9

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES MADISON COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	966	155	811	1,010	180	830	30.7	19.4	35.1
Fetal deaths	55	12	43	59	15	44	58.4	83.3	53.0
Deaths from all causes	304	84	220	327	89	238	9.9	9.6	10.1
Deaths under 1 year	33	3	30	43	7	36	42.6	38.9	43.4
Deaths under 28 days	15	1	14	22	5	17	21.8	27.8	20.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	3.0	10.8	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	3.0	10.8	-
Syphilis (020-029)	1	0	1	1	0	1	3.0	-	4.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	31	11	20	36	16	20	109.4	172.6	84.6
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	36	11	25	39	12	27	118.5	129.5	114.2
Heart diseases (400-402,410-443)	88	36	52	93	35	58	282.6	377.7	245.4
Hypertension (444-447)	6	1	5	4	0	4	12.2	-	16.9
Arteriosclerosis (450)	1	0	1	1	0	1	3.0	-	4.2
Influenza (480-483)	1	0	1	1	0	1	3.0	-	4.2
#Pneumonia (490-493)	11	2	9	12	2	10	36.5	21.6	42.3
#Diarrhea & enteritis (571)	2	0	2	3	0	3	9.1	-	12.7
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	2	0	3	2	1	9.1	21.6	4.2
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	4	1	3	6	2	4	18.2	21.6	16.9
Birth inj., atelect. (760-762)	1	0	1	3	2	1	3.0	11.1	1.2
#Other dis. early inf. (763-776)	10	0	10	14	1	13	13.9	5.6	15.7
Violent & acc. deaths (800-999)	27	5	22	24	4	20	72.9	43.2	84.6
Acc. deaths, total (800-962)	20	4	16	16	3	13	48.6	32.4	55.0
Motor vehicle acc. (810-835)	4	1	3	4	1	3	12.2	10.8	12.7
Ill-defined (780-795)	59	3	56	60	3	57	182.3	32.4	241.1

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
MARION COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	483	275	208	564	336	228	24.2	21.8	28.9
Fetal deaths	11	7	4	14	8	6	24.8	23.8	26.3
Deaths from all causes	229	149	80	249	166	83	10.7	10.8	10.5
Deaths under 1 year	10	4	6	14	7	7	24.8	20.8	30.7
Deaths under 28 days	9	4	5	12	6	6	21.3	17.8	26.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	1	1	0	4.3	6.5	-
Tbc., respiratory (001-008)	1	1	0	1	1	0	4.3	6.5	-
Syphilis (020-029)	1	0	1	1	0	1	4.3	-	12.7
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	25	17	8	32	22	10	137.4	142.8	126.8
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	41	34	7	37	28	9	158.8	181.7	114.1
Heart diseases (400-402,410-443)	72	47	25	79	51	28	339.2	331.0	355.1
Hypertension (444-447)	2	0	2	2	0	2	8.6	-	25.4
Arteriosclerosis (450)	4	2	2	4	2	2	17.2	13.0	25.4
Influenza (480-483)	1	0	1	1	0	1	4.3	-	12.7
#Pneumonia (490-493)	4	3	1	5	4	1	21.5	26.0	12.7
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	4.3	-	12.7
Maternal deaths (640-689)	2	1	1	3	2	1	53.2	59.5	43.8
Cong. malformations (750-759)	1	0	1	4	2	2	17.2	13.0	25.4
Birth inj., atelect. (760-762)	2	1	1	2	1	1	3.5	3.0	4.4
#Other dis. early inf. (763-776)	4	2	2	5	3	2	8.9	8.9	8.8
Violent & acc. deaths (800-999)	30	17	13	25	18	7	107.3	116.8	88.8
Acc. deaths, total (800-962)	23	13	10	18	14	4	77.3	90.9	50.7
Motor vehicle acc. (810-835)	19	10	9	12	9	3	51.5	58.4	38.0
Ill-defined (780-795)	21	12	9	22	12	10	94.4	77.9	126.8

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
MARSHALL COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	730	126	604	746	145	601	30.4	20.0	34.9
Fetal deaths	39	5	34	36	7	29	48.2	48.3	48.2
Deaths from all causes	204	55	149	232	70	162	9.5	9.6	9.4
Deaths under 1 year	33	3	30	27	3	24	36.2	20.7	39.9
Deaths under 28 days	26	3	23	22	3	19	29.5	20.7	31.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	18	10	8	23	13	10	93.9	179.0	58.0
Diabetes (260)	4	0	4	3	0	3	12.2	-	17.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	26	4	22	31	6	25	126.5	82.6	145.0
Heart diseases (400-402,410-443)	79	30	49	87	34	53	355.0	468.1	307.4
Hypertension (444-447)	2	0	2	1	0	1	4.1	-	5.8
Arteriosclerosis (450)	0	0	0	1	0	1	4.1	-	5.8
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	5	2	3	7	4	3	28.6	55.1	17.4
#Diarrhea & enteritis (571)	3	0	3	2	0	2	8.2	-	11.6
Acute nephritis (590,591)	1	0	1	1	0	1	4.1	-	5.8
Chronic nephritis, etc. (592-594)	2	0	2	3	1	2	12.2	13.8	11.6
Maternal deaths (640-689)	0	0	0	1	0	1	13.4	-	16.6
Cong. malformations (750-759)	4	2	2	5	2	3	20.4	27.5	17.4
Birth inj., atelect. (760-762)	1	0	1	0	0	0	-	-	-
#Other dis. early inf. (763-776)	19	1	18	17	1	16	22.8	6.9	26.6
Violent & acc. deaths (800-999)	15	2	13	14	3	11	57.1	41.3	63.8
Acc. deaths, total (800-962)	11	2	9	8	2	6	32.6	27.5	34.8
Motor vehicle acc. (810-835)	6	1	5	5	1	4	20.4	13.8	23.2
Ill-defined (780-795)	14	2	12	19	2	17	77.5	27.5	98.6

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES MONROE COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	748	379	369	786	422	364	23.1	19.2	30.3
Fetal deaths	22	12	10	24	14	10	30.5	33.2	27.5
Deaths from all causes	271	149	122	296	165	131	8.7	7.5	10.9
Deaths under 1 year	31	10	21	32	10	22	40.7	23.7	60.4
Deaths under 28 days	18	10	8	19	10	9	24.2	23.7	24.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	3	2	1	8.8	9.1	8.3
Tbc., respiratory (001-008)	2	1	1	3	2	1	8.8	9.1	8.3
Syphilis (020-029)	1	1	0	1	1	0	2.9	4.6	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	39	28	11	49	33	16	144.3	150.5	133.1
Diabetes (260)	5	4	1	6	4	2	17.7	18.2	16.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	41	21	20	41	20	21	120.8	91.2	174.7
Heart diseases (400-402,410-443)	69	48	21	75	55	20	220.9	250.8	166.4
Hypertension (444-447)	0	0	0	1	1	0	2.9	4.6	-
Arteriosclerosis (450)	5	4	1	5	4	1	14.7	18.2	8.3
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	6	1	5	7	2	5	20.6	9.1	41.6
#Diarrhea & enteritis (571)	6	0	6	6	0	6	17.7	-	49.9
Acute nephritis (590,591)	1	0	1	1	0	1	2.9	-	8.3
Chronic nephritis, etc. (592-594)	13	3	10	14	4	10	41.2	18.2	83.2
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	2	0	2	2	0	5.9	9.1	-
Birth inj., atelect. (760-762)	8	5	3	8	5	3	10.2	11.8	8.2
#Other dis. early inf. (763-776)	9	3	6	10	3	7	12.7	7.1	19.2
Violent & acc. deaths (800-999)	34	14	20	32	12	20	94.2	54.7	166.4
Acc. deaths, total (800-962)	29	14	15	25	11	14	73.6	50.2	116.5
Motor vehicle acc. (810-835)	13	7	6	12	7	5	35.3	31.9	41.6
Ill-defined (780-795)	10	2	8	11	3	8	32.4	13.7	66.6

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES MONTGOMERY COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place or Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	359	146	213	349	132	217	26.2	18.0	36.3
Fetal deaths	12	3	9	12	4	8	34.4	30.3	36.9
Deaths from all causes	163	90	73	176	101	75	13.2	13.7	12.6
Deaths under 1 year	16	3	13	15	3	12	43.0	22.7	55.3
Deaths under 28 days	9	2	7	7	2	5	20.0	15.2	23.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	26	17	9	29	20	9	217.7	272.1	150.7
Diabetes (260)	2	1	1	3	2	1	22.5	27.2	16.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	35	16	19	36	18	18	270.3	244.9	301.4
Heart diseases (400-402,410-443)	39	27	12	49	36	13	367.9	489.9	217.7
Hypertension (444-447)	3	1	2	3	1	2	22.5	13.6	33.5
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	6	2	4	9	3	6	67.6	40.8	100.5
#Diarrhea & enteritis (571)	1	0	1	1	0	1	7.5	-	16.7
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	7.5	-	16.7
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	2	0	2	2	0	15.0	27.2	-
Birth inj., atelect. (760-762)	3	1	2	3	1	2	8.6	7.6	9.2
#Other dis. early inf. (763-776)	5	1	4	3	1	2	8.6	7.6	9.2
Violent & acc. deaths (800-999)	17	9	8	14	6	8	105.1	81.6	134.0
Acc. deaths, total (800-962)	11	7	4	8	4	4	60.1	54.4	67.0
Motor vehicle acc. (810-835)	4	4	0	3	2	1	22.5	27.2	16.7
Ill-defined (780-795)	9	2	7	11	2	9	82.6	27.2	150.7

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
NESHOBA COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	434	238	196	439	250	189	21.0	16.6	32.0
Fetal deaths	12	8	4	11	7	4	25.0	28.0	21.2
Deaths from all causes	144	94	50	183	134	49	8.7	8.9	8.3
Deaths under 1 year	10	3	7	14	7	7	31.9	28.0	37.0
Deaths under 28 days	5	2	3	9	6	3	20.5	24.0	15.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	4.8	-	16.9
Tbc., respiratory (001-008)	1	0	1	1	0	1	4.8	-	16.9
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	21	16	5	29	23	6	138.6	153.1	101.7
Diabetes (260)	2	2	0	3	3	0	14.3	20.0	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	31	21	10	31	24	7	148.1	159.7	118.6
Heart diseases (400-402,410-443)	38	28	10	48	41	7	229.4	272.9	118.6
Hypertension (444-447)	3	2	1	2	1	1	9.6	6.6	16.9
Arteriosclerosis (450)	4	3	1	3	2	1	14.3	13.3	16.9
Influenza (480-483)	0	0	0	1	0	1	4.8	-	16.9
#Pneumonia (490-493)	0	0	0	1	1	0	4.8	6.6	-
#Diarrhea & enteritis (571)	2	0	2	2	0	2	9.6	-	33.9
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	2	2	0	9.6	13.3	-
Birth inj., atelect. (760-762)	2	1	1	2	2	0	4.6	8.0	-
#Other dis. early inf. (763-776)	3	1	2	7	4	3	15.9	16.0	15.9
Violent & acc. deaths (800-999)	13	9	4	20	15	5	95.6	99.8	84.7
Acc. deaths, total (800-962)	7	5	2	14	11	3	66.9	73.2	50.8
Motor vehicle acc. (810-835)	2	1	1	5	3	2	23.9	20.0	33.9
Ill-defined (780-795)	11	3	8	15	5	10	71.7	33.3	169.5

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES NEWTON COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	404	242	162	434	221	213	22.2	17.1	32.4
Fetal deaths	35	19	16	36	16	20	82.9	72.4	93.9
Deaths from all causes	186	133	53	216	152	64	11.1	11.7	9.7
Deaths under 1 year	13	4	9	16	7	9	36.9	31.7	42.2
Deaths under 28 days	9	4	5	12	7	5	27.6	31.7	23.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	2	1	1	10.2	7.7	15.2
Tbc., respiratory (001-008)	0	0	0	2	1	1	10.2	7.7	15.2
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	19	16	3	21	18	3	107.6	139.0	45.7
Diabetes (260)	4	3	1	3	2	1	15.4	15.4	15.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	26	19	7	31	23	8	158.8	177.6	121.8
Heart diseases (400-402,410-443)	59	46	13	67	52	15	343.3	401.5	228.4
Hypertension (444-447)	3	2	1	3	3	0	15.4	23.2	-
Arteriosclerosis (450)	3	3	0	4	4	0	20.5	30.9	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	6	4	2	7	3	4	35.9	23.2	60.9
#Diarrhea & enteritis (571)	1	0	1	1	0	1	5.1	-	15.2
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	2	0	2	2	0	10.2	15.4	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	5.1	7.7	-
Birth inj., atelect. (760-762)	0	0	0	0	0	0	-	-	-
#Other dis. early inf. (763-776)	7	3	4	8	4	4	18.4	18.1	18.8
Violent & acc. deaths (800-999)	15	10	5	27	19	8	138.3	146.7	121.8
Acc. deaths, total (800-962)	14	9	5	23	16	7	117.8	123.6	106.6
Motor vehicle acc. (810-835)	7	4	3	15	10	5	76.8	77.2	76.1
Ill-defined (780-795)	22	9	13	22	7	15	112.7	54.0	228.4

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
NOXUBEE COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	516	110	406	501	106	395	29.8	22.4	32.6
Fetal deaths	30	2	28	28	1	27	55.9	9.4	68.4
Deaths from all causes	164	44	120	181	51	130	10.8	10.8	10.7
Deaths under 1 year	18	3	15	21	4	17	41.9	37.7	43.0
Deaths under 28 days	10	2	8	12	3	9	24.0	28.3	22.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	19	5	14	21	5	16	124.8	105.8	132.2
Diabetes (260)	1	0	1	2	0	2	11.9	-	16.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	21	5	16	24	6	18	142.6	127.0	148.7
Heart diseases (400-402,410-443)	46	20	26	48	21	27	285.3	444.5	223.1
Hypertension (444-447)	2	0	2	1	0	1	5.9	-	8.3
Arteriosclerosis (450)	8	4	4	7	4	3	41.6	84.7	24.8
Influenza (480-483)	1	0	1	1	0	1	5.9	-	8.3
#Pneumonia (490-493)	5	1	4	7	2	5	41.6	42.3	41.3
#Diarrhea & enteritis (571)	4	0	4	5	0	5	29.7	-	41.3
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	0	2	2	0	2	11.9	-	16.5
Maternal deaths (640-689)	1	0	1	1	0	1	20.0	-	25.3
Cong. malformations (750-759)	3	0	3	4	1	3	23.8	21.2	24.8
Birth inj., atelect. (760-762)	1	0	1	1	0	1	2.0	-	2.5
#Other dis. early inf. (763-776)	4	2	2	4	1	3	8.0	9.4	7.6
Violent & acc. deaths (800-999)	14	5	9	16	6	10	95.1	127.0	82.6
Acc. deaths, total (800-962)	11	4	7	13	5	8	77.3	105.8	66.1
Motor vehicle acc. (810-835)	4	2	2	5	3	2	29.7	63.5	16.5
Ill-defined (780-795)	20	1	19	21	1	20	124.8	21.2	165.3

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES OKTIBBEHA COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	787	340	447	745	310	435	28.5	21.0	38.0
Fetal deaths	22	6	16	25	7	18	33.6	22.6	41.4
Deaths from all causes	195	102	93	215	111	104	8.2	7.5	9.1
Deaths under 1 year	23	8	15	26	10	16	34.9	32.2	36.8
Deaths under 28 days	12	6	6	13	6	7	17.4	19.4	16.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	0	3	3	0	3	11.5	-	26.2
Tbc., respiratory (001-008)	3	0	3	3	0	3	11.5	-	26.2
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	21	14	7	27	16	11	103.2	108.6	96.1
Diabetes (260)	6	6	0	6	6	0	22.9	40.7	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	28	18	10	24	13	11	91.7	88.3	96.1
Heart diseases (400-402,410-443)	67	37	30	73	41	32	278.9	278.4	279.5
Hypertension (444-447)	1	0	1	1	0	1	3.8	-	8.7
Arteriosclerosis (450)	2	2	0	2	2	0	7.6	13.6	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	6	2	4	6	2	4	22.9	13.6	34.9
#Diarrhea & enteritis (571)	3	0	3	3	0	3	11.5	-	26.2
Acute nephritis (590,591)	0	0	0	1	1	0	3.8	6.8	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	3.8	6.8	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	5	3	2	6	4	2	22.9	27.2	17.5
Birth inj., atelect. (760-762)	3	2	1	3	2	1	4.0	6.4	2.3
#Other dis. early inf. (763-776)	6	2	4	6	2	4	8.0	6.4	9.2
Violent & acc. deaths (800-999)	18	9	9	23	13	10	87.9	88.3	87.4
Acc. deaths, total (800-962)	13	8	5	18	12	6	68.8	81.5	52.4
Motor vehicle acc. (810-835)	7	4	3	8	6	2	30.6	40.7	17.5
Ill-defined (780-795)	9	1	8	10	1	9	38.2	6.8	78.6

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
PANOLA COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	821	227	594	848	245	603	29.4	19.5	37.2
Fetal deaths	25	2	23	29	2	27	34.2	8.2	44.8
Deaths from all causes	245	100	145	293	123	170	10.2	9.8	10.5
Deaths under 1 year	36	4	32	44	8	36	51.9	32.6	59.7
Deaths under 28 days	20	3	17	27	6	21	31.8	24.5	34.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	0	2	4	1	3	13.9	8.0	18.5
Tbc., respiratory (001-008)	2	0	2	3	1	2	10.4	8.0	12.3
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	3	1	2	3	1	2	10.4	8.0	12.3
Scarlet fever & streptococcal sore throat (050,051)	1	0	1	1	0	1	3.5	-	6.2
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	25	12	13	31	14	17	107.7	111.4	104.8
Diabetes (260)	2	0	2	2	0	2	6.9	-	12.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	40	15	25	47	20	27	163.2	159.2	166.4
Heart diseases (400-402,410-443)	87	48	39	91	49	42	316.1	390.0	258.8
Hypertension (444-447)	1	0	1	1	0	1	3.5	-	6.2
Arteriosclerosis (450)	2	2	0	4	3	1	13.9	23.9	6.2
Influenza (480-483)	1	0	1	1	0	1	3.5	-	6.2
#Pneumonia (490-493)	7	0	7	9	2	7	31.2	15.9	43.1
#Diarrhea & enteritis (571)	5	1	4	5	1	4	17.4	8.0	24.6
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	3	1	2	3	1	2	10.4	8.0	12.3
Birth inj., atelect. (760-762)	1	0	1	1	0	1	1.2	-	1.6
#Other dis. early inf. (763-776)	14	2	12	20	5	15	23.6	20.4	24.9
Violent & acc. deaths (800-999)	11	6	5	16	9	7	55.6	71.6	43.1
Acc. deaths, total (800-962)	9	6	3	14	9	5	48.6	71.6	30.8
Motor vehicle acc. (810-835)	6	5	1	7	6	1	24.3	47.8	6.2
Ill-defined (780-795)	27	5	22	29	5	24	100.7	39.8	147.9

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES PEARL RIVER COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	589	447	142	514	343	171	22.9	19.9	32.9
Fetal deaths	11	10	1	7	6	1	13.6	17.5	5.8
Deaths from all causes	190	147	43	213	163	50	9.5	9.5	9.6
Deaths under 1 year	15	6	9	19	10	9	37.0	29.2	52.6
Deaths under 28 days	11	6	5	15	10	5	29.2	29.2	29.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	4.5	5.8	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	4.5	5.8	-
Syphilis (020-029)	1	1	0	2	1	1	8.9	5.8	19.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	24	22	2	27	25	2	120.5	145.2	38.5
Diabetes (260)	3	2	1	4	3	1	17.8	17.4	19.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	21	16	5	21	16	5	93.7	92.9	96.3
Heart diseases (400-402,410-443)	79	63	16	82	66	16	365.9	383.2	308.3
Hypertension (444-447)	0	0	0	1	1	0	4.5	5.8	-
Arteriosclerosis (450)	4	4	0	4	4	0	17.8	23.2	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	1	1	4	3	1	17.8	17.4	19.3
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	3	0	2	2	0	8.9	11.6	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	3	3	0	2	2	0	8.9	11.6	-
Birth inj., atelect. (760-762)	4	2	2	6	4	2	11.7	11.7	11.7
#Other dis. early inf. (763-776)	5	2	3	7	4	3	13.6	11.7	17.5
Violent & acc. deaths (800-999)	17	12	5	26	15	11	116.0	87.1	211.9
Acc. deaths, total (800-962)	14	10	4	23	13	10	102.6	75.5	192.7
Motor vehicle acc. (810-835)	6	5	1	14	8	6	62.5	46.4	115.6
Ill-defined (780-795)	8	6	2	5	3	2	22.3	17.4	38.5

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
PERRY COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	45	39	6	211	154	57	24.1	24.3	23.6
Fetal deaths	0	0	0	8	2	6	37.9	13.0	105.3
Deaths from all causes	34	26	8	72	55	17	8.2	8.7	7.0
Deaths under 1 year	0	0	0	8	5	3	37.9	32.5	52.6
Deaths under 28 days	0	0	0	7	5	2	33.2	32.5	35.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	3	1	2	6	4	2	68.6	63.2	82.9
Diabetes (260)	1	0	1	1	0	1	11.4	-	41.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	7	7	0	9	8	1	102.9	126.3	41.4
Heart diseases (400-402,410-443)	12	10	2	24	21	3	274.4	331.6	124.4
Hypertension (444-447)	0	0	0	1	0	1	11.4	-	41.4
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	1	0	1	1	0	11.4	15.8	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	2	2	0	22.9	31.6	-
Maternal deaths (640-689)	0	0	0	1	0	1	47.4	-	175.4
Cong. malformations (750-759)	1	1	0	1	1	0	11.4	15.8	-
Birth inj., atelect. (760-762)	0	0	0	3	2	1	14.2	13.0	17.5
#Other dis. early inf. (763-776)	0	0	0	4	3	1	19.0	19.5	17.5
Violent & acc. deaths (800-999)	4	3	1	11	8	3	125.8	126.3	124.4
Acc. deaths, total (800-962)	3	2	1	10	7	3	114.4	110.5	124.4
Motor vehicle acc. (810-835)	1	1	0	4	3	1	45.7	47.4	41.4
Ill-defined (780-795)	0	0	0	0	0	0	-	-	-

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
PIKE COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	959	457	502	857	369	488	24.4	18.8	31.7
Fetal deaths	35	14	21	27	12	15	31.5	32.5	30.7
Deaths from all causes	375	212	163	401	224	177	11.4	11.4	11.5
Deaths under 1 year	31	8	23	34	12	22	39.7	32.5	45.1
Deaths under 28 days	19	8	11	22	11	11	25.7	29.8	22.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	3	1	2	4	2	2	11.4	10.2	13.0
Tbc., respiratory (001-008)	2	0	2	3	1	2	8.6	5.1	13.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	39	26	13	41	25	16	116.9	127.2	103.8
Diabetes (260)	5	1	4	4	0	4	11.4	-	26.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	59	33	26	59	33	26	168.3	167.9	168.7
Heart diseases (400-402,410-443)	124	87	37	134	95	39	382.2	483.3	253.1
Hypertension (444-447)	3	2	1	5	3	2	14.3	15.3	13.0
Arteriosclerosis (450)	11	8	3	10	7	3	28.5	35.6	19.5
Influenza (480-483)	1	1	0	1	1	0	2.8	5.1	-
#Pneumonia (490-493)	3	2	1	2	1	1	5.7	5.1	6.5
#Diarrhea & enteritis (571)	5	2	3	4	2	2	11.4	10.2	13.0
Acute nephritis (590,591)	1	0	1	2	0	2	5.7	-	13.0
Chronic nephritis, etc. (592-594)	2	1	1	1	0	1	2.8	-	6.5
Maternal deaths (640-689)	2	1	1	2	1	1	23.3	27.1	20.5
Cong. malformations (750-759)	5	1	4	5	1	4	14.3	5.1	26.0
Birth inj., atelect. (760-762)	9	3	6	9	3	6	10.5	8.1	12.3
#Other dis. early inf. (763-776)	5	4	1	8	7	1	9.3	19.0	2.0
Violent & acc. deaths (800-999)	26	10	16	30	14	16	85.6	71.2	103.8
Acc. deaths, total (800-962)	22	9	13	27	13	14	77.0	66.1	90.9
Motor vehicle acc. (810-835)	13	8	5	15	9	6	42.8	45.8	38.9
Ill-defined (780-795)	45	9	36	43	6	37	122.6	30.5	240.1

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
PONTOTOC COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	329	245	84	337	251	86	19.6	18.0	26.2
Fetal deaths	11	8	3	15	11	4	44.5	43.8	46.5
Deaths from all causes	178	151	27	192	165	27	11.1	11.8	8.2
Deaths under 1 year	6	4	2	6	5	1	17.8	19.9	11.6
Deaths under 28 days	5	4	1	4	4	0	11.9	15.9	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	5.8	7.2	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	5.8	7.2	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	23	21	2	28	26	2	162.5	186.4	60.9
Diabetes (260)	1	1	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	20	15	5	23	17	6	133.5	121.9	182.6
Heart diseases (400-402,410-443)	80	74	6	77	71	6	446.8	509.1	182.6
Hypertension (444-447)	1	1	0	1	1	0	5.8	7.2	-
Arteriosclerosis (450)	1	1	0	2	2	0	11.6	14.3	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	9	8	1	11	10	1	63.8	71.7	30.4
#Diarrhea & enteritis (571)	1	0	1	1	0	1	5.8	-	30.4
Acute nephritis (590,591)	1	0	1	1	0	1	5.8	-	30.4
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	1	0	1	1	0	1	29.7	-	116.3
Cong. malformations (750-759)	1	1	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	1	1	0	2	2	0	5.9	8.0	-
#Other dis. early inf. (763-776)	3	2	1	2	2	0	5.9	8.0	-
Violent & acc. deaths (800-999)	16	10	6	18	12	6	104.4	86.0	182.6
Acc. deaths, total (800-962)	8	5	3	9	6	3	52.2	43.0	91.3
Motor vehicle acc. (810-835)	2	2	0	1	1	0	5.8	7.2	-
Ill-defined (780-795)	5	4	1	5	4	1	29.0	28.7	30.4

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
PRENTISS COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	463	381	82	407	334	73	22.7	21.2	33.4
Fetal deaths	7	6	1	4	3	1	9.8	9.0	13.7
Deaths from all causes	178	152	26	171	144	27	9.5	9.1	12.4
Deaths under 1 year	12	8	4	12	8	4	29.5	24.0	54.8
Deaths under 28 days	10	7	3	10	7	3	24.6	21.0	41.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	1	0	1	1	0	1	5.6	-	45.7
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	29	27	2	28	26	2	156.0	164.9	91.5
Diabetes (260)	3	3	0	1	1	0	5.6	6.3	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	11	4	12	9	3	66.8	57.1	137.2
Heart diseases (400-402,410-443)	68	59	9	61	51	10	339.8	323.5	457.4
Hypertension (444-447)	1	1	0	2	1	1	11.1	6.3	45.7
Arteriosclerosis (450)	7	7	0	6	6	0	33.4	38.1	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	6	6	0	3	3	0	16.7	19.0	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	1	1	2	1	1	11.1	6.3	45.7
Birth inj., atelect. (760-762)	1	1	0	1	1	0	2.4	3.0	-
#Other dis. early inf. (763-776)	10	7	3	10	7	3	24.6	21.0	41.1
Violent & acc. deaths (800-999)	7	5	2	13	11	2	72.4	69.8	91.5
Acc. deaths, total (800-962)	7	5	2	13	11	2	72.4	69.8	91.5
Motor vehicle acc. (810-835)	4	3	1	7	6	1	39.0	38.1	45.7
Ill-defined (780-795)	14	12	2	13	11	2	72.4	69.8	91.5

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
QUITMAN COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	779	212	567	697	170	527	33.2	22.0	39.6
Fetal deaths	18	1	17	23	0	23	33.0	-	43.6
Deaths from all causes	172	42	130	218	62	156	10.4	8.0	11.7
Deaths under 1 year	37	11	26	39	10	29	56.0	58.8	55.0
Deaths under 28 days	19	7	12	20	6	14	28.7	35.3	26.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	1	1	2	1	1	9.5	13.0	7.5
Tbc., respiratory (001-008)	2	1	1	2	1	1	9.5	13.0	7.5
Syphilis (020-029)	1	0	1	2	0	2	9.5	-	15.0
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	17	5	12	24	7	17	114.2	90.7	127.8
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	23	6	17	33	12	21	157.0	155.5	157.8
Heart diseases (400-402,410-443)	47	8	39	58	14	44	275.9	181.5	330.7
Hypertension (444-447)	3	0	3	3	0	3	14.3	-	22.5
Arteriosclerosis (450)	1	1	0	1	1	0	4.8	13.0	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	7	2	5	8	3	5	38.1	38.9	37.6
#Diarrhea & enteritis (571)	8	1	7	8	1	7	38.1	13.0	52.6
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	0	3	3	0	3	14.3	-	22.5
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	4	3	1	6	4	2	28.5	51.8	15.0
Birth inj., atelect. (760-762)	1	0	1	2	0	2	2.9	-	3.8
#Other dis. early inf. (763-776)	16	5	11	16	4	12	23.0	23.5	22.8
Violent & acc. deaths (800-999)	18	7	11	23	8	15	109.4	103.7	112.7
Acc. deaths, total (800-962)	15	6	9	20	7	13	95.2	90.7	97.7
Motor vehicle acc. (810-835)	7	4	3	10	5	5	47.6	64.8	37.6
Ill-defined (780-795)	14	0	14	15	0	15	71.4	-	112.7

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES RANKIN COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	239	85	154	858	517	341	25.0	24.0	26.6
Fetal deaths	8	3	5	40	27	13	46.6	52.2	38.1
Deaths from all causes	335	178	157	238	150	88	6.9	7.0	6.9
Deaths under 1 year	9	2	7	28	16	12	32.6	30.9	35.2
Deaths under 28 days	7	2	5	16	9	7	18.6	17.4	20.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	2.9	-	7.8
Tbc., respiratory (001-008)	0	0	0	1	0	1	2.9	-	7.8
Syphilis (020-029)	2	0	2	1	0	1	2.9	-	7.8
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	7	1	6	1	0	1	2.9	-	7.8
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	36	27	9	32	28	4	93.2	130.2	31.2
Diabetes (260)	5	4	1	5	1	4	14.6	4.6	31.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	59	30	29	33	21	12	96.1	97.6	93.6
Heart diseases (400-402,410-443)	94	52	42	68	46	22	198.1	213.9	171.6
Hypertension (444-447)	3	0	3	3	2	1	8.7	9.3	7.8
Arteriosclerosis (450)	6	4	2	5	3	2	14.6	14.0	15.6
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	16	9	7	8	5	3	23.3	23.2	23.4
#Diarrhea & enteritis (571)	1	1	0	2	2	0	5.8	9.3	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	2.9	-	7.8
Maternal deaths (640-689)	1	1	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	1	1	5	4	1	14.6	18.6	7.8
Birth inj., atelect. (760-762)	1	1	0	1	1	0	1.2	1.9	-
#Other dis. early inf. (763-776)	1	0	1	9	5	4	10.5	9.7	11.7
Violent & acc. deaths (800-999)	31	18	13	22	12	10	64.1	55.8	78.0
Acc. deaths, total (800-962)	25	17	8	19	12	7	55.4	55.8	54.6
Motor vehicle acc. (810-835)	4	1	3	6	4	2	17.5	18.6	15.6
Ill-defined (780-795)	20	3	17	23	9	14	67.0	41.8	109.2

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES SCOTT COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	558	335	223	534	289	245	25.2	22.1	30.1
Fetal deaths	21	3	18	25	6	19	46.8	20.8	77.6
Deaths from all causes	169	118	51	196	133	63	9.2	10.2	7.7
Deaths under 1 year	18	9	9	21	9	12	39.3	31.1	49.0
Deaths under 28 days	10	7	3	13	8	5	24.3	27.7	20.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	21	15	6	23	14	9	108.6	107.3	110.6
Diabetes (260)	3	2	1	3	2	1	14.2	15.3	12.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	28	21	7	32	23	9	151.0	176.2	110.6
Heart diseases (400-402,410-443)	61	48	13	63	49	14	297.4	375.5	172.0
Hypertension (444-447)	3	2	1	4	3	1	18.9	23.0	12.3
Arteriosclerosis (450)	2	1	1	3	2	1	14.2	15.3	12.3
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	6	3	3	9	4	5	42.5	30.6	61.4
#Diarrhea & enteritis (571)	1	1	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	2	0	3	3	0	14.2	23.0	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	2	0	4	3	1	18.9	23.0	12.3
Birth inj., atelect. (760-762)	2	2	0	3	3	0	5.6	10.4	-
#Other dis. early inf. (763-776)	10	3	7	10	2	8	18.7	6.9	32.6
Violent & acc. deaths (800-999)	11	9	.2	12	9	3	56.6	69.0	36.9
Acc. deaths, total (800-962)	8	8	0	9	8	1	42.5	61.3	12.3
Motor vehicle acc. (810-835)	8	8	0	7	7	0	33.0	53.6	-
Ill-defined (780-795)	5	1	4	6	2	4	28.3	15.3	49.2

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
SHARKEY COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	233	29	204	356	52	304	33.2	16.0	40.6
Fetal deaths	6	0	6	9	0	9	25.3	-	29.6
Deaths from all causes	105	31	74	133	39	94	12.4	12.0	12.5
Deaths under 1 year	14	1	13	22	0	22	61.8	-	72.4
Deaths under 28 days	8	1	7	12	0	12	33.7	-	39.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	12	4	8	17	7	10	158.3	215.6	133.5
Diabetes (260)	1	1	0	2	1	1	18.6	30.8	13.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	12	3	9	12	3	9	111.8	92.4	120.1
Heart diseases (400-402,410-443)	34	12	22	40	15	25	372.5	462.0	333.7
Hypertension (444-447)	0	0	0	0	0	0	-	-	-
Arteriosclerosis (450)	1	0	1	2	1	1	18.6	30.8	13.3
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	3	0	3	3	0	3	27.9	-	40.0
#Diarrhea & enteritis (571)	0	0	0	1	0	1	9.3	-	13.3
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	5	0	5	6	0	6	55.9	-	80.1
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	2	0	2	18.6	-	26.7
Birth inj., atelect. (760-762)	3	1	2	2	0	2	5.6	-	6.6
#Other dis. early inf. (763-776)	6	0	6	12	0	12	33.7	-	39.5
Violent & acc. deaths (800-999)	18	10	8	19	9	10	176.9	277.2	133.5
Acc. deaths, total (800-962)	11	8	3	12	8	4	111.8	246.4	53.4
Motor vehicle acc. (810-835)	1	1	0	1	1	0	9.3	30.8	-
Ill-defined (780-795)	6	0	6	6	0	6	55.9	-	80.1

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES SIMPSON COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	440	231	209	503	253	250	24.6	19.1	34.7
Fetal deaths	9	5	4	15	8	7	29.8	31.6	28.0
Deaths from all causes	193	121	72	174	104	70	8.5	7.8	9.7
Deaths under 1 year	18	5	13	19	6	13	37.8	23.7	52.0
Deaths under 28 days	7	3	4	10	5	5	19.9	19.8	20.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	30	16	14	1	1	0	4.9	7.5	-
Tbc., respiratory (001-008)	30	16	14	1	1	0	4.9	7.5	-
Syphilis (020-029)	1	1	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	23	16	7	28	20	8	136.9	150.9	111.1
Diabetes (260)	0	0	0	2	1	1	9.8	7.5	13.9
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	20	16	4	21	16	5	102.7	120.7	69.4
Heart diseases (400-402,410-443)	64	43	21	57	36	21	278.7	271.6	291.7
Hypertension (444-447)	0	0	0	1	0	1	4.9	-	13.9
Arteriosclerosis (450)	4	2	2	4	2	2	19.6	15.1	27.8
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	6	2	4	6	2	4	29.3	15.1	55.6
#Diarrhea & enteritis (571)	3	0	3	2	0	2	9.8	-	27.8
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	2	0	1	1	0	4.9	7.5	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	4	3	1	3	2	1	14.7	15.1	13.9
Birth inj., atelect. (760-762)	2	2	0	3	3	0	6.0	11.8	-
#Other dis. early inf. (763-776)	5	0	5	7	1	6	13.9	4.0	24.0
Violent & acc. deaths (800-999)	13	8	5	13	7	6	63.6	52.8	83.3
Acc. deaths, total (800-962)	10	7	3	8	4	4	39.1	30.2	55.6
Motor vehicle acc. (810-835)	5	3	2	1	0	1	4.9	-	13.9
Ill-defined (780-795)	3	0	3	4	1	3	19.6	7.5	41.7

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
SMITH COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	75	42	33	330	190	140	23.1	17.2	43.1
Fetal deaths	2	0	2	8	4	4	24.2	21.0	28.6
Deaths from all causes	66	52	14	123	97	26	8.6	8.8	8.0
Deaths under 1 year	2	0	2	13	5	8	39.4	26.3	57.1
Deaths under 28 days	2	0	2	7	3	4	21.2	15.8	28.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	10	10	0	17	17	0	118.8	153.8	-
Diabetes (260)	0	0	0	1	1	0	7.0	9.0	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	7	5	2	19	16	3	132.8	144.7	92.4
Heart diseases (400-402,410-443)	31	24	7	41	32	9	286.6	289.4	277.2
Hypertension (444-447)	0	0	0	0	0	0	-	-	-
Arteriosclerosis (450)	1	1	0	3	3	0	21.0	27.1	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	0	0	0	2	1	1	14.0	9.0	30.8
#Diarrhea & enteritis (571)	0	0	0	1	0	1	7.0	-	30.8
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	2	0	4	4	0	28.0	36.2	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	4	3	1	28.0	27.1	30.8
Birth inj., atelect. (760-762)	1	0	1	3	1	2	9.1	5.3	14.3
#Other dis. early inf. (763-776)	0	0	0	4	2	2	12.1	10.5	14.3
Violent & acc. deaths (800-999)	9	6	3	14	8	6	97.9	72.4	184.8
Acc. deaths, total (800-962)	6	4	2	12	7	5	83.9	63.3	154.0
Motor vehicle acc. (810-835)	3	3	0	7	5	2	48.9	45.2	61.6
Ill-defined (780-795)	2	1	1	2	1	1	14.0	9.0	30.8

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
STONE COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	149	107	42	160	105	55	22.8	19.8	32.1
Fetal deaths	5	4	1	7	4	3	43.8	38.1	54.5
Deaths from all causes	55	41	14	61	43	18	8.7	8.1	10.5
Deaths under 1 year	3	1	2	4	2	2	25.0	19.0	36.4
Deaths under 28 days	1	1	0	2	2	0	12.5	19.0	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	1	0	1	14.2	-	58.4
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	8	5	3	8	5	3	114.1	94.3	175.3
Diabetes (260)	2	1	1	1	0	1	14.2	-	58.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	3	2	1	4	2	2	57.0	37.7	116.9
Heart diseases (400-402,410-443)	32	27	5	31	23	8	442.0	433.8	467.6
Hypertension (444-447)	0	0	0	0	0	0	-	-	-
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	0	2	2	0	2	28.5	-	116.9
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	14.2	18.9	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	1	1	0	14.2	18.9	-
Birth inj., atelect. (760-762)	1	1	0	1	1	0	6.2	9.5	-
#Other dis. early inf. (763-776)	0	0	0	1	1	0	6.2	9.5	-
Violent & acc. deaths (800-999)	2	2	0	4	4	0	57.0	75.4	-
Acc. deaths, total (800-962)	2	2	0	4	4	0	57.0	75.4	-
Motor vehicle acc. (810-835)	1	1	0	2	2	0	28.5	37.7	-
Ill-defined (780-795)	1	0	1	1	0	1	14.2	-	58.4

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES SUNFLOWER COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,287	241	1,046	1,398	271	1,127	30.6	18.4	36.3
Fetal deaths	41	5	36	46	9	37	32.9	33.2	32.8
Deaths from all causes	379	100	279	456	140	316	10.0	9.5	10.2
Deaths under 1 year	44	5	39	57	11	46	40.8	40.6	40.8
Deaths under 28 days	17	3	14	26	7	19	18.6	25.8	16.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	4	1	3	6	2	4	13.1	13.6	12.9
Tbc., respiratory (001-008)	4	1	3	6	2	4	13.1	13.6	12.9
Syphilis (020-029)	1	0	1	1	0	1	2.2	-	3.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	2.2	-	3.2
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	44	18	26	53	22	31	115.8	149.4	99.9
Diabetes (260)	2	1	1	2	1	1	4.4	6.8	3.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	61	16	45	77	23	54	168.3	156.1	174.1
Heart diseases (400-402,410-443)	97	31	66	112	38	74	244.8	258.0	238.6
Hypertension (444-447)	2	1	1	3	1	2	6.6	6.8	6.4
Arteriosclerosis (450)	7	2	5	8	3	5	17.5	20.4	16.1
Influenza (480-483)	3	1	2	2	1	1	4.4	6.8	3.2
#Pneumonia (490-493)	27	3	24	28	4	24	61.2	27.2	77.4
#Diarrhea & enteritis (571)	6	0	6	7	1	6	15.3	6.8	19.3
Acute nephritis (590,591)	2	0	2	1	0	1	2.2	-	3.2
Chronic nephritis, etc. (592-594)	5	1	4	7	3	4	15.3	20.4	12.9
Maternal deaths (640-689)	4	0	4	4	0	4	28.6	-	35.5
Cong. malformations (750-759)	2	0	2	5	1	4	10.9	6.8	12.9
Birth inj., atelect. (760-762)	7	2	5	8	1	7	5.7	3.7	6.2
#Other dis. early inf. (763-776)	10	1	9	18	6	12	12.9	22.1	10.6
Violent & acc. deaths (800-999)	34	9	25	35	11	24	76.5	74.7	77.4
Acc. deaths, total (800-962)	21	8	13	26	10	16	56.8	67.9	51.6
Motor vehicle acc. (810-835)	8	6	2	7	5	2	15.3	33.9	6.4
Ill-defined (780-795)	37	3	34	36	3	33	78.7	20.4	106.4

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES TALLAHATCHIE COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	735	125	610	782	159	623	32.5	18.5	40.2
Fetal deaths	37	11	26	40	12	28	51.2	75.5	44.9
Deaths from all causes	234	77	157	263	88	175	10.9	10.2	11.3
Deaths under 1 year	26	3	23	28	4	24	35.8	25.2	38.5
Deaths under 28 days	15	2	13	17	3	14	21.7	18.9	22.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	2	0	2	1	0	1	4.2	-	6.4
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	4.2	-	6.4
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	1	0	1	1	0	4.2	11.6	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	25	13	12	27	14	13	112.1	163.2	83.9
Diabetes (260)	2	1	1	2	1	1	8.3	11.6	6.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	18	4	14	28	11	17	116.3	128.2	109.7
Heart diseases (400-402,410-443)	71	27	44	78	31	47	323.9	361.3	303.2
Hypertension (444-447)	3	0	3	3	0	3	12.4	-	19.4
Arteriosclerosis (450)	11	3	8	11	3	8	45.7	35.0	51.6
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	7	2	5	9	4	5	37.4	46.6	32.2
#Diarrhea & enteritis (571)	2	0	2	2	0	2	8.3	-	12.9
Acute nephritis (590,591)	1	1	0	1	1	0	4.2	11.6	-
Chronic nephritis, etc. (592-594)	4	2	2	3	1	2	12.4	11.6	12.9
Maternal deaths (640-689)	0	0	0	2	0	2	25.6	-	32.1
Cong. malformations (750-759)	3	2	1	4	2	2	16.6	23.3	12.9
Birth inj., atelect. (760-762)	2	1	1	2	1	1	2.6	6.3	1.6
#Other dis. early inf. (763-776)	10	0	10	11	1	10	14.1	6.3	16.0
Violent & acc. deaths (800-999)	31	11	20	30	8	22	124.6	93.2	141.9
Acc. deaths, total (800-962)	21	10	11	20	7	13	83.0	81.6	83.9
Motor vehicle acc. (810-835)	4	3	1	3	1	2	12.4	11.6	12.9
Ill-defined (780-795)	18	1	17	20	1	19	83.0	11.6	122.6

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES TATE COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	426	23	403	590	147	443	32.5	19.1	42.4
Fetal deaths	17	1	16	24	2	22	40.7	13.6	49.7
Deaths from all causes	108	50	58	142	67	75	7.8	8.7	7.2
Deaths under 1 year	10	1	9	17	2	15	28.8	13.6	33.9
Deaths under 28 days	5	0	5	11	1	10	18.6	6.8	22.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	1	0	1	1	0	1	5.5	-	9.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	5.5	-	9.6
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	14	10	4	19	14	5	104.8	181.9	47.9
Diabetes (260)	0	0	0	1	0	1	5.5	-	9.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	8	0	8	14	2	12	77.2	26.0	114.9
Heart diseases (400-402,410-443)	53	25	28	62	31	31	341.8	402.8	296.9
Hypertension (444-447)	1	0	1	1	0	1	5.5	-	9.6
Arteriosclerosis (450)	1	1	0	1	1	0	5.5	13.0	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	0	2	4	1	3	22.0	13.0	28.7
#Diarrhea & enteritis (571)	1	0	1	1	0	1	5.5	-	9.6
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	2	0	1	1	0	5.5	13.0	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	1	0	1	1.7	-	2.2
#Other dis. early inf. (763-776)	6	0	6	9	1	8	15.2	6.8	18.0
Violent & acc. deaths (800-999)	9	9	0	8	8	0	44.1	104.0	-
Acc. deaths, total (800-962)	8	8	0	7	7	0	38.6	91.0	-
Motor vehicle acc. (810-835)	5	5	0	4	4	0	22.0	52.0	-
Ill-defined (780-795)	6	1	5	8	2	6	44.1	26.0	57.5

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
TIPPAH COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	360	269	91	318	237	81	21.1	19.2	29.4
Fetal deaths	35	23	12	31	19	12	97.5	80.2	148.1
Deaths from all causes	141	114	27	147	118	29	9.7	9.6	10.5
Deaths under 1 year	4	3	1	3	2	1	9.4	8.4	12.3
Deaths under 28 days	4	3	1	3	2	1	9.4	8.4	12.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	2	2	0	2	2	0	13.2	16.2	-
Tbc., respiratory (001-008)	2	2	0	2	2	0	13.2	16.2	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	24	20	4	30	24	6	198.8	194.5	217.7
Diabetes (260)	1	0	1	1	0	1	6.6	-	36.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	13	8	5	12	7	5	79.5	56.7	181.4
Heart diseases (400-402,410-443)	64	55	9	65	56	9	430.7	453.9	326.6
Hypertension (444-447)	0	0	0	0	0	0	-	-	-
Arteriosclerosis (450)	3	3	0	2	2	0	13.2	16.2	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	1	0	1	1	0	1	6.6	-	36.3
#Diarrhea & enteritis (571)	1	1	0	1	1	0	6.6	8.1	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	6.6	-	36.3
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	4	3	1	3	2	1	9.4	8.4	12.3
#Other dis. early inf. (763-776)	0	0	0	0	0	0	-	-	-
Violent & acc. deaths (800-999)	10	9	1	9	8	1	59.6	64.8	36.3
Acc. deaths, total (800-962)	8	7	1	7	6	1	46.4	48.6	36.3
Motor vehicle acc. (810-835)	5	4	1	4	3	1	26.5	24.3	36.3
Ill-defined (780-795)	6	2	4	7	3	4	46.4	24.3	145.1

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
TISHOMINGO COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	179	160	19	251	229	22	18.1	17.3	32.4
Fetal deaths	2	2	0	4	4	0	15.9	17.5	-
Deaths from all causes	122	114	8	151	143	8	10.9	10.8	11.8
Deaths under 1 year	5	5	0	7	7	0	27.9	30.6	-
Deaths under 28 days	3	3	0	5	5	0	19.9	21.8	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	19	18	1	22	21	1	158.4	159.0	147.3
Diabetes (260)	2	2	0	3	3	0	21.6	22.7	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	42	40	2	43	41	2	309.6	310.4	294.6
Heart diseases (400-402,410-443)	32	28	4	44	41	3	316.8	310.4	441.8
Hypertension (444-447)	2	2	0	1	1	0	7.2	7.6	-
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	2	2	0	3	3	0	21.6	22.7	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	7.2	7.6	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	2	2	0	2	2	0	14.4	15.1	-
Birth inj., atelect. (760-762)	1	1	0	2	2	0	8.0	8.7	-
#Other dis. early inf. (763-776)	1	1	0	3	3	0	12.0	13.1	-
Violent & acc. deaths (800-999)	9	9	0	11	11	0	79.2	83.3	-
Acc. deaths, total (800-962)	9	9	0	11	11	0	79.2	83.3	-
Motor vehicle acc. (810-835)	3	3	0	6	6	0	43.2	45.4	-
Ill-defined (780-795)	4	4	0	4	4	0	28.8	30.3	-

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
TUNICA COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	648	64	584	623	69	554	37.0	19.7	41.6
Fetal deaths	33	2	31	34	3	31	54.6	43.5	56.0
Deaths from all causes	201	38	163	209	39	170	12.4	11.1	12.8
Deaths under 1 year	29	3	26	27	3	24	43.3	43.5	43.3
Deaths under 28 days	17	3	14	16	3	13	25.7	43.5	23.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	14	3	11	19	3	16	112.9	85.6	120.1
Diabetes (260)	1	0	1	1	0	1	5.9	-	7.5
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	22	4	18	27	7	20	160.5	199.7	150.1
Heart diseases (400-402,410-443)	72	17	55	72	15	57	427.9	428.0	427.9
Hypertension (444-447)	2	0	2	2	0	2	11.9	-	15.0
Arteriosclerosis (450)	1	0	1	1	0	1	5.9	-	7.5
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	9	1	8	9	2	7	53.5	57.1	52.5
#Diarrhea & enteritis (571)	3	0	3	3	0	3	17.8	-	22.5
Acute nephritis (590,591)	0	0	0	1	0	1	5.9	-	7.5
Chronic nephritis, etc. (592-594)	3	0	3	3	0	3	17.8	-	22.5
Maternal deaths (640-689)	4	0	4	3	0	3	48.2	-	54.2
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	3	0	3	3	0	3	4.8	-	5.4
#Other dis. early inf. (763-776)	10	3	7	9	3	6	14.4	43.5	10.8
Violent & acc. deaths (800-999)	19	3	16	19	4	15	112.9	114.1	112.6
Acc. deaths, total (800-962)	18	3	15	18	4	14	107.0	114.1	105.1
Motor vehicle acc. (810-835)	7	2	5	8	4	4	47.5	114.1	30.0
Ill-defined (780-795)	27	4	23	25	2	23	148.6	57.1	172.6

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
UNION COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	397	294	103	388	285	103	20.5	18.3	31.1
Fetal deaths	32	31	1	26	24	2	67.0	84.2	19.4
Deaths from all causes	171	138	33	174	142	32	9.2	9.1	9.7
Deaths under 1 year	9	8	1	12	11	1	30.9	38.6	9.7
Deaths under 28 days	8	8	0	11	11	0	28.4	38.6	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	1	0	5.3	6.4	-
Tbc., respiratory (001-008)	0	0	0	1	1	0	5.3	6.4	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	23	18	5	28	23	5	148.1	147.5	151.0
Diabetes (260)	1	1	0	1	1	0	5.3	6.4	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	32	23	9	30	21	9	158.7	134.7	271.7
Heart diseases (400-402,410-443)	66	56	10	64	52	12	338.6	333.5	362.3
Hypertension (444-447)	1	0	1	0	0	0	-	-	-
Arteriosclerosis (450)	4	3	1	6	6	0	31.7	38.5	-
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	3	3	0	1	1	0	5.3	6.4	-
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	5.3	-	30.2
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	2	1	1	10.6	6.4	30.2
Birth inj., atelect. (760-762)	3	3	0	3	3	0	7.7	10.5	-
#Other dis. early inf. (763-776)	5	5	0	8	8	0	20.6	28.1	-
Violent & acc. deaths (800-999)	13	12	1	9	8	1	47.6	51.3	30.2
Acc. deaths, total (800-962)	11	10	1	7	6	1	37.0	38.5	30.2
Motor vehicle acc. (810-835)	9	8	1	5	4	1	26.4	25.6	30.2
Ill-defined (780-795)	11	7	4	12	9	3	63.5	57.7	90.6

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
WALTHALL COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	469	267	202	303	128	175	22.4	17.3	28.7
Fetal deaths	29	14	15	20	5	15	66.0	39.1	85.7
Deaths from all causes	132	76	56	123	67	56	9.1	9.0	9.2
Deaths under 1 year	17	8	9	13	5	8	42.9	39.1	45.7
Deaths under 28 days	10	6	4	6	3	3	19.8	23.4	17.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	7.4	-	16.4
Tbc., respiratory (001-008)	1	0	1	1	0	1	7.4	-	16.4
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	13	7	6	16	9	7	118.4	121.4	114.8
Diabetes (260)	0	0	0	0	0	0	-	-	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	22	11	11	21	11	10	155.4	148.4	163.9
Heart diseases (400-402,410-443)	51	37	14	48	33	15	355.2	445.2	245.9
Hypertension (444-447)	0	0	0	0	0	0	-	-	-
Arteriosclerosis (450)	4	0	4	4	0	4	29.6	-	65.6
Influenza (480-483)	0	0	0	0	0	0	-	-	-
#Pneumonia (490-493)	5	1	4	5	1	4	37.0	13.5	65.6
#Diarrhea & enteritis (571)	1	0	1	1	0	1	7.4	-	16.4
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	1	1	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	0	1	1	0	1	7.4	-	16.4
Birth inj., atelect. (760-762)	3	2	1	1	1	0	3.3	7.8	-
#Other dis. early inf. (763-776)	7	4	3	5	2	3	16.5	15.6	17.1
Violent & acc. deaths (800-999)	11	6	5	8	3	5	59.2	40.5	82.0
Acc. deaths, total (800-962)	11	6	5	7	3	4	51.8	40.5	65.6
Motor vehicle acc. (810-835)	6	3	3	5	3	2	37.0	40.5	32.8
Ill-defined (780-795)	5	4	1	5	4	1	37.0	54.0	16.4

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
WARREN COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,452	620	832	1,091	513	578	25.8	22.8	29.2
Fetal deaths	74	23	51	47	18	29	43.1	35.1	50.2
Deaths from all causes	694	250	444	505	200	305	12.0	8.9	15.4
Deaths under 1 year	94	15	79	45	12	33	41.2	23.4	57.1
Deaths under 28 days	58	10	48	27	8	19	24.7	15.6	32.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	2.4	-	5.1
Tbc., respiratory (001-008)	1	0	1	1	0	1	2.4	-	5.1
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	1	1	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	105	54	51	72	38	34	170.6	169.3	172.1
Diabetes (260)	13	2	11	7	2	5	16.6	8.9	25.3
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	86	27	59	77	25	52	182.4	111.4	263.2
Heart diseases (400-402,410-443)	172	74	98	147	64	83	348.3	285.1	420.1
Hypertension (444-447)	9	1	8	6	1	5	14.2	4.4	25.3
Arteriosclerosis (450)	12	5	7	12	6	6	28.4	26.7	30.4
Influenza (480-483)	1	0	1	0	0	0	-	-	-
#Pneumonia (490-493)	8	3	5	7	1	6	16.6	4.4	30.4
#Diarrhea & enteritis (571)	7	1	6	3	1	2	7.1	4.4	10.1
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	1	3	3	1	2	7.1	4.4	10.1
Maternal deaths (640-689)	6	0	6	4	0	4	36.7	-	69.2
Cong. malformations (750-759)	11	4	7	7	5	2	16.6	22.3	10.1
Birth inj., atelect. (760-762)	12	3	9	6	3	3	5.5	5.8	5.2
#Other dis. early inf. (763-776)	47	5	42	21	4	17	19.2	7.8	29.4
Violent & acc. deaths (800-999)	48	22	26	31	14	17	73.4	62.4	86.0
Acc. deaths, total (800-962)	39	17	22	24	10	14	56.9	44.5	70.8
Motor vehicle acc. (810-835)	13	5	8	7	2	5	16.6	8.9	25.3
Ill-defined (780-795)	65	13	52	54	9	45	127.9	40.1	227.7

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES WASHINGTON COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	2,568	948	1,620	2,460	882	1,578	31.3	25.0	36.4
Fetal deaths	189	96	93	177	82	95	72.0	93.0	60.2
Deaths from all causes	818	236	582	854	232	622	10.8	6.6	14.3
Deaths under 1 year	129	25	104	129	23	106	52.4	26.1	67.2
Deaths under 28 days	72	20	52	72	20	52	29.3	22.7	33.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	5	2	3	6	2	4	7.6	5.7	9.2
Tbc., respiratory (001-008)	3	2	1	4	2	2	5.1	5.7	4.6
Syphilis (020-029)	3	0	3	3	0	3	3.8	-	6.9
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	100	32	68	103	29	74	131.0	82.3	170.5
Diabetes (260)	7	3	4	6	2	4	7.6	5.7	9.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	85	17	68	91	18	73	115.7	51.1	168.2
Heart diseases (400-402,410-443)	202	68	134	216	73	143	274.7	207.2	329.5
Hypertension (444-447)	7	2	5	6	2	4	7.6	5.7	9.2
Arteriosclerosis (450)	7	4	3	8	3	5	10.2	8.5	11.5
Influenza (480-483)	6	1	5	6	1	5	7.6	2.8	11.5
#Pneumonia (490-493)	24	10	14	27	12	15	34.3	34.0	34.6
#Diarrhea & enteritis (571)	25	2	23	25	1	24	31.8	2.8	55.3
Acute nephritis (590,591)	3	0	3	3	0	3	3.8	-	6.9
Chronic nephritis, etc. (592-594)	13	6	7	15	5	10	19.1	14.2	23.0
Maternal deaths (640-689)	1	0	1	2	0	2	8.1	-	12.7
Cong. malformations (750-759)	10	4	6	12	4	8	15.2	11.4	18.4
Birth inj., atelect. (760-762)	12	6	6	12	6	6	4.9	6.8	3.8
#Other dis. early inf. (763-776)	50	9	41	49	9	40	19.9	10.2	25.3
Violent & acc. deaths (800-999)	79	28	51	76	27	49	96.6	76.6	112.9
Acc. deaths, total (800-962)	66	24	42	63	23	40	80.1	65.3	92.2
Motor vehicle acc. (810-835)	30	14	16	23	10	13	29.2	28.4	30.0
Ill-defined (780-795)	127	18	109	123	15	108	156.4	42.6	248.8

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
WAYNE COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	438	313	125	433	259	174	26.6	24.8	30.0
Fetal deaths	8	6	2	13	8	5	30.0	30.9	28.7
Deaths from all causes	141	85	56	162	95	67	10.0	9.1	11.5
Deaths under 1 year	18	8	10	19	8	11	43.9	30.9	63.2
Deaths under 28 days	14	6	8	14	6	8	32.3	23.2	46.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	1	0	3	2	1	18.4	19.1	17.2
Tbc., respiratory (001-008)	0	0	0	2	1	1	12.3	9.6	17.2
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	1	1	0	1	1	0	6.2	9.6	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	16	15	1	18	16	2	110.7	153.1	34.4
Diabetes (260)	5	4	1	5	4	1	30.8	38.3	17.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	12	4	8	13	5	8	80.0	47.8	137.7
Heart diseases (400-402,410-443)	55	40	15	55	42	13	338.3	402.0	223.8
Hypertension (444-447)	2	0	2	2	0	2	12.3	-	34.4
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	1	0	1	1	0	1	6.2	-	17.2
#Pneumonia (490-493)	4	3	1	5	3	2	30.8	28.7	34.4
#Diarrhea & enteritis (571)	1	0	1	1	0	1	6.2	-	17.2
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	0	3	3	0	3	18.4	-	51.6
Maternal deaths (640-689)	0	0	0	1	0	1	23.1	-	57.5
Cong. malformations (750-759)	1	0	1	2	1	1	12.3	9.6	17.2
Birth inj., atelect. (760-762)	3	0	3	3	0	3	6.9	-	17.2
#Other dis. early inf. (763-776)	9	5	4	10	5	5	23.1	19.3	28.7
Violent & acc. deaths (800-999)	7	3	4	13	5	8	80.0	47.8	137.7
Acc. deaths, total (800-962)	5	3	2	9	5	4	55.4	47.8	68.8
Motor vehicle acc. (810-835)	2	0	2	4	1	3	24.6	9.6	51.6
Ill-defined (780-795)	8	2	6	9	2	7	55.4	19.1	120.5

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
WEBSTER COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	168	90	78	204	123	81	19.3	15.5	30.6
Fetal deaths	3	0	3	4	1	3	19.6	8.1	37.0
Deaths from all causes	102	77	25	126	96	30	11.9	12.1	11.4
Deaths under 1 year	4	1	3	5	1	4	24.5	8.1	49.4
Deaths under 28 days	2	1	1	2	1	1	9.8	8.1	12.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	12	10	2	13	11	2	122.9	138.6	75.7
Diabetes (260)	0	0	0	1	1	0	9.4	12.6	-
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	30	26	4	30	27	3	283.6	340.1	113.6
Heart diseases (400-402,410-443)	34	30	4	44	38	6	415.9	478.7	227.1
Hypertension (444-447)	0	0	0	0	0	0	-	-	-
Arteriosclerosis (450)	8	5	3	7	4	3	66.2	50.4	113.6
Influenza (480-483)	1	0	1	1	0	1	9.4	-	37.8
#Pneumonia (490-493)	2	0	2	2	0	2	18.9	-	75.7
#Diarrhea & enteritis (571)	0	0	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	1	1	0	4.9	8.1	-
#Other dis. early inf. (763-776)	1	1	0	0	0	0	-	-	-
Violent & acc. deaths (800-999)	3	1	2	4	2	2	37.8	25.2	75.7
Acc. deaths, total (800-962)	2	0	2	4	2	2	37.8	25.2	75.7
Motor vehicle acc. (810-835)	0	0	0	1	1	0	9.4	12.6	-
Ill-defined (780-795)	3	0	3	8	4	4	75.6	50.4	151.4

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
WILKINSON COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	430	194	236	371	78	293	28.0	20.5	31.1
Fetal deaths	38	15	23	24	4	20	64.7	51.3	68.2
Deaths from all causes	140	52	88	145	43	102	11.0	11.3	10.8
Deaths under 1 year	14	2	12	18	0	18	48.5	-	61.4
Deaths under 28 days	6	1	5	8	0	8	21.6	-	27.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	12	4	8	11	3	8	83.1	78.8	84.8
Diabetes (260)	2	0	2	2	0	2	15.1	-	21.2
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	23	6	17	26	6	20	196.4	157.6	212.1
Heart diseases (400-402,410-443)	40	22	18	38	17	21	287.1	446.5	222.7
Hypertension (444-447)	0	0	0	0	0	0	-	-	-
Arteriosclerosis (450)	1	0	1	0	0	0	-	-	-
Influenza (480-483)	1	0	1	1	0	1	7.6	-	10.6
#Pneumonia (490-493)	7	3	4	6	1	5	45.3	26.3	53.0
#Diarrhea & enteritis (571)	2	0	2	1	0	1	7.6	-	10.6
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	7.6	-	10.6
Maternal deaths (640-689)	1	0	1	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	0	0	0	2	0	2	5.4	-	6.8
#Other dis. early inf. (763-776)	3	1	2	3	0	3	8.1	-	10.2
Violent & acc. deaths (800-999)	10	4	6	13	3	10	98.2	78.8	106.1
Acc. deaths, total (800-962)	8	3	5	10	1	9	75.6	26.3	95.5
Motor vehicle acc. (810-835)	2	1	1	3	1	2	22.7	26.3	21.2
Ill-defined (780-795)	18	2	16	19	2	17	143.6	52.5	180.3

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
WINSTON COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	404	173	231	420	174	246	21.8	16.0	29.3
Fetal deaths	12	2	10	18	6	12	42.8	34.5	48.8
Deaths from all causes	143	86	57	165	99	66	8.6	9.1	7.9
Deaths under 1 year	12	2	10	13	3	10	31.0	17.2	40.6
Deaths under 28 days	6	1	5	8	2	6	19.0	11.5	24.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	1	0	1	1	0	1	5.2	-	11.9
Tbc., respiratory (001-008)	1	0	1	1	0	1	5.2	-	11.9
Syphilis (020-029)	0	0	0	1	0	1	5.2	-	11.9
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	29	24	5	35	28	7	181.8	258.0	83.4
Diabetes (260)	5	3	2	5	2	3	26.0	18.4	35.7
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	24	15	9	28	18	10	145.5	165.8	119.1
Heart diseases (400-402,410-443)	41	21	20	43	21	22	223.4	193.5	262.1
Hypertension (444-447)	1	0	1	1	0	1	5.2	-	11.9
Arteriosclerosis (450)	2	2	0	2	2	0	10.4	18.4	-
Influenza (480-483)	2	1	1	2	1	1	10.4	9.2	11.9
#Pneumonia (490-493)	4	3	1	4	3	1	20.8	27.6	11.9
#Diarrhea & enteritis (571)	2	1	1	2	1	1	10.4	9.2	11.9
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	1	1	0	1	1	0	5.2	9.2	-
Birth inj., atelect. (760-762)	1	1	0	1	1	0	2.4	5.7	-
#Other dis. early inf. (763-776)	6	0	6	8	1	7	19.0	5.7	28.4
Violent & acc. deaths (800-999)	8	2	6	9	4	5	46.8	36.8	59.6
Acc. deaths, total (800-962)	8	2	6	9	4	5	46.8	36.8	59.6
Motor vehicle acc. (810-835)	1	1	0	2	2	0	10.4	18.4	-
Ill-defined (780-795)	5	3	2	7	4	3	36.4	36.8	35.7

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
YALOBUSHA COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	288	117	171	279	112	167	22.3	16.1	30.1
Fetal deaths	7	2	5	5	2	3	17.9	17.8	18.0
Deaths from all causes	128	73	55	159	97	62	12.7	13.9	11.2
Deaths under 1 year	6	1	5	8	3	5	28.7	26.8	29.9
Deaths under 28 days	5	1	4	7	3	4	25.1	26.8	24.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	0	0	0	-	-	-
Tbc., respiratory (001-008)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	22	13	9	31	22	9	248.0	316.0	162.4
Diabetes (260)	2	1	1	1	0	1	8.0	-	18.0
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	18	13	5	19	14	5	152.0	201.1	90.2
Heart diseases (400-402,410-443)	42	29	13	48	32	16	383.9	459.6	288.8
Hypertension (444-447)	9	4	5	8	3	5	64.0	43.1	90.2
Arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Influenza (480-483)	1	1	0	1	1	0	8.0	14.4	-
#Pneumonia (490-493)	7	3	4	9	5	4	72.0	71.8	72.2
#Diarrhea & enteritis (571)	1	1	0	0	0	0	-	-	-
Acute nephritis (590,591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	2	0	2	16.0	-	36.1
Maternal deaths (640-689)	0	0	0	0	0	0	-	-	-
Cong. malformations (750-759)	0	0	0	0	0	0	-	-	-
Birth inj., atelect. (760-762)	2	1	1	3	2	1	10.8	17.8	6.0
#Other dis. early inf. (763-776)	3	0	3	4	1	3	14.3	8.9	18.0
Violent & acc. deaths (800-999)	5	2	3	13	7	6	104.0	100.5	108.3
Acc. deaths, total (800-962)	4	1	3	11	6	5	88.0	86.2	90.2
Motor vehicle acc. (810-835)	2	1	1	6	5	1	48.0	71.8	18.0
Ill-defined (780-795)	5	0	5	5	0	5	40.0	-	90.2

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES
YAZOO COUNTY, MISSISSIPPI, 1961

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	844	237	607	900	263	637	28.4	20.4	33.9
Fetal deaths	32	7	25	42	10	32	46.7	38.0	50.2
Deaths from all causes	283	102	181	314	117	197	9.9	9.1	10.5
Deaths under 1 year	24	8	16	33	9	24	36.7	34.2	37.7
Deaths under 28 days	14	7	7	23	9	14	25.6	34.2	22.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	0	0	0	1	0	1	3.2	-	5.3
Tbc., respiratory (001-008)	0	0	0	1	0	1	3.2	-	5.3
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	3.2	-	5.3
Scarlet fever & streptococcal sore throat (050,051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	3.2	-	5.3
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	33	14	19	36	14	22	113.7	108.8	117.1
Diabetes (260)	5	3	2	6	4	2	19.0	31.1	10.6
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	63	26	37	65	29	36	205.4	225.5	191.6
Heart diseases (400-402,410-443)	57	25	32	65	32	33	205.4	248.8	175.6
Hypertension (444-447)	2	0	2	3	0	3	9.5	-	16.0
Arteriosclerosis (450)	7	3	4	6	3	3	19.0	23.3	16.0
Influenza (480-483)	1	0	1	0	0	0	-	-	-
#Pneumonia (490-493)	6	1	5	7	1	6	22.1	7.8	31.9
#Diarrhea & enteritis (571)	2	0	2	2	0	2	6.3	-	10.6
Acute nephritis (590,591)	4	0	4	4	0	4	12.6	-	21.3
Chronic nephritis, etc. (592-594)	7	1	6	5	1	4	15.8	7.8	21.3
Maternal deaths (640-689)	1	0	1	1	0	1	11.1	-	15.7
Cong. malformations (750-759)	1	0	1	4	2	2	12.6	15.5	10.6
Birth inj., atelect. (760-762)	3	1	2	4	1	3	4.4	3.8	4.7
#Other dis. early inf. (763-776)	9	6	3	13	6	7	14.4	22.8	11.0
Violent & acc. deaths (800-999)	30	11	19	31	10	21	97.9	77.7	111.8
Acc. deaths, total (800-962)	24	9	15	24	8	16	75.8	62.2	85.1
Motor vehicle acc. (810-835)	13	8	5	11	5	6	34.8	38.9	31.9
Ill-defined (780-795)	34	4	30	35	4	31	110.6	31.1	165.0

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES STATE OF MISSISSIPPI, 1961

By Place of Occurrence, by Place of Residence, by Race, with Rates

	Occurrence			Residence					
				Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	60002	27797	32205	59523	27435	32088	27.3	21.8	34.8
Fetal deaths	2562	1105	1457	2502	1055	1447	42.0	38.4	45.1
Deaths from all causes	21032	10961	10071	21232	11063	10169	9.7	8.8	11.0
Deaths under 1 year	2241	639	1602	2250	645	1605	37.8	23.5	50.0
Deaths under 28 days	1396	507	889	1401	509	892	23.5	18.6	27.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tbc., all forms (001-019)	109	51	58	113	54	59	5.2	4.3	6.4
Tbc., respiratory (001-008)	101	48	53	103	50	53	4.7	4.0	5.8
Syphilis (020-029)	29	3	26	31	4	27	1.4	0.3	2.9
Typhoid fever (040)	2	0	2	2	0	2	0.1	-	0.2
Dysentery (045-048)	17	4	13	17	4	13	0.8	0.3	1.4
Scarlet fever & streptococcal sore throat (050,051)	2	0	2	2	0	2	0.1	-	0.2
Diphtheria (055)	3	0	3	3	0	3	0.1	-	0.3
Whooping cough (056)	4	0	4	5	0	5	0.2	-	0.5
Meningococcal infections (057)	3	2	1	3	2	1	0.1	0.2	0.1
Acute poliomyelitis (080)	4	2	2	4	2	2	0.2	0.2	0.2
Smallpox (084)	0	0	0	0	0	0	-	-	-
Measles (085)	9	4	5	9	4	5	0.4	0.3	0.5
Malaria (110-117)	0	0	0	0	0	0	-	-	-
Cancer, incl. leukemia (140-205)	2639	1652	987	2748	1732	1016	126.2	137.7	110.4
Diabetes (260)	292	150	142	291	149	142	13.4	11.8	15.4
Pellagra (281)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	2872	1459	1413	2891	1464	1427	132.7	116.4	155.0
Heart diseases (400-402,410-443)	6553	4035	2518	6561	4022	2539	301.2	319.8	275.8
Hypertension (444-447)	201	66	135	200	64	136	9.2	5.1	14.8
Arteriosclerosis (450)	326	207	119	327	208	119	15.0	16.5	12.9
Influenza (480-483)	59	20	39	59	19	40	2.7	1.5	4.3
#Pneumonia (490-493)	662	294	368	662	288	374	30.4	22.9	40.6
#Diarrhea & enteritis (571)	173	22	151	173	23	150	7.9	1.8	16.3
Acute nephritis (590,591)	47	12	35	47	12	35	2.2	1.0	3.8
Chronic nephritis, etc. (592-594)	234	75	159	237	79	158	10.9	6.3	17.2
Maternal deaths (640-689)	49	9	40	49	9	40	8.2	3.3	12.5
Cong. malformations (750-759)	229	130	99	239	138	101	11.0	11.0	11.0
Birth inj., atelect. (760-762)	307	162	145	307	166	141	5.2	6.0	4.4
#Other dis. early inf. (763-776)	888	249	639	894	250	644	15.0	9.1	20.1
Violent & acc. deaths (800-999)	1794	922	872	1804	929	875	82.8	73.9	95.0
Acc. deaths, total (800-962)	1417	755	662	1427	760	667	65.5	60.4	72.4
Motor vehicle acc. (810-835)	599	391	208	585	380	205	26.8	30.2	22.3
Ill-defined (780-795)	1748	465	1283	1730	444	1286	79.4	35.3	139.7

* Total birth and death rates per 1,000 population. Rates for fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from infant causes (760-776) per 1,000 live births. Maternal death rates per 10,000 live births. Death rates for specific causes per 100,000 population.

\# Pneumonia and diarrhea of the newborn (-28 days) included in other diseases peculiar to early infancy.