


VITAL STATISTICS
MISSISSIPPI
1963



Public Health Statistics
Mississippi State Board of Health
Jackson, Mississippi
A. L. Gray, M. D., Executive Officer

VITAL STATISTICS

MISSISSIPPI

1963

MISSISSIPPI STATE BOARD OF HEALTH
JACKSON, MISSISSIPPI

A. L. GRAY, M. D., EXECUTIVE OFFICER

TABLE OF CONTENTS

SYNOPSIS OF VITAL STATISTICS, MISSISSIPPI, 1963	Page
Introduction	I
Live Births	II
Fetal Deaths	V
Deaths	VIII
Marriages	XVIII
Divorces	XX

POPULATION

Table 1. Estimated Population, by County, by Race, Mississippi, July 1, 1963	1
--	---

LIVE BIRTHS

Table 2. By Place of Occurrence, and by Place of Residence of Mother, by Race, with Rates, Mississippi, 1963	3
Table 3. By Place of Residence of Mother, by Race and Attendant, and by Sex, Mississippi, 1963	6

FETAL DEATHS

Table 4. By Place of Occurrence; by Place of Residence of Mother, by Race and Attendant, and by Sex, Mississippi, 1963.....	10
---	----

DEATHS

Table 5. By Cause of Death, by Race, Residence Data, Mississippi, 1963 .	14
Table 6. By Place of Occurrence, and by Place of Residence, by Race, with Rates, Mississippi, 1963	26
Table 7. By Cause Groups and Certain Selected Causes, by Race, Sex, and Age, with Total Rates, Residence Data, Mississippi, 1963	29

DEATHS UNDER 1 YEAR

Table 8. By Selected Causes, by Race and Age, with Total Rates, Residence Data, Mississippi, 1963.....	38
--	----

MARRIAGES AND DIVORCES

Table 9. Marriages by Place of Licensure, and by Place of Residence of Bride, by Race of Bride; Divorces by Place of Occurrence, by Race, Mississippi, 1963	42
Table 10. Divorces by Cause of Divorce, by Race, Mississippi, 1963	44
Divorces by Number of Minor Children, by Race, Mississippi, 1963	44
Divorces by Number of Years Married, by Race, Mississippi, 1963	44

SUMMARY TABLES

Table 11. <u>Live Births, Mississippi, 1913-1963</u> , by Year, by Race, with Rates, Occurrence Data 1913-1943, Residence Data 1944-1963	45
Table 12. <u>Deaths, Mississippi, 1913-1963</u> , by Year, by Race, with Rates, Occurrence Data 1913-1943, Residence Data 1944-1963 ..	47
Table 13. <u>Maternal Deaths, Mississippi, 1913-1963</u> , by Year, by Race, with Rates, Occurrence Data 1913-1943, Residence Data 1944-1963	49
Table 14. <u>Deaths Under 1 Year, Mississippi, 1917-1963</u> , by Year, by Race, with Rates, Occurrence Data 1917-1943, Residence Data 1944-1963	51
Table 15. <u>Deaths Under 1 Month, Mississippi, 1924-1963</u> , by Year, by Race, with Rates, Occurrence Data 1924-1944, Residence Data 1945-1963	53
Table 16. <u>Deaths from Heart Diseases, Vascular Lesions, Malignant Neoplasms, Accidents, and Influenza and Pneumonia, Mississippi, 1954-1963</u> , by Year, by Race, with Rates, Residence Data	54
Table 17. <u>Marriages and Divorces, Mississippi, 1926-1963</u> , by Year, by Race, Occurrence Data	55

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH, by Place of Occurrence, and by Place of Residence, by Race, with Rates, 1963 ...

County Tables	59
State Table	141

SYNOPSIS OF 1963 VITAL STATISTICS

INTRODUCTION

The 1963 Vital Statistics Bulletin is the twenty-first consecutive issue prepared from data obtained from records of live births, fetal deaths, deaths, marriages, and divorces collected by the Division of Public Health Statistics

Population and Rates (See Table 1.)

Population figures used in this bulletin are estimates as of July 1, 1963, prepared by the Division of Sociology and Rural Life, Mississippi State University. According to these new estimates, there has been an increase of about 112,000 or 5.1 percent since the 1960 census. The increase for the whites has amounted to about 91,000 or 7.3 percent and for the nonwhites, about 21,000 or 2.2 percent. Figures for the counties indicate that 56 of the 82 have gained population, 19 have lost population, and in 7 the difference was less than 0.6 percent and thus considered as no change.

Rates for the state and counties by race were calculated using these figures, but no age or sex-specific rates (except the infant and maternal death rates which are based on live births) and no city rates were calculated because current estimates by age and sex and by city are not available.

Urban-Rural Classification

Incorporated towns with populations of 2,500 or more according to the 1960 census were considered as urban areas. All other areas were classified as rural.

Occurrence and Residence Allocation

Births, fetal deaths, and deaths were classified by place of occurrence and by place of residence. Some of the tables contain both occurrence and residence figures, but most of the tables are based on residence data only, as these data are considered more suitable for studying public health problems.

Occurrence data were obtained from certificates filed in Mississippi at the place of the event. Residence data include transcripts of records of events that occurred in other states to Mississippi residents and exclude records filed in this state for nonresidents. Residence assignment for births and fetal deaths was determined by the usual residence of the mother. For deaths it was determined by the usual residence of the decedent, except that deaths of inmates of institutions were reallocated to place of previous residence regardless of length of stay in the institution.

In Table A it may be seen that about 2,100 transcripts were received for vital events that occurred out of Mississippi to citizens of this state and that nearly 2,200 such events were registered here for residents of other states. The transcripts were sent from 35 states and the District of Columbia, and one birth record came from outside the United States. The first four states listed in Table A border on Mississippi and reported 87 percent of the out-of-state events.

Vital Statistics
Mississippi - 1963

Table A. Transcripts received, by place of occurrence,
and nonresident certificates

Place of Occurrence	Total Records	Live Births	Fetal Deaths	Deaths
<u>Transcripts</u>	<u>2,101</u>	<u>1,069</u>	<u>31</u>	<u>1,001</u>
Tennessee	1,109	542	21	546
Louisiana	331	151	3	177
Alabama	329	218	5	106
Arkansas	52	23	0	29
Florida	36	16	0	20
Illinois	36	14	0	22
Texas	33	9	0	24
Missouri	29	20	0	9
Kentucky	25	20	0	5
Other places	121	56	2	63
<u>Nonresident certificates</u>	<u>2,178</u>	<u>1,402</u>	<u>71</u>	<u>705</u>

Marriages were classified according to place of issuance of marriage license and also place of residence of bride. Since there is no exchange among the states of transcripts of marriage records for nonresidents, out-of-state marriages of Mississippi brides were not included in the tabulations. Divorces were classified only according to place where the decree was granted.

Classification of Causes of Death

Causes of death were assigned according to the seventh revision (1955) of the International Statistical Classification of Diseases, Injuries, and Causes of Death. Only the underlying cause of death was coded. Since the seventh revision differs only slightly from the sixth revision used from 1949 through 1957, figures for 1949-1963 are comparable, but comparison of data by cause for these years with those for the years prior to 1949 should be made with caution.

LIVE BIRTHS (See Tables 2, 3, 11.)

The live birth total for Mississippi residents in 1963 was 57,316, a figure 1,605 or 2.7 percent lower than that for 1962 and the smallest since 1945. An even greater decline occurred in the crude birth rate which dropped by 4.2 percent to 25.0 per 1,000 population in 1963, making it the lowest rate since 1941. (The birth rate for the country as a whole has been declining steadily for several years and in 1963 reached its lowest point since 1945.) Live births among the white population numbered 26,823; and in the 51 years (1913-1963) for which birth data are available, the low rate of 19.9 for 1963 was second only to the 19.7 experienced in the depression year of 1933. In the counties, rates for whites varied from 9.2 in Carroll County to 28.8 in Harrison County. The birth total for nonwhites was 30,493 and the rate was 32.4. Both of these figures were below those for 1962, and the rate was the smallest since 1948. County rates for nonwhites ranged from 18.5 in Itawamba County to 40.9 in Calhoun County.

Vital Statistics
Mississippi - 1963

Urban-Rural Distribution

During 1963, 21,693 children were born to urban residents and 35,623 to rural residents. Comparison of occurrence and residence data shows that, although only 38 percent of the total births were to urban residents, 64 percent occurred in urban places. The difference between the two sets of figures indicates that about 9,000 white mothers and about 6,000 nonwhite mothers from rural areas went to cities and towns for delivery.

Attendant

Table B which gives the distribution of live births to Mississippi residents by attendant and race shows that, in 1963, 75 percent occurred in hospitals or physicians' offices and clinics which provide in-patient care. An additional 2 percent were delivered by physicians in homes and places other than hospitals or clinics, leaving 23 percent with no medical attention at delivery. There was a tremendous difference between the two races in the proportion of mothers without a physician in attendance: less than one percent for whites, but 43 percent for nonwhites.

Table B. Live births by attendant and race

Attendant	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
<u>Total</u>	<u>57,316</u>	<u>100.0</u>	<u>26,823</u>	<u>100.0</u>	<u>30,493</u>	<u>100.0</u>
Physician in hospital	42,917	74.9	26,648	99.3	16,269	53.4
Physician in home	1,224	2.1	83	0.3	1,141	3.7
Midwife	13,063	22.8	74	0.3	12,989	42.6
Other	112	0.2	18	0.1	94	0.3

As shown by Table C, much progress has been made in obtaining hospitalization at time of delivery. Twenty years ago, in 1943, slightly over a fourth of the babies were born in a hospital; by 1953 the percentage had increased to 58, and in 1963 it was 75. The rate of increase for nonwhites was rapid through 1960, but has slowed considerably since then.

Table C. Percentage of live births in hospitals and clinics,
by race, 1943, 1953, 1959-1963

Year	Total	White	Nonwhite
1943	26.5	52.2	4.6
1953	58.2	93.3	30.5
1959	72.1	98.9	49.1
1960	73.2	99.0	51.3
1961	73.2	99.2	51.0
1962	74.6	99.3	52.9
1963	74.9	99.3	53.4

Vital Statistics
Mississippi - 1963

Month

The distribution of live births by month is given in Table D. August was the month with the largest number and also the highest daily average. April ranked lowest in number and daily average.

Table D. Live births and average per day, by month

Month	Births	Average per Day	Month	Births	Average per Day
January	4,968	160	July	5,221	168
February	4,516	161	August	5,636	182
March	4,767	154	September	5,276	176
April	3,923	131	October	5,003	161
May	4,080	132	November	4,650	155
June	4,446	148	December	4,830	156

Age of Mother

More mothers were in the age group 20-24 years than in any other. (See Table E.) There also were large numbers 15-19 and 25-29 years old, and 76 percent of all births were to mothers 15-29 years of age. Considerably more of the nonwhite than of the white mothers were either teenagers or 35 and over. Declines, some of which were quite large, were observed in the numbers of nonwhite mothers of all ages and white mothers of nearly all ages in 1963. In contrast, there was an increase in the number of white mothers at ages 20-24 and also at ages 45-49. The totals for the second group are small and therefore more subject to chance variation; but the increase in the younger group, though numerically small, is noteworthy.

Table E. Live births by age of mother and race,
with percent of total births

Age of Mother	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
All Ages	57,316	100.0	26,823	100.0	30,493	100.0
10-14	304	0.5	25	0.1	279	0.9
15-19	11,603	20.2	4,793	17.9	6,810	22.3
20-24	19,664	34.3	10,387	38.7	9,277	30.4
25-29	12,452	21.7	6,300	23.5	6,152	20.2
30-34	7,387	12.9	3,255	12.1	4,132	13.6
35-39	4,363	7.6	1,567	5.8	2,796	9.2
40-44	1,369	2.4	463	1.7	906	3.0
45-49	98	0.2	28	0.1	70	0.2
Unknown	76	0.1	5	*	71	0.2

* Less than 0.06 percent.

Vital Statistics
Mississippi - 1963

Sex

Liveborn males numbered 29,356 and females, 27,960. These totals yielded a sex ratio of 1,050 males per 1,000 females. As is generally true, the sex ratio for whites (1,068) was higher than that for nonwhites (1,034).

Plurality

Tabulation of the 57,649 births that occurred in Mississippi in 1963 showed that 1,345 or 2.3 percent were from plural deliveries. Among these was a set of white quadruplets, all females, born in Hinds County. There was a higher rate of plurality among nonwhites, 2.9 percent, than among whites, 1.6 percent. Among nonwhites the ratio of males to females for births from plural deliveries (1,091) was considerably higher than that for single deliveries (1,032), but among whites not only was the sex ratio for births from plural deliveries (867) lower than that for single deliveries (1,069), but it was also unusual in that it was less than 1,000.

Immaturity

Defining immaturity as birth weight of 5 1/2 pounds or less, 5,711 infants or 9.9 percent of those born in the state in 1963 were immature. The rate of immaturity was much greater among nonwhites than among whites, 12.6 percent in comparison with 6.8 percent.

Legitimacy

Mississippi women gave birth to 8,699 illegitimate children in 1963. This was 15.2 percent of the total births. According to the statements on birth certificates, 464 or 1.7 percent of the white infants and 8,235 or 27.0 percent of the non-white infants were born to unmarried mothers. The rate of illegitimate births among both whites and nonwhites has gone up steadily in recent years.

FETAL DEATHS (See Table 4.)

Regulations of the Mississippi State Board of Health specify that all fetal deaths, regardless of the length of gestation, shall be registered. Prior to 1952, only fetal deaths of 20 or more weeks gestation were reported; therefore, data on the all-inclusive registration are available for only 12 years, 1952-1963. Registration of both early and late fetal deaths is known to be incomplete; and because of the presumably sizeable number unreported, any or all of the data presented here could give a false picture regarding fetal deaths.

There were 2,795 fetal death records filed in the state in 1963, and 2,755 records for Mississippi residents. Both of these totals were slightly larger than in 1962. Fetal death ratios (fetal deaths per 1,000 live births) for the total population, 48.1, the whites, 46.9, and the nonwhites, 49.1, are all the highest recorded in the 12 years of expanded reporting. In the counties, ratios for whites varied from zero in Greene, Kemper, Tunica, and Winston Counties to 155.6 in Washington County. Ratios for nonwhites ranged from zero in Itawamba and Tishomingo Counties to 99.0 in Lawrence County.

Vital Statistics
Mississippi - 1963

The wide ranges in county ratios and the variations from year to year in individual counties probably can be attributed in large measure to variations in completeness of reporting rather than to actual differences. If it is assumed that the high ratio of 155.6 for whites in Washington County is not exaggerated, but represents complete reporting, it can be estimated from this ratio that one out of every seven or eight pregnancies terminates in fetal death.

Attendant

In Table F fetal deaths by attendant and race are shown. Deliveries in hospitals and clinics comprised 79 percent of the total; for 14.5 percent no medical attendant was present. The proportion of white fetal deaths reported from hospitals, 94.6 percent, was smaller than the proportion of white live births in hospitals, 99.3 percent. The reverse was true for nonwhites: 66 percent of fetal deaths, but only 53 percent of live births were delivered in hospitals. The probable underreporting of fetal deaths in the home and the tendency to transfer difficult and premature deliveries to a hospital may account for the difference noted for nonwhites.

Table F. Fetal deaths by attendant and race

Attendant	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
<u>Total</u>	<u>2,755</u>	<u>100.0</u>	<u>1,257</u>	<u>100.0</u>	<u>1,498</u>	<u>100.0</u>
Physician in hospital	2,176	79.0	1,189	94.6	987	65.9
Physician in home	179	6.5	66	5.2	113	7.5
Midwife	386	14.0	0	-	386	25.8
Other	14	0.5	2	0.2	12	0.8

Table G shows how the proportion of fetal deaths reported by hospitals has increased recently. The nonwhites have gained steadily, but the figures for whites have been somewhat erratic in the past 3 years. The gap between the two races is not so great for hospital deliveries of fetal deaths as for hospital deliveries of live births.

Table G. Percentage of fetal deaths delivered in hospitals or clinics, by race, 1953, 1959-1963

Year	Total	White	Nonwhite
1953	56.4	89.6	42.6
1959	73.5	94.4	59.6
1960	74.2	95.5	59.9
1961	75.2	93.4	61.9
1962	79.5	97.3	64.2
1963	79.0	94.6	65.9

Age of Mother

Fetal death ratios by age of mother and race are presented in Table H. The ratios for ages 15-19 and 20-24 were almost equal and the lowest on the table. From

Vital Statistics
Mississippi - 1963

age 25 on, the ratios increased in each successive 5-year age group, and in the groups 35 and over were more than twice those for ages 15-24. The ratio for mothers 10-14 years of age was approximately the same as that for mothers 30-34. There were no large differences by race.

Table H. Fetal deaths by age of mother and race, with fetal death ratios

Age of Mother	Total		White		Nonwhite	
	No.	Ratio*	No.	Ratio*	No.	Ratio*
<u>All Ages</u>	<u>2,755</u>	<u>48.1</u>	<u>1,257</u>	<u>46.9</u>	<u>1,498</u>	<u>49.1</u>
10-14	18	59.2	1	#	17	60.9
15-19	427	36.8	158	33.0	269	39.5
20-24	736	37.4	405	39.0	331	35.7
25-29	594	47.7	319	50.6	275	44.7
30-34	455	61.6	189	58.1	266	64.4
35-39	365	83.6	135	86.2	230	82.3
40-44	130	95.0	42	90.7	88	97.1
45-49	11	#	2	#	9	#
Unknown	19		6		13	

* Fetal deaths per 1,000 live births to mothers in specific age-race group.

Ratio not calculated on less than 100 live births.

Sex

Using the certificates with sex stated, a sex ratio of 1,293 males per 1,000 females was calculated. The ratio for nonwhites was considerably higher than the one for whites, 1,317 in comparison with 1,229. This is the reverse of the occurrence among live births. However, sex ratios determined from fetal death certificates vary widely from year to year in Mississippi, and in some years the ratio for whites is greater than that for nonwhites.

Plurality

Among fetal deaths occurring in the state the percentage from plural deliveries (4.4) was greater than the corresponding percentage of live births (2.3). For white fetal deaths the rate was only 2.4 percent, but for nonwhites it was 6.1 percent.

Length of Gestation

The stated length of gestation was less than 20 weeks on 1,269 of the certificates for Mississippi residents; it was 20 weeks or more on 1,357; and on 129 records the gestation period was not stated. The proportion of white fetal deaths at less than 20 weeks was 68 percent, but for nonwhites it was only 28 percent. This difference may be due partly to better reporting of early fetal deaths among whites and also to the fact that the late (20 weeks or more) fetal death ratio for nonwhites is more than 2 1/2 times that for whites.

Vital Statistics
Mississippi - 1963

Legitimacy

A larger percentage of fetal deaths than of live births was reported as illegitimate, 17.2 percent in comparison with 15.2 percent. The difference was small for nonwhites, but the rate for white fetal deaths was 2 1/2 times the rate for white live births.

DEATHS (See Tables 5-8, 12-16.)

In 1963 the resident death total was 23,339 and the crude death rate was 10.2 per 1,000 population. Both the number and the rate were above those for the preceding year, and the rate was the highest since 1943. The numbers for both races increased, and the total of 12,118 deaths among white persons was the largest ever recorded in Mississippi. The rate for nonwhites increased also, but the rate for whites was the same as in 1962.

Considerable variation was shown in the rates by county. The range for whites was from 6.2 in Jackson County to 16.3 in Kemper County, and the state death rate was 9.0. For nonwhites the state death rate was 11.9, with the lowest, 7.9, in Calhoun County and the highest, 17.4, in Issaquena County. Fifty-four percent of the deaths occurred in urban areas, but only 37 percent of those who died were urban residents.

Natural Increase (Excess of Birth Rate over Death Rate)

As a result of the continued decline in the birth rate and the continued rise in the death rate, the natural increase in population also dropped again in 1963, and the rate of 14.8 per 1,000 population was the smallest since 1941. For whites the rate of 10.9 was the lowest on record, even lower than in 1918 the year of the great influenza pandemic. The rate calculated for nonwhites was 20.5. There were three counties in the state (Carroll, Jefferson, and Kemper) in which the number of deaths of white residents was greater than the number of live births to white residents; in Noxubee County there was only one more birth than deaths; and in Franklin County only three more births than deaths among the whites. This indicates that the age distribution of the white populations of these counties is abnormal with few young adults and a high proportion of old people.

Month

According to Table I, March was the month in which the largest number of deaths occurred and also the highest average per day. In July the smallest number and the lowest daily average occurred.

Vital Statistics
Mississippi - 1963

Table I. Deaths and average per day, by month

Month	Deaths	Average per Day	Month	Deaths	Average per Day
January	2,251	73	July	1,619	52
February	2,174	78	August	1,751	56
March	2,503	81	September	1,681	56
April	1,703	57	October	1,815	58
May	1,822	59	November	1,842	61
June	1,851	62	December	2,327	75

Age at Death

The median age at death in Mississippi in 1963 was 67.2. The figure for whites was 70.3 and for nonwhites, 63.1. As shown by the data in Table J, the proportion of deaths among children under one year of age was almost as large as the proportion in the next 44 years of life (1-44 years of age), 10.2 percent in comparison with 12.0 percent respectively. There were fewer deaths in the decade 5-14 years than in any other period. Nearly 62 percent of the whites but only 47 percent of the nonwhites who died in 1963 had reached 65, or what is generally considered retirement age.

Table J. Deaths by age and race

Age	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
All Ages	23,339	100.0	12,118	100.0	11,221	100.0
Under 1	2,372	10.2	610	5.0	1,762	15.7
1-4	394	1.7	102	0.8	292	2.6
5-9	169	0.7	63	0.5	106	0.9
10-14	132	0.6	49	0.4	83	0.7
15-24	510	2.2	226	1.9	284	2.5
25-34	530	2.3	212	1.7	318	2.8
35-44	1,060	4.5	494	4.1	566	5.0
45-54	1,993	8.5	1,060	8.7	933	8.3
55-64	3,391	14.5	1,833	15.1	1,558	13.9
65-74	4,912	21.0	2,669	22.0	2,243	20.0
75-84	5,144	22.0	3,184	26.3	1,960	17.5
85 & over	2,714	11.6	1,615	13.3	1,099	9.8
Unknown	18	0.1	1	*	17	0.2

* Less than 0.06 percent.

Leading Causes of Death

In Table K are listed the ten leading causes of death in Mississippi in 1963. The first eight of these causes are the same and in the same order as in the preceding year. Congenital malformations (tenth place) was in ninth place in 1962. Homicide (ninth place) was not on the list in 1962; gastritis, duodenitis, enteritis, etc. was in tenth place in that year. In 1963 the ten leading causes accounted for 79.5 percent of the total deaths.

Vital Statistics
Mississippi - 1963

Table K. Ten leading causes of death by race, with percent of total deaths

Cause and International List Number	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
1. Diseases of heart (400-443)	7,300	31.3	4,396	36.3	2,904	25.9
2. Vascular lesions affecting central nervous system (330-334)	3,154	13.5	1,650	13.6	1,504	13.4
3. Malignant neoplasms (140-205)	2,945	12.6	1,859	15.3	1,086	9.7
4. Accidents (800-962)	1,571	6.7	851	7.0	720	6.4
5. Certain diseases of early infancy (760-776)	1,262	5.4	389	3.2	873	7.8
6. Influenza and pneumonia (480-493)	1,113	4.8	420	3.5	693	6.2
7. General arteriosclerosis (450)	364	1.6	250	2.1	114	1.0
8. Diabetes mellitus (260)	326	1.4	132	1.1	194	1.7
9. Homicide (964, 980-985)	267	1.1	48	0.4	219	2.0
10. Congenital malformations (750-759)	256	1.1	147	1.2	109	1.0

Leading Causes by Age

The leading causes of death in each age group except infants (see infant mortality section) are given in Table L. Although no two lists were alike, several causes appeared a number of times and malignant neoplasms was on every list. Accidents was the leading cause in all age groups under 45, was fourth at 45-64, and was sixth at 65 and over. The list for the 1-4 year group was the only one on which diseases of the heart did not appear, and the 15-24 and 25-44 year groups were the only ones without influenza and pneumonia. Vascular lesions affecting the central nervous system showed up on the list for all groups 25 and over. Among persons 15-24 years of age, 64 percent of the deaths were due to external causes (accidents, homicide, and suicide).

Ill-defined and Unknown Causes

In 1963, 2,498 or 10.8 percent of the death certificates filed in Mississippi contained no satisfactory statement of cause. The percentages by race were 5.3 for whites and 16.8 for nonwhites. There were 69 certificates which showed senility as the only cause; on 708, ill-defined diseases of early infancy appeared; and the remaining 1,721 had other ill-defined causes or no cause at all. Over 20 percent of the death records filed in 4 counties (Benton, Carroll, Holmes, and Madison) and from 15 to 20 percent of those filed in 11 counties (Bolivar, Leflore, Marion, Marshall, Noxubee, Perry, Quitman, Tallahatchie, Tunica, Warren, and Washington) lacked a satisfactory statement of cause.

If it were possible to know and tabulate the true cause of death for persons whose certificates lacked this information, the order of the leading causes might not be the same as presently shown and even the causes themselves might be different. The group of deaths from ill-defined and unknown causes, if placed on the list of leading causes, would rank fourth in the total population and would appear as one of the five leading causes in every age group.

Vital Statistics
Mississippi - 1963

Table L. Five leading causes of death by age group

Cause of Death	Age Group	No. of Deaths	% of Total
1-4 years			
<u>All Causes</u>		<u>394</u>	<u>100.0</u>
1. Accidents		127	32.2
2. Influenza and pneumonia		88	22.3
3. Congenital malformations		29	7.4
4. Malignant neoplasms		17	4.3
5. <u>Gastritis, duodenitis, enteritis, and colitis</u>		15	3.8
5-14 years			
<u>All Causes</u>		<u>301</u>	<u>100.0</u>
1. Accidents		136	45.2
2. Malignant neoplasms		20	6.6
3. Congenital malformations		16	5.3
4. Diseases of heart		15	5.0
5. Influenza and pneumonia		13	4.3
15-24 years			
<u>All Causes</u>		<u>510</u>	<u>100.0</u>
1. Accidents		252	49.4
2. Homicide		56	11.0
3. Malignant neoplasms		31	6.1
4. Diseases of heart		20	3.9
5. Suicide		17	3.3
25-44 years			
<u>All Causes</u>		<u>1,590</u>	<u>100.0</u>
1. Accidents		355	22.3
2. Diseases of heart		294	18.5
3. Malignant neoplasms		208	13.1
4. Homicide		138	8.7
5. <u>Vascular lesions affecting central nervous system</u>		100	6.3
45-64 years			
<u>All Causes</u>		<u>5,384</u>	<u>100.0</u>
1. Diseases of heart		1,871	34.8
2. Malignant neoplasms		1,096	20.4
3. Vascular lesions affecting central nervous system		612	11.4
4. Accidents		296	5.5
5. Influenza and pneumonia		112	2.1
65 years and over			
<u>All Causes</u>		<u>12,770</u>	<u>100.0</u>
1. Diseases of heart		5,087	39.8
2. Vascular lesions affecting central nervous system		2,432	19.0
3. Malignant neoplasms		1,568	12.3
4. Influenza and pneumonia		593	4.6
5. <u>General arteriosclerosis</u>		343	2.7

Vital Statistics
Mississippi - 1963

Specific Causes

Heart disease, at the top of the list of principal causes of death in Mississippi beginning in 1925 and every year since then except 1929, was responsible for 7,300 deaths or more than 31 percent of the total in 1963. The death rate for whites (325.9 per 100,000 population) was higher than that for nonwhites (308.5), but the difference was not large. Arteriosclerotic and degenerative heart disease was the type which most frequently caused death and hypertensive heart disease was the second most fatal type. (See Table M.)

Table M. Deaths from diseases of the heart, by type and race

Cause and International List Number	Total	White	Nonwhite
<u>Diseases of the heart (400-443)</u>	<u>7,300</u>	<u>4,396</u>	<u>2,904</u>
Rheumatic fever (400-402)	10	0	10
Chronic rheumatic heart disease (410-416)	93	41	52
Arteriosclerotic and degenerative heart disease (420-422)	5,359	3,691	1,668
Other diseases of the heart (430-434)	631	295	336
<u>Hypertensive heart disease (440-443)</u>	<u>1,207</u>	<u>369</u>	<u>838</u>

Vascular lesions affecting the central nervous system was the second ranking cause of death, taking the lives of 3,154 Mississippians. This figure was 13.5 percent of the total, but less than half the toll from heart disease. There was a notable difference in the death rates for the two races, 122.3 for whites and 159.8 for nonwhites. According to the data in Table N, cerebral hemorrhage was the most important type of fatal vascular lesion.

Table N. Deaths from vascular lesions affecting central nervous system, by type of lesion and race

Cause and International List Number	Total	White	Nonwhite
<u>Vascular lesions affecting central nervous system (330-334)</u>	<u>3,154</u>	<u>1,650</u>	<u>1,504</u>
Subarachnoid hemorrhage (330)	111	58	53
Cerebral hemorrhage (331)	1,954	965	989
Cerebral embolism and thrombosis (332)	705	446	259
<u>Other and ill-defined (333,334)</u>	<u>384</u>	<u>181</u>	<u>203</u>

Malignant neoplasms, in third place among the leading causes, accounted for 2,945 deaths in 1963. Both this number and the rate of 128.6 were larger than in 1962 and continued the upward trend which has prevailed for many years. The age-race-sex distribution of cancer deaths is shown in Table O. Less than 10 percent occurred among persons under 45 years of age, and there were more deaths at 65-74 years than in any other age period. For white females, however, the number of deaths at 75-84 years was slightly higher than the number at 65-74. The two principal sites of fatal cancers were the digestive organs and peritoneum and the genito-urinary organs and breast.

Vital Statistics
Mississippi - 1963

Table 0. Deaths from malignant neoplasms, by age, sex, and race

Age	Number of Deaths					Percent				
	Total	Male		Female		Total	Male		Female	
		White	Non-white	White	Non-white		White	Non-white	White	Non-white
All Ages	2,945	1,080	559	779	527	100.0	100.0	100.0	100.0	100.0
Under 10	30	12	8	7	3	1.0	1.1	1.4	0.9	0.6
10-14	9	3	1	1	4	0.3	0.3	0.2	0.1	0.8
15-24	31	13	6	7	5	1.0	1.2	1.1	0.9	0.9
25-34	47	14	5	15	13	1.6	1.3	0.9	1.9	2.5
35-44	161	48	27	47	39	5.5	4.4	4.8	6.0	7.4
45-54	445	154	67	131	93	15.1	14.2	12.0	16.8	17.6
55-64	651	240	115	161	135	22.1	22.2	20.6	20.7	25.6
65-74	800	309	175	173	143	27.2	28.6	31.3	22.2	27.1
75-84	577	219	111	181	66	19.6	20.3	19.8	23.2	12.5
85 & over	191	68	42	56	25	6.5	6.3	7.5	7.2	4.7
Unknown	3	0	2	0	1	0.1	-	0.4	-	0.2

Chronic respiratory diseases have been increasingly prominent as causes of both morbidity and mortality in recent years. The most important of these diseases are listed in Table P with mortality figures for certain years for which comparable data are available. A significant decline has occurred in deaths from respiratory tuberculosis, and no marked trend in either direction has been observed for asthma or bronchiectasis. All of the other causes listed, however, show definite upward trends. Deaths from malignant neoplasms of the respiratory system, the most prominent of these chronic respiratory diseases, rose by 210 percent from 1949 to 1963 when the total was 499, and the increase in deaths from cancer of the lung, bronchus, and trachea was ever larger. Emphysema, another chronic respiratory disease for which a remarkable increment was recorded (6 deaths listed in 1949 and 152 in 1963) is now the second most important of these diseases. Reported mortality from chronic and unqualified bronchitis (including bronchitis with emphysema) and from other chronic interstitial pneumonia has shown a generally upward trend, although there has been considerable year to year variation.

Table P. Deaths from chronic respiratory diseases, 1949, 1955, 1960 1963

Cause and International List Number	1949	1955	1960	1963
Tuberculosis of respiratory system (001-008)	552	189	135	106
Malignant neoplasms of respiratory system (160-165)	161	267	357	499
Malignant neoplasms of lung, bronchus, and trachea (162,163)	140	229	317	443
Asthma (241)	55	56	74	45
Bronchitis, chronic and unqualified (501,502)	22	23	30	49
Bronchitis with emphysema (502.0)	0	4	10	23
Other chronic interstitial pneumonia (525)	4	9	20	27
Bronchiectasis (526)	27	21	25	19
Emphysema (527.1)	6	25	89	152

Vital Statistics
Mississippi - 1963

Influenza and pneumonia, acute respiratory diseases which rank in sixth place among the leading causes of death, were directly responsible for the loss of 1,113 lives in Mississippi in 1963, and the death rate was 48.6 per 100,000 population. Both of these figures were much larger than any others recorded in the 15-year period 1949-1963 for which comparable data are available. The total for 1963 included 109 deaths due to influenza alone, 144 due to influenza with pneumonia, and 860 due to pneumonia. As has been true for most of the past fifteen years, the rate for nonwhites was more than double the rate for whites, 73.6 in comparison with 31.1 per 100,000 population.

Scarlet fever and diphtheria are acute communicable diseases which merit special mention as causes of death: scarlet fever because a death from this disease was reported in 1963 for the first time since 1952; and diphtheria because no deaths from this cause occurred in 1963, and 1957 is the only other year on record in which there were no diphtheria deaths.

Accidents caused 1,571 deaths among Mississippians in 1963, and the death rate was 68.6 per 100,000 population. The number was the largest since 1940, and the rate was one of the highest although 3 other years in that period were slightly higher. Accidents ranked fourth among the leading causes of death in the state and was the leading cause for persons 1-44 years of age. There were more accidental deaths in December than in any other month, but the peak for motor vehicle accidents alone came in November. Deaths caused by accidents averaged 4.3 per day.

Table Q lists the five chief causes of accidental death. These are the same ones which have appeared on the list for a number of years and in 1963 caused 76 percent of all accidental deaths. There was an increase in the number of deaths from motor vehicle accidents, and both the number (675) and the rate of 29.5 per 100,000 population for 1963 (residence data) were the highest ever recorded in the state. The number of motor vehicle accident deaths that occurred in the state, 705, was also the largest on record.

Table Q. Five leading causes of accidental deaths by race,
with percent of total accidental deaths

Type of Accident and International List Number	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
1. Motor vehicle (810-835)	675	43.0	435	51.1	240	33.3
2. Fire and explosion of combustible material (916)	223	14.2	69	8.1	154	21.4
3. Falls (900-904)	120	7.6	88	10.3	32	4.4
4. Drowning, except in water transport (929)	91	5.8	31	3.6	60	8.3
5. Firearm (919)	83	5.3	47	5.5	36	5.0

Deaths occurring in Mississippi from nontransport accidents (International Codes 870-936) numbered 778. Of these, 419 or 54 percent were due to accidents that took place in the home, and the two principal causes of the home accidents were fire and explosion of combustible material (191 deaths) and falls (71 deaths). Nearly three-fourths of the victims of fatal home accidents were either children under 15 or persons 65 and over.

Vital Statistics
Mississippi - 1963

Suicide and homicide were the causes of death stated on the certificates of 126 and 267 Mississippians in 1963. Suicide was not one of the ten leading causes of death, but did show up in eleventh place among the white population. A large increase in homicides occurred in 1963 with the result that this cause ranked ninth in the total population and seventh among nonwhites. The most frequently used method for both suicides and homicides was firearms and explosives. There were more deaths from suicide in the age group 45-54 years than in any other period; for homicides, ages 25-34 and 35-44 were the periods of highest incidence. The suicide rate for white persons was over 3 times the rate for nonwhites, but the relationship between the homicide rates was reversed with the rate for nonwhites more than 6 times that for whites.

Maternal deaths (deaths from deliveries and complications of pregnancy, child-birth, and the puerperium) totaled 44 and the rate was 7.7 deaths per 10,000 live births. Both the number and the rate for 1963 were the lowest ever recorded in this state. There was a small increase for white mothers, but, for nonwhite mothers, the number and rate were the lowest on record. Toxemia was the cause of 12 and hemorrhage of 9 maternal deaths.

Age and race-specific maternal death rates are given in Table R. The safest period for childbearing was the decade 15-24 years of age, and in this period the rates for whites and nonwhites were nearly equal. At all other ages, however, the rates for nonwhite mothers were far higher than those for white mothers, and the rates for nonwhites in the youngest group (10-14 years) and in the 35-44 year group were extremely large.

Table R. Maternal deaths by age and race, with rates


Age	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
All Ages	44	7.7	9	3.4	35	11.5
10-14	1	32.9	0	-	1	35.8
15-24	7	2.2	3	2.0	4	2.5
25-34	19	9.6	3	3.1	16	15.6
35-44	17	29.6	3	14.8	14	37.8

* Deaths per 10,000 live births to mothers in specific age-race group.

Infant Mortality

The infant death total for 1963 was 2,372 and the infant mortality rate was 41.4 per 1,000 live births. Infant deaths among whites declined by 95 between 1962 and 1963, and the rate of 22.7 in 1963 was the lowest on record in this state. There was a marked contrast between these figures and the ones for nonwhites. The total for nonwhites increased by 114 during 1963, and the rate of 57.8, which was 2 1/2 times the rate for whites, was the highest since 1941. Study of the accompanying graph of infant mortality rates shows that the downward trend for whites has slowed considerably since 1955 and that the curve for the nonwhites has been going up since 1946.

Vital Statistics
Mississippi - 1963



Mississippi had the highest infant mortality rate in the United States in 1962 (latest year for which national data are available). The rate was 15 percent larger than that for the District of Columbia which ranked second and 17 percent above that for Alaska which ranked third. The high rate in Mississippi was due not only to the extremely bad situation among the nonwhites but also to the fact that, although progress has been made in reducing the white infant mortality rate, it is still one of the highest in the country.

Information on infant mortality by age and race is given in Table S. These figures show that the highest rate for white infants was in the first day, and there was a substantial drop after the first 28 days. The reverse was true for nonwhite infants with the lowest rate under one day and the highest at 28 days to one year. All the rates for nonwhite infants were notably larger than those for white infants; in the neonatal period (first 28 days of life) they were about twice as high, and after the neonatal period the rate for nonwhites was 5 times the rate for whites. Study of the trend of neonatal death rates shows that the rate of 17.5 per 1,000 live births for whites was the smallest on record, but the rate of 32.1 for nonwhites was the largest since 1934.

Vital Statistics
Mississippi - 1963

Table S. Infant deaths by age and race, with rates

Age	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
<u>Total under one year</u>	<u>2,372</u>	<u>41.4</u>	<u>610</u>	<u>22.7</u>	<u>1,762</u>	<u>57.8</u>
Under one day	711	12.4	238	8.9	473	15.5
One to 28 days	738	12.9	232	8.6	506	16.6
<u>28 days to one year</u>	<u>923</u>	<u>16.1</u>	<u>140</u>	<u>5.2</u>	<u>783</u>	<u>25.7</u>

* Deaths per 1,000 live births.

Table T presents the five leading causes of infant deaths. These five causes were responsible for 38 percent of the total infant deaths in 1963 and for 66 percent of those with a definite cause given. The first four causes are the same ones which have been listed every year for which comparable data are available, 1949-1963, but there have been variations in the order. The fifth-ranking cause, accidents, appeared on the list for the first time in 1963 and replaced birth injuries which had been on the list for 14 years. However, there was a difference of only two in the number of deaths from these two causes in 1963. Marked contrasts were noted in the rates by race for some of the causes; the most pronounced was in the gastritis group, 0.2 for whites and 4.3 for nonwhites.

Table T. Leading causes of infant deaths by race, with rates

Cause of Death	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
<u>All Causes</u>	<u>2,372</u>	<u>41.4</u>	<u>610</u>	<u>22.7</u>	<u>1,762</u>	<u>57.8</u>
1. Influenza and pneumonia, except pneumonia of newborn	260	4.5	33	1.2	227	7.4
2. Postnatal asphyxia and atelectasis	229	4.0	104	3.9	125	4.1
3. Congenital malformations	170	3.0	101	3.8	69	2.3
4. Gastritis, duodenitis, enteritis, & colitis, except diarrhea of newborn	136	2.4	6	0.2	130	4.3
5. Accidents	102	1.8	23	0.8	79	2.6

* Deaths per 1,000 live births.

Tabulation of the leading causes by age reveals that postnatal asphyxia and atelectasis ranked first among those who died in the first day and caused almost as many deaths as the next three leading causes for this age group. Birth injuries ranked second as a cause of these early deaths. The leading cause at 1-27 days was the same as for under one day, and pneumonia of the newborn was second. Influenza and pneumonia was at the top of the list for children 28 days to one year, and the gastritis group was next in importance.

Immaturity, though no longer considered a distinct classification in ranking causes of infant deaths, remains an important factor. There were 855 deaths, 44 percent of those for white children and 33 percent of those for nonwhites, attributed to immaturity alone or immaturity in combination with another condition.

Vital Statistics
Mississippi - 1963

Because 1,007 or 42 percent of the infant death records in 1963 either had no cause of death or had ill-defined causes, the data previously presented regarding cause must be considered as incomplete or possibly erroneous. If correct information could be obtained for this large number of records, the true number of deaths from each cause could be ascertained, thereby making it possible to determine the most effective preventive measures to be taken.

MARRIAGES (See Tables 9, 17.)

Marriages in Mississippi in 1963 numbered 22,140; this was 133 more than in 1962. Although the increase was small, 1963 was the fourth consecutive year in which a gain occurred. Both the total of 14,508 for whites and that of 7,632 for nonwhites were slightly higher than those for 1962.

Month

As usual, more marriages occurred in June than in any other month, and this month had the highest daily average. (See Table U.) December ranked second with August a close third.

Table U. Marriages and average per day, by month

Month	Marriages	Average per Day	Month	Marriages	Average per Day
<u>Total</u>	<u>22,140</u>	<u>61</u>			
January	1,495	48	July	1,954	63
February	1,395	50	August	2,199	71
March	1,730	56	September	1,653	55
April	1,548	52	October	1,621	52
May	1,716	55	November	1,850	62
June	2,721	91	December	2,258	73

Officiant

Ministers performed 72 percent of the marriage ceremonies in the state. Justices of the peace officiated at nearly all of the others, though a few were performed by other qualified officiants.

Age of Bride and Groom

According to Table V which shows the age distribution of brides and grooms, 9,542 or 43 percent of the brides were teenagers and 4,351 or one out of every five brides was under 18 years of age. There were 3 brides who were only 12 years old, 18 who were 13 years old, and 112 who were 14 years old. Grooms in their teens numbered 3,635 (16.4 percent of the total), and 3 were only 14 years of age. In contrast with these young persons, there were 24 brides and 130 grooms who were 75 years of age or older. The oldest bride

Vital Statistics
Mississippi - 1963

was 84 and the oldest groom 96. Although in most instances the difference between the ages of the two parties to the marriage was not large, five of the couples who married in 1963 were 50 years or more apart in age, and the greatest difference was in the marriage of a bride of 20 and a groom of 75.

Table V. Brides and grooms by age

Age	Bride		Groom	
	No.	%	No.	%
<u>Total</u>	<u>22,140</u>	<u>100.0</u>	<u>22,140</u>	<u>100.0</u>
Under 15	133	0.6	3	*
15	800	3.6	17	0.1
16	1,499	6.8	68	0.3
17	1,919	8.7	465	2.1
18	2,747	12.4	1,195	5.4
19	2,444	11.0	1,887	8.5
20-24	6,281	28.4	9,353	42.2
25-34	2,830	12.8	4,517	20.4
35-44	1,643	7.4	1,925	8.7
45-54	1,070	4.8	1,189	5.4
55-64	560	2.5	804	3.6
65-74	180	0.8	570	2.6
75 and over	24	0.1	130	0.6
Unknown	10	*	17	0.1

* Less than 0.06 percent.

Previous Marriages

Both the bride and the groom were being married for the first time in 65 percent of the weddings in 1963. Table W shows data on those who had previous marriages. Among the 27.6 percent of the brides and the 28.0 percent of the grooms who had been married before, nearly three-fourths were divorced persons. There were 1,475 brides (6.7 percent of the total) and 1,511 grooms (6.8 percent) who had two or more previous marriages. Two of the grooms had been married 7 times before and 2 of the brides had been married 9 times before.

Table W. Number of previous marriages of brides and grooms

Previous Marriages	Bride		Groom	
	No.	%	No.	%
<u>Total</u>	<u>22,140</u>	<u>100.0</u>	<u>22,140</u>	<u>100.0</u>
None	15,992	72.2	15,891	71.8
One	4,627	20.9	4,685	21.2
Two	1,240	5.6	1,205	5.4
Three or more	235	1.1	306	1.4
Not stated	46	0.2	53	0.2

Vital Statistics
Mississippi - 1963

Nonresident Brides

Out-of-state brides who were married in Mississippi in 1963 numbered 3,625, and 3,074 of them married nonresident grooms. Brides from other states comprised 16 percent of the total, a sizeable drop from the 66 percent reported in 1957 before the passage of a stricter marriage law in this state. In 1963, these brides came from all but 5 of the other 49 states and from the District of Columbia, and 12 came from foreign countries. However, eighty percent of these nonresident brides were from the bordering states of Alabama, Arkansas, Louisiana, and Tennessee, and 78 percent of the licenses were issued in Mississippi counties that are adjacent to these states. DeSoto County which is about 5 miles from Memphis, Tennessee, issued 809 licenses or 22 percent of the total number issued to nonresident brides.

Out-of-State Use of Mississippi Certificates

On July 1, 1956, a new marriage record containing more detailed information was put into use in Mississippi. These records have shown that, in every year since that time, some Mississippi licenses have been used for marriages performed in other states. These marriages have not been included in the annual tabulations because they did not occur in this state and are of questionable legality. In 1963, according to a special count, 107 records showed the place of marriage as another state; 74 of these were used in Louisiana and 19 in Tennessee. Two-thirds of the licenses were issued by DeSoto, Pearl River, and Wilkinson Counties. The out-of-state place of residence of either the bride or the groom or of both was the place of occurrence of 96 of these 107 marriages. Nine of the other 11 marriages were between persons who were residents of Mississippi but used their licenses in another state. The remaining 2 couples were residents of other states who used Mississippi licenses in yet another state.

DIVORCES (See Tables 9, 10, 17.)

There were 5,452 divorces in Mississippi in 1963. This total was the second largest since 1951 and only 15 less than that for 1962 which was the largest since 1951. Although there was little difference between the totals for 1962 and 1963, an increase among the whites brought the figure of 3,930 for that race up to the third highest on record, exceeded only in the postwar years 1946 and 1947. The number of divorces among nonwhites declined to 1,522.

Cause and Complainant

According to the information on the statistical records submitted by Chancery Clerks, 90 percent of the divorces were granted because of either cruel and inhuman treatment or desertion. The leading cause listed for whites was cruel and inhuman treatment with 67 percent of the total. For nonwhites, desertion ranked first and was the reason stated on 64 percent of the records. The wife was the complainant in 71 percent of the cases involving white persons, but the husband was the complainant in slightly over half of those among nonwhites.

Vital Statistics
Mississippi - 1963

Minor Children

Nearly 42 percent of the white couples and 56 percent of the nonwhites had no minor children. There were 2,885 couples who did have minor children, and the total number of such children was 6,378. Five or more children each were involved in 203 cases, and in 4 of these there were 10 or more children each.

Duration of Marriage

In 1963, 374 or nearly 7 percent of the couples obtained divorces after less than a year of marriage, and 37 percent had been married less than five years. At the other end of the range were 12 couples who divorced after 50 or more years of marriage and one of these had been married 61 years. Among the divorced couples there was a notable difference in duration of marriage by race. The average duration for nonwhites (16.7 years) was nearly twice as long as that for whites (8.5 years). Only 10 percent of the white couples had been married as long as 20 years, but 33 percent of the nonwhite couples were in this category.



TABLE 1
ESTIMATED POPULATION*
MISSISSIPPI
JULY 1, 1963

By County, by Race

<u>County</u>	<u>Total</u>	<u>White</u>	<u>Nonwhite</u>
<u>STATE</u>	<u>2,290,180</u>	<u>1,348,880</u>	<u>941,300</u>
Adams	37,520	18,560	18,960
Alcorn	26,840	23,380	3,460
Amite	17,100	8,150	8,950
Attala	22,520	13,420	9,100
Benton	8,160	3,710	4,450
Bolivar	57,620	19,570	38,050
Calhoun	16,630	12,840	3,790
Carroll	12,030	5,550	6,480
Chickasaw	17,610	10,620	6,990
Choctaw	7,850	5,310	2,540
Claiborne	9,800	2,840	6,960
Clarke	17,070	11,580	5,490
Clay	19,760	9,980	9,780
Coahoma	48,500	15,160	33,340
Copiah	27,250	12,790	14,460
Covington	16,190	10,740	5,450
DeSoto	27,310	12,710	14,600
Forrest	56,590	41,770	14,820
Franklin	9,230	5,710	3,520
George	11,130	9,450	1,680
Greene	8,350	6,780	1,570
Grenada	19,120	10,320	8,800
Hancock	16,100	13,140	2,960
Harrison	125,390	105,210	20,180
Hinds	199,680	117,020	82,660
Holmes	27,040	7,860	19,180
Humphreys	19,380	6,090	13,290
Issaquena	2,690	850	1,840
Itawamba	17,350	16,430	920
Jackson	66,990	55,350	11,640
Jasper	17,130	9,030	8,100
Jefferson	10,150	3,250	6,900
Jeff Davis	13,430	6,800	6,630
Jones	59,070	43,890	15,180
Kemper	11,670	4,900	6,770
Lafayette	22,170	14,710	7,460

Table 1 - Page 2
 Estimated Population - Mississippi - July 1, 1963

County	Total	White	Nonwhite
Lamar	14,890	12,720	2,170
Lauderdale	73,620	50,340	23,280
Lawrence	9,360	6,120	3,240
Leake	17,390	9,850	7,540
Lee	44,340	32,670	11,670
Leflore	48,100	17,400	30,700
Lincoln	26,170	18,230	7,940
Lowndes	52,500	33,520	18,980
Madison	33,310	10,480	22,830
Marion	21,440	13,840	7,600
Marshall	25,120	7,550	17,570
Monroe	36,500	25,260	11,240
Montgomery	13,350	8,080	5,270
Neshoba	21,460	15,800	5,660
Newton	20,620	13,700	6,920
Noxubee	17,540	5,130	12,410
Oktibbeha	28,220	15,700	12,520
Panola	31,220	14,130	17,090
Pearl River	25,910	20,480	5,430
Perry	8,530	5,820	2,710
Pike	34,870	19,870	15,000
Pontotoc	18,820	15,360	3,460
Prentiss	19,280	16,830	2,450
Quitman	23,070	9,200	13,870
Rankin	38,050	23,540	14,510
Scott	23,050	14,640	8,410
Sharkey	10,550	2,980	7,570
Simpson	22,240	15,160	7,080
Smith	15,860	12,060	3,800
Stone	6,630	4,870	1,760
Sunflower	48,190	14,820	33,370
Tallahatchie	24,920	8,930	15,990
Tate	21,470	9,280	12,190
Tippah	16,720	13,500	3,220
Tishomingo	16,640	15,930	710
Tunica	16,960	3,520	13,440
Union	20,800	17,510	3,290
Walthall	13,450	7,420	6,030
Warren	44,160	24,440	19,720
Washington	77,180	33,710	43,470
Wayne	16,180	9,950	6,230
Webster	10,390	7,540	2,850
Wilkinson	12,920	3,990	8,930
Winston	19,560	10,680	8,880
Yalobusha	12,730	7,080	5,650
Yazoo	33,480	13,780	19,700

* Prepared by Division of Sociology and Rural Life,
 Mississippi State University.

TABLE 2
LIVE BIRTHS
MISSISSIPPI
1963

By Place of Occurrence, and by Place of Residence of Mother, by Race, with Rates

County	Place of Occurrence			Place of Residence					
	Number		Non-white	Number			Total	Rate*	
	Total	White		Total	White	Non-white		White	Non-white
<u>STATE</u>	<u>57,649</u>	<u>27,061</u>	<u>30,588</u>	<u>57,316</u>	<u>26,823</u>	<u>30,493</u>	<u>25.0</u>	<u>19.9</u>	<u>32.4</u>
Adams	1,491	694	797	1,046	464	582	27.9	25.0	30.7
Alcorn	568	464	104	525	421	104	19.6	18.0	30.0
Amite	153	0	153	337	108	229	19.7	13.2	25.6
Attala	477	202	275	481	201	280	21.4	15.0	30.8
Benton	115	1	114	187	55	132	22.9	14.8	29.7
Bolivar	1,936	352	1,584	1,802	351	1,451	31.3	17.9	38.1
Calhoun	334	148	186	360	205	155	21.6	16.0	40.9
Carroll	142	9	133	212	51	161	17.6	9.2	24.8
Chickasaw	408	232	176	412	215	197	23.4	20.2	28.2
Choctaw	187	101	86	173	94	79	22.0	17.7	31.1
Claiborne	173	25	148	243	46	197	24.8	16.2	28.3
Clarke	125	64	61	335	167	168	19.6	14.4	30.6
Clay	458	193	265	492	211	281	24.9	21.1	28.7
Coahoma	1,520	370	1,150	1,509	327	1,182	31.1	21.6	35.4
Copiah	840	253	587	668	233	435	24.5	18.2	30.1
Covington	173	117	56	313	163	150	19.3	15.2	27.5
DeSoto	589	70	519	750	265	485	27.5	20.8	33.2
Forrest	1,583	1,160	423	1,281	826	455	22.6	19.8	30.7
Franklin	138	62	76	175	77	98	19.0	13.5	27.8
George	343	277	66	316	248	68	28.4	26.2	40.5
Greene	122	72	50	178	125	53	21.3	18.4	33.8
Grenada	471	154	317	488	174	314	25.5	16.9	35.7
Hancock	219	153	66	344	272	72	21.4	20.7	24.3
Harrison	4,100	3,353	747	3,751	3,029	722	29.9	28.8	35.8
Hinds	6,028	3,055	2,973	5,249	2,492	2,757	26.3	21.3	33.4
Holmes	786	130	656	756	113	643	28.0	14.4	33.5
Humphreys	550	66	484	574	95	479	29.6	15.6	36.0
Issaquena	38	0	38	80	15	65	29.7	17.6	35.3
Itawamba	64	45	19	254	237	17	14.6	14.4	18.5
Jackson	1,454	1,040	414	1,822	1,370	452	27.2	24.8	38.8
Jasper	177	99	78	399	148	251	23.3	16.4	31.0
Jefferson	105	0	105	247	41	206	24.3	12.6	29.8
Jeff Davis	291	159	132	268	117	151	20.0	17.2	22.8
Jones	2,114	1,017	1,097	1,263	788	475	21.4	18.0	31.3
Kemper	202	38	164	251	58	193	21.5	11.8	28.5
Lafayette	486	246	240	514	259	255	23.2	17.6	34.2

Table 2 - Page 2
Live Births - Mississippi - 1963

County	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Lamar	111	81	30	293	236	57	19.7	18.6	26.3
Lauderdale	2,092	1,206	886	1,698	999	699	23.1	19.8	30.0
Lawrence	137	49	88	210	109	101	22.4	17.8	31.2
Leake	251	97	154	333	137	196	19.1	13.9	26.0
Lee	1,353	1,006	347	1,062	717	345	24.0	21.9	29.6
Leflore	1,461	421	1,040	1,422	374	1,048	29.6	21.5	34.1
Lincoln	600	357	243	596	323	273	22.8	17.7	34.4
Lowndes	1,526	872	654	1,476	821	655	28.1	24.5	34.5
Madison	796	132	664	903	177	726	27.1	16.9	31.8
Marion	444	237	207	525	291	234	24.5	21.0	30.8
Marshall	642	95	547	743	165	578	29.6	21.8	32.9
Monroe	807	466	341	799	449	350	21.9	17.8	31.1
Montgomery	397	175	222	340	131	209	25.5	16.2	39.6
Neshoba	394	205	189	401	215	186	18.7	13.6	32.9
Newton	370	206	164	378	185	193	18.3	13.5	27.9
Noxubee	449	82	367	438	74	364	25.0	14.4	29.3
Oktibbeha	803	344	459	741	309	432	26.2	19.7	34.5
Panola	882	282	600	879	288	591	28.2	20.4	34.6
Pearl River	647	500	147	578	426	152	22.3	20.8	28.0
Perry	25	20	5	198	143	55	23.2	24.6	20.3
Pike	814	393	421	766	331	435	22.0	16.6	29.0
Pontotoc	361	267	94	327	246	81	17.4	16.0	23.4
Prentiss	424	348	76	382	313	69	19.8	18.6	28.2
Quitman	790	215	575	687	174	513	29.8	18.9	37.0
Rankin	246	74	172	847	515	332	22.3	21.9	22.9
Scott	505	325	180	507	273	234	22.0	18.6	27.8
Sharkey	181	27	154	341	56	285	32.3	18.8	37.6
Simpson	390	233	157	456	242	214	20.5	16.0	30.2
Smith	45	16	29	322	176	146	20.3	14.6	38.4
Stone	148	89	59	141	80	61	21.3	16.4	34.6
Sunflower	1,214	193	1,021	1,344	237	1,107	27.9	16.0	33.2
Tallahatchie	670	105	565	735	148	587	29.5	16.6	36.7
Tate	344	3	341	564	150	414	26.3	16.2	34.0
Tippah	388	286	102	349	252	97	20.9	18.7	30.1
Tishomingo	221	207	14	311	290	21	18.7	18.2	29.6
Tunica	598	55	543	601	60	541	35.4	17.0	40.2
Union	315	247	68	367	297	70	17.6	17.0	21.3
Walthall	453	254	199	289	115	174	21.5	15.5	28.8
Warren	1,471	570	901	1,088	482	606	24.6	19.7	30.7
Washington	2,440	863	1,577	2,328	797	1,531	30.2	23.6	35.2
Wayne	482	306	176	413	228	185	25.5	22.9	29.7
Webster	118	55	63	197	116	81	19.0	15.4	28.4
Wilkinson	380	151	229	323	74	249	25.0	18.5	27.9
Winston	386	167	219	404	166	238	20.6	15.5	26.8
Yalobusha	288	115	173	287	113	174	22.5	16.0	30.8
Yazoo	830	243	587	872	262	610	26.0	19.0	31.0
Out-of-State	1,069	929	140	1,402	1,167	235	-	-	-

Table 2 - Page 3
Live Births - Mississippi - 1963

City**	Place of Occurrence			Place of Residence		
	Number		Non-white	Number		Non-white
	Total	White		Total	White	
Biloxi	2,499	2,248	251	1,499	1,295	204
Clarksdale	961	369	592	567	225	342
Cleveland	408	173	235	300	130	170
Columbus	980	538	442	814	406	408
Corinth	553	462	91	237	163	74
Greenville	1,414	702	712	1,196	511	685
Greenwood	824	419	405	604	254	350
Gulfport	1,484	1,097	387	810	553	257
Hattiesburg	1,534	1,159	375	832	496	336
Jackson	5,715	3,054	2,661	3,979	2,000	1,979
Laurel	2,082	1,003	1,079	630	307	323
McComb	609	353	256	294	156	138
Meridian	2,013	1,188	825	1,175	689	486
Natchez	1,484	694	790	627	255	372
Pascagoula	83	4	79	557	412	145
Tupelo	1,097	858	239	401	285	116
Vicksburg	1,453	569	884	748	315	433
Yazoo City	443	242	201	319	133	186
Total Urban #	36,716	21,007	15,709	21,693	11,782	9,911
Total Rural	20,933	6,054	14,879	35,623	15,041	20,582

* Births per 1,000 population.

** Cities having 10,000 population and over according to 1960 census.

Includes all incorporated towns of 2,500 population or more according to 1960 census.

TABLE 3
LIVE BIRTHS
MISSISSIPPI
1963

By Place of Residence of Mother, by Race and Attendant, and by Sex

County	TOTAL	WHITE			NONWHITE			SEX			
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female
STATE	<u>57,316</u>	<u>26,823</u>	<u>26,648</u>	<u>83</u>	<u>92</u>	<u>30,493</u>	<u>16,269</u>	<u>1,141</u>	<u>13,083</u>	<u>29,356</u>	<u>27,960</u>
Adams	1,046	464	464	0	0	582	572	0	10	531	515
Alcorn	525	421	419	0	2	104	70	1	33	279	246
Anite	337	108	108	0	0	229	76	2	151	174	163
Attala	481	201	200	0	1	280	80	6	194	282	199
Benton	187	55	54	1	0	132	18	0	114	95	92
Bolivar	1,802	351	347	2	2	1,451	623	82	746	922	880
Calhoun	360	205	204	0	1	155	103	2	50	175	185
Carroll	212	51	50	0	1	161	44	2	115	98	114
Chickasaw	412	215	212	1	2	197	116	20	61	205	207
Choctaw	173	94	94	0	0	79	32	2	45	88	85
Claiborne	243	46	46	0	0	197	71	3	123	121	122
Clarke	335	167	167	0	0	168	127	6	35	174	161
Clay	492	211	211	0	0	281	177	8	96	247	245
Coahoma	1,509	327	325	1	1	1,182	566	5	611	770	739
Copiah	668	233	233	0	0	435	362	18	55	332	336
Covington	313	163	163	0	0	150	115	5	30	136	177
DeSoto	750	265	265	0	0	485	284	4	197	362	388
Forrest	1,281	826	824	1	1	455	297	6	152	631	650
Franklin	175	77	77	0	0	98	79	0	19	88	87
George	316	248	246	1	1	68	53	0	15	163	153
Greene	178	125	124	0	1	53	32	1	20	89	89
Grenada	488	174	174	0	0	314	52	77	185	253	235
Hancock	344	272	268	3	1	72	34	0	38	179	165
Harrison	3,751	3,029	3,012	2	15	722	428	44	250	1,916	1,835

Table 3 - Page 2
Live Births - Mississippi - 1963

County	TOTAL	WHITE			NONWHITE			SEX			
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female
Hinds	5,249	2,492	2,489	3	0	2,757	2,215	62	480	2,663	2,586
Holmes	756	113	112	0	1	643	100	7	536	401	355
Humphreys	574	95	94	0	1	479	94	2	383	309	265
Issaquena	80	15	15	0	0	65	30	0	35	47	33
Itawamba	254	237	233	2	2	17	16	1	0	136	118
Jackson	1,822	1,370	1,361	2	7	452	243	77	132	929	893
Jasper	399	148	148	0	0	251	200	7	44	203	196
Jefferson	247	41	41	0	0	206	102	0	104	116	131
Jeff Davis	268	117	117	0	0	151	82	1	68	145	123
Jones	1,263	788	784	3	1	475	451	4	20	645	618
Kemper	251	58	57	0	1	193	63	2	128	131	120
Lafayette	514	259	256	0	3	255	50	12	193	248	266
Lamar	295	236	236	0	0	57	52	2	3	151	142
Lauderdale	1,698	999	999	0	0	699	578	15	105	866	832
Lawrence	210	109	108	0	1	101	49	4	48	109	101
Leake	333	137	135	2	0	196	102	4	90	172	161
Lee	1,062	717	714	2	1	345	230	26	89	562	500
Leflore	1,422	374	368	4	2	1,048	151	24	873	774	648
Lincoln	596	323	322	0	1	273	241	3	29	308	288
Lowndes	1,476	821	814	1	6	655	276	7	372	721	755
Madison	903	177	177	0	0	726	216	14	496	490	413
Marion	525	291	286	4	1	234	76	45	113	265	260
Marshall	743	165	164	0	1	578	187	27	364	383	360
Monroe	799	449	447	1	1	350	166	21	163	412	387
Montgomery	340	131	131	0	0	209	85	6	118	164	176
Neshoba	401	215	215	0	0	186	133	0	53	213	188
Newton	378	182	182	2	0	193	125	53	15	197	181
Noxubee	438	74	73	0	1	364	129	6	229	220	218
Oktibbeha	744	369	369	1	0	432	215	6	211	355	386
Panola	875	332	287	1	1	391	363	76	147	467	412

Table 3 - Page 3
Live Births - Mississippi - 1963

County	TOTAL	WHITE			NONWHITE			SEX			
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female
Pearl River	578	426	423	3	0	152	131	0	21	296	282
Perry	198	143	142	0	1	55	49	1	5	89	109
Pike	766	331	328	0	3	435	266	1	168	395	371
Pontotoc	327	246	244	2	0	81	41	4	36	176	151
Prentiss	382	313	313	0	0	69	52	1	16	206	176
Quitman	687	174	171	3	0	513	353	14	146	350	337
Rankin	847	515	514	0	1	332	267	1	64	439	408
Scott	507	273	273	0	0	234	146	1	87	259	248
Sharkey	341	56	55	0	1	285	170	3	112	186	155
Simpson	456	242	242	0	0	214	176	4	34	218	238
Smith	322	176	176	0	0	146	122	11	13	175	147
Stone	141	80	79	1	0	61	16	0	45	83	58
Sunflower	1,344	237	230	5	2	1,107	362	103	642	665	679
Tallahatchie	735	148	142	5	1	587	107	24	456	396	339
Tate	564	150	147	1	2	414	72	48	294	281	283
Tippah	349	252	243	7	2	97	58	23	16	190	159
Tishomingo	311	290	288	2	0	21	21	0	0	165	146
Tunica	601	60	58	0	2	541	197	12	332	312	289
Union	367	297	285	4	8	70	25	3	42	189	178
Walton	289	115	111	1	3	174	75	1	98	150	139
Warren	1,088	482	481	0	1	606	528	2	76	565	523
Washington	2,328	797	795	1	1	1,531	1,033	32	466	1,207	1,121
Wayne	413	228	227	1	0	185	136	3	46	210	203
Webster	197	116	111	5	0	81	43	9	29	112	85
Wilkinson	323	74	74	0	0	249	81	1	167	156	167
Winston	404	166	163	1	2	238	89	1	148	195	209
Yalobusha	287	113	113	0	0	174	115	28	31	150	137
Yazoo	872	262	259	1	2	610	102	2	506	429	443

Table 3 - Page 4
Live Births - Mississippi - 1963

City*	TOTAL	WHITE			NONWHITE			SEX			
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female
Biloxi	1,499	1,295	1,290	1	4	204	159	13	32	783	716
Clarksdale	567	225	224	1	0	342	284	3	55	292	275
Cleveland	300	130	130	0	0	170	79	22	69	153	147
Columbus	814	406	400	1	5	408	170	5	233	394	420
Corinth	237	163	163	0	0	74	53	0	21	121	116
Greenville	1,196	511	510	1	0	685	485	16	184	611	585
Greenwood	604	254	253	0	1	350	90	4	256	328	276
Gulfport	810	553	549	1	3	257	155	25	77	416	394
Hattiesburg	832	496	495	0	1	336	227	4	105	395	437
Jackson	3,979	2,000	1,999	1	0	1,979	1,717	55	207	2,049	1,930
Laurel	630	307	306	1	0	323	318	2	3	301	329
McComb	294	156	156	0	0	138	98	1	39	151	143
Meridian	1,175	689	689	0	0	486	432	12	42	589	586
Natchez	627	255	255	0	0	372	368	0	4	324	303
Pascagoula	557	412	409	0	3	145	71	26	48	278	279
Tupelo	401	285	285	0	0	116	91	5	20	215	186
Vicksburg	748	315	315	0	0	433	379	1	53	401	347
Yazoo City	319	133	131	1	1	186	49	0	137	160	159

* Cities having 10,000 population and over according to 1960 census.

TABLE 4
FETAL DEATHS
MISSISSIPPI
1963

By Place of Occurrence; by Place of Residence of Mother, by Race and Attendant, and by Sex

Table 4 - Page 2
Fetal Deaths - Mississippi - 1963

County	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total	Phys. in Hosp.	Other	Total	Phys. in Hosp.	Other	Male	Female	Unde-termined
Hinds	533	451	293	255	37	1	158	135	8	78	45
Holmes	33	34	3	0	0	31	12	4	15	18	9
Humphreys	14	22	6	6	0	16	8	0	8	10	5
Issaquena	1	2	1	1	0	0	0	0	1	1	1
Itawamba	1	5	5	5	0	0	0	0	2	3	0
Jackson	81	101	71	68	3	0	30	26	4	0	22
Jasper	9	18	11	11	0	0	7	7	0	0	3
Jefferson	2	10	2	2	0	0	8	6	0	2	5
Jeff Davis	12	17	3	3	0	0	14	10	1	3	10
Jones	99	59	30	22	8	0	29	25	3	1	16
Kemper	8	11	0	0	0	0	11	7	0	4	7
Lafayette	22	21	3	3	0	0	18	5	1	12	10
Lamar	2	9	6	6	0	0	3	3	0	0	3
Lauderdale	61	44	13	13	0	0	31	22	4	5	23
Lawrence	4	12	2	1	1	0	10	7	0	3	7
Leake	7	17	3	3	0	0	14	12	1	1	9
Lee	33	25	12	12	0	0	13	11	1	1	15
Leflore	135	129	47	43	4	0	82	38	8	36	35
Lincoln	20	28	9	9	0	0	19	16	1	2	14
Lowndes	58	56	27	27	0	0	29	19	2	8	24
Madison	28	37	8	8	0	0	29	21	0	8	11
Marion	15	16	7	7	0	0	9	5	3	1	8
Marshall	43	38	2	2	0	0	36	17	1	18	16
Monroe	28	32	14	13	1	0	18	15	2	1	7
Montgomery	21	21	7	7	0	0	14	8	1	5	8
Neshoba	12	18	5	5	0	0	13	13	0	0	6
Newton	33	31	14	13	1	0	17	16	1	0	6
Noxubee	16	15	2	2	0	0	13	10	0	3	9
Oktibbeha	19	23	7	7	0	0	16	5	0	11	7
Panola	22	30	6	6	0	0	24	14	6	4	17
											0

Table 4 - Page 3
Fetal Deaths - Mississippi - 1963

County	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total	Phys. in Hosp.	Other	Total	Phys. in Hosp.	Other	Male	Female	Unde-termined
Pearl River	8	9	6	0	0	3	2	1	0	4	1
Perry	1	3	1	0	0	17	9	0	0	2	0
Pike	25	25	8	0	0	0	2	1	0	1	0
Pontotoc	3	7	4	0	0	3	2	0	1	2	5
Prentiss	13	12	9	0	0	3	2	0	1	2	2
Quitman	19	24	2	1	0	22	15	2	5	5	3
Rankin	11	36	23	20	3	0	13	10	0	3	10
Scott	11	14	3	3	0	0	11	7	0	4	3
Sharkey	6	17	4	4	0	0	13	9	3	1	4
Simpson	14	17	13	13	0	0	4	3	0	1	7
Smith	2	12	4	4	0	0	8	5	2	1	6
Stone	4	4	3	3	0	0	1	0	1	5	3
Sunflower	55	63	7	7	0	0	56	25	5	26	25
Tallahatchie	21	25	3	3	0	0	22	7	2	13	15
Tate	9	17	1	1	0	0	16	7	1	8	6
Tippah	33	27	19	18	1	0	8	6	2	0	5
Tishomingo	3	3	3	3	0	0	0	0	0	1	0
Tunica	21	20	0	0	0	0	20	8	2	10	8
Union	28	25	20	17	3	0	5	4	0	1	9
Walthall	24	9	4	4	0	0	5	3	0	2	5
Warren	112	63	38	38	0	0	25	23	0	2	2
Washington	232	222	124	123	1	0	98	77	5	16	48
Wayne	7	7	2	2	0	0	5	3	1	4	1
Webster	7	8	4	4	0	0	4	4	0	0	5
Wilkinson	17	15	1	1	0	0	14	8	1	5	7
Winston	4	8	0	0	0	0	8	6	0	2	6
Yalobusha	7	8	4	4	0	0	4	4	0	0	1
Yazoo	27	32	7	6	1	0	25	9	1	15	13

Table 4 - Page 4
Fetal Deaths - Mississippi - 1963

City*	Total by Occurrence	Total by Residence	WHITE			NONWHITE			SEX		
			Total	Phys. in Hosp.	Other	Total	Phys. in Hosp.	Other	Male	Female	Unde-termined
Biloxi	212	150	135	135	0	15	14	0	1	10	18
Clarksdale	56	32	12	12	0	20	18	0	2	8	12
Cleveland	11	9	3	3	0	6	5	0	1	5	4
Columbus	43	36	16	16	0	20	13	2	5	16	15
Corinth	8	3	0	0	0	3	2	0	1	3	0
Greenville	179	144	95	95	0	49	45	1	3	23	11
Greenwood	104	70	36	33	3	0	34	21	1	12	14
Gulfport	43	43	20	20	0	0	23	14	2	7	17
Hattiesburg	32	24	10	10	0	0	14	9	2	3	5
Jackson	523	399	268	231	37	0	131	117	7	7	61
Laurel	94	31	16	12	4	0	15	14	1	0	10
McComb	15	9	3	3	0	0	6	2	0	4	2
Meridian	59	36	13	13	0	0	23	16	4	3	17
Natchez	60	19	8	8	0	0	11	11	0	0	12
Pascagoula	3	47	32	31	1	0	15	13	2	0	11
Tupelo	24	10	5	5	0	0	5	3	1	1	8
Vicksburg	112	46	30	30	0	0	16	14	0	2	17
Yazoo City	16	15	3	3	0	0	12	6	0	6	4

* Cities having 10,000 population and over according to 1960 census.

TABLE 5
DEATHS
MISSISSIPPI
1963

By Cause of Death, by Race, Residence Data

International List Number*	Cause of Death**	Total	White	Non-white
<u>001-999</u>	<u>TOTAL ALL CAUSES</u>	<u>23,339</u>	<u>12,118</u>	<u>11,221</u>
<u>001-019</u>	<u>Tuberculosis, all forms</u>	<u>113</u>	<u>40</u>	<u>73</u>
002	Pulmonary	101	39	62
003	Pleural	4	0	4
008	Unspecified site	1	0	1
010	Meninges and central nervous system	4	0	4
011	Intestines, peritoneum, and mesenteric glands	1	0	1
016	Genito-urinary system	1	1	0
019	Disseminated	1	0	1
<u>020-029</u>	<u>Syphilis and its sequelae</u>	<u>34</u>	<u>9</u>	<u>25</u>
020	Congenital syphilis	3	1	2
022	Aneurysm of aorta	15	7	8
023	Other cardiovascular syphilis	10	0	10
025	General paralysis of insane	1	1	0
026	Other syphilis of central nervous system	3	0	3
029	Syphilis, unqualified	2	0	2
<u>045-048</u>	<u>Dysentery, all forms</u>	<u>14</u>	<u>3</u>	<u>11</u>
045	Bacillary	5	1	4
046	Amebiasis	6	2	4
048	Unspecified forms	3	0	3
049	Food poisoning (infection and intoxication)	2	0	2
<u>050,051</u>	<u>Scarlet fever and streptococcal sore throat</u>	<u>4</u>	<u>1</u>	<u>3</u>
050	Scarlet fever	1	0	1
051	Streptococcal sore throat	3	1	2
052	Erysipelas	1	0	1
053	Septicemia and pyemia	32	17	15
056	Whooping cough	5	1	4
057	Meningococcal infections	6	4	2
061	Tetanus	5	1	4
063	Gas gangrene	1	0	1
064	Other bacterial diseases	2	1	1
072	Leptospirosis	1	0	1
080	Acute poliomyelitis	4	4	0
082	Acute infectious encephalitis	10	5	5
083	Late effects of acute infectious encephalitis	1	1	0
085	Measles	7	3	4
087	Chickenpox	1	1	0
092	Infectious hepatitis	5	4	1
096	Other diseases attributable to viruses	7	4	3
104	Tick-borne typhus	1	0	1
130	Infestation with worms of other, mixed, and unspecified type	2	0	2
132	Actinomycosis	1	0	1
133	Coccidioidomycosis	1	0	1

Table 5 - Page 2
Deaths - Mississippi - 1963

International List Number*	Cause of Death**	Total	White	Non-white
134	Other fungus infections	5	3	2
138	Other infective and parasitic diseases	3	2	1
<u>140-205</u>	<u>Malignant neoplasms</u>	<u>2,945</u>	<u>1,859</u>	<u>1,086</u>
<u>140-148</u>	<u>Buccal cavity and pharynx</u>	<u>52</u>	<u>42</u>	<u>10</u>
140	Lip	2	2	0
141	Tongue	9	7	2
142	Salivary gland	4	3	1
143	Floor of mouth	3	2	1
144	Other parts of mouth and mouth unspecified	6	5	1
145	Oral mesopharynx	3	3	0
146	Nasopharynx	9	8	1
148	Pharynx, unspecified	16	12	4
<u>150-159</u>	<u>Digestive organs and peritoneum</u>	<u>909</u>	<u>543</u>	<u>366</u>
150	Esophagus	41	23	18
151	Stomach	255	115	140
152	Small intestine, including duodenum	6	4	2
153	Large intestine, except rectum	249	173	76
154	Rectum	54	29	25
155	Biliary passages and liver (stated to be primary site)	40	31	9
156	Liver (secondary and unspecified)	61	31	30
157	Pancreas	172	118	54
158	Peritoneum	8	4	4
159	Unspecified digestive organs	23	15	8
<u>160-165</u>	<u>Respiratory system</u>	<u>499</u>	<u>375</u>	<u>124</u>
160	Nose, nasal cavities, middle ear, and accessory sinuses	4	4	0
161	Larynx	31	25	6
162	Bronchus and trachea, and lung specified as primary	190	143	47
163	Lung, unspecified as to whether primary or secondary	253	184	69
164	Mediastinum	2	2	0
165	Thoracic organs (secondary)	19	17	2
<u>170-181</u>	<u>Breast and genito-urinary organs</u>	<u>823</u>	<u>433</u>	<u>390</u>
170	Breast	193	123	70
171	Cervix uteri	158	72	86
172	Corpus uteri	13	5	8
173	Other parts of uterus, including choriocarcinoma	1	0	1
174	Uterus, unspecified	65	26	39
175	Ovary, Fallopian tube, and broad ligament	44	26	18
176	Other and unspecified female genital organs	9	4	5
177	Prostate	222	110	112
178	Testis	7	4	3
179	Other and unspecified male genital organs	4	2	2
180	Kidney	28	19	9
181	Bladder and other urinary organs	79	42	37

Table 5 - Page 3
 Deaths - Mississippi - 1963

International List Number*	Cause of Death**	Total	White	Non-white
<u>190-199</u>	<u>Other and unspecified sites</u>	<u>364</u>	<u>253</u>	<u>111</u>
190	Melanoma of skin	17	15	2
191	Skin, other	40	35	5
192	Eye	4	4	0
193	Brain and other parts of nervous system	88	70	18
194	Thyroid gland	7	4	3
196	Bone	24	15	9
197	Connective tissue	10	8	2
198	Lymph nodes (secondary and unspecified)	8	5	3
199	Other and unspecified sites	166	97	69
<u>200-205</u>	<u>Lymphatic and hematopoietic tissues</u>	<u>298</u>	<u>213</u>	<u>85</u>
200	Lymphosarcoma and reticulosarcoma	57	48	9
201	Hodgkin's disease	34	24	10
202	Other forms of lymphoma (reticulososis)	10	10	0
203	Multiple myeloma (plasmacytoma)	31	19	12
204	Leukemia and aleukemia	164	110	54
205	Mycosis fungoides	2	2	0
<u>210-229</u>	<u>Benign neoplasms</u>	<u>26</u>	<u>13</u>	<u>13</u>
211	Other parts of digestive system	1	1	0
214	Uterine fibromyoma	7	2	5
215	Uterus, other	1	0	1
216	Ovary	2	0	2
219	Kidney and other urinary organs	1	1	0
223	Brain and other parts of nervous system	7	4	3
224	Endocrine glands	4	2	2
228	Hemangioma and lymphangioma	3	3	0
<u>230-239</u>	<u>Neoplasms of unspecified nature</u>	<u>19</u>	<u>8</u>	<u>11</u>
237	Brain and other parts of nervous system	13	7	6
239	Other and unspecified organs	6	1	5
240	Hay fever	1	1	0
241	Asthma	45	31	14
245	Other allergic disorders	2	1	1
250	Simple goiter	3	1	2
252	Thyrotoxicosis with or without goiter	6	1	5
253	Myxedema and cretinism	2	2	0
254	Other diseases of thyroid gland	1	1	0
260	Diabetes mellitus	326	132	194
270	Disorders of pancreatic internal secretion other than diabetes mellitus	2	1	1
272	Diseases of pituitary gland	5	2	3
273	Diseases of thymus gland	1	1	0
274	Diseases of adrenal glands	2	2	0
280	Beriberi	1	1	0
281	Pellagra	1	0	1
286	Other avitaminoses and nutritional deficiency states	24	8	16
287	Obesity, not specified as of endocrine origin	7	2	5

Table 5 - Page 4
 Deaths - Mississippi - 1963

International List Number*	Cause of Death**	Total	White	Non-white
288	Gout	1	0	1
289	Other metabolic diseases	13	9	4
<u>290-293</u>	<u>Anemias</u>	<u>60</u>	<u>16</u>	<u>44</u>
290	Pernicious and other hyperchromic	6	2	4
291	Iron deficiency (hypochromic)	2	0	2
292	Other specified type	30	6	24
293	Unspecified type	22	8	14
294	Polycythemia	3	2	1
295	Hemophilia	1	1	0
296	Purpura and other hemorrhagic conditions	15	10	5
297	Agranulocytosis	2	1	1
298	Diseases of spleen	1	1	0
299	Other diseases of blood and blood-forming organs	1	1	0
304	Senile psychosis	9	7	2
305	Presenile psychosis	1	1	0
307	Alcoholic psychosis	1	1	0
309	Other and unspecified psychoses	2	2	0
315	Psychoneurosis with somatic symptoms (somatization reaction) affecting circulatory system	1	0	1
322	Alcoholism	11	7	4
323	Other drug addiction	1	1	0
325	Mental deficiency	2	2	0
<u>330-334</u>	<u>Vascular lesions affecting central nervous system</u>	<u>3,154</u>	<u>1,650</u>	<u>1,504</u>
330	Subarachnoid hemorrhage	111	58	53
331	Cerebral hemorrhage	1,954	965	989
332	Cerebral embolism and thrombosis	705	446	259
334	Other and ill-defined	384	181	203
340	Meningitis, except meningococcal and tuberculous	50	13	37
341	Phlebitis and thrombophlebitis of intracranial venous sinuses	2	1	1
342	Intracranial and intraspinal abscess	3	0	3
343	Encephalitis, myelitis, and encephalomyelitis (except acute infectious)	7	2	5
344	Late effects of intracranial abscess or pyogenic infection	9	3	6
345	Multiple sclerosis	6	3	3
350	Paralysis agitans	27	25	2
351	Cerebral spastic infantile paralysis	10	4	6
352	Other cerebral paralysis	10	4	6
353	Epilepsy	35	6	29
355	Other diseases of brain	13	11	2
356	Motor neurone disease and muscular atrophy	8	7	1
357	Other diseases of spinal cord	1	0	1
364	Polyneuritis and polyradiculitis	3	3	0
385	Cataract	2	1	1
392	Otitis media with mastoiditis	1	0	1
393	Mastoiditis without mention of otitis media	1	1	0

Table 5 - Page 5
 Deaths - Mississippi - 1963

International List Number*	Cause of Death**	Total	White	Non-white
<u>400-443</u>	<u>Diseases of the heart</u>	<u>7,300</u>	<u>4,396</u>	<u>2,904</u>
401	Rheumatic fever with heart involvement	10	0	10
410	Diseases of mitral valve	37	17	20
411	Diseases of aortic valve specified as rheumatic	7	4	3
414	Other endocarditis specified as rheumatic	5	2	3
415	Other myocarditis specified as rheumatic	1	0	1
416	Other heart disease specified as rheumatic	43	18	25
420	Arteriosclerotic heart disease, including coronary disease	4,751	3,361	1,390
421	Chronic endocarditis not specified as rheumatic	77	39	38
422	Other myocardial degeneration	531	291	240
430	Acute and subacute endocarditis	6	4	2
431	Acute myocarditis not specified as rheumatic	12	6	6
432	Acute pericarditis specified as nonrheumatic	1	0	1
433	Functional disease of heart	101	75	26
434	Other and unspecified diseases of heart	511	210	301
441	Essential malignant hypertensive heart disease	12	0	12
442	Hypertensive heart disease with arteriolar nephrosclerosis	214	65	149
443	Other and unspecified hypertensive heart disease	981	304	677
<u>444-447</u>	<u>Other hypertensive disease</u>	<u>203</u>	<u>77</u>	<u>126</u>
444	Essential benign hypertension	63	21	42
445	Essential malignant hypertension	59	13	46
446	Hypertension with arteriolar nephrosclerosis	79	42	37
447	Other hypertensive disease without mention of heart	2	1	1
450	General arteriosclerosis	364	250	114
<u>451-468</u>	<u>Other diseases of circulatory system</u>	<u>227</u>	<u>134</u>	<u>93</u>
451	Aortic aneurysm, nonsyphilitic, and dissecting aneurysm	77	58	19
452	Other aneurysm, except of heart and aorta	4	1	3
453	Peripheral vascular disease	2	1	1
454	Arterial embolism and thrombosis	4	4	0
455	Gangrene of unspecified cause	5	3	2
456	Other diseases of arteries	10	5	5
460	Varicose veins of lower extremities	6	3	3
461	Hemorrhoids	2	0	2
462	Varicose veins of other specified sites	5	3	2
463	Phlebitis and thrombophlebitis of lower extremities	12	8	4
464	Phlebitis and thrombophlebitis of other sites	13	7	6
465	Pulmonary embolism and infarction	44	21	23
466	Other venous embolism and thrombosis	16	8	8
467	Other diseases of circulatory system	27	12	15
470	Acute nasopharyngitis (common cold)	13	1	12

Table 5 - Page 6
 Deaths - Mississippi - 1963

International List Number*	Cause of Death**	Total	White	Non-white
472	Acute pharyngitis	5	3	2
474	Acute laryngitis and tracheitis	1	0	1
475	Acute upper respiratory infection of multiple or unspecified sites	11	2	9
<u>480-483</u>	<u>Influenza</u>	<u>253</u>	<u>75</u>	<u>178</u>
480	With pneumonia	144	35	109
481	With other respiratory manifestations, and influenza unqualified	99	37	62
482	With digestive manifestations, but without respiratory symptoms	2	0	2
483	With nervous manifestations, but without digestive or respiratory symptoms	8	3	5
<u>490-493</u>	<u>Pneumonia (excluding pneumonia of newborn)</u>	<u>860</u>	<u>345</u>	<u>515</u>
490	Lobar	161	49	112
491	Broncho	351	154	197
492	Primary atypical	57	27	30
493	Other and unspecified	291	115	176
<u>500-502</u>	<u>Bronchitis</u>	<u>58</u>	<u>43</u>	<u>15</u>
500	Acute	9	5	4
501	Unqualified	13	7	6
502	Chronic	36	31	5
517	Other diseases of upper respiratory tract	5	0	5
518	Empyema	4	2	2
519	Pleurisy	5	2	3
521	Abscess of lung	3	1	2
522	Pulmonary congestion and hypostasis	25	17	8
525	Other chronic interstitial pneumonia	27	21	6
526	Bronchiectasis	19	17	2
527	Other diseases of lung and pleural cavity	188	152	36
537	Diseases of salivary glands	1	1	0
539	Diseases of esophagus	3	2	1
<u>540,541</u>	<u>Ulcer of stomach and duodenum</u>	<u>68</u>	<u>46</u>	<u>22</u>
540	Stomach	33	24	9
541	Duodenum	35	22	13
542	Gastrojejunal ulcer	1	1	0
543	Gastritis and duodenitis	5	2	3
544	Disorders of function of stomach	4	2	2
545	Other diseases of stomach and duodenum	2	2	0
<u>550-553</u>	<u>Appendicitis</u>	<u>21</u>	<u>10</u>	<u>11</u>
550	Acute	19	8	11
551	Unqualified	2	2	0
<u>560,561,570</u>	<u>Hernia and intestinal obstruction</u>	<u>117</u>	<u>60</u>	<u>57</u>
560	Hernia of abdominal cavity without mention of obstruction	11	3	8
561	Hernia of abdominal cavity with obstruction	27	14	13
570	Intestinal obstruction without mention of hernia	79	43	36

Table 5 - Page 7
 Deaths - Mississippi - 1963

International List Number*	Cause of Death**	Total	White	Non-white
571	Gastritis and colitis, except ulcerative, age 4 weeks and over	177	24	153
572	Chronic enteritis and ulcerative colitis	22	17	5
575	Abscess of anal and rectal regions	1	0	1
576	Peritonitis	11	6	5
578	Other diseases of intestines and peritoneum	46	27	19
580	Acute and subacute yellow atrophy of liver	7	4	3
581	Cirrhosis of liver	105	86	19
583	Other diseases of liver	12	7	5
<u>584,585</u>	<u>Cholelithiasis, cholecystitis, and cholangitis</u>	<u>39</u>	<u>29</u>	<u>10</u>
584	Cholelithiasis	18	16	2
585	Cholecystitis and cholangitis, without mention of calculi	21	13	8
586	Other diseases of gallbladder and biliary ducts	6	6	0
587	Diseases of pancreas	15	11	4
<u>590,591</u>	<u>Acute nephritis and nephritis with edema, including nephrosis</u>	<u>39</u>	<u>13</u>	<u>26</u>
590	Acute	23	8	15
591	With edema, including nephrosis	16	5	11
<u>592-594</u>	<u>Chronic and unspecified nephritis and other renal sclerosis</u>	<u>222</u>	<u>90</u>	<u>132</u>
592	Chronic nephritis	166	65	101
593	Nephritis not specified as acute or chronic	56	25	31
600	Infections of kidney	170	79	91
601	Hydronephrosis	5	3	2
602	Calculi of kidney and ureter	15	11	4
603	Other diseases of kidney and ureter	18	9	9
604	Calculi of other parts of urinary system	2	1	1
606	Other diseases of bladder	4	1	3
607	Urethritis (nonvenereal)	1	0	1
608	Stricture of urethra	4	0	4
609	Other diseases of urethra	12	7	5
610	Hyperplasia of prostate	66	24	42
611	Prostatitis	5	1	4
612	Other diseases of prostate	2	2	0
617	Other diseases of male genital organs	1	0	1
622	Acute salpingitis and oophoritis	1	0	1
623	Chronic salpingitis and oophoritis	1	0	1
624	Salpingitis and oophoritis, unqualified	1	0	1
625	Other diseases of ovary and Fallopian tube	1	0	1
626	Diseases of parametrium and pelvic peritoneum (female)	7	1	6
631	Uterovaginal prolapse	1	1	0
633	Other diseases of uterus	3	0	3
<u>640-689</u>	<u>Deliveries and complications of pregnancy, childbirth, and the puerperium</u>	<u>44</u>	<u>9</u>	<u>35</u>
642	Toxemias of pregnancy	10	0	10
645	Ectopic pregnancy	2	0	2

Table 5 - Page 8
 Deaths - Mississippi - 1963

International List Number*	Cause of Death**	Total	White	Non-white
647	Pregnancy with malposition of fetus in uterus	1	0	1
648	Other complications arising from pregnancy	3	2	1
651	Abortion with sepsis	1	0	1
660	Delivery without mention of complication	4	2	2
670	Delivery complicated by placenta previa or antepartum hemorrhage	5	0	5
671	Delivery complicated by retained placenta	1	0	1
672	Delivery complicated by other postpartum hemorrhage	3	1	2
674	Delivery complicated by disproportion or mal-position of fetus	1	0	1
675	Delivery complicated by prolonged labor of other origin	1	0	1
677	Delivery with other trauma	3	1	2
681	Sepsis of childbirth and the puerperium	3	1	2
682	Puerperal phlebitis and thrombosis	1	0	1
684	Puerperal pulmonary embolism	2	1	1
686	Other forms of puerperal toxemia	2	1	1
688	Other and unspecified complications of the puerperium	1	0	1
690	Boil and carbuncle	3	2	1
692	Other cellulitis and abscess without mention of lymphangitis	6	2	4
698	Other local infections of skin and subcutaneous tissue	1	1	0
704	Pemphigus	1	0	1
705	Erythematous conditions	6	3	3
710	Other hypertrophic and atrophic conditions of skin	8	4	4
715	Chronic ulcer of skin	4	1	3
716	Other diseases of skin	1	0	1
722	Rheumatoid arthritis and allied conditions	16	12	4
723	Osteo-arthritis (arthrosis) and allied conditions	2	2	0
725	Arthritis, unspecified	6	4	2
727	Rheumatism, unspecified	2	1	1
730	Osteomyelitis and periostitis	4	1	3
731	Osteitis deformans	1	0	1
733	Other diseases of bone	4	3	1
744	Other diseases of muscle, tendon, and fascia	8	7	1
745	Curvature of spine	1	1	0
<u>750-759</u>	<u>Congenital malformations</u>	<u>256</u>	<u>147</u>	<u>109</u>
750	Monstrosity	8	7	1
751	Spina bifida and meningocele	16	13	3
752	Congenital hydrocephalus	12	7	5
753	Other, nervous system and sense organs	6	3	3
754	Circulatory system	134	70	64
755	Cleft palate and harelip	1	1	0
756	Digestive system	19	8	11
757	Genito-urinary system	18	13	5

Table 5 - Page 9
 Deaths - Mississippi - 1963

International List Number*	Cause of Death**	Total	White	Non-white
758	Bone and joint	7	5	2
759	Other and unspecified, not elsewhere classified	35	20	15
<u>760-776</u>	<u>Certain diseases of early infancy</u>	<u>1,262</u>	<u>389</u>	<u>873</u>
760	Intracranial and spinal injury at birth	29	11	18
761	Other birth injury	71	37	34
762	Postnatal asphyxia and atelectasis	229	104	125
763	Pneumonia of newborn	93	24	69
764	Diarrhea of newborn	32	4	28
766	Pemphigus neonatorum	1	0	1
767	Umbilical sepsis	3	0	3
768	Other sepsis of newborn	18	4	14
769	Neonatal disorders arising from certain diseases of the mother during pregnancy	13	1	12
770	Hemolytic disease of newborn (erythroblastosis)	15	8	7
771	Hemorrhagic disease of newborn	14	4	10
772	Nutritional maladjustment	31	1	30
773	Ill-defined diseases peculiar to early infancy	208	85	123
774	Immaturity with mention of any other subsidiary condition	14	4	10
776	Immaturity, unqualified	491	102	389
<u>780-795</u>	<u>Symptoms, senility, and ill-defined conditions</u>	<u>1,770</u>	<u>435</u>	<u>1,335</u>
780	Certain symptoms referable to nervous system and special senses	6	3	3
782	Symptoms referable to cardiovascular and lymphatic system	53	25	28
783	Symptoms referable to respiratory system	7	4	3
784	Symptoms referable to upper gastro-intestinal tract	8	3	5
785	Symptoms referable to abdomen and lower gastro-intestinal tract	12	1	11
786	Symptoms referable to genito-urinary system	5	0	5
787	Symptoms referable to limbs and back	2	0	2
788	Other general symptoms	8	4	4
790	Nervousness and debility	1	0	1
792	Uremia	18	10	8
794	Senility without mention of psychosis	71	14	57
795	Ill-defined and unknown causes	1,579	371	1,208
<u>800-962</u>	<u>Accidents</u>	<u>1,571</u>	<u>851</u>	<u>720</u>
<u>800-802</u>	<u>Railway</u>	<u>16</u>	<u>9</u>	<u>7</u>
800	Involving railroad employee	2	2	0
802	Involving other and unspecified person	14	7	7
<u>810-835</u>	<u>Motor vehicle</u>	<u>675</u>	<u>435</u>	<u>240</u>
810	Traffic, collision with railway train	27	19	8
812	Traffic, to pedestrian	111	48	63
813	Traffic, to pedal cyclist	12	10	2
814	Traffic, to rider or passenger of motorcycle in collision with nonmotor vehicle or object	1	1	0
815	Traffic, to rider or passenger of motorcycle in collision with other motor vehicle	6	6	0

Table 5 - Page 10
Deaths - Mississippi - 1963

International List Number*	Cause of Death**	Total	White	Non-white
816	Traffic, other involving two or more motor vehicles	243	177	66
818	Traffic, collision with animal or animal-drawn vehicle	1	1	0
819	Traffic, collision with fixed or unspecified object	30	21	9
821	Traffic, to rider of motorcycle without antecedent collision	4	4	0
822	Traffic, overturning in roadway	19	9	10
823	Traffic, running off roadway	159	105	54
824	Traffic, other noncollision	13	5	8
825	Traffic, unspecified nature	28	16	12
830	Nontraffic, to pedestrian	10	5	5
835	Nontraffic, other and unspecified nature	11	8	3
<u>840-845</u>	<u>Other road vehicle</u>	1	0	1
845	Other nonmotor	1	0	1
<u>850-858</u>	<u>Water transport</u>	32	21	11
850	Submersion of occupant of small boat	27	17	10
851	Other injury by submersion	3	3	0
853	Other falls from one level to another	1	0	1
857	Other specified	1	1	0
<u>860-866</u>	<u>Aircraft</u>	19	19	0
860	To personnel in military aircraft	10	10	0
863	To occupant of other specified aircraft	1	1	0
866	Other and unspecified	8	8	0
<u>870-895</u>	<u>Poisoning</u>	33	15	18
871	Barbituric acid and derivatives	3	2	1
872	Aspirin and salicylates	3	2	1
874	Other analgesic and soporific drugs	1	1	0
878	Other and unspecified drugs	3	2	1
881	Petroleum products	4	0	4
883	Corrosive aromatics, acids, and caustic alkalies	1	1	0
884	Mercury and its compounds	1	1	0
885	Lead and its compounds	1	0	1
886	Arsenic and antimony, and their compounds	1	1	0
888	Other and unspecified solid and liquid substances	2	0	2
890	Utility (illuminating) gas	1	0	1
891	Motor vehicle exhaust gas	7	3	4
892	Other carbon monoxide gas	2	2	0
894	Other specified gases and vapors	1	0	1
895	Unspecified gases and vapors	2	0	2
<u>900-904</u>	<u>Falls</u>	120	88	32
900	On stairs	4	4	0
901	From ladders	2	1	1
902	Other, from one level to another	16	10	6
903	On same level	28	23	5
904	Unspecified	70	50	20
910	Blow from falling or projected object or missile	29	14	15
911	Caused by vehicle (not classified elsewhere)	1	0	1
912	Caused by machinery	38	19	19

Table 5 - Page 11
 Deaths - Mississippi - 1963

International List Number*	Cause of Death**	Total	White	Non-white
913	Caused by cutting and piercing instruments	2	0	2
914	Caused by electric current	19	13	6
915	Caused by explosion of pressure vessel	2	2	0
916	Caused by fire and explosion of combustible material	223	69	154
917	Caused by hot substance, corrosive liquid, and steam	6	3	3
919	Caused by firearm	83	47	36
921	Inhalation and ingestion of food causing obstruction or suffocation	19	6	13
922	Inhalation and ingestion of other object causing obstruction or suffocation	2	0	2
923	Foreign body entering other orifice	10	5	5
924	Mechanical suffocation in bed and cradle	37	10	27
925	Mechanical suffocation in other and unspecified circumstances	12	3	9
926	Lack of care of infants under 1 year of age	5	0	5
928	Other, caused by animals	2	2	0
929	Drowning and submersion	91	31	60
931	Excessive heat and insolation	2	1	1
932	Excessive cold	22	4	18
933	Hunger, thirst, and exposure	7	3	4
934	Cataclysm	8	3	5
935	Lightning	4	2	2
936	Other and unspecified	24	13	11
946	Other complications due to nontherapeutic medical and surgical procedures	2	2	0
950	Therapeutic misadventure in surgical treatment	3	2	1
951	Therapeutic misadventure in infusion or transfusion	1	1	0
953	Therapeutic misadventure in administration of drugs or biologicals	1	0	1
954	Therapeutic misadventure in anesthesia	1	0	1
955	Other and unspecified therapeutic misadventure	1	0	1
956	Late complication of surgical operation	1	0	1
957	Late complication of amputation stump	2	0	2
960	Late effect of motor vehicle accident	1	0	1
962	Late effect of other accidental injury	6	3	3
<u>963, 970-979</u>	<u>Suicide</u>	<u>126</u>	<u>104</u>	<u>22</u>
970	By poisoning with analgesic and soporific substances	6	5	1
971	By poisoning with other solid and liquid substances	4	3	1
973	By poisoning with other gases	4	4	0
974	By hanging and strangulation	8	6	2
975	By submersion (drowning)	2	1	1
976	By firearms and explosives	97	81	16
977	By cutting and piercing instruments	4	4	0

Table 5 - Page 12
 Deaths - Mississippi - 1963

International List Number*	Cause of Death**	Total	White	Non-white
978	By jumping from high place	1	0	1
<u>964,980-985</u>	<u>Homicide</u>	<u>267</u>	<u>48</u>	<u>219</u>
980	Nonaccidental poisoning by another person	3	0	3
981	Assault by firearm and explosive	169	38	131
982	Assault by cutting and piercing instruments	67	5	62
983	Assault by other means	20	4	16
984	Injury by intervention of police	6	0	6
985	Execution	2	1	1

* International List Numbers are those of the International Statistical Classification of Diseases, Injuries, and Causes of Death, Seventh Revision, 1955.

** When cause of death and International List Number are omitted, no deaths were reported.

TABLE 6
DEATHS
MISSISSIPPI
1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

County	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
<u>STATE</u>	<u>23,043</u>	<u>11,960</u>	<u>11,083</u>	<u>23,339</u>	<u>12,118</u>	<u>11,221</u>	<u>10.2</u>	<u>9.0</u>	<u>11.9</u>
Adams	500	179	321	391	128	263	10.4	6.9	13.9
Alcorn	273	226	47	277	229	48	10.3	9.8	13.9
Amite	96	28	68	182	87	95	10.6	10.7	10.6
Attala	219	127	92	253	153	100	11.2	11.4	11.0
Benton	46	18	28	78	36	42	9.6	9.7	9.4
Bolivar	646	135	511	659	171	488	11.4	8.7	12.8
Calhoun	143	104	39	163	133	30	9.8	10.4	7.9
Carroll	64	24	40	123	58	65	10.2	10.4	10.0
Chickasaw	192	103	89	198	105	93	11.2	9.9	13.3
Choctaw	94	50	44	95	55	40	12.1	10.4	15.7
Claiborne	86	18	68	113	29	84	11.5	10.2	12.1
Clarke	130	95	35	208	147	61	12.2	12.7	11.1
Clay	177	86	91	193	97	96	9.8	9.7	9.8
Coahoma	594	152	442	596	129	467	12.3	8.5	14.0
Copiah	272	122	150	324	154	170	11.9	12.0	11.8
Covington	125	80	45	193	124	69	11.9	11.5	12.7
DeSoto	156	38	118	237	84	153	8.7	6.6	10.5
Forrest	563	387	176	568	366	202	10.0	8.8	13.6
Franklin	93	53	40	118	74	44	12.8	13.0	12.5
George	81	65	16	92	77	15	8.3	8.1	8.9
Greene	58	46	12	82	64	18	9.8	9.4	11.5
Grenada	221	113	108	205	93	112	10.7	9.0	12.7
Hancock	144	113	31	158	126	32	9.8	9.6	10.8
Harrison	1,129	892	237	948	716	232	7.6	6.8	11.5
Hinds	2,502	1,334	1,168	1,684	813	871	8.4	6.9	10.5
Holmes	307	81	226	335	101	234	12.4	12.8	12.2
Humphreys	188	47	141	219	63	156	11.3	10.3	11.7
Issaquena	25	2	23	39	7	32	14.5	8.2	17.4
Itawamba	111	101	10	152	142	10	8.8	8.6	10.9
Jackson	362	267	95	449	343	106	6.7	6.2	9.1
Jasper	124	66	58	165	84	81	9.6	9.3	10.0
Jefferson	83	19	64	136	44	92	13.4	13.5	13.3
Jeff Davis	107	51	56	136	69	67	10.1	10.1	10.1
Jones	743	448	295	587	412	175	9.9	9.4	11.5
Kemper	105	55	50	143	80	63	12.2	16.3	9.3
Lafayette	162	89	73	202	122	80	9.1	8.3	10.7

Table 6 - Page 2
Deaths - Mississippi - 1963

County	Place of Occurrence			Place of Residence					
	Number		Non-white	Number			Rate*		Non-white
	Total	White		Total	White	Non-white	Total	White	
Lamar	86	69	17	136	112	24	9.1	8.8	11.0
Lauderdale	996	617	379	804	454	350	10.9	9.0	15.0
Lawrence	89	53	36	124	80	44	13.2	13.1	13.6
Leake	151	99	52	181	116	65	10.4	11.8	8.6
Lee	448	314	134	419	287	132	9.4	8.8	11.3
Leflore	638	230	408	575	166	409	12.0	9.5	13.3
Lincoln	233	157	76	265	169	96	10.1	9.3	12.1
Lowndes	465	230	235	482	243	239	9.2	7.2	12.6
Madison	319	86	233	372	107	265	11.2	10.2	11.6
Marion	193	128	65	207	132	75	9.6	9.5	9.9
Marshall	238	76	162	254	68	186	10.1	9.0	10.6
Monroe	352	229	123	366	238	128	10.0	9.4	11.4
Montgomery	126	83	43	141	95	46	10.6	11.8	8.7
Neshoba	192	141	51	224	167	57	10.4	10.6	10.1
Newton	198	109	89	248	149	99	12.0	10.9	14.3
Noxubee	214	61	153	230	73	157	13.1	14.2	12.6
Oktibbeha	215	95	120	232	103	129	8.2	6.6	10.3
Panola	252	107	145	320	128	192	10.2	9.0	11.2
Pearl River	219	162	57	243	176	67	9.4	8.6	12.3
Perry	40	25	15	85	52	33	10.0	8.9	12.2
Pike	390	210	180	407	222	185	11.7	11.2	12.3
Pontotoc	193	156	37	209	173	36	11.1	11.3	10.4
Prentiss	195	167	28	195	165	30	10.1	9.8	12.2
Quitman	208	49	159	258	83	175	11.2	9.0	12.6
Rankin	395	196	199	275	158	117	7.2	6.7	8.1
Scott	245	141	104	256	151	105	11.1	10.3	12.5
Sharkey	112	22	90	123	28	95	11.6	9.4	12.5
Simpson	278	187	91	235	164	71	10.6	10.8	10.0
Smith	77	60	17	155	121	34	9.8	10.0	8.9
Stone	66	51	15	60	43	17	9.0	8.8	9.6
Sunflower	436	99	337	496	131	365	10.3	8.8	10.9
Tallahatchie	257	77	180	298	101	197	12.0	11.3	12.3
Tate	146	52	94	209	91	118	9.7	9.8	9.7
Tippah	145	110	35	163	131	32	9.7	9.7	9.9
Tishomingo	136	129	7	161	153	8	9.7	9.6	11.3
Tunica	212	33	179	223	33	190	13.1	9.4	14.1
Union	215	162	53	237	180	57	11.4	10.3	17.3
Walthall	138	75	63	135	73	62	10.0	9.8	10.3
Warren	707	286	421	548	249	299	12.4	10.2	15.2
Washington	838	257	581	893	249	644	11.6	7.4	14.8
Wayne	154	84	70	172	90	82	10.6	9.0	13.2
Webster	96	77	19	115	90	25	11.1	11.9	8.8
Wilkinson	161	81	80	160	54	106	12.4	13.5	11.9
Winston	171	97	74	191	111	80	9.8	10.4	9.0
Yalobusha	149	90	59	163	100	63	12.8	14.1	11.2
Yazoo	338	127	211	393	144	249	11.7	10.4	12.6
Out-of-State	1,001	699	302	705	541	164	-	-	-

Table 6 - Page 3
 Deaths - Mississippi - 1963

<u>City**</u>	Place of Occurrence			Place of Residence		
	Number		Non-white	Number		Non-white
	Total	White		Total	White	
Biloxi	290	239	51	270	214	56
Clarksdale	397	138	259	271	85	186
Cleveland	100	51	49	90	41	49
Columbus	330	183	147	280	139	141
Corinth	200	158	42	149	110	39
Greenville	528	204	324	466	156	310
Greenwood	424	157	267	299	108	191
Gulfport	482	359	123	330	228	102
Hattiesburg	473	330	143	360	206	154
Jackson	2,215	1,231	984	1,265	669	596
Laurel	608	339	269	291	165	126
McComb	238	145	93	175	105	70
Meridian	852	527	325	580	324	256
Natchez	437	164	273	281	92	189
Pascagoula	32	20	12	121	81	40
Tupelo	283	204	79	158	108	50
Vicksburg	606	244	362	382	181	201
Yazoo City	195	94	101	153	60	93
Total Urban#	12,458	6,939	5,519	8,692	4,584	4,108
Total Rural	10,585	5,021	5,564	14,647	7,534	7,113

* Deaths per 1,000 population.

** Cities having 10,000 population and over according to 1960 census.

Includes all incorporated towns of 2,500 population or more according to 1960 census.

TABLE 7
DEATHS
MISSISSIPPI
1963

By Cause Groups and Certain Selected Causes, by Race, Sex, and Age,
with Total Rates, Residence Data

International List Number#	Cause of Death	Total		Race Rate*	Total by Race	Sex	Total by Race and Sex	Age							Unknown Over 65	
		Total Number	Rate*					1-4	5-9	10-14	15-24	25-34	35-44	45-54		
001-999	<u>ALL CAUSES</u>	<u>23,339</u>	<u>10.2</u>	W <u>12,118</u>	M <u>7,271</u>	M	<u>3,711</u>	<u>53</u>	<u>39</u>	<u>37</u>	<u>157</u>	<u>153</u>	<u>356</u>	<u>761</u>	<u>1,280</u>	<u>4,063</u>
		C <u>11,221</u>		C <u>11,221</u>	F <u>4,847</u>	F	<u>239</u>	<u>49</u>	<u>24</u>	<u>12</u>	<u>69</u>	<u>59</u>	<u>138</u>	<u>299</u>	<u>553</u>	<u>3,405</u>
001-138	<u>Infective and parasitic diseases</u>	<u>268</u>	<u>11.7</u>	W <u>104</u>	M <u>61</u>	M	<u>2</u>	<u>1</u>	<u>4</u>	<u>2</u>	<u>0</u>	<u>2</u>	<u>6</u>	<u>1</u>	<u>15</u>	<u>24</u>
		C <u>164</u>		C <u>164</u>	F <u>89</u>	F	<u>2</u>	<u>1</u>	<u>4</u>	<u>1</u>	<u>0</u>	<u>1</u>	<u>3</u>	<u>8</u>	<u>6</u>	<u>0</u>
001-019	Tuberculosis (all forms)	113	4.9	W <u>40</u>	M <u>30</u>	M	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>4</u>	<u>3</u>	<u>14</u>
		C <u>73</u>		C <u>73</u>	F <u>45</u>	F	<u>1</u>	<u>1</u>	<u>1</u>	<u>1</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>0</u>	<u>1</u>	<u>8</u>
001-008	Tuberculosis (respiratory system)	106	4.6	W <u>39</u>	M <u>29</u>	M	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>4</u>	<u>2</u>	<u>8</u>
		C <u>67</u>		C <u>67</u>	F <u>43</u>	F	<u>1</u>	<u>1</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>0</u>	<u>1</u>	<u>8</u>
020-029	Syphilis & sequelae	34	1.5	W <u>9</u>	M <u>6</u>	M	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>0</u>	<u>3</u>
		C <u>25</u>		C <u>25</u>	F <u>17</u>	F	<u>2</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>10</u>
							<u>8</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>2</u>	<u>4</u>

Table 7 - Page 2
Deaths - Mississippi - 1963

International List Number#	Cause of Death	Total		Race*	Total by Race	Total by Sex	Age	Unknown Over 65 &
		Number	Rate*					
045-048	Dysentery (all forms)	14	0.6	W	3	M 2 F 1	0 0	0 0
050,051	Scarlet fever and streptococcal sore throat	4	0.2	W	1	M 0 F 1	0 1	0 0
055	Diphtheria	0	-	C	3	M 0 F 3	0 3	0 0
056	Whooping cough	5	0.2	W	1	M 0 F 1	0 1	0 0
057	Meningococcal infections	6	0.3	W	4	M 1 F 3	0 2	0 0
080	Acute poliomyelitis	4	0.2	W	4	M 2 F 2	1 1	0 0
085	Measles	7	0.3	W	3	M 0 F 3	0 2	0 0
				C	4	M 2 F 2	1 1	0 0

Table 7 - Page 3
Deaths - Mississippi - 1963

Inter-national List Number#	Cause of Death	Total		Race by Sex	Total by Race and Sex	Under 1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	Unknown Over 65	
		Number	Rate*												
140-205	Malignant neoplasms	2,945	128.6	W	1,859	M	1,080	F	779	M	559	F	527	M	410 328 234
				C	1,086	C	790	M	527	C	390	M	311	F	596 410 328 234
210-239	Benign and unspecified neoplasms	45	2.0	W	21	M	12	F	9	M	5	F	13	M	161 115 135
				C	24	M	5	F	5	C	13	M	13	F	161 115 135
240-289	Allergic, endocrine system, metabolic, & nutritional diseases	443	19.3	W	196	M	110	F	86	M	83	F	164	M	240 188 39
				C	247	M	31	F	83	C	31	M	31	F	240 188 39
241	Asthma	45	2.0	W	31	M	22	F	0	M	0	F	0	M	60 57 83
260	Diabetes mellitus	326	14.2	W	132	M	72	F	60	M	61	F	133	M	26 24 27
290-299	Diseases of the blood and blood- forming organs	83	3.6	W	32	M	21	F	11	M	11	F	21	M	12 7 9

Table 7 - Page 4
Deaths - Mississippi - 1963

Table 7 - Page 5
Deaths - Mississippi - 1963

International List Number#	Cause of Death	Total		Total by Race and Sex	Total by Race	Number	Rate*	Age						Unknown					
		Total	Rate*					10-14	15-19	20-24	25-34	35-44	45-54						
420-422	Arteriosclerotic and degenerative heart disease	5,359	234.0	W	3,691	M	2,348	1	0	0	0	96	245	484	1,517	0			
				F	1,343	M	1,343	0	0	1	5	10	35	143	1,148	0			
				C	1,668	M	927	0	0	2	3	24	82	155	659	1			
				F	741	F	741	0	0	0	2	13	20	65	138	502	1		
430-434	Other diseases of the heart	631	27.6	W	295	M	174	0	1	0	0	5	22	37	106	0			
				C	336	M	121	1	1	0	0	2	2	8	14	93	0		
				F	141	F	195	0	0	0	1	1	5	5	25	46	112	0	
							3	2	0	1	2	7	12	21	29	64	0		
440-443	Hypertensive heart disease	1,207	52.7	W	369	M	171	0	0	0	0	0	0	1	10	28	132	0	
				C	838	M	198	0	0	0	0	0	0	2	4	12	180	0	
				F	376	F	462	0	0	0	0	0	5	13	33	78	246	1	
							0	0	0	0	0	2	5	28	43	82	302	0	
444-447	Other hypertensive disease	203	8.9	W	77	M	37	0	0	0	0	0	0	0	4	6	27	0	
				C	126	M	40	0	0	0	0	0	0	1	2	3	34	0	
				F	71	F	55	0	0	0	0	0	3	8	6	11	27	0	
							0	0	0	0	0	0	0	0	1	3	132	0	
450	General arteriosclerosis	364	15.9	W	250	M	112	0	0	0	0	0	0	0	1	5	106	0	
				C	114	M	138	0	0	0	0	0	0	0	0	4	56	0	
				F	53	F	61	0	0	0	0	0	0	0	0	1	3	49	0
							0	0	0	0	0	0	0	0	2	26	303	0	
	Diseases of the respiratory system	1,477	64.5	W	681	M	430	23	6	4	2	2	2	7	15	192	0		
				C	796	M	251	15	7	1	5	3	6	7	19	168	1		
				F	344	F	452	152	47	3	3	0	0	0	0	8	139	3	
							0	0	0	0	0	0	0	0	0	0	0	0	

Table 7 - Page 6
Deaths - Mississippi - 1963

Table 7 - Page 7
Deaths - Mississippi - 1963

Inter-national List Number#	Cause of Death	Total Number	Rate*	Race	Total by Race	Total by Sex	Total by Race and Sex	Age	Unknown	
									65 & Over	55-64
581	Cirrhosis of liver	105	4.6	W	86	M	60	0	27	0
				C	19	M	26	5	5	0
				F	7	F	12	5	5	0
590-637	<u>Diseases of the genito-urinary system</u>	581	25.4	W	243	M	150	0	100	0
				C	338	M	93	61	61	0
				F	0	F	209	141	141	0
					0		129	58	58	0
590-594	Nephritis and nephrosis	261	11.4	W	103	M	57	1	11	25
				C	158	M	46	0	8	0
				F	0	F	100	1	14	28
					3		58	5	7	0
610	Hyperplasia of prostate	66	2.9	W	24	M	24	0	0	59
				C	42	M	0	0	8	0
				F	0	F	42	1	1	31
					0		0	5	7	0
640-689	<u>Complications of pregnancy, child- birth, and the puerperium</u>	44	7.7*	W	9	M	0	0	0	0
				C	35	M	0	0	0	0
				F	0	F	35	0	0	0
690-716	<u>Diseases of the skin and cellular tissue</u>	30	1.3	W	13	M	7	6	2	0
				C	17	M	8	9	1	0
				F	0	F	9	0	0	0

Table 7 - Page 8
Deaths - Mississippi - 1963

International List Number#	Cause of Death	Total Rate*	Race	Total by Race	Total by Sex	Race and Sex	Age												
							Number	Rate*	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65-74	Over 75	Unknown
<u>720-749</u>	<u>Diseases of the bones and organs of movement</u>	<u>44</u>	<u>1.9</u>	<u>W</u>	<u>31</u>	<u>M</u>	<u>16</u>	<u>15</u>	<u>F</u>	<u>5</u>	<u>M</u>	<u>13</u>	<u>F</u>	<u>8</u>	<u>1</u>	<u>1</u>	<u>1</u>	<u>1</u>	<u>1</u>
<u>750-759</u>	<u>Congenital malformations</u>	<u>256</u>	<u>11.2</u>	<u>W</u>	<u>147</u>	<u>M</u>	<u>73</u>	<u>51</u>	<u>F</u>	<u>74</u>	<u>M</u>	<u>109</u>	<u>F</u>	<u>61</u>	<u>50</u>	<u>39</u>	<u>10</u>	<u>8</u>	<u>1</u>
<u>760-776</u>	<u>Certain diseases of early infancy</u>	<u>1,262</u>	<u>22.0*</u>	<u>W</u>	<u>389</u>	<u>M</u>	<u>248</u>	<u>248</u>	<u>F</u>	<u>141</u>	<u>M</u>	<u>873</u>	<u>F</u>	<u>48</u>	<u>141</u>	<u>10</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>780-795</u>	<u>Symptoms, senility, and ill-defined conditions</u>	<u>1,770</u>	<u>77.3</u>	<u>W</u>	<u>435</u>	<u>M</u>	<u>272</u>	<u>13</u>	<u>F</u>	<u>163</u>	<u>M</u>	<u>1,335</u>	<u>F</u>	<u>737</u>	<u>12</u>	<u>21</u>	<u>0</u>	<u>1</u>	<u>1</u>
<u>800-999</u>	<u>Accidents, poisonings, and violence</u>	<u>1,964</u>	<u>85.8</u>	<u>W</u>	<u>1,003</u>	<u>M</u>	<u>744</u>	<u>12</u>	<u>F</u>	<u>259</u>	<u>M</u>	<u>961</u>	<u>F</u>	<u>598</u>	<u>115</u>	<u>19</u>	<u>4</u>	<u>6</u>	<u>1</u>
<u>810-835</u>	<u>Motor vehicle accidents</u>	<u>675</u>	<u>29.5</u>	<u>W</u>	<u>435</u>	<u>M</u>	<u>312</u>	<u>1</u>	<u>F</u>	<u>682</u>	<u>M</u>	<u>240</u>	<u>F</u>	<u>123</u>	<u>6</u>	<u>12</u>	<u>9</u>	<u>3</u>	<u>1</u>
<u>800-802, 840-962</u>	<u>All other accidents</u>	<u>896</u>	<u>39.1</u>	<u>W</u>	<u>416</u>	<u>M</u>	<u>304</u>	<u>11</u>	<u>F</u>	<u>112</u>	<u>F</u>	<u>480</u>	<u>M</u>	<u>188</u>	<u>0</u>	<u>18</u>	<u>5</u>	<u>10</u>	<u>25</u>
																	<u>5</u>	<u>6</u>	<u>6</u>
																	<u>14</u>	<u>37</u>	<u>41</u>
																	<u>4</u>	<u>44</u>	<u>33</u>
																	<u>8</u>	<u>4</u>	<u>13</u>
																	<u>23</u>	<u>38</u>	<u>53</u>
																	<u>6</u>	<u>6</u>	<u>63</u>
																	<u>2</u>	<u>6</u>	<u>46</u>
																	<u>8</u>	<u>8</u>	<u>47</u>

Table 7 - Page 9
Deaths - Mississippi - 1963

Inter-national List Number#	Cause of Death	Total		Race*	Total by Race	Sex	Total by Race and Sex	Age									
		Total Number	Rate*					10-14	15-24	25-34	35-44	45-54	55-64	65 & Over	Unknown		
963, 970-979	Suicide	126	5.5	W	104	M	86	0	0	0	8	17	26	15	12	0	
				C	22	F	18	0	0	0	3	2	3	2	1	0	
964, 980-985	Homicide	267	11.6	W	48	M	42	0	1	0	4	11	15	4	6	0	
				C	219	F	6	0	0	0	1	0	0	1	2	0	
					177	M	177	1	0	0	2	45	46	23	14	4	0
					42	F	42	1	1	3	1	6	14	10	3	2	1

International List Numbers are those of the International Statistical Classification of Diseases, Injuries, and Causes of Death, Seventh Revision, 1955.

* Total deaths per 1,000 population. Deaths from specific causes per 100,000 population. Deaths from maternal causes per 10,000 live births. Deaths from diseases of early infancy per 1,000 live births.

TABLE 8
DEATHS UNDER 1 YEAR
MISSISSIPPI
1963

By Selected Causes, by Race and Age, with Total Rates, Residence Data

Cause of Death and International List Number#	Total	Rate*	Race	DAYS OF AGE						MONTHS OF AGE										
				Under 1 day	1-	7-	14-	20	27	28 days	1	2	3	4	5	6	7	8		
<u>ALL CAUSES</u>	<u>2,372</u>	<u>41.4</u>	W C	<u>610</u> <u>1,762</u>	<u>238</u> <u>473</u>	<u>188</u> <u>317</u>	<u>23</u> <u>75</u>	<u>12</u> <u>68</u>	<u>9</u> <u>46</u>	<u>470</u> <u>979</u>	<u>29</u> <u>164</u>	<u>28</u> <u>131</u>	<u>13</u> <u>101</u>	<u>10</u> <u>90</u>	<u>14</u> <u>77</u>	<u>9</u> <u>69</u>	<u>16</u> <u>45</u>	<u>7</u> <u>30</u>	<u>3</u> <u>27</u>	<u>4</u> <u>25</u>
Dysentery, all forms (045-048)	5	0.1	W C	0 5	0 0	0 0	0 0	0 0	0 0	0 0	0 1	0 1	0 0	0 2	0 0	0 0	0 0	0 1	0 0	0 0
Septicemia and pyemia (053)	5	0.1	W C	1 4	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 1	0 2	0 0	0 0	0 0	0 1	0 0	0 0
Whooping cough (056)	2	0.03	W C	0 2	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 1	0 0	0 0	0 0	0 1	0 0	0 0	0 0
Meningococcal infections (057)	1	0.02	W C	0 1	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0
Tetanus (061)	3	0.05	W C	0 3	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0
Other infective and parasitic diseases	16	0.3	W C	3 13	0 0	0 1	0 0	2 0	1 0	0 0	3 0	0 0	0 0	0 1	0 0	0 0	0 0	0 0	0 0	0 0
Malignant neoplasms (140-205)	2	0.03	W C	2 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 1	0 2	0 0	2 1	0 0	1 1	0 0
Benign and unspecified neoplasms (210-239)	1	0.02	W C	1 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0
Diseases of thymus gland (273)	0	-	W C	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0
Meningitis, except meningococcal and tuberculous (340)	24	0.4	W C	6 18	0 1	2 1	1 0	1 2	1 0	1 2	4 1	0 2	1 1	0 0	0 0	0 0	0 0	0 3	0 0	0 1

Table 8 - Page 2
Deaths Under 1 Year - Mississippi - 1963

Cause of Death and International List Number#	Total	Rate*	Case Race	Total by Race	MONTHS OF AGE															
					Under 1 day	1-	6	7-	13	20	27	28 days	1	2	3	4	5	6	7	
Other diseases of nervous system and sense organs (330-334, 341-398)	11	0.2	W C	3 8	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	
Acute upper respiratory infections (470-475)	17	0.3	W C	1 16	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0
Influenza & pneumonia, except pneumonia of newborn (480-493)	260	4.5	W C	33 227	0	0	0	0	0	0	8	5	5	2	4	5	2	0	1	0
Bronchitis (500-502)	5	0.1	W C	2 3	0	0	0	0	1	1	0	0	0	0	1	0	0	0	0	0
Other diseases of respiratory system (510-522, 525-527)	13	0.2	W C	2 11	0	0	0	0	0	1	1	0	0	1	0	1	0	0	0	0
Hernia & intestinal obstruction (560, 561, 570)	14	0.2	W C	1 13	0	0	0	0	1	0	4	2	1	0	1	0	0	0	0	0
Gastritis, duodenitis, enteritis, & colitis, except diarrhea of newborn (543, 571, 572)	136	2.4	W C	6 130	0	0	0	0	0	0	0	19	19	0	0	0	2	0	1	0
Other diseases of digestive system (530-542, 544-553, 573-587)	2	0.03	W C	1 1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0
Congenital malformations (750-759)	170	3.0	W C	101 69	33 15	15 13	9 5	3 5	63 40	10 7	5 6	4 2	2 1	4 3	1 2	4 2	2 0	2 0	0 0	0 0

Table 8 - Page 3
Deaths Under 1 Year - Mississippi - 1963

Table 8 - Page 4
Deaths Under 1 Year - Mississippi - 1963

Cause of Death and International List Number#	Total	Rate*	Total by Race	Days Under 1 day	Days of Age						Months of Age							
					Under 1 6	7- 13	14- 20	21- 27	Under 28 days	1	2	3	4	5	6	7		
Immaturity, unqualified (776)	491	8.6	W C	102 389	72 220	27 87	2 15	0 4	101 352	1 14	0 6	0 4	0 6	0 2	0 1	0 1	0 1	
Symptoms & ill-defined conditions (780-793, 795)	294	5.1	W C	25 269	6 42	1 37	1 6	0 8	1 9	9 99	2 45	1 33	3 20	1 17	1 17	1 13	1 6	0 7
All other diseases	25	0.4	W C	10 15	1 1	2 2	0 0	3 0	6 0	0 3	3 0	1 2	0 2	0 2	0 2	0 3	0 3	0 0
Accidents (800-962)	102	1.8	W C	23 79	0 4	0 6	0 3	1 2	0 6	1 21	3 9	7 12	3 4	0 7	2 10	2 3	1 4	1 2
Homicide (964, 980-984)	2	0.03	W C	0 2	0 0	0 0	0 1	0 0	0 1	0 1	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0

International List Numbers are those of the International Statistical Classification of Diseases, Injuries, and Causes of Death, Seventh Revision, 1955.

* Deaths per 1,000 live births.

TABLE 9
MARRIAGES AND DIVORCES
MISSISSIPPI
1963

Marriages by Place of Licensure, and by Place of Residence of Bride, by Race of Bride
Divorces by Place of Occurrence, by Race

County	Marriages						Divorces		
	Place of Licensure			Residence of Bride					
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
<u>STATE</u>	<u>22,140</u>	<u>14,508</u>	<u>7,632</u>	<u>18,515</u>	<u>11,548</u>	<u>6,967</u>	<u>5,452</u>	<u>3,930</u>	<u>1,522</u>
Adams	365	236	129	292	170	122	94	59	35
Alcorn	508	466	42	260	233	27	75	68	7
Amite	124	70	54	100	56	44	26	21	5
Attala	194	127	67	194	124	70	50	22	28
Benton	57	22	35	60	34	26	18	11	7
Bolivar	483	196	287	424	165	259	89	53	36
Calhoun	145	118	27	131	104	27	43	34	9
Carroll	70	36	34	62	27	35	13	10	3
Chickasaw	195	113	82	181	109	72	31	22	9
Choctaw	78	50	28	64	48	16	6	5	1
Claiborne	71	28	43	65	25	40	19	1	18
Clarke	125	89	36	110	76	34	40	22	18
Clay	160	89	71	138	78	60	43	24	19
Coahoma	373	138	235	351	120	231	93	40	53
Copiah	216	125	91	230	133	97	70	29	41
Covington	202	137	65	140	93	47	44	37	7
DeSoto	963	529	434	149	60	89	48	33	15
Forrest	558	424	134	499	360	139	181	148	33
Franklin	85	64	21	85	65	20	19	9	10
George	165	153	12	95	88	7	23	22	1
Greene	92	80	12	47	36	11	12	12	0
Grenada	201	121	80	188	109	79	44	24	20
Hancock	445	406	39	107	82	25	39	33	6
Harrison	1,531	1,311	220	1,159	957	202	643	559	84
Hinds	1,779	1,057	722	1,797	1,098	699	612	470	142
Holmes	188	69	119	218	71	147	40	13	27
Humphreys	158	66	92	154	59	95	27	19	8
Issaquena	12	5	7	21	7	14	6	4	2
Itawamba	116	108	8	72	66	6	18	18	0
Jackson	774	623	151	562	451	111	233	194	39
Jasper	103	64	39	119	69	50	21	14	7
Jefferson	75	35	40	69	23	46	9	4	5
Jeff Davis	124	59	65	102	44	58	23	16	7
Jones	583	473	110	554	458	96	133	108	25
Kemper	84	37	47	66	29	37	18	7	11
Lafayette	134	86	48	124	72	52	35	21	14

Table 9 - Page 2
Marriages and Divorces - Mississippi - 1963

County	Marriages						Divorces		
	Place of Licensure			Residence of Bride			Total	White	Non-white
	Total	White	Non-white	Total	White	Non-white			
Lamar	117	104	13	142	126	16	32	30	2
Lauderdale	558	382	176	565	380	185	203	153	50
Lawrence	73	54	19	138	88	50	26	22	4
Leake	150	107	43	152	104	48	31	18	13
Lee	351	268	83	374	295	79	122	110	12
Leflore	464	187	277	454	189	265	97	60	37
Lincoln	252	190	62	221	168	53	49	36	13
Lowndes	385	234	151	355	195	160	118	85	33
Madison	306	112	194	255	87	168	67	14	53
Marion	249	185	64	228	171	57	49	41	8
Marshall	225	110	115	130	57	73	36	19	17
Monroe	273	210	63	271	195	76	88	72	16
Montgomery	122	73	49	113	63	50	32	20	12
Neshoba	175	126	49	193	142	51	55	46	9
Newton	160	118	42	141	101	40	29	21	8
Noxubee	112	23	89	113	26	87	25	13	12
Oktibbeha	192	104	88	173	90	83	36	19	17
Panola	241	122	119	231	118	113	46	31	15
Pearl River	362	316	46	257	220	37	93	83	10
Perry	79	61	18	90	73	17	21	16	5
Pike	290	175	115	299	175	124	105	69	36
Pontotoc	179	152	27	157	132	25	37	30	7
Prentiss	214	200	14	190	176	14	50	45	5
Quitman	209	95	114	208	89	119	40	20	20
Rankin	292	212	80	259	172	87	58	42	16
Scott	199	144	55	204	149	55	35	29	6
Sharkey	101	42	59	81	31	50	21	9	12
Simpson	214	151	63	167	122	45	46	33	13
Smith	93	77	16	112	83	29	27	22	5
Stone	92	73	19	77	63	14	23	18	5
Sunflower	317	109	208	362	111	251	68	43	25
Tallahatchie	217	85	132	198	81	117	50	30	20
Tate	162	89	73	148	68	80	22	16	6
Tippah	154	127	27	143	119	24	28	25	3
Tishomingo	136	130	6	109	105	4	20	20	0
Tunica	128	40	88	109	24	85	19	7	12
Union	182	141	41	161	122	39	38	29	9
Walthall	160	108	52	83	47	36	24	18	6
Warren	397	229	168	357	203	154	127	78	49
Washington	851	471	380	690	348	342	212	129	83
Wayne	156	103	53	116	78	38	43	33	10
Webster	73	57	16	83	64	19	17	16	1
Wilkinson	407	331	76	90	28	62	32	8	24
Winston	148	85	63	131	74	57	28	18	10
Yalobusha	138	82	56	129	82	47	28	19	9
Yazoo	249	104	145	267	115	152	61	29	32
Out-of-State				3,625	2,960	665			

TABLE 10
DIVORCES
MISSISSIPPI
1963

By Cause of Divorce, by Race

Cause	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>ALL CAUSES</u>	<u>5,452</u>	<u>3,930</u>	<u>1,522</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
Cruel and inhuman treatment	3,035	2,617	418	55.7	66.6	27.5
Desertion	1,867	892	975	34.2	22.7	64.1
Adultery	219	134	85	4.0	3.4	5.6
Drunkenness	230	210	20	4.2	5.3	1.3
Penitentiary	51	42	9	0.9	1.1	0.6
Insanity	5	5	0	0.1	0.1	-
Annulment	12	11	1	0.2	0.3	0.1
Bigamy	16	7	9	0.3	0.2	0.6
Miscellaneous and other	5	5	0	0.1	0.1	-
Not stated	12	7	5	0.2	0.2	0.3

By Number of Minor Children, by Race

Number of Minor Children	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>TOTAL DIVORCES</u>	<u>5,452</u>	<u>3,930</u>	<u>1,522</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
None	2,505	1,645	860	45.9	41.8	56.5
One	1,138	889	249	20.9	22.6	16.4
Two	806	666	140	14.8	16.9	9.2
Three or more	941	690	251	17.2	17.6	16.5
Not stated	62	40	22	1.1	1.0	1.4

By Number of Years Married, by Race

Number of Years Married	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>TOTAL DIVORCES</u>	<u>5,452</u>	<u>3,930</u>	<u>1,522</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
<u>Under five</u>	<u>2,005</u>	<u>1,713</u>	<u>292</u>	<u>36.8</u>	<u>43.6</u>	<u>19.2</u>
Under one	374	347	27	6.8	8.8	1.8
One	507	445	62	9.3	11.3	4.1
Two	421	369	52	7.7	9.4	3.4
Three	385	315	70	7.1	8.0	4.6
Four	318	237	81	5.8	6.0	5.3
Five - nine	1,238	922	316	22.7	23.5	20.8
Ten - nineteen	1,197	810	387	22.0	20.6	25.4
Twenty and over	912	408	504	16.7	10.4	33.1
Not stated	100	77	23	1.8	2.0	1.5

TABLE 11
LIVE BIRTHS
MISSISSIPPI
1913-1963

By Year, by Race, with Rates, Occurrence Data 1913-1943,
Residence Data 1944-1963

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1913	41,103	21,874	19,229	21.9	26.6	18.2
1914	40,623	21,360	19,263	21.4	25.7	18.0
1915	39,484	20,999	18,485	20.5	23.8	17.1
1916	39,033	20,130	18,903	20.0	23.6	17.2
1917	37,141	19,056	18,085	18.8	22.0	16.3
1918	39,391	20,005	19,386	19.7	22.8	17.2
1919	42,491	22,028	20,463	21.0	24.9	18.0
1920	43,743	22,438	21,305	24.4	26.3	22.7
1921	46,482	23,703	22,779	25.6	27.3	24.1
1922	43,723	21,286	22,437	23.4	24.1	23.6
1923	43,266	21,140	22,126	23.3	23.6	23.1
1924	43,960	22,180	21,780	23.4	24.3	22.5
1925	45,563	22,265	23,298	24.0	24.1	23.9
1926	50,449	24,600	25,849	26.2	26.2	26.3
1927	49,204	24,163	25,041	25.3	25.3	25.3
1928	48,126	23,642	24,484	24.5	24.4	24.5
1929	45,570	22,105	23,465	22.9	22.5	23.3
1930	47,982	23,192	24,790	23.8	23.2	24.0
1931	45,123	21,637	23,486	22.2	21.4	22.9
1932	46,481	21,883	24,598	22.6	21.4	23.9
1933	44,184	20,407	23,777	21.3	19.7	22.9
1934	47,905	22,342	25,563	23.0	21.4	24.6
1935	48,291	21,972	26,319	23.0	20.8	25.1
1936	49,261	22,194	27,067	23.3	20.8	25.7
1937	52,170	23,285	28,885	24.4	21.6	27.3
1938	53,729	24,046	29,683	25.0	22.1	27.8
1939	52,288	23,505	28,783	24.1	21.4	26.8
1940	52,526	23,084	29,442	24.0	20.9	27.3
1941	54,561	24,167	30,394	24.7	21.6	28.0
1942	56,521	25,306	31,215	25.4	22.4	28.6
1943	59,408	27,611	31,797	28.9	26.4	31.6
1944#	56,586	25,647	30,939	28.3	25.2	31.6
1945#	53,540	23,426	30,114	26.9	23.1	30.9
1946#	61,430	29,715	31,715	26.8	25.4	28.2
1947#	66,291	32,423	33,868	28.6	27.5	29.9
1948#	65,291	29,666	35,625	28.0	24.9	31.2
1949#	66,128	28,786	37,342	28.1	24.0	32.5

Table 11 - Page 2
Live Births - Mississippi - 1913-1963

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1950#	64,427	27,188	37,239	29.6	22.9	37.6
1951#	65,512	27,798	37,714	30.1	23.4	38.1
1952#	64,509	28,073	36,436	29.6	23.6	36.8
1953#	64,061	28,312	35,749	29.4	23.8	36.1
1954#	64,748	28,781	35,967	29.7	24.2	36.3
1955#	63,244	27,546	35,698	29.0	23.2	36.0
1956#	62,178	27,278	34,900	29.5	23.8	36.2
1957#	60,712	26,804	33,908	28.8	23.4	35.2
1958#	59,189	26,812	32,377	27.1	20.9	35.9
1959#	60,562	27,979	32,583	27.2	20.9	36.6
1960#	59,210	27,198	32,012	27.2	21.6	34.8
1961#	59,523	27,435	32,088	27.3	21.8	34.8
1962#	58,921	27,512	31,409	26.1	20.8	33.5
1963#	57,316	26,823	30,493	25.0	19.9	32.4

* Births per 1,000 population.

1913-1920, population estimates based on U.S. Censuses of 1910 and 1920.

1921-1930, estimates based on U.S. Censuses of 1920 and 1930.

1931-1940, estimates based on U.S. Censuses of 1930 and 1940.

1941 and 1942, projected estimates based on U.S. Censuses of 1930 and 1940.

1943 and 1944, estimates of civilian population obtained from Ration Book Registration.

1945, estimate of civilian population, July 1, 1945, obtained from Bureau of Census Bulletin, Series P-46, No. 3, dated February 12, 1946.

1946-1949, projected estimates as of July 1, each year based on U.S. Censuses of 1930 and 1940.

1950-1955, U.S. Census population of April 1, 1950.

1956 and 1957, estimate as of July 1, 1956, prepared by Dr. Harald A. Pedersen, Sociology Department, Mississippi State University, with adjustments by Statistical Service, Mississippi State Board of Health.

1958, estimate as of January 1, prepared by Dr. George L. Wilber and Mr. Carl T. Robbins, Sociology Department, Mississippi State University.

1959, estimate as of January 1, prepared by Dr. George L. Wilber and Ellen Bryant, Sociology Department, Mississippi State University.

1960 and 1961, U.S. Census population of April 1, 1960.

1962 and 1963, estimates as of July 1, prepared by the Division of Sociology and Rural Life, Mississippi State University.

On basis of place of residence.

TABLE 12
DEATHS
MISSISSIPPI
1913-1963

By Year, by Race, with Rates, Occurrence Data 1913-1943,
Residence Data 1944-1963

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1913	21,305	7,643	13,662	11.4	9.3	13.0
1914	21,485	7,442	14,043	11.3	8.9	13.1
1915	24,001	8,088	15,913	12.5	9.6	14.7
1916	21,794	7,588	14,206	11.2	8.9	13.0
1917	23,579	8,048	15,531	11.9	9.3	14.0
1918	30,437	10,177	20,260	15.2	11.6	18.0
1919	22,667	8,142	14,525	12.7	9.5	15.5
1920	22,072	7,916	14,156	12.3	9.3	15.1
1921	19,991	7,308	12,683	11.0	8.4	13.4
1922	19,278	7,312	11,966	10.5	8.3	12.6
1923	20,363	7,688	12,675	11.0	8.6	13.2
1924	21,155	7,809	13,346	11.3	8.6	13.8
1925	22,229	8,143	14,086	11.7	8.8	14.4
1926	23,929	9,133	14,796	12.5	9.7	15.1
1927	23,257	8,765	14,492	12.0	9.2	14.6
1928	25,965	10,111	15,854	13.2	10.4	15.9
1929	25,868	10,048	15,820	13.0	10.2	15.7
1930	24,098	9,072	15,026	11.9	9.0	14.8
1931	22,007	8,641	13,366	10.8	8.5	13.0
1932	20,322	8,295	12,027	9.9	8.1	11.6
1933	21,617	8,898	12,719	10.4	8.6	12.2
1934	21,786	9,350	12,436	10.5	9.0	12.0
1935	21,285	9,186	12,099	10.1	8.7	11.6
1936	24,127	10,367	13,760	11.4	9.7	13.1
1937	24,005	10,055	13,950	11.2	9.3	13.2
1938	22,814	9,471	13,343	10.6	8.7	12.5
1939	22,701	9,464	13,237	10.5	8.6	12.3
1940	23,185	9,565	13,620	10.6	8.6	12.6
1941	22,573	9,442	13,131	10.2	8.4	12.1
1942	20,693	8,750	11,943	9.3	7.7	10.9
1943	20,850	9,139	11,711	10.2	8.7	11.6
1944#	20,199	8,830	11,369	10.1	8.7	11.6
1945#	19,024	8,452	10,572	9.6	8.3	10.8
1946#	19,321	8,995	10,326	8.4	7.7	9.2
1947#	20,007	9,238	10,769	8.6	7.8	9.5
1948#	20,183	9,216	10,967	8.6	7.7	9.6
1949#	20,489	9,233	11,256	8.7	7.7	9.8

Table 12 - Page 2
 Deaths - Mississippi - 1913-1963

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1950#	20,567	9,466	11,101	9.4	8.0	11.2
1951#	21,007	9,593	11,414	9.6	8.1	11.5
1952#	20,657	9,387	11,270	9.5	7.9	11.4
1953#	20,492	9,495	10,997	9.4	8.0	11.1
1954#	19,353	9,453	9,900	8.9	8.0	10.0
1955#	19,403	9,501	9,902	8.9	8.0	10.0
1956#	19,780	9,969	9,811	9.4	8.7	10.2
1957#	20,476	10,339	10,137	9.7	9.0	10.5
1958#	21,541	10,811	10,730	9.9	8.4	11.9
1959#	20,890	10,742	10,148	9.4	8.0	11.4
1960#	21,713	11,204	10,509	10.0	8.9	11.4
1961#	21,232	11,063	10,169	9.7	8.8	11.0
1962#	22,713	11,868	10,845	10.0	9.0	11.6
1963#	23,339	12,118	11,221	10.2	9.0	11.9

* Deaths per 1,000 population. See footnote on Table 11 for source of population estimates.

On basis of place of residence.

TABLE 13
MATERNAL DEATHS
MISSISSIPPI
1913-1963

By Year, by Race, with Rates, Occurrence Data 1913-1943,
Residence Data 1944-1963

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1913	447	162	285	108.8	74.1	148.2
1914	413	124	289	101.7	58.0	150.0
1915	289	88	201	73.2	41.9	108.7
1916	326	102	224	83.5	50.7	118.5
1917	297	105	192	80.0	55.1	106.2
1918	302	82	220	76.7	41.0	113.5
1919	407	173	234	95.8	78.5	114.4
1920	393	127	266	89.8	56.6	124.8
1921	391	145	246	84.1	61.2	108.0
1922	313	110	203	71.6	51.7	90.5
1923	321	108	213	74.2	51.1	96.3
1924	413	137	276	93.9	61.8	126.7
1925	435	143	292	95.5	64.2	125.3
1926	388	153	235	76.9	62.2	90.9
1927	409	134	275	83.1	55.4	109.8
1928	421	161	260	87.5	68.1	106.2
1929	410	152	258	90.0	68.8	110.0
1930	489	171	318	101.9	73.7	128.3
1931	352	131	221	78.0	60.5	94.1
1932	279	103	176	60.0	47.1	71.6
1933	315	116	199	71.3	56.8	83.7
1934	306	116	190	63.9	51.9	74.3
1935	334	137	197	69.2	62.4	74.8
1936	358	134	224	72.7	60.4	82.8
1937	370	150	220	70.9	64.4	76.2
1938	333	116	217	62.0	48.2	73.1
1939	311	97	214	59.5	41.3	74.3
1940	327	106	221	62.2	45.9	75.1
1941	293	93	200	53.7	38.5	65.8
1942	248	77	171	43.9	30.4	54.8
1943	233	66	167	39.2	23.9	52.5
1944#	225	72	153	39.8	28.1	49.4
1945#	216	64	152	40.3	27.3	50.5
1946#	199	67	132	32.4	22.5	41.6
1947#	175	50	125	26.4	15.4	36.9
1948#	177	38	139	27.1	12.8	39.0
1949#	147	32	115	22.2	11.1	30.8

Table 13 - Page 2
 Maternal Deaths - Mississippi - 1913-1963

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1950#	165	30	135	25.6	11.0	36.2
1951#	143	28	115	21.8	10.1	30.5
1952#	119	15	104	18.4	5.3	28.5
1953#	119	24	95	18.6	8.5	26.6
1954#	104	11	93	16.1	3.8	25.8
1955#	76	12	64	12.0	4.4	17.9
1956#	74	12	62	11.9	4.4	17.8
1957#	75	14	61	12.4	5.2	18.0
1958#	53	10	43	9.0	3.7	13.3
1959#	55	9	46	9.1	3.2	14.1
1960#	59	8	51	10.0	2.9	15.9
1961#	49	9	40	8.2	3.3	12.5
1962#	55	7	48	9.3	2.5	15.3
1963#	44	9	35	7.7	3.4	11.5

* Deaths per 10,000 live births.

On basis of place of residence.

TABLE 14
DEATHS UNDER 1 YEAR
MISSISSIPPI
1917-1963

By Year, by Race, with Rates, Occurrence Data 1917-1943,
Residence Data 1944-1963

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1917	3,689	1,296	2,393	99.3	68.0	132.3
1918	**	**	**	**	**	**
1919	3,554	**	**	83.6	**	**
1920	3,499	1,344	2,155	80.0	59.9	101.1
1921	3,157	1,243	1,914	67.9	52.4	84.0
1922	2,826	1,067	1,759	64.6	50.1	78.4
1923	2,908	1,104	1,804	67.2	52.2	81.5
1924	3,137	1,188	1,949	71.4	53.6	89.5
1925	3,176	1,179	1,997	69.7	53.0	85.7
1926	3,578	1,447	2,131	70.9	58.8	82.4
1927	3,291	1,303	1,988	66.9	54.0	79.4
1928	3,567	1,416	2,151	74.1	59.9	87.9
1929	3,275	1,268	2,007	71.9	57.4	85.5
1930	3,276	1,182	2,094	68.3	51.0	84.5
1931	2,547	961	1,586	56.4	44.4	67.5
1932	2,482	965	1,517	53.4	44.1	61.7
1933	2,800	1,035	1,765	63.4	50.7	74.2
1934	3,070	1,199	1,871	64.1	53.7	73.2
1935	2,587	1,044	1,543	53.6	47.5	58.6
1936	2,883	1,105	1,778	58.5	49.8	65.7
1937	3,098	1,174	1,924	59.4	50.4	66.6
1938	3,051	1,205	1,846	56.8	50.1	62.2
1939	2,908	1,135	1,773	55.6	48.3	61.6
1940	2,864	1,070	1,794	54.5	46.4	60.9
1941	2,969	1,113	1,856	54.4	46.1	61.1
1942	2,654	957	1,697	47.0	37.8	54.4
1943	2,774	1,137	1,637	46.7	41.2	51.5
1944#	2,489	974	1,515	44.0	38.0	49.0
1945#	2,192	846	1,346	40.9	36.1	44.7
1946#	2,298	1,005	1,293	37.4	33.8	40.8
1947#	2,428	1,009	1,419	36.6	31.1	41.9
1948#	2,470	988	1,482	37.8	33.3	41.6
1949#	2,612	900	1,712	39.5	31.3	45.8

Table 14 - Page 2
 Deaths Under 1 Year - Mississippi - 1917-1963

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1950#	2,348	763	1,585	36.4	28.1	42.6
1951#	2,589	814	1,775	39.5	29.3	47.1
1952#	2,592	760	1,832	40.2	27.1	50.3
1953#	2,534	807	1,727	39.6	28.5	48.3
1954#	2,360	744	1,616	36.4	25.8	44.9
1955#	2,338	680	1,658	37.0	24.7	46.4
1956#	2,353	734	1,619	37.8	26.9	46.4
1957#	2,257	694	1,563	37.2	25.9	46.1
1958#	2,424	706	1,718	41.0	26.3	53.1
1959#	2,355	677	1,678	38.9	24.2	51.5
1960#	2,457	714	1,743	41.5	26.2	54.4
1961#	2,250	645	1,605	37.8	23.5	50.0
1962#	2,353	705	1,648	39.9	25.6	52.5
1963#	2,372	610	1,762	41.4	22.7	57.8

* Deaths per 1,000 live births.

** Not available.

On basis of place of residence.

TABLE 15
DEATHS UNDER 1 MONTH*
MISSISSIPPI
1924-1963

By Year, by Race, with Rates, Occurrence Data 1924-1944,
Residence Data 1945-1963

Year	Number			Rate**		
	Total	White	Non-white	Total	White	Non-white
1924	1,598	730	868	36.4	32.9	39.8
1925	1,639	701	938	36.0	31.5	40.3
1926	1,731	823	908	34.3	33.4	35.1
1927	1,644	786	858	33.4	32.5	34.3
1928	1,788	804	984	37.2	34.0	40.2
1929	1,662	773	889	36.5	35.0	37.9
1930	1,692	715	977	35.3	30.8	39.4
1931	1,339	590	749	29.7	27.3	31.9
1932	1,361	572	789	29.3	26.1	32.1
1933	1,540	636	904	34.8	31.2	38.0
1934	1,505	644	861	31.4	28.8	33.7
1935	1,446	642	804	29.9	29.2	30.5
1936	1,494	676	818	30.3	30.4	30.2
1937	1,678	755	923	32.2	32.4	32.0
1938	1,688	747	941	31.4	31.1	31.7
1939	1,637	735	902	31.3	31.3	31.3
1940	1,613	680	933	30.7	29.4	31.7
1941	1,663	734	929	30.5	30.4	30.6
1942	1,544	638	906	27.3	25.2	29.0
1943	1,652	791	861	27.8	28.6	27.1
1944	1,436	661	775	25.2	25.4	25.1
1945#	1,351	624	727	25.2	26.6	24.1
1946#	1,561	794	767	25.4	26.7	24.2
1947#	1,595	750	845	24.1	23.1	24.9
1948#	1,674	773	901	25.6	26.0	25.3
1949#	1,594	661	933	24.1	23.0	25.0
1950#	1,446	560	886	22.4	20.6	23.8
1951#	1,679	626	1,053	25.6	22.5	27.9
1952#	1,658	581	1,077	25.7	20.7	29.6
1953#	1,630	635	995	25.4	22.4	27.8
1954#	1,521	601	920	23.5	20.9	25.6
1955#	1,440	527	913	22.8	19.1	25.6
1956#	1,451	581	870	23.3	21.3	24.9
1957#	1,425	540	885	23.5	20.1	26.1
1958#	1,475	540	935	24.9	20.1	28.9
1959#	1,450	515	935	23.9	18.4	28.7
1960#	1,497	550	947	25.3	20.2	29.6
1961#	1,401	509	892	23.5	18.6	27.8
1962#	1,456	550	906	24.7	20.0	28.8
1963#	1,449	470	979	25.3	17.5	32.1

* Figures for 1949-1963 include only the deaths under 28 days.

** Deaths per 1,000 live births.

#On basis of place of residence.

TABLE 16
DEATHS FROM HEART DISEASES, VASCULAR LESIONS,
MALIGNANT NEOPLASMS, ACCIDENTS, AND INFLUENZA AND PNEUMONIA
MISSISSIPPI
1954-1963

By Year, by Race, with Rates, Residence Data

Year	HEART DISEASES (400-443)			VASCULAR LESIONS (330-334)			MALIGNANT NEOPLASMS (140-205)			ACCIDENTS (800-962)			INFLUENZA & PNEUMONIA (480-493)			
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	
1954 No. Rate*	5,738 263.3	3,333 280.4	2,405 242.9	2,422 111.2	1,176 98.9	1,246 125.8	2,332 107.0	1,420 119.5	912 92.1	1,292 59.3	684 57.5	608 61.4	585 26.8	209 17.6	376 38.0	
1955 No. Rate*	5,994 275.1	3,525 296.6	2,469 249.3	2,447 112.3	1,169 98.3	1,278 129.0	2,317 106.3	1,402 118.0	915 92.4	1,349 61.9	696 58.6	653 65.9	634 29.1	220 18.5	414 41.8	
1956 No. Rate*	6,117 289.9	3,709 323.3	2,408 250.2	2,550 120.9	1,229 107.1	1,321 137.2	2,432 115.3	1,439 125.4	993 103.2	1,302 61.7	700 61.0	602 62.5	678 32.1	250 21.8	428 44.5	
1957 No. Rate*	6,378 302.3	3,711 323.5	2,667 277.1	2,800 132.7	1,420 123.8	1,380 143.9	2,496 118.3	1,482 129.2	1,014 105.3	1,302 61.7	707 61.6	595 61.8	794 37.6	301 37.6	493 51.2	
1958 No. Rate*	6,528 298.9	3,929 306.3	2,599 288.4	2,867 131.3	1,426 111.2	1,441 159.9	2,525 115.6	1,524 118.8	1,001 111.1	1,379 63.1	729 56.8	650 72.1	909 41.6	344 26.8	565 62.7	
1959 No. Rate*	6,233 279.9	3,774 282.5	2,459 276.0	2,683 120.5	1,386 103.7	1,297 145.6	2,600 116.7	1,580 118.3	1,020 114.5	1,501 67.4	837 62.6	664 74.5	768 34.5	328 24.6	440 49.4	
1960 No. Rate*	6,321 290.2	3,882 308.7	2,439 264.9	2,960 135.9	1,513 120.3	1,447 157.2	2,635 121.0	1,644 130.7	991 107.6	1,504 69.0	827 65.8	677 73.5	958 44.0	389 30.9	569 61.8	
1961 No. Rate*	6,561 301.2	4,022 319.8	2,539 275.8	2,891 132.7	1,464 116.4	1,427 155.0	2,748 126.2	1,732 137.7	1,016 110.4	1,427 65.5	760 60.4	667 72.4	721 33.1	307 24.4	414 45.0	
1962 No. Rate*	7,077 313.4	4,206 318.4	2,871 306.4	3,112 137.8	1,552 117.5	1,560 166.5	2,827 125.2	1,793 135.7	1,034 110.4	1,557 69.0	846 64.0	711 75.9	879 38.9	362 27.4	517 55.2	
1963 No. Rate*	7,300 318.8	4,396 325.9	2,904 308.5	3,154 137.7	1,650 122.3	1,504 159.8	2,945 128.6	1,859 137.8	1,086 115.4	1,571 68.6	851 63.1	720 76.5	1,113 76.1	420 48.6	693 31.1	420 73.6

* Deaths per 100,000 population.

TABLE 17
MARRIAGES AND DIVORCES
MISSISSIPPI
1926-1963

By Year, by Race, Occurrence Data

Year	MARRIAGES			DIVORCES		
	Total	White	Non-white	Total	White	Non-white
1926*	28,712			2,766		
1927*	29,048#	11,402	17,383	2,895		
1928	30,231#	11,662	18,257	3,009	1,435	1,574
1929	31,495#	12,864	18,388	3,172	1,598	1,574
1930	25,728#	10,815	14,751	2,743	1,487	1,256
1931	21,589#	9,952	11,629	2,015	1,308	707
1932	22,688#	10,147	12,453	1,918	1,308	610
1933	27,567#	12,201	15,300	1,955	1,396	559
1934	30,149#	13,497	16,541	2,446	1,704	742
1935	28,722#	12,955	15,635	2,425	1,681	744
1936	32,480#	14,120	18,196	2,520	1,690	830
1937	32,055#	14,802	17,246	2,977	1,984	993
1938	31,495	14,224	17,271	2,916	2,069	847
1939	31,956	15,370	16,586	3,034	2,166	868
1940	34,088	16,834	17,254	3,251	2,295	956
1941	42,142	21,611	20,531	3,831	2,444	1,387
1942	49,565	27,227	22,338	4,326	2,606	1,720
1943	46,030	23,445	22,585	5,851	3,062	2,789
1944	41,559	21,173	20,386	6,742	3,425	3,317
1945	44,871	24,738	20,133	6,727	3,919	2,808
1946	64,204	37,099	27,105	9,495	5,867	3,628
1947	53,584	30,514	23,070	7,004	4,323	2,681
1948	53,645	31,504	22,141	6,697	3,864	2,833
1949	52,765	32,191	20,574	6,285	3,621	2,664
1950	56,738	37,871	18,867	6,065	3,711	2,354
1951	57,321	37,857	19,464	5,464	3,601	1,863
1952	58,116	39,162	18,954	5,419	3,607	1,812
1953	61,751	42,961	18,790	5,204	3,515	1,689
1954	61,381	43,268	18,113	5,001	3,424	1,577
1955	66,423	47,618	18,805	4,845	3,372	1,473
1956	66,223	48,140	18,083	5,027	3,452	1,575
1957	62,495	46,868	15,627	4,974	3,350	1,624
1958	36,402	26,998	9,404	5,016	3,485	1,531
1959	20,447	13,547	6,900	5,108	3,557	1,551
1960	21,233	14,029	7,204	5,260	3,553	1,707
1961	21,610	14,438	7,172	5,213	3,545	1,668
1962	22,007	14,432	7,575	5,467	3,827	1,640
1963	22,140	14,508	7,632	5,452	3,930	1,522

* Total number of marriages for 1926 and total number of divorces for 1926 and 1927 furnished by U.S. Bureau of Census; distribution by race not available.

Includes some records on which the race of the couple was not stated.



BIRTHS, FETAL DEATHS, DEATHS

THE COUNTIES AND THE STATE OF MISSISSIPPI, 1963

Footnotes for Tables on Pages 59-141

- * Total births and deaths per 1,000 population. Fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from diseases of early infancy per 1,000 live births. Maternal deaths per 10,000 live births. Deaths from specific causes per 100,000 population.
- # Pneumonia and diarrhea of the newborn (under 28 days) included in diseases of early infancy.



BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
ADAMS COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	1,491	694	797	1,046	464	582	27.9	25.0	30.7
Fetal deaths	60	20	40	27	11	16	25.8	23.7	27.5
Total deaths	500	179	321	391	128	263	10.4	6.9	13.9
Deaths under 1 year	51	12	39	35	10	25	33.5	21.6	43.0
Deaths under 28 days	32	8	24	21	7	14	20.1	15.1	24.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	1	0	1	1	1	0	2.7	-	5.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	1	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	67	29	38	53	20	33	141.2	107.8	174.0
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	2	1	1	1	0	1	2.7	-	5.3
Diabetes mellitus (260)	9	3	6	6	3	3	16.0	16.2	15.8
Anemias (290-293)	4	0	4	3	0	3	8.0	-	15.8
Intracranial lesions (330-334)	69	27	42	56	21	35	149.2	113.1	184.6
Nonmeningococcal meningitis (340)	1	0	1	1	0	1	2.7	-	5.3
Heart diseases (400-443)	170	58	112	147	46	101	391.8	247.8	532.7
Other hypertensive disease (444-447)	0	0	0	1	0	1	2.7	-	5.3
General arteriosclerosis (450)	10	5	5	8	3	5	21.3	16.2	26.4
Other circulatory diseases (451-468)	5	5	0	2	2	0	5.3	10.8	-
#Influenza & pneumonia (480-493)	18	5	13	11	3	8	29.3	16.2	42.2
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	1	0	1	1	0	1	2.7	-	5.3
#Gastritis, etc. (543, 571, 572)	2	1	1	2	1	1	5.3	5.4	5.3
Hernia & intestinal obstruction (560, 561, 570)	7	3	4	2	0	2	5.3	-	10.5
Cirrhosis of liver (581)	1	1	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	1	3	2	1	1	5.3	5.4	5.3
Infections of kidney (600)	10	4	6	4	1	3	10.7	5.4	15.8
Hyperplasia of prostate (610)	6	1	5	5	1	4	13.3	5.4	21.1
Maternal causes (640-689)	1	0	1	0	0	0	-	-	-
Congenital malformations (750-759)	8	4	4	7	3	4	18.6	16.2	21.1
Diseases of early infancy (760-776)	31	7	24	20	6	14	19.1	12.9	24.0
Accidents (800-962)	31	14	17	26	12	14	69.3	64.6	73.8
Motor vehicle acc. (810-835)	14	4	10	12	4	8	32.0	21.6	42.2
Suicide (963, 970-979)	3	2	1	2	1	1	5.3	5.4	5.3
Homicide (964, 980-985)	5	0	5	5	0	5	13.3	-	26.4
Ill-defined (780-795)	19	4	15	14	3	11	37.3	16.2	58.0

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
ALCORN COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	568	464	104	525	421	104	19.6	18.0	30.0
Fetal deaths	8	5	3	7	3	4	13.3	7.1	38.5
Total deaths	273	226	47	277	229	48	10.3	9.8	13.9
Deaths under 1 year	17	11	6	13	7	6	24.8	16.6	57.7
Deaths under 28 days	10	7	3	7	4	3	13.3	9.5	28.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	33	31	2	35	33	2	130.4	141.1	57.8
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	5	3	2	4	2	2	14.9	8.6	57.8
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	64	50	14	65	51	14	242.2	218.1	404.6
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	96	80	16	92	77	15	342.8	329.3	433.5
Other hypertensive disease (444-447)	5	3	2	4	2	2	14.9	8.6	57.8
General arteriosclerosis (450)	1	1	0	1	1	0	3.7	4.3	-
Other circulatory diseases (451-468)	1	1	0	3	3	0	11.2	12.8	-
#Influenza & pneumonia (480-493)	10	7	3	11	8	3	41.0	34.2	86.7
Bronchitis (500-502)	2	2	0	2	2	0	7.4	8.6	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	2	2	0	3	3	0	11.2	12.8	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	3	0	3	3	0	11.2	12.8	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	1	1	0	1	1	0	3.7	4.3	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	3	3	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	8	5	3	7	4	3	13.3	9.5	28.8
Accidents (800-962)	11	11	0	13	11	2	48.4	47.0	57.8
Motor vehicle acc. (810-835)	5	5	0	3	2	1	11.2	8.6	28.9
Suicide (963, 970-979)	2	2	0	2	2	0	7.4	8.6	-
Homicide (964, 980-985)	2	1	1	3	2	1	11.2	8.6	28.9
Ill-defined (780-795)	16	12	4	16	12	4	59.6	51.3	115.6

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
AMITE COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	153	0	153	337	108	229	19.7	13.2	25.6
Fetal deaths	9	0	9	21	4	17	62.3	37.0	74.2
Total deaths	96	28	68	182	87	95	10.6	10.7	10.6
Deaths under 1 year	8	0	8	20	3	17	59.3	27.8	74.2
Deaths under 28 days	2	0	2	10	2	8	29.7	18.5	34.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	1	0	1	1	0	5.8	12.3	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	5.8	-	11.2
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	15	6	9	28	15	13	163.7	184.0	145.2
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	0	2	3	0	3	17.5	-	33.5
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	3	12	26	10	16	152.0	122.7	178.8
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	26	11	15	48	33	15	280.7	404.9	167.6
Other hypertensive disease (444-447)	1	0	1	1	0	1	5.8	-	11.2
General arteriosclerosis (450)	0	0	0	2	2	0	11.7	24.5	-
Other circulatory diseases (451-468)	1	0	1	3	2	1	17.5	24.5	11.2
#Influenza & pneumonia (480-493)	8	1	7	18	5	13	105.3	61.3	145.2
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	1	0	1	2	0	2	11.7	-	22.3
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	1	0	1	5.8	-	11.2
Hyperplasia of prostate (610)	1	0	1	1	0	1	5.8	-	11.2
Maternal causes (640-689)	1	0	1	1	0	1	29.7	-	43.7
Congenital malformations (750-759)	1	0	1	1	0	1	5.8	-	11.2
Diseases of early infancy (760-776)	1	0	1	9	2	7	26.7	18.5	30.6
Accidents (800-962)	9	2	7	15	7	8	87.7	85.9	89.4
Motor vehicle acc. (810-835)	2	1	1	6	4	2	35.1	49.1	22.3
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	0	0	0	1	1	0	5.8	12.3	-
Ill-defined (780-795)	10	1	9	13	2	11	76.0	24.5	122.9

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
ATTALA COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	477	202	275	481	201	280	21.4	15.0	30.8
Fetal deaths	13	7	6	13	7	6	27.0	34.8	21.4
Total deaths	219	127	92	253	153	100	11.2	11.4	11.0
Deaths under 1 year	20	2	18	22	4	18	45.7	19.9	64.3
Deaths under 28 days	14	2	12	16	4	12	33.3	19.9	42.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	31	22	9	35	27	8	155.4	201.2	87.9
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	4	3	1	4	2	2	17.8	14.9	22.0
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	31	23	8	34	24	10	151.0	178.8	109.9
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	82	44	38	94	51	43	417.4	380.0	472.5
Other hypertensive disease (444-447)	2	1	1	3	2	1	13.3	14.9	11.0
General arteriosclerosis (450)	3	3	0	5	5	0	22.2	37.2	-
Other circulatory diseases (451-468)	0	0	0	2	2	0	8.9	14.9	-
#Influenza & pneumonia (480-493)	4	1	3	5	1	4	22.2	7.4	44.0
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	1	0	1	1	0	1	4.4	-	11.0
#Gastritis, etc. (543, 571, 572)	2	0	2	2	0	2	8.9	-	22.0
Hernia & intestinal obstruction (560, 561, 570)	3	3	0	3	3	0	13.3	22.4	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	3	3	0	13.3	22.4	-
Infections of kidney (600)	2	1	1	3	1	2	13.3	7.4	22.0
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	3	1	2	3	1	2	13.3	7.4	22.0
Diseases of early infancy (760-776)	15	2	13	17	4	13	35.3	19.9	46.4
Accidents (800-962)	11	7	4	12	8	4	53.3	59.6	44.0
Motor vehicle acc. (810-835)	5	3	2	5	3	2	22.2	22.4	22.0
Suicide (963, 970-979)	2	2	0	2	2	0	8.9	14.9	-
Homicide (964, 980-985)	1	0	1	1	0	1	4.4	-	11.0
Ill-defined (780-795)	16	8	8	16	9	7	71.0	67.1	76.9

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
BENTON COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	115	1	114	187	55	132	22.9	14.8	29.7
Fetal deaths	6	0	6	17	7	10	90.9	127.3	75.8
Total deaths	46	18	28	78	36	42	9.6	9.7	9.4
Deaths under 1 year	5	0	5	8	1	7	42.8	18.2	53.0
Deaths under 28 days	3	0	3	6	1	5	32.1	18.2	37.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	1	1	0	7	5	2	85.8	134.8	44.9
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	1	0	1	12.2	-	22.5
Diabetes mellitus (260)	1	0	1	2	0	2	24.5	-	44.9
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	7	3	4	14	6	8	171.6	161.7	179.8
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	14	9	5	22	14	8	269.6	377.4	179.8
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	1	1	0	1	1	0	12.2	27.0	-
#Influenza & pneumonia (480-493)	4	1	3	5	2	3	61.3	53.9	67.4
Bronchitis (500-502)	1	0	1	1	0	1	12.2	-	22.5
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	2	2	0	24.5	53.9	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	2	1	1	24.5	27.0	22.5
Infections of kidney (600)	1	0	1	1	0	1	12.2	-	22.5
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	2	0	2	5	1	4	26.7	18.2	30.3
Accidents (800-962)	2	1	1	2	1	1	24.5	27.0	22.5
Motor vehicle acc. (810-835)	0	0	0	0	0	0	-	-	-
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	9	1	8	9	1	8	110.3	27.0	179.8

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
BOLIVAR COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,936	352	1,584	1,802	351	1,451	31.3	17.9	38.1
Fetal deaths	60	6	54	58	7	51	32.2	19.9	35.1
Total deaths	646	135	511	659	171	488	11.4	8.7	12.8
Deaths under 1 year	95	8	87	97	6	91	53.8	17.1	62.7
Deaths under 28 days	55	5	50	54	4	50	30.0	11.4	34.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	0	1	4	1	3	6.9	5.1	7.9
Syphilis (020-029)	1	0	1	1	0	1	1.7	-	2.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	65	15	50	76	28	48	131.9	143.1	126.1
Benign & unspec. neoplasms (210-239)	0	0	0	2	0	2	3.5	-	5.2
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	6	0	6	6	0	6	10.4	-	15.8
Anemias (290-293)	2	0	2	1	0	1	1.7	-	2.6
Intracranial lesions (330-334)	90	18	72	88	22	66	152.7	112.4	173.4
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	196	51	145	196	61	135	340.2	311.7	354.8
Other hypertensive disease (444-447)	3	1	2	4	2	2	6.9	10.2	5.2
General arteriosclerosis (450)	6	1	5	6	2	4	10.4	10.2	10.5
Other circulatory diseases (451-468)	3	0	3	5	1	4	8.7	5.1	10.5
#Influenza & pneumonia (480-493)	30	5	25	33	7	26	57.3	35.8	68.3
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	3	3	0	3	3	0	5.2	15.3	-
#Gastritis, etc. (543, 571, 572)	10	1	9	10	0	10	17.4	-	26.3
Hernia & intestinal obstruction (560, 561, 570)	5	0	5	8	1	7	13.9	5.1	18.4
Cirrhosis of liver (581)	1	1	0	2	2	0	3.5	10.2	-
Acute nephritis (590, 591)	1	0	1	1	0	1	1.7	-	2.6
Chronic nephritis, etc. (592-594)	10	2	8	7	2	5	12.1	10.2	13.1
Infections of kidney (600)	0	0	0	4	2	2	6.9	10.2	5.2
Hyperplasia of prostate (610)	2	0	2	2	0	2	3.5	-	5.2
Maternal causes (640-689)	2	0	2	1	0	1	5.5	-	6.9
Congenital malformations (750-759)	9	3	6	10	2	8	17.4	10.2	21.0
Diseases of early infancy (760-776)	48	3	45	46	2	44	25.5	5.7	30.3
Accidents (800-962)	40	12	28	34	12	22	59.0	61.3	57.8
Motor vehicle acc. (810-835)	17	6	11	15	7	8	26.0	35.8	21.0
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	6	2	4	6	2	4	10.4	10.2	10.5
Ill-defined (780-795)	79	8	71	77	8	69	133.6	40.9	181.3

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
CALHOUN COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	334	148	186	360	205	155	21.6	16.0	40.9
Fetal deaths	17	6	11	10	5	5	27.8	24.4	32.2
Total deaths	143	104	39	163	133	30	9.8	10.4	7.9
Deaths under 1 year	22	8	14	21	9	12	58.3	43.9	77.4
Deaths under 28 days	14	8	6	12	7	5	33.3	34.1	32.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	16	12	4	16	15	1	96.2	116.8	26.4
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	1	1	0	1	1	0	6.0	7.8	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	9	8	1	18	16	2	108.2	124.6	52.8
Nonmeningococcal meningitis (340)	1	1	0	1	1	0	6.0	7.8	-
Heart diseases (400-443)	62	48	14	68	59	9	408.9	459.5	237.5
Other hypertensive disease (444-447)	1	1	0	1	1	0	6.0	7.8	-
General arteriosclerosis (450)	8	6	2	8	6	2	48.1	46.7	52.8
Other circulatory diseases (451-468)	1	0	1	1	0	1	6.0	-	26.4
#Influenza & pneumonia (480-493)	9	6	3	13	10	3	78.2	77.9	79.2
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	1	1	0	6.0	7.8	-
#Gastritis, etc. (543, 571, 572)	2	0	2	1	0	1	6.0	-	26.4
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	1	1	0	1	1	0	6.0	7.8	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	0	1	6.0	-	26.4
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	2	2	0	12.0	15.6	-
Diseases of early infancy (760-776)	12	7	5	10	6	4	27.8	29.3	25.8
Accidents (800-962)	5	3	2	6	4	2	36.1	31.2	52.8
Motor vehicle acc. (810-835)	1	1	0	2	2	0	12.0	15.6	-
Suicide (963, 970-979)	1	1	0	1	1	0	6.0	7.8	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	5	1	4	5	1	4	30.1	7.8	105.5

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
CARROLL COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	142	9	133	212	51	161	17.6	9.2	24.8
Fetal deaths	5	0	5	11	3	8	51.9	58.8	49.7
Total deaths	64	24	40	123	58	65	10.2	10.4	10.0
Deaths under 1 year	3	0	3	8	1	7	37.7	19.6	43.5
Deaths under 28 days	0	0	0	1	1	0	4.7	19.6	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	6	3	3	18	13	5	149.6	234.2	77.2
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	1	0	1	2	0	2	16.6	-	30.9
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	6	3	3	16	8	8	133.0	144.1	123.4
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	12	8	4	34	22	12	282.6	396.4	185.2
Other hypertensive disease (444-447)	6	2	4	6	3	3	49.9	54.0	46.3
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	2	2	0	2	2	0	16.6	36.0	-
#Influenza & pneumonia (480-493)	2	1	1	6	3	3	49.9	54.0	46.3
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	2	0	2	16.6	-	30.9
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	1	0	1	2	1	1	8.3	-	15.4
Maternal causes (640-689)	0	0	0	1	0	1	16.6	18.0	15.4
Congenital malformations (750-759)	1	0	1	1	0	1	8.3	-	15.4
Diseases of early infancy (760-776)	1	0	1	3	1	2	14.2	19.6	12.4
Accidents (800-962)	10	2	8	7	2	5	58.2	36.0	77.2
Motor vehicle acc. (810-835)	6	2	4	3	2	1	24.9	36.0	15.4
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	1	1	0	1	0	1	8.3	-	15.4
Ill-defined (780-795)	13	2	11	18	3	15	149.6	54.0	231.5

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
CHICKASAW COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	408	232	176	412	215	197	23.4	20.2	28.2
Fetal deaths	27	16	11	32	17	15	77.7	79.1	76.1
Total deaths	192	103	89	198	105	93	11.2	9.9	13.3
Deaths under 1 year	16	5	11	13	4	9	31.6	18.6	45.7
Deaths under 28 days	13	5	8	11	4	7	26.7	18.6	35.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	0	1	5.7	-	14.3
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	1	1	0	5.7	9.4	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	24	17	7	27	15	12	153.3	141.2	171.7
Benign & unspec. neoplasms (210-239)	3	2	1	2	1	1	11.4	9.4	14.3
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	0	0	0	0	0	0	-	-	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	36	19	17	34	18	16	193.1	169.5	228.9
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	62	38	24	65	39	26	369.1	367.2	372.0
Other hypertensive disease (444-447)	4	1	3	3	0	3	17.0	-	42.9
General arteriosclerosis (450)	4	2	2	5	3	2	28.4	28.2	28.6
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	9	3	6	10	5	5	56.8	47.1	71.5
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	1	0	1	2	1	1	11.4	9.4	14.3
Hernia & intestinal obstruction (560, 561, 570)	2	1	1	1	1	0	5.7	9.4	-
Cirrhosis of liver (581)	1	0	1	1	0	1	5.7	-	14.3
Acute nephritis (590, 591)	1	0	1	1	0	1	5.7	-	14.3
Chronic nephritis, etc. (592-594)	2	1	1	4	2	2	22.7	18.8	28.6
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	1	0	1	1	0	1	24.3	-	50.8
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	9	3	6	8	3	5	19.4	14.0	25.4
Accidents (800-962)	14	7	7	13	7	6	73.8	65.9	85.8
Motor vehicle acc. (810-835)	7	4	3	6	4	2	34.1	37.7	28.6
Suicide (963, 970-979)	1	1	0	1	1	0	5.7	9.4	-
Homicide (964, 980-985)	1	0	1	1	0	1	5.7	-	14.3
Ill-defined (780-795)	9	1	8	11	3	8	62.5	28.2	114.4

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
CHOCTAW COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	187	101	86	173	94	79	22.0	17.7	31.1
Fetal deaths	6	3	3	4	2	2	23.1	21.3	25.3
Total deaths	94	50	44	95	55	40	12.1	10.4	15.7
Deaths under 1 year	8	2	6	9	3	6	52.0	31.9	75.9
Deaths under 28 days	5	2	3	6	3	3	34.7	31.9	38.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	1	0	1	1	0	1	12.7	-	39.4
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	7	3	4	12	8	4	152.9	150.6	157.5
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	0	0	0	0	0	0	-	-	-
Anemias (290-293)	1	1	0	1	1	0	12.7	18.8	-
Intracranial lesions (330-334)	15	8	7	15	8	7	191.1	150.6	275.6
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	33	22	11	29	20	9	369.4	376.6	354.3
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	5	3	2	3	2	1	38.2	37.7	39.4
Other circulatory diseases (451-468)	0	0	0	1	1	0	12.7	18.8	-
#Influenza & pneumonia (480-493)	6	2	4	6	3	3	76.4	56.5	118.1
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	1	1	0	12.7	18.8	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	1	2	3	1	2	38.2	18.8	78.7
Infections of kidney (600)	2	0	2	1	0	1	12.7	-	39.4
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	6	2	4	7	3	4	40.5	31.9	50.6
Accidents (800-962)	9	4	5	10	5	5	127.4	94.2	196.8
Motor vehicle acc. (810-835)	2	0	2	2	0	2	25.5	-	78.7
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	2	0	2	3	0	3	38.2	-	118.1

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
CLAIBORNE COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	173	25	148	243	46	197	24.8	16.2	28.3
Fetal deaths	10	1	9	17	3	14	70.0	65.2	71.1
Total deaths	86	18	68	113	29	84	11.5	10.2	12.1
Deaths under 1 year	9	0	9	13	0	13	53.5	-	66.0
Deaths under 28 days	8	0	8	12	0	12	49.4	-	60.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	10.2	-	14.4
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	9	1	8	16	4	12	163.3	140.8	172.4
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	1	1	4	2	2	40.8	70.4	28.7
Anemias (290-293)	1	1	0	1	1	0	10.2	35.2	-
Intracranial lesions (330-334)	8	2	6	9	3	6	91.8	105.6	86.2
Nonmeningococcal meningitis (340)	0	0	0	1	0	1	10.2	-	14.4
Heart diseases (400-443)	38	10	28	44	15	29	449.0	528.2	416.7
Other hypertensive disease (444-447)	4	1	3	4	1	3	40.8	35.2	43.1
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	4	0	4	4	0	4	40.8	-	57.5
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	3	1	2	30.6	35.2	28.7
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	1	1	0	3	1	2	123.4	217.4	101.5
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	6	0	6	10	0	10	41.2	-	50.8
Accidents (800-962)	6	0	6	6	0	6	61.2	-	86.2
Motor vehicle acc. (810-835)	2	0	2	2	0	2	20.4	-	28.7
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	1	0	1	1	0	1	10.2	-	14.4
III-defined (780-795)	4	0	4	5	0	5	51.0	-	71.8

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
CLARKE COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	125	64	61	335	167	168	19.6	14.4	30.6
Fetal deaths	5	3	2	10	4	6	29.8	24.0	35.7
Total deaths	130	95	35	208	147	61	12.2	12.7	11.1
Deaths under 1 year	4	1	3	11	3	8	32.8	18.0	47.6
Deaths under 28 days	1	1	0	7	3	4	20.9	18.0	23.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	0	1	1	0	1	5.8	-	18.2
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	15	14	1	24	22	2	140.6	190.0	36.4
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	1	0	1	1	0	5.8	8.6	-
Diabetes mellitus (260)	3	2	1	6	3	3	35.1	25.9	54.6
Anemias (290-293)	1	0	1	0	0	0	-	-	-
Intracranial lesions (330-334)	14	8	6	22	17	5	128.9	146.8	91.1
Nonmeningococcal meningitis (340)	0	0	0	1	0	1	5.8	-	18.2
Heart diseases (400-443)	55	44	11	79	61	18	462.8	526.8	327.9
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	1	0	1	2	0	2	11.7	-	36.4
Other circulatory diseases (451-468)	0	0	0	2	2	0	11.7	17.3	-
#Influenza & pneumonia (480-493)	7	4	3	9	6	3	52.7	51.8	54.6
Bronchitis (500-502)	1	1	0	1	1	0	5.8	8.6	-
Stomach & duodenal ulcers (540, 541)	1	1	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	1	0	1	5.8	-	18.2
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	2	1	1	11.7	8.6	18.2
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	4	2	2	23.4	17.3	36.4
Infections of kidney (600)	0	0	0	1	0	1	5.8	-	18.2
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	2	1	1	59.7	59.9	59.5
Congenital malformations (750-759)	0	0	0	1	1	0	5.8	8.6	-
Diseases of early infancy (760-776)	1	1	0	6	2	4	17.9	12.0	23.8
Accidents (800-962)	10	8	2	14	10	4	82.0	86.4	72.8
Motor vehicle acc. (810-835)	5	4	1	6	5	1	35.1	43.2	18.2
Suicide (963, 970-979)	2	2	0	2	2	0	11.7	17.3	-
Homicide (964, 980-985)	4	1	3	8	2	6	46.9	17.3	109.3
III-defined (780-795)	9	5	4	12	7	5	70.3	60.4	91.1

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
 AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
 CLAY COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	458	193	265	492	211	281	24.9	21.1	28.7
Fetal deaths	14	1	13	16	2	14	32.5	9.5	49.8
Total deaths	177	86	91	193	97	96	9.8	9.7	9.8
Deaths under 1 year	13	5	8	16	6	10	32.5	28.4	35.6
Deaths under 28 days	9	5	4	11	6	5	22.4	28.4	17.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	0	1	1	0	1	5.1	-	10.2
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	1	0	5.1	10.0	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	29	16	13	31	18	13	156.9	180.4	132.9
Benign & unspec. neoplasms (210-239)	1	1	0	1	1	0	5.1	10.0	-
Asthma (241)	1	1	0	1	1	0	5.1	10.0	-
Diabetes mellitus (260)	2	2	0	2	2	0	10.1	20.0	-
Anemias (290-293)	1	0	1	1	0	1	5.1	-	10.2
Intracranial lesions (330-334)	32	13	19	35	14	21	177.1	140.3	214.7
Nonmeningococcal meningitis (340)	0	0	0	2	0	2	10.1	-	20.4
Heart diseases (400-443)	61	33	28	60	34	26	303.6	340.7	265.8
Other hypertensive disease (444-447)	1	1	0	1	1	0	5.1	10.0	-
General arteriosclerosis (450)	6	4	2	7	5	2	35.4	50.1	20.4
Other circulatory diseases (451-468)	1	1	0	2	2	0	10.1	20.0	-
#Influenza & pneumonia (480-493)	5	2	3	5	2	3	25.3	20.0	30.7
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	1	0	1	1	0	1	5.1	-	10.2
#Gastritis, etc. (543, 571, 572)	0	0	0	2	1	1	10.1	10.0	10.2
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	1	1	0	5.1	10.0	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	1	0	1	5.1	-	10.2
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	2	0	2	2	0	2	10.1	-	20.4
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	1	0	1	1	0	1	20.3	-	35.6
Congenital malformations (750-759)	1	1	0	2	2	0	10.1	20.0	-
Diseases of early infancy (760-776)	8	4	4	10	5	5	20.3	23.7	17.8
Accidents (800-962)	10	3	7	11	4	7	55.7	40.1	71.6
Motor vehicle acc. (810-835)	5	2	3	6	3	3	30.4	30.1	30.7
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	6	1	5	5	0	5	25.3	-	51.1

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
COAHOMA COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,520	370	1,150	1,509	327	1,182	31.1	21.6	35.4
Fetal deaths	77	16	61	72	14	58	47.7	42.8	49.1
Total deaths	594	152	442	596	129	467	12.3	8.5	14.0
Deaths under 1 year	104	6	98	107	6	101	70.9	18.3	85.4
Deaths under 28 days	66	3	63	67	4	63	44.4	12.2	53.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	0	1	3	0	3	6.2	-	9.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	59	29	30	57	22	35	117.5	145.1	105.0
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	1	0	1	1	0	2.1	6.6	-
Diabetes mellitus (260)	10	1	9	10	1	9	20.6	6.6	27.0
Anemias (290-293)	2	0	2	2	0	2	4.1	-	6.0
Intracranial lesions (330-334)	71	25	46	69	19	50	142.3	125.3	150.0
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	164	53	111	161	47	114	332.0	310.0	341.9
Other hypertensive disease (444-447)	7	2	5	9	2	7	18.6	13.2	21.0
General arteriosclerosis (450)	3	2	1	3	2	1	6.2	13.2	3.0
Other circulatory diseases (451-468)	5	4	1	5	3	2	10.3	19.8	6.0
#Influenza & pneumonia (480-493)	42	4	38	45	3	42	92.8	19.8	126.0
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	1	0	1	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	8	0	8	9	0	9	18.6	-	27.0
Hernia & intestinal obstruction (560, 561, 570)	2	0	2	3	1	2	6.2	6.6	6.0
Cirrhosis of liver (581)	3	2	1	4	3	1	8.2	19.8	3.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	5	0	5	5	0	5	10.3	-	15.0
Infections of kidney (600)	17	1	16	14	0	14	28.9	-	42.0
Hyperplasia of prostate (610)	1	0	1	3	1	2	6.2	6.6	6.0
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	4	1	3	4	1	3	8.2	6.6	9.0
Diseases of early infancy (760-776)	59	3	56	59	3	56	39.1	9.2	47.4
Accidents (800-962)	37	11	26	31	9	22	63.9	59.4	66.0
Motor vehicle acc. (810-835)	12	5	7	6	2	4	12.4	13.2	12.0
Suicide (963, 970-979)	2	2	0	1	1	0	2.1	6.6	-
Homicide (964, 980-985)	8	1	7	10	1	9	20.6	6.6	27.0
Ill-defined (780-795)	47	0	47	52	0	52	107.2	-	156.0

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
COPIAH COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	840	253	587	668	233	435	24.5	18.2	30.1
Fetal deaths	17	5	12	14	3	11	21.0	12.9	25.3
Total deaths	272	122	150	324	154	170	11.9	12.0	11.8
Deaths under 1 year	26	6	20	24	4	20	35.9	17.2	46.0
Deaths under 28 days	17	6	11	14	4	10	21.0	17.2	23.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	2	1	1	2	1	1	7.3	7.8	6.9
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	24	16	8	27	16	11	99.1	125.1	76.1
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	4	1	3	5	1	4	18.3	7.8	27.7
Anemias (290-293)	2	0	2	2	0	2	7.3	-	13.8
Intracranial lesions (330-334)	34	13	21	47	20	27	172.5	156.4	186.7
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	120	61	59	132	69	63	484.4	539.5	435.7
Other hypertensive disease (444-447)	1	0	1	3	0	3	11.0	-	20.7
General arteriosclerosis (450)	3	2	1	7	6	1	25.7	46.9	6.9
Other circulatory diseases (451-468)	4	2	2	6	3	3	22.0	23.4	20.7
#Influenza & pneumonia (480-493)	16	5	11	19	8	11	69.7	62.5	76.1
Bronchitis (500-502)	2	2	0	2	2	0	7.3	15.6	-
Stomach & duodenal ulcers (540, 541)	0	0	0	3	2	1	11.0	15.6	6.9
#Gastritis, etc. (543, 571, 572)	2	0	2	2	0	2	7.3	-	13.8
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	2	2	0	7.3	15.6	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	1	3	5	1	4	18.3	7.8	27.7
Infections of kidney (600)	3	0	3	4	1	3	14.7	7.8	20.7
Hyperplasia of prostate (610)	1	0	1	1	0	1	3.7	-	6.9
Maternal causes (640-689)	1	0	1	1	0	1	15.0	-	23.0
Congenital malformations (750-759)	5	2	3	4	2	2	14.7	15.6	13.8
Diseases of early infancy (760-776)	13	4	9	12	3	9	18.0	12.9	20.7
Accidents (800-962)	16	8	8	18	8	10	66.0	62.5	69.2
Motor vehicle acc. (810-835)	7	5	2	5	3	2	18.3	23.4	13.8
Suicide (963, 970-979)	1	1	0	1	1	0	3.7	7.8	-
Homicide (964, 980-985)	3	0	3	2	0	2	7.3	-	13.8
Ill-defined (780-795)	7	1	6	7	1	6	25.7	7.8	41.5

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
 AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
 COVINGTON COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	173	117	56	313	163	150	19.3	15.2	27.5
Fetal deaths	4	4	0	13	8	5	41.5	49.1	33.3
Total deaths	125	80	45	193	124	69	11.9	11.5	12.7
Deaths under 1 year	8	2	6	16	4	12	51.1	24.5	80.0
Deaths under 28 days	2	1	1	7	2	5	22.4	12.3	33.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	1	0	1	6.2	-	18.3
Malignant neoplasms (140-205)	13	8	5	22	16	6	135.9	149.0	110.1
Benign & unspec. neoplasms (210-239)	2	1	1	3	2	1	18.5	18.6	18.3
Asthma (241)	1	1	0	1	1	0	6.2	9.3	-
Diabetes mellitus (260)	0	0	0	2	1	1	12.4	9.3	18.3
Anemias (290-293)	1	0	1	1	0	1	6.2	-	18.3
Intracranial lesions (330-334)	24	17	7	34	21	13	210.0	195.5	238.5
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	43	33	10	59	48	11	364.4	446.9	201.8
Other hypertensive disease (444-447)	0	0	0	1	0	1	6.2	-	18.3
General arteriosclerosis (450)	5	3	2	5	3	2	30.9	27.9	36.7
Other circulatory diseases (451-468)	0	0	0	1	1	0	6.2	9.3	-
#Influenza & pneumonia (480-493)	4	2	2	6	3	3	37.0	27.9	55.0
Bronchitis (500-502)	0	0	0	1	1	0	6.2	9.3	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	3	1	2	3	1	2	18.5	9.3	36.7
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	1	0	6.2	9.3	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	2	1	1	2	1	1	12.4	9.3	18.3
Chronic nephritis, etc. (592-594)	4	2	2	6	4	2	37.0	37.2	36.7
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	2	1	1	12.4	9.3	18.3
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	1	0	3	2	1	18.5	18.6	18.3
Diseases of early infancy (760-776)	0	0	0	4	1	3	12.8	6.1	20.0
Accidents (800-962)	14	5	9	19	6	13	117.4	55.9	238.5
Motor vehicle acc. (810-835)	11	4	7	14	4	10	86.5	37.2	183.5
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	4	1	3	6	2	4	37.0	18.6	73.4

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
DESOTO COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	589	70	519	750	265	485	27.5	20.8	33.2
Fetal deaths	14	2	12	18	1	17	24.0	3.8	35.0
Total deaths	156	38	118	237	84	153	8.7	6.6	10.5
Deaths under 1 year	9	0	9	22	4	18	29.3	15.1	37.1
Deaths under 28 days	7	0	7	17	4	13	22.7	15.1	26.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	0	1	2	0	2	7.3	-	13.7
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	21	8	13	34	17	17	124.5	133.8	116.4
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	3	2	1	11.0	15.7	6.8
Diabetes mellitus (260)	1	0	1	3	2	1	11.0	15.7	6.8
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	21	3	18	32	11	21	117.2	86.5	143.8
Nonmeningococcal meningitis (340)	0	0	0	2	0	2	7.3	-	13.7
Heart diseases (400-443)	38	17	21	53	28	25	194.1	220.3	171.2
Other hypertensive disease (444-447)	2	1	1	3	2	1	11.0	15.7	6.8
General arteriosclerosis (450)	8	1	7	9	2	7	33.0	15.7	47.9
Other circulatory diseases (451-468)	0	0	0	1	1	0	3.7	7.9	-
#Influenza & pneumonia (480-493)	18	2	16	21	3	18	76.9	23.6	123.3
Bronchitis (500-502)	1	0	1	1	0	1	3.7	-	6.8
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	2	1	1	7.3	7.9	6.8
Hernia & intestinal obstruction (560, 561, 570)	1	0	1	1	1	0	3.7	7.9	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	1	0	1	3.7	-	6.8
Chronic nephritis, etc. (592-594)	1	0	1	3	0	3	11.0	-	20.5
Infections of kidney (600)	0	0	0	1	1	0	3.7	7.9	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	1	0	1	13.3	-	20.6
Congenital malformations (750-759)	0	0	0	3	2	1	11.0	15.7	6.8
Diseases of early infancy (760-776)	5	0	5	13	3	10	17.3	11.3	20.6
Accidents (800-962)	13	3	10	16	3	13	58.6	23.6	89.0
Motor vehicle acc. (810-835)	5	2	3	5	0	5	18.3	-	34.2
Suicide (963, 970-979)	0	0	0	1	1	0	3.7	7.9	-
Homicide (964, 980-985)	4	0	4	7	0	7	25.6	-	47.9
Ill-defined (780-795)	18	3	15	18	2	16	65.9	15.7	109.6

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
 AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
 FORREST COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,583	1,160	423	1,281	826	455	22.6	19.8	30.7
Fetal deaths	35	21	14	35	14	21	27.3	16.9	46.2
Total deaths	563	387	176	568	366	202	10.0	8.8	13.6
Deaths under 1 year	36	22	14	41	16	25	32.0	19.4	54.9
Deaths under 28 days	26	18	8	29	15	14	22.6	18.2	30.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	4	3	1	7.1	7.2	6.7
Syphilis (020-029)	1	0	1	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	0	1	1.8	-	6.7
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	76	58	18	80	58	22	141.4	138.8	148.4
Benign & unspec. neoplasms (210-239)	2	0	2	4	1	3	7.1	2.4	20.2
Asthma (241)	3	2	1	3	2	1	5.3	4.8	6.7
Diabetes mellitus (260)	10	5	5	13	6	7	23.0	14.4	47.2
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	54	39	15	55	34	21	97.2	81.4	141.7
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	227	161	66	207	141	66	365.8	337.6	445.3
Other hypertensive disease (444-447)	7	2	5	6	3	3	10.6	7.2	20.2
General arteriosclerosis (450)	16	12	4	13	9	4	23.0	21.5	27.0
Other circulatory diseases (451-468)	8	5	3	12	7	5	21.2	16.8	33.7
#Influenza & pneumonia (480-493)	21	10	11	26	12	14	45.9	28.7	94.5
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	2	2	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	4	3	1	6	3	3	10.6	7.2	20.2
Hernia & intestinal obstruction (560, 561, 570)	4	3	1	1	0	1	1.8	-	6.7
Cirrhosis of liver (581)	4	3	1	4	3	1	7.1	7.2	6.7
Acute nephritis (590, 591)	2	1	1	2	1	1	3.5	2.4	6.7
Chronic nephritis, etc. (592-594)	10	8	2	10	8	2	17.7	19.2	13.5
Infections of kidney (600)	3	1	2	4	2	2	7.1	4.8	13.5
Hyperplasia of prostate (610)	2	1	1	3	1	2	5.3	2.4	13.5
Maternal causes (640-689)	1	0	1	0	0	0	-	-	-
Congenital malformations (750-759)	6	6	0	4	3	1	7.1	7.2	6.7
Diseases of early infancy (760-776)	18	13	5	23	11	12	18.0	13.3	26.4
Accidents (800-962)	24	17	7	27	19	8	47.7	45.5	54.0
Motor vehicle acc. (810-835)	6	5	1	8	7	1	14.1	16.8	6.7
Suicide (963, 970-979)	3	2	1	3	2	1	5.3	4.8	6.7
Homicide (964, 980-985)	6	0	6	6	0	6	10.6	-	40.5
Ill-defined (780-795)	26	12	14	25	15	10	44.2	35.9	67.5

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
FRANKLIN COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	138	62	76	175	77	98	19.0	13.5	27.8
Fetal deaths	5	2	3	8	1	7	45.7	13.0	71.4
Total deaths	93	53	40	118	74	44	12.8	13.0	12.5
Deaths under 1 year	5	0	5	7	1	6	40.0	13.0	61.2
Deaths under 28 days	1	0	1	3	1	2	17.1	13.0	20.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	11	8	3	17	12	5	184.2	210.2	142.0
Benign & unspec. neoplasms (210-239)	1	1	0	1	1	0	10.8	17.5	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	2	0	2	2	0	21.7	35.0	-
Anemias (290-293)	1	0	1	1	0	1	10.8	-	28.4
Intracranial lesions (330-334)	17	10	7	15	9	6	162.5	157.6	170.4
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	47	28	19	54	35	19	585.0	613.0	539.8
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	1	1	0	2	2	0	21.7	35.0	-
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	3	1	2	4	1	3	43.3	17.5	85.2
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	2	0	2	3	1	2	32.5	17.5	56.8
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	0	1	10.8	-	28.4
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	2	1	1	21.7	17.5	28.4
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	1	0	1	3	1	2	17.1	13.0	20.4
Accidents (800-962)	2	0	2	5	4	1	54.2	70.0	28.4
Motor vehicle acc. (810-835)	2	0	2	5	4	1	54.2	70.0	28.4
Suicide (963, 970-979)	1	1	0	1	1	0	10.8	17.5	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	4	1	3	5	2	3	54.2	35.0	85.2

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
 AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
 GEORGE COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	343	277	66	316	248	68	28.4	26.2	40.5
Fetal deaths	11	8	3	14	12	2	44.3	48.4	29.4
Total deaths	81	65	16	92	77	15	8.3	8.1	8.9
Deaths under 1 year	2	2	0	3	3	0	9.5	12.1	-
Deaths under 28 days	1	1	0	2	2	0	6.3	8.1	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	1	0	9.0	10.6	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	11	9	2	15	13	2	134.8	137.6	119.0
Benign & unspec. neoplasms (210-239)	1	1	0	1	1	0	9.0	10.6	-
Asthma (241)	0	0	0	1	1	0	9.0	10.6	-
Diabetes mellitus (260)	0	0	0	0	0	0	-	-	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	19	12	7	17	11	6	152.7	116.4	357.1
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	25	19	6	28	22	6	251.6	232.8	357.1
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	1	1	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	3	3	0	2	2	0	18.0	21.2	-
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	2	0	2	2	0	18.0	21.2	-
Infections of kidney (600)	2	2	0	2	2	0	18.0	21.2	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	1	1	0	9.0	10.6	-
Diseases of early infancy (760-776)	1	1	0	2	2	0	6.3	8.1	-
Accidents (800-962)	7	6	1	7	7	0	62.9	74.1	-
Motor vehicle acc. (810-835)	2	2	0	2	2	0	18.0	21.2	-
Suicide (963, 970-979)	1	1	0	1	1	0	9.0	10.6	-
Homicide (964, 980-985)	1	1	0	1	1	0	9.0	10.6	-
III-defined (780-795)	2	2	0	3	3	0	27.0	31.7	-

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
GREENE COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	122	72	50	178	125	53	21.3	18.4	33.8
Fetal deaths	1	0	1	4	0	4	22.5	-	75.5
Total deaths	58	46	12	82	64	18	9.8	9.4	11.5
Deaths under 1 year	1	1	0	3	2	1	16.8	16.0	18.9
Deaths under 28 days	0	0	0	1	1	0	5.6	8.0	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	1	1	0	12.0	14.7	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	11	11	0	13	13	0	155.7	191.7	-
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	3	2	1	2	1	1	24.0	14.7	63.7
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	11	9	2	14	12	2	167.7	177.0	127.4
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	20	13	7	25	17	8	299.4	250.7	509.6
Other hypertensive disease (444-447)	0	0	0	2	1	1	24.0	14.7	63.7
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	2	2	0	4	4	0	47.9	59.0	-
#Influenza & pneumonia (480-493)	2	2	0	4	3	1	47.9	44.2	63.7
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	1	1	0	12.0	14.7	-
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	1	1	0	12.0	14.7	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	12.0	14.7	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	1	0	1	56.2	-	188.7
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	0	0	0	1	1	0	5.6	8.0	-
Accidents (800-962)	2	2	0	4	3	1	47.9	44.2	63.7
Motor vehicle acc. (810-835)	1	1	0	2	2	0	24.0	29.5	-
Suicide (963, 970-979)	1	1	0	1	1	0	12.0	14.7	-
Homicide (964, 980-985)	2	1	1	1	1	0	12.0	14.7	-
Ill-defined (780-795)	0	0	0	0	0	0	-	-	-

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
GRENADA COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	471	154	317	488	174	314	25.5	16.9	35.7
Fetal deaths	33	6	27	35	8	27	71.7	46.0	86.0
Total deaths	221	113	108	205	93	112	10.7	9.0	12.7
Deaths under 1 year	25	1	24	25	1	24	51.2	5.7	76.4
Deaths under 28 days	14	1	13	15	1	14	30.7	5.7	44.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	1	0	1	1	0	1	5.2	-	11.4
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	20	13	7	21	11	10	109.8	106.6	113.6
Benign & unspec. neoplasms (210-239)	1	0	1	1	0	1	5.2	-	11.4
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	3	1	2	1	1	0	5.2	9.7	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	38	18	20	34	14	20	177.8	135.6	227.3
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	64	40	24	58	35	23	303.3	339.1	261.4
Other hypertensive disease (444-447)	5	4	1	4	2	2	20.9	19.4	22.7
General arteriosclerosis (450)	4	3	1	5	3	2	26.2	29.1	22.7
Other circulatory diseases (451-468)	2	2	0	3	2	1	15.7	19.4	11.4
#Influenza & pneumonia (480-493)	18	10	8	12	4	8	62.8	38.8	90.9
Bronchitis (500-502)	0	0	0	1	0	1	5.2	-	11.4
Stomach & duodenal ulcers (540, 541)	1	1	0	1	1	0	5.2	9.7	-
#Gastritis, etc. (543, 571, 572)	3	1	2	4	2	2	20.9	19.4	22.7
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	1	1	0	5.2	9.7	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	5.2	-	11.4
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	1	0	1	1	0	1	5.2	-	-
Maternal causes (640-689)	1	0	1	1	0	1	20.5	-	11.4
Congenital malformations (750-759)	1	1	0	1	1	0	5.2	9.7	-
Diseases of early infancy (760-776)	13	1	12	14	1	13	28.7	5.7	41.4
Accidents (800-962)	16	9	7	16	9	7	83.7	87.2	79.5
Motor vehicle acc. (810-835)	3	2	1	4	3	1	20.9	29.1	11.4
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	1	0	1	1	0	1	5.2	-	11.4
Ill-defined (780-795)	20	3	17	18	2	16	94.1	19.4	181.8

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
HANCOCK COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	219	153	66	344	272	72	21.4	20.7	24.3
Fetal deaths	3	1	2	6	2	4	17.4	7.4	55.6
Total deaths	144	113	31	158	126	32	9.8	9.6	10.8
Deaths under 1 year	11	6	5	12	7	5	34.9	25.7	69.4
Deaths under 28 days	8	3	5	8	3	5	23.2	11.0	69.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	0	1	1	0	1	6.2	-	33.8
Syphilis (020-029)	0	0	0	1	0	1	6.2	-	33.8
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	18	17	1	28	26	2	173.9	197.9	67.6
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	1	0	1	1	0	6.2	7.6	-
Diabetes mellitus (260)	3	2	1	4	3	1	24.8	22.8	33.8
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	26	20	6	26	20	6	161.5	152.2	202.7
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	44	40	4	46	42	4	285.7	319.6	135.1
Other hypertensive disease (444-447)	1	0	1	2	1	1	12.4	7.6	33.8
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	5	4	1	5	4	1	31.0	30.4	33.8
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	1	1	0	6.2	7.6	-
Cirrhosis of liver (581)	0	0	0	2	2	0	12.4	15.2	-
Acute nephritis (590, 591)	0	0	0	1	0	1	6.2	-	33.8
Chronic nephritis, etc. (592-594)	2	2	0	2	2	0	12.4	15.2	-
Infections of kidney (600)	3	3	0	3	3	0	18.6	22.8	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	2	1	1	5	4	1	31.0	30.4	33.8
Diseases of early infancy (760-776)	7	3	4	7	3	4	20.3	11.0	55.6
Accidents (800-962)	11	8	3	6	5	1	37.3	38.0	33.8
Motor vehicle acc. (810-835)	8	5	3	3	2	1	18.6	15.2	33.8
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	18	10	8	15	7	8	93.2	53.3	270.3

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
HARRISON COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	4,100	3,353	747	3,751	3,029	722	29.9	28.8	35.8
Fetal deaths	259	206	53	229	183	46	61.0	60.4	63.7
Total deaths	1,129	892	237	948	716	232	7.6	6.8	11.5
Deaths under 1 year	98	62	36	90	54	36	24.0	17.8	49.9
Deaths under 28 days	69	48	21	63	42	21	16.8	13.9	29.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	0	1	4	3	1	3.2	2.8	5.0
Syphilis (020-029)	4	1	3	3	1	2	2.4	1.0	9.9
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	0.8	-	5.0
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	194	167	27	151	131	20	120.4	124.5	99.1
Benign & unspec. neoplasms (210-239)	2	2	0	2	2	0	1.6	1.9	-
Asthma (241)	2	1	1	1	0	1	0.8	-	5.0
Diabetes mellitus (260)	9	7	2	11	8	3	8.8	7.6	14.9
Anemias (290-293)	1	1	0	1	1	0	0.8	1.0	-
Intracranial lesions (330-334)	107	79	28	104	76	28	82.9	72.2	138.8
Nonmeningococcal meningitis (340)	1	1	0	0	0	0	-	-	-
Heart diseases (400-443)	345	293	52	278	228	50	221.7	216.7	247.8
Other hypertensive disease (444-447)	9	8	1	6	5	1	4.8	4.8	5.0
General arteriosclerosis (450)	7	6	1	4	4	0	3.2	3.8	-
Other circulatory diseases (451-468)	10	8	2	9	6	3	7.2	5.7	14.9
#Influenza & pneumonia (480-493)	58	47	11	35	24	11	27.9	22.8	54.5
Bronchitis (500-502)	7	6	1	8	7	1	6.4	6.6	5.0
Stomach & duodenal ulcers (540, 541)	10	10	0	5	5	0	4.0	4.8	-
#Gastritis, etc. (543, 571, 572)	4	1	3	4	1	3	3.2	1.0	14.9
Hernia & intestinal obstruction (560, 561, 570)	2	0	2	4	2	2	3.2	1.9	9.9
Cirrhosis of liver (581)	15	12	3	7	5	2	5.6	4.8	9.9
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	4	0	5	5	0	4.0	4.8	-
Infections of kidney (600)	5	4	1	7	5	2	5.6	4.8	9.9
Hyperplasia of prostate (610)	1	1	0	1	1	0	0.8	1.0	-
Maternal causes (640-689)	1	0	1	1	0	1	2.7	-	13.8
Congenital malformations (750-759)	15	10	5	13	9	4	10.4	8.6	19.8
Diseases of early infancy (760-776)	57	40	17	53	36	17	14.1	11.9	23.5
Accidents (800-962)	90	71	19	80	58	22	63.8	55.1	109.0
Motor vehicle acc. (810-835)	44	34	10	43	31	12	34.3	29.5	59.5
Suicide (963, 970-979)	8	8	0	8	8	0	6.4	7.6	-
Homicide (964, 980-985)	10	2	8	11	1	10	8.8	1.0	49.6
Ill-defined (780-795)	90	52	38	86	47	39	68.6	44.7	193.3

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
HINDS COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	6,028	3,055	2,973	5,249	2,492	2,757	26.3	21.3	33.4
Fetal deaths	533	331	202	451	293	158	85.9	117.6	57.3
Total deaths	2,502	1,334	1,168	1,684	813	871	8.4	6.9	10.5
Deaths under 1 year	252	89	163	186	56	130	35.4	22.5	47.2
Deaths under 28 days	171	73	98	123	48	75	23.4	19.3	27.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	9	2	7	9	3	6	4.5	2.6	7.2
Syphilis (020-029)	5	2	3	4	1	3	2.0	0.8	3.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	0.5	-	1.2
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	1	1	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	430	281	149	248	165	83	124.2	141.0	100.4
Benign & unspec. neoplasms (210-239)	8	4	4	1	0	1	0.5	-	1.2
Asthma (241)	4	2	2	3	1	2	1.5	0.8	2.4
Diabetes mellitus (260)	34	18	16	23	13	10	11.5	11.1	12.1
Anemias (290-293)	7	1	6	5	1	4	2.5	0.8	4.8
Intracranial lesions (330, 334)	266	142	124	177	82	95	88.6	70.1	114.9
Nonmeningococcal meningitis (340)	16	0	16	11	0	11	5.5	-	13.3
Heart diseases (400-443)	682	395	287	502	274	228	251.4	234.1	275.8
Other hypertensive disease (444-447)	21	4	17	12	1	11	6.0	0.8	13.3
General arteriosclerosis (450)	25	21	4	19	15	4	9.5	12.8	4.8
Other circulatory diseases (451-468)	42	26	16	23	11	12	11.5	9.4	14.5
#Influenza & pneumonia (480-493)	73	28	45	63	20	43	31.6	17.1	52.0
Bronchitis (500-502)	9	8	1	5	3	2	2.5	2.6	2.4
Stomach & duodenal ulcers (540, 541)	13	9	4	6	4	2	3.0	3.4	2.4
#Gastritis, etc. (543, 571, 572)	23	10	13	13	3	10	6.5	2.6	12.1
Hernia & intestinal obstruction (560, 561, 570)	9	7	2	4	3	1	2.0	2.6	1.2
Cirrhosis of liver (581)	22	20	2	13	12	1	6.5	10.2	1.2
Acute nephritis (590, 591)	4	0	4	4	0	4	2.0	-	4.8
Chronic nephritis, etc. (592-594)	35	11	24	16	5	11	8.0	4.3	13.3
Infections of kidney (600)	26	12	14	14	7	7	7.0	6.0	8.5
Hyperplasia of prostate (610)	8	2	6	3	0	3	1.5	-	3.6
Maternal causes (640-689)	2	1	1	1	0	1	1.9	-	3.6
Congenital malformations (750-759)	65	35	30	42	22	20	21.0	18.8	24.2
Diseases of early infancy (760-776)	140	53	87	100	33	67	19.0	13.2	24.3
Accidents (800-962)	182	106	76	116	59	57	58.1	50.4	69.0
Motor vehicle acc. (810-835)	84	63	21	50	36	14	25.0	30.8	16.9
Suicide (963, 970-979)	8	6	2	4	3	1	2.0	2.6	1.2
Homicide (964, 980-985)	36	3	33	34	2	32	17.0	1.7	38.7
Ill-defined (780-795)	133	28	105	124	26	98	62.1	22.2	118.6

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
HOLMES COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	786	130	656	756	113	643	28.0	14.4	33.5
Fetal deaths	33	6	27	34	3	31	45.0	26.5	48.2
Total deaths	307	81	226	335	101	234	12.4	12.8	12.2
Deaths under 1 year	27	1	26	28	3	25	37.0	26.5	38.9
Deaths under 28 days	9	1	8	11	2	9	14.6	17.7	14.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	0	1	2	1	1	7.4	12.7	5.2
Syphilis (020-029)	1	1	0	1	1	0	3.7	12.7	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	38	8	30	47	13	34	173.8	165.4	177.3
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	0	1	1	0	1	3.7	-	5.2
Diabetes mellitus (260)	5	1	4	5	1	4	18.5	12.7	20.8
Anemias (290-293)	1	0	1	2	1	1	7.4	12.7	5.2
Intracranial lesions (330-334)	34	6	28	34	6	28	125.7	76.3	146.0
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	87	35	52	91	41	50	336.5	521.6	260.7
Other hypertensive disease (444-447)	2	1	1	3	1	2	11.1	12.7	10.4
General arteriosclerosis (450)	5	3	2	7	4	3	25.9	50.9	15.6
Other circulatory diseases (451-468)	3	0	3	4	0	4	14.8	-	20.8
#Influenza & pneumonia (480-493)	23	6	17	27	10	17	99.8	127.2	88.6
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	1	0	1	2	2	0	7.4	25.4	-
Hernia & intestinal obstruction (560, 561, 570)	1	0	1	1	0	1	3.7	-	5.2
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	1	0	1	3.7	-	5.2
Chronic nephritis, etc. (592-594)	5	1	4	6	1	5	22.2	12.7	26.1
Infections of kidney (600)	3	2	1	4	2	2	14.8	25.4	10.4
Hyperplasia of prostate (610)	1	0	1	2	1	1	7.4	12.7	5.2
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	2	0	2	2	0	2	7.4	-	10.4
Diseases of early infancy (760-776)	10	1	9	10	2	8	13.2	17.7	12.4
Accidents (800-962)	15	5	10	14	4	10	51.8	50.9	52.1
Motor vehicle acc. (810-835)	6	3	3	2	0	2	7.4	-	10.4
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	4	1	3	4	1	3	14.8	12.7	15.6
Ill-defined (780-795)	57	7	50	55	6	49	203.4	76.3	255.5

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
HUMPHREYS COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	550	66	484	574	95	479	29.6	15.6	36.0
Fetal deaths	14	0	14	22	6	16	38.3	63.2	33.4
Total deaths	188	47	141	219	63	156	11.3	10.3	11.7
Deaths under 1 year	34	4	30	39	3	36	67.9	31.6	75.2
Deaths under 28 days	18	2	16	21	3	18	36.6	31.6	37.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	1	0	1	1	0	1	5.2	-	7.5
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	1	0	5.2	16.4	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	21	8	13	27	15	12	139.3	246.3	90.3
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	1	0	1	1	0	5.2	16.4	-
Diabetes mellitus (260)	2	1	1	4	1	3	20.6	16.4	22.6
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	25	3	22	24	4	20	123.8	65.7	150.5
Nonmeningococcal meningitis (340)	1	1	0	1	0	1	5.2	-	7.5
Heart diseases (400-443)	44	16	28	53	17	36	273.5	279.1	270.9
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	4	1	3	3	1	2	15.5	16.4	15.0
Other circulatory diseases (451-468)	1	0	1	2	0	2	10.3	-	15.0
#Influenza & pneumonia (480-493)	18	2	16	16	1	15	82.6	16.4	112.9
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	2	0	2	4	0	4	20.6	-	30.1
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	0	1	5.2	-	7.5
Cirrhosis of liver (581)	0	0	0	1	0	1	5.2	-	7.5
Acute nephritis (590, 591)	1	0	1	2	1	1	10.3	16.4	7.3
Chronic nephritis, etc. (592-594)	6	0	6	8	1	7	41.3	16.4	52.7
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	1	0	1	1	0	1	17.4	-	20.9
Congenital malformations (750-759)	4	0	4	3	0	3	15.5	-	22.6
Diseases of early infancy (760-776)	14	2	12	17	3	14	29.6	31.6	29.2
Accidents (800-962)	15	6	9	18	10	8	92.9	164.2	60.2
Motor vehicle acc. (810-835)	5	2	3	10	7	3	51.6	114.9	22.6
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	4	2	2	4	2	2	20.6	32.8	15.0
III-defined (780-795)	19	1	18	20	1	19	103.2	16.4	143.0

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
ISSAQUNA COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	38	0	38	80	15	65	29.7	17.6	35.3
Fetal deaths	1	0	1	2	1	1	25.0	66.7	15.4
Total deaths	25	2	23	39	7	32	14.5	8.2	17.4
Deaths under 1 year	4	0	4	4	0	4	50.0	-	61.5
Deaths under 28 days	1	0	1	1	0	1	12.5	-	15.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	5	2	3	9	3	6	334.6	352.9	326.1
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	1	0	1	1	0	1	37.2	-	54.3
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	1	0	1	3	0	3	111.5	-	163.0
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	9	0	9	14	4	10	520.4	470.6	543.5
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	1	0	1	1	0	1	37.2	-	54.3
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	1	0	1	1	0	1	12.5	-	15.4
Accidents (800-962)	2	0	2	3	0	3	111.5	-	163.0
Motor vehicle acc. (810-835)	1	0	1	1	0	1	37.2	-	54.3
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	3	0	3	4	0	4	148.7	-	217.4

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
ITAWAMBA COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	64	45	19	254	237	17	14.6	14.4	18.5
Fetal deaths	1	1	0	5	5	0	19.7	21.1	-
Total deaths	111	101	10	152	142	10	8.8	8.6	10.9
Deaths under 1 year	6	2	4	7	4	3	27.6	16.9	176.5
Deaths under 28 days	3	1	2	5	3	2	19.7	12.6	117.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	1	0	1	1	0	5.8	6.1	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	19	18	1	25	24	1	144.1	146.1	108.7
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	1	1	0	5.8	6.1	-
Diabetes mellitus (260)	1	1	0	1	1	0	5.8	6.1	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	18	17	1	26	25	1	149.8	152.2	108.7
Nonmeningococcal meningitis (340)	0	0	0	2	2	0	11.5	12.2	-
Heart diseases (400-443)	48	44	4	58	53	5	334.3	322.6	543.5
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	1	1	0	3	3	0	17.3	18.2	-
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	1	1	0	3	3	0	17.3	18.2	-
Bronchitis (500-502)	1	1	0	1	1	0	5.8	6.1	-
Stomach & duodenal ulcers (540, 541)	1	1	0	1	1	0	5.8	6.1	-
#Gastritis, etc. (543, 571, 572)	1	0	1	0	0	0	5.8	6.1	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	1	0	5.8	6.1	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	1	0	1	1	0	5.8	6.1	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	2	2	0	3	3	0	17.3	18.2	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	1	0	2	2	0	11.5	12.2	-
Diseases of early infancy (760-776)	3	1	2	4	2	2	15.7	8.4	117.6
Accidents (800-962)	7	6	1	12	11	1	69.2	67.0	108.7
Motor vehicle acc. (810-835)	3	3	0	5	5	0	28.8	30.4	-
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	2	2	0	2	2	0	11.5	12.2	-

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
JACKSON COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,454	1,040	414	1,822	1,370	452	27.2	24.8	38.8
Fetal deaths	81	54	27	101	71	30	55.4	51.8	66.4
Total deaths	362	267	95	449	343	106	6.7	6.2	9.1
Deaths under 1 year	45	23	22	59	36	23	32.4	26.3	50.9
Deaths under 28 days	31	16	15	42	26	16	23.0	19.0	35.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	1	0	2	1	1	3.0	1.8	8.6
Syphilis (020-029)	2	1	1	3	2	1	4.5	3.6	8.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	1	0	1	1	0	1.5	1.8	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	39	27	12	66	48	18	98.5	86.7	154.6
Benign & unspec. neoplasms (210-239)	2	1	1	4	3	1	6.0	5.4	8.6
Asthma (241)	1	1	0	3	3	0	4.5	5.4	-
Diabetes mellitus (260)	8	7	1	8	7	1	11.9	12.6	8.6
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	28	21	7	36	31	5	53.7	56.0	43.0
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	108	88	20	124	102	22	185.1	184.3	189.0
Other hypertensive disease (444-447)	5	3	2	5	3	2	7.5	5.4	17.2
General arteriosclerosis (450)	5	5	0	6	5	1	9.0	9.0	8.6
Other circulatory diseases (451-468)	4	4	0	5	4	1	7.5	7.2	8.6
#Influenza & pneumonia (480-493)	18	11	7	23	16	7	34.3	28.9	60.1
Bronchitis (500-502)	1	1	0	2	2	0	3.0	3.6	-
Stomach & duodenal ulcers (540, 541)	2	2	0	3	3	0	4.5	5.4	-
#Gastritis, etc. (543, 571, 572)	1	1	0	1	1	0	1.5	1.8	-
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	1	1	0	1.5	1.8	-
Cirrhosis of liver (581)	3	3	0	3	3	0	4.5	5.4	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	1	0	1	1.5	-	8.6
Infections of kidney (600)	4	3	1	4	3	1	6.0	5.4	8.6
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	6	4	2	10	7	3	14.9	12.6	25.8
Diseases of early infancy (760-776)	24	13	11	34	22	12	18.7	16.0	26.5
Accidents (800-962)	46	32	14	48	35	13	71.6	63.2	111.7
Motor vehicle acc. (810-835)	22	14	8	22	15	7	32.8	27.1	60.1
Suicide (963, 970-979)	2	2	0	3	3	0	4.5	5.4	-
Homicide (964, 980-985)	9	2	7	9	3	6	13.4	5.4	51.5
Ill-defined (780-795)	31	25	6	30	23	7	44.8	41.6	60.1

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
JASPER COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	177	99	78	399	148	251	23.3	16.4	31.0
Fetal deaths	9	7	2	18	11	7	45.1	74.3	27.9
Total deaths	124	66	58	165	84	81	9.6	9.3	10.0
Deaths under 1 year	5	0	5	13	1	12	32.6	6.8	47.8
Deaths under 28 days	2	0	2	8	0	8	20.0	-	31.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	0	1	5.8	-	12.3
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	25	9	16	32	14	18	186.8	155.0	222.2
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	1	1	0	5.8	11.1	-
Diabetes mellitus (260)	3	3	0	5	3	2	29.2	33.2	24.7
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	13	8	5	14	8	6	81.7	88.6	74.1
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	41	26	15	49	30	19	286.0	332.2	234.6
Other hypertensive disease (444-447)	0	0	0	2	0	2	11.7	-	24.7
General arteriosclerosis (450)	2	2	0	3	3	0	17.5	33.2	-
Other circulatory diseases (451-468)	1	1	0	2	1	1	11.7	11.1	12.3
#Influenza & pneumonia (480-493)	7	4	3	8	3	5	46.7	33.2	61.7
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	1	0	1	2	0	2	11.7	-	24.7
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	1	1	0	5.8	11.1	-
Cirrhosis of liver (581)	0	0	0	2	2	0	11.7	22.1	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	5.8	11.1	-
Infections of kidney (600)	2	1	1	4	2	2	23.4	22.1	24.7
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	0	1	2	0	2	11.7	-	24.7
Diseases of early infancy (760-776)	2	0	2	7	0	7	17.5	-	27.9
Accidents (800-962)	10	4	6	11	6	5	64.2	66.4	61.7
Motor vehicle acc. (810-835)	5	2	3	7	4	3	40.9	44.3	37.0
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	8	2	6	8	2	6	46.7	22.1	74.1

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
JEFFERSON COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	105	0	105	247	41	206	24.3	12.6	29.8
Fetal deaths	2	0	2	10	2	8	40.5	48.8	38.8
Total deaths	83	19	64	136	44	92	13.4	13.5	13.3
Deaths under 1 year	4	0	4	6	0	6	24.3	-	29.1
Deaths under 28 days	1	0	1	1	0	1	4.0	-	4.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	7	2	5	14	8	6	137.9	246.2	87.0
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	0	0	0	1	0	1	9.8	-	14.5
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	11	2	9	15	2	13	147.8	61.5	188.4
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	27	5	22	49	14	35	482.8	430.8	507.2
Other hypertensive disease (444-447)	1	0	1	1	0	1	9.8	-	14.5
General arteriosclerosis (450)	4	2	2	5	3	2	49.3	92.3	29.0
Other circulatory diseases (451-468)	2	0	2	2	0	2	19.7	-	29.0
#Influenza & pneumonia (480-493)	9	1	8	11	1	10	108.4	30.8	144.9
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	4	3	1	39.4	92.3	14.5
Cirrhosis of liver (581)	1	0	1	2	1	1	19.7	30.8	14.5
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	4	2	2	39.4	61.5	29.0
Infections of kidney (600)	0	0	0	2	0	2	19.7	-	29.0
Hyperplasia of prostate (610)	1	0	1	3	1	2	29.6	30.8	29.0
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	1	0	1	1	0	1	4.0	-	4.8
Accidents (800-962)	5	2	3	7	3	4	69.0	92.3	58.0
Motor vehicle acc. (810-835)	1	0	1	1	1	0	9.8	30.8	-
Suicide (963, 970-979)	1	1	0	1	1	0	9.8	30.8	-
Homicide (964, 980-985)	2	1	1	2	1	1	19.7	30.8	14.5
Ill-defined (780-795)	6	0	6	7	0	7	69.0	-	101.4

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
JEFF DAVIS COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	291	159	132	268	117	151	20.0	17.2	22.8
Fetal deaths	12	2	10	17	3	14	63.4	25.6	92.7
Total deaths	107	51	56	136	69	67	10.1	10.1	10.1
Deaths under 1 year	11	1	10	12	1	11	44.8	8.5	72.8
Deaths under 28 days	5	1	4	6	1	5	22.4	8.5	33.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	1	0	7.4	14.7	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	9	7	2	11	9	2	81.9	132.4	30.2
Benign & unspec. neoplasms (210-239)	2	0	2	2	0	2	14.9	-	30.2
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	4	1	3	4	1	3	29.8	14.7	45.2
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	21	9	12	25	11	14	186.2	161.8	211.2
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	39	23	16	48	28	20	357.4	411.8	301.6
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	1	0	1	1	0	1	7.4	-	15.1
Other circulatory diseases (451-468)	1	0	1	3	1	2	22.3	14.7	30.2
#Influenza & pneumonia (480-493)	4	0	4	4	0	4	29.8	-	60.3
Bronchitis (500-502)	1	0	1	1	0	1	7.4	-	15.1
Stomach & duodenal ulcers (540, 541)	1	1	0	1	1	0	7.4	14.7	-
#Gastritis, etc. (543, 571, 572)	1	0	1	1	0	1	7.4	-	15.1
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	1	0	2	1	1	14.9	14.7	15.1
Diseases of early infancy (760-776)	3	1	2	4	1	3	14.9	8.5	19.9
Accidents (800-962)	5	3	2	11	7	4	81.9	102.9	60.3
Motor vehicle acc. (810-835)	1	1	0	2	2	0	14.9	29.4	-
Suicide (963, 970-979)	1	1	0	1	1	0	7.4	14.7	-
Homicide (964, 980-985)	3	0	3	4	0	4	29.8	-	60.3
Ill-defined (780-795)	7	1	6	7	2	5	52.1	29.4	75.4

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
JONES COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	2,114	1,017	1,097	1,263	788	475	21.4	18.0	31.3
Fetal deaths	99	40	59	59	30	29	46.7	38.1	61.0
Total deaths	743	448	295	587	412	175	9.9	9.4	11.5
Deaths under 1 year	97	22	75	44	15	29	34.8	19.0	61.0
Deaths under 28 days	66	17	49	28	12	16	22.2	15.2	33.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	4	2	2	5	4	1	8.5	9.1	6.6
Syphilis (020-029)	1	1	0	1	1	0	1.7	2.3	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	1.7	-	6.6
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	2	1	1	1	1	0	1.7	2.3	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	3	2	1	1	1	0	1.7	2.3	-
Malignant neoplasms (140-205)	86	64	22	75	58	17	127.0	132.1	112.0
Benign & unspec. neoplasms (210-239)	2	1	1	1	1	0	1.7	2.3	-
Asthma (241)	3	3	0	1	1	0	1.7	2.3	-
Diabetes mellitus (260)	17	9	8	10	8	2	16.9	18.2	13.2
Anemias (290-293)	3	1	2	3	1	2	5.1	2.3	13.2
Intracranial lesions (330-334)	105	69	36	84	59	25	142.2	134.4	164.7
Nonmeningococcal meningitis (340)	3	1	2	2	1	1	3.4	2.3	6.6
Heart diseases (400-443)	174	125	49	172	135	37	291.2	307.6	243.7
Other hypertensive disease (444-447)	12	4	8	5	2	3	8.5	4.6	19.8
General arteriosclerosis (450)	8	8	0	9	9	0	15.2	20.5	-
Other circulatory diseases (451-468)	10	5	5	7	5	2	11.8	11.4	13.2
#Influenza & pneumonia (480-493)	43	17	26	26	15	11	44.0	34.2	72.5
Bronchitis (500-502)	2	2	0	1	1	0	1.7	2.3	-
Stomach & duodenal ulcers (540, 541)	1	0	1	1	0	1	1.7	-	6.6
#Gastritis, etc. (543, 571, 572)	10	1	9	4	1	3	6.8	2.3	19.8
Hernia & intestinal obstruction (560, 561, 570)	4	2	2	1	1	0	1.7	2.3	-
Cirrhosis of liver (581)	6	6	0	7	7	0	11.8	15.9	-
Acute nephritis (590, 591)	4	1	3	3	1	2	5.1	2.3	13.2
Chronic nephritis, etc. (592-594)	4	4	0	5	4	1	8.5	9.1	6.6
Infections of kidney (600)	10	6	4	5	4	1	8.5	9.1	6.6
Hyperplasia of prostate (610)	2	0	2	1	0	1	8.5	9.1	6.6
Maternal causes (640-689)	3	0	3	1	0	1	1.7	-	6.6
Congenital malformations (750-759)	11	6	5	10	7	3	16.9	15.9	21.0
Diseases of early infancy (760-776)	60	15	45	26	8	18	20.6	10.2	37.9
Accidents (800-962)	56	33	23	38	25	13	64.3	57.0	85.6
Motor vehicle acc. (810-835)	30	16	14	18	10	8	30.5	22.8	52.7
Suicide (963, 970-979)	3	2	1	3	2	1	5.1	4.6	6.6
Homicide (964, 980-985)	9	1	8	8	2	6	13.5	4.6	39.5
Ill-defined (780-795)	31	22	9	30	19	11	50.8	43.3	72.5

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
KEMPER COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	202	38	164	251	58	193	21.5	11.8	28.5
Fetal deaths	8	1	7	11	0	11	43.8	-	57.0
Total deaths	105	55	50	143	80	63	12.2	16.3	9.3
Deaths under 1 year	7	0	7	7	0	7	27.9	-	36.3
Deaths under 28 days	4	0	4	4	0	4	15.9	-	20.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	7	3	4	14	4	10	120.0	81.6	147.7
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	4	3	1	5	4	1	42.8	81.6	14.8
Anemias (290-293)	2	2	0	2	2	0	17.1	40.8	-
Intracranial lesions (330-334)	9	5	4	20	15	5	171.4	306.1	73.8
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	47	27	20	49	28	21	419.9	571.4	310.2
Other hypertensive disease (444-447)	0	0	0	1	0	1	8.6	-	14.8
General arteriosclerosis (450)	0	0	0	2	1	1	17.1	20.4	14.8
Other circulatory diseases (451-468)	1	1	0	1	1	0	8.6	20.4	-
#Influenza & pneumonia (480-493)	6	4	2	11	9	2	94.2	183.7	29.5
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	1	0	1	8.6	-	14.8
#Gastritis, etc, (543, 571, 572)	0	0	0	1	0	1	8.6	-	14.8
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	1	0	8.6	20.4	-
Cirrhosis of liver (581)	0	0	0	1	1	0	8.6	20.4	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	8.6	-	14.8
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	1	1	0	8.6	20.4	-
Maternal causes (640-689)	1	0	1	1	0	1	39.8	-	51.8
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	2	0	2	2	0	2	8.0	-	10.4
Accidents (800-962)	8	4	4	7	3	4	60.0	61.2	59.1
Motor vehicle acc. (810-835)	1	0	1	1	0	1	8.6	-	14.8
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	3	0	3	2	0	2	17.1	-	29.5
III-defined (780-795)	12	5	7	14	6	8	120.0	122.4	118.2

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
LAFAYETTE COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	486	246	240	514	259	255	23.2	17.6	34.2
Fetal deaths	22	4	18	21	3	18	40.8	11.6	70.6
Total deaths	162	89	73	202	122	80	9.1	8.3	10.7
Deaths under 1 year	17	4	13	20	6	14	38.9	23.2	54.9
Deaths under 28 days	10	3	7	11	4	7	21.4	15.4	27.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	0	1	1	0	1	4.5	-	13.4
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	16	9	7	22	14	8	99.2	95.2	107.2
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	2	0	4	3	1	18.0	20.4	13.4
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	24	9	15	32	16	16	144.3	108.8	214.5
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	52	37	15	59	44	15	266.1	299.1	201.1
Other hypertensive disease (444-447)	2	0	2	4	2	2	18.0	13.6	26.8
General arteriosclerosis (450)	6	6	0	7	7	0	31.6	47.6	-
Other circulatory diseases (451-468)	1	1	0	4	3	1	18.0	20.4	13.4
#Influenza & pneumonia (480-493)	9	4	5	10	4	6	45.1	27.2	80.4
Bronchitis (500-502)	0	0	0	2	2	0	9.0	13.6	-
Stomach & duodenal ulcers (540, 541)	1	1	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	1	1	0	2	1	1	9.0	6.8	13.4
Hernia & intestinal obstruction (560, 561, 570)	3	2	1	3	2	1	13.5	13.6	13.4
Cirrhosis of liver (581)	0	0	0	1	0	1	4.5	-	13.4
Acute nephritis (590, 591)	1	0	1	1	0	1	4.5	-	13.4
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	1	0	3	2	1	13.5	13.6	13.4
Diseases of early infancy (760-776)	12	2	10	13	3	10	25.3	11.6	39.2
Accidents (800-962)	10	5	5	13	9	4	58.6	61.2	53.6
Motor vehicle acc. (810-835)	6	3	3	7	4	3	31.6	27.2	40.2
Suicide (963, 970-979)	2	2	0	1	1	0	4.5	6.8	-
Homicide (964, 980-985)	3	1	2	3	1	2	13.5	6.8	26.8
Ill-defined (780-795)	8	1	7	8	1	7	36.1	6.8	93.8

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
LAMAR COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	111	81	30	293	236	57	19.7	18.6	26.3
Fetal deaths	2	1	1	9	6	3	30.7	25.4	52.6
Total deaths	86	69	17	136	112	24	9.1	8.8	11.0
Deaths under 1 year	3	2	1	7	5	2	23.9	21.2	35.1
Deaths under 28 days	1	0	1	5	3	2	17.1	12.7	35.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	1	0	6.7	7.9	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	17	15	2	27	22	5	181.3	173.0	230.4
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	1	0	0	0	0	-	-	-
Diabetes mellitus (260)	0	0	0	1	1	0	6.7	7.9	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	19	14	5	27	22	5	181.3	173.0	230.4
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	24	20	4	38	33	5	255.2	259.4	230.4
Other hypertensive disease (444-447)	0	0	0	1	0	1	6.7	-	46.1
General arteriosclerosis (450)	2	2	0	4	4	0	26.9	31.4	-
Other circulatory diseases (451-468)	1	1	0	1	1	0	6.7	7.9	-
#Influenza & pneumonia (480-493)	4	4	0	6	5	1	40.3	39.3	46.1
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	2	2	0	13.4	15.7	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	2	2	0	13.4	15.7	-
Cirrhosis of liver (581)	1	1	0	1	1	0	6.7	7.9	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	6.7	7.9	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	1	1	0	6.7	7.9	-
Diseases of early infancy (760-776)	1	0	1	3	2	1	10.2	8.5	17.5
Accidents (800-962)	10	7	3	9	7	2	60.4	55.0	92.2
Motor vehicle acc. (810-835)	7	5	2	7	6	1	47.0	47.2	46.1
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	1	0	1	2	1	1	13.4	7.9	46.1

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
LAUDERDALE COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	2,092	1,206	886	1,698	999	699	23.1	19.8	30.0
Fetal deaths	61	17	44	44	13	31	25.9	13.0	44.3
Total deaths	996	617	379	804	454	350	10.9	9.0	15.0
Deaths under 1 year	87	34	53	72	33	39	42.4	33.0	55.8
Deaths under 28 days	61	32	29	47	29	18	27.7	29.0	25.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	1	0	3	1	2	4.1	2.0	8.6
Syphilis (020-029)	1	0	1	1	0	1	1.4	-	4.3
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	2	1	1	1	0	1	1.4	-	4.3
Malignant neoplasms (140-205)	109	74	35	91	58	33	123.6	115.2	141.8
Benign & unspec. neoplasms (210-239)	1	0	1	1	0	1	1.4	-	4.3
Asthma (241)	1	1	0	1	1	0	1.4	2.0	-
Diabetes mellitus (260)	22	8	14	16	5	11	21.7	9.9	47.2
Anemias (290-293)	1	1	0	1	1	0	1.4	2.0	-
Intracranial lesions (330-334)	143	111	32	102	65	37	138.5	129.1	158.9
Nonmeningococcal meningitis (340)	3	1	2	2	1	1	2.7	2.0	4.3
Heart diseases (400-443)	294	190	104	250	148	102	339.6	294.0	438.1
Other hypertensive disease (444-447)	5	2	3	5	2	3	6.8	4.0	12.9
General arteriosclerosis (450)	19	16	3	14	11	3	19.0	21.8	12.9
Other circulatory diseases (451-468)	7	5	2	6	3	3	8.1	6.0	12.9
#Influenza & pneumonia (480-493)	48	34	14	30	16	14	40.7	31.8	60.1
Bronchitis (500-502)	2	2	0	1	1	0	1.4	2.0	-
Stomach & duodenal ulcers (540, 541)	2	2	0	2	2	0	2.7	4.0	-
#Gastritis, etc. (543, 571, 572)	5	0	5	3	0	3	4.1	-	12.9
Hernia & intestinal obstruction (560, 561, 570)	13	6	7	9	4	5	12.2	7.9	21.5
Cirrhosis of liver (581)	6	6	0	5	5	0	6.8	9.9	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	10	4	6	8	4	4	10.9	7.9	17.2
Infections of kidney (600)	11	7	4	8	4	4	10.9	7.9	17.2
Hyperplasia of prostate (610)	3	2	1	3	2	1	4.1	4.0	4.3
Maternal causes (640-689)	3	1	2	2	0	2	11.8	-	28.6
Congenital malformations (750-759)	6	5	1	4	3	1	5.4	6.0	4.3
Diseases of early infancy (760-776)	52	26	26	40	26	14	23.6	26.0	20.0
Accidents (800-962)	62	44	18	53	36	17	72.0	71.5	73.0
Motor vehicle acc. (810-835)	30	26	4	22	18	4	29.9	35.8	17.2
Suicide (963, 970-979)	7	6	1	7	6	1	9.5	11.9	4.3
Homicide (964, 980-985)	5	1	4	7	3	4	9.5	6.0	17.2
Ill-defined (780-795)	113	38	75	104	33	71	141.3	65.6	305.0

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
LAWRENCE COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	137	49	88	210	109	101	22.4	17.8	31.2
Fetal deaths	4	1	3	12	2	10	57.1	18.3	99.0
Total deaths	89	53	36	124	80	44	13.2	13.1	13.6
Deaths under 1 year	6	2	4	8	2	6	38.1	18.3	59.4
Deaths under 28 days	3	2	1	5	2	3	23.8	18.3	29.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	11	9	2	16	14	2	170.9	228.8	61.7
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	4	3	1	4	3	1	42.7	49.0	30.9
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	16	6	10	22	12	10	235.0	196.1	308.6
Nonmeningococcal meningitis (340)	0	0	0	1	0	1	10.7	-	30.9
Heart diseases (400-443)	31	20	11	41	30	11	438.0	490.2	339.5
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	1	1	0	1	1	0	10.7	16.3	-
#Influenza & pneumonia (480-493)	1	0	1	6	3	3	64.1	49.0	92.6
Bronchitis (500-502)	0	0	0	1	1	0	10.7	16.3	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	1	0	1	1	0	1	10.7	-	30.9
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	1	1	0	1	1	0	10.7	16.3	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	5	4	1	5	4	1	53.4	65.4	30.9
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	1	0	1	47.6	-	99.0
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	4	2	2	5	2	3	23.8	18.3	29.7
Accidents (800-962)	4	2	2	8	4	4	85.5	65.4	123.4
Motor vehicle acc. (810-835)	3	2	1	3	2	1	32.0	32.7	30.9
Suicide (963, 970-979)	1	0	1	1	0	1	10.7	-	30.9
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
III-defined (780-795)	3	0	3	3	0	3	32.0	-	92.6

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
LEAKE COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	251	97	154	333	137	196	19.1	13.9	26.0
Fetal deaths	7	3	4	17	3	14	51.0	21.9	71.4
Total deaths	151	99	52	181	116	65	10.4	11.8	8.6
Deaths under 1 year	9	3	6	14	6	8	42.0	43.8	40.8
Deaths under 28 days	4	2	2	8	5	3	24.0	36.5	15.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	20	16	4	20	14	6	115.0	142.1	79.6
Benign & unspec. neoplasms (210-239)	1	0	1	1	0	1	5.8	-	13.3
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	0	2	3	1	2	17.2	10.2	26.5
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	32	24	8	35	25	10	201.3	253.8	132.6
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	46	32	14	55	38	17	316.3	385.8	225.5
Other hypertensive disease (444-447)	3	1	2	3	1	2	17.2	10.2	26.5
General arteriosclerosis (450)	6	4	2	6	4	2	34.5	40.6	26.5
Other circulatory diseases (451-468)	2	1	1	2	1	1	11.5	10.2	13.3
#Influenza & pneumonia (480-493)	6	2	4	8	4	4	46.0	40.6	53.0
Bronchitis (500-502)	2	2	0	2	2	0	11.5	20.3	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	1	0	1	2	0	2	11.5	-	26.5
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	0	1	5.8	-	13.3
Cirrhosis of liver (581)	1	1	0	1	1	0	5.8	10.2	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	2	1	1	11.5	10.2	13.3
Infections of kidney (600)	0	0	0	2	1	1	11.5	10.2	13.3
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	1	1	0	1	1	0	30.0	73.0	-
Congenital malformations (750-759)	0	0	0	1	1	0	5.8	10.2	-
Diseases of early infancy (760-776)	3	2	1	6	4	2	18.0	29.2	10.2
Accidents (800-962)	10	7	3	13	9	4	74.8	91.4	53.0
Motor vehicle acc. (810-835)	7	5	2	7	4	3	40.2	40.6	39.8
Suicide (963, 970-979)	2	1	1	3	2	1	17.2	20.3	13.3
Homicide (964, 980-985)	2	1	1	2	1	1	11.5	10.2	13.3
Ill-defined (780-795)	8	2	6	9	3	6	51.8	30.4	79.6

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
 AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
 LEE COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,353	1,006	347	1,062	717	345	24.0	21.9	29.6
Fetal deaths	33	20	13	25	12	13	23.5	16.7	37.7
Total deaths	448	314	134	419	287	132	9.4	8.8	11.3
Deaths under 1 year	45	26	19	34	17	17	32.0	23.7	49.3
Deaths under 28 days	31	21	10	21	13	8	19.8	18.1	23.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	2	1	1	2	1	1	4.5	3.1	8.6
Syphilis (020-029)	1	0	1	1	0	1	2.2	-	8.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	1	1	0	2.2	3.1	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	75	58	17	65	49	16	146.6	150.0	137.1
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	1	0	1	1	0	2.2	3.1	-
Diabetes mellitus (260)	4	3	1	3	2	1	6.8	6.1	8.6
Anemias (290-293)	1	0	1	1	0	1	2.2	-	8.6
Intracranial lesions (330-334)	56	39	17	53	36	17	119.5	110.2	145.7
Nonmeningococcal meningitis (340)	1	1	0	0	0	0	-	-	-
Heart diseases (400-443)	166	120	46	159	115	44	358.6	352.0	377.0
Other hypertensive disease (444-447)	1	1	0	3	3	0	6.8	9.2	-
General arteriosclerosis (450)	8	6	2	5	3	2	11.3	9.2	17.1
Other circulatory diseases (451-468)	5	3	2	7	4	3	15.8	12.2	25.7
#Influenza & pneumonia (480-493)	17	9	8	17	9	8	38.3	27.5	68.6
Bronchitis (500-502)	0	0	0	1	1	0	2.2	3.1	-
Stomach & duodenal ulcers (540, 541)	1	0	1	1	0	1	2.2	-	8.6
#Gastritis, etc. (543, 571, 572)	1	1	0	2	1	1	4.5	3.1	8.6
Hernia & intestinal obstruction (560, 561, 570)	3	2	1	2	1	1	4.5	3.1	8.6
Cirrhosis of liver (581)	2	2	0	2	2	0	4.5	6.1	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	2	1	2	1	1	4.5	3.1	8.6
Infections of kidney (600)	2	2	0	1	1	0	2.2	3.1	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	2	0	2	2	0	2	18.8	-	58.0
Congenital malformations (750-759)	7	5	2	3	2	1	6.8	6.1	8.6
Diseases of early infancy (760-776)	28	18	10	21	12	9	19.8	16.7	26.1
Accidents (800-962)	31	25	6	31	25	6	69.9	76.5	51.4
Motor vehicle acc. (810-835)	16	16	0	17	17	0	38.3	52.0	-
Suicide (963, 970-979)	5	3	2	5	3	2	11.3	9.2	17.1
Homicide (964, 980-985)	5	2	3	3	0	3	6.8	-	25.7
Ill-defined (780-795)	11	5	6	12	6	6	27.1	18.4	51.4

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
 AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
 LEFLORE COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,461	421	1,040	1,422	374	1,048	29.6	21.5	34.1
Fetal deaths	135	55	80	129	47	82	90.7	125.7	78.2
Total deaths	638	230	408	575	166	409	12.0	9.5	13.3
Deaths under 1 year	105	17	88	95	12	83	66.8	32.1	79.2
Deaths under 28 days	58	13	45	54	10	44	38.0	26.7	42.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	3	1	2	6	2	4	12.5	11.5	13.0
Syphilis (020-029)	0	0	0	1	0	1	2.1	-	3.2
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	64	31	33	61	26	35	126.8	149.4	114.0
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	6	0	6	6	0	6	12.5	-	19.5
Anemias (290-293)	4	0	4	4	0	4	8.3	-	13.0
Intracranial lesions (330-334)	72	31	41	63	22	41	131.0	126.4	133.6
Nonmeningococcal meningitis (340)	5	1	4	4	0	4	8.3	-	13.0
Heart diseases (400-443)	209	103	106	164	65	99	341.0	373.6	322.5
Other hypertensive disease (444-447)	2	0	2	2	0	2	4.2	-	6.5
General arteriosclerosis (450)	13	7	6	7	1	6	14.6	5.7	19.5
Other circulatory diseases (451-468)	8	3	5	7	1	6	14.6	5.7	19.5
#Influenza & pneumonia (480-493)	24	5	19	22	3	19	45.7	17.2	61.9
Bronchitis (500-502)	1	0	1	1	0	1	2.1	-	3.2
Stomach & duodenal ulcers (540, 541)	2	1	1	1	0	1	2.1	-	3.2
#Gastritis, etc. (543, 571, 572)	13	2	11	11	1	10	22.9	5.7	32.6
Hernia & intestinal obstruction (560, 561, 570)	3	1	2	4	1	3	8.3	5.7	9.8
Cirrhosis of liver (581)	5	3	2	7	5	2	14.6	28.7	6.5
Acute nephritis (590, 591)	3	1	2	2	1	1	4.2	5.7	3.2
Chronic nephritis, etc. (592-594)	3	0	3	6	0	6	12.5	-	19.5
Infections of kidney (600)	3	1	2	3	1	2	6.2	5.7	6.5
Hyperplasia of prostate (610)	3	2	1	3	2	1	6.2	11.5	3.2
Maternal causes (640-689)	1	0	1	2	0	2	14.1	-	19.1
Congenital malformations (750-759)	5	1	4	4	0	4	8.3	-	13.0
Diseases of early infancy (760-776)	61	12	49	58	9	49	40.8	24.1	46.8
Accidents (800-962)	35	7	28	39	9	30	81.1	51.7	97.7
Motor vehicle acc. (810-835)	14	4	10	18	7	11	37.4	40.2	35.8
Suicide (963, 970-979)	6	4	2	6	4	2	12.5	23.0	6.5
Homicide (964, 980-985)	5	0	5	4	0	4	8.3	-	13.0
Ill-defined (780-795)	64	8	56	60	7	53	124.7	40.2	172.6

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
LINCOLN COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	600	357	243	596	323	273	22.8	17.7	34.4
Fetal deaths	20	9	11	28	9	19	47.0	27.9	69.6
Total deaths	233	157	76	265	169	96	10.1	9.3	12.1
Deaths under 1 year	21	5	16	24	5	19	40.3	15.5	69.6
Deaths under 28 days	18	5	13	20	5	15	33.6	15.5	54.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	2	1	1	4	1	3	15.3	5.5	37.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	1	1	0	3.8	5.5	-
Malignant neoplasms (140-205)	25	20	5	28	21	7	107.0	115.2	88.2
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	1	0	1	1	0	3.8	5.5	-
Diabetes mellitus (260)	4	1	3	4	1	3	15.3	5.5	37.8
Anemias (290-293)	1	0	1	1	0	1	3.8	-	12.6
Intracranial lesions (330-334)	31	23	8	31	21	10	118.4	115.2	125.9
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	108	86	22	111	85	26	424.1	466.3	327.4
Other hypertensive disease (444-447)	1	0	1	1	0	1	3.8	-	12.6
General arteriosclerosis (450)	2	0	2	3	1	2	11.5	5.5	25.2
Other circulatory diseases (451-468)	0	0	0	1	1	0	3.8	5.5	-
#Influenza & pneumonia (480-493)	5	4	1	4	3	1	15.3	16.4	12.6
Bronchitis (500-502)	0	0	0	1	1	0	3.8	5.5	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	1	0	1	3.8	-	12.6
Chronic nephritis, etc. (592-594)	3	1	2	2	1	1	7.6	5.5	12.6
Infections of kidney (600)	1	0	1	4	2	2	15.3	11.0	25.2
Hyperplasia of prostate (610)	0	0	0	1	0	1	3.8	-	12.6
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	2	1	1	5	3	2	19.1	16.4	25.2
Diseases of early infancy (760-776)	14	5	9	15	4	11	25.2	12.4	40.3
Accidents (800-962)	9	5	4	19	13	6	72.6	71.3	75.6
Motor vehicle acc. (810-835)	4	3	1	9	6	3	34.4	32.9	37.8
Suicide (963, 970-979)	1	1	0	1	1	0	3.8	5.5	-
Homicide (964, 980-985)	2	1	1	1	0	1	3.8	-	12.6
Ill-defined (780-795)	12	3	9	12	2	10	45.8	11.0	125.9

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
LOWNDES COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,526	872	654	1,476	821	655	28.1	24.5	34.5
Fetal deaths	58	27	31	56	27	29	37.9	32.9	44.3
Total deaths	465	230	235	482	243	239	9.2	7.2	12.6
Deaths under 1 year	48	19	29	47	19	28	31.8	23.1	42.7
Deaths under 28 days	33	16	17	33	16	17	22.4	19.5	26.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	6	1	5	6	1	5	11.4	3.0	26.3
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	1	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	2	0	2	2	0	2	3.8	-	10.5
Malignant neoplasms (140-205)	58	41	17	61	40	21	116.2	119.3	110.6
Benign & unspec. neoplasms (210-239)	2	1	1	2	1	1	3.8	3.0	5.3
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	3	1	2	2	0	2	3.8	-	10.5
Anemias (290-293)	1	0	1	2	1	1	3.8	3.0	5.3
Intracranial lesions (330-334)	53	23	30	55	25	30	104.8	74.6	158.1
Nonmeningococcal meningitis (340)	1	0	1	1	0	1	1.9	-	5.3
Heart diseases (400-443)	147	77	70	146	79	67	278.1	235.7	353.0
Other hypertensive disease (444-447)	2	1	1	3	1	2	5.7	3.0	10.5
General arteriosclerosis (450)	17	14	3	18	15	3	34.3	44.7	15.8
Other circulatory diseases (451-468)	1	0	1	2	1	1	3.8	3.0	5.3
#Influenza & pneumonia (480-493)	22	5	17	21	5	16	40.0	14.9	84.3
Bronchitis (500-502)	1	1	0	1	1	0	1.9	3.0	-
Stomach & duodenal ulcers (540, 541)	1	1	0	1	1	0	1.9	3.0	-
#Gastritis, etc. (543, 571, 572)	1	0	1	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	3	2	1	4	2	2	7.6	6.0	10.5
Cirrhosis of liver (581)	2	2	0	2	2	0	3.8	6.0	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	2	2	5	3	2	9.5	8.9	10.5
Infections of kidney (600)	3	1	2	3	1	2	5.7	3.0	10.5
Hyperplasia of prostate (610)	1	0	1	1	0	1	1.9	-	5.3
Maternal causes (640-689)	3	0	3	2	0	2	13.6	-	30.5
Congenital malformations (750-759)	5	5	0	5	5	0	9.5	14.9	-
Diseases of early infancy (760-776)	26	13	13	26	13	13	17.6	15.8	19.8
Accidents (800-962)	31	18	13	41	26	15	78.1	77.6	79.0
Motor vehicle acc. (810-835)	15	12	3	15	13	2	28.6	38.8	10.5
Suicide (963, 970-979)	6	4	2	7	5	2	13.3	14.9	10.5
Homicide (964, 980-985)	3	2	1	2	1	1	3.8	3.0	5.3
Ill-defined (780-795)	36	2	34	37	2	35	70.5	6.0	184.4

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
MADISON COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	796	132	664	903	177	726	27.1	16.9	31.8
Fetal deaths	28	4	24	37	8	29	41.0	45.2	39.9
Total deaths	319	86	233	372	107	265	11.2	10.2	11.6
Deaths under 1 year	39	1	38	45	3	42	49.8	16.9	57.8
Deaths under 28 days	29	1	28	33	2	31	36.5	11.3	42.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	0	1	1	0	1	3.0	-	4.4
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	3.0	-	4.4
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	27	9	18	39	14	25	117.1	133.6	109.5
Benign & unspec. neoplasms (210-239)	0	0	0	1	0	1	3.0	-	4.4
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	5	1	4	5	1	4	15.0	9.5	17.5
Anemias (290-293)	1	0	1	3	0	3	9.0	-	13.1
Intracranial lesions (330-334)	48	6	42	56	11	45	168.1	105.0	197.1
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	83	37	46	95	43	52	285.2	410.3	227.8
Other hypertensive disease (444-447)	5	1	4	4	1	3	12.0	9.5	13.1
General arteriosclerosis (450)	7	2	5	8	3	5	24.0	28.6	21.9
Other circulatory diseases (451-468)	1	0	1	1	0	1	3.0	-	4.4
#Influenza & pneumonia (480-493)	8	2	6	8	2	6	24.0	19.1	26.3
Bronchitis (500-502)	1	0	1	1	0	1	3.0	-	4.4
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	4	0	4	4	0	4	12.0	-	17.5
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	3	2	1	3	2	1	9.0	19.1	4.4
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	6.0	9.5	4.4
Infections of kidney (600)	1	0	1	2	0	2	6.0	-	8.8
Hyperplasia of prostate (610)	1	0	1	1	0	1	3.0	-	4.4
Maternal causes (640-689)	1	0	1	1	0	1	11.1	-	13.8
Congenital malformations (750-759)	3	0	3	8	2	6	24.0	19.1	26.3
Diseases of early infancy (760-776)	21	1	20	26	2	24	28.8	11.3	33.0
Accidents (800-962)	31	14	17	34	13	21	102.1	124.0	92.0
Motor vehicle acc. (810-835)	14	9	5	15	9	6	45.0	85.9	26.3
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	4	0	4	7	1	6	21.0	9.5	26.3
Ill-defined (780-795)	51	6	45	47	4	43	141.1	38.2	188.3

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
MARION COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	444	237	207	525	291	234	24.5	21.0	30.8
Fetal deaths	15	6	9	16	7	9	30.5	24.0	38.5
Total deaths	193	128	65	207	132	75	9.6	9.5	9.9
Deaths under 1 year	14	4	10	17	5	12	32.4	17.2	51.3
Deaths under 28 days	9	3	6	10	3	7	19.0	10.3	29.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	4	2	2	5	2	3	23.3	14.4	39.5
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	30	22	8	32	24	8	149.2	173.4	105.3
Benign & unspec. neoplasms (210-239)	1	1	0	1	1	0	4.7	7.2	-
Asthma (241)	0	0	0	1	1	0	4.7	7.2	-
Diabetes mellitus (260)	1	0	1	3	0	3	14.0	-	39.5
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	33	24	9	30	21	9	139.9	151.7	118.4
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	50	37	13	51	37	14	237.9	267.3	184.2
Other hypertensive disease (444-447)	2	0	2	2	0	2	9.3	-	26.3
General arteriosclerosis (450)	7	6	1	7	6	1	32.6	43.4	13.2
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	7	1	6	9	2	7	42.0	14.4	92.1
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	1	1	0	1	1	0	4.7	7.2	-
#Gastritis, etc. (543, 571, 572)	0	0	0	1	0	1	4.7	-	13.2
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	1	0	1	1	0	4.7	7.2	-
Chronic nephritis, etc. (592-594)	1	1	0	2	2	0	9.3	14.4	-
Infections of kidney (600)	0	0	0	1	1	0	4.7	7.2	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	1	1	0	4.7	7.2	-
Diseases of early infancy (760-776)	8	2	6	8	2	6	15.2	6.9	25.6
Accidents (800-962)	12	9	3	13	10	3	60.6	72.2	39.5
Motor vehicle acc. (810-835)	10	7	3	9	6	3	42.0	43.4	39.5
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	1	0	1	1	0	1	4.7	-	13.2
Ill-defined (780-795)	25	13	12	26	12	14	121.3	86.7	184.2

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
MARSHALL COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	642	95	547	743	165	578	29.6	21.8	32.9
Fetal deaths	43	1	42	38	2	36	51.1	12.1	62.3
Total deaths	238	76	162	254	68	186	10.1	9.0	10.6
Deaths under 1 year	37	3	34	41	4	37	55.2	24.2	64.0
Deaths under 28 days	24	2	22	25	3	22	33.6	18.2	38.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	5	0	5	6	0	6	23.9	-	34.1
Syphilis (020-029)	1	1	0	1	1	0	4.0	13.2	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	1	0	1	1	0	1	4.0	-	5.7
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	14	5	9	19	5	14	75.6	66.2	79.7
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	5	2	3	3	1	2	11.9	13.2	11.4
Anemias (290-293)	1	0	1	1	0	1	4.0	-	5.7
Intracranial lesions (330-334)	44	15	29	40	9	31	159.2	119.2	176.4
Nonmeningococcal meningitis (340)	0	0	0	1	0	1	4.0	-	5.7
Heart diseases (400-443)	70	28	42	73	28	45	290.6	370.9	256.1
Other hypertensive disease (444-447)	1	1	0	0	0	0	-	-	-
General arteriosclerosis (450)	2	0	2	3	1	2	11.9	13.2	11.4
Other circulatory diseases (451-468)	1	0	1	4	0	4	15.9	-	22.8
#Influenza & pneumonia (480-493)	9	1	8	13	2	11	51.8	26.5	62.6
Bronchitis (500-502)	0	0	0	1	1	0	4.0	13.2	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	2	0	2	3	0	3	11.9	-	17.1
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	1	0	4.0	13.2	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	0	1	1	0	1	4.0	-	5.7
Chronic nephritis, etc. (592-594)	4	3	1	7	2	5	27.9	26.5	28.4
Infections of kidney (600)	1	1	0	1	1	0	4.0	13.2	-
Hyperplasia of prostate (610)	1	0	1	1	0	1	4.0	-	5.7
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	0	1	2	0	2	8.0	-	11.4
Diseases of early infancy (760-776)	19	1	18	19	2	17	25.6	12.1	29.4
Accidents (800-962)	18	8	10	13	6	7	51.8	79.5	39.8
Motor vehicle acc. (810-835)	13	6	7	7	3	4	27.9	39.7	22.8
Suicide (963, 970-979)	5	4	1	2	1	1	8.0	13.2	5.7
Homicide (964, 980-985)	2	0	2	3	0	3	11.9	-	17.1
Ill-defined (780-795)	23	3	20	25	4	21	99.5	53.0	119.5

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
MONROE COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	807	466	341	799	449	350	21.9	17.8	31.1
Fetal deaths	28	11	17	32	14	18	40.0	31.2	51.4
Total deaths	352	229	123	366	238	128	10.0	9.4	11.4
Deaths under 1 year	37	8	29	40	8	32	50.1	17.8	91.4
Deaths under 28 days	22	6	16	22	5	17	27.5	11.1	48.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	0	1	1	0	1	2.7	-	8.9
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	1	1	0	1	1	0	2.7	4.0	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	44	35	9	48	39	9	131.5	154.4	80.1
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	7	2	5	8	2	6	21.9	7.9	53.4
Anemias (290-293)	1	0	1	1	0	1	2.7	-	8.9
Intracranial lesions (330-334)	54	39	15	53	37	16	145.2	146.5	142.3
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	117	87	30	120	91	29	328.8	360.2	258.0
Other hypertensive disease (444-447)	1	0	1	1	0	1	2.7	-	8.9
General arteriosclerosis (450)	6	4	2	8	6	2	21.9	23.8	17.8
Other circulatory diseases (451-468)	4	3	1	5	3	2	13.7	11.9	17.8
#Influenza & pneumonia (480-493)	17	8	9	21	10	11	57.5	39.6	97.9
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	3	1	2	3	1	2	8.2	4.0	17.8
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	1	0	2.7	4.0	-
Cirrhosis of liver (581)	1	0	1	0	0	0	-	-	-
Acute nephritis (590, 591)	2	0	2	2	0	2	5.5	-	17.8
Chronic nephritis, etc. (592-594)	10	2	8	9	1	8	24.6	4.0	71.2
Infections of kidney (600)	3	2	1	3	2	1	8.2	7.9	8.9
Hyperplasia of prostate (610)	0	0	0	1	1	0	2.7	4.0	-
Maternal causes (640-689)	1	1	0	1	1	0	12.5	22.3	-
Congenital malformations (750-759)	3	1	2	5	3	2	13.7	11.9	17.8
Diseases of early infancy (760-776)	20	6	14	20	5	15	25.0	11.1	42.8
Accidents (800-962)	26	22	4	20	16	4	54.8	63.3	35.6
Motor vehicle acc. (810-835)	12	9	3	11	9	2	30.1	35.6	17.8
Suicide (963, 970-979)	3	3	0	3	3	0	8.2	11.9	-
Homicide (964, 980-985)	4	2	2	6	4	2	16.4	15.8	17.8
Ill-defined (780-795)	12	3	9	13	3	10	35.6	11.9	89.0

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
MONTGOMERY COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	397	175	222	340	131	209	25.5	16.2	39.6
Fetal deaths	21	6	15	21	7	14	61.8	53.4	67.0
Total deaths	126	83	43	141	95	46	10.6	11.8	8.7
Deaths under 1 year	12	3	9	13	3	10	38.2	22.9	47.8
Deaths under 28 days	6	2	4	7	3	4	20.6	22.9	19.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	1	0	1	1	0	1	7.5	-	19.0
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	20	18	2	21	17	4	157.3	210.4	75.9
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	1	0	1	1	0	1	7.5	-	19.0
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	10	5	18	14	4	134.8	173.3	75.9
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	53	38	15	54	41	13	404.5	507.4	246.7
Other hypertensive disease (444-447)	3	1	2	1	0	1	7.5	-	19.0
General arteriosclerosis (450)	1	1	0	1	1	0	7.5	12.4	-
Other circulatory diseases (451-468)	2	2	0	3	2	1	22.5	24.8	19.0
#Influenza & pneumonia (480-493)	4	2	2	7	4	3	52.4	49.5	56.9
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	1	1	0	7.5	12.4	-
#Gastritis, etc. (543, 571, 572)	1	0	1	2	1	1	15.0	12.4	19.0
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	1	1	0	1	1	0	7.5	12.4	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	7.5	12.4	-
Infections of kidney (600)	0	0	0	1	0	1	7.5	-	19.0
Hyperplasia of prostate (610)	0	0	0	1	1	0	7.5	12.4	-
Maternal causes (640-689)	1	0	1	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	1	0	1	7.5	-	19.0
Diseases of early infancy (760-776)	5	2	3	6	3	3	17.6	22.9	14.4
Accidents (800-962)	5	3	2	7	3	4	52.4	37.1	75.9
Motor vehicle acc. (810-835)	4	3	1	3	2	1	22.5	24.8	19.0
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	0	0	0	1	0	1	7.5	-	19.0
Ill-defined (780-795)	7	2	5	6	2	4	44.9	24.8	75.9

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
NESHOBIA COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	394	205	189	401	215	186	18.7	13.6	32.9
Fetal deaths	12	1	11	18	5	13	44.9	23.2	69.9
Total deaths	192	141	51	224	167	57	10.4	10.6	10.1
Deaths under 1 year	17	7	10	24	8	16	59.8	37.2	86.0
Deaths under 28 days	6	3	3	12	5	7	29.9	23.2	37.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	0	1	4.6	-	17.7
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	29	21	8	31	23	8	144.4	145.6	141.3
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	0	2	1	0	1	4.6	-	17.7
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	39	31	8	51	44	7	237.6	278.5	123.7
Nonmeningococcal meningitis (340)	0	0	0	1	0	1	4.6	-	17.7
Heart diseases (400-443)	51	41	10	52	42	10	242.3	265.8	176.7
Other hypertensive disease (444-447)	2	1	1	2	1	1	9.3	6.3	17.7
General arteriosclerosis (450)	3	3	0	3	3	0	14.0	19.0	-
Other circulatory diseases (451-468)	3	3	0	4	4	0	18.6	25.3	-
#Influenza & pneumonia (480-493)	3	2	1	8	6	2	37.3	38.0	35.3
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	1	1	0	2	2	0	9.3	12.6	-
#Gastritis, etc. (543, 571, 572)	3	1	2	3	1	2	14.0	6.3	35.3
Hernia & intestinal obstruction (560, 561, 570)	1	0	1	1	0	1	4.6	-	17.7
Cirrhosis of liver (581)	1	1	0	1	1	0	4.6	6.3	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	-	-	-
Infections of kidney (600)	1	0	1	0	1	0	4.6	6.3	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	3	2	1	4	3	1	18.6	19.0	17.7
Diseases of early infancy (760-776)	5	3	2	10	3	7	24.9	14.0	37.6
Accidents (800-962)	17	14	3	20	16	4	93.2	101.3	70.7
Motor vehicle acc. (810-835)	6	4	2	7	5	2	32.6	31.6	35.3
Suicide (963, 970-979)	2	1	1	1	0	1	4.6	-	17.7
Homicide (964, 980-985)	4	2	2	3	1	2	14.0	6.3	35.3
Ill-defined (780-795)	9	4	5	11	6	5	51.2	38.0	88.3

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
NEWTON COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	370	206	164	378	185	193	18.3	13.5	27.9
Fetal deaths	33	18	15	31	14	17	82.0	75.7	88.1
Total deaths	198	109	89	248	149	99	12.0	10.9	14.3
Deaths under 1 year	14	5	9	20	8	12	52.9	43.2	62.2
Deaths under 28 days	8	4	4	11	6	5	29.1	32.4	25.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	1	0	1	4.8	-	14.4
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	18	9	9	26	17	9	126.1	124.1	130.0
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	3	1	2	4	1	3	19.4	7.3	43.4
Anemias (290-293)	1	0	1	1	0	1	4.8	-	14.4
Intracranial lesions (330-334)	42	22	20	41	23	18	198.8	167.9	260.1
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	69	49	20	85	62	23	412.2	452.6	332.4
Other hypertensive disease (444-447)	3	2	1	2	1	1	9.7	7.3	14.4
General arteriosclerosis (450)	2	1	1	4	3	1	19.4	21.9	14.4
Other circulatory diseases (451-468)	2	1	1	2	1	1	9.7	7.3	14.4
#Influenza & pneumonia (480-493)	4	0	4	5	1	4	24.2	7.3	57.8
Bronchitis (500-502)	1	0	1	1	0	1	4.8	-	14.4
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	2	0	2	4	1	3	19.4	7.3	43.4
Hernia & intestinal obstruction (560, 561, 570)	2	1	1	2	1	1	9.7	7.3	14.4
Cirrhosis of liver (581)	1	1	0	1	1	0	4.8	7.3	-
Acute nephritis (590, 591)	1	1	0	1	1	0	4.8	7.3	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	2	2	0	6	5	1	29.1	36.5	14.4
Diseases of early infancy (760-776)	7	4	3	7	5	2	18.5	27.0	10.4
Accidents (800-962)	14	6	8	19	11	8	92.1	80.3	115.6
Motor vehicle acc. (810-835)	7	3	4	12	8	4	58.2	58.4	57.8
Suicide (963, 970-979)	2	2	0	4	4	0	19.4	29.2	-
Homicide (964, 980-985)	3	1	2	4	1	3	19.4	7.3	43.4
Ill-defined (780-795)	11	2	9	15	3	12	72.7	21.9	173.4

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
NOXUBEE COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	449	82	367	438	74	364	25.0	14.4	29.3
Fetal deaths	16	2	14	15	2	13	34.2	27.0	35.7
Total deaths	214	61	153	230	73	157	13.1	14.2	12.6
Deaths under 1 year	24	0	24	24	0	24	54.8	-	65.9
Deaths under 28 days	11	0	11	10	0	10	22.8	-	27.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	22	11	11	24	13	11	136.8	253.4	88.6
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	1	0	1	1	0	5.7	19.5	-
Diabetes mellitus (260)	4	0	4	4	0	4	22.8	-	32.2
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	32	10	22	34	12	22	193.8	233.9	177.3
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	57	24	33	60	28	32	342.1	545.8	257.8
Other hypertensive disease (444-447)	0	0	0	2	1	1	11.4	19.5	8.0
General arteriosclerosis (450)	7	4	3	6	4	2	34.2	78.0	16.1
Other circulatory diseases (451-468)	2	0	2	3	1	2	17.1	19.5	16.1
#Influenza & pneumonia (480-493)	12	1	11	14	1	13	79.8	19.5	104.8
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	2	0	2	1	0	1	5.7	-	8.0
#Gastritis, etc. (543, 571, 572)	2	1	1	3	1	2	17.1	19.5	16.1
Hernia & intestinal obstruction (560, 561, 570)	3	1	2	3	1	2	17.1	19.5	16.1
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	0	3	3	0	3	17.1	-	24.2
Infections of kidney (600)	1	0	1	1	0	1	5.7	-	8.0
Hyperplasia of prostate (610)	1	0	1	1	0	1	5.7	-	8.0
Maternal causes (640-689)	1	0	1	2	0	2	45.7	-	54.9
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	11	0	11	11	0	11	25.1	-	30.2
Accidents (800-962)	14	3	11	13	1	12	74.1	19.5	96.7
Motor vehicle acc. (810-835)	3	1	2	3	0	3	17.1	-	24.2
Suicide (963, 970-979)	0	0	0	1	0	1	5.7	-	8.0
Homicide (964, 980-985)	2	0	2	2	0	2	11.4	-	16.1
<u>Ill-defined (780-795)</u>	30	2	28	31	3	28	176.7	58.5	225.6

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
OKTIBBEHA COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	803	344	459	741	309	432	26.2	19.7	34.5
Fetal deaths	19	5	14	23	7	16	31.0	22.6	37.0
Total deaths	215	95	120	232	103	129	8.2	6.6	10.3
Deaths under 1 year	25	7	18	21	3	18	28.3	9.7	41.7
Deaths under 28 days	17	7	10	13	3	10	17.5	9.7	23.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	1	0	1	1	0	1	3.5	-	8.0
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	22	9	13	24	11	13	85.0	70.1	103.8
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	0	1	1	0	1	3.5	-	8.0
Diabetes mellitus (260)	2	2	0	1	1	0	3.5	6.4	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	39	19	20	38	18	20	134.6	114.6	159.7
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	73	36	37	89	46	43	315.4	293.0	343.4
Other hypertensive disease (444-447)	1	0	1	3	0	3	10.6	-	24.0
General arteriosclerosis (450)	6	4	2	6	4	2	21.3	25.5	16.0
Other circulatory diseases (451-468)	2	1	1	1	0	1	3.5	-	8.0
#Influenza & pneumonia (480-493)	11	4	7	11	4	7	39.0	25.5	55.9
Bronchitis (500-502)	1	0	1	1	0	1	3.5	-	8.0
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	5	0	5	6	1	5	21.3	6.4	39.9
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	1	0	1	1	0	1	3.5	-	8.0
Acute nephritis (590, 591)	1	1	0	1	1	0	3.5	6.4	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	3.5	-	8.0
Infections of kidney (600)	2	1	1	3	2	1	10.6	12.7	8.0
Hyperplasia of prostate (610)	1	0	1	1	0	1	3.5	-	8.0
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	3	1	2	3	1	2	10.6	6.4	16.0
Diseases of early infancy (760-776)	14	6	8	10	2	8	13.5	6.5	18.5
Accidents (800-962)	7	6	1	8	6	2	28.3	38.2	16.0
Motor vehicle acc. (810-835)	1	0	1	1	0	1	3.5	-	8.0
Suicide (963, 970-979)	1	1	0	1	1	0	3.5	6.4	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	12	2	10	12	2	10	42.5	12.7	79.9

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
PANOLA COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	882	282	600	879	288	591	28.2	20.4	34.6
Fetal deaths	22	4	18	30	6	24	34.1	20.8	40.6
Total deaths	252	107	145	320	128	192	10.2	9.0	11.2
Deaths under 1 year	27	5	22	37	6	31	42.1	20.8	52.4
Deaths under 28 days	17	3	14	24	4	20	27.3	13.9	33.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	0	1	5	2	3	16.0	14.2	17.6
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	1	0	2	1	1	6.4	7.1	5.8
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	2	0	2	2	0	2	6.4	-	11.7
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	30	16	14	43	23	20	137.7	162.8	117.0
Benign & unspec. neoplasms (210-239)	1	1	0	1	1	0	3.2	7.1	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	5	1	4	5	0	5	16.0	-	29.2
Anemias (290-293)	2	2	0	2	1	1	6.4	7.1	5.8
Intracranial lesions (330-334)	34	15	19	41	20	21	131.3	141.5	122.9
Nonmeningococcal meningitis (340)	1	0	1	2	0	2	6.4	-	11.7
Heart diseases (400-443)	87	43	44	97	46	51	310.7	325.5	298.4
Other hypertensive disease (444-447)	3	1	2	5	1	4	16.0	7.1	23.4
General arteriosclerosis (450)	4	3	1	4	3	1	12.8	21.2	5.8
Other circulatory diseases (451-468)	3	0	3	4	0	4	12.8	-	23.4
#Influenza & pneumonia (480-493)	11	2	9	16	4	12	51.2	28.3	70.2
Bronchitis (500-502)	0	0	0	1	1	0	3.2	7.1	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	2	0	2	4	0	4	12.8	-	23.4
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	3	0	3	9.6	-	17.6
Cirrhosis of liver (581)	0	0	0	1	1	0	3.2	7.1	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	2	0	2	6.4	-	11.7
Infections of kidney (600)	1	0	1	2	1	1	6.4	7.1	5.8
Hyperplasia of prostate (610)	1	1	0	1	1	0	3.2	7.1	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	3	0	3	7	3	4	22.4	21.2	23.4
Diseases of early infancy (760-776)	12	3	9	16	2	14	18.2	6.9	23.7
Accidents (800-962)	17	9	8	15	5	10	48.0	35.4	58.5
Motor vehicle acc. (810-835)	5	4	1	6	3	3	19.2	21.2	17.6
Suicide (963, 970-979)	3	1	2	3	1	2	9.6	7.1	11.7
Homicide (964, 980-985)	4	1	3	8	2	6	25.6	14.2	35.1
Ill-defined (780-795)	13	3	10	17	4	13	54.4	28.3	76.1

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
PEARL RIVER COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	647	500	147	578	426	152	22.3	20.8	28.0
Fetal deaths	8	5	3	9	6	3	15.6	14.1	19.7
Total deaths	219	162	57	243	176	67	9.4	8.6	12.3
Deaths under 1 year	21	12	9	21	11	10	36.3	25.8	65.8
Deaths under 28 days	16	9	7	17	9	8	29.4	21.1	52.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	2	2	0	0	0	0	-	-	-
Syphilis (020-029)	1	0	1	1	0	1	3.8	-	18.4
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	1	1	0	1	1	0	3.8	4.9	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	26	22	4	34	25	9	131.2	122.1	165.7
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	3	3	0	4	4	0	15.4	19.5	-
Diabetes mellitus (260)	3	2	1	3	2	1	11.6	9.8	18.4
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	9	6	22	15	7	84.9	73.2	128.9
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	81	60	21	84	61	23	324.2	297.8	423.6
Other hypertensive disease (444-447)	1	0	1	1	0	1	3.8	-	18.4
General arteriosclerosis (450)	10	10	0	10	10	0	38.6	48.8	-
Other circulatory diseases (451-468)	2	1	1	2	1	1	7.7	4.9	18.4
#Influenza & pneumonia (480-493)	12	11	1	16	14	2	61.8	68.4	36.8
Bronchitis (500-502)	1	1	0	1	1	0	3.8	4.9	-
Stomach & duodenal ulcers (540, 541)	1	1	0	1	1	0	3.8	4.9	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	2	1	1	2	1	1	7.7	4.9	18.4
Acute nephritis (590, 591)	0	0	0	1	0	1	3.8	-	18.4
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	3.8	4.9	-
Infections of kidney (600)	2	2	0	3	3	0	11.6	14.6	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	2	2	0	2	2	0	7.7	9.8	-
Diseases of early infancy (760-776)	13	8	5	14	8	6	24.2	18.8	39.5
Accidents (800-962)	21	15	6	19	14	5	73.3	68.4	92.1
Motor vehicle acc. (810-835)	13	10	3	11	9	2	42.4	43.9	36.8
Suicide (963, 970-979)	1	1	0	1	1	0	3.8	4.9	-
Homicide (964, 980-985)	2	0	2	2	0	2	7.7	-	36.8
Ill-defined (780-795)	12	7	5	11	6	5	42.4	29.3	92.1

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
PERRY COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	25	20	5	198	143	55	23.2	24.6	20.3
Fetal deaths	1	0	1	3	1	2	15.2	7.0	36.4
Total deaths	40	25	15	85	52	33	10.0	8.9	12.2
Deaths under 1 year	2	0	2	5	0	5	25.2	-	90.9
Deaths under 28 days	0	0	0	3	0	3	15.2	-	54.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	4	3	1	8	7	1	93.8	120.3	36.9
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	0	0	0	0	0	0	-	-	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	7	4	3	16	11	5	187.6	189.0	184.5
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	14	11	3	24	19	5	281.4	326.5	184.5
Other hypertensive disease (444-447)	0	0	0	1	0	1	11.7	-	36.9
General arteriosclerosis (450)	1	1	0	1	1	0	11.7	17.2	-
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	1	0	1	4	1	3	46.9	17.2	110.7
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	1	1	0	1	1	0	11.7	17.2	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	1	1	0	11.7	17.2	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	2	1	1	23.4	17.2	36.9
Hyperplasia of prostate (610)	1	0	1	1	0	1	11.7	-	36.9
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	0	0	0	3	0	3	15.2	-	54.5
Accidents (800-962)	3	1	2	12	5	7	140.7	85.9	258.3
Motor vehicle acc. (810-835)	2	1	1	6	2	4	70.3	34.4	147.6
Suicide (963, 970-979)	1	1	0	1	1	0	11.7	17.2	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	6	3	3	6	3	3	70.3	51.5	110.7

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
PIKE COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	814	393	421	766	331	435	22.0	16.6	29.0
Fetal deaths	25	8	17	25	8	17	32.6	24.2	39.1
Total deaths	390	210	180	407	222	185	11.7	11.2	12.3
Deaths under 1 year	40	9	31	34	7	27	44.4	21.1	62.1
Deaths under 28 days	27	8	19	21	6	15	27.4	18.1	34.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	0	1	1	0	1	2.9	-	6.7
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	1	0	1	1	0	2.9	5.0	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	39	20	19	50	25	25	143.4	125.8	166.7
Benign & unspec. neoplasms (210-239)	1	1	0	1	1	0	2.9	5.0	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	7	3	4	8	3	5	22.9	15.1	33.3
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	72	44	28	68	44	24	195.0	221.4	160.0
Nonmeningococcal meningitis (340)	1	1	0	1	1	0	2.9	5.0	-
Heart diseases (400-443)	128	81	47	134	85	49	384.3	427.8	326.7
Other hypertensive disease (444-447)	6	4	2	6	4	2	17.2	20.1	13.3
General arteriosclerosis (450)	6	3	3	4	1	3	11.5	5.0	20.0
Other circulatory diseases (451-468)	4	3	1	5	4	1	14.3	20.1	6.7
#Influenza & pneumonia (480-493)	19	7	12	17	6	11	48.8	30.2	73.3
Bronchitis (500-502)	1	0	1	1	0	1	2.9	-	6.7
Stomach & duodenal ulcers (540, 541)	1	0	1	3	1	2	8.6	5.0	13.3
#Gastritis, etc. (543, 571, 572)	5	0	5	4	0	4	11.5	-	26.7
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	1	0	2.9	5.0	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	1	0	1	1	0	2.9	5.0	-
Chronic nephritis, etc. (592-594)	2	2	0	2	2	0	5.7	10.1	-
Infections of kidney (600)	1	0	1	2	0	2	5.7	-	13.3
Hyperplasia of prostate (610)	1	0	1	1	0	1	2.9	-	6.7
Maternal causes (640-689)	1	0	1	1	0	1	13.0	-	23.0
Congenital malformations (750-759)	2	2	0	2	2	0	5.7	10.1	-
Diseases of early infancy (760-776)	24	7	17	18	5	13	23.5	15.1	29.9
Accidents (800-962)	18	8	10	24	14	10	68.8	70.4	66.7
Motor vehicle acc. (810-835)	6	1	5	7	2	5	20.1	10.1	33.3
Suicide (963, 970-979)	2	2	0	2	2	0	5.7	10.1	-
Homicide (964, 980-985)	2	1	1	3	0	3	8.6	-	20.0
Ill-defined (780-795)	29	10	19	28	9	19	80.3	45.3	126.7

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
PONTOTOC COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	361	267	94	327	246	81	17.4	16.0	23.4
Fetal deaths	3	0	3	7	4	3	21.4	16.3	37.0
Total deaths	193	156	37	209	173	36	11.1	11.3	10.4
Deaths under 1 year	8	5	3	9	6	3	27.5	24.4	37.0
Deaths under 28 days	6	4	2	7	5	2	21.4	20.3	24.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	2	2	0	10.6	13.0	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	19	15	4	25	22	3	132.8	143.2	86.7
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	3	1	2	4	2	2	21.2	13.0	57.8
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	23	17	6	26	21	5	138.2	136.7	144.5
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	92	81	11	93	82	11	494.2	533.8	317.9
Other hypertensive disease (444-447)	3	2	1	3	2	1	15.9	13.0	28.9
General arteriosclerosis (450)	6	5	1	5	4	1	26.6	26.0	28.9
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	7	5	2	7	5	2	37.2	32.6	57.8
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	1	1	0	1	1	0	5.3	6.5	-
#Gastritis, etc. (543, 571, 572)	2	2	0	1	1	0	5.3	6.5	-
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	1	1	0	5.3	6.5	-
Cirrhosis of liver (581)	1	1	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	1	1	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	1	0	3	3	0	15.9	19.5	-
Diseases of early infancy (760-776)	6	4	2	6	4	2	18.3	16.3	24.7
Accidents (800-962)	13	11	2	14	11	3	74.4	71.6	86.7
Motor vehicle acc. (810-835)	12	10	2	12	9	3	63.8	58.6	86.7
Suicide (963, 970-979)	1	1	0	1	1	0	5.3	6.5	-
Homicide (964, 980-985)	6	2	4	6	2	4	31.9	13.0	115.6
Ill-defined (780-795)	1	0	1	1	0	1	5.3	-	28.9

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
PRENTISS COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	424	348	76	382	313	69	19.8	18.6	28.2
Fetal deaths	13	9	4	12	9	3	31.4	28.8	43.5
Total deaths	195	167	28	195	165	30	10.1	9.8	12.2
Deaths under 1 year	7	4	3	7	4	3	18.3	12.8	43.5
Deaths under 28 days	4	3	1	4	3	1	10.5	9.6	14.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	21	18	3	25	21	4	129.7	124.8	163.3
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	3	3	0	2	2	0	10.4	11.9	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	33	27	6	32	26	6	166.0	154.5	244.9
Nonmeningococcal meningitis (340)	1	1	0	1	1	0	5.2	5.9	-
Heart diseases (400-443)	89	77	12	81	68	13	420.1	404.0	530.6
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	11	10	1	10	9	1	51.9	53.5	40.8
Other circulatory diseases (451-468)	1	1	0	2	2	0	10.4	11.9	-
#Influenza & pneumonia (480-493)	6	5	1	7	6	1	36.3	35.6	40.8
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	1	1	0	1	1	0	5.2	5.9	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	1	1	0	5.2	5.9	-
Cirrhosis of liver (581)	1	1	0	1	1	0	5.2	5.9	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	1	1	0	1	1	0	5.2	5.9	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	4	3	1	4	3	1	10.5	9.6	14.5
Accidents (800-962)	9	8	1	13	12	1	67.4	71.3	40.8
Motor vehicle acc. (810-835)	4	4	0	5	5	0	25.9	29.7	-
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	8	6	2	8	6	2	41.5	35.6	81.6

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
QUITMAN COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	790	215	575	687	174	513	29.8	18.9	37.0
Fetal deaths	19	1	18	24	2	22	34.9	11.5	42.9
Total deaths	208	49	159	258	83	175	11.2	9.0	12.6
Deaths under 1 year	40	7	33	37	6	31	53.8	34.5	60.4
Deaths under 28 days	18	4	14	17	3	14	24.7	17.2	27.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	2	0	2	2	0	2	8.7	-	14.4
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	24	7	17	31	12	19	134.4	130.4	137.0
Benign & unspec. neoplasms (210-239)	1	0	1	2	1	1	8.7	10.9	7.2
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	3	0	3	5	0	5	21.7	-	36.0
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	16	4	12	27	11	16	117.0	119.6	115.4
Nonmeningococcal meningitis (340)	1	0	1	0	0	0	-	-	-
Heart diseases (400-443)	67	24	43	84	36	48	364.1	391.3	346.1
Other hypertensive disease (444-447)	3	1	2	3	1	2	13.0	10.9	14.4
General arteriosclerosis (450)	0	0	0	1	1	0	4.3	10.9	-
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	12	0	12	13	1	12	56.4	10.9	86.5
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	1	0	1	4.3	-	7.2
#Gastritis, etc. (543, 571, 572)	4	1	3	3	1	2	13.0	10.9	14.4
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	1	1	0	4.3	10.9	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	1	0	1	1	0	1	14.6	-	19.5
Congenital malformations (750-759)	5	3	2	5	3	2	21.7	32.6	14.4
Diseases of early infancy (760-776)	14	2	12	12	0	12	17.5	-	23.4
Accidents (800-962)	18	5	13	26	8	18	112.7	87.0	129.8
Motor vehicle acc. (810-835)	4	2	2	7	4	3	30.3	43.5	21.6
Suicide (963, 970-979)	0	0	0	1	1	0	4.3	10.9	-
Homicide (964, 980-985)	4	1	3	2	0	2	8.7	-	14.4
Ill-defined (780-795)	26	0	26	26	1	25	112.7	10.9	180.2

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
RANKIN COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	246	74	172	847	515	332	22.3	21.9	22.9
Fetal deaths	11	3	8	36	23	13	42.5	44.7	39.2
Total deaths	395	196	199	275	158	117	7.2	6.7	8.1
Deaths under 1 year	11	4	7	32	17	15	37.8	33.0	45.2
Deaths under 28 days	5	1	4	26	14	12	30.7	27.2	36.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	3	2	1	0	0	0	-	-	-
Syphilis (020-029)	1	0	1	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	6	2	4	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	41	13	28	36	19	17	94.6	80.7	117.2
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	2	2	0	1	1	0	2.6	4.2	-
Diabetes mellitus (260)	7	4	3	5	2	3	13.1	8.5	20.7
Anemias (290-293)	2	0	2	1	0	1	2.6	-	6.9
Intracranial lesions (330-334)	59	25	34	32	17	15	84.1	72.2	103.4
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	100	57	43	90	51	39	236.5	216.6	268.8
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	5	5	0	2	2	0	5.2	8.5	-
Other circulatory diseases (451-468)	15	5	10	3	2	1	7.9	8.5	6.9
#Influenza & pneumonia (480-493)	25	10	15	6	4	2	15.8	17.0	13.8
Bronchitis (500-502)	1	1	0	1	1	0	2.6	4.2	-
Stomach & duodenal ulcers (540, 541)	2	2	0	2	2	0	5.2	8.5	-
#Gastritis, etc. (543, 571, 572)	5	1	4	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	6	5	1	2	2	0	5.2	8.5	-
Cirrhosis of liver (581)	2	2	0	1	1	0	2.6	4.2	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	1	2	0	0	0	-	-	-
Infections of kidney (600)	6	2	4	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	4	2	2	5	3	2	13.1	12.7	13.8
Diseases of early infancy (760-776)	5	0	5	23	11	12	27.2	21.4	36.1
Accidents (800-962)	25	13	12	28	16	12	73.6	68.0	82.7
Motor vehicle acc. (810-835)	4	2	2	11	7	4	28.9	29.7	27.6
Suicide (963, 970-979)	3	3	0	3	3	0	7.9	12.7	-
Homicide (964, 980-985)	7	3	4	5	2	3	13.1	8.5	20.7
Ill-defined (780-795)	31	14	17	16	8	8	42.0	34.0	55.1

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
SCOTT COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	505	325	180	507	273	234	22.0	18.6	27.8
Fetal deaths	11	2	9	14	3	11	27.6	11.0	47.0
Total deaths	245	141	104	256	151	105	11.1	10.3	12.5
Deaths under 1 year	12	6	6	14	5	9	27.6	18.3	38.5
Deaths under 28 days	8	5	3	9	3	6	17.8	11.0	25.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	1	0	4.3	6.8	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	32	18	14	33	21	12	143.2	143.4	142.7
Benign & unspec. neoplasms (210-239)	0	0	0	1	1	0	4.3	6.8	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	1	0	1	2	1	1	8.7	6.8	11.9
Anemias (290-293)	1	1	0	0	0	0	-	-	-
Intracranial lesions (330-334)	23	11	12	26	15	11	112.8	102.4	130.8
Nonmeningococcal meningitis (340)	0	0	0	1	0	1	4.3	-	11.9
Heart diseases (400-443)	120	74	46	112	71	41	485.9	485.0	487.5
Other hypertensive disease (444-447)	2	0	2	4	2	2	17.4	13.7	23.8
General arteriosclerosis (450)	1	0	1	3	2	1	13.0	13.7	11.9
Other circulatory diseases (451-468)	1	1	0	2	1	1	8.7	6.8	11.9
#Influenza & pneumonia (480-493)	4	2	2	4	2	2	17.4	13.7	23.8
Bronchitis (500-502)	0	0	0	1	1	0	4.3	6.8	-
Stomach & duodenal ulcers (540, 541)	1	1	0	1	1	0	4.3	6.8	-
#Gastritis, etc. (543, 571, 572)	0	0	0	2	2	0	8.7	13.7	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	1	1	0	1	1	0	4.3	6.8	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	0	1	4.3	-	11.9
Infections of kidney (600)	0	0	0	1	1	0	4.3	6.8	-
Hyperplasia of prostate (610)	1	0	1	1	0	1	4.3	-	11.9
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	1	0	1	1	0	4.3	6.8	-
Diseases of early infancy (760-776)	9	5	4	9	3	6	17.8	11.0	25.6
Accidents (800-962)	25	19	6	24	16	8	104.1	109.3	95.1
Motor vehicle acc. (810-835)	15	12	3	11	8	3	47.7	54.6	35.7
Suicide (963, 970-979)	3	3	0	2	2	0	8.7	13.7	-
Homicide (964, 980-985)	3	0	3	4	1	3	17.4	6.8	35.7
Ill-defined (780-795)	11	1	10	11	0	11	47.7	-	130.8

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
SHARKEY COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	181	27	154	341	56	285	32.3	18.8	37.6
Fetal deaths	6	0	6	17	4	13	49.8	71.4	45.6
Total deaths	112	22	90	123	28	95	11.6	9.4	12.5
Deaths under 1 year	10	0	10	15	1	14	44.0	17.8	49.1
Deaths under 28 days	4	0	4	6	1	5	17.6	17.8	17.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	0	1	1	0	1	9.5	-	13.2
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	9	2	7	11	4	7	104.3	134.2	92.5
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	1	0	1	0	0	0	-	-	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	8	0	8	10	1	9	94.8	33.6	118.9
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	38	16	22	32	13	19	303.3	436.2	251.0
Other hypertensive disease (444-447)	1	0	1	1	0	1	9.5	-	13.2
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	0	0	0	1	1	0	9.5	33.6	-
#Influenza & pneumonia (480-493)	12	2	10	14	2	12	132.7	67.1	158.5
Bronchitis (500-502)	1	1	0	1	1	0	9.5	33.6	-
Stomach & duodenal ulcers (540, 541)	0	0	0	1	1	0	9.5	33.6	-
#Gastritis, etc. (543, 571, 572)	1	0	1	2	0	2	19.0	-	26.4
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	0	1	9.5	-	13.2
Cirrhosis of liver (581)	1	1	0	1	1	0	9.5	33.6	-
Acute nephritis (590, 591)	0	0	0	1	1	0	9.5	33.6	-
Chronic nephritis, etc. (592-594)	5	0	5	5	0	5	47.4	-	66.0
Infections of kidney (600)	2	0	2	2	0	2	19.0	-	26.4
Hyperplasia of prostate (610)	0	0	0	1	1	0	9.5	33.6	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	3	0	3	5	1	4	14.7	17.8	14.0
Accidents (800-962)	15	0	15	19	1	18	180.1	33.6	237.8
Motor vehicle acc. (810-835)	0	0	0	1	1	0	9.5	33.6	-
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	12	0	12	12	0	12	113.7	-	158.5

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
SIMPSON COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	390	233	157	456	242	214	20.5	16.0	30.2
Fetal deaths	14	9	5	17	13	4	37.3	53.7	18.7
Total deaths	278	187	91	235	164	71	10.6	10.8	10.0
Deaths under 1 year	15	5	10	15	5	10	32.9	20.7	46.7
Deaths under 28 days	2	1	1	6	3	3	13.2	12.4	14.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	34	17	17	1	1	0	4.5	6.6	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	28	22	6	27	23	4	121.4	151.7	56.5
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	3	2	1	4	3	1	18.0	19.8	14.1
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	43	31	12	42	30	12	188.8	197.9	169.5
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	82	62	20	74	55	19	332.7	362.8	268.4
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	3	1	2	4	2	2	18.0	13.2	28.2
Other circulatory diseases (451-468)	2	2	0	2	2	0	9.0	13.2	-
#Influenza & pneumonia (480-493)	19	12	7	15	6	9	67.4	39.6	127.1
Bronchitis (500-502)	0	0	0	1	1	0	4.5	6.6	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	2	0	2	2	0	2	9.0	-	28.2
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	2	1	2	1	1	9.0	6.6	14.1
Infections of kidney (600)	2	2	0	1	1	0	4.5	6.6	-
Hyperplasia of prostate (610)	2	2	0	1	1	0	4.5	6.6	-
Maternal causes (640-689)	0	0	0	1	1	0	21.9	41.3	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	2	1	1	6	3	3	13.2	12.4	14.0
Accidents (800-962)	14	9	5	16	12	4	71.9	79.2	56.5
Motor vehicle acc. (810-835)	6	3	3	8	5	3	36.0	33.0	42.4
Suicide (963, 970-979)	2	2	0	2	2	0	9.0	13.2	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	23	10	13	21	9	12	94.4	59.4	169.5

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
SMITH COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	45	16	29	322	176	146	20.3	14.6	38.4
Fetal deaths	2	0	2	12	4	8	37.3	22.7	54.8
Total deaths	77	60	17	155	121	34	9.8	10.0	8.9
Deaths under 1 year	5	2	3	19	10	9	59.0	56.8	61.6
Deaths under 28 days	0	0	0	7	4	3	21.7	22.7	20.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	1	0	6.3	8.3	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	10	9	1	22	18	4	138.7	149.2	105.3
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	0	0	0	1	1	0	6.3	8.3	-
Anemias (290-293)	0	0	0	1	1	0	6.3	8.3	-
Intracranial lesions (330-334)	10	8	2	23	19	4	145.0	157.5	105.3
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	34	28	6	52	44	8	327.9	364.8	210.5
Other hypertensive disease (444-447)	1	1	0	1	1	0	6.3	8.3	-
General arteriosclerosis (450)	3	3	0	3	3	0	18.9	24.9	-
Other circulatory diseases (451-468)	0	0	0	1	1	0	6.3	8.3	-
#Influenza & pneumonia (480-493)	3	1	2	9	6	3	56.7	49.8	78.9
Bronchitis (500-502)	0	0	0	1	1	0	6.3	8.3	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	1	1	0	3	1	2	18.9	8.3	52.6
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	0	1	6.3	-	26.3
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	6.3	8.3	-
Infections of kidney (600)	0	0	0	1	0	1	6.3	-	26.3
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	1	1	0	1	1	0	31.0	56.8	-
Congenital malformations (750-759)	0	0	0	2	2	0	12.6	16.6	-
Diseases of early infancy (760-776)	0	0	0	7	5	2	21.7	28.4	13.7
Accidents (800-962)	8	5	3	11	9	2	69.4	74.6	52.6
Motor vehicle acc. (810-835)	4	3	1	4	4	0	25.2	33.2	-
Suicide (963, 970-979)	1	1	0	1	1	0	6.3	8.3	-
Homicide (964, 980-985)	2	1	1	2	1	1	12.6	8.3	26.3
Ill-defined (780-795)	0	0	0	3	1	2	18.9	8.3	52.6

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
STONE COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	148	89	59	141	80	61	21.3	16.4	34.6
Fetal deaths	4	3	1	4	3	1	28.4	37.5	16.4
Total deaths	66	51	15	60	43	17	9.0	8.8	9.6
Deaths under 1 year	5	4	1	5	2	3	35.5	25.0	49.2
Deaths under 28 days	4	4	0	3	2	1	21.3	25.0	16.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	8	7	1	9	8	1	135.7	164.3	56.8
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	1	1	0	0	0	0	-	-	-
Anemias (290-293)	1	1	0	1	1	0	15.1	20.5	-
Intracranial lesions (330-334)	9	6	3	8	5	3	120.7	102.7	170.4
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	23	22	1	22	20	2	331.8	410.7	113.6
Other hypertensive disease (444-447)	1	0	1	1	0	1	15.1	-	56.8
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	5	4	1	6	4	2	90.5	82.1	113.6
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	1	0	1	1	0	1	15.1	-	56.8
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	4	4	0	3	2	1	21.3	25.0	16.4
Accidents (800-962)	8	2	6	7	2	5	105.6	41.1	284.1
Motor vehicle acc. (810-835)	3	2	1	3	2	1	45.2	41.1	56.8
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	1	0	1	0	0	0	-	-	-
Ill-defined (780-795)	2	2	0	0	0	0	-	-	-

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
SUNFLOWER COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,214	193	1,021	1,344	237	1,107	27.9	16.0	33.2
Fetal deaths	55	5	50	63	7	56	46.9	29.5	50.6
Total deaths	436	99	337	496	131	365	10.3	8.8	10.9
Deaths under 1 year	70	8	62	66	7	59	49.1	29.5	53.3
Deaths under 28 days	28	5	23	26	4	22	19.3	16.9	19.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	0	1	2	1	1	4.2	6.7	3.0
Syphilis (020-029)	1	0	1	1	0	1	2.1	-	3.0
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	42	6	36	56	12	44	116.2	81.0	131.8
Benign & unspec. neoplasms (210-239)	0	0	0	1	1	0	2.1	6.7	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	6	2	4	7	3	4	14.5	20.2	12.0
Anemias (290-293)	0	0	0	1	0	1	2.1	-	3.0
Intracranial lesions (330-334)	81	13	68	86	19	67	178.5	128.2	200.8
Nonmeningococcal meningitis (340)	0	0	0	1	0	1	2.1	-	3.0
Heart diseases (400-443)	114	44	70	134	56	78	278.1	377.9	233.7
Other hypertensive disease (444-447)	4	0	4	4	0	4	8.3	-	12.0
General arteriosclerosis (450)	4	1	3	5	1	4	10.4	6.7	12.0
Other circulatory diseases (451-468)	0	0	0	1	0	1	2.1	-	3.0
#Influenza & pneumonia (480-493)	35	2	33	39	4	35	80.9	27.0	104.9
Bronchitis (500-502)	1	0	1	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	1	0	1	1	0	1	2.1	-	3.0
#Gastritis, etc. (543, 571, 572)	7	1	6	8	2	6	16.6	13.5	18.0
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	2	1	1	4.2	6.7	3.0
Cirrhosis of liver (581)	1	0	1	2	1	1	4.2	6.7	3.0
Acute nephritis (590, 591)	1	1	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	2	2	6	2	4	12.4	13.5	12.0
Infections of kidney (600)	2	0	2	2	0	2	4.2	-	6.0
Hyperplasia of prostate (610)	1	0	1	3	0	3	6.2	-	9.0
Maternal causes (640-689)	0	0	0	1	0	1	7.4	-	9.0
Congenital malformations (750-759)	3	2	1	6	3	3	12.4	20.2	9.0
Diseases of early infancy (760-776)	25	4	21	22	3	19	16.4	12.6	17.2
Accidents (800-962)	30	9	21	36	11	25	74.7	74.2	74.9
Motor vehicle acc. (810-835)	14	6	8	19	8	11	39.4	54.0	33.0
Suicide (963, 970-979)	3	3	0	3	3	0	6.2	20.2	-
Homicide (964, 980-985)	10	1	9	4	0	4	8.3	-	12.0
Ill-defined (780-795)	47	3	44	49	2	47	101.7	13.5	140.8

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
TALLAHATCHIE COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	670	105	565	735	148	587	29.5	16.6	36.7
Fetal deaths	21	1	20	25	3	22	34.0	20.3	37.5
Total deaths	257	77	180	298	101	197	12.0	11.3	12.3
Deaths under 1 year	41	4	37	46	6	40	62.6	40.5	68.1
Deaths under 28 days	24	2	22	26	3	23	35.4	20.3	39.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	2	0	2	2	0	2	8.0	-	12.5
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	4.0	-	6.2
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	1	1	0	1	1	0	4.0	11.2	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	25	11	14	35	15	20	140.4	168.0	125.1
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	2	1	1	2	1	1	8.0	11.2	6.2
Diabetes mellitus (260)	5	0	5	4	0	4	16.0	-	25.0
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	26	6	20	32	8	24	128.4	89.6	150.1
Nonmeningococcal meningitis (340)	0	0	0	2	1	1	8.0	11.2	6.2
Heart diseases (400-443)	79	37	42	90	44	46	361.2	492.7	287.7
Other hypertensive disease (444-447)	5	1	4	5	2	3	20.1	22.4	18.8
General arteriosclerosis (450)	4	1	3	7	4	3	28.1	44.8	18.8
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	12	1	11	15	4	11	60.2	44.8	68.8
Bronchitis (500-502)	0	0	0	1	0	1	4.0	-	6.2
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	5	0	5	5	0	5	20.1	-	31.3
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	1	1	0	4.0	11.2	-
Cirrhosis of liver (581)	2	2	0	3	3	0	12.0	33.6	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	1	0	0	-	-	-
Hyperplasia of prostate (610)	2	0	2	2	0	2	8.0	-	12.5
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	2	1	1	2	1	1	8.0	11.2	6.2
Diseases of early infancy (760-776)	22	2	20	23	3	20	31.3	20.3	34.1
Accidents (800-962)	19	6	13	18	5	13	72.2	56.0	81.3
Motor vehicle acc. (810-835)	5	3	2	4	2	2	16.0	22.4	12.5
Suicide (963, 970-979)	2	2	0	2	2	0	8.0	22.4	-
Homicide (964, 980-985)	6	0	6	7	0	7	28.1	-	43.8
Ill-defined (780-795)	27	3	24	28	3	25	112.4	33.6	156.3

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
TATE COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	344	3	341	564	150	414	26.3	16.2	34.0
Fetal deaths	9	0	9	17	1	16	30.1	6.7	38.6
Total deaths	146	52	94	209	91	118	9.7	9.8	9.7
Deaths under 1 year	14	1	13	25	7	18	44.3	46.7	43.5
Deaths under 28 days	7	0	7	13	4	9	23.0	26.7	21.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	2	0	2	2	0	2	9.3	-	16.4
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	18	10	8	25	14	11	116.4	150.9	90.2
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	0	2	2	0	2	9.3	-	16.4
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	21	8	13	32	15	17	149.0	161.6	139.4
Nonmeningococcal meningitis (340)	0	0	0	1	0	1	4.6	-	8.2
Heart diseases (400-443)	59	25	34	78	37	41	363.3	398.7	336.3
Other hypertensive disease (444-447)	1	0	1	1	0	1	4.6	-	8.2
General arteriosclerosis (450)	1	1	0	1	1	0	4.6	10.8	-
Other circulatory diseases (451-468)	0	0	0	4	2	2	18.6	21.6	16.4
#Influenza & pneumonia (480-493)	5	1	4	6	1	5	27.9	10.8	41.0
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	1	0	1	4.6	-	8.2
#Gastritis, etc. (543, 571, 572)	0	0	0	1	0	1	4.6	-	8.2
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	2	2	0	9.3	21.6	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	4.6	-	8.2
Infections of kidney (600)	0	0	0	1	1	0	4.6	10.8	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	1	1	0	1	1	0	17.7	66.7	-
Congenital malformations (750-759)	1	0	1	4	2	2	18.6	21.6	16.4
Diseases of early infancy (760-776)	5	0	5	9	2	7	16.0	13.3	16.9
Accidents (800-962)	6	3	3	8	5	3	37.3	53.9	24.6
Motor vehicle acc. (810-835)	1	1	0	1	1	0	4.6	10.8	-
Suicide (963, 970-979)	1	1	0	1	1	0	4.6	10.8	-
Homicide (964, 980-985)	2	0	2	2	0	2	9.3	-	16.4
Ill-defined (780-795)	16	0	16	15	1	14	69.9	10.8	114.8

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
TIPPAH COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	388	286	102	349	252	97	20.9	18.7	30.1
Fetal deaths	33	25	8	27	19	8	77.4	75.4	82.5
Total deaths	145	110	35	163	131	32	9.7	9.7	9.9
Deaths under 1 year	12	6	6	15	8	7	43.0	31.7	72.2
Deaths under 28 days	6	5	1	7	6	1	20.0	23.8	10.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	21	19	2	26	24	2	155.5	177.8	62.1
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	0	0	0	1	1	0	6.0	7.4	-
Anemias (290-293)	0	0	0	1	1	0	6.0	7.4	-
Intracranial lesions (330-334)	24	18	6	25	20	5	149.5	148.1	155.3
Nonmeningoococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	61	49	12	64	55	9	382.8	407.4	279.5
Other hypertensive disease (444-447)	1	1	0	1	1	0	6.0	7.4	-
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	1	1	0	4	3	1	23.9	22.2	31.0
#Influenza & pneumonia (480-493)	7	3	4	9	5	4	53.8	37.0	124.2
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	1	0	1	1	0	1	6.0	-	31.0
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	3	2	1	1	0	1	6.0	-	31.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	1	0	1	1	0	1	6.0	-	31.0
Maternal causes (640-689)	1	1	0	1	1	0	6.0	7.4	-
Congenital malformations (750-759)	2	2	0	3	3	0	17.9	22.2	-
Diseases of early infancy (760-776)	7	5	2	9	6	3	25.8	23.8	30.9
Accidents (800-962)	10	5	5	8	5	3	47.8	37.0	93.2
Motor vehicle acc. (810-835)	5	4	1	4	4	0	23.9	29.6	-
Suicide (963, 970-979)	0	0	0	1	1	0	6.0	7.4	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	1	0	1	1	0	1	6.0	-	31.0

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
TISHOMINGO COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	221	207	14	311	290	21	18.7	18.2	29.6
Fetal deaths	3	3	0	3	3	0	9.6	10.3	-
Total deaths	136	129	7	161	153	8	9.7	9.6	11.3
Deaths under 1 year	2	2	0	8	8	0	25.7	27.6	-
Deaths under 28 days	0	0	0	6	6	0	19.3	20.7	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	1	0	1	1	0	6.0	6.3	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	21	21	0	24	24	0	144.2	150.6	-
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	0	0	0	0	0	0	-	-	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	30	29	1	30	29	1	180.3	182.0	140.8
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	51	48	3	65	61	4	390.6	382.9	563.4
Other hypertensive disease (444-447)	1	1	0	1	1	0	6.0	6.3	-
General arteriosclerosis (450)	1	1	0	2	2	0	12.0	12.6	-
Other circulatory diseases (451-468)	1	1	0	3	3	0	18.0	18.8	-
#Influenza & pneumonia (480-493)	5	4	1	5	4	1	30.0	25.1	140.8
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	0	0	0	-	-	-
Cirrhosis of liver (581)	2	2	0	1	1	0	6.0	6.3	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	6.0	6.3	-
Infections of kidney (600)	1	0	1	1	0	1	6.0	-	140.8
Hyperplasia of prostate (610)	1	1	0	1	1	0	6.0	6.3	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	0	0	0	6	6	0	19.3	20.7	-
Accidents (800-962)	11	10	1	13	12	1	78.1	75.3	140.8
Motor vehicle acc. (810-835)	6	5	1	9	8	1	54.1	50.2	140.8
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	1	1	0	1	1	0	6.0	6.3	-
Ill-defined (780-795)	5	5	0	3	3	0	18.0	18.8	-

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
TUNICA COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	598	55	543	601	60	541	35.4	17.0	40.2
Fetal deaths	21	0	21	20	0	20	33.3	-	37.0
Total deaths	212	33	179	223	33	190	13.1	9.4	14.1
Deaths under 1 year	42	0	42	43	1	42	71.5	16.7	77.6
Deaths under 28 days	23	0	23	23	0	23	38.3	-	42.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	2	0	2	3	0	3	17.7	-	22.3
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	2	0	2	2	0	2	11.8	-	14.9
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	14	7	7	18	5	13	106.1	142.0	96.7
Benign & unspec. neoplasms (210-239)	0	0	0	1	0	1	5.9	-	7.4
Asthma (241)	1	0	1	1	0	1	5.9	-	7.4
Diabetes mellitus (260)	3	1	2	3	1	2	17.7	28.4	14.9
Anemias (290-293)	0	0	0	1	1	0	5.9	28.4	-
Intracranial lesions (330-334)	29	3	26	29	4	25	171.0	113.6	186.0
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	51	8	43	56	14	42	330.2	397.7	312.5
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	25	1	24	26	1	25	153.3	28.4	186.0
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	1	0	1	1	0	1	5.9	-	7.4
#Gastritis, etc. (543, 571, 572)	2	0	2	3	0	3	17.7	-	22.3
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	5.9	-	7.4
Infections of kidney (600)	1	0	1	2	0	2	11.8	-	14.9
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	23	0	23	24	1	23	39.9	16.7	42.5
Accidents (800-962)	19	9	10	14	3	11	82.5	85.2	81.8
Motor vehicle acc. (810-835)	10	5	5	6	2	4	35.4	56.8	29.8
Suicide (963, 970-979)	1	1	0	0	0	0	-	-	-
Homicide (964, 980-985)	2	0	2	2	0	2	11.8	-	14.9
Ill-defined (780-795)	27	3	24	26	2	24	153.3	56.8	178.6

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
UNION COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	315	247	68	367	297	70	17.6	17.0	21.3
Fetal deaths	28	21	7	25	20	5	68.1	67.3	71.4
Total deaths	215	162	53	237	180	57	11.4	10.3	17.3
Deaths under 1 year	16	9	7	17	10	7	46.3	33.7	100.0
Deaths under 28 days	8	5	3	9	6	3	24.5	20.2	42.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	0	1	4.8	-	30.4
Syphilis (020-029)	1	0	1	1	0	1	4.8	-	30.4
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	18	16	2	22	20	2	105.8	114.2	60.8
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	1	1	0	1	1	0	4.8	5.7	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	34	23	11	38	25	13	182.7	142.8	395.1
Nonmeningococcal meningitis (340)	1	1	0	1	1	0	4.8	5.7	-
Heart diseases (400-443)	82	67	15	85	69	16	408.6	394.1	486.3
Other hypertensive disease (444-447)	5	2	3	5	2	3	24.0	11.4	91.2
General arteriosclerosis (450)	0	0	0	1	1	0	4.8	5.7	-
Other circulatory diseases (451-468)	2	2	0	2	2	0	9.6	11.4	-
#Influenza & pneumonia (480-493)	21	13	8	21	13	8	101.0	74.2	243.2
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	2	2	0	3	3	0	14.4	17.1	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	1	1	0	4.8	5.7	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	2	0	3	3	0	14.4	17.1	-
Infections of kidney (600)	1	1	0	3	3	0	14.4	17.1	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	1	1	0	1	1	0	27.2	33.7	-
Congenital malformations (750-759)	1	1	0	2	2	0	9.6	11.4	-
Diseases of early infancy (760-776)	6	4	2	7	5	2	19.1	16.8	28.6
Accidents (800-962)	12	8	4	16	12	4	76.9	68.5	121.6
Motor vehicle acc. (810-835)	5	5	0	9	9	0	43.3	51.4	-
Suicide (963, 970-979)	3	3	0	3	3	0	14.4	17.1	-
Homicide (964, 980-985)	2	0	2	2	0	2	9.6	-	60.8
Ill-defined (780-795)	15	10	5	14	9	5	67.3	51.4	152.0

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
WALTHALL COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	453	254	199	289	115	174	21.5	15.5	28.8
Fetal deaths	24	15	9	9	4	5	31.1	34.8	28.7
Total deaths	138	75	63	135	73	62	10.0	9.8	10.3
Deaths under 1 year	20	4	16	18	4	14	62.3	34.8	80.4
Deaths under 28 days	9	2	7	8	2	6	27.7	17.4	34.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	1	0	7.4	13.5	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	18	12	6	22	14	8	163.6	188.7	132.7
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	1	0	1	0	0	0	-	-	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	25	15	10	23	13	10	171.0	175.2	165.8
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	41	30	11	32	24	8	237.9	323.4	132.7
Other hypertensive disease (444-447)	2	1	1	2	1	1	14.9	13.5	16.6
General arteriosclerosis (450)	3	0	3	3	0	3	22.3	-	49.8
Other circulatory diseases (451-468)	1	1	0	1	1	0	7.4	13.5	-
#Influenza & pneumonia (480-493)	5	0	5	4	0	4	29.7	-	66.3
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	2	1	1	2	1	1	14.9	13.5	16.6
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	0	1	1	0	1	7.4	-	16.6
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	2	1	1	2	1	1	14.9	13.5	16.6
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	3	1	2	3	1	2	22.3	13.5	33.2
Accidents (800-962)	8	2	6	8	2	6	27.7	17.4	34.5
Motor vehicle acc. (810-835)	8	4	4	10	4	6	74.3	53.9	99.5
Suicide (963, 970-979)	5	3	2	7	4	3	52.0	53.9	49.8
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	11	2	9	12	3	9	89.2	40.4	149.2

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
WARREN COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,471	570	901	1,088	482	606	24.6	19.7	30.7
Fetal deaths	112	56	56	63	38	25	57.9	78.8	41.2
Total deaths	707	286	421	548	249	299	12.4	10.2	15.2
Deaths under 1 year	83	15	68	55	15	40	50.6	31.1	66.0
Deaths under 28 days	64	12	52	46	13	33	42.3	27.0	54.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	1	0	1	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	102	56	46	76	46	30	172.1	188.2	152.1
Benign & unspec. neoplasms (210-239)	2	0	2	1	0	1	2.3	-	5.1
Asthma (241)	1	0	1	1	0	1	2.3	-	5.1
Diabetes mellitus (260)	16	3	13	11	3	8	24.9	12.3	40.6
Anemias (290-293)	1	0	1	1	0	1	2.3	-	5.1
Intracranial lesions (330-334)	95	38	57	87	35	52	197.0	143.2	263.7
Nonmeningococcal meningitis (340)	4	0	4	1	0	1	2.3	-	5.1
Heart diseases (400-443)	172	84	88	142	72	70	321.6	294.6	355.0
Other hypertensive disease (444-447)	10	7	3	7	6	1	15.8	24.5	5.1
General arteriosclerosis (450)	11	7	4	9	6	3	20.4	24.5	15.2
Other circulatory diseases (451-468)	3	0	3	3	0	3	6.8	-	15.2
#Influenza & pneumonia (480-493)	17	6	11	17	8	9	38.5	32.7	45.6
Bronchitis (500-502)	4	2	2	2	2	0	4.5	8.2	-
Stomach & duodenal ulcers (540, 541)	2	1	1	2	1	1	4.5	4.1	5.1
#Gastritis, etc. (543, 571, 572)	5	0	5	2	0	2	4.5	-	10.1
Hernia & intestinal obstruction (560, 561, 570)	18	5	13	10	4	6	22.6	16.4	30.4
Cirrhosis of liver (581)	3	1	2	3	2	1	6.8	8.2	5.1
Acute nephritis (590, 591)	3	1	2	1	0	1	2.3	-	5.1
Chronic nephritis, etc. (592-594)	8	4	4	4	1	3	9.0	4.1	15.2
Infections of kidney (600)	4	2	2	6	3	3	13.6	12.3	15.2
Hyperplasia of prostate (610)	5	1	4	1	0	1	2.3	-	5.1
Maternal causes (640-689)	3	0	3	0	0	0	-	-	-
Congenital malformations (750-759)	8	4	4	5	4	1	11.3	16.4	5.1
Diseases of early infancy (760-776)	58	9	49	42	10	32	38.6	20.7	52.8
Accidents (800-962)	49	26	23	23	17	6	52.1	69.6	30.4
Motor vehicle acc. (810-835)	20	10	10	12	8	4	27.2	32.7	20.3
Suicide (963, 970-979)	3	3	0	3	3	0	6.8	12.3	-
Homicide (964, 980-985)	3	0	3	3	0	3	6.8	-	15.2
Ill-defined (780-795)	67	8	59	63	12	51	142.7	49.1	258.6

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
WASHINGTON COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	2,440	863	1,577	2,328	797	1,531	30.2	23.6	35.2
Fetal deaths	232	141	91	222	124	98	95.4	155.6	64.0
Total deaths	838	257	581	893	249	644	11.6	7.4	14.8
Deaths under 1 year	90	13	77	95	12	83	40.8	15.0	54.2
Deaths under 28 days	61	11	50	64	9	55	27.5	11.3	35.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	4	0	4	5.2	-	9.2
Syphilis (020-029)	2	1	1	4	1	3	5.2	3.0	6.9
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	1	0	1	1.3	-	2.3
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	0	1	1	0	1	1.3	-	2.3
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	1	1	0	1	1	0	1.3	3.0	-
Malignant neoplasms (140-205)	83	37	46	89	34	55	115.3	100.9	126.5
Benign & unspec. neoplasms (210-239)	3	1	2	4	1	3	5.2	3.0	6.9
Asthma (241)	1	0	1	1	0	1	1.3	-	2.3
Diabetes mellitus (260)	8	3	5	9	2	7	11.7	5.9	16.1
Anemias (290-293)	5	0	5	6	0	6	7.8	-	13.8
Intracranial lesions (330-334)	106	29	77	108	28	80	139.9	83.1	184.0
Nonmeningococcal meningitis (340)	1	1	0	1	1	0	1.3	3.0	-
Heart diseases (400-443)	234	84	150	247	83	164	320.0	246.2	377.3
Other hypertensive disease (444-447)	7	2	5	8	1	7	10.4	3.0	16.1
General arteriosclerosis (450)	12	.9	3	9	5	4	11.7	14.8	9.2
Other circulatory diseases (451-468)	3	0	3	8	4	4	10.4	11.9	9.2
#Influenza & pneumonia (480-493)	58	14	44	53	11	42	68.7	32.6	96.6
Bronchitis (500-502)	1	1	0	2	1	1	2.6	3.0	2.3
Stomach & duodenal ulcers (540, 541)	3	1	2	2	0	2	2.6	-	4.6
#Gastritis, etc. (543, 571, 572)	12	4	8	13	5	8	16.8	14.8	18.4
Hernia & intestinal obstruction (560, 561, 570)	3	0	3	5	0	5	6.5	-	11.5
Cirrhosis of liver (581)	0	0	0	1	1	0	1.3	3.0	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	10	2	8	14	2	12	18.1	5.9	27.6
Infections of kidney (600)	4	0	4	4	0	4	5.2	-	9.2
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	1	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	53	10	43	58	9	49	24.9	11.3	32.0
Accidents (800-962)	51	16	35	54	15	39	70.0	44.5	89.7
Motor vehicle acc. (810-835)	20	10	10	17	9	8	22.0	26.7	18.4
Suicide (963, 970-979)	5	2	3	6	3	3	7.8	8.9	6.9
Homicide (964, 980-985)	13	1	12	15	1	14	19.4	3.0	32.2
Ill-defined (780-795)	121	21	100	123	22	101	159.4	65.3	232.3

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
WAYNE COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	482	306	176	413	228	185	25.5	22.9	29.7
Fetal deaths	7	2	5	7	2	5	16.9	8.8	27.0
Total deaths	154	84	70	172	90	82	10.6	9.0	13.2
Deaths under 1 year	16	6	10	18	5	13	43.6	21.9	70.3
Deaths under 28 days	10	6	4	10	5	5	24.2	21.9	27.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	0	1	6.2	-	16.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	11	8	3	15	11	4	92.7	110.6	64.2
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	3	0	3	4	0	4	24.7	-	64.2
Anemias (290-293)	0	0	0	1	0	1	6.2	-	16.0
Intracranial lesions (330-334)	19	6	13	18	5	13	111.2	50.2	208.7
Nonmeningococcal meningitis (340)	2	2	0	2	2	0	12.4	20.1	-
Heart diseases (400-443)	58	39	19	56	36	20	346.1	361.8	321.0
Other hypertensive disease (444-447)	2	2	0	2	2	0	12.4	20.1	-
General arteriosclerosis (450)	3	2	1	4	3	1	24.7	30.2	16.0
Other circulatory diseases (451-468)	1	1	0	2	1	1	12.4	10.0	16.0
#Influenza & pneumonia (480-493)	2	0	2	3	0	3	18.5	-	48.2
Bronchitis (500-502)	0	0	0	1	1	0	6.2	10.0	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	2	0	2	2	0	2	12.4	-	32.1
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	0	1	6.2	-	16.0
Cirrhosis of liver (581)	1	1	0	1	1	0	6.2	10.0	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	1	1	12.4	10.0	16.0
Infections of kidney (600)	1	1	0	1	1	0	6.2	10.0	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	6	4	2	6	3	3	14.5	13.2	16.2
Accidents (800-962)	10	5	5	18	11	7	111.2	110.6	112.4
Motor vehicle acc. (810-835)	5	4	1	9	8	1	55.6	80.4	16.0
Suicide (963, 970-979)	3	3	0	3	3	0	18.5	30.2	-
Homicide (964, 980-985)	3	0	3	4	0	4	24.7	-	64.2
<u>III-defined (780-795)</u>	17	4	13	16	3	13	98.9	30.2	208.7

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
WEBSTER COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	118	55	63	197	116	81	19.0	15.4	28.4
Fetal deaths	7	3	4	8	4	4	40.6	34.5	49.4
Total deaths	96	77	19	115	90	25	11.1	11.9	8.8
Deaths under 1 year	8	0	8	14	4	10	71.1	34.5	123.4
Deaths under 28 days	5	0	5	10	4	6	50.8	34.5	74.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	11	11	0	14	12	2	134.7	159.2	70.2
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	1	0	1	1	0	9.6	13.3	-
Diabetes mellitus (260)	0	0	0	1	1	0	9.6	13.3	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	19	17	2	24	22	2	231.0	291.8	70.2
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	28	26	2	32	28	4	308.0	371.4	140.4
Other hypertensive disease (444-447)	1	0	1	0	0	0	-	-	-
General arteriosclerosis (450)	1	1	0	1	1	0	9.6	13.3	-
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	10	8	2	6	4	2	57.7	53.0	70.2
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	1	1	0	9.6	13.3	-
#Gastritis, etc. (543, 571, 572)	1	0	1	1	0	1	9.6	-	35.1
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	0	1	1	0	1	9.6	-	35.1
Diseases of early infancy (760-776)	5	0	5	10	4	6	50.8	34.5	74.1
Accidents (800-962)	7	6	1	7	6	1	67.4	79.6	35.1
Motor vehicle acc. (810-835)	6	5	1	5	4	1	48.1	53.0	35.1
Suicide (963, 970-979)	1	0	1	1	0	1	9.6	-	35.1
Homicide (964, 980-985)	1	0	1	1	0	1	9.6	-	35.1
Ill-defined (780-795)	6	4	2	6	4	2	57.7	53.0	70.2

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
WILKINSON COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	380	151	229	323	74	249	25.0	18.5	27.9
Fetal deaths	17	7	10	15	1	14	46.4	13.5	56.2
Total deaths	161	81	80	160	54	106	12.4	13.5	11.9
Deaths under 1 year	8	2	6	14	1	13	43.3	13.5	52.2
Deaths under 28 days	4	1	3	10	0	10	31.0	-	40.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	0	1	7.7	-	11.2
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	17	12	5	13	4	9	100.6	100.2	100.8
Benign & unspec. neoplasms (210-239)	1	0	1	1	0	1	7.7	-	11.2
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	1	1	5	1	4	38.7	25.1	44.8
Anemias (290-293)	0	0	0	1	0	1	7.7	-	11.2
Intracranial lesions (330-334)	20	6	14	23	6	17	178.0	150.4	190.4
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	48	32	16	35	17	18	270.9	426.1	201.6
Other hypertensive disease (444-447)	1	0	1	1	0	1	7.7	-	11.2
General arteriosclerosis (450)	1	1	0	1	1	0	7.7	25.1	-
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	12	7	5	9	5	4	69.6	125.3	44.8
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	2	1	1	2	1	1	15.5	25.1	11.2
#Gastritis, etc. (543, 571, 572)	1	0	1	1	0	1	7.7	-	11.2
Hernia & intestinal obstruction (560, 561, 570)	1	0	1	1	0	1	7.7	-	11.2
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	2	1	1	15.5	25.1	11.2
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	1	0	1	7.7	-	11.2
Maternal causes (640-689)	1	0	1	2	0	2	61.9	-	80.3
Congenital malformations (750-759)	3	2	1	2	1	1	15.5	25.1	11.2
Diseases of early infancy (760-776)	5	1	4	11	0	11	34.0	-	44.2
Accidents (800-962)	11	3	8	13	3	10	100.6	75.2	112.0
Motor vehicle acc. (810-835)	0	0	0	1	0	1	7.7	-	11.2
Suicide (963, 970-979)	1	1	0	1	1	0	7.7	25.1	-
Homicide (964, 980-985)	1	0	1	1	0	1	7.7	-	11.2
III-defined (780-795)	19	2	17	20	3	17	154.8	75.2	190.4

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
WINSTON COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	386	167	219	404	166	238	20.6	15.5	26.8
Fetal deaths	4	0	4	8	0	8	19.8	-	33.6
Total deaths	171	97	74	191	111	80	9.8	10.4	9.0
Deaths under 1 year	20	7	13	20	6	14	49.5	36.1	58.8
Deaths under 28 days	12	7	5	12	6	6	29.7	36.1	25.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	2	1	1	2	1	1	10.2	9.4	11.3
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	1	0	1	1	0	1	5.1	-	11.3
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	14	10	4	18	15	3	92.0	140.4	33.8
Benign & unspec. neoplasms (210-239)	0	0	0	1	0	1	5.1	-	11.3
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	1	1	2	1	1	10.2	9.4	11.3
Anemias (290-293)	1	0	1	1	0	1	5.1	-	11.3
Intracranial lesions (330-334)	32	19	13	31	17	14	158.5	159.2	157.6
Nonmeningococcal meningitis (340)	1	0	1	1	0	1	5.1	-	11.3
Heart diseases (400-443)	62	40	22	69	47	22	352.8	440.1	247.7
Other hypertensive disease (444-447)	1	0	1	1	0	1	5.1	-	11.3
General arteriosclerosis (450)	3	1	2	5	2	3	25.6	18.7	33.8
Other circulatory diseases (451-468)	2	2	0	3	3	0	15.3	28.1	-
#Influenza & pneumonia (480-493)	6	2	4	8	3	5	40.9	28.1	56.3
Bronchitis (500-502)	1	0	1	1	0	1	5.1	-	11.3
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	1	1	0	5.1	9.4	-
Cirrhosis of liver (581)	0	0	0	1	1	0	5.1	9.4	-
Acute nephritis (590, 591)	1	0	1	1	0	1	5.1	-	11.3
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	1	0	1	5.1	-	11.3
Hyperplasia of prostate (610)	1	1	0	1	1	0	5.1	9.4	-
Maternal causes (640-689)	1	0	1	1	0	1	24.8	-	42.0
Congenital malformations (750-759)	1	1	0	1	1	0	5.1	9.4	-
Diseases of early infancy (760-776)	13	7	6	13	6	7	32.2	36.1	29.4
Accidents (800-962)	9	6	3	8	5	3	40.9	46.8	33.8
Motor vehicle acc. (810-835)	4	3	1	3	2	1	15.3	18.7	11.3
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	2	0	2	2	0	2	10.2	-	22.5
Ill-defined (780-795)	9	1	8	9	1	8	46.0	9.4	90.1

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
YALOBUSHA COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	288	115	173	287	113	174	22.5	16.0	30.8
Fetal deaths	7	3	4	8	4	4	27.9	35.4	23.0
Total deaths	149	90	59	163	100	63	12.8	14.1	11.2
Deaths under 1 year	11	3	8	15	6	9	52.3	53.1	51.7
Deaths under 28 days	8	3	5	10	5	5	34.8	44.2	28.7
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	1	0	7.8	14.1	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	24	17	7	28	20	8	220.0	282.5	141.6
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	0	1	1	0	1	7.8	-	17.7
Diabetes mellitus (260)	2	0	2	2	0	2	15.7	-	35.4
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	21	15	6	17	11	6	133.5	155.4	106.2
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	44	29	15	46	29	17	361.4	409.6	300.9
Other hypertensive disease (444-447)	0	0	0	1	1	0	7.8	14.1	-
General arteriosclerosis (450)	1	0	1	1	1	0	7.8	14.1	-
Other circulatory diseases (451-468)	2	2	0	4	4	0	31.4	56.5	-
#Influenza & pneumonia (480-493)	7	4	3	10	7	3	78.6	98.9	53.1
Bronchitis (500-502)	4	4	0	1	1	0	7.8	14.1	-
Stomach & duodenal ulcers (540, 541)	1	1	0	1	1	0	7.8	14.1	-
#Gastritis, etc. (543, 571, 572)	2	1	1	2	1	1	15.7	14.1	17.7
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	2	1	1	2	1	1	15.7	14.1	17.7
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	7.8	14.1	-
Infections of kidney (600)	2	2	0	1	1	0	7.8	14.1	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	1	1	0	-	-	-
Diseases of early infancy (760-776)	6	2	4	8	4	4	27.9	35.4	23.0
Accidents (800-962)	10	5	5	11	6	5	86.4	84.7	88.5
Motor vehicle acc. (810-835)	7	5	2	7	5	2	55.0	70.6	35.4
Suicide (963, 970-979)	1	1	0	1	1	0	7.8	14.1	-
Homicide (964, 980-985)	2	1	1	2	1	1	15.7	14.1	17.7
I11-defined (780-795)	11	1	10	12	1	11	94.3	14.1	194.7

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
YAZOO COUNTY, MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	830	243	587	872	262	610	26.0	19.0	31.0
Fetal deaths	27	4	23	32	7	25	36.7	26.7	41.0
Total deaths	338	127	211	393	144	249	11.7	10.4	12.6
Deaths under 1 year	35	4	31	46	5	41	52.8	19.1	67.2
Deaths under 28 days	13	3	10	22	3	19	25.2	11.4	31.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	0	1	3.0	-	5.1
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	33	14	19	44	21	23	131.4	152.4	116.8
Benign & unspec. neoplasms (210-239)	1	0	1	0	0	0	-	-	-
Asthma (241)	1	1	0	1	1	0	3.0	7.2	-
Diabetes mellitus (260)	2	0	2	2	0	2	6.0	-	10.2
Anemias (290-293)	1	1	0	1	1	0	3.0	7.2	-
Intracranial lesions (330-334)	50	16	34	60	17	43	179.2	123.4	218.3
Nonmeningococcal meningitis (340)	1	1	0	1	1	0	3.0	7.2	-
Heart diseases (440-443)	92	43	49	100	48	52	298.7	348.3	264.0
Other hypertensive disease (444-447)	3	0	3	6	1	5	17.9	7.2	25.4
General arteriosclerosis (450)	5	3	2	5	3	2	14.9	21.8	10.2
Other circulatory diseases (451-468)	2	2	0	2	2	0	6.0	14.5	-
#Influenza & pneumonia (480-493)	17	3	14	20	5	15	59.7	36.3	76.1
Bronchitis (500-502)	1	1	0	1	1	0	3.0	7.2	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	6	0	6	6	0	6	17.9	-	30.4
Hernia & intestinal obstruction (560, 561, 570)	1	0	1	1	0	1	3.0	-	5.1
Cirrhosis of liver (581)	1	1	0	2	1	1	6.0	7.2	5.1
Acute nephritis (590, 591)	1	0	1	1	0	1	3.0	-	5.1
Chronic nephritis, etc. (592-594)	5	0	5	7	1	6	20.9	7.2	30.4
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	3	2	1	3	1	2	9.0	7.2	10.2
Maternal causes (640-689)	2	1	1	1	1	0	11.5	38.2	-
Congenital malformations (750-759)	1	1	0	5	2	3	14.9	14.5	15.2
Diseases of early infancy (760-776)	14	4	10	21	4	17	24.1	15.3	27.9
Accidents (800-962)	26	9	17	30	9	21	89.6	65.3	106.6
Motor vehicle acc. (810-835)	11	5	6	13	5	8	38.8	36.3	40.6
Suicide (963, 970-979)	2	2	0	2	2	0	6.0	14.5	-
Homicide (964, 980-985)	6	2	4	7	2	5	20.9	14.5	25.4
Ill-defined (780-795)	39	8	31	40	8	32	119.5	58.0	162.4

See page 57 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
STATE OF MISSISSIPPI, 1963

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	57649	27061	30588	57316	26823	30493	25.0	19.9	32.4
Fetal deaths	2795	1302	1493	2755	1257	1498	48.1	46.9	49.1
Total deaths	23043	11960	11083	23339	12118	11221	10.2	9.0	11.9
Deaths under 1 year	2339	601	1738	2372	610	1762	41.4	22.7	57.8
Deaths under 28 days	1432	461	971	1449	470	979	25.3	17.5	32.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	101	37	64	113	40	73	4.9	3.0	7.8
Syphilis (020-029)	32	8	24	34	9	25	1.5	0.7	2.6
Typhoid fever (040)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	15	4	11	14	3	11	0.6	0.2	1.2
Scarlet fever & streptococcal sore throat (050, 051)	4	1	3	4	1	3	0.2	0.1	0.3
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	5	1	4	5	1	4	0.2	0.1	0.4
Meningococcal infections (057)	6	4	2	6	4	2	0.3	0.3	0.2
Acute poliomyelitis (080)	3	3	0	4	4	0	0.2	0.3	-
Measles (085)	8	4	4	7	3	4	0.3	0.2	0.4
Malignant neoplasms (140-205)	2846	1794	1052	2945	1859	1086	128.6	137.8	115.4
Benign & unspec. neoplasms (210-239)	42	19	23	45	21	24	2.0	1.6	2.5
Asthma (241)	41	29	12	45	31	14	2.0	2.3	1.5
Diabetes mellitus (260)	327	135	192	326	132	194	14.2	9.8	20.6
Anemias (290-293)	56	13	43	60	16	44	2.6	1.2	4.7
Intracranial lesions (330-334)	3125	1630	1495	3154	1650	1504	137.7	122.3	159.8
Nonmeningococcal meningitis (340)	47	14	33	50	13	37	2.2	1.0	3.9
Heart diseases (400-443)	7281	4381	2900	7300	4396	2904	318.8	325.9	308.5
Other hypertensive disease (444-447)	199	77	122	203	77	126	8.9	5.7	13.4
General arteriosclerosis (450)	360	247	113	364	250	114	15.9	18.5	12.1
Other circulatory diseases (451-468)	201	121	80	227	134	93	9.9	9.9	9.9
#Influenza & pneumonia (480-493)	1100	420	680	1113	420	693	48.6	31.1	73.6
Bronchitis (500-502)	53	39	14	58	43	15	2.5	3.2	1.6
Stomach & duodenal ulcers (540, 541)	71	50	21	68	46	22	3.0	3.4	2.3
#Gastritis, etc. (543, 571, 572)	197	39	158	204	43	161	8.9	3.2	17.1
Hernia & intestinal obstruction (560, 561, 570)	115	59	56	117	60	57	5.1	4.4	6.0
Cirrhosis of liver (581)	107	87	20	105	86	19	4.6	6.4	2.0
Acute nephritis (590, 591)	36	13	23	39	13	26	1.7	1.0	2.8
Chronic nephritis, etc. (592-594)	212	84	128	222	90	132	9.7	6.7	14.0
Infections of kidney (600)	164	74	90	170	79	91	7.4	5.8	9.7
Hyperplasia of prostate (610)	62	21	41	66	24	42	2.9	1.8	4.5
Maternal causes (640-689)	44	9	35	44	9	35	7.7	3.4	11.5
Congenital malformations (750-759)	237	132	105	256	147	109	11.2	10.9	11.6
Diseases of early infancy (760-776)	1245	381	864	1262	389	873	22.0	14.5	28.6
Accidents (800-962)	1579	855	724	1571	851	720	68.6	63.1	76.5
Motor vehicle acc. (810-835)	705	447	258	675	435	240	29.5	32.2	25.5
Suicide (963, 970-979)	129	107	22	126	104	22	5.5	7.7	2.3
Homicide (964, 980-985)	256	48	208	267	48	219	11.6	3.6	23.3
Ill-defined (780-795)	1790	449	1341	1770	435	1335	77.3	32.2	141.8

See page 57 for footnotes.