


VITAL STATISTICS

MISSISSIPPI

1968



Mississippi State Board of Health
Jackson, Mississippi

VITAL STATISTICS

MISSISSIPPI

1968

STATISTICAL SERVICES UNIT
MISSISSIPPI STATE BOARD OF HEALTH
P. O. BOX 1700
JACKSON, MISSISSIPPI 39205

TABLE OF CONTENTS

SYNOPSIS OF VITAL STATISTICS, MISSISSIPPI, 1968

	Page
Introduction	I
Description of Bulletin	I
Definitions of Live Birth and Fetal Death	I
Revised Certificate Forms and New Information Items	I
Occurrence and Residence Allocation	II
Urban-Rural Classification	III
Population	III
Rates	III
Classification of Cause of Death	IV
Live Births	IV
Trend	IV
Race	V
County	V
Urban-Rural Variation	V
Natural Increase	V
Attendant	VI
Month	VII
Age of Mother	VII
Education of Parents	VIII
Legitimacy	IX
Total-Birth Order and Average Number of Births per Mother	X
Last Prior Delivery	XII
Prenatal Care	XIII
Complications during Pregnancy and Labor	XIV
Length of Gestation and Birth Weight	XVI
Sex	XVIII
Plurality	XVIII
Birth Injuries and Congenital Malformations	XIX
Fetal Deaths	XX
Reporting and Reliability	XX
Trend and Race	XX
County	XXI
Attendant	XXI
Age of Mother	XXII
Education of Parents	XXII
Legitimacy	XXIII
Total-Birth Order	XXIII
Last Prior Delivery	XXIV
Prenatal Care	XXV
Length of Gestation and Birth Weight	XXVI
Sex	XXVII
Plurality	XXVII
Cause of Fetal Death, Complications of Pregnancy and Labor, Birth Injuries, and Congenital Malformations	XXVIII
Deaths	XXVIII
Trend	XXVIII
Race	XXVIII
County and Urban-Rural Variation	XXVIII
Month	XXVIII

Deaths, continued

Age at Death	XXIX
Leading Causes of Death	XXX
Leading Causes by Age	XXX
Ill-defined and Unknown Causes	XXXII
Specific Causes	XXXII
Heart Disease	XXXII
Malignant Neoplasms	XXXII
Cerebrovascular Disease	XXXIII
Chronic Respiratory Diseases	XXXIII
Influenza and Pneumonia	XXXIV
Accidents	XXXIV
Suicide and Homicide	XXXIV
Injury Undetermined Whether Accidentally or Purposely Inflicted ..	XXXV
Maternal Deaths	XXXV
Infant Mortality	XXXV
Marriages	XXXVIII
Trend	XXXVIII
Month	XXXVIII
Officiant	XXXIX
Age of Bride and Groom	XXXIX
Marriage Order and Previous Marital Status	XL
Nonresident Brides	XLI
Out-of-State Use of Mississippi Licenses	XLI
Divorces	XLII
Trend	XLII
Cause and Complainant	XLII
Minor Children	XLII
Duration of Marriage	XLII
Additional Information Needed	XLIII

POPULATION

Table 1. Estimated, by County, by Race, Mississippi, July 1, 1968	1
---	---

LIVE BIRTHS

Table 2. By Place of Occurrence, and by Place of Residence of Mother, by Race, with Rates, Mississippi, 1968	3
Table 3. By Place of Residence of Mother, by Race and Attendant, and by Sex, Mississippi, 1968	6

FETAL DEATHS

Table 4. By Place of Occurrence, by Place of Residence of Mother, by Race and Attendant, and by Sex, Mississippi, 1968	10
--	----

DEATHS

Table 5.	By Cause of Death, by Race, Residence Data, Mississippi, 1968	14
Table 6.	By Place of Occurrence, and by Place of Residence, by Race, with Rates, Mississippi, 1968	29
Table 7.	By Cause Groups and Certain Selected Causes, by Race, Sex, and Age, with Total Rates, Residence Data, Mississippi, 1968	32

DEATHS UNDER 1 YEAR

Table 8.	By Selected Causes, by Race and Age, with Total Rates, Residence Data, Mississippi, 1968	40
----------	--	----

MARRIAGES AND DIVORCES

Table 9.	Marriages by Place of Licensure, by Place of Occurrence, and by Place of Residence of Bride, by Race of Bride; Divorces by Place of Occurrence, by Race, Mississippi, 1968	44
Table 10.	Divorces by Cause of Divorce, by Race, Mississippi, 1968	47
	Divorces by Number of Minor Children, by Race, Mississippi, 1968 .	47
	Divorces by Number of Years Married, by Race, Mississippi, 1968 ..	47

TIME SERIES

Table 11.	<u>Live Births, Mississippi, 1913-1968</u> , by Year, by Race, with Rates, Occurrence Data 1913-1943, Residence Data 1944-1968	48
Table 12.	<u>Deaths, Mississippi, 1913-1968</u> , by Year, by Race, with Rates, Occurrence Data 1913-1943, Residence Data 1944-1968	50
Table 13.	<u>Maternal Deaths, Mississippi, 1913-1968</u> , by Year, by Race, with Rates, Occurrence Data 1913-1943, Residence Data 1944-1968	52
Table 14.	<u>Deaths Under 1 Year, Mississippi, 1917-1968</u> , by Year, by Race, with Rates, Occurrence Data 1917-1943, Residence Data 1944-1968 ..	54
Table 15.	<u>Deaths Under 1 Month, Mississippi, 1924-1968</u> , by Year, by Race, with Rates, Occurrence Data 1924-1944, Residence Data 1945-1968 ..	56
Table 16.	<u>Deaths from Heart Diseases, Malignant Neoplasms, Cerebrovascular Disease, Accidents, and Influenza and Pneumonia, Mississippi, 1959-1968</u> , by Year, by Race, with Rates, Residence Data	57
Table 17.	<u>Marriages and Divorces, Mississippi, 1926-1968</u> , by Year, by Race, Occurrence Data	58

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH,
by Place of Occurrence, and by Place of Residence, by Race, with Rates, 1968

The Counties (in alphabetical order)	61
State of Mississippi	143

SYNOPSIS OF VITAL STATISTICS MISSISSIPPI, 1968

INTRODUCTION

Description of Bulletin

The 1968 Vital Statistics Bulletin is the twenty-sixth annual report prepared by Statistical Services from information contained on records of live births, fetal deaths, deaths, marriages, and divorces collected by the office of Vital Records Registration. This synopsis provides certain explanations and definitions necessary for understanding the data presented and also summarizes the most noteworthy facts regarding the vital events of the year. Following the synopsis are ten tables (Numbers 1-10) containing statistics for 1968 and seven tables (Numbers 11-17) showing comparable data over a span of years, in most cases the entire period for which the information is available. The last section of the book contains a one-page summary table for each county and also for the state.

Definitions of Live Birth and Fetal Death

The definitions adopted by the Mississippi State Board of Health for use in Mississippi are those formulated by the World Health Organization and recommended by the National Center for Health Statistics. They are as follows:

Live birth - the complete expulsion or extraction from its mother of a product of human conception, irrespective of the duration of pregnancy, which, after such expulsion or extraction, breathes or shows any other evidence of life such as beating of the heart, pulsation of the umbilical cord, or definite movement of voluntary muscles, whether or not the umbilical cord has been cut or the placenta is attached.

Fetal death - death prior to the complete expulsion or extraction from its mother of a product of human conception, irrespective of the duration of pregnancy; the death is indicated by the fact that after such expulsion or extraction the fetus does not breathe or show any other evidence of life such as beating of the heart, pulsation of the umbilical cord, or definite movement of voluntary muscles.

Revised Certificate Forms and New Information Items

Mississippi, as well as most other states, began using revised certificates for live births, fetal deaths, and deaths in 1968. Each state decided upon the format of and items to be included on its own certificates, but all were either modeled after or the same as standard certificates designed by the National Center for Health Statistics in cooperation with the various states. The changes in 1968 were part of a periodic review and revision program undertaken to insure that these certificates adequately serve current legal and statistical needs of individuals and public health programs. The last revision in Mississippi prior to 1968 was in 1956.

Although the new death certificates contain only a relatively small amount of information not on the old certificates, the contents of the live birth and fetal death certificates were greatly expanded. The new information collected was considered to be of special value for planning public health activities directed toward

Vital Statistics
Mississippi - 1968

lowering Mississippi's excessively high infant mortality rate which is one of the state's most serious health problems.

Most of the new data items provide information never before available for Mississippi. However, considerable difficulty was experienced in collecting and tabulating this information. It was found that answers for new items were missing from many certificates; the proportion lacking answers for a certain item ranged from less than one percent of the live birth records to nearly 13 percent and from 6 to 22 percent of the fetal death records. In addition, there were numerous obvious inaccuracies. Because of these deficiencies, data compiled from the new items will have to be considered with some reservations.

Occurrence and Residence Allocation

Live births, fetal deaths, and deaths were classified by place of occurrence and by place of residence. Some of the tables contain both occurrence and residence figures, but most of the tables are based on residence data only, as these data are considered more suitable for studying public health problems. All figures given in this synopsis are residence counts unless otherwise stated.

Occurrence data were obtained from certificates filed in Mississippi at the place of the event. Residence data include transcripts of records of events that occurred in other states to Mississippi residents and exclude records filed in this state for non-residents. Residence assignment for live births and fetal deaths was determined by the usual residence of the mother. For deaths it was determined by the usual residence of decedent, except that deaths of inmates of institutions were reallocated to place of previous residence regardless of length of stay in the institution.

Table A provides information on vital events occurring out of the state to Mississippi residents and in Mississippi to nonresidents. The counts showed that 2,724 events to Mississippians were recorded in other places and that 2,423 records were filed in this state for events to nonresidents. States are listed on the table in order according to the total number of records involved. The largest number was exchanged with Tennessee, which is the only one of the states with an exchange of over 50 records from which Mississippi received more records than it sent. This state, together with the other three states which border on Mississippi (Alabama, Arkansas, and Louisiana), provided 90 percent of the transcripts received; these border states were also listed as the place of residence on 83 percent of the certificates filed in Mississippi for non-residents. However, although most of the vital record interchange was with neighboring states, Mississippi received transcripts from a total of 38 other states and the District of Columbia as well as one from out of the country and filed certificates for residents of 40 other states plus three for foreign residents.

Marriages were classified according to place of issuance of marriage license, place of occurrence of marriage, and place of residence of bride. However, the figures on brides who were Mississippi residents are incomplete since there is no exchange among the states of transcripts of marriage records for nonresidents and the out-of-state marriages of Mississippi brides could not be included. Divorces were classified only according to place where the decree was granted, as there is no information regarding place of residence on the record.

Vital Statistics
Mississippi - 1968

Table A. Transcripts of certificates filed in other places for Mississippi residents, by place of occurrence, and certificates filed in Mississippi for nonresidents, by place of residence, by type of certificate

Place	Total		Live Births		Fetal Deaths		Deaths	
	Transcripts	Certificates, Nonresidents	Transcripts	Certificates, Nonresidents	Transcripts	Certificates, Nonresidents	Transcripts	Certificates, Nonresidents
Total	2,724	2,423	1,497	1,484	30	81	1,197	858
Tennessee	1,667	267	981	163	26	7	660	97
Louisiana	420	888	217	620	0	32	203	236
Alabama	300	739	181	466	2	19	117	254
Arkansas	64	110	23	69	0	2	41	39
Florida	33	76	13	22	0	1	20	53
Illinois	39	65	9	22	1	6	29	37
Texas	48	48	7	20	0	2	41	26
Other places	153	230	66	102	1	12	86	116

Urban-Rural Classification

Incorporated towns with populations of 2,500 or more according to the 1960 census were considered as urban areas, and a total of 62 towns were included in this category. All other areas were classified as rural. These categories were used for live births, fetal deaths, and deaths only. Information on marriage records is inadequate for proper classification, and divorce records do not contain this item.

Population

The Mississippi State Board of Health does not make population estimates but uses those prepared by the Division of Sociology and Anthropology of Mississippi State University. However, these estimates are based in part on live birth and death figures provided by the State Board of Health for this purpose.

Rates

Rates and ratios were calculated from residence data on live births, fetal deaths, and deaths. Estimated populations for the state and counties by race were used to prepare rates for live births, total deaths, and deaths from specific causes (except maternal and certain causes of mortality in early infancy). Although compilation of rates by age and sex and for cities is desirable, this could not be done because of the lack of population data for these categories. Figures on live births were used to calculate fetal death ratios and rates for maternal deaths, infant deaths, neonatal deaths, and deaths from certain causes of mortality in early infancy, as they approximate the groups at risk more closely than do general population figures.

The base for each type of rate used in this bulletin is given in the footnote on page 59.

IV

Vital Statistics Mississippi - 1968

Classification of Cause of Death

Development of a system for classifying diseases began in Europe in the 1700's. By the end of the nineteenth century a standard classification of causes of death was being used in several countries and had been adopted for use in the United States. This classification has been revised approximately every ten years since 1900 in order to keep pace with new knowledge in the field of medicine. The latest version, Eighth Revision International Classification of Diseases, Adapted for Use in the United States, went into effect in Mississippi and a majority of the other states in 1968.

A list of the international classification numbers and group titles for the new revision is presented here to show the organization of the system.

000-136	Infective and parasitic diseases
140-239	Neoplasms
240-279	Endocrine, nutritional, and metabolic diseases
280-289	Diseases of the blood and blood-forming organs
290-315	Mental disorders
320-389	Diseases of the nervous system and sense organs
390-458	Diseases of the circulatory system
460-519	Diseases of the respiratory system
520-577	Diseases of the digestive system
580-629	Diseases of the genitourinary system
630-678	Complications of pregnancy, childbirth, and the puerperium
680-709	Diseases of the skin and subcutaneous tissue
710-738	Diseases of the musculoskeletal system and connective tissue
740-759	Congenital anomalies
760-779	Certain causes of mortality in early infancy
780-796	Symptoms and ill-defined conditions
800-999	Accidents, poisonings, and violence

On certificates with more than one cause entered, only that indicated as the underlying cause was classified, but the system can be used for studying all conditions listed on death certificates.

Although standard classification enables comparison of statistics from one country or state with those of another country or state regardless of language or terminology differences, the periodic revisions which have taken place in the past have resulted in an unavoidable lack of comparability over the years for some causes of death. The greatest change was in the Sixth Revision in 1949, and only minor changes were incorporated in the Seventh Revision which went into use in 1958. Differences between the Seventh and Eighth Revisions have not been fully analyzed yet; but it appears that, although they are of more consequence than those between the Sixth and Seventh Revisions, there was a relatively small number of causes with major changes. Because of these revisions, comparisons of data over a period of years should be made with caution. However, the National Center for Health Statistics has prepared and published comparability ratios which show the effect revisions have had on each cause and which may be used to adjust figures for the purpose of analyzing trends.

LIVE BIRTHS (See Tables 2, 3, 11)

Trend

Mississippi in 1968 experienced its seventh consecutive year of decline in

Vital Statistics
Mississippi - 1968

number of live births, as did the country as a whole. The total count of 45,719 births to Mississippi residents was 23 percent lower than the figure of 59,523 reported for 1961, the last year before the decrease began. However, the rate of decline slowed considerably after reaching a peak of 9.7 percent in 1965 and in 1968 amounted to only 1.9 percent. In 1968 the live birth rate was 19.4 per 1,000 population, but it should be noted that population estimates made this long after the decennial census are subject to sizeable inaccuracies which would, of course, affect any figures based on them. This rate was the second lowest on record, exceeding only that for 1917, and the number of births was the smallest since 1933.

Race

Counts by race showed a difference of only 105 between the figure of 22,807 for whites and that of 22,912 for nonwhites. In every year since 1924 the nonwhite total has been the larger; but if the trend continues in 1969 as it has in recent years, the reverse will be true. Although the numbers for the two groups were nearly the same, there was still a large gap between the rates; for whites the rate was only 15.7 per 1,000 population, but for nonwhites it was 25.4. The figure for whites was the lowest ever calculated, and that for nonwhites was the smallest since 1935.

County

Among both races there was marked variation in the county rates. The range for whites was from 6.1 in Jefferson County to 21.0 in Jackson County; for nonwhites the lowest was 11.7 in Tishomingo County and the highest, 39.3 in Greene County. However, the previously mentioned difficulties with population estimates could be responsible for some of this variation.

Urban-Rural Variation

Children born to urban residents numbered 18,879 and to rural residents 26,840. Comparison of counts by place of occurrence and place of residence showed that although only 41 percent of the births were to urban residents, 73 percent occurred in urban places, indicating that many mothers from rural areas went to towns and cities for delivery. This figure was estimated to be over 8,500 for whites and over 5,500 for nonwhites.

Natural Increase (Excess of Birth Rate over Death Rate)

The continued decline in the birth rate coupled with a rise in the death rate produced the lowest rate of natural increase since 1929. The actual excess of the birth rate over the death rate was 9.2 per 1,000 population in 1968 and less than half the rate of 19.1 only 11 years earlier in 1957. In number of people the natural increase amounted to 21,733 during 1968. The rate of gain among the nonwhite population was more than twice that for the whites, 13.8 in comparison with 6.4, but this was the lowest rate the nonwhites had experienced since 1936. However, the rate for whites was the smallest on record for this race. Deaths outnumbered births among the white population in 13 counties (Amite, Benton, Carroll, Choctaw, Franklin, Holmes, Jefferson, Kemper, Newton, Tallahatchie, Webster, Wilkinson, and Yalobusha), and the two figures were equal in Noxubee County. There was also one county, Tishomingo, where the nonwhites experienced one more death than birth; however, both figures were small. Jackson

Vital Statistics
Mississippi - 1968

County had the highest rate of natural increase noted for whites and Smith County the highest for nonwhites.

Attendant

Information presented in Table B1 shows that 84 percent of the live births to residents of Mississippi took place in hospitals or clinics which provide in-patient care. One percent were delivered by physicians in locations other than a hospital or clinic, and the remaining 15 percent had no medical attendant. The difference by race in care at delivery was quite pronounced. All but a fraction of one percent of the white mothers were delivered in hospitals and clinics but 6,643 or 29 percent of the non-white mothers had no medical care at delivery.

Table B1. Live births and percentage distribution, by attendant and race

Attendant	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>45,719</u>	<u>22,807</u>	<u>22,912</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
Hospital or clinic	38,525	22,704	15,821	84.3	99.5	69.0
Physician, other place	482	34	448	1.0	0.1	2.0
Midwife	6,621	54	6,567	14.5	0.2	28.7
Other	91	15	76	0.2	0.1	0.3

Impressive gains have been made in the last twenty years in obtaining hospital care for delivery with the proportion hospitalized almost doubling (refer to Table B2). The increase from 14 percent in 1948 to 69 percent in 1968 for the nonwhites has been even more spectacular. However, Mississippi still compares unfavorably with the nation in this regard, as the national figure for nonwhite deliveries in hospitals in 1968 was 94 percent.

Table B2. Percentage of live births in hospitals or clinics, by year and race, 1948, 1958, 1964-1968

Year	Total	White	Non-white
1948	42.8	77.9	13.6
1958	70.4	98.6	47.0
1964	76.0	99.4	56.1
1965	77.3	99.4	58.8
1966	79.3	99.4	61.4
1967	82.1	99.5	65.5
1968	84.3	99.5	69.0

Vital Statistics
Mississippi - 1968

Month

The seasonal variation in births is readily apparent in Table B3. These figures show that there were considerably fewer births in the first six months of the year than in the second half and that the lowest level was reached in the spring months of March, April, and May. August was the month with the largest total number and the highest daily average, but December was a close second.

Table B3. Live births and average per day, by month

Month	Births	Average per Day	Month	Births	Average per Day
<u>Total</u>	<u>45,719</u>	<u>125</u>			
January	3,826	123	July	4,045	130
February	3,541	122	August	4,234	136
March	3,548	114	September	3,931	131
April	3,356	112	October	3,905	126
May	3,572	115	November	3,931	131
June	3,668	122	December	4,162	134

Age of Mother

Eighty percent of the babies born in 1968 were born to mothers between the ages of 15 and 29. The greatest concentration was in the ages 20-24, as shown in Table B4, with this group accounting for 36 percent of the total births. Age distributions of white and nonwhite mothers were considerably different; there were relatively fewer nonwhites than whites 20-29 years of age and higher proportions of nonwhites in the younger and older groups. As a result of the excess of nonwhite mothers under age 20, their median age (22.8 years) was less than that of 23.7 years for white mothers. The youngest Mississippi mother in 1968 was only 10 years old there were two others who were only 11 years of age. There were also 13 who were 12 years old and 60 whose age was 13. Although the total number of births continued to decline in 1968, the number to white mother ages 20-29 and to nonwhite mothers under 20 increased. It is likely that at least part of this rise is due to the fact that the unusually large number of babies born in the years immediately after World War II have now reached the age to have children themselves.

Vital Statistics
Mississippi - 1968

Table B4. Live births and percentage distribution, by age of mother and race

Age of Mother	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>45,719</u>	<u>22,807</u>	<u>22,912</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
10-14	311	24	287	0.7	0.1	1.2
15-19	10,858	4,183	6,675	23.7	18.3	29.1
20-24	16,326	9,272	7,054	35.7	40.6	30.8
25-29	9,628	5,647	3,981	21.0	24.8	17.4
30-34	4,868	2,368	2,500	10.6	10.4	10.9
35-39	2,713	1,023	1,690	5.9	4.5	7.4
40-44	907	271	636	2.0	1.2	2.8
45-49	68	10	58	0.1	*	0.2
50-54	2	2	0	*	*	-
Unknown	38	7	31	0.1	*	0.1

* Less than 0.06 percent.

Education of Parents (New information)

Information on education of mother and father was included on Mississippi certificates and those of many other states beginning in 1968. The highest grade of school completed by each parent is considered to be one of the best indicators of socio-economic status that can be obtained and is much simpler to define and tabulate than the information previously collected regarding occupation. Education level of the parents is of particular relevance in the consideration of type of prenatal care sought by and provided for mothers, care the infant receives after birth, and family planning.

The first data available for Mississippi on this subject are summarized in Table B5. Median years of school completed by each parent are shown along with numerical and percentage distributions according to total years completed. Since no information about fathers of illegitimate children is collected, illegitimate births were excluded from the figures on education of the father. On the right hand side of the table, percentages are cumulated downward so that the figure opposite each level of education represents the sum of those who did not advance beyond the highest year in that level. These figures show that 48 percent of the fathers and 54 percent of the mothers had not graduated from high school. Although the number who had not completed even one year of school was extremely small, those who had gone no farther than the sixth grade numbered in the thousands and nearly one-third of both fathers and mothers had not gone beyond the ninth grade. Tabulations by race revealed that nonwhite parents had significantly less education than white parents. Differences in age distribution probably are partly responsible for this; for example, nonwhite mothers under the age of 18 made up 14 percent of the total but only 6 percent of the white mothers were this young. There was a difference of almost three years in the median years of school completed by white and nonwhite fathers and a difference of 2.2 years for the mothers. Almost three-fourths of the nonwhite fathers and mothers had not completed high school.

Vital Statistics
Mississippi - 1968

Table B5. Live births and percentage distribution,
by years of school completed by father and mother and race

Years of School Completed by Father and Mother	Number			Percent*			Cumulative Percent*		
	Total	White	Non- white	Total	White	Non- white	Total	White	Non- white
FATHER									
Median years completed*	12.1	12.5	9.6						
Total legitimate births	37,142	22,121	15,021	100.0	100.0	100.0			
Less than 1 year	129	36	93	0.4	0.2	0.6	0.4	0.2	0.6
1-3 years	905	166	739	2.5	0.8	5.2	2.9	0.9	5.9
4-6 years	3,162	738	2,424	8.8	3.4	17.2	11.7	4.3	23.1
7-9 years	7,012	2,741	4,271	19.6	12.6	30.3	31.3	17.0	53.4
10, 11 years	5,929	3,072	2,857	16.6	14.2	20.2	47.8	31.1	73.6
12 years	10,389	7,795	2,594	29.0	35.9	18.4	76.8	67.0	92.0
1-3 years college	4,491	3,888	603	12.5	17.9	4.3	89.4	84.9	96.3
4 or more years college	3,798	3,274	524	10.6	15.1	3.7	100.0	100.0	100.0
Unknown	1,327	411	916						
MOTHER									
Median years completed*	11.6	12.4	10.2						
Total births	45,719	22,807	22,912	100.0	100.0	100.0			
Less than 1 year	67	15	52	0.1	0.1	0.2	0.1	0.1	0.2
1-3 years	487	71	416	1.1	0.3	1.8	1.2	0.4	2.1
4-6 years	2,842	421	2,421	6.3	1.9	10.8	7.6	2.2	12.8
7-9 years	10,750	3,130	7,620	23.9	13.9	33.9	31.4	16.2	46.7
10, 11 years	10,315	4,326	5,989	22.9	19.2	26.6	54.4	35.4	73.3
12 years	13,557	9,132	4,425	30.1	40.6	19.7	84.5	76.1	93.0
1-3 years college	4,532	3,565	967	10.1	15.9	4.3	94.6	91.9	97.3
4 or more years college	2,417	1,813	604	5.4	8.1	2.7	100.0	100.0	100.0
Unknown	752	334	418						

* Calculations exclude births to parents with education unknown; calculations for fathers also exclude illegitimate births.

Data on education of parents and legitimacy were combined to produce a more complete picture of the socio-economic status into which Mississippi children were born. These figures showed that only 53 percent of the white and 10 percent of the nonwhite children were legitimate and also born into families in which both parents had at least a high school education, two conditions that might be considered as minimal for normal family life and adequate support of a family.

Legitimacy

Information recorded on birth certificates showed that 8,577 or 18.8 percent of the children born to Mississippi mothers in 1968 were illegitimate. For white mothers the count was 686 and the percentage 3.0; corresponding figures for nonwhites were 7,891 and 34.4 percent. Among both races the trend has been upward for many years - since 1957 for the whites and since 1949 for the nonwhites. The percentage for whites, though small, has tripled in that time and that for nonwhites has more than doubled.

Vital Statistics
Mississippi - 1968

The ratio of illegitimate to total births was highest among mothers under 20 years of age which is, of course, the population group with the largest proportion of unmarried women. It was found that 39 percent of the births to mothers in this age group were illegitimate, and that the proportion has risen steadily in recent years. In other groups the proportion ranged from 9 to 15 percent.

Special new tabulations on legitimacy revealed that the mothers of 22 percent of the white illegitimate children and 52 percent of the nonwhite illegitimate children had had previous deliveries. However, it is not known how many of the previous births were also illegitimate.

Total-Birth Order and Average Number of Births per Mother (New information)

Although information regarding number of previous deliveries to the mother had been on birth certificates prior to 1968, it had not been tabulated in Mississippi since 1954 because of doubts concerning the validity of the entries. It was evident in compiling this information for 1968 that the accuracy of reporting is still questionable, and, therefore, the figures will have to be accepted with reservations. However, it is believed that these items contain fewer errors than some of the new items added to the certificate in 1968. Also, the tabulations showed that the proportion of records with these facts not stated was smaller than the proportions of certificates with newly added items not answered.

In 1968, nearly a third of the live births were the first delivery to that mother; the figures in Table B6 show that these first births numbered 14,868. An additional 22 percent were second in total-birth order; thus 54 percent were included among mothers' first two deliveries. (Total-birth order is based on live births plus all fetal deaths regardless of length of gestation.) As birth order increased the number of births steadily declined. However, there were 10,098 births of the fifth or higher orders (22 percent of the total), and 2,272 or 5 percent were of the tenth or higher orders. Twenty mothers, all nonwhite, reported having 20 or more deliveries, with 23 the largest total number. Comparisons by race revealed that relatively more of the white than of the nonwhite births were among the first through third total-birth orders, that the percentages were approximately the same for fourth births, and that proportionately fewer of the whites were of fifth or higher orders (only 11 percent as compared with 33 percent of the nonwhites). Nine percent or 1 out of every 11 nonwhite births was tenth or higher in delivery order.

As shown in Table B6, the median age of mothers at first delivery was 19.9 years. For each birth order nonwhite mothers were younger than white mothers; at first delivery the difference in the median ages was 2.1 years, and for the second through seventh orders there were differences of three to four years. Closer spacing of children among the nonwhites than among the whites is indicated by the fact that there was a difference of 7.4 years between the median ages of white mothers having their first child and those having their fourth delivery in comparison with only 5.5 years for nonwhite mothers.

Vital Statistics
Mississippi - 1968

Table B6. Live births, percentage distribution, and median age of mother, by total-birth order and race

Total-Birth Order*	Number			Percent			Median Age of Mother		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>45,719</u>	<u>22,807</u>	<u>22,912</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>23.3</u>	<u>23.7</u>	<u>22.8</u>
First	14,868	8,543	6,325	32.5	37.4	27.6	19.9	20.8	18.7
Second	9,926	5,961	3,965	21.7	26.1	17.3	22.2	23.5	20.5
Third	6,202	3,589	2,613	13.6	15.7	11.4	24.4	26.0	22.1
Fourth	4,016	2,028	1,988	8.8	8.9	8.7	26.1	28.2	24.2
Fifth	2,623	1,113	1,510	5.7	4.9	6.6	27.4	29.7	25.8
Sixth	1,924	606	1,318	4.2	2.6	5.8	28.4	31.4	27.4
Seventh	1,379	347	1,032	3.0	1.5	4.5	30.0	32.5	29.4
Eighth	1,063	185	878	2.3	0.8	3.8	31.5	33.2	31.3
Ninth	837	117	720	1.8	0.5	3.1	33.4	33.8	33.4
Tenth and over	2,272	159	2,113	5.0	0.7	9.2	36.5	36.6	36.5
Unknown	609	159	450	1.3	0.7	2.0			

* Includes all live births plus all fetal deaths regardless of length of gestation.

Table B7 provides additional information on age of and deliveries to mothers, but shows number of deliveries per mother by age group rather than ages of mothers by birth order. According to these figures, mothers of children born in 1968 had experienced an average of 3.2 deliveries (including the child born in 1968 and all fetal deaths). The average number increased with the age of the mother, of course, reaching 9.2 at ages 40-44 and 11.3 at 45 years and over. The overall average for white mothers was 2.4 and for nonwhites, 4.1. For all age groups 15 years of age and over the average for nonwhites was higher than that for whites, but for mothers 15-19 the difference between the races was not extreme. However, the size of disparity increased steadily with age up to the age of 45 denoting an accumulation of excess fertility among the nonwhites.

Table B7. Average number of total births and living children per mother, (for mothers who had a live birth in 1968), by age of mother and race

Age of Mother	Average Number of Total Births* per Mother			Average Number of Living Children** per Mother		
	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>3.2</u>	<u>2.4</u>	<u>4.1</u>	<u>2.9</u>	<u>2.2</u>	<u>3.7</u>
Under 15	1.0	1.0	1.0	1.0	1.0	1.0
15-19	1.4	1.3	1.6	1.4	1.2	1.5
20-24	2.3	1.9	2.8	2.1	1.7	2.6
25-29	3.7	2.9	5.1	3.4	2.6	4.6
30-34	5.8	4.2	7.6	5.2	3.6	6.7
35-39	7.6	4.9	9.6	6.8	4.3	8.4
40-44	9.2	5.9	11.0	8.2	5.1	9.6
45 and over	11.3	7.3	12.1	10.0	6.6	10.7

* Includes the child born in 1968, all other live births, and all fetal deaths regardless of length of gestation.

** Includes the child born in 1968 and all other children still living at the time of that delivery.

Vital Statistics
Mississippi - 1968

Table B7 also contains data on average number of living children per mother which is an indication of family size. (Living children include the child born in 1968 and all other children still living at the time of that delivery.) These calculations disclosed that the average was 2.2 for white mothers and 3.7 or 68 percent higher for nonwhites.

Last Prior Delivery (New information)

Specific information regarding the dates of last live birth and last fetal death was collected in Mississippi for the first time in 1968. As was true for other new items added in 1968, many certificates had entries that were obviously incorrect or had no entry; and, therefore, the reliability of the figures presented here is somewhat questionable. The two dates were used to determine the outcome of the last prior delivery and calculate the interval between it and the child born in 1968. Researchers have found the health of both the mother and the child to be adversely correlated with spacing of pregnancies at intervals of less than two years and with a history of previous fetal deaths.

Table B8. Live births to mothers who had had previous pregnancies and percentage distribution, by race, months since last delivery, and outcome of last delivery

Race and Months since Last Delivery	Number			Percent		
	Total*	Outcome of Last Delivery		Total*	Outcome of Last Delivery	
		Live Birth	Fetal Death		Live Birth	Fetal Death
ALL RACES						
Median months**	26.0	26.9	18.2			
Total	28,928	26,174	2,754	100.0	100.0	100.0
Under 24	12,153	10,573	1,580	42.0	40.4	57.4
Under 12	1,345	1,013	332	4.6	3.9	12.0
12-23	10,808	9,560	1,248	37.4	36.5	45.3
24-59	11,632	10,930	702	40.2	41.8	25.5
60 and over	3,661	3,505	156	12.6	13.4	5.7
Unknown	1,482	1,166	316	5.1	4.4	11.5
WHITE						
Median months**	33.1	35.1	19.1			
Total	13,630	11,994	1,636	100.0	100.0	100.0
Under 24	4,211	3,307	904	30.9	27.6	55.2
Under 12	404	224	180	3.0	1.9	11.0
12-23	3,807	3,083	724	27.9	25.7	44.2
24-59	6,352	5,908	444	46.6	49.2	27.1
60 and over	2,548	2,440	108	18.7	20.3	6.6
Unknown	519	339	180	3.8	2.8	11.0
NONWHITE						
Median months**	21.8	22.1	16.9			
Total	15,298	14,180	1,118	100.0	100.0	100.0
Under 24	7,942	7,266	676	51.9	51.2	60.5
Under 12	941	789	152	6.2	5.6	13.6
12-23	7,001	6,477	524	45.8	45.7	46.9
24-59	5,280	5,022	258	34.5	35.4	23.1
60 and over	1,113	1,065	48	7.3	7.5	4.3
Unknown	963	827	136	6.3	5.8	12.2

* Excludes those with outcome of last delivery unknown.

** Calculated from ungrouped data.

Vital Statistics
Mississippi - 1968

Information in Table B8 relating to mothers who had had previous pregnancies shows that the median interval since the last delivery was 26.0 months. However, it was noted from more detailed figures than are given in the table that more births occurred during the second year after a preceding delivery than in any other twelve-month period. This was true for both race groups and whether the preceding delivery was a live birth or a fetal death. Almost 10 percent of the births that were from second or later pregnancies had been preceded by a fetal death at the last pregnancy, and the percentage was higher for whites (12) than for nonwhites (7). Further study of spacing in relation to race and outcome of last prior delivery revealed that spacing patterns for white mothers delivering after a fetal death and nonwhite mothers delivering after either a live birth or a fetal death were similar but that white mothers who delivered after a live birth had a quite different pattern and their median interval was 35.1 months in comparison with 19.1, 22.1, and 16.9 months respectively for the other mothers.

Prenatal Care (New information)

The month of pregnancy in which prenatal care began and the number of prenatal visits to a physician or clinic were among the new items added to birth certificates in 1968. Since a number of certificates had no answers in one or both of these items and on some records the entries were obviously wrong, the validity of the data obtained from these items is subject to question. It was reported that 1,202 or 2.6 percent of the mothers had no prenatal care; and, as was anticipated, proportionately more of the nonwhites (4.4 percent) than of the whites (0.8 percent) lacked care. In addition there were also 2,026 mothers whose records did not reveal whether they had any prenatal care.

Analysis of the time prenatal care began showed that 56 percent of the mothers went to a physician or clinic during the first three months of pregnancy, 25 percent waited until the second trimester to begin care, and 10 percent did not receive any care until the third trimester (see Table B9). Although the difference between the proportions of white and nonwhite mothers who received some prenatal care was not large, there was a sharp contrast between the two races in how early in pregnancy care began. Among whites 74 percent had care in the first trimester, but only about half that proportion, 38 percent, of the nonwhites sought medical advice in the first three months.

Information presented in Table B9 relating to number of prenatal visits shows that one-third of the mothers had 6-10 visits and that approximately the same proportion (29 percent) had under 6 visits as had more than 10. The average number per mother who received care was 8.4. Nonwhite mothers averaged only a little over half as many visits (5.9) as the white mothers (10.8), and this correlates with the fact that the nonwhite mothers did not begin prenatal care as early in pregnancy as the white mothers. An additional illustration of the disparity between the two race groups is that only 9 percent of the white mothers had under 6 visits and 48 percent had more than 10, whereas the reverse was true for nonwhites with 48 percent having less than 6 visits and only 9 percent having more than 10.

Vital Statistics
Mississippi - 1968

Table B9. Live births and percentage distribution,
by whether mother received prenatal care,
time during pregnancy care began,
number of visits to physician or clinic, and race

Prenatal Care, Time Care Began, and Number of Visits	Number			Percent		
	Total	White	Non- white	Total	White	Non- white
<u>Total</u>	<u>45,719</u>	<u>22,807</u>	<u>22,912</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
No prenatal care	1,202	188	1,014	2.6	0.8	4.4
<u>Some prenatal care</u>	<u>42,491</u>	<u>21,881</u>	<u>20,610</u>	<u>92.9</u>	<u>95.9</u>	<u>90.0</u>
First trimester	25,415	16,821	8,594	55.6	73.8	37.5
Second trimester	11,566	3,782	7,784	25.3	16.6	34.0
Third trimester	4,532	998	3,534	9.9	4.4	15.4
Unknown month	978	280	698	2.1	1.2	3.0
Average number of visits per mother who received care	8.4	10.8	5.9			
1-5 visits	13,067	2,140	10,927	28.6	9.4	47.7
6-10 visits	15,312	8,148	7,164	33.5	35.7	31.3
11-15 visits	10,871	9,102	1,769	23.8	39.9	7.7
16 visits or more	2,236	1,922	314	4.9	8.4	1.4
Unknown number	1,005	569	436	2.2	2.5	1.9
Unknown whether any prenatal care	2,026	738	1,288	4.4	3.2	5.6

Complications during Pregnancy and Labor (New information)

The year 1968 was the first in which an attempt was made to have medical information about complications during pregnancy and labor recorded on birth certificates. However, the results were disappointing. For 13 percent or 1 out of every 8 births it was unknown whether the mother had any complications, and much of the information which was provided on the remainder of the records was entered in such a way that its proper classification was extremely doubtful. It is likely that in many of the instances in which there was no information given no complications occurred, but this could not be taken for granted.

Complications were reported on 4,966 records or about 11 percent of the total (see Table B10), but this should be considered as a minimum figure. In addition to the previously mentioned difficulties with the records, there were numerous conditions such as the common cold, pleurisy, virus, itch, blackout, emotional upset, back pain, dehydration, numbness, etc. which were not counted because they were considered insignificant or too vague for classification. However, since no clear definition of what is to be considered as a complication has been established, deciding what to accept and what to reject was a serious problem which was never satisfactorily solved. Finally, on numerous records it was indicated that a cesarean section or other procedure was done, but the complication which necessitated it was not recorded. The proportion of white mothers

Vital Statistics
Mississippi - 1968

with at least one complication was slightly higher than that for nonwhite mothers, 11.7 percent in comparison with 10.0 percent. This difference, however, may be merely a reflection of a difference in number of conditions diagnosed as the white mothers received more medical care during pregnancy and labor.

Table B10. Live births and percentage distribution,
by whether mother had complications during pregnancy and labor,
category of complication, and race

Complications	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>45,719</u>	<u>22,807</u>	<u>22,912</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
Without complications	35,011	17,282	17,729	76.6	75.8	77.4
With complications	4,966	2,672	2,294	10.9	11.7	10.0
Related to pregnancy	(2,338)	(1,124)	(1,214)	(5.1)	(4.9)	(5.3)
During but not related to pregnancy	(1,171)	(614)	(557)	(2.6)	(2.7)	(2.4)
Of labor	(2,258)	(1,303)	(955)	(4.9)	(5.7)	(4.2)
Unknown whether any complications	5,742	2,853	2,889	12.6	12.5	12.6

Table B10 also provides information on number and percentage of records showing complications in each of the three major categories. (These figures add up to more than the total number with complications as some mothers had complications in more than one category.) About 1 out of every 20 or 5 percent of the mothers were known to have had complications directly related to this pregnancy and the same proportion had complications during labor. In addition, 2.6 percent had complications during pregnancy that did not arise from the pregnancy itself.

Data presented in Table B11 pertain to the number of specific complications listed rather than the number of mothers with complications. Some of these complications such as hemorrhages and premature rupture of membranes can occur either during pregnancy or at the time of labor. Others such as hypertension and anemia can be either related to pregnancy or not related to pregnancy depending upon whether they arose during the pregnancy or existed as a chronic condition prior to pregnancy. Determination of the proper category was a matter of guesswork in many such cases. Sections of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States were used for specific classification of complications within the three major categories: codes 630-639 for those related to pregnancy, codes 760 and 761 for those not related to pregnancy, and codes 651-662 for those of labor. However, the index for the Eighth Revision was not available until records for eleven months of the year had already been processed, and it was found after the index was received that a number of conditions had been incorrectly classified during the first eleven months.

All of the complications with specific codes which appeared on as many as 100 certificates are shown in Table B11. Toxemias of pregnancy were listed more frequently than any other complication in any of the three major categories; they appeared on 1,179 records and accounted for 19 percent of the total reported. This was the leading complication only among nonwhite mothers, however, as malpresentation of the fetus at the time of delivery was listed more frequently than any other among white mothers and ranked

Vital Statistics
Mississippi - 1968

second in the total group with 773 cases reported. The third-ranking condition was hemorrhage with a total of 652 either during pregnancy or during labor, and this also was listed more often for whites than for nonwhites.

Table B11. Complications of pregnancy and labor
reported on live birth certificates, by type and race

Type of Complication and International Classification Number*	Total	White	Non- white
<u>Total complications reported</u>	<u>6,274</u>	<u>3,337</u>	<u>2,937</u>
<u>Total related to pregnancy (630-639)</u>	<u>2,499</u>	<u>1,193</u>	<u>1,306</u>
Toxemias (636-639)	1,179	402	777
Hemorrhage (632)	382	249	133
Anemia (633)	203	90	113
Urinary infections (635)	183	117	66
Premature rupture of membrane without delivery (634.4)	136	75	61
All others	416	260	156
<u>Total during but not related to pregnancy (760, 761)</u>	<u>1,242</u>	<u>650</u>	<u>592</u>
Chronic (and unspecified) circulatory diseases (760.0-760.3)	282	81	201
Chronic (and unspecified) genitourinary diseases (760.4, 760.5)	134	88	46
Diabetes mellitus (761.1)	103	47	56
All others	723	434	289
<u>Total of labor (651-662)</u>	<u>2,533</u>	<u>1,494</u>	<u>1,039</u>
Malpresentation of fetus (656)	773	468	305
Obstetrical trauma (658-660)	363	244	119
Fetopelvic disproportion (655)	280	201	79
Hemorrhage (651-653)	270	150	120
Premature rupture of membranes (661.0)	142	63	79
Umbilical cord complication (661.3)	111	63	48
All others	594	305	289

* From the Eighth Revision International Classification of Diseases, Adapted for Use in the United States.

Length of Gestation and Birth Weight

For a number of years information about length of gestation and birth weight has been collected and tabulated in Mississippi. The item on birth weight was added to the certificate in 1949 and has remained unchanged since that time. Information relating to length of gestation has been collected since 1929, but the form in which it has been requested has changed several times. In 1949 this item was revised to show number of weeks of gestation. However, the method of determining the number of weeks was left to the attendant, and it was apparent that in many instances no attempt was made to calculate the correct period. The length of gestation for the vast majority of births was reported as 40 weeks. Among the remainder there was concentration in the even-numbered weeks.

Since the method of collecting gestation information used since 1949 yielded unsatisfactory results, a different method was initiated in Mississippi and most other states

Vital Statistics
Mississippi - 1968

with the certificate revision of 1968. The attendant was requested to give the exact date of the mother's last menstrual period (LMP) prior to delivery so that the number of completed weeks of gestation could be accurately calculated by determining the interval between the LMP and the date of delivery. Because the LMP is a date, often unnoted at the time, that must be recalled by the mother at some later time (in many instances several months later), the accuracy of this item is probably more questionable than that of any other on the birth certificate. Despite this difficulty, however, the figures obtained by this method are superior to those previously collected.

Table B12 reveals the sharp contrast between the two methods. In 1967, the last year in which the attendant was asked to state the number of weeks, nearly 86 percent of the births were recorded with a gestation period of 40 weeks, whereas in 1968, from calculations using the LMP, only 18 percent were found to be at 40 weeks. In addition, there were substantial differences in the proportions of premature deliveries (5 percent under 37 weeks in 1967 and 17 percent in 1968) and of deliveries after more than 40 weeks of gestation (2 percent in 1967 and 21 percent in 1968). Even greater differences were noted in the incidence of prematurity among nonwhites and pregnancies of more than 40 weeks among the whites.

Table B12. Percentage distribution of live births,
by completed weeks of gestation and race, 1967 and 1968

Weeks of Gestation	1967			1968		
	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
<u>Under 37</u>	<u>5.2</u>	<u>4.5</u>	<u>5.8</u>	<u>16.6</u>	<u>10.8</u>	<u>22.4</u>
Under 20	*	*	*	0.1	0.1	0.2
20-27	0.6	0.4	0.7	1.0	0.5	1.4
28-31	0.9	0.6	1.2	2.0	1.2	2.8
32-35	1.7	1.7	1.7	8.4	5.2	11.6
36	1.9	1.7	2.1	5.0	3.8	6.3
37-39	7.1	7.8	6.5	38.7	38.1	39.2
40	85.5	84.3	86.6	18.1	21.8	14.4
<u>Over 40</u>	<u>2.1</u>	<u>3.1</u>	<u>1.0</u>	<u>21.4</u>	<u>26.0</u>	<u>16.8</u>
41-44	2.0	3.1	0.9	19.7	24.2	15.2
45-52	*	*	0.1	1.7	1.9	1.6
Unknown	0.1	0.2	0.1	5.2	3.2	7.2

* Less than 0.06 percent.

The major reason for collecting data on gestation period and birth weight is to study the problem of babies who are not mature when born. The term "premature" is frequently used to indicate those for whom a lack of maturity is indicated by a gestation period of less than 37 weeks, and the term "immature" is used to indicate a birth weight of 5 1/2 pounds (2,500 grams) or less. The two conditions are, of course, correlated; but, as the figures in Table B13 make clear, the correlation was far from perfect. A total of 2,428 (5.3 percent) of the infants born alive in 1968 met both criteria for lack of maturity. An additional 2,065 were classified as immature but not premature, bringing the total proportion that were immature up to 9.8 percent. There were 5,157 others who were premature but not immature, raising the incidence

XVIII

Vital Statistics
Mississippi - 1968

of prematurity to 16.6 percent. The total number of babies affected by either a low birth weight and/or a significantly shortened gestation period was 9,650 or 21.1 percent. Regardless of what measurement of lack of maturity was used, the percentage was higher for nonwhites than for whites. The difference between the races was greater when measured by gestation period, however, than when measured by weight.

Table B13. Live births and percentage distribution, by maturity and race

Maturity*	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>45,719</u>	<u>22,807</u>	<u>22,912</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
Immature only**	2,065	735	1,330	4.5	3.2	5.8
Premature only**	5,157	1,547	3,610	11.3	6.8	15.8
Both immature and premature	2,428	913	1,515	5.3	4.0	6.6
Total immature	4,493	1,648	2,845	9.8	7.2	12.4
Total premature	7,585	2,460	5,125	16.6	10.8	22.4
Total immature and/or premature	9,650	3,195	6,455	21.1	14.0	28.2
Neither immature nor premature**	36,069	19,612	16,457	78.9	86.0	71.8

* Immature - birth weight 5 1/2 pounds (2,500 grams) or less; premature - gestation less than 37 weeks.

** Births classified as immature only or as premature only include some for which the other measurement was unknown. Those classified as neither immature nor premature include some for which one or both measurements were unknown.

Sex

Counts by sex showed 23,428 males and 22,291 females and these figures yielded a sex ratio of 1,051 males per 1,000 females; comparison of the ratios for the past 25 years revealed that they have varied from 1,024 to 1,052. In all years for which sex ratios have been calculated that for the whites was higher than that for nonwhites, and in 1968 the figures were 1,077 and 1,026 respectively.

Plurality (See Plurality in Fetal Death section also)

Live births from plural deliveries numbered 1,127 and comprised 2.5 percent of the total. Figures by race were 2.0 percent for whites and 3.0 percent for nonwhites. Tabulation by age of mother revealed that the proportion of plural births was the same for the two groups under age 20, increased steadily from ages 15-19 through ages 35-39, and then declined. The highest percentage was 5.0 among mothers 35-39 years old and the lowest, 1.6 for the groups under age 20. At all ages except 40-44 the proportion was greater among nonwhite mothers than among whites with the largest difference in the peak interval (ages 35-39) where the figure for nonwhites was more than twice as high as that for whites.

Vital Statistics
Mississippi - 1968

Further analysis showed extremely high proportions of babies with low birth weight and of deliveries before term among the plural deliveries. Seven times as many of the plural births as of the single births weighed 5 1/2 pounds or less (56 percent in comparison with 8 percent), and about three times as many had a gestation period of less than 37 weeks (46 percent in comparison with 15 percent). Differences between single and plural births in birth weight and in weeks of gestation were greater for whites than for nonwhites.

Birth Injuries and Congenital Malformations (New information)

A majority of the states had sections for birth injuries and congenital malformations on their birth certificates in 1968. Many already had one or both items on their records prior to 1968 and others, including Mississippi, added them in that year as a result of the incorporation of these two items into the standard certificate which serves as a model. Most of the remaining states had only a section for congenital malformations. As was true for the new items on complications during pregnancy and labor, there were large numbers of certificates with no entry in the sections relating to the condition of the baby - 13 percent or about 1 out of every 8 (see Table B14). Although it is likely that in many of these instances there was nothing wrong with the child, this was not known to be so and could not be assumed. Malformations and/or injuries were reported on 522 or 1.1 percent of the certificates, with 7 children having both an injury and a malformation; and on nearly 86 percent it was stated that there were none. Slightly more of the whites than of the nonwhites had malformations or injuries listed, but this could be the result of better diagnosis and reporting for white children since over 99 percent were born in hospitals or clinics.

Table B14. Live births and percentage distribution,
by whether child had any birth injuries or congenital malformations,
whether injury or malformation, and race

Birth Injuries and Congenital Malformations	Number			Percent		
	Total	White	Non- white	Total	White	Non- white
<u>Total</u>	<u>45,719</u>	<u>22,807</u>	<u>22,912</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
Without injury or malfor- mation	39,121	19,669	19,452	85.6	86.2	84.9
With injury and/or mal- formation	522	297	225	1.1	1.3	1.0
Injury	(89)	(57)	(32)	(0.2)	(0.2)	(0.1)
Malformation	(440)	(244)	(196)	(1.0)	(1.1)	(0.8)
? Unknown whether injury or malformation	6,076	2,841	3,235	13.3	12.4	14.1

Birth injuries were mentioned on only 89 records which is 0.2 percent of the total. It is probable that there were others which were not recorded because of the fear of blame or even legal action against the attendant, but the degree of under-reporting is not known. No spinal cord injuries and only 10 brain injuries were listed; 36 injuries to bone and nerve were reported and 42 in all other categories. One case of multiple injuries was recorded which involved the brain and another site.

Vital Statistics
Mississippi - 1968

Although more congenital malformations than birth injuries were reported, under-reporting is also considered to be a problem in analyzing the occurrence of congenital malformations because of several reasons. Some types of malformations, especially of internal organs, are not apparent at the time of birth and others, although recognizable by trained physicians, might not be noted by midwives, who delivered 29 percent of the nonwhite babies in 1968. Undoubtedly there were also instances where a diagnosed malformation was not recorded for fear of stigmatizing a child. Table B15 shows that a total of 518 were reported. (The actual total reported probably was slightly higher as there was room on the punch card used for tabulating purposes for only three per child and a few children may have had more than three.) However, only 440 children were affected as 54 had two malformations each and 12 had more than two each. Among body sites the musculoskeletal system ranked first, partly because of ease of recognition; 94 of the 207 reported were polydactyly (extra fingers and toes) and 43 were club foot. The digestive system ranked second; and, again, this was probably due at least in part to the fact that it included a sizeable number of an easily recognized condition - 56 cases of cleft palate and/or cleft lip among the total of 100 for this system. Other systems for which 50 or more malformations were reported were the nervous system (55 cases) and genital organs (54 cases).

Table B15. Congenital malformations reported on live birth certificates, by system or organs affected and race

System or Organs Affected and International Classification Number*	Total	White	Non-white
Total malformations reported	518	300	218
Musculoskeletal system (754-756)	207	91	116
Digestive system (749-751)	100	63	37
Nervous system (740-743)	55	43	12
Genital organs (752)	54	37	17
All others	102	66	36

* From the Eighth Revision International Classification of Diseases, Adapted for Use in the United States.

FETAL DEATHS (See Table 4)

Reporting and Reliability

In Mississippi the State Board of Health has required the registration of all fetal deaths, regardless of the length of gestation, since 1952. (The reporting requirement prior to that year was for stillbirths of 20 or more weeks of gestation.) However, it is obvious from the results obtained that registration of early fetal deaths as well as those that occur after 20 or more weeks gestation is far less complete than that for live births or deaths. The degree of underreporting is not known but is thought to be significant; and, therefore, the information presented herein regarding fetal deaths may not be reliable.

Trend and Race

A total of 2,373 such events were recorded in Mississippi in 1968, and the count

Vital Statistics
Mississippi - 1968

for residents of the state was 2,322. Although these figures were only slightly larger than those for 1966 and 1967 and considerably less than those for the early 1960's, the fetal death ratio of 50.8 per 1,000 live births was the highest recorded in the seventeen years of all-inclusive registration. Only among the whites was an increase noted, however; both the number and ratio for nonwhites went down. These opposite changes brought the ratios for the two groups closer together; but, as usual, that for the nonwhites was higher, 52.5 in comparison with 49.1 for whites.

County

County ratios showed extreme variation. For the white population they ranged from zero in three counties to 142.8 in Issaquena County, and for nonwhites they went from zero in four counties to 285.7 in Tishomingo. Differences such as these, even though partly due to variations in the age distribution of the population and other factors, are probably in large measure the result of differences in reporting. If it is assumed that the high ratios in counties where the number of live births and fetal deaths is large enough to provide a reliable calculation represent complete reporting, it can be estimated that at least ten percent of all deliveries terminate in fetal deaths; special studies done in other places have estimated the true incidence in excess of twenty percent.

Attendant

Nearly 90 percent of the fetal deaths to Mississippi residents in 1968 were delivered in hospitals or clinics (refer to Table C1). Physicians attended 2 percent in homes and other places, and for 192 or 8 percent there was only a midwife or other nonmedical person in attendance. Even though just 2 white mothers did not receive medical attention during delivery, 190 nonwhite mothers (16 percent of the total) did not have a physician in attendance. Relatively more of the nonwhite mothers, however, were hospitalized at delivery of a fetal death than at delivery of a live birth (81 percent vs. 69 percent), but this is to be expected when it is considered that prematurity and complications are commonly associated with death of the fetus.

Table C1. Fetal deaths and percentage distribution, by attendant and race

Attendant	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>2,322</u>	<u>1,119</u>	<u>1,203</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
Hospital or clinic	2,078	1,101	977	89.5	98.4	81.2
Physician, other place	52	16	36	2.2	1.4	3.0
Midwife	183	2	181	7.9	0.2	15.0
Other	9	0	9	0.4	-	0.7

Advances in securing hospital care when a fetal death occurs are evident from the data in Table C2. Significant gains were made by both race groups in the last ten years and the gap between the two narrowed considerably (from 32 percentage points to 17).

Vital Statistics
Mississippi - 1968

Table C2. Percentage of fetal deaths delivered in hospitals or clinics, by year and race, 1958, 1964-1968

Year	Total	White	Non-white
1958	69.0	88.8	56.8
1964	79.7	92.0	70.1
1965	79.4	95.8	67.5
1966	83.7	97.5	71.9
1967	86.2	97.7	76.8
1968	89.5	98.4	81.2

Age of Mother

The fetal death ratio (number of fetal deaths per 1,000 live births) was lowest for mothers 15-19 years of age and rose steadily with increasing age as shown in Table C3. Mothers 40-44 years old had the highest ratio calculated which was 120.2, or more than three times greater than the low figure for those of ages 15-19. Among the very young mothers (ages 10-14) the ratio was higher than for any others under the age of 30 but less than for any 30 and over. Oddly enough, although most of the ratios for nonwhites were higher than those for whites, the lowest ratio of all was that for nonwhite mothers 15-19 years old.

Table C3. Fetal deaths and fetal death ratios, by age of mother and race

Age of Mother	Number			Ratio*		
	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>2,322</u>	<u>1,119</u>	<u>1,203</u>	<u>50.8</u>	<u>49.1</u>	<u>52.5</u>
10-14	19	1	18	61.1	41.7	62.7
15-19	413	180	233	38.0	43.0	34.9
20-24	711	394	317	43.6	42.5	44.9
25-29	454	249	205	47.2	44.1	51.5
30-34	357	166	191	73.3	70.1	76.4
35-39	222	76	146	81.8	74.3	86.4
40-44	109	34	75	120.2	125.5	117.9
45-49	17	7	10	#	#	#
Unknown	20	12	8			

* Fetal deaths per 1,000 live births to mothers in specific age-race group.

Ratio not calculated on less than 100 live births.

Education of Parents (New information)

An item on the highest grade of school completed by each parent was added to fetal death certificates in 1968 and was identical to the item added to live birth certificates. As was true in general in comparing information on live birth and fetal death

Vital Statistics
Mississippi - 1968

certificates, reporting was far less complete for fetal deaths; 21 percent of the records for legitimate children did not have the father's education filled in and 18 percent of the total lacked the number of years of school the mother had completed. However, using the records which did have the information entered, it was found that educational levels of the parents who had fetal deaths were about the same as those who had live births. Forty-five percent of the fathers and 53 percent of the mothers had not finished high school, and nearly a third of both fathers and mothers had not gone beyond the ninth grade. Nonwhite parents had received significantly less education than white parents - fathers 3.5 years less and mothers 2.3 years less.

Legitimacy

Almost one-fifth of the fetal deaths were reported as illegitimate. The actual percentage, 19.7, was a little higher than that of 18.8 for live births, but data by race show that this was true only for the whites. A far greater proportion of nonwhite than of white fetal deaths were illegitimate, 32.8 percent in comparison with 5.5 percent, but both figures were lower than in the previous year.

Total-Birth Order (New information)

Fetal deaths in Mississippi had never been classified by number of previous deliveries to the mother before 1968 although there had been items on the certificates concerning this subject. This information was missing from 10 percent of the 1968 records; and even though there is some doubt concerning the accuracy of that which was entered, it was used to determine total-birth order which includes live births plus all fetal deaths regardless of length of gestation. First deliveries made up 24.5 percent of the total. This percentage as well as those for second and third total-birth orders were somewhat lower than the corresponding proportions among live births; the first three orders made up 68 percent of the live birth total but only 52 percent of the fetal death total. The largest number of previous deliveries reported for a mother who had a fetal death in 1968 was 21, which was only one less than the largest number reported for a mother of a live birth.

Fetal death ratios generally increased with total-birth order, ranging from 38.3 per 1,000 live births for those of the first order to 86.7 for those of tenth or higher orders (see Table C4). This correlates with the fact that fetal death ratios rise as the age of the mother increases. It was also noted that for all orders except the eighth and ninth the ratios for whites were higher than those for nonwhites.

Vital Statistics
Mississippi - 1968

Table C4. Fetal death ratios*, by total-birth order and race

Total-Birth Order**	Total	White	Non- white
First	38.3	38.6	37.8
Second	39.7	40.4	38.6
Third	40.6	41.8	39.0
Fourth	53.3	59.2	47.3
Fifth	48.0	50.3	46.4
Sixth	65.5	72.6	62.2
Seventh	68.9	86.4	63.0
Eighth	53.6	43.2	55.8
Ninth	75.3	59.8	77.8
Tenth and over	86.7	125.8	83.8

* Fetal deaths per 1,000 live births of specified total-birth order and race.

** Includes all live births plus all fetal deaths regardless of length of gestation.

Last Prior Delivery (New information)

A new section relating to the last live birth and the last fetal death was added to fetal death certificates in 1968 as it was to live birth certificates in order to obtain data on outcome of the last prior delivery and the interval between it and the current delivery. However, the information was either missing or inadequate to determine the outcome for 16 percent and the interval for 22 percent of the fetal deaths. Also, the accuracy of data which could be used is subject to some doubt.

Twenty-one percent of the fetal deaths (but only 10 percent of the live births) that were from second or later pregnancies had been preceded by a fetal death at the time of the last pregnancy. In addition, it was found that the fetal death ratio among mothers whose last pregnancy also terminated in a fetal death was 103.1 per 1,000 live births in contrast to a ratio of only 40.4 for those whose last pregnancy terminated in a live birth. Both calculations are indicative of the greater likelihood of an unfavorable pregnancy outcome when there is a history of previous fetal deaths.

The median number of months between a fetal death and the last prior delivery was close to the same as that between a live birth and the last prior delivery, 24.5 and 26.0 months respectively. As the figures in Table C5 show, the only marked difference in the median interval prior to a live birth and that prior to a fetal death was for whites whose last delivery was a fetal death; among whites the median interval was 5 months shorter for a fetal death preceded by fetal death than for a live birth preceded by a fetal death. It was also noted that, among both race groups and regardless of whether a live birth or a fetal death occurred, the median length of time since the last delivery was less if the last delivery was a fetal death than if it was a live birth and that the difference was far greater for whites than for nonwhites.

Vital Statistics
Mississippi - 1968

Table C5. Median number of months since last delivery to mothers who had had previous pregnancies, by race, outcome of last delivery, and outcome of delivery in 1968

Race and Outcome of Last Delivery	Outcome of Delivery in 1968	
	Live Birth	Fetal Death
<u>Total*</u>	<u>26.0</u>	<u>24.5</u>
White: live birth	35.1	35.7
fetal death	19.1	13.8
Nonwhite: live birth	22.1	22.0
fetal death	16.9	15.3

* Excludes those with outcome of last delivery unknown.

Prenatal Care (New information)

Information on the month of pregnancy prenatal care began and the number of prenatal visits was included on fetal death as well as live birth certificates for the first time in 1968. Reporting of these facts on fetal death certificates was especially poor. For 21 percent of the records it could not be determined whether any care was received.

Sixteen percent of the mothers were reported to have had no prenatal care at all (see Table C6); and, if the facts were known for the large number with no information on the record, the proportion would probably be much higher. In either case, it compares unfavorably with the experience of mothers of live births of whom all but 3 percent received care. It was reported for fetal deaths that 62 percent did receive care, 45 percent began care in the first trimester, only 3.5 percent started care as late as the third trimester, and the average number of visits for those who received care was 4.5 in comparison with 8.4 for mothers of live births. However, since many fetal deaths occurred in the first and second trimesters, there was not as long a period of time in which the mothers could have received care and the figures in the preceding sentence are not really comparable to the same type of data on live births. There was a relatively small difference in the average number of visits for white and nonwhite mothers (4.8 and 4.2 respectively), but the nonwhite mothers could have been expected to have averaged more visits than the white mothers as proportionately more of their reported fetal deaths occurred in the third trimester and fewer in the first trimester.

Vital Statistics
Mississippi - 1968

Table C6. Percentage distribution of live births and fetal deaths by whether mother received prenatal care and race, and average number of visits per mother who received care, by race

Prenatal Care and Visits	Live Births			Fetal Deaths		
	Total	White	Non- white	Total	White	Non- white
<u>Total</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
No care	2.6	0.8	4.4	16.4	14.9	17.8
Some care	92.9	95.9	90.0	62.4	64.6	60.3
Unknown whether any care	4.4	3.2	5.6	21.2	20.5	21.9
Average number of visits per mother who received care	8.4	10.8	5.9	4.5	4.8	4.2

Length of Gestation and Birth Weight

The collection and tabulation of information on length of gestation and birth weight for fetal deaths is identical to that for live births (see Live Birth section of Synopsis for details). No birth weight was given for 57 percent of the fetal deaths in 1968. Therefore, no data on that topic are presented here although it might be possible to extract some meaningful material from records of fetal deaths of 28 or more weeks gestation since reporting of weight for those occurring in this period was far better than for early fetal deaths. The change in the method of obtaining information on gestation period in 1968 produced better results than before even though the new method was subject to the same deficiencies as were noted in the discussion on live births, including a larger proportion with the number of weeks unknown, 15 percent in comparison with only 4 percent the year before.

Comparative figures on length of gestation as determined in 1967 and from the revised certificate of 1968 are presented in Table C7. The total proportions of early fetal deaths (under 20 completed weeks), those of 20-36 weeks gestation, and at or near term deliveries (37 weeks or more) did not differ greatly from 1967 to 1968, but when the early and term deliveries were separated into more detailed periods it was found that there were changes in opposite directions within these groups. Among early fetal deaths relatively fewer of less than 10 weeks gestation were reported in 1968 and relatively more (especially among whites) with gestations of 10-19 weeks. For term deliveries there was no longer a peak at exactly 40 weeks in 1968 and there were more in the weeks immediately prior to and after the fortieth. This change was quite pronounced for nonwhites; in fact, the drop in the proportion at exactly 40 weeks was so large that it resulted in a decline of some significance in the total percentage of nonwhite fetal deaths at term.

Vital Statistics
Mississippi - 1968

Table C7. Percentage distribution of fetal deaths
by completed weeks of gestation and race, 1967 and 1968

Weeks of Gestation	1967			1968		
	Total	White	Non- white	Total	White	Non- white
<u>Total</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
<u>Under 20</u>	<u>47.0</u>	<u>62.4</u>	<u>34.4</u>	<u>43.0</u>	<u>55.4</u>	<u>31.5</u>
<u>Under 10</u>	<u>20.6</u>	<u>31.7</u>	<u>11.6</u>	<u>11.0</u>	<u>14.6</u>	<u>7.7</u>
10-19	26.4	30.6	22.8	32.0	40.8	23.8
<u>20-36</u>	<u>30.2</u>	<u>22.1</u>	<u>36.8</u>	<u>28.3</u>	<u>22.1</u>	<u>34.1</u>
20-27	15.0	11.9	17.5	13.4	11.8	14.9
28-36	15.2	10.2	19.4	14.9	10.3	19.2
<u>37 and over</u>	<u>18.8</u>	<u>11.2</u>	<u>25.0</u>	<u>14.0</u>	<u>10.0</u>	<u>17.8</u>
37-39	2.4	2.7	2.1	7.0	4.7	9.1
40	15.9	8.0	22.5	2.7	1.6	3.7
Over 40	0.4	0.5	0.4	4.3	3.7	4.9
<u>Unknown</u>	<u>4.0</u>	<u>4.4</u>	<u>3.7</u>	<u>14.6</u>	<u>12.5</u>	<u>16.6</u>

Sex

A large portion of fetal deaths occur so early in pregnancy that the sex cannot be determined; in 1968, 44 percent of the records did not show sex of the fetus. Those on which the sex was recorded were used to calculate a sex ratio which was 1,496 males per 1,000 females and the highest ever reported for Mississippi. The fetal death sex ratio is always much higher than that for live births (1,051 in 1968). It may be that if the sex could be determined for all fetal deaths a more even distribution would be found, but there is also the possibility that there actually is excessive mortality among the males during pregnancy and labor.

Plurality

In 1968, 3.6 percent of the fetal deaths were from plural deliveries, and the rate of occurrence was greater among nonwhites than among whites, 4.8 and 2.3 percent respectively. As usual, the frequency among fetal deaths was higher than that among live births (only 2.5 percent).

Further analysis of plurality among both live births and fetal deaths revealed that 1 out of every 79 pregnancies resulted in a set of either twins or triplets in 1968. The ratio for whites was only 1 in every 100, but that for nonwhites was 1 in every 65. Twin sets numbered 595 and there were 7 sets of triplets. All the children were born alive in 89.5 percent of the plural deliveries, 7 percent had both live and dead members, and all were born dead in 3.5 percent of the cases. When fetal mortality from single deliveries was compared with the information on the plural sets it was found that 6.9 percent of the fetuses from these plural sets was born dead in contrast to only 4.8 percent of those from single births and that the difference was more pronounced among nonwhites than among whites.

Vital Statistics
Mississippi - 1968

Cause of Fetal Death, Complications of Pregnancy and Labor, Birth Injuries, and Congenital Malformations

Although fetal death certificates contained a section for cause prior to 1968, the information, if any, entered in this section was of too poor quality to classify. This was also true of the records filed in 1968. However, an attempt was made to use the data entered in the new sections on complications of pregnancy and labor, birth injuries, and congenital malformations which were added when the certificate was revised in 1968. It was planned that a comparative study would be made comparing the occurrence of these events when the outcome was a live birth and when it was a fetal death, but the recorded information was so poor that it was not possible to code it. Of particular concern was the fact that entries in the cause section and the new sections were frequently inconsistent and even conflicting.

DEATHS (See Tables 5-8, 12-16)

Trend

The death total of 23,986 Mississippi residents for 1968 was the largest on record since 1937 and represents an increase of 1,377 over the previous year. An increase was also registered in the death rate which went up from 9.6 to 10.2 per 1,000 population. The figure for 1968 was the same as the rate for 1963 but higher than those for any of the years in between. However, as mentioned in the section on live births, population estimates for the latter part of a decade are especially subject to inaccuracies because of the length of time since the last decennial census, and rates based on them would be affected accordingly.

Race

Counts for both whites and nonwhites rose in 1968; in fact, the death toll of 13,468 for whites was the highest ever recorded in this state. The rates for the two groups were 9.3 (whites) and 11.7 (nonwhites), and the difference between them was greater than in 1967 because that for nonwhites went up more than that for whites.

County and Urban-Rural Variation

Death rates varied considerably by county. For the white population the range was from 5.1 in Jackson and Tunica Counties to a high in Wilkinson County of 15.6, which was over three times the lowest rate. Rates for nonwhites ranged from 6.6 in Rankin County to 18.0 in Greene County. The fact that many rural residents go into towns and cities for health care is evident from death figures, as it was from birth data; although only 40 percent of the deaths occurred among urban residents, 62 percent occurred in urban locations.

Month

Data in Table D1 show that far more deaths occurred in January than in any other month and that the daily average was also considerably higher. The smallest number and lowest daily average were recorded in July, but figures for several other months were

Vital Statistics
Mississippi - 1968

nearly as low. The seasonal picture revealed is one of excess mortality during the part of the year when the coldest weather occurs.

Table D1. Deaths and average per day, by month

Month	Deaths	Average per Day	Month	Deaths	Average per Day
<u>Total</u>	<u>23,986</u>	<u>66</u>			
January	2,727	88	July	1,822	59
February	1,998	69	August	1,879	61
March	1,947	63	September	1,827	61
April	1,829	61	October	1,956	63
May	1,875	60	November	1,941	65
June	1,933	64	December	2,252	73

Age at Death

There has been an upward, though not unbroken, trend in the median age at death in recent years. The median age in 1968 was 68.9 which is exactly two years more than it was in 1961; the change between 1967 and 1968 alone amounted to 0.8 of a year. Medians of 70.7 and 66.6 years were calculated for the whites and nonwhites respectively. In both groups the trend has been upward and the gap between the two has narrowed. As may be seen in Table D2, the number of deaths among persons 75-84 years of age was larger than the number for any other ten-year age group with the count for persons 65-74 years old not far behind. The smallest total for a ten-year age span was that for children 5-14 years old. Differences between the age distributions of whites and nonwhites were minor except among the very young and the very old. Of special significance is the fact that only 3.8 percent of the whites who died were under the age of one but the proportion for the nonwhites was 10.4 percent.

Table D2. Deaths and percentage distribution, by age and race

Age	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>23,986</u>	<u>13,468</u>	<u>10,518</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
Under 1	1,613	517	1,096	6.7	3.8	10.4
1-4	264	89	175	1.1	0.7	1.7
5-9	148	69	79	0.6	0.5	0.8
10-14	138	58	80	0.6	0.4	0.8
15-24	621	325	296	2.6	2.4	2.8
25-34	555	245	310	2.3	1.8	2.9
35-44	895	444	451	3.7	3.3	4.3
45-54	1,917	1,125	792	8.0	8.4	7.5
55-64	3,732	2,131	1,601	15.6	15.8	15.2
65-74	5,383	3,030	2,353	22.4	22.5	22.4
75-84	5,560	3,528	2,032	23.2	26.2	19.3
85 and over	3,151	1,906	1,245	13.1	14.2	11.8
Unknown	9	1	8	*	*	*

* Less than 0.06 percent.

Vital Statistics
Mississippi - 1968

Leading Causes of Death

Table D3 lists the ten leading causes of death and shows that 79 percent of the total deaths were attributed to this group of causes. Relatively more of the whites than of the nonwhites died from these causes, but this was principally due to the differences in the proportions assigned to the two top-ranking causes. Heart diseases and malignant neoplasms together accounted for 53 percent of the deaths among the white population but only 39 percent of those among nonwhites. The first eight causes are the same ones which were listed in the previous year, although the order of those in the fifth through the eighth places varied. Homicide, in tenth place in 1968, had appeared on the list in two other years in the 1960's, but the combination of respiratory diseases which ranked ninth is a new grouping being used since the revised system of classifying causes of death went into effect in 1968. In comparing the ranking of certain causes in 1968 with that for prior years, it should be realized that some of the variation probably resulted from the change in the classification system; for example, only about 89 percent of the deaths which would have been assigned to arteriosclerosis in the old system would have been placed in this category under the new system.

Table D3. Deaths and percent of total deaths from ten leading causes,
by cause and race

Cause and International Classification Number	Number			Percent		
	Total	White	Non- white	Total	White	Non- white
<u>Total of ten leading causes</u>	<u>19,001</u>	<u>11,223</u>	<u>7,778</u>	<u>79.2</u>	<u>83.3</u>	<u>73.9</u>
1. Diseases of heart (390-398, 402, 404-429)	7,969	5,024	2,945	33.2	37.3	28.0
2. Malignant neoplasms (140-209)	3,267	2,101	1,166	13.6	15.6	11.1
3. Cerebrovascular disease (430-438)	3,115	1,631	1,484	13.0	12.1	14.1
4. Accidents (800-949)	1,723	999	724	7.2	7.4	6.9
5. Influenza and pneumonia (470-486)	825	414	411	3.4	3.1	3.9
6. Certain causes of mortality in early infancy (760-778)	791	320	471	3.3	2.4	4.5
7. Diabetes mellitus (250)	436	221	215	1.8	1.6	2.0
8. Arteriosclerosis (440)	316	211	105	1.3	1.6	1.0
9. Chronic and unqualified bronchitis, emphysema, asthma (490-493)	303	242	61	1.3	1.8	0.6
10. Homicide (960-978)	256	60	196	1.1	0.4	1.9

Leading Causes by Age

Listed in Table D4 are the five leading causes of death for each age group except infants (infant mortality is discussed in another section). Although each of these lists was different from any of the others, malignant neoplasms and accidents appeared on every one and heart diseases on all except that for children 1-4 years old. However, the number of deaths of children and young adults under the age of 25 that were due to malignant neoplasms and heart diseases was relatively insignificant. The leading cause of death for ages 1-44 was accidents and for ages 45 and over, heart diseases. Of special note is the fact that among persons 15-24 years of age the two leading causes of death were external causes (accidents and homicide) rather than diseases and were responsible for 64 percent of the total in that group.

Vital Statistics
Mississippi - 1968

Table D4. Deaths and percent of total deaths from five leading causes, by age and cause

Cause of Death	Age	Number	Percent
1-4 years			
<u>All causes</u>		<u>264</u>	<u>100.0</u>
1. Accidents		97	36.7
2. Influenza and pneumonia		29	11.0
3. Congenital anomalies		21	8.0
4. Enteritis and other diarrheal diseases		14	5.3
5. Malignant neoplasms		11	4.2
5-14 years			
<u>All causes</u>		<u>286</u>	<u>100.0</u>
1. Accidents		158	55.2
2. Malignant neoplasms		29	10.1
3. Congenital anomalies		18	6.3
4. Diseases of heart		8	2.8
5. Meningitis		6	2.1
5. Influenza and pneumonia		6	2.1
15-24 years			
<u>All causes</u>		<u>621</u>	<u>100.0</u>
1. Accidents		350	56.4
2. Homicide		49	7.9
3. Malignant neoplasms		31	5.0
4. Diseases of heart		19	3.0
5. Influenza and pneumonia		17	2.7
25-44 years			
<u>All causes</u>		<u>1,450</u>	<u>100.0</u>
1. Accidents		351	24.2
2. Malignant neoplasms		197	13.6
3. Diseases of heart		194	13.4
4. Homicide		120	8.3
5. Cerebrovascular disease		102	7.0
45-64 years			
<u>All causes</u>		<u>5,649</u>	<u>100.0</u>
1. Diseases of heart		1,982	35.1
2. Malignant neoplasms		1,181	20.9
3. Cerebrovascular disease		526	9.3
4. Accidents		339	6.0
5. Diabetes mellitus		137	2.4
65 years and over			
<u>All causes</u>		<u>14,094</u>	<u>100.0</u>
1. Diseases of heart		5,738	40.7
2. Cerebrovascular disease		2,473	17.5
3. Malignant neoplasms		1,811	12.8
4. Influenza and pneumonia		509	3.6
5. Accidents		351	2.5

Vital Statistics
Mississippi - 1968

Ill-defined and Unknown Causes

A substantial number of the death certificates filed in Mississippi do not have a satisfactory statement of cause of death. In 1968, there were 1,872 deaths in the state for which the cause could only be classified in a category called "symptoms and ill-defined conditions". This group made up 7.9 percent of the total with far more of the records for nonwhites (13.2) thus classified than for whites (3.7 percent). If the 279 certificates for infants whose stated cause of death was immaturity with no additional information were also included as ill-defined, the overall total would be 2,151 and the percentage 9.1. However, considering only those deaths in the category which is actually labeled as ill-defined, review of this information by county of occurrence of death revealed that 20 percent or more of the certificates lacked a satisfactory statement of cause in five counties - Amite, Benton, Issaquena, Noxubee (a high of 31 percent), and Yazoo. For six other counties (DeSoto, Hancock, Holmes, Madison, Marion, and Tunica) the rate was 15-19 percent. If the cause of this large group of deaths could be determined and tabulated, the order of the leading causes and even the causes themselves might change.

Specific Causes

Heart disease, the leading cause of death in Mississippi as well as in the nation, has been the leading cause in this state in every year from 1925 to the present with the exception of 1929. Accounting for 33 percent of the deaths among residents in 1968, the death toll of 7,969 and the death rate of 339.0 per 100,000 population were the highest ever recorded in Mississippi. Although the rate of 346.4 for the white population was higher than that of 327.1 for nonwhites, the difference between the two was not excessively large. Data in Table D5 show that ischemic heart disease caused most of the deaths due to heart disease; the principal category within this group was acute myocardial infarction which alone accounted for 4,039 deaths.

Table D5. Deaths from diseases of heart, by type and race

Cause and International Classification Number	Total	White	Non-white
<u>Diseases of heart (390-398, 402, 404-429)</u>	<u>7,969</u>	<u>5,024</u>	<u>2,945</u>
Active rheumatic fever and chronic rheumatic heart disease (390-398)	98	61	37
Hypertensive heart disease with or without renal disease (402, 404)	296	91	205
Ischemic heart disease (410-414)	6,818	4,496	2,322
Chronic disease of endocardium and other myocardial insufficiency (424, 428)	166	87	79
Other forms of heart disease (420-423, 425-427, 429)	591	289	302

Malignant neoplasms again ranked second on the list of leading causes, having moved up from third to second place in the previous year. As was true for the leading cause, heart disease, the number of deaths (3,267) and the death rate of 139.0 per 100,000 population were the highest on record in Mississippi; however, cancer was responsible for less than half as many deaths as heart disease. Another similarity between these two causes is that the death rate for cancer was higher for whites than for nonwhites but not extremely so. In the section on leading causes of death by age it was noted that cancer

Vital Statistics
Mississippi - 1968

was one of the five leading causes for every age group except infants. Despite this, comparatively few deaths from malignant neoplasms occurred among young persons; in fact, only 8 percent of the total occurred among persons under the age of 45. Study of the organs or systems of the body where the primary neoplasm was located revealed that the three principal sites were the digestive organs and peritoneum (920 deaths), the respiratory system (634 deaths), and the genital organs (501 deaths).

Cerebrovascular disease was the third leading cause of death with a total count (3,115) and a death rate (132.5) that were not much below those for malignant neoplasms. Figures by race show that this disease is one that was especially serious among the non-white population as it ranked second on their list of leading causes and their rate was 47 percent higher than the rate for whites. No trend, either upward or downward, in the mortality from this disease has been noted in over ten years.

Chronic respiratory diseases have been the object of much concern during the 1960's and even earlier because of the rise in both illnesses and deaths due to some of these conditions. In Table D6 the diseases responsible for the largest numbers of deaths are listed with the death toll from each in 1968 and 5, 10, and 15 years earlier. Figures for 20 or more years earlier are not shown because of a major difference in the classification system for causes of death. Although the system was revised again in 1958 and 1968, the changes in 1958 were very minor and only one of the causes in this table was significantly affected by the revision in 1968. For this one cause, asthma, adjusted data for the years prior to 1968 are also given in order to provide a set of comparable figures.

Mortality from tuberculosis of the respiratory system declined considerably in the last 15 years. In contrast, there was a sharp rise in the death counts for malignant neoplasms of the respiratory system and emphysema. In 1968 cancer of the respiratory system caused more deaths than the total of all of the other diseases listed in the table. The number of deaths attributed to cancer of this site increased by 184 percent since 1953, and those from cancer of just the trachea, bronchus, and lung tripled in that time. The upswing in fatal cases of emphysema was even more dramatic with an increase from 12 to 213 deaths in the past 15 years. For chronic and unqualified bronchitis and for asthma no trends in either direction were maintained, although the general level of mortality from bronchitis was higher in the 1960's than in the 1950's.

Table D6. Deaths from major chronic respiratory diseases, 1953, 1958, 1963, 1968

Cause and International Classification Number	1953	1958	1963	1968
Tuberculosis of respiratory system (010-012)	293	154	106	78
Malignant neoplasms of respiratory system (160-163)	223	310	480	634
Malignant neoplasms of trachea, bronchus, and lung (162)	200	279	443	602
Chronic and unqualified bronchitis (490, 491)	28	31	49	38
Emphysema (492)	12	61	152	213
Asthma (493)	62	58	45	52
(Adjusted figures*)	(43)	(40)	(31)	

* Adjusted according to a comparability ratio prepared by the National Center for Health Statistics in order to compensate for a break in comparability caused by revision of the cause-of-death classification system in 1968.

Vital Statistics
Mississippi - 1968

Influenza and pneumonia moved up to fifth place on the list of leading causes in 1968 after being in sixth place for the previous 19 years during which the ranking of leading causes was comparable. The total of 825 deaths in 1968 was 168 more than occurred in 1967 and the highest number since 1963. However, 1968 was not a year of extremely high mortality as in seven of the last twenty years the death toll from this causes was higher than it was in 1968. Included in the figure for 1968 were 729 deaths from pneumonia, 59 from influenza with pneumonia, and 37 from influenza without mention of pneumonia. Data by race showed that, as usual, the rate for nonwhites was substantially larger than that for whites, 45.6 in comparison with 28.5 per 100,000 population.

Accidents caused the death of 1,723 Mississippians in 1968. This toll and the death rate of 73.3 per 100,000 population have been surpassed in only two other years, 1966 and 1967 when the counts were 1,872 and 1,890 respectively. On the list of leading causes of death, accidents ranked fourth and was the leading cause for persons 1-44 years old. Over half of the deaths among persons 5-24 years of age were due to accidents, and this cause was responsible for several times as many deaths in these age groups as the second-ranking causes. November was the peak month of occurrence with December ranking second.

The five principal types of fatal accidents among Mississippi residents are shown in Table D7 and are the same ones which have been listed for a number of years. Together they accounted for nearly 80 percent of the total and the top-ranking type, motor vehicle accidents, accounted for 48 percent. Although the 821 deaths caused by motor vehicle accidents was the third highest count on record for Mississippians, this number was 110 less than the peak number reported the year before, and the decline which occurred in 1968 followed five successive years in which the total for each year surpassed all previous years.

Table D7. Deaths and percent of total accidental deaths
from five leading causes of accidental death, by cause and race

Type of Accident and International Classification Number	Number			Percent		
	Total	White	Non- white	Total	White	Non- white
<u>Total of five leading causes</u>	<u>1,370</u>	<u>806</u>	<u>564</u>	<u>79.5</u>	<u>80.7</u>	<u>77.9</u>
1. Motor vehicle (810-823)	821	542	279	47.6	54.2	38.5
2. Fires and flames (890-899)	208	68	140	12.1	6.8	19.3
3. Falls (880-887)	157	115	42	9.1	11.5	5.8
4. Drowning, except in water transport (910)	111	42	69	6.4	4.2	9.5
5. Firearm (922)	73	39	34	4.2	3.9	4.7

A special compilation of data on accidental deaths that occurred in Mississippi showed that for ages 5-64 the leading cause of accidental death was motor vehicle accidents. For infants under one year of age it was mechanical suffocation; for ages 1-4 it was fires and flames; and for persons 65 and over it was falls.

Suicide and homicide were the causes of death of 171 and 256 Mississippians in 1968. Suicide was not on the list of ten leading causes but did rank in tenth place among the white population. Homicide was of even greater significance as indicated by the fact that it ranked in tenth place in the total population, in second place among young

Vital Statistics
Mississippi - 1968

persons 15-24 years of age, in fourth place among those 25-34 years old, in eighth place among nonwhites, and in eighth place among males. A preponderance of males was noted for both suicides and homicides, but there was a sharp contrast in the occurrence of these events by race. The suicide rate among whites was nearly four times as great as that among nonwhites, but the homicide rate was over five times greater for nonwhites than for whites. An overwhelming majority of both suicides and homicides were accomplished by firearms and explosives - 82 percent of the suicides and 73 percent of the homicides.

Injury undetermined whether accidentally or purposely inflicted is the title of a new category in the cause-of-death classification system which went into effect in 1968. Formerly all deaths not due to natural causes (except those resulting from war operations) had to be classified as either accident, suicide, or homicide. If the certifier could not or would not make a decision, the death certificate was left as the certifier had prepared it but one of the three categories listed above was assigned for tabulating purposes according to certain prescribed rules. In most instances these deaths were classified as accidents. The new category, which was reflected in the design of the revised death certificate, was intended to eliminate such automatic decisions and also incorrect choices by certifiers who made assignments without substantiating evidence just because a decision was expected of them. The number of deaths assigned to this category in 1968 was 51, and 22 resulted from injuries by firearms and explosives. Counts by race were 24 white persons and 27 nonwhites; approximately half were 25-44 years of age.

Maternal deaths are defined as those resulting from deliveries and complications of pregnancy, childbirth, and the puerperium. In 1968 the number in this category was 21 and the death rate was 4.6 per 10,000 live births. Both of these figures were record lows for Mississippi. From the data in Table D8 it may be seen that mothers 15-24 years of age had the lowest maternal death rate and that there was a marked disparity between the rates for whites and nonwhites, 1.3 and 7.8 respectively. Eight deaths were attributed to toxemia, 5 to hemorrhage, 2 to sepsis, and 6 to various other causes.

Table D8. Maternal deaths and maternal death rates, by age and race

Age	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>21</u>	<u>3</u>	<u>18</u>	<u>4.6</u>	<u>1.3</u>	<u>7.8</u>
10-14	1	0	1	32.2	-	34.8
15-24	8	0	8	2.9	-	5.8
25-34	10	2	8	6.9	2.5	12.3
35-44	2	1	1	5.5	7.7	4.3
45-54	0	0	0	-	-	-


* Deaths per 10,000 live births to mothers in specific age-race group.

Infant Mortality

The number of infant deaths declined slightly in 1968 to reach a new record low

Vital Statistics
Mississippi - 1968

of 1,613. No change occurred in the death rate which remained at 35.3 per 1,000 live births; this was also the lowest on record, however. Despite the fact that the infant mortality rate reached a new low in 1967 which was maintained in 1968, the problem is still serious. The death rate of 35.3 was not much below the rate of 36.6 which was recorded 21 years earlier in 1947. As the accompanying graph shows, 1947 was the last year of a long-term downward trend in the total infant mortality rate. In the intervening years only slight progress was made in reducing the rate for white infants and for nonwhite infants the trend was upward with indication of a reversal of this trend just in the last few years. In addition, Mississippi's rate was the highest in the nation. It was 31 percent above that for South Carolina which had the second highest rate and 62 percent above that for the nation as a whole. The race-specific rates for Mississippi were also excessive - 18 percent above the national level for whites and 39 percent for nonwhites.



Data relating to infant deaths by age and race are presented in Table D9. More deaths occurred in the first of life (35 percent of the total) than in either the next 27 days or the other eleven months of infancy. Total neonatal deaths (those in the first 28 days of life) amounted to 69 percent of those in the first year leaving only 31 percent in the postneonatal period. The infant death rate of 47.8 for nonwhites was more than

Vital Statistics
Mississippi - 1968

double the rate of 22.7 for whites. Although rates for nonwhites were higher than those for whites in each age group, the discrepancy was greatest for children 28 days to one year of age. The excessive mortality of nonwhite babies in this age group is a significant factor in the state's extremely high infant death rate. However, Mississippi's neonatal death rate was also the highest in the country and substantially higher than that for any other state.

Table D9. Infant deaths and infant death rates, by age and race

Age	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>1,613</u>	<u>517</u>	<u>1,096</u>	<u>35.3</u>	<u>22.7</u>	<u>47.8</u>
Under one day	565	225	340	12.4	9.9	14.8
One to 28 days	551	199	352	12.0	8.7	15.4
28 days to one year	497	93	404	10.9	4.1	17.6

* Deaths per 1,000 live births.

In previous years a table which listed the leading causes of infant death in rank order was presented in this section on infant mortality. Since no system of ranking infant causes or even grouping them has been standardized yet for the new cause-of-death classification which went into use in 1968, Table D10 presents some causes and cause groups which appeared to be significant. The category on the table with the largest number of deaths, 323, is anoxic and hypoxic conditions not elsewhere classifiable and has as its three major components hyaline membrane disease (92 deaths), respiratory distress syndrome (71 deaths), and unspecified asphyxia of the newborn (143 deaths). A cause group only partially provided for in the previous classification system is that for infant deaths resulting from complications of pregnancy and childbirth. Deaths assigned to this group totaled 114 with 29 resulting from complications due to a multiple pregnancy and 25 from premature separation of the placenta. A breakdown of the 77 accidental deaths showed that 28 were due to mechanical suffocation and 23 were caused by inhalation or ingestion of food or other objects.

Analysis of age at death and race of the infants who died from the causes listed in Table D10 revealed pronounced contrasts. Death rates from diarrheal disease, influenza and pneumonia, and accidents were respectively 27, 5, and 2 times higher for nonwhites than for whites, but the rate for congenital anomalies was twice as high among whites as among nonwhites. Regarding age at death, all but a very few of the deaths from anoxic and hypoxic conditions and from complications of pregnancy and childbirth occurred among babies less than one week old, 69 percent of the deaths caused by congenital anomalies occurred in the neonatal period, and two-thirds or more of the deaths attributed to diarrheal disease, influenza and pneumonia, and accidents were in the postneonatal period.

Vital Statistics
Mississippi - 1968

Table D10. Infant deaths and infant death rates
for significant causes of infant death, by cause and race

Cause of Death and International Classification Number	Number			Rate*		
	Total	White	Non- white	Total	White	Non- white
<u>All causes</u>	<u>1,613</u>	<u>517</u>	<u>1,096</u>	<u>35.3</u>	<u>22.7</u>	<u>47.8</u>
Diarrheal disease (009)	66	3	63	1.4	0.1	2.7
Influenza and pneumonia (470-486)	150	25	125	3.3	1.1	5.4
Congenital anomalies (740-759)	137	90	47	3.0	3.9	2.0
Complications of pregnancy and childbirth, including conditions of placenta and umbilical cord (760-771)	114	51	63	2.5	2.2	2.7
Anoxic and hypoxic conditions not elsewhere classifiable (776)	323	163	160	7.1	7.1	7.0
Immaturity, unqualified (777)	283	76	207	6.2	3.3	9.0
Accidents (800-949)	77	26	51	1.7	1.1	2.2

* Deaths per 1,000 live births.

If immaturity is mentioned on a death certificate with some other condition(s), it is not classified as the underlying cause because of the vagueness of the term, but some certificates show no other condition except immaturity. In 1968 this total was 283 which is the second largest in Table D10. In addition, there were 233 other deaths with no definite statement of cause which made a total of 516 in the "ill-defined" categories. This was 32 percent of the total infant deaths; 39 percent of those among nonwhites and 41 percent of those among infants under one day of age were thus classified. The magnitude of these figures indicates a major difficulty in the study of the problem of infant mortality.

MARRIAGES (See Tables 9, 17)

Trend

The year 1968 was the ninth in succession in which the number of marriages went up. Weddings occurring in that year numbered 25,355, an increase of nearly eight percent over the figure for 1967, and the size of the increase was much larger than that in any other year of this period of upward trend. The rise in marriages is no doubt due to a shift in the age distribution of the population as the unusually large number of children born in the years immediately after World War II reach marriageable age and also to overall growth of the state's population. For both whites and nonwhites the trend has been upward, though not unbroken, with the latest counts 17,142 and 8,213 respectively.

Month

As expected, marriages in June far outnumbered those in any other month (see Table E1). August ranked second in number and daily average with December in third place. During the 1960's the order in which these two months ranked usually depended on which one had five Saturdays, as more marriages occur on Saturday than any other day of the week. The National Center for Health Statistics reported that, in 1968, 34 percent of the

Vital Statistics
Mississippi - 1968

marriages in Mississippi occurred on Saturday. This concentration and the fact that there can be either four or five Saturdays in any month result in year-to-year variations in the number of marriages occurring in any certain month.

Table E1. Marriages and average per day, by month

Month	Marriages	Average per Day	Month	Marriages	Average per Day
<u>Total</u>	<u>25,355</u>	<u>69</u>			
January	1,507	49	July	2,274	73
February	1,641	56	August	2,983	96
March	1,824	59	September	1,932	64
April	1,732	58	October	1,715	55
May	1,983	64	November	2,032	68
June	3,263	109	December	2,469	80

Officiant

The ceremony was performed by a minister at 72 percent of the marriages. Justices of the peace officiated at nearly all the rest with only a few conducted by mayors, judges, members of boards of supervisors, and other persons.

Age of Bride and Groom

Information in Table E2 shows that 10,444 of the brides married in Mississippi in 1968 were teenagers (or subteenage). They comprised 41 percent of the total, and 4,589 or 18 percent were less than 18 years of age. One bride only 12 years old, 13 who were 13 years old, and 89 who were 14 were included in this group. Teenage grooms numbered 4,697 which was 18 percent of the total. The youngest were only 14 years old and 783 were under age 18. Old age as well as extreme youth was represented among those getting married, however, with 31 brides and 145 grooms in the group 75 years of age or older. The oldest ages were 97 for grooms and 85 for brides. Differences in age distribution by race were not large, but for both brides and grooms the proportions of whites under age 25 were higher than the proportions of nonwhites of that age.

As may be seen from the figures in Table E3 which shows median ages at marriage, brides were younger than grooms (as is to be expected) and white brides and grooms were younger than nonwhites of the same sex. Information is also given for brides and grooms marrying for the first time. These brides and grooms had lower median ages, but the same relationships between the races and the sexes were noted as for all marriages.

Vital Statistics
Mississippi - 1968

Table E2. Brides and grooms and percentage distribution, by age and race

Age	Number						Percent					
	Bride			Groom			Bride			Groom		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Total	25,355	17,142	8,213	25,355	17,149	8,206	100.0	100.0	100.0	100.0	100.0	100.0
Under 20	10,444	7,155	3,289	4,697	3,251	1,446	41.2	41.7	40.0	18.5	19.0	17.6
12	1	0	1	0	0	0	*	-	*	-	-	-
13	13	7	6	0	0	0	*	*	0.1	-	-	-
14	89	47	42	2	1	1	0.4	0.3	0.5	*	*	*
15	827	569	258	18	11	7	3.3	3.3	3.1	0.1	0.1	0.1
16	1,443	945	498	79	50	29	5.7	5.5	6.1	0.3	0.3	0.4
17	2,216	1,501	715	684	493	191	8.7	8.8	8.7	2.7	2.9	2.3
18	3,032	2,113	919	1,569	1,057	512	12.0	12.3	11.2	6.2	6.2	6.2
19	2,823	1,973	850	2,345	1,639	706	11.1	11.5	10.3	9.2	9.6	8.6
20-24	7,876	5,483	2,393	10,901	7,578	3,323	31.1	32.0	29.1	43.0	44.2	40.5
25-34	3,315	2,126	1,189	4,935	3,314	1,621	13.1	12.4	14.5	19.5	19.3	19.8
35-44	1,750	1,165	585	1,992	1,384	608	6.9	6.8	7.1	7.8	8.1	7.4
45-54	1,058	710	348	1,297	837	460	4.2	4.1	4.2	5.1	4.9	5.6
55-64	634	347	287	893	477	416	2.5	2.0	3.5	3.5	2.8	5.1
65-74	229	130	99	480	230	250	0.9	0.8	1.2	1.9	1.3	3.0
75 & over	31	18	13	145	75	70	0.1	0.1	0.2	0.6	0.4	0.8
Unknown	18	8	10	15	3	12	0.1	*	0.1	*	*	0.1

* Less than 0.06 percent.

Table E3. Median age of bride and groom, for all marriages and first marriages, by race

All and First Marriages	Bride		Groom	
	White	Non-white	White	Non-white
All marriages	20.8	21.1	22.8	23.5
First marriages	19.5	20.0	22.1	22.6

Marriage Order and Previous Marital Status

Data presented in Table E4 show that 73 percent of the brides and 73 percent of the grooms who married in 1968 were married for the first time. However, the proportion of weddings in which both bride and groom were married for the first time was somewhat smaller, 66 percent. Most of those who had been married before had only one previous marriage, but 1,622 brides and exactly the same number of grooms undertook a marriage of third or higher order and one bride began her eighth marriage. Far more of both brides and grooms were divorced than were widowed.

Vital Statistics
Mississippi - 1968

Table E4. Brides and grooms and percentage distribution,
by marriage order and previous marital status

Marriage Order and Previous Marital Status	Number		Percent	
	Bride	Groom	Bride	Groom
<u>Total</u>	<u>25,355</u>	<u>25,355</u>	<u>100.0</u>	<u>100.0</u>
First marriage	18,531	18,573	73.1	73.2
<u>Remarriage</u>	<u>6,772</u>	<u>6,748</u>	<u>26.7</u>	<u>26.6</u>
Second	5,150	5,126	20.3	20.2
<u>Third or higher</u>	<u>1,622</u>	<u>1,622</u>	<u>6.4</u>	<u>6.4</u>
Third	1,284	1,271	5.1	5.0
Fourth	286	293	1.1	1.2
Fifth	35	38	0.1	0.1
Sixth	15	13	**	**
Seventh	1	7	**	**
Eighth	1	0	**	-
Widowed	1,612	1,367	6.4	5.4
Divorced*	5,039	5,255	19.9	20.7
Unknown whether widowed or divorced	121	126	0.5	0.5
Unknown whether any previous marriages	52	34	0.2	0.1

* Includes those whose last marriage was annulled.

** Less than 0.06 percent.

Nonresident Brides

Fifteen percent of the brides married in Mississippi in 1968 were not residents of the state, and in 12 percent of the marriages neither the bride nor the groom was a state resident. Although the 3,738 nonresident brides came from 44 of the other 49 states, the District of Columbia, and also from foreign countries, 79 percent were from the neighboring states of Alabama, Arkansas, Louisiana, and Tennessee. DeSoto County, just south of Memphis, Tennessee, issued 20 percent of the licenses to nonresident brides, and all Mississippi border counties together accounted for 76 percent.

Out-of-State Use of Mississippi Licenses

Some marriage licenses issued in Mississippi are used for weddings performed in other states. Such marriages are not included in the annual tabulations for this state because they did not occur here and are of questionable legality. Special counts of these records have been made during a twelve-year period, and the total of 71 in 1968 is the smallest recorded during that time. The bride and/or groom resided in the state

Vital Statistics
Mississippi - 1968

of occurrence in 72 percent of these cases, and in all except one of the remaining cases the bride and/or groom resided in Mississippi. States which border on Mississippi were the site of all but 8 of the weddings with 37 occurring in Louisiana. It was also noted that two-thirds of the licenses were obtained in Mississippi counties adjacent to the state where the wedding took place.

DIVORCES (See Tables 9, 10, 17)

Trend

In 1968 the number of divorces in Mississippi was the second largest on record. The total of 7,020 was exceeded only in 1946 when an extremely high peak was reached following World War II. Counts for both whites and nonwhites went up in 1968, with that for the whites also the second largest on record and almost 1,000 more than occurred only three years earlier.

Cause and Complainant

The most frequently given reason for divorce was cruel and inhuman treatment which was listed for 62 percent of the cases. Desertion ranked second with 29 percent and was the stated cause of more than half of those among nonwhite couples. Together these two causes accounted for 91 percent of the divorces granted in this state in 1968. The wife was the complainant in 71 percent of the cases involving white persons, but in only 53 percent of those among nonwhites.

Minor Children

A large proportion of the couples who were granted divorces stated that they had no minor children - 42 percent of the white and 50 percent of the nonwhite couples. Those who did have minor children averaged 2.2 per family with a total of 8,382 children reported. There were 274 families with five or more children and 5 of these had between ten and thirteen minor children.

Duration of Marriage

The length of time married before obtaining a divorce covered a broad range - from less than one year to 62 years. Eight percent of the divorces occurring in 1968 came after less than one year of marriage and 42 percent were granted during the first five years. However, approximately one-sixth of the couples had been married twenty years or longer and 20 had been married for fifty or more years. There was a substantial disparity between the length of time reported for whites and nonwhites. The median duration for whites was 5.1 years with 49 percent in the first five years, but for nonwhites the median was 12.4 years with only 23 percent in the first five years.

Vital Statistics
Mississippi - 1968

Additional Information Needed

Mississippi's statistical record of divorce is a form which provides only limited information because of the omission of a number of pertinent items which many other states collect and also because of the lack of clarity regarding some of the items which are on the form. No data at all are available concerning place of marriage, place of residence of the couple at time of separation, their ages at marriage and at divorce, previous marriages, and socio-economic factors. Information which is presently tabulated on minor children, cause of divorce, and complainant probably is erroneous because of lack of definitions and instructions regarding how such facts should be entered on the form and also because of the poor arrangement of the form. These deficiencies seriously hamper meaningful study of the divorce problem on a state level as well as on a national level. Since Mississippi's record does not contain even what is considered minimum information, this state is not eligible to become part of the National Divorce Registration Area. Revision of the statistical record of divorce would be of great help in providing material for analysis of this serious social problem.

TABLE 1
ESTIMATED POPULATION*
MISSISSIPPI
July 1, 1968

By County, by Race

County	Total	White	Nonwhite
<u>STATE</u>	<u>2,350,590</u>	<u>1,450,340</u>	<u>900,250</u>
Adams	36,730	18,970	17,760
Alcorn	27,690	24,630	3,060
Amite	15,200	7,040	8,160
Attala	22,660	13,200	9,460
Benton	7,370	4,220	3,150
Bolivar	54,650	19,840	34,810
Calhoun	15,790	12,060	3,730
Carroll	10,680	5,410	5,270
Chickasaw	19,110	12,210	6,900
Choctaw	8,610	6,330	2,280
Claiborne	11,380	3,240	8,140
Clarke	15,520	9,980	5,540
Clay	18,190	8,980	9,210
Coahoma	46,540	16,480	30,060
Copiah	26,360	13,800	12,560
Covington	14,330	10,050	4,280
DeSoto	36,760	22,590	14,170
Forrest	61,570	45,660	15,910
Franklin	9,300	6,240	3,060
George	11,720	10,380	1,340
Greene	8,150	6,370	1,780
Grenada	20,050	10,810	9,240
Hancock	20,920	17,790	3,130
Harrison	141,100	118,910	22,190
Hinds	216,200	129,810	86,390
Holmes	27,550	8,220	19,330
Humphreys	16,700	5,420	11,280
Issaquena	2,880	1,050	1,830
Itawamba	18,930	18,110	820
Jackson	81,700	68,180	13,520
Jasper	16,720	8,980	7,740
Jefferson	10,700	3,300	7,400
Jeff Davis	14,250	6,670	7,580
Jones	58,930	44,620	14,310
Kemper	11,590	5,310	6,280
Lafayette	26,230	19,220	7,010

Table 1 - Page 2
 Estimated Population - Mississippi - July 1, 1968

County	Total	White	Nonwhite
Lamar	15,820	13,090	2,730
Lauderdale	76,880	52,600	24,280
Lawrence	11,240	7,710	3,530
Leake	19,560	11,880	7,680
Lee	48,680	38,360	10,320
Leflore	48,240	18,220	30,020
Lincoln	26,220	18,130	8,090
Lowndes	56,800	37,350	19,450
Madison	32,490	10,830	21,660
Marion	23,190	15,400	7,790
Marshall	24,080	8,680	15,400
Monroe	35,880	24,910	10,970
Montgomery	14,590	8,540	6,050
Neshoba	20,050	14,800	5,250
Newton	20,600	14,560	6,040
Noxubee	15,210	4,590	10,620
Oktibbeha	31,020	19,800	11,220
Panola	29,330	12,960	16,370
Pearl River	29,080	23,970	5,110
Perry	9,120	6,400	2,720
Pike	32,410	18,420	13,990
Pontotoc	18,140	14,670	3,470
Prentiss	21,470	19,110	2,360
Quitman	18,070	7,240	10,830
Rankin	42,690	29,890	12,800
Scott	24,010	16,030	7,980
Sharkey	9,520	3,320	6,200
Simpson	20,230	14,800	5,430
Smith	13,000	9,620	3,380
Stone	8,110	6,210	1,900
Sunflower	41,760	13,260	28,500
Tallahatchie	21,460	7,570	13,890
Tate	20,610	9,670	10,940
Tippah	17,650	14,390	3,260
Tishomingo	16,690	16,090	600
Tunica	15,110	4,300	10,810
Union	19,920	16,760	3,160
Walthall	14,670	8,130	6,540
Warren	45,800	26,210	19,590
Washington	76,550	34,850	41,700
Wayne	17,220	10,820	6,400
Webster	11,140	8,530	2,610
Wilkinson	11,100	3,340	7,760
Winston	18,670	9,650	9,020
Yalobusha	12,470	7,170	5,300
Yazoo	31,280	13,430	17,850

*Prepared by Division of Sociology and Anthropology,
 Mississippi State University.

TABLE 2
LIVE BIRTHS
MISSISSIPPI
1968

By Place of Occurrence, and by Place of Residence of Mother, by Race, with Rates

County	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
<u>STATE</u>	<u>45,706</u>	<u>22,896</u>	<u>22,810</u>	<u>45,719</u>	<u>22,807</u>	<u>22,912</u>	<u>19.4</u>	<u>15.7</u>	<u>25.4</u>
Adams	1,358	666	692	817	391	426	22.2	20.6	24.0
Alcorn	374	293	81	473	387	86	17.1	15.7	28.1
Amite	93	1	92	265	82	183	17.4	11.6	22.4
Attala	375	161	214	387	150	237	17.1	11.4	25.0
Benton	56	0	56	135	51	84	18.3	12.1	26.7
Bolivar	1,595	359	1,236	1,404	337	1,067	25.7	17.0	30.6
Calhoun	204	94	110	270	175	95	17.1	14.5	25.5
Carroll	77	9	68	153	49	104	14.3	9.0	19.7
Chickasaw	305	120	185	362	172	190	18.9	14.1	27.5
Choctaw	141	60	81	135	76	59	15.7	12.0	25.9
Claiborne	92	12	80	184	38	146	16.2	11.7	17.9
Clarke	59	32	27	269	135	134	17.3	13.5	24.2
Clay	271	99	172	375	140	235	20.6	15.6	25.5
Coahoma	871	253	618	902	207	695	19.4	12.6	23.1
Copiah	583	185	398	468	170	298	17.8	12.3	23.7
Covington	185	132	53	295	163	132	20.6	16.2	30.8
DeSoto	290	19	271	781	393	388	21.2	17.4	27.4
Forrest	1,439	1,061	378	1,119	726	393	18.2	15.9	24.7
Franklin	66	29	37	124	60	64	13.3	9.6	20.9
George	282	229	53	239	196	43	20.4	18.9	32.1
Greene	69	26	43	174	104	70	21.3	16.3	39.3
Grenada	384	133	251	427	178	249	21.3	16.5	26.9
Hancock	160	118	42	338	278	60	16.2	15.6	19.2
Harrison	3,407	2,694	713	3,054	2,372	682	21.6	19.9	30.7
Hinds	5,618	2,868	2,750	4,391	2,175	2,216	20.3	16.8	25.6
Holmes	522	78	444	535	88	447	19.4	10.7	23.1
Humphreys	377	52	325	393	98	295	23.5	18.1	26.2
Issaquena	7	0	7	60	14	46	20.8	13.3	25.1
Itawamba	39	36	3	228	214	14	12.0	11.8	17.1
Jackson	1,548	1,142	406	1,864	1,433	431	22.8	21.0	31.9
Jasper	167	111	56	292	106	186	17.5	11.8	24.0
Jefferson	69	8	61	176	20	156	16.4	6.1	21.1
Jeff Davis	160	84	76	260	94	166	18.2	14.1	21.9
Jones	1,860	801	1,059	954	650	304	16.2	14.6	21.2
Kemper	88	10	78	207	46	161	17.9	8.7	25.6
Lafayette	496	310	186	435	251	184	16.6	13.0	26.2

Table 2 - Page 2
Live Births - Mississippi - 1968

County	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Lamar	86	58	28	281	222	59	17.8	17.0	21.6
Lauderdale	1,966	975	991	1,412	724	688	18.4	13.8	28.3
Lawrence	124	69	55	212	128	84	18.9	16.6	23.8
Leake	178	92	86	301	135	166	15.4	11.4	21.6
Lee	1,293	1,017	276	944	678	266	19.4	17.7	25.8
Leflore	1,104	346	758	1,017	262	755	21.1	14.4	25.1
Lincoln	522	338	184	457	279	178	17.4	15.4	22.0
Lowndes	1,131	644	487	992	491	501	17.5	13.1	25.8
Madison	525	76	449	772	159	613	23.8	14.7	28.3
Marion	324	189	135	444	248	196	19.1	16.1	25.2
Marshall	334	31	303	531	155	376	22.0	17.8	24.4
Monroe	787	515	272	646	394	252	18.0	15.8	23.0
Montgomery	321	153	168	254	111	143	17.4	13.0	23.6
Neshoba	351	238	113	376	233	143	18.8	15.7	27.2
Newton	304	181	123	349	181	168	16.9	12.4	27.8
Noxubee	318	67	251	306	61	245	20.1	13.3	23.1
Oktibbeha	606	272	334	543	285	258	17.5	14.4	23.0
Panola	634	192	442	610	209	401	20.8	16.1	24.5
Pearl River	597	476	121	555	387	168	19.1	16.1	32.9
Perry	104	75	29	191	125	66	20.9	19.5	24.3
Pike	624	325	299	583	261	322	18.0	14.2	23.0
Pontotoc	294	214	80	276	211	65	15.2	14.4	18.7
Prentiss	406	324	82	352	277	75	16.4	14.5	31.8
Quitman	459	126	333	385	102	283	21.3	14.1	26.1
Rankin	78	35	43	755	496	259	17.7	16.6	20.2
Scott	411	285	126	437	256	181	18.2	16.0	22.7
Sharkey	88	11	77	224	42	182	23.5	12.6	29.4
Simpson	277	179	98	396	210	186	19.6	14.2	34.2
Smith	13	0	13	271	151	120	20.8	15.7	35.5
Stone	138	97	41	154	102	52	19.0	16.4	27.4
Sunflower	837	136	701	994	192	802	23.8	14.5	28.1
Tallahatchie	441	74	367	457	75	382	21.3	9.9	27.5
Tate	146	1	145	367	119	248	17.8	12.3	22.7
Tippah	342	268	74	309	236	73	17.5	16.4	22.4
Tishomingo	149	143	6	195	188	7	11.7	11.7	11.7
Tunica	324	31	293	344	54	290	22.8	12.6	26.8
Union	281	197	84	324	241	83	16.3	14.4	26.3
Walthall	325	178	147	250	100	150	17.0	12.3	22.9
Warren	1,305	534	771	944	472	472	20.6	18.0	24.1
Washington	1,850	665	1,185	1,726	571	1,155	22.5	16.4	27.7
Wayne	325	222	103	311	158	153	18.1	14.6	23.9
Webster	91	63	28	167	108	59	15.0	12.7	22.6
Wilkinson	221	96	125	220	50	170	19.8	15.0	21.9
Winston	291	136	155	334	145	189	17.9	15.0	21.0
Yalobusha	168	64	104	210	89	121	16.8	12.4	22.8
Yazoo	526	203	323	596	215	381	19.0	16.0	21.3
Out of state	1,497	1,146	351	1,484	1,235	249			

Table 2 - Page 3
Live Births - Mississippi - 1968

City**	Place of Occurrence			Place of Residence		
	Number			Number		
	Total	White	Non-white	Total	White	Non-white
Biloxi	2,000	1,752	248	1,301	1,094	207
Clarksdale	634	253	381	461	159	302
Cleveland	596	289	307	336	159	177
Columbus	822	451	371	557	242	315
Corinth	24	2	22	175	121	54
Greenville	1,156	527	629	922	349	573
Greenwood	757	346	411	485	170	315
Gulfport	1,372	936	436	881	607	274
Hattiesburg	1,421	1,060	361	757	447	310
Jackson	4,959	2,391	2,568	3,283	1,610	1,673
Laurel	1,844	789	1,055	429	216	213
McComb	495	300	195	223	117	106
Meridian	1,943	974	969	935	465	470
Natchez	1,354	666	688	549	233	316
Pascagoula	1,403	1,075	328	654	527	127
Tupelo	1,180	940	240	401	319	82
Vicksburg	1,298	534	764	581	246	335
Yazoo City	336	202	134	264	98	166
Total urban#	33,362	18,793	14,569	18,879	10,152	8,727
Total rural	12,344	4,103	8,241	26,840	12,655	14,185

* Births per 1,000 population.

** Cities having 10,000 population and over according to 1960 census.

Includes all incorporated towns of 2,500 population or more according to 1960 census.

TABLE 3
LIVE BIRTHS
MISSISSIPPI
1968

By Place of Residence of Mother, by Race and Attendant, and by Sex

County	Total	White			Nonwhite			Sex			
		Total	Hosp. or Clinic	Phys., Other Place	Other	Total	Hosp. or Clinic	Phys., Other Place	Other	Male	Female
STATE	45,719	22,807	22,704	34	69	22,912	15,821	448	6,643	23,428	22,291
Adams	817	391	390	1	0	426	421	0	5	413	404
Alcorn	473	387	385	0	2	86	55	0	31	229	244
Amite	265	82	82	0	0	183	94	2	87	154	111
Attala	387	150	149	0	1	237	86	0	151	205	182
Benton	135	51	51	0	0	84	27	0	57	70	65
Bolivar	1,404	337	336	1	0	1,067	820	5	242	720	684
Calhoun	270	175	173	1	1	95	80	1	14	120	150
Carroll	153	49	49	0	0	104	46	1	57	66	87
Chickasaw	362	172	171	1	0	190	146	1	43	174	188
Choctaw	135	76	76	0	0	59	37	0	22	77	58
Claiborne	184	38	36	2	0	146	90	0	56	104	80
Clarke	269	135	135	0	0	134	123	2	9	142	127
Clay	375	140	140	0	0	235	147	17	71	197	178
Coahoma	902	207	207	0	0	695	398	1	296	475	427
Copiah	468	170	169	1	0	298	240	21	37	235	233
Covington	295	163	163	0	0	132	123	0	9	154	141
DeSoto	781	393	391	2	0	388	289	1	98	418	363
Forrest	1,119	726	722	0	4	393	319	0	74	553	566
Franklin	124	60	60	0	0	64	45	1	18	69	55
George	239	196	196	0	0	43	38	0	5	120	119
Greene	174	104	104	0	0	70	62	0	8	94	80
Grenada	427	178	177	0	1	249	87	25	137	215	212
Hancock	338	278	273	0	5	60	35	0	25	184	154
Harrison	3,054	2,372	2,357	2	13	682	486	5	191	1,555	1,499

Table 3 - Page 3

Live Births - Mississippi - 1968

County	Total	White				Nonwhite				Sex	
		Total	Hosp. or Clinic	Phys., Other Place	Other	Total	Hosp. or Clinic	Phys., Other Place	Other	Male	Female
Pearl River	555	387	386	1	0	168	150	1	17	272	283
Perry	191	125	125	0	0	66	64	0	2	95	96
Pike	583	261	261	0	0	322	201	29	92	309	274
Pontotoc	276	211	208	1	2	65	61	0	4	153	123
Prentiss	352	277	277	0	0	75	55	1	19	183	169
Quitman	385	102	102	0	0	283	244	3	36	184	201
Rankin	755	496	491	2	3	259	245	4	10	407	348
Scott	437	256	254	2	0	181	143	9	29	241	196
Sharkey	224	42	42	0	0	182	132	0	50	109	115
Simpson	396	210	210	0	0	186	172	1	13	206	190
Smith	271	151	151	0	0	120	106	2	12	146	125
Stone	154	102	101	1	0	52	35	1	16	75	79
Sunflower	994	192	187	2	3	802	272	6	524	520	474
Tallahatchie	457	75	74	1	0	382	113	8	261	230	227
Tate	367	119	117	2	0	248	104	10	134	184	183
Tippah	309	236	233	3	0	73	65	1	7	168	141
Tishomingo	195	188	188	0	0	7	6	1	0	102	93
Tunica	344	54	53	0	1	290	156	1	133	169	175
Union	324	241	239	0	2	83	38	5	40	173	151
Walthall	250	100	90	0	10	150	83	1	66	128	122
Warren	944	472	472	0	0	472	445	1	26	486	458
Washington	1,726	571	565	1	5	1,155	847	9	299	892	834
Wayne	311	158	158	0	0	153	139	0	14	171	140
Webster	167	108	108	0	0	59	41	3	15	80	87
Wilkinson	220	50	50	0	0	170	95	1	74	101	119
Winston	334	145	145	0	0	189	98	0	91	159	175
Yalobusha	210	89	89	0	0	121	100	1	20	99	111
Yazoo	596	215	213	0	2	381	123	0	258	296	300

Table 3 - Page 4
Live Births - Mississippi - 1968

City*	Total	White				Nonwhite				Sex	
		Total	Hosp. or Clinic	Phys., Other Place	Other	Total	Hosp. or Clinic	Phys., Other Place	Other	Male	Female
Biloxi	1,301	1,094	1,091	1	2	207	183	1	23	675	626
Clarksdale	461	159	159	0	0	302	241	1	60	245	216
Cleveland	336	159	159	0	0	177	139	0	38	167	169
Columbus	557	242	238	0	4	315	144	5	166	284	273
Corinth	175	121	119	0	2	54	32	0	22	79	96
Greenville	922	349	347	0	2	573	432	2	139	469	453
Greenwood	485	170	170	0	0	315	180	1	134	228	257
Gulfport	881	607	603	1	3	274	200	3	71	436	445
Hattiesburg	757	447	446	0	1	310	255	0	55	366	391
Jackson	3,283	1,610	1,607	1	2	1,673	1,629	14	30	1,680	1,603
Laurel	429	216	216	0	0	213	210	2	1	239	190
McComb	223	117	117	0	0	106	75	10	21	112	111
Meridian	935	465	465	0	0	470	465	4	1	461	474
Natchez	549	233	233	0	0	316	315	0	1	279	270
Pascagoula	654	527	527	0	0	127	90	8	29	351	303
Tupelo	401	319	319	0	0	82	76	2	4	216	185
Vicksburg	581	246	246	0	0	335	317	0	18	296	285
Yazoo City	264	98	97	0	1	166	81	0	85	119	145

* Cities having 10,000 population and over according to 1960 census.

TABLE 4
FETAL DEATHS
MISSISSIPPI
1968

By Place of Occurrence; by Place of Residence of Mother, by Race and Attendant, and by Sex

County	Total by Occurrence	Total by Residence	White			Nonwhite			Sex				
			Total	Hosp. or Clinic	Phys., Other Place	Other	Total	Hosp. or Clinic	Phys., Other Place	Other	Male	Female	Unde- terminated
STATE	2,373	2,322	1,119	1,101	16	2	1,203	977	36	190	778	520	1,024
Adams	82	48	20	20	0	0	28	28	0	0	21	9	18
Alcorn	8	8	6	6	0	0	2	1	0	1	5	3	0
Amite	2	10	4	4	0	0	6	4	0	2	5	3	2
Attala	19	20	9	9	0	0	11	8	0	3	4	4	12
Benton	0	4	4	4	0	0	0	0	0	0	0	1	3
Bolivar	40	38	8	8	0	0	30	21	0	9	17	16	5
Calhoun	15	15	7	6	1	0	8	6	2	0	6	3	6
Carroll	4	5	2	2	0	0	3	0	0	3	2	1	2
Chickasaw	40	30	16	16	0	0	14	14	0	0	10	9	11
Choctaw	5	4	2	2	0	0	2	1	0	1	0	4	0
Claiborne	2	11	2	2	0	0	9	8	0	1	3	2	6
Clarke	2	14	8	8	0	0	6	6	0	0	6	1	7
Clay	7	10	4	4	0	0	6	5	1	0	4	5	1
Coahoma	51	41	13	13	0	0	28	22	0	6	16	10	15
Copiah	12	18	3	3	0	0	15	12	0	3	7	6	5
Covington	1	10	2	2	0	0	8	8	0	0	3	3	4
DeSoto	11	27	4	4	0	0	23	18	0	5	12	15	0
Forrest	33	34	17	17	0	0	17	15	0	2	10	13	11
Franklin	3	5	0	0	0	0	5	5	0	0	2	3	0
George	3	3	1	1	0	0	2	2	0	0	1	2	0
Greene	0	2	2	2	0	0	0	0	0	0	1	0	1
Grenada	17	20	7	7	0	0	13	5	4	4	9	8	3
Hancock	1	4	2	2	0	0	2	2	0	0	1	2	1
Harrison	289	258	193	193	0	0	65	59	1	5	53	26	179

Table 4 - Page 2
Fetal Deaths - Mississippi - 1968

County	Total by Occurrence	Total by Residence	White				Nonwhite				Sex		Unde- terminated
			Total	Hosp. or Clinic	Phys., Other Place	Other	Total	Hosp. or Clinic	Phys., Other Place	Other	Male	Female	
Hinds	580	404	227	223	4	0	177	174	0	3	82	49	273
Holmes	22	23	2	2	0	0	21	10	0	11	10	10	3
Humphreys	16	21	6	6	0	0	15	8	0	7	5	12	4
Issaquena	1	3	2	2	0	0	1	1	0	0	1	0	2
Itawamba	0	1	1	1	0	0	0	0	0	0	1	0	0
Jackson	32	58	45	45	0	0	13	12	1	0	24	14	20
Jasper	7	12	3	3	0	0	9	8	0	1	6	5	1
Jefferson	1	3	2	2	0	0	1	1	0	0	1	0	2
Jeff Davis	9	14	7	7	0	0	7	6	0	1	4	8	2
Jones	78	33	8	8	0	0	25	23	1	1	13	9	11
Kemper	2	6	2	2	0	0	4	4	0	0	2	1	3
Lafayette	19	22	8	8	0	0	14	6	0	8	9	9	4
Lamar	1	5	2	2	0	0	3	3	0	0	4	0	1
Lauderdale	100	66	46	46	0	0	20	19	1	0	14	10	42
Lawrence	4	7	1	1	0	0	6	3	2	1	6	1	0
Leake	2	8	4	4	0	0	4	3	0	1	3	2	3
Lee	23	22	12	12	0	0	10	9	0	1	15	6	1
Leflore	80	70	27	23	4	0	43	26	3	14	24	9	37
Lincoln	11	25	8	8	0	0	17	15	2	0	7	12	6
Lowndes	29	29	10	10	0	0	19	15	1	3	16	10	3
Madison	27	62	21	21	0	0	41	32	1	8	25	6	31
Marion	25	34	18	17	1	0	16	13	0	3	7	11	16
Marshall	13	15	0	0	0	0	15	8	0	7	12	2	1
Monroe	19	20	8	8	0	0	12	12	0	0	9	8	3
Montgomery	4	2	0	0	0	0	2	1	0	1	1	1	0
Neshoba	7	15	6	6	0	0	9	8	1	0	5	6	4
Newton	29	28	18	16	2	0	10	6	4	0	7	2	19
Noxubee	11	13	1	1	0	0	12	10	0	2	7	6	0
Oktibbeha	17	19	8	8	0	0	11	10	0	1	12	6	1
Panola	26	25	4	4	0	0	21	19	0	2	15	10	0

Table 4 - Page 3
Fetal Deaths - Mississippi - 1968

County	Total by Occurrence	Total by Residence	White				Nonwhite				Sex		
			Total	Hosp. or Clinic	Phys., Other Place	Other	Total	Hosp. or Clinic	Phys., Other Place	Other	Male	Female	Unde- terminated
Pearl River	8	11	5	5	0	0	6	5	1	0	7	1	3
Perry	2	5	4	4	0	0	1	1	0	0	1	3	1
Pike	16	18	4	4	0	0	14	11	0	3	7	5	6
Pontotoc	8	9	8	8	0	0	1	1	0	0	4	3	2
Prentiss	5	6	6	5	0	1	0	0	0	0	2	4	0
Quitman	18	22	3	3	0	0	19	18	0	1	10	10	2
Rankin	3	50	42	41	1	0	8	7	0	1	8	4	38
Scott	11	14	8	8	0	0	6	4	0	2	10	2	2
Sharkey	2	13	2	2	0	0	11	9	0	2	4	5	4
Simpson	8	18	10	10	0	0	8	7	0	1	8	5	5
Smith	1	6	3	2	1	0	3	3	0	0	2	2	2
Stone	1	4	3	3	0	0	1	1	0	0	2	1	1
Sunflower	24	43	9	8	0	1	34	20	4	10	25	14	4
Tallahatchie	15	20	5	4	1	0	15	8	0	7	11	3	6
Tate	5	12	2	2	0	0	10	5	0	5	8	4	0
Tippah	27	19	11	11	0	0	8	8	0	0	7	1	11
Tishomingo	7	6	4	4	0	0	2	2	0	0	2	2	2
Tunica	15	16	2	2	0	0	14	8	1	5	9	5	2
Union	9	11	7	7	0	0	4	4	0	0	4	3	4
Walthall	16	11	6	6	0	0	5	3	0	2	2	6	3
Warren	142	94	51	51	0	0	43	42	0	1	22	8	64
Washington	117	116	50	50	0	0	66	52	4	10	40	26	50
Wayne	22	21	12	12	0	0	9	9	0	0	7	5	9
Webster	2	7	3	3	0	0	4	2	1	1	3	3	1
Wilkinson	6	10	2	2	0	0	8	7	0	1	4	2	4
Winston	13	15	4	4	0	0	11	7	0	4	8	6	1
Yalobusha	5	5	1	1	0	0	4	3	0	1	2	2	1
Yazoo	23	36	9	8	1	0	27	15	0	12	14	11	11

Table 4 - Page 4
Fetal Deaths - Mississippi - 1968

City*	Total by Occurrence	Total by Residence	White			Nonwhite			Sex				
			Total	Hosp. or Clinic	Phys., Other Place	Other	Total	Hosp. or Clinic	Phys., Other Place	Other	Male	Female	Unde- terminated
Biloxi	212	132	108	108	0	0	24	24	0	0	13	6	113
Clarksdale	45	28	13	13	0	0	15	15	0	0	9	4	15
Cleveland	13	7	4	4	0	0	3	1	0	2	3	2	2
Columbus	24	15	4	4	0	0	11	10	0	1	9	4	2
Corinth	0	2	1	1	0	0	1	1	0	0	2	0	0
Greenville	94	69	33	33	0	0	36	29	3	4	17	20	32
Greenwood	65	36	19	17	2	0	17	13	1	3	10	4	22
Gulfport	75	72	43	43	0	0	29	26	1	2	18	11	43
Hattiesburg	32	20	8	8	0	0	12	11	0	1	6	7	7
Jackson	485	325	179	175	4	0	146	143	0	3	69	38	218
Laurel	77	16	2	2	0	0	14	13	1	0	3	7	6
McComb	5	6	2	2	0	0	4	2	0	2	2	2	2
Meridian	100	50	32	32	0	0	18	17	1	0	11	9	30
Natchez	81	33	14	14	0	0	19	19	0	0	15	7	11
Pascagoula	30	15	12	12	0	0	3	3	0	0	10	4	1
Tupelo	19	7	4	4	0	0	3	3	0	0	3	4	0
Vicksburg	142	65	26	26	0	0	39	38	0	1	15	6	44
Yazoo City	9	13	3	3	0	0	10	8	0	2	3	5	5

* Cities having 10,000 population and over according to 1960 census.

TABLE 5
DEATHS
MISSISSIPPI
1968

By Cause of Death, by Race, Residence Data

International Classification Number*	Cause of Death**	Total	White	Non- white
<u>000-999</u>	<u>ALL CAUSES</u>	<u>23,986</u>	<u>13,468</u>	<u>10,518</u>
<u>000-136</u>	<u>INFECTIVE AND PARASITIC DISEASES</u>	<u>337</u>	<u>130</u>	<u>207</u>
<u>000-009</u>	<u>Intestinal infectious diseases</u>	<u>128</u>	<u>34</u>	<u>94</u>
004,006	<u>Bacillary dysentery and amebiasis</u>	<u>14</u>	<u>5</u>	<u>9</u>
004	Bacillary dysentery	9	3	6
006	Amebiasis	5	2	3
005	Food poisoning (bacterial)	1	0	1
<u>008,009</u>	<u>Enteritis and other diarrheal diseases</u>	<u>113</u>	<u>29</u>	<u>84</u>
008	Enteritis due to other specified organism	6	3	3
009	Diarrheal disease	107	26	81
<u>010-019</u>	<u>Tuberculosis, all forms</u>	<u>89</u>	<u>37</u>	<u>52</u>
<u>010-012</u>	<u>Respiratory</u>	<u>78</u>	<u>33</u>	<u>45</u>
011	Pulmonary	75	31	44
012	Other respiratory	3	2	1
<u>013-018</u>	<u>Other forms</u>	<u>7</u>	<u>2</u>	<u>5</u>
013	Meninges and central nervous system	2	1	1
015	Bones and joints	5	1	4
019	Late effects of tuberculosis	4	2	2
<u>020-027</u>	<u>Zoonotic bacterial diseases</u>	<u>2</u>	<u>1</u>	<u>1</u>
023	Brucellosis	1	1	0
027	Other zoonotic bacterial diseases	1	0	1
<u>030-039</u>	<u>Other bacterial diseases</u>	<u>60</u>	<u>31</u>	<u>29</u>
031	Other diseases due to mycobacteria	1	1	0
032	Diphtheria	1	1	0
033	Whooping cough	1	0	1
034	Streptococcal sore throat and scarlet fever	1	1	0
035	Erysipelas	1	1	0
036	Meningococcal infection	5	5	0
037	Tetanus	1	0	1
038	Septicemia	46	22	24
039	Other bacterial diseases	3	0	3
<u>040-046</u>	<u>Poliomyelitis and other enterovirus diseases of central nervous system</u>	<u>2</u>	<u>1</u>	<u>1</u>
045	Aseptic meningitis due to enterovirus	1	0	1
046	Other enterovirus diseases of central nervous system	1	1	0
<u>050-057</u>	<u>Viral diseases accompanied by exanthem</u>	<u>4</u>	<u>1</u>	<u>3</u>
052	Chickenpox	3	1	2
054	Herpes simplex	1	0	1
<u>060-068</u>	<u>Arthropod-borne viral diseases</u>	<u>6</u>	<u>3</u>	<u>3</u>
<u>062-065</u>	<u>Viral encephalitis</u>	<u>6</u>	<u>3</u>	<u>3</u>
065	Viral encephalitis, unspecified	6	3	3

Table 5 - Page 2
Deaths - Mississippi - 1968

International Classification Number*	Cause of Death**	Total	White	Non- white
<u>070-079</u>	<u>Other viral diseases</u>	<u>26</u>	<u>15</u>	<u>11</u>
070	Infectious hepatitis	18	12	6
079	Other viral diseases	8	3	5
<u>080-089</u>	<u>Rickettsioses and other arthropod-borne diseases</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>090-099</u>	<u>Syphilis and other venereal diseases</u>	<u>13</u>	<u>2</u>	<u>11</u>
090-097	<u>Syphilis and its sequelae</u>	<u>12</u>	<u>2</u>	<u>10</u>
090	Congenital	1	0	1
093	Cardiovascular	8	1	7
094	Central nervous system	2	1	1
095	Other late, with symptoms	1	0	1
099	Other venereal diseases	1	0	1
<u>100-104</u>	<u>Other spirochetal diseases</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>110-117</u>	<u>Mycoses</u>	<u>3</u>	<u>3</u>	<u>0</u>
112	Moniliasis	1	1	0
115	Histoplasmosis	1	1	0
116	Blastomycosis	1	1	0
<u>120-129</u>	<u>Helminthiasis</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>130-136</u>	<u>Other infective and parasitic diseases</u>	<u>4</u>	<u>2</u>	<u>2</u>
135	Sarcoidosis	2	0	2
136	Other and unspecified infective and parasitic diseases	2	2	0
<u>140-239</u>	<u>NEOPLASMS</u>	<u>3,316</u>	<u>2,129</u>	<u>1,187</u>
<u>140-209</u>	<u>Malignant neoplasms</u>	<u>3,267</u>	<u>2,101</u>	<u>1,166</u>
140-149	<u>Buccal cavity and pharynx</u>	<u>72</u>	<u>54</u>	<u>18</u>
140	Lip	3	3	0
141	Tongue	9	7	2
142	Salivary gland	13	10	3
143	Gum	2	0	2
144	Floor of mouth	2	2	0
145	Other and unspecified parts of mouth	9	7	2
146	Oropharynx	5	3	2
147	Nasopharynx	6	5	1
148	Hypopharynx	4	3	1
149	Pharynx, unspecified	19	14	5
<u>150-159</u>	<u>Digestive organs and peritoneum</u>	<u>920</u>	<u>575</u>	<u>345</u>
150	Esophagus	48	25	23
151	Stomach	218	105	113
152	Small intestine, including duodenum	10	7	3
153	Large intestine, except rectum	283	206	77
154	Rectum and rectosigmoid junction	65	33	32
155	Liver and intrahepatic bile ducts, specified as primary	22	13	9
156	Gallbladder and bile ducts	28	26	2
157	Pancreas	217	147	70
158	Peritoneum and retroperitoneal tissue	9	6	3
159	Unspecified digestive organs	20	7	13
<u>160-163</u>	<u>Respiratory system</u>	<u>634</u>	<u>467</u>	<u>167</u>
160	Nose, nasal cavities, middle ear, and accessory sinuses	7	3	4
161	Larynx	22	16	6

Table 5 - Page 3
Deaths - Mississippi - 1968

International Classification Number*	Cause of Death**	Total	White	Non- white
162	Trachea, bronchus, and lung	602	447	155
163	Other and unspecified respiratory organs	3	1	2
<u>170-174</u>	<u>Bone, connective tissue, skin, and breast</u>	<u>359</u>	<u>244</u>	<u>115</u>
170	Bone	22	14	8
171	Connective and other soft tissue	11	7	4
172	Melanoma of skin	40	37	3
173	Skin, other	28	22	6
174	Breast	258	164	94
<u>180-189</u>	<u>Genitourinary organs</u>	<u>623</u>	<u>307</u>	<u>316</u>
<u>(180-187)</u>	<u>(Genital organs)</u>	<u>501</u>	<u>225</u>	<u>276</u>
180	Cervix uteri	111	34	77
182	Uterus, other	79	40	39
183	Ovary, fallopian tube, and broad ligament	68	43	25
184	Other and unspecified female genital organs	9	6	3
185	Prostate	223	99	124
186	Testis	4	3	1
187	Other and unspecified male genital organs	7	0	7
<u>(188,189)</u>	<u>(Urinary organs)</u>	<u>122</u>	<u>82</u>	<u>40</u>
188	Bladder	64	43	21
189	Other and unspecified urinary organs	58	39	19
<u>190-199</u>	<u>Other and unspecified sites</u>	<u>329</u>	<u>205</u>	<u>124</u>
190	Eye	3	2	1
191	Brain	75	61	14
192	Other parts of nervous system	19	14	5
193	Thyroid gland	10	5	5
194	Other endocrine glands	3	2	1
195	Ill-defined sites	35	19	16
196	Lymph nodes, secondary and unspecified	1	1	0
197	Respiratory and digestive systems, secondary	70	40	30
198	Other secondary	14	11	3
199	Site unspecified	99	50	49
<u>200-209</u>	<u>Lymphatic and hematopoietic tissue</u>	<u>330</u>	<u>249</u>	<u>81</u>
200	Lymphosarcoma and reticulum-cell sarcoma	66	58	8
201	Hodgkin's disease	34	27	7
202	Lymphoid tissue, other	17	13	4
203	Multiple myeloma	46	31	15
208	Polycythemia vera	5	5	0
209	Myelofibrosis	4	2	2
<u>(204-207)</u>	<u>(Leukemia)</u>	<u>158</u>	<u>113</u>	<u>45</u>
204	Lymphatic	63	42	21
205	Myeloid	52	40	12
206	Monocytic	6	4	2
207	Other and unspecified	37	27	10
<u>210-228</u>	<u>Benign neoplasms</u>	<u>27</u>	<u>17</u>	<u>10</u>
211	Benign neoplasm of other parts of digestive system	1	1	0
215	Benign neoplasm of muscular and connective tissue, other	1	0	1
218	Uterine fibroma	5	1	4
220	Benign neoplasm of ovary	1	1	0

Table 5 - Page 4
Deaths - Mississippi - 1968

International Classification Number*	Cause of Death**	Total	White	Non- white
221	Benign neoplasm of other female genital organs	1	1	0
223	Benign neoplasm of kidney and other urinary organs	1	1	0
225	Benign neoplasm of brain and other parts of nervous system	13	9	4
226	Benign neoplasm of endocrine glands	2	2	0
227	Hemangioma and lymphangioma	2	1	1
<u>230-239</u>	<u>Neoplasms of unspecified nature</u>	<u>22</u>	<u>11</u>	<u>11</u>
230	Digestive organs	2	2	0
231	Respiratory organs	2	1	1
234	Uterus	2	1	1
235	Ovary	1	1	0
237	Other genitourinary organs	1	0	1
238	Eye, brain, and other parts of nervous system	12	6	6
239	Other and unspecified organs	2	0	2
<u>240-279</u>	<u>ENDOCRINE, NUTRITIONAL, AND METABOLIC DISEASES</u>	<u>521</u>	<u>267</u>	<u>254</u>
<u>240-246</u>	<u>Diseases of thyroid gland</u>	<u>1</u>	<u>0</u>	<u>1</u>
242	Thyrotoxicosis with or without goiter	1	0	1
<u>250-258</u>	<u>Diseases of other endocrine glands</u>	<u>449</u>	<u>231</u>	<u>218</u>
250	Diabetes mellitus	436	221	215
251	Disorders of pancreatic internal secretion other than diabetes mellitus	4	2	2
255	Diseases of adrenal glands	7	6	1
258	Polyglandular dysfunction and other diseases of endocrine glands	2	2	0
<u>260-269</u>	<u>Avitaminoses and other nutritional deficiency</u>	<u>39</u>	<u>15</u>	<u>24</u>
262	Niacin deficiency	1	1	0
266	Other vitamin deficiency states	1	0	1
267	Protein malnutrition	1	1	0
268	Nutritional marasmus	3	2	1
269	Other nutritional deficiency	33	11	22
<u>270-279</u>	<u>Other metabolic diseases</u>	<u>32</u>	<u>21</u>	<u>11</u>
272	Congenital disorders of lipid metabolism	1	1	0
273	Other and unspecified congenital disorders of metabolism	16	12	4
274	Gout	1	1	0
276	Amyloidosis	2	1	1
277	Obesity not specified as of endocrine origin	11	5	6
279	Other and unspecified metabolic diseases	1	1	0
<u>280-289</u>	<u>DISEASES OF THE BLOOD AND BLOOD-FORMING ORGANS</u>	<u>68</u>	<u>25</u>	<u>43</u>
<u>280-285</u>	<u>Anemias</u>	<u>56</u>	<u>15</u>	<u>41</u>
280	Iron deficiency	4	2	2
281	Other deficiency	1	0	1
282	Hereditary hemolytic	22	0	22
284	Aplastic	5	3	2
285	Other and unspecified	24	10	14

Table 5 - Page 5
Deaths - Mississippi - 1968

International Classification Number*	Cause of Death**	Total	White	Non- white
286	Coagulation defects	2	2	0
287	Purpura and other hemorrhagic conditions	5	4	1
289	Other diseases of blood and blood-forming organs	5	4	1
<u>290-315</u>	<u>MENTAL DISORDERS</u>	<u>49</u>	<u>25</u>	<u>24</u>
<u>290-299</u>	<u>Psychoses</u>	<u>10</u>	<u>6</u>	<u>4</u>
290	Senile and presenile dementia	3	2	1
291	Alcoholic psychosis	4	2	2
295	Schizophrenia	2	1	1
299	Unspecified psychosis	1	1	0
<u>300-309</u>	<u>Neuroses, personality disorders, and other non- psychotic mental disorders</u>	<u>39</u>	<u>19</u>	<u>20</u>
300	Neuroses	1	1	0
303	Alcoholism	30	11	19
304	Drug dependence	1	1	0
309	Mental disorders not specified as psychotic associated with physical conditions	7	6	1
<u>310-315</u>	<u>Mental retardation</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>320-389</u>	<u>DISEASES OF THE NERVOUS SYSTEM AND SENSE ORGANS</u>	<u>167</u>	<u>89</u>	<u>78</u>
<u>320-324</u>	<u>Inflammatory diseases of central nervous system</u>	<u>40</u>	<u>13</u>	<u>27</u>
320	Meningitis	32	8	24
322	Intracranial and intraspinal abscess	2	1	1
323	Encephalitis, myelitis, and encephalomyelitis	5	3	2
324	Late effects of intracranial abscess or pyogenic infection	1	1	0
<u>330-333</u>	<u>Hereditary and familial diseases of nervous system</u>	<u>9</u>	<u>9</u>	<u>0</u>
330	Hereditary neuromuscular disorders	5	5	0
331	Hereditary diseases of the striatopallidal system	4	4	0
<u>340-349</u>	<u>Other diseases of central nervous system</u>	<u>110</u>	<u>62</u>	<u>48</u>
340	Multiple sclerosis	11	6	5
341	Other demyelinating diseases of central nervous system	2	1	1
342	Paralysis agitans	23	16	7
343	Cerebral spastic infantile paralysis	10	6	4
344	Other cerebral paralysis	9	3	6
345	Epilepsy	23	8	15
347	Other diseases of brain	18	13	5
348	Motor neurone disease	12	9	3
349	Other diseases of spinal cord	2	0	2
<u>350-358</u>	<u>Diseases of nerves and peripheral ganglia</u>	<u>4</u>	<u>3</u>	<u>1</u>
354	Polyneuritis and polyradiculitis	2	2	0
355	Other and unspecified forms of neuralgia and neuritis	1	1	0
358	Diseases of peripheral autonomic nervous system	1	0	1

Table 5 - Page 6
Deaths - Mississippi - 1968

International Classification Number*	Cause of Death**	Total	White	Non- white
<u>360-369</u>	<u>Inflammatory diseases of the eye</u>	<u>2</u>	<u>1</u>	<u>1</u>
365	Choroiditis	1	0	1
366	Other inflammation of uveal tract	1	1	0
<u>370-379</u>	<u>Other diseases and conditions of eye</u>	<u>1</u>	<u>0</u>	<u>1</u>
375	Glaucoma	1	0	1
<u>380-389</u>	<u>Diseases of the ear and mastoid process</u>	<u>1</u>	<u>1</u>	<u>0</u>
<u>381-383</u>	<u>Otitis media and mastoiditis</u>	<u>1</u>	<u>1</u>	<u>0</u>
381	Otitis media without mention of mastoiditis	1	1	0
<u>390-458</u>	<u>DISEASES OF THE CIRCULATORY SYSTEM</u>	<u>12,000</u>	<u>7,207</u>	<u>4,793</u>
<u>390-392</u>	<u>Active rheumatic fever</u>	<u>5</u>	<u>3</u>	<u>2</u>
391	Rheumatic fever with heart involvement	5	3	2
<u>393-398</u>	<u>Chronic rheumatic heart disease</u>	<u>93</u>	<u>58</u>	<u>35</u>
393	Diseases of pericardium	1	0	1
394	Diseases of mitral valve	26	18	8
395	Diseases of aortic valve	26	19	7
396	Diseases of mitral and aortic valve	8	4	4
397	Diseases of other endocardial structures	6	3	3
398	Other heart disease, specified as rheumatic	26	14	12
<u>400-404</u>	<u>Hypertensive disease</u>	<u>505</u>	<u>169</u>	<u>336</u>
<u>400,401,</u>				
403	Hypertension	209	78	131
400	Malignant hypertension	42	8	34
401	Essential benign hypertension	49	15	34
403	Hypertensive renal disease	118	55	63
<u>402,404</u>	<u>Hypertensive heart disease with or without renal disease</u>	<u>296</u>	<u>91</u>	<u>205</u>
402	Hypertensive heart disease	195	55	140
404	Hypertensive heart and renal disease	101	36	65
<u>410-414</u>	<u>Ischemic heart disease</u>	<u>6,818</u>	<u>4,496</u>	<u>2,322</u>
410	Acute myocardial infarction	4,039	2,976	1,063
411	Other acute and subacute forms of ischemic heart disease	54	27	27
412	Chronic ischemic heart disease	2,709	1,488	1,221
413	Angina pectoris	16	5	11
<u>420-429</u>	<u>Other forms of heart disease</u>	<u>757</u>	<u>376</u>	<u>381</u>
421	Acute and subacute endocarditis	4	2	2
422	Acute myocarditis	4	2	2
423	Chronic disease of pericardium, nonrheumatic	3	2	1
424	Chronic disease of endocardium	25	14	11
425	Cardiomyopathy	4	1	3
426	Pulmonary heart disease	6	1	5
427	Symptomatic heart disease	273	148	125
428	Other myocardial insufficiency	141	73	68
429	Ill-defined heart disease	297	133	164
<u>430-438</u>	<u>Cerebrovascular disease</u>	<u>3,115</u>	<u>1,631</u>	<u>1,484</u>
430	Subarachnoid hemorrhage	110	62	48
431	Cerebral hemorrhage	693	282	411
432	Occlusion of precerebral arteries	39	30	9
433	Cerebral thrombosis	702	432	270

Table 5 - Page 7
Deaths - Mississippi - 1968

International Classification Number*	Cause of Death**	Total	White	Non- white
434	Cerebral embolism	17	6	11
435	Transient cerebral ischemia	1	1	0
436	Acute but ill-defined cerebrovascular disease	1,221	589	632
437	Generalized ischemic cerebrovascular disease	307	215	92
438	Other and ill-defined cerebrovascular disease	25	14	11
<u>440-448</u>	<u>Diseases of arteries, arterioles, and capillaries</u>	<u>545</u>	<u>374</u>	<u>171</u>
440	Arteriosclerosis	316	211	105
<u>441-448</u>	<u>Other diseases of arteries, arterioles, and capillaries</u>	<u>229</u>	<u>163</u>	<u>66</u>
441	Aortic aneurysm (nonsyphilitic)	110	83	27
442	Other aneurysm	11	8	3
443	Other peripheral vascular disease	6	4	2
444	Arterial embolism and thrombosis	60	48	12
445	Gangrene	33	13	20
446	Polyarteritis nodosa and allied conditions	7	6	1
447	Other diseases of arteries and arterioles	2	1	1
<u>450-458</u>	<u>Diseases of veins and lymphatics, and other diseases of circulatory system</u>	<u>162</u>	<u>100</u>	<u>62</u>
<u>450-453</u>	<u>Venous thrombosis and embolism</u>	<u>139</u>	<u>91</u>	<u>48</u>
450	Pulmonary embolism and infarction	79	55	24
451	Phlebitis and thrombophlebitis	38	24	14
453	Other venous embolism and thrombosis	22	12	10
<u>454-458</u>	<u>Other diseases of the circulatory system</u>	<u>23</u>	<u>9</u>	<u>14</u>
454	Varicose veins of lower extremities	3	1	2
455	Hemorrhoids	2	2	0
456	Varicose veins of other sites	1	1	0
458	Other diseases of circulatory system	17	5	12
<u>460-519</u>	<u>DISEASES OF THE RESPIRATORY SYSTEM</u>	<u>1,282</u>	<u>738</u>	<u>544</u>
<u>460-466</u>	<u>Acute respiratory infections, except influenza</u>	<u>46</u>	<u>14</u>	<u>32</u>
460	Acute nasopharyngitis (common cold)	16	2	14
462	Acute pharyngitis	1	0	1
463	Acute tonsillitis	1	1	0
464	Acute laryngitis and tracheitis	1	1	0
465	Acute upper respiratory infection of multiple or unspecified sites	11	5	6
466	Acute bronchitis and bronchiolitis	16	5	11
<u>470-474</u>	<u>Influenza</u>	<u>96</u>	<u>45</u>	<u>51</u>
470	Unqualified	34	12	22
471	With pneumonia	59	31	28
472	With other respiratory manifestations	2	2	0
474	With nervous manifestations	1	0	1
<u>480-486</u>	<u>Pneumonia</u>	<u>729</u>	<u>369</u>	<u>360</u>
480	Viral	12	8	4
<u>481,482</u>	<u>Bacterial</u>	<u>108</u>	<u>48</u>	<u>60</u>
481	Pneumococcal	93	39	54
482	Other bacterial	15	9	6
484	Acute interstitial, organism unspecified	9	4	5
485	Broncho, organism unspecified	228	122	106
486	Unspecified	372	187	185

Table 5 - Page 8
Deaths - Mississippi - 1968

International Classification Number*	Cause of Death**	Total	White	Non- white
<u>490-493</u>	<u>Bronchitis, emphysema, and asthma</u>	<u>303</u>	<u>242</u>	<u>61</u>
490,491	Chronic and unqualified bronchitis	38	31	7
490	Unqualified	12	6	6
491	Chronic	26	25	1
492	Emphysema	213	190	23
493	Asthma	52	21	31
<u>500-508</u>	<u>Other diseases of upper respiratory tract</u>	<u>3</u>	<u>0</u>	<u>3</u>
503	Chronic sinusitis	1	0	1
508	Other diseases of upper respiratory tract	2	0	2
<u>510-519</u>	<u>Other diseases of respiratory system</u>	<u>105</u>	<u>68</u>	<u>37</u>
510	Empyema	1	1	0
511	Pleurisy	2	0	2
512	Spontaneous pneumothorax	1	1	0
513	Abscess of lung	9	3	6
514	Pulmonary congestion and pypostasis	15	6	9
517	Other chronic interstitial pneumonia	33	30	3
518	Bronchiectasis	13	11	2
519	Other diseases of respiratory system	31	16	15
<u>520-577</u>	<u>DISEASES OF THE DIGESTIVE SYSTEM</u>	<u>566</u>	<u>378</u>	<u>188</u>
<u>520-529</u>	<u>Diseases of oral cavity, salivary glands, and jaws</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>530-537</u>	<u>Diseases of esophagus, stomach, and duodenum</u>	<u>110</u>	<u>76</u>	<u>34</u>
530	Diseases of esophagus	8	6	2
<u>531-533</u>	<u>Peptic ulcer</u>	<u>82</u>	<u>59</u>	<u>23</u>
531	Ulcer of stomach	31	17	14
532	Ulcer of duodenum	33	27	6
533	Peptic ulcer, site unspecified	18	15	3
534	Gastrojejunal ulcer	4	2	2
535	Gastritis and duodenitis	4	3	1
536	Disorders of function of stomach	6	3	3
537	Other diseases of stomach and duodenum	6	3	3
<u>540-543</u>	<u>Appendicitis</u>	<u>15</u>	<u>7</u>	<u>8</u>
540	Acute appendicitis	12	6	6
541	Appendicitis, unqualified	2	1	1
543	Other diseases of appendix	1	0	1
<u>550-553</u>	<u>Hernia of abdominal cavity</u>	<u>29</u>	<u>13</u>	<u>16</u>
550	Inguinal, without mention of obstruction	3	3	0
551	Other, of abdominal cavity without mention of obstruction	14	5	9
552	Inguinal, with obstruction	3	0	3
553	Other, of abdominal cavity with obstruction	9	5	4
<u>560-569</u>	<u>Other diseases of intestine and peritoneum</u>	<u>154</u>	<u>97</u>	<u>57</u>
560	Intestinal obstruction without mention of hernia	59	30	29
562	Diverticula of intestine	24	21	3
563	Chronic enteritis and ulcerative colitis	11	8	3
566	Abscess of anal and rectal regions	1	0	1
567	Peritonitis	6	4	2
569	Other diseases of intestines and peritoneum	53	34	19
<u>570-577</u>	<u>Diseases of liver, gallbladder, and pancreas</u>	<u>258</u>	<u>185</u>	<u>73</u>
570	Acute and subacute necrosis of liver	4	0	4
571	Cirrhosis of liver	146	113	33

Table 5 - Page 9
Deaths - Mississippi - 1968

International Classification Number*	Cause of Death**	Total	White	Non- white
572	Suppurative hepatitis and liver abscess	3	0	3
573	Other diseases of liver	17	9	8
<u>574,575</u>	<u>Cholelithiasis, cholecystitis, and cholangitis</u>	<u>42</u>	<u>29</u>	<u>13</u>
574	Cholelithiasis	20	16	4
575	Cholecystitis and cholangitis, without mention of calculus	22	13	9
576	Other diseases of gallbladder and biliary ducts	7	5	2
577	Diseases of pancreas	39	29	10
<u>580-629</u>	<u>DISEASES OF THE GENITOURINARY SYSTEM</u>	<u>518</u>	<u>252</u>	<u>266</u>
<u>580-584</u>	<u>Nephritis and nephrosis</u>	<u>181</u>	<u>58</u>	<u>123</u>
<u>580,581</u>	<u>Acute nephritis and nephrotic syndrome</u>	<u>21</u>	<u>6</u>	<u>15</u>
580	Acute nephritis	8	3	5
581	Nephrotic syndrome	13	3	10
<u>582-584</u>	<u>Chronic and unqualified nephritis and renal sclerosis</u>	<u>160</u>	<u>52</u>	<u>108</u>
582	Chronic nephritis	138	44	94
583	Nephritis, unqualified	22	8	14
<u>590-599</u>	<u>Other diseases of urinary system</u>	<u>273</u>	<u>169</u>	<u>104</u>
590	Infections of kidney	199	133	66
591	Hydronephrosis	1	1	0
<u>592,594</u>	<u>Calculus of urinary system</u>	<u>7</u>	<u>7</u>	<u>0</u>
592	Of kidney and ureter	7	7	0
593	Other diseases of kidney and ureter	22	9	13
595	Cystitis	3	1	2
596	Other diseases of bladder	4	1	3
597	Urethritis (nonvenereal)	1	1	0
598	Stricture of urethra	7	1	6
599	Other diseases of urinary tract	29	15	14
<u>600-607</u>	<u>Diseases of male genital organs</u>	<u>55</u>	<u>23</u>	<u>32</u>
600	Hyperplasia of prostate	49	21	28
601	Prostatitis	5	2	3
602	Other diseases of prostate	1	0	1
<u>610-616</u>	<u>Diseases of breast, ovary, fallopian tube, and parametrium</u>	<u>5</u>	<u>1</u>	<u>4</u>
614	Salpingitis and oophoritis, unqualified	1	1	0
615	Other diseases of ovary and fallopian tube	1	0	1
616	Diseases of parametrium and pelvic peritoneum (female)	3	0	3
<u>620-629</u>	<u>Diseases of uterus and other female genital organs</u>	<u>4</u>	<u>1</u>	<u>3</u>
622	Infective diseases of uterus (except cervix), vagina, and vulva	1	0	1
623	Uterovaginal prolapse	1	0	1
629	Other diseases of female genital organs	2	1	1
<u>630-678</u>	<u>COMPLICATIONS OF PREGNANCY, CHILDBIRTH, AND THE PUERPERIUM</u>	<u>21</u>	<u>3</u>	<u>18</u>
<u>630-634</u>	<u>Complications of pregnancy</u>	<u>3</u>	<u>1</u>	<u>2</u>
632	Hemorrhage of pregnancy	2	1	1

Table 5 - Page 10
Deaths - Mississippi - 1968

International Classification Number*	Cause of Death**	Total	White	Non- white
634	Other complications of pregnancy	1	0	1
<u>635-639</u>	<u>Urinary infections and toxemias of pregnancy and puerperium</u>	<u>8</u>	<u>1</u>	<u>7</u>
<u>636-639</u>	<u>Toxemias of pregnancy and the puerperium, except abortion with toxemia</u>	<u>8</u>	<u>1</u>	<u>7</u>
637	Pre-eclampsia, eclampsia, and toxemia, unspecified	8	1	7
<u>640-645</u>	<u>Abortion</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>650-662</u>	<u>Delivery</u>	<u>7</u>	<u>0</u>	<u>7</u>
651	Delivery complicated by placenta previa or antepartum hemorrhage	1	0	1
653	Delivery complicated by other postpartum hemorrhage	2	0	2
655	Delivery complicated by fetopelvic disproportion	1	0	1
660	Delivery with other obstetrical trauma	1	0	1
661	Delivery with other complications	2	0	2
<u>670-678</u>	<u>Complications of the puerperium</u>	<u>3</u>	<u>1</u>	<u>2</u>
<u>670,671,</u>				
<u>673</u>	<u>Sepsis of childbirth and the puerperium</u>	<u>2</u>	<u>1</u>	<u>1</u>
673	Puerperal pulmonary embolism	2	1	1
675	Puerperal blood dyscrasias	1	0	1
<u>680-709</u>	<u>DISEASES OF THE SKIN AND SUBCUTANEOUS TISSUE</u>	<u>26</u>	<u>14</u>	<u>12</u>
<u>680-686</u>	<u>Infections of skin and subcutaneous tissue</u>	<u>8</u>	<u>6</u>	<u>2</u>
682	Other cellulitis and abscess	5	3	2
686	Other local infections of skin and subcutaneous tissue	3	3	0
<u>690-698</u>	<u>Other inflammatory conditions of skin and subcutaneous tissue</u>	<u>10</u>	<u>3</u>	<u>7</u>
692	Other eczema and dermatitis	1	0	1
694	Pemphigus	2	1	1
695	Erythematous conditions	7	2	5
<u>700-709</u>	<u>Other diseases of skin and subcutaneous tissue</u>	<u>8</u>	<u>5</u>	<u>3</u>
707	Chronic ulcer of skin	7	5	2
709	Other diseases of skin	1	0	1
<u>710-738</u>	<u>DISEASES OF THE MUSCULOSKELETAL SYSTEM AND CONNECTIVE TISSUE</u>	<u>60</u>	<u>38</u>	<u>22</u>
<u>710-718</u>	<u>Arthritis and rheumatism, except rheumatic fever</u>	<u>35</u>	<u>21</u>	<u>14</u>
<u>710-715</u>	<u>Arthritis and spondylitis</u>	<u>33</u>	<u>19</u>	<u>14</u>
712	Rheumatoid arthritis and allied conditions	20	13	7
713	Osteoarthritis and allied conditions	5	2	3
715	Arthritis, unspecified	8	4	4
<u>716-718</u>	<u>Nonarticular rheumatism and rheumatism unspecified</u>	<u>2</u>	<u>2</u>	<u>0</u>
716	Polymyositis and dermatomyositis	1	1	0
718	Rheumatism, unspecified	1	1	0
<u>720-729</u>	<u>Osteomyelitis and other diseases of bone and joint</u>	<u>4</u>	<u>4</u>	<u>0</u>

Table 5 - Page 11
Deaths - Mississippi - 1968

International Classification Number*	Cause of Death**	Total	White	Non- white
721	Osteitis deformans	2	2	0
723	Other diseases of bone	2	2	0
<u>730-738</u>	<u>Other diseases of musculoskeletal system</u>	<u>21</u>	<u>13</u>	<u>8</u>
733	Other diseases of muscle, tendon, and fascia	6	3	3
734	Diffuse diseases of connective tissue	11	6	5
735	Curvature of spine	4	4	0
<u>740-759</u>	<u>CONGENITAL ANOMALIES</u>	<u>205</u>	<u>135</u>	<u>70</u>
<u>740-743</u>	<u>Nervous system</u>	<u>48</u>	<u>39</u>	<u>9</u>
740	Anencephalus	11	10	1
741	Spina bifida	17	16	1
742	Congenital hydrocephalus	11	7	4
743	Other congenital anomalies of nervous system	9	6	3
<u>746,747</u>	<u>Circulatory system</u>	<u>99</u>	<u>63</u>	<u>36</u>
746	Heart	85	51	34
747	Other of circulatory system	14	12	2
748	Respiratory system	5	2	3
<u>749-751</u>	<u>Digestive system</u>	<u>19</u>	<u>10</u>	<u>9</u>
749	Cleft palate and cleft lip	1	0	1
750	Other congenital anomalies of upper ali- mentary tract	7	3	4
751	Other congenital anomalies of digestive system	11	7	4
<u>752,753</u>	<u>Genitourinary system</u>	<u>8</u>	<u>6</u>	<u>2</u>
752	Genital organs	1	1	0
753	Urinary system	7	5	2
<u>754-756</u>	<u>Musculoskeletal system</u>	<u>4</u>	<u>4</u>	<u>0</u>
756	Other congenital anomalies of musculo- skeletal system	4	4	0
757	Skin, hair, and nails	1	1	0
758	Other and unspecified	6	1	5
759	Congenital syndromes affecting multiple systems	15	9	6
<u>760-778</u>	<u>CERTAIN CAUSES OF MORTALITY IN EARLY INFANCY</u>	<u>791</u>	<u>320</u>	<u>471</u>
760	Chronic circulatory and genitourinary diseases in mother	1	0	1
761	Other maternal conditions unrelated to pregnancy	3	0	3
762	Toxemia of pregnancy	6	3	3
763	Maternal ante- and intrapartum infection	2	1	1
<u>764-768</u>	<u>Difficult labor</u>	<u>12</u>	<u>4</u>	<u>8</u>
764	With abnormality of bones, organs, or tissues of pelvis	1	0	1
766	With malposition of fetus	7	2	5
768	With other and unspecified complications	4	2	2
769	Other complications of pregnancy and childbirth	43	21	22
<u>770,771</u>	<u>Conditions of placenta and cord</u>	<u>47</u>	<u>22</u>	<u>25</u>
770	Of placenta	39	17	22
771	Of umbilical cord	8	5	3
772	Birth injury without mention of cause	31	15	16

Table 5 - Page 12
Deaths - Mississippi - 1968

International Classification Number*	Cause of Death**	Total	White	Non- white
<u>774,775</u>	<u>Hemolytic disease of newborn</u>	<u>8</u>	<u>5</u>	<u>3</u>
774	With kernicterus	3	1	2
775	Without mention of kernicterus	5	4	1
776	Anoxic and hypoxic conditions not elsewhere classifiable	323	163	160
777	Immaturity, unqualified	283	76	207
778	Other conditions of fetus or newborn	32	10	22
<u>780-796</u>	<u>SYMPTOMS AND ILL-DEFINED CONDITIONS</u>	<u>1,858</u>	<u>487</u>	<u>1,371</u>
<u>780-789</u>	<u>Symptoms referable to systems or organs</u>	<u>83</u>	<u>32</u>	<u>51</u>
780	Certain symptoms referable to nervous system and special senses	5	2	3
781	Other symptoms referable to nervous system and special senses	1	1	0
782	Symptoms referable to cardiovascular and lymphatic system	56	23	33
783	Symptoms referable to respiratory system	10	5	5
784	Symptoms referable to upper gastrointestinal tract	2	0	2
785	Symptoms referable to abdomen and lower gastrointestinal tract	4	1	3
788	Other general symptoms	5	0	5
<u>790-796</u>	<u>Senility and ill-defined diseases</u>	<u>1,775</u>	<u>455</u>	<u>1,320</u>
791	Headache	1	0	1
792	Uremia	10	2	8
794	Senility without mention of psychosis	68	11	57
795	Sudden death (cause unknown)	18	11	7
796	Other ill-defined and unknown causes	1,678	431	1,247
<u>800-999</u>	<u>ACCIDENTS, POISONINGS, AND VIOLENCE (EXTERNAL CAUSE)</u>	<u>2,201</u>	<u>1,231</u>	<u>970</u>
<u>800-949</u>	<u>Accidents</u>	<u>1,723</u>	<u>999</u>	<u>724</u>
<u>800-807</u>	<u>Railway</u>	<u>10</u>	<u>5</u>	<u>5</u>
804	Fall in, on, or from train	1	1	0
805	Hit by rolling stock	9	4	5
<u>810-823</u>	<u>Motor vehicle</u>	<u>821</u>	<u>542</u>	<u>279</u>
<u>(810-819)</u>	<u>(Traffic)</u>	<u>804</u>	<u>533</u>	<u>271</u>
810	Involving collision with train	29	20	9
812	Involving collision with another motor vehicle	310	236	74
813	Involving collision with other vehicle	9	6	3
814	Involving collision with pedestrian	91	32	59
815	Other involving collision	32	23	9
816	Noncollision due to loss of control	234	155	79
818	Other noncollision	11	6	5
819	Of unspecified nature	88	55	33
<u>(820-823)</u>	<u>(Nontraffic)</u>	<u>17</u>	<u>9</u>	<u>8</u>
820	Involving collision with moving object	11	6	5
823	Of other and unspecified nature	6	3	3

Table 5 - Page 13
Deaths - Mississippi - 1968

International Classification Number*	Cause of Death**	Total	White	Non- white
<u>825-827</u>	<u>Other road vehicle</u>	<u>4</u>	<u>3</u>	<u>1</u>
827	Other nonmotor road vehicle accident	4	3	1
<u>830-838</u>	<u>Water transport</u>	<u>35</u>	<u>27</u>	<u>8</u>
830	Accident to watercraft causing submersion	11	8	3
832	Other submersion or drowning in water transport	19	14	5
837	Explosion, fire, burning in water transport	2	2	0
838	Other and unspecified water transport accident	3	3	0
<u>840-845</u>	<u>Air and space transport</u>	<u>12</u>	<u>12</u>	<u>0</u>
841	Accident to powered aircraft, other and unspecified	11	11	0
843	Fall in, on, or from aircraft	1	1	0
<u>850-877</u>	<u>Poisoning</u>	<u>26</u>	<u>15</u>	<u>11</u>
(850-859)	(By drugs and medicaments)	<u>5</u>	<u>4</u>	<u>1</u>
853	Analgesics and antipyretics	1	0	1
854	Other sedatives and hypnotics	3	3	0
859	Other and unspecified drugs and medicaments	1	1	0
(860-869)	(By other solid and liquid substances)	<u>11</u>	<u>7</u>	<u>4</u>
860	Alcohol	3	1	2
861	Cleansing and polishing agents	1	1	0
864	Petroleum products and other solvents	1	1	0
865	Pesticides, fertilizers, or plant food	1	0	1
869	Other and unspecified solid and liquid substances	5	4	1
(870-877)	(By gases and vapors)	<u>10</u>	<u>4</u>	<u>6</u>
870	Gas distributed by pipeline	2	1	1
873	Motor vehicle exhaust gas	6	1	5
874	Carbon monoxide from incomplete combustion of domestic fuels	1	1	0
877	Unspecified gases and vapors	1	1	0
<u>880-887</u>	<u>Falls</u>	<u>157</u>	<u>115</u>	<u>42</u>
880	On or from stairs and steps	7	4	3
881	On or from ladders or scaffolding	4	4	0
882	From or out of building or other structure	10	5	5
884	Other from one level to another	17	12	5
885	On same level from slipping, stumbling, or tripping	9	6	3
887	Other and unspecified fall	110	84	26
<u>890-899</u>	<u>Caused by fires and flames</u>	<u>208</u>	<u>68</u>	<u>140</u>
890	Conflagration in private dwelling	113	26	87
891	Conflagration in other building or structure	4	1	3
892	Conflagration not in building or structure	3	1	2
893	Ignition of clothing	36	14	22
894	Ignition of highly inflammable material	6	5	1
895	Controlled fire in private dwelling	20	9	11
897	Controlled fire not in building or structure	2	2	0
898	Other specified fires and flames	8	6	2
899	Unspecified fire	16	4	12
<u>900-909</u>	<u>Due to natural and environmental factors</u>	<u>25</u>	<u>7</u>	<u>18</u>
900	Excessive heat	3	2	1
901	Excessive cold	9	0	9

Table 5 - Page 14
Deaths - Mississippi - 1968

International Classification Number*	Cause of Death**	Total	White	Non- white
904	Hunger, thirst, exposure, and neglect	5	1	4
905	Bites and stings of venomous animals and insects	1	1	0
906	Other accidents caused by animals	2	2	0
907	Lightning	4	1	3
908	Cataclysm	1	0	1
<u>910-929</u>	<u>Other accidents</u>	<u>407</u>	<u>192</u>	<u>215</u>
910	Drowning and submersion	111	42	69
911	Inhalation and ingestion of food causing obstruction or suffocation	51	29	22
912	Inhalation and ingestion of other object causing obstruction or suffocation	4	2	2
913	Mechanical suffocation	36	14	22
915	Foreign body entering other orifice	1	1	0
916	Struck by falling object	32	20	12
917	Struck against or struck by objects	18	7	11
918	Caught in or between objects	11	3	8
920	Caused by cutting or piercing instruments	2	1	1
921	Caused by explosion of pressure vessel	2	1	1
922	Caused by firearm missiles	73	39	34
923	Caused by explosive material	10	4	6
924	Caused by hot substance, corrosive liquid, and steam	2	2	0
925	Caused by electric current	18	9	9
928	Machinery accidents not elsewhere classifiable	17	9	8
929	Other and unspecified accidents	19	9	10
<u>930-936</u>	<u>Surgical and medical complications and mis- adventures</u>	<u>8</u>	<u>6</u>	<u>2</u>
930	In operative therapeutic procedures	2	1	1
931	In other and unspecified therapeutic pro- cedures	3	3	0
932	In diagnostic procedures	3	2	1
<u>940-949</u>	<u>Late effects of accidental injury</u>	<u>10</u>	<u>7</u>	<u>3</u>
940	Of motor vehicle accident	3	2	1
943	Of accidental fall	5	4	1
946	Of other accident	2	1	1
<u>950-959</u>	<u>Suicide and self-inflicted injury</u>	<u>171</u>	<u>148</u>	<u>23</u>
<u>950-952</u>	<u>By poisoning</u>	<u>14</u>	<u>13</u>	<u>1</u>
950	By solid or liquid substances	10	9	1
952	By other gases	4	4	0
953	By hanging, strangulation, and suffocation	11	8	3
954	By submersion (drowning)	3	3	0
955	By firearms and explosives	140	121	19
956	By cutting and piercing instruments	2	2	0
957	By jumping from high place	1	1	0
<u>960-978</u>	<u>Homicide and injury purposely inflicted by other persons; legal intervention</u>	<u>256</u>	<u>60</u>	<u>196</u>
<u>960-969</u>	<u>Homicide and injury purposely inflicted by other persons</u>	<u>252</u>	<u>59</u>	<u>193</u>
960	Fight, brawl, rape	2	1	1

Table 5 - Page 15
Deaths - Mississippi - 1968

International Classification Number*	Cause of Death**	Total	White	Non- white
962	Assault by poisoning	1	1	0
963	Assault by hanging and strangulation	1	0	1
964	Assault by submersion (drowning)	1	0	1
965	Assault by firearms and explosives	188	46	142
966	Assault by cutting and piercing instruments	39	6	33
968	Assault by other and unspecified means	20	5	15
<u>970-978</u>	<u>Legal intervention</u>	<u>4</u>	<u>1</u>	<u>3</u>
970	Injury by firearms	4	1	3
<u>980-989</u>	<u>Injury undetermined whether accidentally or purposely inflicted</u>	<u>51</u>	<u>24</u>	<u>27</u>
980	Poisoning by solid or liquid substances	6	5	1
982	Poisoning by other gases	1	1	0
983	Hanging and strangulation	2	1	1
984	Submersion (drowning)	3	2	1
985	Injury by firearms and explosives	22	12	10
986	Injury by cutting and piercing instruments	7	0	7
988	Injury by other and unspecified means	9	2	7
989	Late effect of injury	1	1	0
<u>990-999</u>	<u>Injury resulting from operations of war</u>	<u>0</u>	<u>0</u>	<u>0</u>

* International Classification numbers are those of the Eighth Revision International Classification of Diseases, Adapted for Use in the United States.

** When cause of death and International Classification number are omitted, no deaths were reported.

TABLE 6
DEATHS
MISSISSIPPI
1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

County	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
<u>STATE</u>	<u>23,647</u>	<u>13,261</u>	<u>10,386</u>	<u>23,986</u>	<u>13,468</u>	<u>10,518</u>	<u>10.2</u>	<u>9.3</u>	<u>11.7</u>
Adams	493	203	290	368	139	229	10.0	7.3	12.9
Alcorn	294	248	46	324	273	51	11.7	11.1	16.7
Amite	70	25	45	158	83	75	10.4	11.8	9.2
Attala	228	130	98	254	141	113	11.2	10.7	11.9
Benton	29	12	17	86	56	30	11.7	13.3	9.5
Bolivar	568	118	450	564	151	413	10.3	7.6	11.9
Calhoun	130	105	25	183	155	28	11.6	12.8	7.5
Carroll	47	10	37	130	62	68	12.2	11.5	12.9
Chickasaw	198	137	61	219	149	70	11.4	12.2	10.1
Choctaw	115	72	43	122	87	35	14.2	13.7	15.4
Claiborne	96	23	73	123	34	89	10.8	10.5	10.9
Clarke	113	68	45	193	122	71	12.4	12.2	12.8
Clay	191	90	101	207	95	112	11.4	10.6	12.2
Coahoma	582	224	358	555	184	371	11.9	11.2	12.3
Copiah	251	115	136	318	163	155	12.1	11.8	12.3
Covington	103	69	34	166	117	49	11.6	11.6	11.4
DeSoto	144	59	85	279	148	131	7.6	6.6	9.2
Forrest	728	504	224	650	409	241	10.6	9.0	15.1
Franklin	86	59	27	114	78	36	12.2	12.5	11.8
George	77	65	12	95	79	16	8.1	7.6	11.9
Greene	65	33	32	82	50	32	10.1	7.8	18.0
Grenada	253	148	105	220	110	110	11.0	10.2	11.9
Hancock	150	126	24	170	145	25	8.1	8.2	8.0
Harrison	1,317	1,033	284	1,110	863	247	7.9	7.2	11.1
Hinds	2,724	1,532	1,192	1,797	1,000	797	8.3	7.7	9.2
Holmes	311	84	227	356	107	249	12.9	13.0	12.9
Humphreys	147	33	114	206	56	150	12.3	10.3	13.3
Issaquena	16	2	14	32	9	23	11.1	8.6	12.6
Itawamba	80	77	3	145	136	9	7.6	7.5	11.0
Jackson	430	310	120	468	346	122	5.7	5.1	9.0
Jasper	158	74	84	200	104	96	12.0	11.6	12.4
Jefferson	71	15	56	121	40	81	11.3	12.1	10.9
Jeff Davis	139	67	72	158	71	87	11.1	10.6	11.5
Jones	750	466	284	600	419	181	10.2	9.4	12.6
Kemper	80	32	48	116	54	62	10.0	10.2	9.9
Lafayette	207	135	72	220	142	78	8.4	7.4	11.1

Table 6 - Page 2

Deaths - Mississippi - 1968

County	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Lamar	85	70	15	150	126	24	9.5	9.6	8.8
Lauderdale	1,109	740	369	821	520	301	10.7	9.9	12.4
Lawrence	103	46	57	147	80	67	13.1	10.4	19.0
Leake	154	90	64	222	133	89	11.3	11.2	11.6
Lee	571	409	162	466	325	141	9.6	8.5	13.7
Leflore	597	246	351	547	209	338	11.3	11.5	11.2
Lincoln	337	249	88	319	216	103	12.2	11.9	12.7
Lowndes	489	261	228	498	256	242	8.8	6.8	12.4
Madison	337	125	212	354	118	236	10.9	10.9	10.9
Marion	204	144	60	237	162	75	10.2	10.5	9.6
Marshall	192	61	131	255	92	163	10.6	10.6	10.6
Monroe	373	226	147	387	242	145	10.8	9.7	13.2
Montgomery	194	127	67	170	108	62	11.6	12.6	10.2
Neshoba	213	144	69	233	167	66	11.6	11.3	12.6
Newton	223	147	76	270	188	82	13.1	12.9	13.6
Noxubee	167	42	125	210	61	149	13.8	13.3	14.0
Oktibbeha	233	109	124	249	119	130	8.0	6.0	11.6
Panola	260	109	151	329	140	189	11.2	10.8	11.5
Pearl River	218	165	53	278	210	68	9.6	8.8	13.3
Perry	73	54	19	103	70	33	11.3	10.9	12.1
Pike	414	234	180	454	251	203	14.0	13.6	14.5
Pontotoc	203	171	32	209	173	36	11.5	11.8	10.4
Prentiss	240	212	28	245	219	26	11.4	11.4	11.0
Quitman	131	32	99	223	80	143	12.3	11.0	13.2
Rankin	439	259	180	268	183	85	6.3	6.1	6.6
Scott	223	144	79	252	163	89	10.5	10.2	11.2
Sharkey	88	20	68	108	27	81	11.3	8.1	13.1
Simpson	226	152	74	238	169	69	11.8	11.4	12.7
Smith	68	52	16	155	120	35	11.9	12.5	10.4
Stone	67	49	18	83	56	27	10.2	9.0	14.2
Sunflower	375	111	264	455	157	298	10.9	11.8	10.4
Tallahatchie	209	69	140	240	88	152	11.2	11.6	10.9
Tate	112	50	62	189	90	99	9.2	9.3	9.0
Tippah	170	138	32	181	149	32	10.2	10.4	9.8
Tishomingo	119	112	7	164	156	8	9.8	9.7	13.3
Tunica	124	20	104	144	22	122	9.5	5.1	11.3
Union	193	155	38	212	170	42	10.6	10.1	13.3
Walthall	117	72	45	145	90	55	9.9	11.1	8.4
Warren	710	305	405	560	234	326	12.2	8.9	16.6
Washington	860	303	557	897	294	603	11.7	8.4	14.5
Wayne	169	101	68	183	107	76	10.6	9.9	11.9
Webster	108	94	14	134	116	18	12.0	13.6	6.9
Wilkinson	111	56	55	119	52	67	10.7	15.6	8.6
Winston	176	110	66	226	138	88	12.1	14.3	9.8
Yalobusha	120	67	53	160	97	63	12.8	13.5	11.9
Yazoo	302	106	196	388	148	240	12.4	11.0	13.4
Out of state	1,197	863	334	858	656	202			

Table 6 - Page 3
Deaths - Mississippi - 1968

City**	Place of Occurrence			Place of Residence		
	Number			Number		
	Total	White	Non-white	Total	White	Non-white
Biloxi	437	363	74	376	305	71
Clarksdale	466	213	253	306	129	177
Cleveland	172	78	94	128	58	70
Columbus	358	212	146	280	150	130
Corinth	78	63	15	156	121	35
Greenville	633	268	365	530	198	332
Greenwood	472	198	274	334	151	183
Gulfport	558	416	142	402	292	110
Hattiesburg	660	463	197	430	248	182
Jackson	2,203	1,234	969	1,396	822	574
Laurel	612	353	259	297	173	124
McComb	233	143	90	171	110	61
Meridian	1,027	685	342	595	369	226
Natchez	451	191	260	269	94	175
Pascagoula	336	245	91	179	137	42
Tupelo	453	334	119	184	132	52
Vicksburg	629	284	345	406	163	243
Yazoo City	191	84	107	186	63	123
Total urban#	14,639	8,712	5,927	9,678	5,407	4,271
Total rural	9,008	4,549	4,459	14,308	8,061	6,247

* Deaths per 1,000 population.

** Cities having 10,000 population and over according to 1960 census.

Includes all incorporated towns of 2,500 population or more according to 1960 census.

TABLE 7
DEATHS
MISSISSIPPI
1968

By Cause Groups and Certain Selected Causes, by Race, Sex, and Age,
with Total Rates, Residence Data

Inter- national Classi- fication Number#	Cause of Death	Total		Race	Total by Race	Sex	Total by Race and Sex	Age								65 and Over	Unknown	
		Number	Rate*					Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54			55-64
000-999	ALL CAUSES	23,986	10.2	W	13,468	M	7,939	295	50	41	43	246	170	307	795	1,496	4,495	1
				C	10,518	F	5,529	222	39	28	15	79	75	137	330	635	3,969	0
						M	5,766	621	94	49	52	199	195	225	451	916	2,959	5
						F	4,752	475	81	30	28	97	115	226	341	685	2,671	3
000-136	Infective and parasitic diseases	337	14.3	W	130	M	78	9	2	3	1	6	3	2	8	18	26	0
				C	207	F	52	8	5	0	0	2	1	2	1	7	26	0
						M	111	48	4	1	1	4	4	4	5	16	24	0
						F	96	42	10	2	1	4	2	8	4	7	16	0
004,006	Bacillary dysentery and amebiasis	14	0.6	W	5	M	4	0	0	0	0	2	0	1	0	0	1	0
				C	9	F	1	1	0	0	0	0	0	0	0	0	0	0
						M	5	2	0	1	0	0	0	0	0	1	1	0
						F	4	2	1	0	1	0	0	0	0	0	0	0
008,009	Enteritis and other diarrheal diseases	113	4.8	W	29	M	16	3	2	2	0	0	0	0	0	1	7	0
				C	84	F	13	2	2	0	0	0	0	0	0	1	8	0
						M	42	35	3	0	0	0	1	0	1	1	1	0
						F	42	31	7	0	0	0	1	0	0	0	3	0
010-019	Tuberculosis (all forms)	89	3.8	W	37	M	28	0	0	0	0	0	1	0	4	11	12	0
				C	52	F	9	0	1	0	0	0	0	0	0	3	5	0
						M	32	0	0	0	0	1	0	3	3	9	16	0
						F	20	0	0	0	0	2	1	5	2	4	6	0

Table 7 - Page 2
Deaths - Mississippi - 1968

Inter-national Classification Number#	Cause of Death	Total		Race	Total by Race	Sex	Total by Race and Sex	Age								65 & Over	Unknown
		Number	Rate*					Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	
036	Meningococcal infection	5	0.2	W	5	M	1	0	0	0	0	1	0	0	0	0	0
				C	0	F	4	0	1	0	0	0	0	0	0	0	0
						M	0	0	0	0	0	0	0	0	0	0	0
						F	0	0	0	0	0	0	0	0	0	0	0
070	Infectious hepatitis	18	0.8	W	12	M	5	0	0	0	0	1	0	0	2	2	0
				C	6	F	7	1	0	0	0	1	1	0	0	1	0
						M	1	0	0	0	0	1	0	0	0	0	0
						F	5	0	0	0	0	1	0	2	0	1	0
090-097	Syphilis and its sequelae	12	0.5	W	2	M	2	0	0	0	0	0	0	0	0	1	0
				C	10	F	0	0	0	0	0	0	0	0	0	0	0
						M	7	1	0	0	0	0	0	0	1	2	0
						F	3	0	0	0	0	0	0	1	0	1	0
140-209	Malignant neoplasms	3,267	139.0	W	2,101	M	1,213	0	4	7	4	10	13	44	148	323	660
				C	1,166	F	888	2	1	4	3	7	16	41	130	189	498
						M	609	0	3	4	3	7	8	14	55	139	375
						F	557	3	4	3	3	7	12	49	68	129	278
210-239	Benign and unspecified neoplasms	49	2.1	W	28	M	9	0	0	0	0	0	2	1	2	2	0
				C	21	F	19	1	0	0	0	1	1	1	3	5	0
						M	9	2	0	0	0	1	1	0	1	3	0
						F	12	1	0	0	0	1	0	3	2	2	0
240-279	Endocrine, nutritional, and metabolic diseases	521	22.2	W	267	M	114	2	3	0	0	1	1	5	10	20	72
				C	254	F	153	3	2	3	0	1	3	5	5	25	106
						M	88	9	0	2	0	2	0	4	4	24	43
						F	166	5	1	0	0	3	6	3	18	42	88
250	Diabetes mellitus	436	18.5	W	221	M	92	0	0	0	0	0	1	3	9	18	61
				C	215	F	129	0	0	0	0	0	3	4	4	25	93
						M	64	0	0	0	0	1	0	3	4	22	32
						F	151	0	0	2	0	3	5	3	16	39	85

Inter-national Classification Number#	Cause of Death	Total		Sex	Total by Race	Total by Race and Sex	Age								65 & Over	Unknown	
		Number	Rate*				Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54			55-64
260-269	Avitaminoses and other nutritional deficiency	39	1.6	W	15	M 7 F 8	1 0 8 5	0 1 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	1 0 1 1	5 7 7 2		
280-289	Diseases of blood and blood-forming organs	68	2.9	W C	25 43	M 16 F 9 M 23 F 20	0 0 2 1	1 0 3 3	0 0 1 1	0 0 2 3	1 0 1 2	1 1 3 1	1 1 3 1	1 0 2 2	11 7 4 2		
280-285	Anemias	56	2.4	W C	15 41	M 9 F 6 M 22 F 19	0 0 2 1	0 0 3 3	0 0 1 1	0 0 2 3	0 1 1 2	0 0 3 1	0 0 3 1	1 0 2 2	7 5 4 1		
290-315	Mental disorders	49	2.1	W C	25 24	M 22 F 3 M 15 F 9	0 0 0 0	1 0 0 0	0 0 1 2	0 0 3 1	1 1 6 3	1 0 1 1	1 0 1 1	6 1 3 1	9 1 1 1		
303	Alcoholism	30	1.3	W C	11 19	M 11 F 0 M 12 F 7	0 0 0 0	0 0 0 0	0 0 0 0	0 0 3 1	0 0 4 3	4 0 1 1	4 0 1 1	5 0 2 1	2 0 1 0		
320-389	Diseases of the nervous system and sense organs	167	7.1	W C	89 78	M 51 F 38 M 44 F 34	4 2 7 4	3 1 6 3	3 1 5 0	1 0 2 4	3 2 4 1	8 1 4 1	9 5 8 3	9 5 8 3	13 20 5 13		
320	Meningitis	32	1.4	W C	8 24	M 6 F 2 M 18 F 6	1 0 6 3	0 1 5 0	1 1 2 0	0 0 0 2	0 0 2 0	2 0 1 0	2 0 1 0	2 0 1 1	0 0 0 0		

Table 7 - Page 4
Deaths - Mississippi - 1968

Inter-national Classification Number#	Cause of Death	Total		Total by Race	Sex	Total by Race and Sex	Age								55-64	65 and Over	Unknown
		Number	Rate*	Race	Sex		Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54			
340	Multiple sclerosis	11	0.5	W	M	2	0	0	0	0	0	0	1	0	0	1	0
				C	F	4	0	0	0	0	0	0	1	0	1	2	0
					M	4	0	0	0	0	0	0	1	1	2	0	0
					F	1	0	0	0	0	0	0	0	0	0	1	0
345	Epilepsy	23	1.0	W	M	5	0	0	0	1	0	1	1	2	0	0	0
				C	F	3	0	0	0	0	1	0	0	1	1	0	0
					M	8	0	0	1	1	1	1	0	1	2	1	0
					F	7	0	0	0	0	3	1	0	0	0	3	0
390-458	Diseases of the circulatory system	12,000	510.5	W	M	4,078	0	0	0	1	6	16	87	351	784	2,833	0
				C	F	3,129	4	0	1	1	4	8	33	93	282	2,703	0
					M	2,520	12	4	1	5	12	29	62	214	456	1,725	3
					F	2,273	11	1	1	1	11	28	72	146	357	1,646	2
390-398, 402, 404-429	Diseases of heart	7,969	339.0	W	M	3,028	0	0	0	0	5	9	66	302	663	1,983	0
				C	F	1,996	3	0	1	1	1	3	15	59	209	1,704	0
					M	1,584	9	4	0	5	4	16	36	140	295	1,072	3
					F	1,361	8	1	0	1	9	15	34	96	218	979	0
400,401, 403	Hypertension	209	8.9	W	M	43	0	0	0	0	0	0	3	3	7	30	0
				C	F	35	0	0	0	0	0	0	2	1	5	27	0
					M	64	0	0	0	0	1	4	3	13	9	34	0
					F	67	0	0	1	0	0	0	9	9	15	33	0
430-438	Cerebrovascular disease	3,115	132.5	W	M	756	0	0	0	1	1	5	14	31	84	620	0
				C	F	875	1	0	0	0	1	4	12	24	52	781	0
					M	747	0	0	1	0	7	8	21	50	136	524	0
					F	737	0	0	0	0	1	11	27	36	113	548	1
440	Arteriosclerosis	316	13.4	W	M	95	0	0	0	0	0	0	0	3	1	91	0
				C	F	116	0	0	0	0	0	0	0	1	5	110	0
					M	54	0	0	0	0	0	0	0	2	2	50	0
					F	51	0	0	0	0	0	0	0	1	4	45	1

Table 7 - Page 5
Deaths - Mississippi - 1968

Inter- national Classi- fication Number#	Cause of Death	Total		Race	Total by Race	Sex	Total by Race and Sex	Age								65 and Over	Unknown	
		Number	Rate*					Under- 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54			55-64
460-519	<u>Diseases of the respiratory system</u>	1,282	54.5	W	738	M	478	9	9	2	0	7	5	7	28	69	342	0
				C	544	F	260	18	2	1	1	1	3	10	11	23	196	0
						M	290	73	22	2	4	8	5	17	17	28	121	0
						F	254	84	13	1	1	1	7	5	7	17	115	0
470-474	Influenza	96	4.1	W	45	M	27	0	0	0	0	1	0	0	0	1	25	0
				C	51	F	18	0	0	0	0	1	0	0	2	1	14	0
						M	21	1	0	0	1	2	0	0	0	2	15	0
						F	30	2	1	0	0	0	1	1	0	5	20	0
480-486	Pneumonia	729	31.0	W	369	M	208	9	6	0	0	3	4	4	13	19	150	0
				C	360	F	161	16	1	0	0	3	1	1	3	4	132	0
						M	187	58	15	2	2	4	5	6	12	11	72	0
						F	173	64	6	0	1	3	4	0	4	10	81	0
490,491	Chronic and unquali- fied bronchitis	38	1.6	W	31	M	24	0	2	0	0	0	0	1	1	4	16	0
				C	7	F	7	0	0	0	0	0	0	1	1	2	3	0
						M	6	2	1	0	0	0	0	1	0	1	1	0
						F	1	0	0	0	0	0	0	0	0	0	1	0
492	Emphysema	213	9.1	W	190	M	157	0	0	0	0	0	0	2	7	32	116	0
				C	23	F	33	0	0	0	0	0	0	0	3	9	21	0
						M	20	0	0	0	0	0	0	0	1	4	15	0
						F	3	0	0	0	0	0	0	1	0	0	2	0
493	Asthma	52	2.2	W	21	M	14	0	1	0	0	0	1	0	3	3	6	0
				C	31	F	7	0	0	0	0	0	2	0	1	3	1	0
						M	16	0	0	0	1	0	0	1	2	5	7	0
						F	15	1	3	0	0	1	1	2	3	1	3	0

Table 7 - Page 6
Deaths - Mississippi - 1968

Inter-national Classification Number#	Cause of Death	Total		Race	Total by Race	Sex	Total by Race and Sex	Age								65 and Over	Unknown
		Number	Rate*					Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	
<u>520-577</u>	<u>Diseases of the digestive system</u>	<u>566</u>	<u>24.1</u>	W	378	M	208	0	0	1	0	2	9	16	34	53	<u>93</u>
				C	188	F	170	2	1	1	1	5	2	11	20	25	<u>101</u>
531-533	Peptic ulcer	82	3.5	W	59	M	43	0	0	0	0	0	1	3	4	12	23
				C	23	F	16	0	0	0	0	0	0	0	3	1	12
550-553, 560	Hernia and intestinal obstruction	88	3.7	W	43	M	20	0	0	1	0	0	0	3	3	2	11
				C	45	F	23	2	0	0	0	0	0	2	1	4	14
571	Cirrhosis of liver	146	6.2	W	113	M	79	0	0	0	0	0	2	6	22	25	24
				C	33	F	34	0	1	0	2	0	0	2	8	7	14
<u>580-629</u>	<u>Diseases of the genitourinary system</u>	<u>518</u>	<u>22.0</u>	W	252	M	130	0	1	1	0	3	3	3	10	14	98
				C	266	F	122	1	0	0	1	3	4	2	7	14	90
580-584	Nephritis and nephrosis	181	7.7	W	58	M	30	0	1	1	0	0	2	0	6	7	13
				C	123	F	28	0	0	0	0	2	2	1	4	1	18
590	Infections of kidney	199	8.5	W	133	M	58	0	0	0	0	0	0	1	3	5	49
				C	66	F	75	0	0	0	1	0	2	0	3	12	57
						M	40	0	0	0	0	1	0	2	2	7	28
						F	26	0	0	0	0	1	0	3	3	5	14

Table 7 - Page 7
Deaths - Mississippi - 1968

Inter-national Classi- fication Number#	Cause of Death	Total		Race	Total by Race	Sex	Total by Race and Sex	Age								65 and Over	Unknown
		Number	Rate*					Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	
630-678	Complications of pregnancy, childbirth, and the puerperium	21	4.6	W	3	M	0	0	0	0	0	0	0	0	0	0	0
				C	18	F	3	0	0	0	0	0	0	0	0	0	0
680-709	Diseases of the skin and subcutaneous tissue	26	1.1	W	14	M	6	0	0	0	0	0	0	0	0	1	0
				C	12	F	8	0	0	0	0	0	0	0	0	1	0
710-738	Diseases of the musculoskeletal system and connective tissue	60	2.6	W	38	M	13	0	0	0	0	0	0	0	0	0	0
				C	22	F	25	0	0	0	0	0	0	0	0	0	0
740-759	Congenital anomalies	205	8.7	W	135	M	78	49	7	6	1	4	2	0	0	0	0
				C	70	F	57	41	7	2	1	3	0	0	0	0	0
760-778	Certain causes of mortality in early infancy	791	17.3	W	320	M	191	191	0	0	0	0	0	0	0	0	0
				C	471	F	129	129	0	0	0	0	0	0	0	0	0
780-796	Symptoms and ill-defined conditions	1,858	79.0	W	487	M	327	11	0	1	0	2	4	19	46	83	163
				C	1,371	F	160	5	0	0	1	9	1	5	15	25	106
						M	752	128	11	0	2	8	14	23	56	126	381
						F	619	89	12	0	2	0	13	25	47	71	352

Table 7 - Page 8
Deaths - Mississippi - 1968

Inter- national Classi- fication Number#	Cause of Death	Total		Total by Race	Sex	Total by Race and Sex	Age								65 and Over	Unknown
		Number	Rate*	Race			Under 1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64		
300-999	Accidents, poison- ings, and violence	2,201	93.6	W	M	927	20	17	35	205	108	115	140	107	160	1
				C	F	304	6	15	8	42	30	28	38	29	88	0
					M	702	24	25	32	139	118	88	73	64	102	0
					F	268	27	20	11	34	21	31	20	19	56	0
310-823	Motor vehicle accidents	821	34.9	W	M	406	5	7	18	130	56	43	55	48	41	0
				C	F	136	1	7	5	31	20	12	16	14	21	0
					M	207	0	8	7	56	37	28	12	19	31	0
					F	72	1	9	6	14	5	10	6	5	8	0
300-807, 325-949	All other accidents	902	38.4	W	M	336	15	10	17	57	27	38	46	29	83	0
				C	F	121	5	8	2	5	6	3	8	9	66	0
					M	308	24	17	22	50	28	25	33	25	56	0
					F	137	26	11	4	7	4	9	5	9	45	0
950-959	Suicide	171	7.3	W	M	112	0	0	0	9	9	13	30	19	32	0
				C	F	36	0	0	1	5	4	9	12	4	1	0
					M	14	0	0	1	2	2	1	1	3	4	0
					F	9	0	0	1	0	1	3	0	2	2	0
960-978	Homicide	256	10.9	W	M	53	0	0	0	7	12	13	8	10	1	1
				C	F	7	0	0	0	1	0	2	1	2	0	0
					M	157	0	0	0	29	46	32	24	15	11	0
					F	39	0	0	0	12	9	6	8	1	0	0

* Numbers are those of the Eighth Revision International Classification of Diseases, Adapted for Use in the United States which went into use January 1, 1968.

Total deaths per 1,000 population. Deaths from specific causes per 100,000 population, except deaths from maternal causes (classification numbers 630-678) are per 10,000 live births and deaths from certain causes of mortality in early infancy (classification numbers 760-778) are per 1,000 live births.

TABLE 8
DEATHS UNDER 1 YEAR
MISSISSIPPI
1968

By Selected Causes, by Race and Age, with Total Rates, Residence Data

Cause of Death and International Classi- fication Number#	Total		Total by Race	Days of Age						Months of Age										
	Number	Rate*		Under 1 Day	1- 6	7- 13	14- 20	21- 27	Under 28 Days	1	2	3	4	5	6	7	8	9	10	11
ALL CAUSES	1,613	35.3	W 517 C 1,096	225 340	162 240	18 33	13 36	6 43	424 692	19 123	23 73	12 52	11 38	6 29	5 24	6 17	5 11	3 18	2 8	1 11
Diarrheal disease (009)	66	1.4	W 3 C 63	0 0	3 5	0 4	0 3	0 3	15	13	9	3	5	4	0	1	0	0	0	0
Septicemia (038)	23	0.5	W 9 C 14	2 0	4 6	2 1	0 1	0 1	8 9	1 1	0 1	0 1	0 0	0 1	0 0	0 0	0 0	0 1	0 0	0
Other infective and parasitic diseases	18	0.4	W 5 C 13	1 0	0 1	0 0	0 1	0 1	1 3	0 2	1 1	1 2	1 0	1 2	0 1	0 0	0 0	0 1	0 1	0
Malignant neoplasms (140-209)	5	0.1	W 2 C 3	0 0	1 1	0 0	1 0	0 0	2 1	0 0	0 0	0 1	0 0	0 0	0 0	0 0	0 0	0 1	0 0	0
Benign and unspecified neoplasms (210-239)	4	0.1	W 1 C 3	0 1	0 1	0 0	0 0	0 0	0 2	0 0	0 0	0 0	1 0	0 0	0 0	0 0	0 0	0 0	0 0	0
Cystic fibrosis (273.0)	4	0.1	W 3 C 1	0 0	1 1	0 0	0 0	0 0	1 1	0 0	0 0	1 0	0 0	1 0	0 0	0 0	0 0	0 0	0 0	0
Diseases of blood and blood-forming organs (280-289)	3	0.1	W 0 C 3	0 0	0 0	0 0	0 0	0 0	0 0	0 1	0 0	0 0	0 0	0 0	0 2	0 0	0 0	0 0	0 0	0
Meningitis (320)	10	0.2	W 1 C 9	0 0	0 0	0 0	0 0	0 1	0 1	0 2	0 3	0 0	1 1	0 0	0 0	0 0	0 0	0 1	0 0	0
Acute upper respiratory infections (460-465)	16	0.3	W 1 C 15	0 0	0 0	0 0	0 1	0 2	0 3	0 2	1 3	0 2	0 2	0 1	0 2	0 0	0 0	0 0	0 0	0
Bronchitis and bronchio- litis (466, 490, 491)	8	0.2	W 0 C 8	0 0	0 3	0 0	0 0	0 0	0 3	0 2	0 0	0 1	0 0	0 0	0 1	0 0	0 1	0 0	0 0	0

Table 8 - Page 2
Deaths Under 1 Year - Mississippi - 1968

Cause of Death and International Classification Number#	Total		Total by Race	Days of Age							Months of Age										
	Number	Rate*		Under 1 Day	1-6	7-13	14-20	21-27	Under 28 Days	1	2	3	4	5	6	7	8	9	10	11	
Influenza and pneumonia (470-486) Other diseases of respiratory system (492-519) Hernia and intestinal obstruction (550-553, 560) Congenital anomalies (740-759) Maternal conditions unrelated to pregnancy (760, 761) Hypoxemia of pregnancy (762) Maternal antepartum and intrapartum infection (763) Difficult labor (764-768) Other complications of pregnancy and childbirth (769) Conditions of placenta (770) Conditions of umbilical cord (771)	150	3.3	W 25 C 125	2	6	0	2	0	10	5	2	1	0	2	3	1	1	0	0	11	
	10	0.2	W 1 C 9	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	
	11	0.2	W 2 C 9	0	1	1	0	0	2	0	0	0	0	0	0	0	0	0	0	0	
	137	3.0	W 90 C 47	32	19	4	5	3	63	4	7	5	3	0	2	2	2	0	1	1	
	4	0.1	W 0 C 4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
6	0.1	W 3 C 3	2	1	0	0	0	3	0	0	0	0	0	0	0	0	0	0	0	0	
2	**	W 1 C 1	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	
12	0.3	W 4 C 8	3	1	0	0	0	4	0	0	0	0	0	0	0	0	0	0	0	0	
43	0.9	W 21 C 22	18	3	0	0	0	21	0	0	0	0	0	0	0	0	0	0	0	0	
39	0.8	W 17 C 22	12	5	0	0	0	17	0	0	0	0	0	0	0	0	0	0	0	0	
8	0.2	W 5 C 3	4	1	0	0	0	5	0	0	0	0	0	0	0	0	0	0	0	0	

Table 8 - Page 3
Deaths Under 1 Year - Mississippi - 1968

Cause of Death and International Classification Number#	Total		Total by Race	Days of Age						Months of Age											
	Number	Rate*		Under 1 Day	1-6	7-13	14-20	21-27	Under 28 Days	1	2	3	4	5	6	7	8	9	10	11	
Birth injury without mention of cause (772)	31	0.7	W 15 C 16	5 5	8 7	2 1	0 2	0 1	15 16	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	
Hemolytic disease of newborn (774, 775)	8	0.2	W 5 C 3	3 0	2 2	0 0	0 0	0 0	5 2	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	1 0	
Hyaline membrane disease (776.1)	92	2.0	W 50 C 42	19 15	31 25	0 0	0 1	0 0	50 41	0 1	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	
Respiratory distress syndrome (776.2)	71	1.6	W 33 C 38	14 19	19 17	0 2	0 0	0 0	33 38	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	
Asphyxia of newborn, unspecified (776.9)	143	3.1	W 75 C 68	44 40	27 25	1 1	2 1	1 1	75 68	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	
Other anoxic and hypoxic conditions not elsewhere classifiable (776.0, 776.3, 776.4)	17	0.4	W 5 C 12	3 6	2 5	0 0	0 1	0 0	5 12	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	
Immaturity, unqualified (777)	283	6.2	W 76 C 207	52 122	18 58	3 8	2 4	0 5	75 197	1 9	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	1 0	0 0	
Hemorrhagic disease of newborn (778.2)	15	0.3	W 4 C 11	1 2	3 9	0 0	0 0	0 0	4 11	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	
Other conditions of newborn (778.0, 778.1, 778.3, 778.9)	17	0.4	W 6 C 11	3 7	2 1	1 2	0 0	0 1	6 11	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	
Symptoms and ill-defined conditions (780-796)	233	5.1	W 16 C 217	3 54	3 21	0 3	0 10	2 7	8 95	2 40	4 34	1 13	1 12	0 5	0 6	0 5	0 2	0 3	0 0	2 2	

Table 8 - Page 4
Deaths Under 1 Year - Mississippi - 1968

Cause of Death and International Classi- fication Number#	Total		Total by Race	Days of Age					Months of Age											
	Number	Rate*		Under 1 Day	1- 6	7- 13	14- 20	21- 27	Under 28 Days	1	2	3	4	5	6	7	8	9	10	11
Accidents (800-949)	77	1.7	W	26	0	1	2	0	0	3	5	6	2	3	1	0	2	2	0	0
			C	51	0	3	1	2	3	9	16	2	8	4	1	2	2	1	3	1
All other causes	47	1.0	W	12	1	3	1	1	0	6	1	1	1	1	0	0	0	1	1	0
			C	35	6	5	1	1	1	14	8	1	4	2	2	0	3	0	1	0

Numbers are those of the Eighth Revision International Classification of Diseases, Adapted for Use in the United States which went into use January 1, 1968.

* Deaths per 1,000 live births.

* Rate less than 0.06.

TABLE 9
MARRIAGES AND DIVORCES
MISSISSIPPI
1968

Marriages by Place of Licensure, by Place of Occurrence, and by Place of Residence of Bride, by Race of Bride;
Divorces by Place of Occurrence, by Race

County	Marriages												Divorces		
	Place of Licensure			Place of Occurrence			Residence of Bride								
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
STATE	25,355	17,142	8,213	25,355	17,142	8,213	21,617	14,065	7,552	7,020	5,202	1,818			
Adams	426	264	162	420	263	157	345	209	136	144	79	65			
Alcorn	482	432	50	500	450	50	269	233	36	89	83	6			
Amite	122	73	49	118	73	45	117	62	55	20	14	6			
Attala	195	115	80	203	126	77	200	119	81	48	38	10			
Benton	56	37	19	67	47	20	67	45	22	11	6	5			
Bolivar	528	257	271	504	239	265	481	220	261	111	57	54			
Calhoun	129	101	28	146	116	30	141	110	31	33	24	9			
Carroll	27	20	7	45	38	7	42	32	10	11	7	4			
Chickasaw	237	167	70	218	150	68	208	144	64	51	40	11			
Choctaw	78	61	17	72	55	17	58	40	18	13	10	3			
Claiborne	71	25	46	81	27	54	68	19	49	17	1	16			
Clarke	169	112	57	171	116	55	156	102	54	45	33	12			
Clay	174	109	65	153	92	61	162	97	65	57	38	19			
Coahoma	391	163	228	386	160	226	354	139	215	114	55	59			
Copiah	241	121	120	249	133	116	264	138	126	43	26	17			
Covington	145	95	50	144	96	48	168	114	54	37	26	11			
DeSoto	947	589	358	943	583	360	191	124	67	121	108	13			
Forrest	700	533	167	727	555	172	679	501	178	312	239	73			
Franklin	72	46	26	75	55	20	67	47	20	17	11	6			
George	221	195	26	201	179	22	124	106	18	30	26	4			
Greene	68	55	13	99	83	16	69	46	23	22	18	4			
Grenada	240	158	82	217	134	83	219	137	82	59	36	23			
Hancock	362	341	21	348	329	19	127	112	15	66	64	2			
Harrison	1,968	1,697	271	2,003	1,731	272	1,604	1,336	268	734	632	102			

Table 9 - Page 2
Marriages and Divorces - Mississippi - 1968

County	Marriages												Divorces		
	Place of Licensure			Place of Occurrence			Residence of Bride								
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white			
Hinds	2,337	1,487	850	2,144	1,308	836	2,276	1,461	815	756	563	193			
Holmes	234	80	154	227	70	157	259	90	169	48	24	24			
Humphreys	132	64	68	128	57	71	129	58	71	23	15	8			
Issaquena	22	15	7	23	11	12	16	10	6	8	6	2			
Itawamba	178	168	10	184	177	7	136	126	10	35	34	1			
Jackson	980	805	175	952	774	178	828	676	152	335	286	49			
Jasper	120	69	51	138	76	62	163	96	67	17	11	6			
Jefferson	83	25	58	73	20	53	71	19	52	23	4	19			
Jeff Davis	138	77	61	127	70	57	99	55	44	19	8	11			
Jones	596	467	129	612	487	125	555	438	117	211	183	28			
Kemper	90	35	55	104	45	59	79	28	51	23	12	11			
Lafayette	159	102	57	154	93	61	142	86	56	65	35	30			
Lamar	131	114	17	160	140	20	148	128	20	40	34	6			
Lauderdale	699	457	242	683	444	239	668	434	234	225	191	34			
Lawrence	88	60	28	97	67	30	126	80	46	26	18	8			
Leake	163	96	67	168	101	67	167	91	76	52	38	14			
Lee	499	394	105	485	380	105	550	433	117	158	143	15			
Leflore	497	232	265	494	228	266	484	234	250	132	88	44			
Lincoln	286	210	76	288	214	74	249	177	72	55	42	13			
Lowndes	478	321	157	492	337	155	417	272	145	131	105	26			
Madison	298	106	192	333	128	205	282	114	168	97	44	53			
Marion	287	199	88	279	191	88	272	193	79	99	81	18			
Marshall	399	189	210	376	174	202	257	104	153	83	42	41			
Monroe	297	213	84	300	209	91	286	209	77	98	83	15			
Montgomery	134	83	51	142	88	54	134	84	50	30	21	9			
Neshoba	252	171	81	237	160	77	204	127	77	98	73	25			
Newton	153	100	53	181	120	61	187	127	60	35	23	12			
Noxubee	99	32	67	108	42	66	109	39	70	24	14	10			
Oktibbeha	270	146	124	256	132	124	249	135	114	55	34	21			
Panola	290	160	130	288	153	135	256	131	125	60	41	19			

Table 9 - Page 3
 Marriages and Divorces - Mississippi - 1968

County	Marriages												Divorces		
	Place of Licensure			Place of Occurrence			Residence of Bride								
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white			
Pearl River	369	315	54	369	316	53	265	221	44	146	129	17			
Perry	109	89	20	98	80	18	104	86	18	22	19	3			
Pike	386	242	144	409	260	149	359	229	130	118	85	33			
Pontotoc	183	149	34	184	154	30	162	131	31	46	40	6			
Prentiss	228	209	19	222	204	18	174	164	10	65	55	10			
Quitman	163	79	84	176	96	80	168	83	85	39	20	19			
Rankin	344	248	96	396	303	93	286	192	94	88	65	23			
Scott	256	153	103	241	147	94	272	170	102	54	35	19			
Sharkey	76	29	47	68	32	36	71	32	39	16	6	10			
Simpson	216	151	65	225	160	65	192	128	64	61	49	12			
Smith	104	80	24	126	98	28	108	75	33	33	26	7			
Stone	104	84	20	92	75	17	78	63	15	30	17	13			
Sunflower	281	112	169	320	138	182	309	126	183	53	25	28			
Tallahatchie	200	90	110	205	93	112	184	88	96	47	26	21			
Tate	231	153	78	218	142	76	136	78	58	27	17	10			
Tippah	164	138	26	167	141	26	169	138	31	46	43	3			
Tishomingo	291	272	19	277	257	20	176	167	9	35	35	0			
Tunica	151	57	94	135	46	89	130	38	92	27	13	14			
Union	230	186	44	234	189	45	207	168	39	40	33	7			
Walthall	130	88	42	134	92	42	82	48	34	35	29	6			
Warren	468	283	185	460	283	177	415	249	166	178	125	53			
Washington	824	478	346	837	489	348	733	393	340	285	177	108			
Wayne	200	139	61	181	124	57	137	97	40	66	51	15			
Webster	82	73	9	100	90	10	84	70	14	25	24	1			
Wilkinson	560	473	87	567	475	92	96	33	63	16	7	9			
Winston	177	98	79	178	98	80	157	89	68	41	24	17			
Yalobusha	120	83	37	123	85	38	126	85	41	34	23	11			
Yazoo	270	148	122	262	143	119	259	137	122	81	37	44			
Not reported				28	6	22									
Out of state							3,738	3,077	661						

TABLE 10
DIVORCES
MISSISSIPPI
1968

By Cause of Divorce, by Race

Cause	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>ALL CAUSES</u>	<u>7,020</u>	<u>5,202</u>	<u>1,818</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
Cruel and inhuman treatment	4,371	3,771	600	62.3	72.5	33.0
Desertion	2,009	950	1,059	28.6	18.3	58.2
Adultery	302	210	92	4.3	4.0	5.1
Drunkenness	228	194	34	3.2	3.7	1.9
Penitentiary	39	28	11	0.6	0.5	0.6
Insanity	9	6	3	0.1	0.1	0.2
Annulment	19	15	4	0.3	0.3	0.2
Bigamy	15	4	11	0.2	0.1	0.6
Miscellaneous and other	15	14	1	0.2	0.3	*
Not stated	13	10	3	0.2	0.2	0.2

* Less than 0.06 percent.

By Number of Minor Children, by Race

Number of Minor Children	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>TOTAL DIVORCES</u>	<u>7,020</u>	<u>5,202</u>	<u>1,818</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
None	3,090	2,186	904	44.0	42.0	49.7
One	1,523	1,229	294	21.7	23.6	16.2
Two	1,090	864	226	15.5	16.6	12.4
Three or more	1,198	840	358	17.1	16.1	19.7
Not stated	119	83	36	1.7	1.6	2.0

By Number of Years Married, by Race

Number of Years Married	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>TOTAL DIVORCES</u>	<u>7,020</u>	<u>5,202</u>	<u>1,818</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
<u>Under five</u>	<u>2,958</u>	<u>2,544</u>	<u>414</u>	<u>42.1</u>	<u>48.9</u>	<u>22.8</u>
Under one	546	507	39	7.8	9.7	2.1
One	735	652	83	10.5	12.5	4.6
Two	647	555	92	9.2	10.7	5.1
Three	583	472	111	8.3	9.1	6.1
Four	447	358	89	6.4	6.9	4.9
Five - nine	1,400	1,055	345	19.9	20.3	19.0
Ten - nineteen	1,420	963	457	20.2	18.5	25.1
Twenty and over	1,174	591	583	16.7	11.4	32.1
Not stated	68	49	19	1.0	0.9	1.0

TABLE 11
LIVE BIRTHS
MISSISSIPPI
1913-1968

By Year, by Race, with Rates, Occurrence Data 1913-1943,
Residence Data 1944-1968

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1913	41,103	21,874	19,229	21.9	26.6	18.2
1914	40,623	21,360	19,263	21.4	25.7	18.0
1915	39,484	20,999	18,485	20.5	23.8	17.1
1916	39,033	20,130	18,903	20.0	23.6	17.2
1917	37,141	19,056	18,085	18.8	22.0	16.3
1918	39,391	20,005	19,386	19.7	22.8	17.2
1919	42,491	22,028	20,463	21.0	24.9	18.0
1920	43,743	22,438	21,305	24.4	26.3	22.7
1921	46,482	23,703	22,779	25.6	27.3	24.1
1922	43,723	21,286	22,437	23.8	24.1	23.6
1923	43,266	21,140	22,126	23.3	23.6	23.1
1924	43,960	22,180	21,780	23.4	24.3	22.5
1925	45,563	22,265	23,298	24.0	24.1	23.9
1926	50,449	24,600	25,849	26.2	26.2	26.3
1927	49,204	24,163	25,041	25.3	25.3	25.3
1928	48,126	23,642	24,484	24.5	24.4	24.5
1929	45,570	22,105	23,465	22.9	22.5	23.3
1930	47,982	23,192	24,790	23.8	23.2	24.0
1931	45,123	21,637	23,486	22.2	21.4	22.9
1932	46,481	21,883	24,598	22.6	21.4	23.9
1933	44,184	20,407	23,777	21.3	19.7	22.9
1934	47,905	22,342	25,563	23.0	21.4	24.6
1935	48,291	21,972	26,319	23.0	20.8	25.1
1936	49,261	22,194	27,067	23.3	20.8	25.7
1937	52,170	23,285	28,885	24.4	21.6	27.3
1938	53,729	24,046	29,683	25.0	22.1	27.8
1939	52,288	23,505	28,783	24.1	21.4	26.8
1940	52,526	23,084	29,442	24.0	20.9	27.3
1941	54,561	24,167	30,394	24.7	21.6	28.0
1942	56,521	25,306	31,215	25.4	22.4	28.6
1943	59,408	27,611	31,797	28.9	26.4	31.6
1944#	56,586	25,647	30,939	28.3	25.2	31.6
1945#	53,540	23,426	30,114	26.9	23.1	30.9
1946#	61,430	29,715	31,715	26.8	25.4	28.2
1947#	66,291	32,423	33,868	28.6	27.5	29.9
1948#	65,291	29,666	35,625	28.0	24.9	31.2
1949#	66,128	28,786	37,342	28.1	24.0	32.5

Table 11 - Page 2
Live Births - Mississippi - 1913-1968

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1950#	64,427	27,188	37,239	29.6	22.9	37.6
1951#	65,512	27,798	37,714	30.1	23.4	38.1
1952#	64,509	28,073	36,436	29.6	23.6	36.8
1953#	64,061	28,312	35,749	29.4	23.8	36.1
1954#	64,748	28,781	35,967	29.7	24.2	36.3
1955#	63,244	27,546	35,698	29.0	23.2	36.0
1956#	62,178	27,278	34,900	29.5	23.8	36.2
1957#	60,712	26,804	33,908	28.8	23.4	35.2
1958#	59,189	26,812	32,377	27.1	20.9	35.9
1959#	60,562	27,979	32,583	27.2	20.9	36.6
1960#	59,210	27,198	32,012	27.2	21.6	34.8
1961#	59,523	27,435	32,088	27.3	21.8	34.8
1962#	58,921	27,512	31,409	26.1	20.8	33.5
1963#	57,316	26,823	30,493	25.0	19.9	32.4
1964#	56,650	25,942	30,708	24.5	18.9	32.5
1965#	51,171	23,369	27,802	22.0	17.4	28.2
1966#	48,230	22,736	25,494	20.4	16.5	25.9
1967#	46,594	22,675	23,919	19.7	15.9	25.6
1968#	45,719	22,807	22,912	19.4	15.7	25.4

* Births per 1,000 population.

1913-1920, population estimates based on U.S. Censuses of 1910 and 1920.

1921-1930, estimates based on U.S. Censuses of 1920 and 1930.

1931-1940, estimates based on U.S. Censuses of 1930 and 1940.

1941 and 1942, projected estimates based on U.S. Censuses of 1930 and 1940.

1943 and 1944, estimates of civilian population obtained from Ration Book Registration.

1945, estimate of civilian population, July 1, 1945, obtained from Bureau of Census Bulletin, Series P-46, No. 3, dated February 12, 1946.

1946-1949, projected estimates as of July 1, each year based on U.S. Censuses of 1930 and 1940.

1950-1955, U.S. Census population of April 1, 1950.

1956 and 1957, estimate as of July 1, 1956, prepared by Dr. Harald A. Pedersen, Sociology Department, Mississippi State University, with adjustments by Statistical Service, Mississippi State Board of Health.

1958, estimate as of January 1, prepared by Dr. George L. Wilber and Mr. Carl T. Robbins, Sociology Department, Mississippi State University.

1959, estimate as of January 1, prepared by Dr. George L. Wilber and Ellen Bryant, Sociology Department, Mississippi State University.

1960 and 1961, U.S. Census population of April 1, 1960.

1962-1968, estimates as of July 1, prepared by the Division of Sociology and Anthropology, Mississippi State University.

On basis of place of residence.

TABLE 12
DEATHS
MISSISSIPPI
1913-1968

By Year, by Race, with Rates, Occurrence Data 1913-1943,
Residence Data 1944-1968

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1913	21,305	7,643	13,662	11.4	9.3	13.0
1914	21,485	7,442	14,043	11.3	8.9	13.1
1915	24,001	8,088	15,913	12.5	9.6	14.7
1916	21,794	7,588	14,206	11.2	8.9	13.0
1917	23,579	8,048	15,531	11.9	9.3	14.0
1918	30,437	10,177	20,260	15.2	11.6	18.0
1919	22,667	8,142	14,525	12.7	9.5	15.5
1920	22,072	7,916	14,156	12.3	9.3	15.1
1921	19,991	7,308	12,683	11.0	8.4	13.4
1922	19,278	7,312	11,966	10.5	8.3	12.6
1923	20,363	7,688	12,675	11.0	8.6	13.2
1924	21,155	7,809	13,346	11.3	8.6	13.8
1925	22,229	8,143	14,086	11.7	8.8	14.4
1926	23,929	9,133	14,796	12.5	9.7	15.1
1927	23,257	8,765	14,492	12.0	9.2	14.6
1928	25,965	10,111	15,854	13.2	10.4	15.9
1929	25,868	10,048	15,820	13.0	10.2	15.7
1930	24,098	9,072	15,026	11.9	9.0	14.8
1931	22,007	8,641	13,366	10.8	8.5	13.0
1932	20,322	8,295	12,027	9.9	8.1	11.6
1933	21,617	8,898	12,719	10.4	8.6	12.2
1934	21,786	9,350	12,436	10.5	9.0	12.0
1935	21,285	9,186	12,099	10.1	8.7	11.6
1936	24,127	10,367	13,760	11.4	9.7	13.1
1937	24,005	10,055	13,950	11.2	9.3	13.2
1938	22,814	9,471	13,343	10.6	8.7	12.5
1939	22,701	9,464	13,237	10.5	8.6	12.3
1940	23,185	9,565	13,620	10.6	8.6	12.6
1941	22,573	9,442	13,131	10.2	8.4	12.1
1942	20,693	8,750	11,943	9.3	7.7	10.9
1943	20,850	9,139	11,711	10.2	8.7	11.6
1944#	20,199	8,830	11,369	10.1	8.7	11.6
1945#	19,024	8,452	10,572	9.6	8.3	10.8
1946#	19,321	8,995	10,326	8.4	7.7	9.2
1947#	20,007	9,238	10,769	8.6	7.8	9.5
1948#	20,183	9,216	10,967	8.6	7.7	9.6
1949#	20,489	9,233	11,256	8.7	7.7	9.8

Table 12 - Page 2
Deaths - Mississippi - 1913-1968

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1950#	20,567	9,466	11,101	9.4	8.0	11.2
1951#	21,007	9,593	11,414	9.6	8.1	11.5
1952#	20,657	9,387	11,270	9.5	7.9	11.4
1953#	20,492	9,495	10,997	9.4	8.0	11.1
1954#	19,353	9,453	9,900	8.9	8.0	10.0
1955#	19,403	9,501	9,902	8.9	8.0	10.0
1956#	19,780	9,969	9,811	9.4	8.7	10.2
1957#	20,476	10,339	10,137	9.7	9.0	10.5
1958#	21,541	10,811	10,730	9.9	8.4	11.9
1959#	20,890	10,742	10,148	9.4	8.0	11.4
1960#	21,713	11,204	10,509	10.0	8.9	11.4
1961#	21,232	11,063	10,169	9.7	8.8	11.0
1962#	22,713	11,868	10,845	10.0	9.0	11.6
1963#	23,339	12,118	11,221	10.2	9.0	11.9
1964#	22,559	12,039	10,520	9.7	8.8	11.1
1965#	22,652	12,376	10,276	9.7	9.2	10.4
1966#	23,350	12,976	10,374	9.9	9.4	10.5
1967#	22,609	12,816	9,793	9.6	9.0	10.5
1968#	23,986	13,468	10,518	10.2	9.3	11.7

* Deaths per 1,000 population. See footnote on Table 11 for source of population estimates.

On basis of place of residence.

TABLE 13
MATERNAL DEATHS
MISSISSIPPI
1913-1968

By Year, by Race, with Rates, Occurrence Data 1913-1943,
Residence Data 1944-1968

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1913	447	162	285	108.8	74.1	148.2
1914	413	124	289	101.7	58.0	150.0
1915	289	88	201	73.2	41.9	108.7
1916	326	102	224	83.5	50.7	118.5
1917	297	105	192	80.0	55.1	106.2
1918	302	82	220	76.7	41.0	113.5
1919	407	173	234	95.8	78.5	114.4
1920	393	127	266	89.8	56.6	124.8
1921	391	145	246	84.1	61.2	108.0
1922	313	110	203	71.6	51.7	90.5
1923	321	108	213	74.2	51.1	96.3
1924	413	137	276	93.9	61.8	126.7
1925	435	143	292	95.5	64.2	125.3
1926	388	153	235	76.9	62.2	90.9
1927	409	134	275	83.1	55.4	109.8
1928	421	161	260	87.5	68.1	106.2
1929	410	152	258	90.0	68.8	110.0
1930	489	171	318	101.9	73.7	128.3
1931	352	131	221	78.0	60.5	94.1
1932	279	103	176	60.0	47.1	71.6
1933	315	116	199	71.3	56.8	83.7
1934	306	116	190	63.9	51.9	74.3
1935	334	137	197	69.2	62.4	74.8
1936	358	134	224	72.7	60.4	82.8
1937	370	150	220	70.9	64.4	76.2
1938	333	116	217	62.0	48.2	73.1
1939	311	97	214	59.5	41.3	74.3
1940	327	106	221	62.2	45.9	75.1
1941	293	93	200	53.7	38.5	65.8
1942	248	77	171	43.9	30.4	54.8
1943	233	66	167	39.2	23.9	52.5
1944#	225	72	153	39.8	28.1	49.4
1945#	216	64	152	40.3	27.3	50.5
1946#	199	67	132	32.4	22.5	41.6
1947#	175	50	125	26.4	15.4	36.9
1948#	177	38	139	27.1	12.8	39.0
1949#	147	32	115	22.2	11.1	30.8

Table 13 - Page 2
Maternal Deaths - Mississippi - 1913-1968

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1950#	165	30	135	25.6	11.0	36.2
1951#	143	28	115	21.8	10.1	30.5
1952#	119	15	104	18.4	5.3	28.5
1953#	119	24	95	18.6	8.5	26.6
1954#	104	11	93	16.1	3.8	25.8
1955#	76	12	64	12.0	4.4	17.9
1956#	74	12	62	11.9	4.4	17.8
1957#	75	14	61	12.4	5.2	18.0
1958#	53	10	43	9.0	3.7	13.3
1959#	55	9	46	9.1	3.2	14.1
1960#	59	8	51	10.0	2.9	15.9
1961#	49	9	40	8.2	3.3	12.5
1962#	55	7	48	9.3	2.5	15.3
1963#	44	9	35	7.7	3.4	11.5
1964#	57	9	48	10.1	3.5	15.6
1965#	40	3	37	7.8	1.3	13.3
1966#	27	6	21	5.6	2.6	8.2
1967#	36	10	26	7.7	4.4	10.9
1968#	21	3	18	4.6	1.3	7.8

* Deaths per 10,000 live births.

On basis of place of residence.

TABLE 14
DEATHS UNDER 1 YEAR
MISSISSIPPI
1917-1968

By Year, by Race, with Rates, Occurrence Data 1917-1943,
Residence Data 1944-1968

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1917	3,689	1,296	2,393	99.3	68.0	132.3
1918	**	**	**	**	**	**
1919	3,554	**	**	83.6	**	**
1920	3,499	1,344	2,155	80.0	59.9	101.1
1921	3,157	1,243	1,914	67.9	52.4	84.0
1922	2,826	1,067	1,759	64.6	50.1	78.4
1923	2,908	1,104	1,804	67.2	52.2	81.5
1924	3,137	1,188	1,949	71.4	53.6	89.5
1925	3,176	1,179	1,997	69.7	53.0	85.7
1926	3,578	1,447	2,131	70.9	58.8	82.4
1927	3,291	1,303	1,988	66.9	54.0	79.4
1928	3,567	1,416	2,151	74.1	59.9	87.9
1929	3,275	1,268	2,007	71.9	57.4	85.5
1930	3,276	1,182	2,094	68.3	51.0	84.5
1931	2,547	961	1,586	56.4	44.4	67.5
1932	2,482	965	1,517	53.4	44.1	61.7
1933	2,800	1,035	1,765	63.4	50.7	74.2
1934	3,070	1,199	1,871	64.1	53.7	73.2
1935	2,587	1,044	1,543	53.6	47.5	58.6
1936	2,883	1,105	1,778	58.5	49.8	65.7
1937	3,098	1,174	1,924	59.4	50.4	66.6
1938	3,051	1,205	1,846	56.8	50.1	62.2
1939	2,908	1,135	1,773	55.6	48.3	61.6
1940	2,864	1,070	1,794	54.5	46.4	60.9
1941	2,969	1,113	1,856	54.4	46.1	61.1
1942	2,654	957	1,697	47.0	37.8	54.4
1943	2,774	1,137	1,637	46.7	41.2	51.5
1944#	2,489	974	1,515	44.0	38.0	49.0
1945#	2,192	846	1,346	40.9	36.1	44.7
1946#	2,298	1,005	1,293	37.4	33.8	40.8
1947#	2,428	1,009	1,419	36.6	31.1	41.9
1948#	2,470	988	1,482	37.8	33.3	41.6
1949#	2,612	900	1,712	39.5	31.3	45.8

Table 14 - Page 2
Deaths Under 1 Year - Mississippi - 1917-1968

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1950#	2,348	763	1,585	36.4	28.1	42.6
1951#	2,589	814	1,775	39.5	29.3	47.1
1952#	2,592	760	1,832	40.2	27.1	50.3
1953#	2,534	807	1,727	39.6	28.5	48.3
1954#	2,360	744	1,616	36.4	25.8	44.9
1955#	2,338	680	1,658	37.0	24.7	46.4
1956#	2,353	734	1,619	37.8	26.9	46.4
1957#	2,257	694	1,563	37.2	25.9	46.1
1958#	2,424	706	1,718	41.0	26.3	53.1
1959#	2,355	677	1,678	38.9	24.2	51.5
1960#	2,457	714	1,743	41.5	26.2	54.4
1961#	2,250	645	1,605	37.8	23.5	50.0
1962#	2,353	705	1,648	39.9	25.6	52.5
1963#	2,372	610	1,762	41.4	22.7	57.8
1964#	2,230	599	1,631	39.4	23.1	53.1
1965#	2,126	593	1,533	41.5	25.4	55.1
1966#	1,856	520	1,336	38.5	22.9	52.4
1967#	1,645	512	1,133	35.3	22.6	47.4
1968#	1,613	517	1,096	35.3	22.7	47.8

* Deaths per 1,000 live births.

** Not available.

On basis of place of residence.

TABLE 15
DEATHS UNDER 1 MONTH*
MISSISSIPPI
1924-1968

By Year, by Race, with Rates, Occurrence Data 1924-1944,
Residence Data 1945-1968

Year	Number			Rate**		
	Total	White	Non-white	Total	White	Non-white
1924	1,598	730	868	36.4	32.9	39.8
1925	1,639	701	938	36.0	31.5	40.3
1926	1,731	823	908	34.3	33.4	35.1
1927	1,644	786	858	33.4	32.5	34.3
1928	1,788	804	984	37.2	34.0	40.2
1929	1,662	773	889	36.5	35.0	37.9
1930	1,692	715	977	35.3	30.8	39.4
1931	1,339	590	749	29.7	27.3	31.9
1932	1,361	572	789	29.3	26.1	32.1
1933	1,540	636	904	34.8	31.2	38.0
1934	1,505	644	861	31.4	28.8	33.7
1935	1,446	642	804	29.9	29.2	30.5
1936	1,494	676	818	30.3	30.4	30.2
1937	1,678	755	923	32.2	32.4	32.0
1938	1,688	747	941	31.4	31.1	31.7
1939	1,637	735	902	31.3	31.3	31.3
1940	1,613	680	933	30.7	29.4	31.7
1941	1,663	734	929	30.5	30.4	30.6
1942	1,544	638	906	27.3	25.2	29.0
1943	1,652	791	861	27.8	28.6	27.1
1944	1,436	661	775	25.2	25.4	25.1
1945#	1,351	624	727	25.2	26.6	24.1
1946#	1,561	794	767	25.4	26.7	24.2
1947#	1,595	750	845	24.1	23.1	24.9
1948#	1,674	773	901	25.6	26.0	25.3
1949#	1,594	661	933	24.1	23.0	25.0
1950#	1,446	560	886	22.4	20.6	23.8
1951#	1,679	626	1,053	25.6	22.5	27.9
1952#	1,658	581	1,077	25.7	20.7	29.6
1953#	1,630	635	995	25.4	22.4	27.8
1954#	1,521	601	920	23.5	20.9	25.6
1955#	1,440	527	913	22.8	19.1	25.6
1956#	1,451	581	870	23.3	21.3	24.9
1957#	1,425	540	885	23.5	20.1	26.1
1958#	1,475	540	935	24.9	20.1	28.9
1959#	1,450	515	935	23.9	18.4	28.7
1960#	1,497	550	947	25.3	20.2	29.6
1961#	1,401	509	892	23.5	18.6	27.8
1962#	1,456	550	906	24.7	20.0	28.8
1963#	1,449	470	979	25.3	17.5	32.1
1964#	1,372	460	912	24.2	17.7	29.7
1965#	1,283	470	813	25.1	20.1	29.2
1966#	1,123	386	737	23.3	17.0	28.9
1967#	1,053	388	665	22.6	17.1	27.8
1968#	1,116	424	692	24.4	18.6	30.2

* Figures for 1949-1968 include only the deaths under 28 days.

** Deaths per 1,000 live births.

On basis of place of residence.

TABLE 16
DEATHS FROM HEART DISEASES, MALIGNANT NEOPLASMS,
CEREBROVASCULAR DISEASE, ACCIDENTS, AND INFLUENZA AND PNEUMONIA
(WITH INTERNATIONAL CLASSIFICATION NUMBERS, EIGHTH REVISION)
MISSISSIPPI
1959-1968

By Year, by Race, with Rates, Residence Data

Year	Heart Diseases (390-398, 402, 404-429)			Malignant Neoplasms (140-209)			Cerebrovascular Disease (430-438)			Accidents (800-949)			Influenza & Pneumonia (470-486)		
	Total	White	Non- white	Total	White	Non- white	Total	White	Non- white	Total	White	Non- white	Total	White	Non- white
1959 Number	6,233	3,774	2,459	2,600	1,580	1,020	2,683	1,386	1,297	1,501	837	664	768	328	440
1959 Rate*	279.9	282.5	276.0	116.7	118.3	114.5	120.5	103.7	145.6	67.4	62.6	74.5	34.5	24.6	49.4
1960 Number	6,321	3,882	2,439	2,635	1,644	991	2,960	1,513	1,447	1,504	827	677	958	389	569
1960 Rate*	290.2	308.7	264.9	121.0	130.7	107.6	135.9	120.3	157.2	69.0	65.8	73.5	44.0	30.9	61.8
1961 Number	6,561	4,022	2,539	2,748	1,732	1,016	2,891	1,464	1,427	1,427	760	667	721	307	414
1961 Rate*	301.2	319.8	275.8	126.2	137.7	110.4	132.7	116.4	155.0	65.5	60.4	72.4	33.1	24.4	45.0
1962 Number	7,077	4,206	2,871	2,827	1,793	1,034	3,112	1,552	1,560	1,557	846	711	879	362	517
1962 Rate*	313.4	318.4	306.4	125.2	135.7	110.4	137.8	117.5	166.5	69.0	64.0	75.9	38.9	27.4	55.2
1963 Number	7,300	4,396	2,904	2,945	1,859	1,086	3,154	1,650	1,504	1,571	851	720	1,113	420	693
1963 Rate*	318.8	325.9	308.5	128.6	137.8	115.4	137.7	122.3	159.8	68.6	63.1	76.5	48.6	31.1	73.6
1964 Number	6,981	4,313	2,668	2,968	1,861	1,107	3,077	1,588	1,489	1,612	898	714	791	308	483
1964 Rate*	301.4	314.7	282.2	128.2	135.8	117.1	132.9	115.9	157.5	69.6	65.5	75.5	34.2	22.5	51.1
1965 Number	7,038	4,429	2,609	2,931	1,867	1,064	3,064	1,639	1,425	1,610	881	729	764	350	414
1965 Rate*	302.1	329.4	264.8	125.8	138.9	108.0	131.5	121.9	144.6	69.1	65.5	74.0	32.8	26.0	42.0
1966 Number	7,468	4,729	2,739	3,098	1,990	1,108	3,132	1,681	1,451	1,872	1,072	800	773	341	432
1966 Rate*	316.5	343.8	278.2	131.3	144.7	112.6	132.7	122.2	147.4	79.3	77.9	81.3	32.8	24.8	43.9
1967 Number	7,300	4,640	2,660	3,136	2,061	1,075	2,981	1,632	1,349	1,890	1,166	724	657	310	347
1967 Rate*	309.2	325.2	284.7	132.8	144.4	115.1	126.2	114.4	144.4	80.0	81.7	77.5	27.8	21.7	37.1
1968 Number	7,969	5,024	2,945	3,267	2,101	1,166	3,115	1,631	1,484	1,723	999	724	825	414	411
1968 Rate*	339.0	346.4	327.1	139.0	144.9	129.5	132.5	112.4	164.8	73.3	68.9	80.4	35.1	28.5	45.6

* Deaths per 100,000 population.

TABLE 17
MARRIAGES AND DIVORCES
MISSISSIPPI
1926-1968

By Year, by Race, Occurrence Data

Year	Marriages			Divorces		
	Total	White	Non-white	Total	White	Non-white
1926*	28,712			2,766		
1927*	29,048#	11,402	17,383	2,895		
1928	30,231#	11,662	18,257	3,009	1,435	1,574
1929	31,495#	12,864	18,388	3,172	1,598	1,574
1930	25,728#	10,815	14,751	2,743	1,487	1,256
1931	21,589#	9,952	11,629	2,015	1,308	707
1932	22,688#	10,147	12,453	1,918	1,308	610
1933	27,567#	12,201	15,300	1,955	1,396	559
1934	30,149#	13,497	16,541	2,446	1,704	742
1935	28,722#	12,955	15,635	2,425	1,681	744
1936	32,480#	14,120	18,196	2,520	1,690	830
1937	32,055#	14,802	17,246	2,977	1,984	993
1938	31,495	14,224	17,271	2,916	2,069	847
1939	31,956	15,370	16,586	3,034	2,166	868
1940	34,088	16,834	17,254	3,251	2,295	956
1941	42,142	21,611	20,531	3,831	2,444	1,387
1942	49,565	27,227	22,338	4,326	2,606	1,720
1943	46,030	23,445	22,585	5,851	3,062	2,789
1944	41,559	21,173	20,386	6,742	3,425	3,317
1945	44,871	24,738	20,133	6,727	3,919	2,808
1946	64,204	37,099	27,105	9,495	5,867	3,628
1947	53,584	30,514	23,070	7,004	4,323	2,681
1948	53,645	31,504	22,141	6,697	3,864	2,833
1949	52,765	32,191	20,574	6,285	3,621	2,664
1950	56,738	37,871	18,867	6,065	3,711	2,354
1951	57,321	37,857	19,464	5,464	3,601	1,863
1952	58,116	39,162	18,954	5,419	3,607	1,812
1953	61,751	42,961	18,790	5,204	3,515	1,689
1954	61,381	43,268	18,113	5,001	3,424	1,577
1955	66,423	47,618	18,805	4,845	3,372	1,473
1956	66,223	48,140	18,083	5,027	3,452	1,575
1957	62,495	46,868	15,627	4,974	3,350	1,624
1958	36,402	26,998	9,404	5,016	3,485	1,531
1959	20,447	13,547	6,900	5,108	3,557	1,551
1960	21,233	14,029	7,204	5,260	3,553	1,707
1961	21,610	14,438	7,172	5,213	3,545	1,668
1962	22,007	14,432	7,575	5,467	3,827	1,640
1963	22,140	14,508	7,632	5,452	3,930	1,522
1964	22,784	15,159	7,625	5,631	4,077	1,554
1965	23,343	15,613	7,730	5,730	4,220	1,510
1966	23,476	15,860	7,616	6,306	4,650	1,656
1967	23,535	15,752	7,783	6,455	4,704	1,751
1968	25,355	17,142	8,213	7,020	5,202	1,818


* Total number of marriages for 1926 and total number of divorces for 1926 and 1927 furnished by U.S. Bureau of Census; distribution by race not available.

Includes some records on which the race of the couple was not stated.

BIRTHS, FETAL DEATHS, DEATHS
THE COUNTIES AND THE STATE OF MISSISSIPPI, 1968

Footnote for Tables on Pages 61-143

- * Total births and deaths per 1,000 population. Fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from certain causes of mortality in early infancy per 1,000 live births. Maternal deaths per 10,000 live births. Deaths from other specific causes per 100,000 population.



LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
ADAMS COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,358	666	692	817	391	426	22.2	20.6	24.0
Fetal deaths	82	39	43	48	20	28	58.8	51.2	65.7
Total deaths	493	203	290	368	139	229	10.0	7.3	12.9
Deaths under 1 year	35	10	25	20	6	14	24.5	15.3	32.9
Deaths under 28 days	27	9	18	13	5	8	15.9	12.8	18.8
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	3	0	3	1	0	1	2.7	-	5.6
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	1	0	1	2.7	-	5.6
Malignant neoplasms (140-209)	79	36	43	60	23	37	163.4	121.2	208.3
Benign and unspecified neoplasms (210-239)	1	0	1	1	0	1	2.7	-	5.6
Diabetes mellitus (250)	10	3	7	6	2	4	16.3	10.5	22.5
Avitaminoses and other nutritional deficiencies (260-269)	1	0	1	1	0	1	2.7	-	5.6
Anemias (280-285)	3	0	3	3	0	3	8.2	-	16.9
Diseases of heart (390-398, 402, 404-429)	171	65	106	138	50	88	375.7	263.6	495.5
Hypertension (400, 401, 403)	3	1	2	1	0	1	2.7	-	5.6
Cerebrovascular disease (430-438)	71	31	40	56	21	35	152.5	110.7	197.1
Arteriosclerosis (440)	3	1	2	3	1	2	8.2	5.3	11.3
Influenza and pneumonia (470-486)	18	7	11	15	7	8	40.8	36.9	45.0
Bronchitis, emphysema, asthma (490-493)	7	1	6	5	1	4	13.6	5.3	22.5
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	3	1	2	2	1	1	5.4	5.3	5.6
Cirrhosis of liver (571)	3	3	0	2	2	0	5.4	10.5	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	2	0	2	1	0	1	2.7	-	5.6
Infections of kidney (590)	6	2	4	2	0	2	5.4	-	11.3
Hyperplasia of prostate (600)	0	0	0	1	0	1	2.7	-	5.6
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	7	3	4	4	2	2	10.9	10.5	11.3
Certain causes of mortality in early infancy (760-778)	17	5	12	8	3	5	9.8	7.7	11.7
Accidents (800-949)	35	20	15	20	11	9	54.4	58.0	50.7
Motor vehicle accidents (810-823)	21	14	7	8	6	2	21.8	31.6	11.3
Suicide (950-959)	6	5	1	6	5	1	16.3	26.4	5.6
Homicide (960-978)	8	0	8	9	1	8	24.5	5.3	45.0
Ill-defined and unknown (780-796)	3	0	3	4	0	4	10.9	-	22.5

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
ALCORN COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live Births, Fetal Deaths, Deaths									
Live births	374	293	81	473	387	86	17.1	15.7	28.1
Fetal deaths	8	6	2	8	6	2	16.9	15.5	23.2
Total deaths	294	248	46	324	273	51	11.7	11.1	16.7
Deaths under 1 year	12	3	9	15	6	9	31.7	15.5	104.6
Deaths under 28 days	10	2	8	13	5	8	27.5	12.9	93.0
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	2	2	0	2	2	0	7.2	8.1	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	1	1	0	1	1	0	3.6	4.1	-
Malignant neoplasms (140-209)	34	30	4	41	36	5	148.1	146.2	163.4
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	3	3	0	5	4	1	18.0	16.2	32.7
Avitaminoses and other nutritional deficiencies (260-269)	2	0	2	1	0	1	3.6	-	32.7
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	147	131	16	139	122	17	502.0	495.3	555.6
Hypertension (400, 401, 403)	1	1	0	1	1	0	3.6	4.1	-
Cerebrovascular disease (430-438)	30	25	5	37	32	5	133.6	129.9	163.4
Arteriosclerosis (440)	4	3	1	6	5	1	21.7	20.3	32.7
Influenza and pneumonia (470-486)	6	6	0	7	6	1	25.3	24.4	32.7
Bronchitis, emphysema, asthma (490-493)	3	2	1	4	3	1	14.4	12.2	32.7
Peptic ulcer (531-533)	1	1	0	2	2	0	7.2	8.1	-
Hernia and intestinal obstruction (550-553, 560)	1	1	0	2	1	1	7.2	4.1	32.7
Cirrhosis of liver (571)	4	3	1	6	5	1	21.7	20.3	32.7
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	3	2	1	3	2	1	10.8	8.1	32.7
Infections of kidney (590)	1	1	0	1	1	0	3.6	4.1	-
Hyperplasia of prostate (600)	1	1	0	2	2	0	7.2	8.1	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	1	1	0	3.6	4.1	-
Certain causes of mortality in early infancy (760-778)	8	2	6	11	5	6	23.2	12.9	69.8
Accidents (800-949)	20	16	4	25	22	3	90.3	89.3	98.0
Motor vehicle accidents (810-823)	8	8	0	12	12	0	43.3	48.7	-
Suicide (950-959)	1	1	0	2	2	0	7.2	8.1	-
Homicide (960-978)	2	1	1	3	1	2	10.8	4.1	65.4
Ill-defined and unknown (780-796)	10	7	3	10	7	3	36.1	28.4	98.0

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
AMITE COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	93	1	92	265	82	183	17.4	11.6	22.4
Fetal deaths	2	0	2	10	4	6	37.7	48.8	32.8
Total deaths	70	25	45	158	83	75	10.4	11.8	9.2
Deaths under 1 year	6	0	6	10	2	8	37.7	24.4	43.7
Deaths under 28 days	1	0	1	2	0	2	7.5	-	10.9
Bacillary dysentery amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	1	0	1	6.6	-	12.2
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	3	1	2	17	11	6	111.8	156.2	73.5
Benign and unspecified neoplasms (210-239)	0	0	0	1	1	0	6.6	14.2	-
Diabetes mellitus (250)	0	0	0	1	0	1	6.6	-	12.2
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	25	9	16	49	26	23	322.4	369.3	281.9
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	5	2	3	19	13	6	125.0	184.6	73.5
Arteriosclerosis (440)	0	0	0	2	2	0	13.2	28.4	-
Influenza and pneumonia (470-486)	5	0	5	9	4	5	59.2	56.8	61.3
Bronchitis, emphysema, asthma (490-493)	0	0	0	3	2	1	19.7	28.4	12.2
Peptic ulcer (531-533)	0	0	0	1	1	0	6.6	14.2	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	2	0	2	13.2	-	24.5
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	1	0	1	6.6	-	12.2
Infections of kidney (590)	0	0	0	2	2	0	13.2	28.4	-
Hyperplasia of prostate (600)	0	0	0	1	0	1	6.6	-	12.2
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	0	1	1	0	1	6.6	-	12.2
Certain causes of mortality in early infancy (760-778)	0	0	0	1	0	1	3.8	-	5.5
Accidents (800-949)	8	4	4	14	8	6	92.1	113.6	73.5
Motor vehicle accidents (810-823)	3	2	1	4	3	1	26.3	42.6	12.2
Suicide (950-959)	3	2	1	3	2	1	19.7	28.4	12.2
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	17	6	11	19	6	13	125.0	85.2	159.3

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
ATTALA COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	375	161	214	387	150	237	17.1	11.4	25.0
Fetal deaths	19	11	8	20	9	11	51.7	60.0	46.4
Total deaths	228	130	98	254	141	113	11.2	10.7	11.9
Deaths under 1 year	13	1	12	17	5	12	43.9	33.3	50.6
Deaths under 28 days	6	0	6	11	3	8	28.4	20.0	33.8
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	1	1	0	4.4	7.6	-
Tuberculosis (010-019)	1	0	1	1	0	1	4.4	-	10.6
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	1	0	1	1	0	1	4.4	-	10.6
Malignant neoplasms (140-209)	33	23	10	39	26	13	172.1	197.0	137.4
Benign and unspecified neoplasms (210-239)	1	0	1	1	0	1	4.4	-	10.6
Diabetes mellitus (250)	10	6	4	9	3	6	39.7	22.7	63.4
Avitaminoses and other nutritional deficiencies (260-269)	1	0	1	0	0	0	-	-	-
Anemias (280-285)	3	1	2	3	1	2	13.2	7.6	21.1
Diseases of heart (390-398, 402, 404-429)	85	56	29	88	57	31	388.3	431.8	327.7
Hypertension (400, 401, 403)	1	0	1	2	0	2	8.8	-	21.1
Cerebrovascular disease (430-438)	31	15	16	31	16	15	136.8	121.2	158.6
Arteriosclerosis (440)	5	3	2	5	3	2	22.1	22.7	21.1
Influenza and pneumonia (470-486)	9	4	5	9	3	6	39.7	22.7	63.4
Bronchitis, emphysema, asthma (490-493)	0	0	0	1	1	0	4.4	7.6	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	1	0	1	1	0	1	4.4	-	10.6
Cirrhosis of liver (571)	0	0	0	1	0	1	4.4	-	10.6
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	1	0	1	4.4	-	10.6
Infections of kidney (590)	1	1	0	2	2	0	8.8	15.2	-
Hyperplasia of prostate (600)	1	1	0	0	0	0	-	-	-
Maternal causes (630-678)	1	1	0	1	1	0	25.8	66.7	-
Congenital anomalies (740-759)	1	1	0	3	3	0	13.2	22.7	-
Certain causes of mortality in early infancy (760-778)	4	0	4	7	2	5	18.1	13.3	21.1
Accidents (800-949)	13	9	4	9	4	5	39.7	30.3	52.8
Motor vehicle accidents (810-823)	10	7	3	5	2	3	22.1	15.2	31.7
Suicide (950-959)	3	1	2	3	1	2	13.2	7.6	21.1
Homicide (960-978)	3	0	3	3	0	3	13.2	-	31.7
Ill-defined and unknown (780-796)	11	3	8	15	6	9	66.2	45.4	95.1

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
BENTON COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	56	0	56	135	51	84	18.3	12.1	26.7
Fetal deaths	0	0	0	4	4	0	29.6	78.4	-
Total deaths	29	12	17	86	56	30	11.7	13.3	9.5
Deaths under 1 year	1	0	1	4	1	3	29.6	19.6	35.7
Deaths under 28 days	1	0	1	3	1	2	22.2	19.6	23.8
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	2	1	1	16	14	2	217.1	331.8	63.5
Benign and unspecified neoplasms (210-239)	1	0	1	1	0	1	13.6	-	31.7
Diabetes mellitus (250)	0	0	0	4	4	0	54.3	94.8	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	12	7	5	25	19	6	339.2	450.2	190.5
Hypertension (400, 401, 403)	0	0	0	1	1	0	13.6	23.7	-
Cerebrovascular disease (430-438)	2	0	2	9	3	6	122.1	71.1	190.5
Arteriosclerosis (440)	0	0	0	1	1	0	13.6	23.7	-
Influenza and pneumonia (470-486)	0	0	0	1	1	0	13.6	23.7	-
Bronchitis, emphysema, asthma (490-493)	0	0	0	1	0	1	13.6	-	31.7
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	1	0	1	13.6	-	31.7
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	1	1	0	13.6	23.7	-
Infections of kidney (590)	0	0	0	1	0	1	13.6	-	31.7
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	2	2	0	27.1	47.4	-
Certain causes of mortality in early infancy (760-778)	0	0	0	2	1	1	14.8	19.6	11.9
Accidents (800-949)	3	1	2	5	3	2	67.8	71.1	63.5
Motor vehicle accidents (810-823)	2	1	1	3	2	1	40.7	47.4	31.7
Suicide (950-959)	1	1	0	2	2	0	27.1	47.4	-
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	7	1	6	8	1	7	108.5	23.7	222.2

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
BOLIVAR COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,595	359	1,236	1,404	337	1,067	25.7	17.0	30.6
Fetal deaths	40	4	36	38	8	30	27.1	23.7	28.1
Total deaths	568	118	450	564	151	413	10.3	7.6	11.9
Deaths under 1 year	47	8	39	48	7	41	34.2	20.8	38.4
Deaths under 28 days	32	6	26	31	6	25	22.1	17.8	23.4
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	6	2	4	7	2	5	12.8	10.1	14.4
Tuberculosis (010-019)	2	1	1	2	1	1	3.6	5.0	2.9
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	59	19	40	61	27	34	111.6	136.1	97.7
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	7	0	7	8	2	6	14.6	10.1	17.2
Avitaminoses and other nutritional deficiencies (260-269)	1	0	1	1	0	1	1.8	-	2.9
Anemias (280-285)	2	0	2	1	0	1	1.8	-	2.9
Diseases of heart (390-398, 402, 404-429)	164	42	122	162	50	112	296.4	252.0	321.7
Hypertension (400, 401, 403)	2	0	2	3	1	2	5.5	5.0	5.7
Cerebrovascular disease (430-438)	107	17	90	98	18	80	179.3	90.7	229.8
Arteriosclerosis (440)	8	3	5	7	2	5	12.8	10.1	14.4
Influenza and pneumonia (470-486)	21	3	18	22	4	18	40.2	20.2	51.7
Bronchitis, emphysema, asthma (490-493)	5	1	4	6	3	3	11.0	15.1	8.6
Peptic ulcer (531-533)	0	0	0	1	1	0	1.8	5.0	-
Hernia and intestinal obstruction (550-553, 560)	4	0	4	1	1	0	1.8	5.0	-
Cirrhosis of liver (571)	4	0	4	5	1	4	9.1	5.0	11.5
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	0	1	1	0	1	1.8	-	2.9
Nephritis and nephrosis (580-584)	13	0	13	8	0	8	14.6	-	23.0
Infections of kidney (590)	2	1	1	5	3	2	9.1	15.1	5.7
Hyperplasia of prostate (600)	5	1	4	1	0	1	1.8	-	2.9
Maternal causes (630-678)	1	0	1	1	0	1	7.1	-	9.4
Congenital anomalies (740-759)	4	1	3	4	1	3	7.3	5.0	8.6
Certain causes of mortality in early infancy (760-778)	23	2	21	23	3	20	16.4	8.9	18.7
Accidents (800-949)	41	14	27	46	18	28	84.2	90.7	80.4
Motor vehicle accidents (810-823)	18	6	12	24	11	13	43.9	55.4	37.3
Suicide (950-959)	4	3	1	5	4	1	9.1	20.2	2.9
Homicide (960-978)	8	0	8	7	0	7	12.8	-	20.1
Ill-defined and unknown (780-796)	55	3	52	56	2	54	102.5	10.1	155.1

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
CALHOUN COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	204	94	110	270	175	95	17.1	14.5	25.5
Fetal deaths	15	7	8	15	7	8	55.6	40.0	84.2
Total deaths	130	105	25	183	155	28	11.6	12.8	7.5
Deaths under 1 year	6	3	3	10	7	3	37.0	40.0	31.6
Deaths under 28 days	3	2	1	7	5	2	25.9	28.6	21.0
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	11	9	2	19	16	3	120.3	132.7	80.4
Benign and unspecified neoplasms (210-239)	1	0	1	2	1	1	12.7	8.3	26.8
Diabetes mellitus (250)	6	3	3	2	1	1	12.7	8.3	26.8
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	1	1	0	6.3	8.3	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	60	52	8	73	65	8	462.3	539.0	214.5
Hypertension (400, 401, 403)	2	2	0	1	1	0	6.3	8.3	-
Cerebrovascular disease (430-438)	11	11	0	25	22	3	158.3	182.4	80.4
Arteriosclerosis (440)	3	3	0	3	3	0	19.0	24.9	-
Influenza and pneumonia (470-486)	9	6	3	11	9	2	69.7	74.6	53.6
Bronchitis, emphysema, asthma (490-493)	4	4	0	2	2	0	12.7	16.6	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	0	1	3	2	1	19.0	16.6	26.8
Infections of kidney (590)	0	0	0	1	0	1	6.3	-	26.8
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	1	1	0	6.3	8.3	-
Certain causes of mortality in early infancy (760-778)	2	1	1	5	3	2	18.5	17.1	21.0
Accidents (800-949)	10	9	1	16	15	1	101.3	124.4	26.8
Motor vehicle accidents (810-823)	4	4	0	7	6	1	44.3	49.8	26.8
Suicide (950-959)	1	1	0	1	1	0	6.3	8.3	-
Homicide (960-978)	2	0	2	2	0	2	12.7	-	53.6
Ill-defined and unknown (780-796)	6	3	3	7	4	3	44.3	33.2	80.4

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
CARROLL COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	77	9	68	153	49	104	14.3	9.0	19.7
Fetal deaths	4	1	3	5	2	3	32.7	40.8	28.8
Total deaths	47	10	37	130	62	68	12.2	11.5	12.9
Deaths under 1 year	2	0	2	8	3	5	52.3	61.2	48.1
Deaths under 28 days	0	0	0	3	1	2	19.6	20.4	19.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	6	0	6	21	9	12	196.6	166.4	227.7
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	1	0	1	2	0	2	18.7	-	38.0
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	11	7	4	40	28	12	374.5	517.6	227.7
Hypertension (400, 401, 403)	4	0	4	5	0	5	46.8	-	94.9
Cerebrovascular disease (430-438)	4	1	3	16	10	6	149.8	184.8	113.8
Arteriosclerosis (440)	0	0	0	0	0	0	-	-	-
Influenza and pneumonia (470-486)	0	0	0	3	2	1	28.1	37.0	19.0
Bronchitis, emphysema, asthma (490-493)	0	0	0	0	0	0	-	-	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	1	0	9.4	18.5	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	1	1	0	9.4	18.5	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	1	1	0	9.4	18.5	-
Certain causes of mortality in early infancy (760-778)	0	0	0	3	1	2	19.6	20.4	19.2
Accidents (800-949)	13	1	12	19	3	16	177.9	55.4	303.6
Motor vehicle accidents (810-823)	4	0	4	6	2	4	56.2	37.0	75.9
Suicide (950-959)	1	1	0	1	1	0	9.4	18.5	-
Homicide (960-978)	1	0	1	2	0	2	18.7	-	38.0
Ill-defined and unknown (780-796)	5	0	5	7	2	5	65.5	37.0	94.9

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
CHICKASAW COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
Live Births, Fetal Deaths, Deaths				Total	White	Non-white	Total	White	Non-white
Live births	305	120	185	362	172	190	18.9	14.1	27.5
Fetal deaths	40	20	20	30	16	14	82.9	93.0	73.7
Total deaths	198	137	61	219	149	70	11.4	12.2	10.1
Deaths under 1 year	12	4	8	12	5	7	33.1	29.1	36.8
Deaths under 28 days	7	3	4	6	3	3	16.6	17.4	15.8
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	2	1	1	10.5	8.2	14.5
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	21	17	4	22	16	6	115.1	131.0	87.0
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	0	0	0	2	2	0	10.5	16.4	-
Avitaminoses and other nutritional deficiencies (260-269)	1	1	0	1	1	0	5.2	8.2	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	77	56	21	83	61	22	434.3	499.6	318.8
Hypertension (400, 401, 403)	4	3	1	3	2	1	15.7	16.4	14.5
Cerebrovascular disease (430-438)	39	23	16	39	22	17	204.1	180.2	246.4
Arteriosclerosis (440)	0	0	0	3	2	1	15.7	16.4	14.5
Influenza and pneumonia (470-486)	13	8	5	11	6	5	57.6	49.1	72.5
Bronchitis, emphysema, asthma (490-493)	1	1	0	2	2	0	10.5	16.4	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	0	1	5.2	-	14.5
Cirrhosis of liver (571)	0	0	0	1	1	0	5.2	8.2	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	1	0	2	1	1	10.5	8.2	14.5
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	1	1	0	1	1	0	5.2	8.2	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	2	1	1	4	3	1	20.9	24.6	14.5
Certain causes of mortality in early infancy (760-778)	6	2	4	5	3	2	13.8	17.4	10.5
Accidents (800-949)	17	13	4	17	13	4	89.0	106.5	58.0
Motor vehicle accidents (810-823)	7	5	2	7	5	2	36.6	41.0	29.0
Suicide (950-959)	4	4	0	3	3	0	15.7	24.6	-
Homicide (960-978)	1	0	1	1	0	1	5.2	-	14.5
Ill-defined and unknown (780-796)	4	1	3	5	2	3	26.2	16.4	43.5

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
CHOCTAW COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	141	60	81	135	76	59	15.7	12.0	25.9
Fetal deaths	5	3	2	4	2	2	29.6	26.3	33.9
Total deaths	115	72	43	122	87	35	14.2	13.7	15.4
Deaths under 1 year	3	0	3	3	0	3	22.2	-	50.8
Deaths under 28 days	2	0	2	2	0	2	14.8	-	33.9
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	1	1	0	11.6	15.8	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	11	7	4	13	12	1	151.0	189.6	43.8
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	2	0	2	2	0	2	23.2	-	87.7
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	1	0	1	1	0	1	11.6	-	43.8
Diseases of heart (390-398, 402, 404-429)	49	32	17	47	34	13	545.9	537.1	570.2
Hypertension (400, 401, 403)	1	1	0	4	2	2	46.4	31.6	87.7
Cerebrovascular disease (430-438)	12	9	3	17	12	5	197.4	189.6	219.3
Arteriosclerosis (440)	6	1	5	4	1	3	46.4	15.8	131.6
Influenza and pneumonia (470-486)	7	3	4	8	5	3	92.9	79.0	131.6
Bronchitis, emphysema, asthma (490-493)	1	1	0	1	1	0	11.6	15.8	-
Peptic ulcer (531-533)	1	1	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	1	0	1	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	1	0	1	2	1	1	23.2	15.8	43.8
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	0	1	1	0	1	11.6	-	43.8
Certain causes of mortality in early infancy (760-778)	1	0	1	1	0	1	7.4	-	16.9
Accidents (800-949)	6	6	0	8	8	0	92.9	126.4	-
Motor vehicle accidents (810-823)	2	2	0	2	2	0	23.2	31.6	-
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	3	2	1	3	2	1	34.8	31.6	43.8

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
CLAIBORNE COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	92	12	80	184	38	146	16.2	11.7	17.9
Fetal deaths	2	1	1	11	2	9	59.8	52.6	61.6
Total deaths	96	23	73	123	34	89	10.8	10.5	10.9
Deaths under 1 year	10	2	8	14	2	12	76.1	52.6	82.2
Deaths under 28 days	5	1	4	9	1	8	48.9	26.3	54.8
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	7	2	5	15	7	8	131.8	216.0	98.3
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	3	0	3	5	1	4	43.9	30.9	49.1
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	37	10	27	40	13	27	351.5	401.2	331.7
Hypertension (400, 401, 403)	1	0	1	2	0	2	17.6	-	24.6
Cerebrovascular disease (430-438)	9	4	5	10	4	6	87.9	123.4	73.7
Arteriosclerosis (440)	2	1	1	2	1	1	17.6	30.9	12.3
Influenza and pneumonia (470-486)	8	2	6	7	2	5	61.5	61.7	61.4
Bronchitis, emphysema, asthma (490-493)	0	0	0	0	0	0	-	-	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	1	0	1	1	0	1	8.8	-	12.3
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	0	1	8.8	-	12.3
Nephritis and nephrosis (580-584)	7	2	5	7	2	5	61.5	61.7	61.4
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	1	0	1	1	0	1	8.8	-	12.3
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	0	1	2	0	2	17.6	-	24.6
Certain causes of mortality in early infancy (760-778)	2	1	1	6	1	5	32.6	26.3	34.2
Accidents (800-949)	9	0	9	10	0	10	87.9	-	122.8
Motor vehicle accidents (810-823)	4	0	4	6	0	6	52.7	-	73.7
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	1	0	1	1	0	1	8.8	-	12.3
Ill-defined and unknown (780-796)	5	1	4	6	1	5	52.7	30.9	61.4

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
CLARKE COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live Births, Fetal Deaths, Deaths									
Live births	59	32	27	269	135	134	17.3	13.5	24.2
Fetal deaths	2	1	1	14	8	6	52.0	59.2	44.8
Total deaths	113	68	45	193	122	71	12.4	12.2	12.8
Deaths under 1 year	5	2	3	9	5	4	33.4	37.0	29.8
Deaths under 28 days	3	1	2	5	3	2	18.6	22.2	14.9
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	1	1	0	6.4	10.0	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	13	6	7	29	15	14	186.8	150.3	252.7
Benign and unspecified neoplasms (210-239)	0	0	0	2	1	1	12.9	10.0	18.0
Diabetes mellitus (250)	1	1	0	2	2	0	12.9	20.0	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	1	1	0	6.4	10.0	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	36	23	13	51	35	16	328.6	350.7	288.8
Hypertension (400, 401, 403)	0	0	0	1	0	1	6.4	-	18.0
Cerebrovascular disease (430-438)	25	16	9	41	27	14	264.2	270.5	252.7
Arteriosclerosis (440)	6	4	2	7	4	3	45.1	40.1	54.2
Influenza and pneumonia (470-486)	3	1	2	8	4	4	51.5	40.1	72.2
Bronchitis, emphysema, asthma (490-493)	0	0	0	0	0	0	-	-	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	1	0	6.4	10.0	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	2	1	1	12.9	10.0	18.0
Infections of kidney (590)	0	0	0	2	1	1	12.9	10.0	18.0
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	1	1	0	6.4	10.0	-
Certain causes of mortality in early infancy (760-778)	3	1	2	4	2	2	14.9	14.8	14.9
Accidents (800-949)	8	5	3	16	10	6	103.1	100.2	108.3
Motor vehicle accidents (810-823)	5	3	2	12	7	5	77.3	70.1	90.2
Suicide (950-959)	3	3	0	2	2	0	12.9	20.0	-
Homicide (960-978)	1	0	1	1	0	1	6.4	-	18.0
Ill-defined and unknown (780-796)	9	4	5	10	6	4	64.4	60.1	72.2

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
CLAY COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	271	99	172	375	140	235	20.6	15.6	25.5
Fetal deaths	7	2	5	10	4	6	26.7	28.6	25.5
Total deaths	191	90	101	207	95	112	11.4	10.6	12.2
Deaths under 1 year	11	0	11	12	1	11	32.0	7.1	46.8
Deaths under 28 days	5	0	5	7	0	7	18.7	-	29.8
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	5	0	5	4	1	3	22.0	11.1	32.6
Tuberculosis (010-019)	1	1	0	1	1	0	5.5	11.1	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	19	11	8	28	14	14	153.9	155.9	152.0
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	3	0	3	3	0	3	16.5	-	32.6
Avitaminoses and other nutritional deficiencies (260-269)	1	0	1	1	0	1	5.5	-	10.8
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	78	43	35	73	39	34	401.3	434.3	369.2
Hypertension (400, 401, 403)	3	1	2	6	2	4	33.0	22.3	43.4
Cerebrovascular disease (430-438)	39	18	21	40	18	22	219.9	200.4	238.9
Arteriosclerosis (440)	3	1	2	3	1	2	16.5	11.1	21.7
Influenza and pneumonia (470-486)	5	1	4	6	1	5	33.0	11.1	54.3
Bronchitis, emphysema, asthma (490-493)	1	1	0	1	1	0	5.5	11.1	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	1	1	0	1	1	0	5.5	11.1	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	5	1	4	5	1	4	27.5	11.1	43.4
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	1	0	1	2	1	1	11.0	11.1	10.8
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	1	1	0	5.5	11.1	-
Certain causes of mortality in early infancy (760-778)	1	0	1	3	0	3	8.0	-	12.8
Accidents (800-949)	11	4	7	11	3	8	60.5	33.4	86.9
Motor vehicle accidents (810-823)	5	2	3	6	2	4	33.0	22.3	43.4
Suicide (950-959)	2	1	1	2	1	1	11.0	11.1	10.8
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	9	3	6	9	3	6	49.5	33.4	65.1

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
COAHOMA COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	871	253	618	902	207	695	19.4	12.6	23.1
Fetal deaths	51	20	31	41	13	28	45.4	62.8	40.3
Total deaths	582	224	358	555	184	371	11.9	11.2	12.3
Deaths under 1 year	52	9	43	49	6	43	54.3	29.0	61.9
Deaths under 28 days	37	8	29	35	5	30	38.8	24.2	43.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	1	0	1	2.1	-	3.3
Enteritis and other diarrheal diseases (008, 009)	4	1	3	2	0	2	4.3	-	6.6
Tuberculosis (010-019)	1	0	1	3	1	2	6.4	6.1	6.6
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	74	38	36	76	39	37	163.3	236.6	123.1
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	11	4	7	7	0	7	15.0	-	23.3
Avitaminoses and other nutritional deficiencies (260-269)	1	0	1	2	0	2	4.3	-	6.6
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	193	82	111	179	61	118	384.6	370.1	392.5
Hypertension (400, 401, 403)	5	1	4	5	1	4	10.7	6.1	13.3
Cerebrovascular disease (430-438)	80	31	49	74	20	54	159.0	121.4	179.6
Arteriosclerosis (440)	6	3	3	4	1	3	8.6	6.1	10.0
Influenza and pneumonia (470-486)	17	12	5	17	12	5	36.5	72.8	16.6
Bronchitis, emphysema, asthma (490-493)	7	5	2	6	4	2	12.9	24.3	6.6
Peptic ulcer (531-533)	5	2	3	5	3	2	10.7	18.2	6.6
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	4	3	1	4	3	1	8.6	18.2	3.3
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	0	1	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	4	0	4	4	0	4	8.6	-	13.3
Infections of kidney (590)	13	6	7	10	6	4	21.5	36.4	13.3
Hyperplasia of prostate (600)	1	0	1	1	1	0	2.1	6.1	-
Maternal causes (630-678)	0	0	0	1	0	1	11.1	-	14.4
Congenital anomalies (740-759)	1	0	1	1	0	1	2.1	-	3.3
Certain causes of mortality in early infancy (760-778)	27	8	19	25	5	20	27.7	24.2	28.8
Accidents (800-949)	31	14	17	30	11	19	64.5	66.7	63.2
Motor vehicle accidents (810-823)	19	12	7	12	5	7	25.8	30.3	23.3
Suicide (950-959)	1	1	0	1	1	0	2.1	6.1	-
Homicide (960-978)	3	1	2	5	1	4	10.7	6.1	13.3
Ill-defined and unknown (780-796)	51	0	51	51	0	51	109.6	-	169.7

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
COPIAH COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	583	185	398	468	170	298	17.8	12.3	23.7
Fetal deaths	12	3	9	18	3	15	38.5	17.6	50.3
Total deaths	251	115	136	318	163	155	12.1	11.8	12.3
Deaths under 1 year	12	1	11	18	4	14	38.5	23.5	47.0
Deaths under 28 days	8	1	7	14	4	10	29.9	23.5	33.6
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal disease (008, 009)	4	0	4	4	0	4	15.2	-	31.8
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	31	20	11	49	32	17	185.9	231.9	135.4
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	3	2	1	4	3	1	15.2	21.7	8.0
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	2	0	2	2	0	2	7.6	-	15.9
Diseases of heart (390-398, 402, 404-429)	97	53	44	112	67	45	424.9	485.5	358.3
Hypertension (400, 401, 403)	0	0	0	1	0	1	3.8	-	8.0
Cerebrovascular disease (430-438)	39	15	24	43	18	25	163.1	130.4	199.0
Arteriosclerosis (440)	1	0	1	3	1	2	11.4	7.2	15.9
Influenza and pneumonia (470-486)	9	7	2	15	10	5	56.9	72.5	39.8
Bronchitis, emphysema, asthma (490-493)	2	2	0	2	2	0	7.6	14.5	-
Peptic ulcer (531-533)	2	1	1	2	1	1	7.6	7.2	8.0
Hernia and intestinal obstruction (550-553, 560)	1	0	1	2	1	1	7.6	7.2	8.0
Cirrhosis of liver (571)	1	0	1	1	0	1	3.8	-	8.0
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	1	0	3.8	7.2	-
Nephritis and nephrosis (580-584)	4	2	2	4	2	2	15.2	14.5	15.9
Infections of kidney (590)	2	0	2	2	0	2	7.6	-	15.9
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	2	2	0	7.6	14.5	-
Certain causes of mortality in early infancy (760-778)	4	1	3	7	2	5	15.0	11.8	16.8
Accidents (800-949)	17	5	12	25	11	14	94.8	79.7	111.5
Motor vehicle accidents (810-823)	10	5	5	12	7	5	45.5	50.7	39.8
Suicide (950-959)	1	0	1	1	0	1	3.8	-	8.0
Homicide (960-978)	4	1	3	4	1	3	15.2	7.2	23.9
Ill-defined and unknown (780-796)	18	2	16	19	2	17	72.1	14.5	135.4

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
COVINGTON COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	185	132	53	295	163	132	20.6	16.2	30.8
Fetal deaths	1	1	0	10	2	8	33.9	12.3	60.6
Total deaths	103	69	34	166	117	49	11.6	11.6	11.4
Deaths under 1 year	5	3	2	10	2	8	33.9	12.3	60.6
Deaths under 28 days	3	3	0	7	2	5	23.7	12.3	37.9
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	3	1	2	20.9	10.0	46.7
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	8	6	2	15	12	3	104.7	119.4	70.1
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	0	0	0	2	2	0	14.0	19.9	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	39	32	7	54	46	8	376.8	457.7	186.9
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	17	10	7	27	17	10	188.4	169.2	233.6
Arteriosclerosis (440)	1	1	0	3	2	1	20.9	19.9	23.4
Influenza and pneumonia (470-486)	4	3	1	8	6	2	55.8	59.7	46.7
Bronchitis, emphysema, asthma (490-493)	1	1	0	1	1	0	7.0	10.0	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	2	1	1	14.0	10.0	23.4
Cirrhosis of liver (571)	0	0	0	2	2	0	14.0	19.9	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	2	2	0	2	2	0	14.0	19.9	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	3	2	1	20.9	19.9	23.4
Certain causes of mortality in early infancy (760-778)	3	3	0	5	2	3	16.9	12.3	22.7
Accidents (800-949)	11	6	5	14	9	5	97.7	89.6	116.8
Motor vehicle accidents (810-823)	8	5	3	7	6	1	48.8	59.7	23.4
Suicide (950-959)	1	1	0	1	1	0	7.0	10.0	-
Homicide (960-978)	1	0	1	1	0	1	7.0	-	23.4
Ill-defined and unknown (780-796)	11	1	10	13	2	11	90.7	19.9	257.0

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
DESOTO COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
Live Births, Fetal Deaths, Deaths				Total	White	Non-white	Total	White	Non-white
Live births	290	19	271	781	393	388	21.2	17.4	27.4
Fetal deaths	11	0	11	27	4	23	34.6	10.2	59.3
Total deaths	144	59	85	279	148	131	7.6	6.6	9.2
Deaths under 1 year	7	0	7	25	10	15	32.0	25.4	38.6
Deaths under 28 days	2	0	2	16	9	7	20.5	22.9	18.0
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	2	1	1	3	1	2	8.2	4.4	14.1
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	18	14	4	40	32	8	108.8	141.6	56.4
Benign and unspecified neoplasms (210-239)	1	0	1	1	0	1	2.7	-	7.0
Diabetes mellitus (250)	2	1	1	4	1	3	10.9	4.4	21.2
Avitaminoses and other nutritional deficiencies (260-269)	1	0	1	1	0	1	2.7	-	7.0
Anemias (280-285)	0	0	0	2	1	1	5.4	4.4	7.0
Diseases of heart (390-398, 402, 404-429)	42	17	25	81	49	32	220.3	216.9	225.8
Hypertension (400, 401, 403)	1	0	1	3	0	3	8.2	-	21.2
Cerebrovascular disease (430-438)	15	4	11	40	15	25	108.8	66.4	176.4
Arteriosclerosis (440)	1	0	1	1	0	1	2.7	-	7.0
Influenza and pneumonia (470-486)	6	1	5	6	2	4	16.3	8.8	28.2
Bronchitis, emphysema, asthma (490-493)	0	0	0	5	5	0	13.6	22.1	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	1	1	0	2.7	4.4	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	1	1	0	2	2	0	5.4	8.8	-
Hyperplasia of prostate (600)	1	1	0	1	1	0	2.7	4.4	-
Maternal causes (630-678)	1	0	1	2	1	1	25.6	25.4	25.8
Congenital anomalies (740-759)	0	0	0	5	3	2	13.6	13.3	14.1
Certain causes of mortality in early infancy (760-778)	0	0	0	10	6	4	12.8	15.3	10.3
Accidents (800-949)	20	10	10	22	10	12	59.8	44.3	84.7
Motor vehicle accidents (810-823)	7	6	1	9	6	3	24.5	26.6	21.2
Suicide (950-959)	1	1	0	2	2	0	5.4	8.8	-
Homicide (960-978)	2	2	0	6	4	2	16.3	17.7	14.1
Ill-defined and unknown (780-796)	25	5	20	27	6	21	73.4	26.6	148.2

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
FORREST COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live Births, Fetal Deaths, Deaths									
Live births	1,439	1,061	378	1,119	726	393	18.2	15.9	24.7
Fetal deaths	33	18	15	34	17	17	30.4	23.4	43.2
Total deaths	728	504	224	650	409	241	10.6	9.0	15.1
Deaths under 1 year	40	21	19	39	14	25	34.8	19.3	63.6
Deaths under 28 days	27	17	10	29	11	18	25.9	15.2	45.8
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	3	2	1	2	2	0	3.2	4.4	-
Tuberculosis (010-019)	0	0	0	1	0	1	1.6	-	6.3
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	1	0	1	1	0	1	1.6	-	6.3
Malignant neoplasms (140-209)	124	93	31	100	65	35	162.4	142.4	220.0
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	24	17	7	18	10	8	29.2	21.9	50.3
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	4	3	1	6.5	6.6	6.3
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	262	189	73	234	163	71	380.0	357.0	446.3
Hypertension (400, 401, 403)	9	0	9	8	0	8	13.0	-	50.3
Cerebrovascular disease (430-438)	75	64	11	65	54	11	105.6	118.3	69.1
Arteriosclerosis (440)	9	7	2	6	3	3	9.7	6.6	18.8
Influenza and pneumonia (470-486)	20	11	9	23	9	14	37.4	19.7	88.0
Bronchitis, emphysema, asthma (490-493)	9	8	1	9	8	1	14.6	17.5	6.3
Peptic ulcer (531-533)	2	1	1	2	1	1	3.2	2.2	6.3
Hernia and intestinal obstruction (550-553, 560)	3	2	1	3	2	1	4.9	4.4	6.3
Cirrhosis of liver (571)	2	2	0	3	3	0	4.9	6.6	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	2	1	1	2	1	1	3.2	2.2	6.3
Nephritis and nephrosis (580-584)	6	2	4	8	2	6	13.0	4.4	37.7
Infections of kidney (590)	8	4	4	8	4	4	13.0	8.8	25.1
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	4	3	1	3	2	1	4.9	4.4	6.3
Certain causes of mortality in early infancy (760-778)	19	11	8	22	8	14	19.7	11.0	35.6
Accidents (800-949)	52	37	15	33	19	14	53.6	41.6	88.0
Motor vehicle accidents (810-823)	33	24	9	18	13	5	29.2	28.5	31.4
Suicide (950-959)	5	5	0	6	6	0	9.7	13.1	-
Homicide (960-978)	5	1	4	7	2	5	11.4	4.4	31.4
Ill-defined and unknown (780-796)	35	10	25	39	11	28	63.3	24.1	176.0

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
FRANKLIN COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	66	29	37	124	60	64	13.3	9.6	20.9
Fetal deaths	3	0	3	5	0	5	40.3	-	78.1
Total deaths	86	59	27	114	78	36	12.2	12.5	11.8
Deaths under 1 year	3	2	1	4	2	2	32.2	33.3	31.2
Deaths under 28 days	1	0	1	3	1	2	24.2	16.7	31.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	1	0	1	1	0	10.8	16.0	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	12	5	7	20	10	10	215.0	160.2	326.8
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	1	1	0	3	2	1	32.2	32.0	32.7
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	39	28	11	48	35	13	516.1	560.9	424.8
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	13	8	5	14	9	5	150.5	144.2	163.4
Arteriosclerosis (440)	1	0	1	2	1	1	21.5	16.0	32.7
Influenza and pneumonia (470-486)	7	4	3	8	5	3	86.0	80.1	98.0
Bronchitis, emphysema, asthma (490-493)	2	2	0	3	3	0	32.2	48.1	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	1	1	0	10.8	16.0	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	2	2	0	1	1	0	10.8	16.0	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	2	1	1	21.5	16.0	32.7
Certain causes of mortality in early infancy (760-778)	0	0	0	2	1	1	16.1	16.7	15.6
Accidents (800-949)	3	3	0	3	3	0	32.2	48.1	-
Motor vehicle accidents (810-823)	2	2	0	3	3	0	32.2	48.1	-
Suicide (950-959)	3	3	0	1	1	0	10.8	16.0	-
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	0	0	0	0	0	0	-	-	-

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
GEORGE COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	282	229	53	239	196	43	20.4	18.9	32.1
Fetal deaths	3	2	1	3	1	2	12.6	5.1	46.5
Total deaths	77	65	12	95	79	16	8.1	7.6	11.9
Deaths under 1 year	5	5	0	6	5	1	25.1	25.5	23.2
Deaths under 28 days	4	4	0	4	3	1	16.7	15.3	23.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	15	11	4	20	16	4	170.6	154.1	298.5
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	3	2	1	3	2	1	25.6	19.3	74.6
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	1	1	0	8.5	9.6	-
Anemias (280-285)	2	2	0	2	2	0	17.1	19.3	-
Diseases of heart (390-398, 402, 404-429)	21	18	3	24	21	3	204.8	202.3	223.9
Hypertension (400, 401, 403)	1	1	0	1	1	0	8.5	9.6	-
Cerebrovascular disease (430-438)	7	7	0	10	10	0	85.3	96.3	-
Arteriosclerosis (440)	0	0	0	0	0	0	-	-	-
Influenza and pneumonia (470-486)	1	1	0	1	1	0	8.5	9.6	-
Bronchitis, emphysema, asthma (490-493)	2	2	0	2	2	0	17.1	19.3	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	1	1	0	1	1	0	8.5	9.6	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	1	1	0	8.5	9.6	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	2	2	0	2	2	0	17.1	19.3	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	2	1	1	17.1	9.6	74.6
Certain causes of mortality in early infancy (760-778)	4	4	0	4	3	1	16.7	15.3	23.2
Accidents (800-949)	11	9	2	12	8	4	102.4	77.1	298.5
Motor vehicle accidents (810-823)	4	3	1	5	3	2	42.7	28.9	149.2
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	4	2	2	4	2	2	34.1	19.3	149.2

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
GREENE COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	69	26	43	174	104	70	21.3	16.3	39.3
Fetal deaths	0	0	0	2	2	0	11.5	19.2	-
Total deaths	65	33	32	82	50	32	10.1	7.8	18.0
Deaths under 1 year	3	1	2	7	3	4	40.2	28.8	57.1
Deaths under 28 days	2	0	2	6	2	4	34.5	19.2	57.1
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	1	1	0	12.3	15.7	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	7	4	3	11	7	4	135.0	109.9	224.7
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	0	0	0	2	2	0	24.5	31.4	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	33	18	15	34	19	15	417.2	298.3	842.7
Hypertension (400, 401, 403)	2	1	1	2	1	1	24.5	15.7	56.2
Cerebrovascular disease (430-438)	5	3	2	5	4	1	61.3	62.8	56.2
Arteriosclerosis (440)	0	0	0	0	0	0	-	-	-
Influenza and pneumonia (470-486)	2	2	0	5	4	1	61.3	62.8	56.2
Bronchitis, emphysema, asthma (490-493)	1	0	1	1	0	1	12.3	-	56.2
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	1	0	12.3	15.7	-
Nephritis and nephrosis (580-584)	2	0	2	1	0	1	12.3	-	56.2
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	1	1	0	1	1	0	12.3	15.7	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	2	1	1	24.5	15.7	56.2
Certain causes of mortality in early infancy (760-778)	2	0	2	4	2	2	23.0	19.2	28.6
Accidents (800-949)	4	2	2	7	5	2	85.9	78.5	112.4
Motor vehicle accidents (810-823)	0	0	0	3	1	2	36.8	15.7	112.4
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	1	0	1	1	0	1	12.3	-	56.2
Ill-defined and unknown (780-796)	0	0	0	0	0	0	-	-	-

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
GRENADA COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	384	133	251	427	178	249	21.3	16.5	26.9
Fetal deaths	17	6	11	20	7	13	46.8	39.3	52.2
Total deaths	253	148	105	220	110	110	11.0	10.2	11.9
Deaths under 1 year	18	5	13	17	5	12	39.8	28.1	48.2
Deaths under 28 days	16	5	11	14	4	10	32.8	22.5	40.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	41	23	18	41	20	21	204.5	185.0	227.3
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	2	2	0	3	2	1	15.0	18.5	10.8
Avitaminoses and other nutritional deficiencies (260-269)	1	1	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	62	46	16	52	32	20	259.4	296.0	216.4
Hypertension (400, 401, 403)	4	1	3	3	1	2	15.0	9.2	21.6
Cerebrovascular disease (430-438)	63	26	37	49	12	37	244.4	111.0	400.4
Arteriosclerosis (440)	6	5	1	3	3	0	15.0	27.8	-
Influenza and pneumonia (470-486)	7	7	0	2	2	0	10.0	18.5	-
Bronchitis, emphysema, asthma (490-493)	6	6	0	5	5	0	24.9	46.2	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	1	0	1	5.0	-	10.8
Infections of kidney (590)	3	3	0	3	3	0	15.0	27.8	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	2	1	1	3	2	1	15.0	18.5	10.8
Certain causes of mortality in early infancy (760-778)	10	4	6	8	3	5	18.7	16.8	20.1
Accidents (800-949)	14	8	6	23	17	6	114.7	157.3	64.9
Motor vehicle accidents (810-823)	7	5	2	14	12	2	69.8	111.0	21.6
Suicide (950-959)	1	1	0	0	0	0	-	-	-
Homicide (960-978)	2	1	1	2	1	1	10.0	9.2	10.8
Ill-defined and unknown (780-796)	20	6	14	16	5	11	79.8	46.2	119.0

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
HANCOCK COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	160	118	42	338	278	60	16.2	15.6	19.2
Fetal deaths	1	0	1	4	2	2	11.8	7.2	33.3
Total deaths	150	126	24	170	145	25	8.1	8.2	8.0
Deaths under 1 year	6	5	1	9	8	1	26.6	28.8	16.7
Deaths under 28 days	4	3	1	7	6	1	20.7	21.6	16.7
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	2	2	0	2	2	0	9.6	11.2	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	18	14	4	26	21	5	124.3	118.0	159.7
Benign and unspecified neoplasms (210-239)	0	0	0	1	1	0	4.8	5.6	-
Diabetes mellitus (250)	5	5	0	5	5	0	23.9	28.1	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	1	1	0	2	2	0	9.6	11.2	-
Diseases of heart (390-398, 402, 404-429)	43	38	5	50	43	7	239.0	241.7	223.6
Hypertension (400, 401, 403)	0	0	0	1	1	0	4.8	5.6	-
Cerebrovascular disease (430-438)	18	11	7	19	14	5	90.8	78.7	159.7
Arteriosclerosis (440)	3	3	0	2	2	0	9.6	11.2	-
Influenza and pneumonia (470-486)	2	2	0	2	2	0	9.6	11.2	-
Bronchitis, emphysema, asthma (490-493)	1	0	1	1	0	1	4.8	-	31.9
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	2	1	1	3	1	2	14.3	5.6	63.9
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	1	1	0	4.8	5.6	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	2	2	0	9.6	11.2	-
Certain causes of mortality in early infancy (760-778)	4	3	1	6	5	1	17.8	18.0	16.7
Accidents (800-949)	15	12	3	10	8	2	47.8	45.0	63.9
Motor vehicle accidents (810-823)	5	5	0	4	3	1	19.1	16.9	31.9
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	0	0	0	1	1	0	4.8	5.6	-
Ill-defined and unknown (780-796)	26	24	2	20	18	2	95.6	101.2	63.9

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
HARRISON COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	3,407	2,694	713	3,054	2,372	682	21.6	19.9	30.7
Fetal deaths	289	221	68	258	193	65	84.5	81.4	95.3
Total deaths	1,317	1,033	284	1,110	863	247	7.9	7.2	11.1
Deaths under 1 year	81	55	26	77	50	27	25.2	21.1	39.6
Deaths under 28 days	63	47	16	58	42	16	19.0	17.7	23.5
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	1	0	1	0.7	-	4.5
Tuberculosis (010-019)	4	0	4	4	1	3	2.8	0.8	13.5
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	1	0	1	0	0	0	-	-	-
Malignant neoplasms (140-209)	217	174	43	173	142	31	122.6	119.4	139.7
Benign and unspecified neoplasms (210-239)	5	4	1	5	4	1	3.5	3.4	4.5
Diabetes mellitus (250)	20	16	4	19	15	4	13.5	12.6	18.0
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	3	0	3	3	0	3	2.1	-	13.5
Diseases of heart (390-398, 402, 404-429)	505	431	74	432	371	61	306.2	312.0	274.9
Hypertension (400, 401, 403)	10	6	4	7	4	3	5.0	3.4	13.5
Cerebrovascular disease (430-438)	121	85	36	107	73	34	75.8	61.4	153.2
Arteriosclerosis (440)	10	7	3	11	8	3	7.8	6.7	13.5
Influenza and pneumonia (470-486)	46	37	9	25	17	8	17.7	14.3	36.0
Bronchitis, emphysema, asthma (490-493)	28	25	3	19	16	3	13.5	13.4	13.5
Peptic ulcer (531-533)	7	6	1	4	3	1	2.8	2.5	4.5
Hernia and intestinal obstruction (550-553, 560)	3	1	2	2	1	1	1.4	0.8	4.5
Cirrhosis of liver (571)	24	18	6	11	8	3	7.8	6.7	13.5
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	4	1	3	3	2	1	2.1	1.7	4.5
Infections of kidney (590)	14	11	3	12	9	3	8.5	7.6	13.5
Hyperplasia of prostate (600)	1	1	0	1	1	0	0.7	0.8	-
Maternal causes (630-678)	1	0	1	1	0	1	3.3	-	14.7
Congenital anomalies (740-759)	20	16	4	19	15	4	13.5	12.6	18.0
Certain causes of mortality in early infancy (760-778)	44	34	10	41	31	10	13.4	13.1	14.7
Accidents (800-949)	88	66	22	90	63	27	63.8	53.0	121.7
Motor vehicle accidents (810-823)	39	30	9	43	32	11	30.5	26.9	49.6
Suicide (950-959)	9	9	0	9	9	0	6.4	7.6	-
Homicide (960-978)	14	5	9	12	4	8	8.5	3.4	36.0
Ill-defined and unknown (780-796)	36	22	14	35	20	15	24.8	16.8	67.6

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
HINDS COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	5,618	2,868	2,750	4,391	2,175	2,216	20.3	16.8	25.6
Fetal deaths	580	317	263	404	227	177	92.0	104.4	79.9
Total deaths	2,724	1,532	1,192	1,797	1,000	797	8.3	7.7	9.2
Deaths under 1 year	206	76	130	124	45	79	28.2	20.7	35.6
Deaths under 28 days	154	59	95	99	39	60	22.5	17.9	27.1
Bacillary dysentery and amebiasis (004, 006)	0	0	0	1	0	1	0.5	-	1.2
Enteritis and other diarrheal diseases (008, 009)	12	6	6	8	5	3	3.7	3.8	3.5
Tuberculosis (010-019)	15	6	9	10	4	6	4.6	3.1	6.9
Meningococcal infection (036)	1	1	0	1	1	0	0.5	0.8	-
Syphilis (090-097)	4	1	3	2	0	2	0.9	-	2.3
Malignant neoplasms (140-209)	500	317	183	266	175	91	123.0	134.8	105.3
Benign and unspecified neoplasms (210-239)	15	10	5	8	5	3	3.7	3.8	3.5
Diabetes mellitus (250)	34	15	19	25	13	12	11.6	10.0	13.9
Avitaminoses and other nutritional deficiencies (260-269)	4	1	3	1	0	1	0.5	-	1.2
Anemias (280-285)	7	1	6	4	1	3	1.8	0.8	3.5
Diseases of heart (390-398, 402, 404-429)	667	435	232	475	314	161	219.7	241.9	186.4
Hypertension (400, 401, 403)	23	8	15	11	5	6	5.1	3.8	6.9
Cerebrovascular disease (430-438)	267	140	127	201	99	102	93.0	76.3	118.1
Arteriosclerosis (440)	31	27	4	21	16	5	9.7	12.3	5.8
Influenza and pneumonia (470-486)	80	41	39	63	35	28	29.1	27.0	32.4
Bronchitis, emphysema, asthma (490-493)	44	39	5	33	31	2	15.3	23.9	2.3
Peptic ulcer (531-533)	12	8	4	7	4	3	3.2	3.1	3.5
Hernia and intestinal obstruction (550-553, 560)	12	7	5	7	3	4	3.2	2.3	4.6
Cirrhosis of liver (571)	30	22	8	16	13	3	7.4	10.0	3.5
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	9	7	2	3	2	1	1.4	1.5	1.2
Nephritis and nephrosis (580-584)	27	7	20	16	2	14	7.4	1.5	16.2
Infections of kidney (590)	21	13	8	13	7	6	6.0	5.4	6.9
Hyperplasia of prostate (600)	6	1	5	3	0	3	1.4	-	3.5
Maternal causes (630-678)	2	0	2	1	0	1	2.3	-	4.5
Congenital anomalies (740-759)	43	31	12	18	13	5	8.3	10.0	5.8
Certain causes of mortality in early infancy (760-778)	104	40	64	73	30	43	16.6	13.8	19.4
Accidents (800-949)	213	118	95	144	79	65	66.6	60.8	75.2
Motor vehicle accidents (810-823)	89	62	27	62	44	18	28.7	33.9	20.8
Suicide (950-959)	21	18	3	15	13	2	6.9	10.0	2.3
Homicide (960-978)	32	6	26	25	4	21	11.6	3.1	24.3
Ill-defined and unknown (780-796)	244	73	171	214	63	151	99.0	48.5	174.8

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
HOLMES COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	522	78	444	535	88	447	19.4	10.7	23.1
Fetal deaths	22	3	19	23	2	21	43.0	22.7	47.0
Total deaths	311	84	227	356	107	249	12.9	13.0	12.9
Deaths under 1 year	23	1	22	21	1	20	39.2	11.4	44.7
Deaths under 28 days	15	1	14	15	1	14	28.0	11.4	31.3
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	3	1	2	4	1	3	14.5	12.2	15.5
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	29	11	18	34	14	20	123.4	170.3	103.5
Benign and unspecified neoplasms (210-239)	1	0	1	1	0	1	3.6	-	5.2
Diabetes mellitus (250)	5	0	5	6	1	5	21.8	12.2	25.9
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	103	31	72	110	33	77	399.3	401.4	398.3
Hypertension (400, 401, 403)	3	1	2	5	2	3	18.1	24.3	15.5
Cerebrovascular disease (430-438)	41	12	29	47	15	32	170.6	182.5	165.5
Arteriosclerosis (440)	1	0	1	4	4	0	14.5	48.7	-
Influenza and pneumonia (470-486)	9	2	7	10	2	8	36.3	24.3	41.4
Bronchitis, emphysema, asthma (490-493)	4	4	0	5	4	1	18.1	48.7	5.2
Peptic ulcer (531-533)	3	0	3	3	0	3	10.9	-	15.5
Hernia and intestinal obstruction (550-553, 560)	2	1	1	2	1	1	7.2	12.2	5.2
Cirrhosis of liver (571)	0	0	0	1	1	0	3.6	12.2	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	0	1	3.6	-	5.2
Nephritis and nephrosis (580-584)	2	0	2	2	0	2	7.2	-	10.3
Infections of kidney (590)	2	0	2	3	0	3	10.9	-	15.5
Hyperplasia of prostate (600)	0	0	0	1	1	0	3.6	12.2	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	5	1	4	5	1	4	18.1	12.2	20.7
Certain causes of mortality in early infancy (760-778)	3	0	3	2	0	2	3.7	-	4.5
Accidents (800-949)	29	11	18	34	12	22	123.4	146.0	113.8
Motor vehicle accidents (810-823)	14	9	5	10	6	4	36.3	73.0	20.7
Suicide (950-959)	2	2	0	2	2	0	7.2	24.3	-
Homicide (960-978)	4	1	3	4	1	3	14.5	12.2	15.5
Ill-defined and unknown (780-796)	52	2	50	50	3	47	181.5	36.5	243.1

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
HUMPHREYS COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	377	52	325	393	98	295	23.5	18.1	26.2
Fetal deaths	16	3	13	21	6	15	53.4	61.2	50.8
Total deaths	147	33	114	206	56	150	12.3	10.3	13.3
Deaths under 1 year	15	1	14	16	1	15	40.7	10.2	50.8
Deaths under 28 days	8	0	8	9	0	9	22.9	-	30.5
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	5	2	3	13	3	10	77.8	55.4	88.6
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	1	0	1	5	1	4	29.9	18.4	35.5
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	1	0	1	6.0	-	8.9
Diseases of heart (390-398, 402, 404-429)	62	18	44	76	27	49	455.1	498.2	434.4
Hypertension (400, 401, 403)	3	1	2	4	2	2	24.0	36.9	17.7
Cerebrovascular disease (430-438)	20	3	17	28	6	22	167.7	110.7	195.0
Arteriosclerosis (440)	1	0	1	2	1	1	12.0	18.4	8.9
Influenza and pneumonia (470-486)	4	0	4	4	0	4	24.0	-	35.5
Bronchitis, emphysema, asthma (490-493)	1	0	1	2	0	2	12.0	-	17.7
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	1	1	0	4	3	1	24.0	55.4	8.9
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	3	0	3	4	0	4	24.0	-	35.5
Infections of kidney (590)	1	1	0	1	1	0	6.0	18.4	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	2	0	2	1	0	1	25.4	-	33.9
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	6	0	6	7	0	7	17.8	-	23.7
Accidents (800-949)	6	1	5	16	5	11	95.8	92.2	97.5
Motor vehicle accidents (810-823)	1	0	1	9	4	5	53.9	73.8	44.3
Suicide (950-959)	1	1	0	1	1	0	6.0	18.4	-
Homicide (960-978)	2	2	0	2	2	0	12.0	36.9	-
Ill-defined and unknown (780-796)	21	2	19	26	2	24	155.7	36.9	212.8

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
ISSAQUENA COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	7	0	7	60	14	46	20.8	13.3	25.1
Fetal deaths	1	0	1	3	2	1	50.0	142.8	21.7
Total deaths	16	2	14	32	9	23	11.1	8.6	12.6
Deaths under 1 year	2	0	2	3	0	3	50.0	-	65.2
Deaths under 28 days	0	0	0	0	0	0	-	-	-
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	1	0	1	34.7	-	54.6
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	2	0	2	7	3	4	243.0	285.7	218.6
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	0	0	0	0	0	0	-	-	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	5	2	3	10	4	6	347.2	381.0	327.9
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	2	0	2	5	1	4	173.6	95.2	218.6
Arteriosclerosis (440)	0	0	0	2	1	1	69.4	95.2	54.6
Influenza and pneumonia (470-486)	1	0	1	3	0	3	104.2	-	163.9
Bronchitis, emphysema, asthma (490-493)	0	0	0	0	0	0	-	-	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	0	0	0	0	0	0	-	-	-
Accidents (800-949)	1	0	1	0	0	0	-	-	-
Motor vehicle accidents (810-823)	0	0	0	0	0	0	-	-	-
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	4	0	4	3	0	3	104.2	-	163.9

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
ITAWAMBA COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	39	36	3	228	214	14	12.0	11.8	17.1
Fetal deaths	0	0	0	1	1	0	4.4	4.7	-
Total deaths	80	77	3	145	136	9	7.6	7.5	11.0
Deaths under 1 year	4	4	0	9	7	2	39.5	32.7	142.8
Deaths under 28 days	4	4	0	7	7	0	30.7	32.7	-
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	8	8	0	17	17	0	89.8	93.9	-
Benign and unspecified neoplasms (210-239)	0	0	0	1	1	0	5.3	5.5	-
Diabetes mellitus (250)	3	3	0	3	3	0	15.8	16.6	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	40	38	2	55	54	1	290.5	298.2	122.0
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	12	11	1	24	21	3	126.8	116.0	365.8
Arteriosclerosis (440)	1	1	0	2	2	0	10.6	11.0	-
Influenza and pneumonia (470-486)	0	0	0	5	3	2	26.4	16.6	243.9
Bronchitis, emphysema, asthma (490-493)	0	0	0	2	2	0	10.6	11.0	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	1	1	0	2	2	0	10.6	11.0	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	2	2	0	10.6	11.0	-
Certain causes of mortality in early infancy (760-778)	4	4	0	6	6	0	26.3	28.0	-
Accidents (800-949)	9	9	0	14	13	1	74.0	71.8	122.0
Motor vehicle accidents (810-823)	7	7	0	9	8	1	47.5	44.2	122.0
Suicide (950-959)	1	1	0	0	0	0	-	-	-
Homicide (960-978)	0	0	0	1	0	1	5.3	-	122.0
Ill-defined and unknown (780-796)	0	0	0	1	1	0	5.3	5.5	-

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
JACKSON COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,548	1,142	406	1,864	1,433	431	22.8	21.0	31.9
Fetal deaths	32	19	13	58	45	13	31.1	31.4	30.2
Total deaths	430	310	120	468	346	122	5.7	5.1	9.0
Deaths under 1 year	52	31	21	58	36	22	31.1	25.1	51.0
Deaths under 28 days	45	28	17	49	31	18	26.3	21.6	41.8
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	4	3	1	4	3	1	4.9	4.4	7.4
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	1	0	1	1	0	1	1.2	-	7.4
Malignant neoplasms (140-209)	51	40	11	62	50	12	75.9	73.3	88.8
Benign and unspecified neoplasms (210-239)	1	0	1	2	1	1	2.4	1.5	7.4
Diabetes mellitus (250)	12	9	3	13	10	3	15.9	14.7	22.2
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	1	0	1	1.2	-	7.4
Diseases of heart (390-398, 402, 404-429)	133	102	31	136	105	31	166.5	154.0	229.3
Hypertension (400, 401, 403)	2	1	1	3	2	1	3.7	2.9	7.4
Cerebrovascular disease (430-438)	43	24	19	38	22	16	46.5	32.3	118.3
Arteriosclerosis (440)	5	4	1	5	4	1	6.1	5.9	7.4
Influenza and pneumonia (470-486)	12	5	7	16	8	8	19.6	11.7	59.2
Bronchitis, emphysema, asthma (490-493)	5	5	0	5	5	0	6.1	7.3	-
Peptic ulcer (531-533)	0	0	0	1	1	0	1.2	1.5	-
Hernia and intestinal obstruction (550-553, 560)	2	2	0	2	2	0	2.4	2.9	-
Cirrhosis of liver (571)	2	2	0	7	6	1	8.6	8.8	7.4
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	1	1	0	1.2	1.5	-
Nephritis and nephrosis (580-584)	1	1	0	3	2	1	3.7	2.9	7.4
Infections of kidney (590)	1	1	0	2	2	0	2.4	2.9	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	5	4	1	8	7	1	9.8	10.3	7.4
Certain causes of mortality in early infancy (760-778)	40	24	16	42	25	17	22.5	17.4	39.4
Accidents (800-949)	38	31	7	44	41	3	53.8	60.1	22.2
Motor vehicle accidents (810-823)	20	18	2	26	25	1	31.8	36.7	7.4
Suicide (950-959)	1	1	0	3	3	0	3.7	4.4	-
Homicide (960-978)	6	0	6	10	2	8	12.2	2.9	59.2
Ill-defined and unknown (780-796)	27	22	5	20	15	5	24.5	22.0	37.0

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
JASPER COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	167	111	56	292	106	186	17.5	11.8	24.0
Fetal deaths	7	4	3	12	3	9	41.1	28.3	48.4
Total deaths	158	74	84	200	104	96	12.0	11.6	12.4
Deaths under 1 year	7	2	5	17	3	14	58.2	28.3	75.3
Deaths under 28 days	4	1	3	12	3	9	41.1	28.3	48.4
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	1	0	1	6.0	-	12.9
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	21	8	13	29	18	11	173.4	200.4	142.1
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	4	1	3	7	1	6	41.9	11.1	77.5
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	69	40	29	73	40	33	436.6	445.4	426.4
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	27	9	18	29	15	14	173.4	167.0	180.9
Arteriosclerosis (440)	3	2	1	3	2	1	17.9	22.3	12.9
Influenza and pneumonia (470-486)	3	2	1	3	1	2	17.9	11.1	25.8
Bronchitis, emphysema, asthma (490-493)	1	0	1	3	1	2	17.9	11.1	25.8
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	1	1	0	6.0	11.1	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	3	3	0	2	2	0	12.0	22.3	-
Nephritis and nephrosis (580-584)	0	0	0	2	0	2	12.0	-	25.8
Infections of kidney (590)	4	1	3	4	2	2	23.9	22.3	25.8
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	2	1	1	12.0	11.1	12.9
Certain causes of mortality in early infancy (760-778)	1	0	1	6	2	4	20.5	18.9	21.5
Accidents (800-949)	9	1	8	14	7	7	83.7	78.0	90.4
Motor vehicle accidents (810-823)	5	1	4	6	3	3	35.9	33.4	38.8
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	2	1	1	2	1	1	12.0	11.1	12.9
Ill-defined and unknown (780-796)	2	1	1	6	1	5	35.9	11.1	64.6

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
JEFFERSON COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live Births, Fetal Deaths, Deaths									
Live births	69	8	61	176	20	156	16.4	6.1	21.1
Fetal deaths	1	0	1	3	2	1	17.0	100.0	6.4
Total deaths	71	15	56	121	40	81	11.3	12.1	10.9
Deaths under 1 year	2	0	2	5	0	5	28.4	-	32.0
Deaths under 28 days	1	0	1	4	0	4	22.7	-	25.6
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	1	0	1	9.3	-	13.5
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	11	1	10	17	5	12	158.9	151.5	162.2
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	0	0	0	3	1	2	28.0	30.3	27.0
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	27	6	21	43	15	28	401.9	454.5	378.4
Hypertension (400, 401, 403)	0	0	0	1	0	1	9.3	-	13.5
Cerebrovascular disease (430-438)	11	2	9	20	7	13	186.9	212.1	175.7
Arteriosclerosis (440)	2	1	1	4	3	1	37.4	90.9	13.5
Influenza and pneumonia (470-486)	1	0	1	3	0	3	28.0	-	40.5
Bronchitis, emphysema, asthma (490-493)	1	1	0	3	3	0	28.0	90.9	-
Peptic ulcer (531-533)	1	0	1	1	0	1	9.3	-	13.5
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	1	1	0	1	1	0	9.3	30.3	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	2	1	1	18.7	30.3	13.5
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	1	0	1	3	0	3	17.0	-	19.2
Accidents (800-949)	7	3	4	9	4	5	84.1	121.2	67.6
Motor vehicle accidents (810-823)	2	1	1	3	2	1	28.0	60.6	13.5
Suicide (950-959)	1	0	1	1	0	1	9.3	-	13.5
Homicide (960-978)	1	0	1	1	0	1	9.3	-	13.5
Ill-defined and unknown (780-796)	2	0	2	2	0	2	18.7	-	27.0

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
JEFF DAVIS COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	160	84	76	260	94	166	18.2	14.1	21.9
Fetal deaths	9	5	4	14	7	7	53.8	74.5	42.2
Total deaths	139	67	72	158	71	87	11.1	10.6	11.5
Deaths under 1 year	8	2	6	14	2	12	53.8	21.3	72.3
Deaths under 28 days	4	1	3	11	2	9	42.3	21.3	54.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	1	0	1	1	0	1	7.0	-	13.2
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	19	13	6	22	13	9	154.4	194.9	118.7
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	5	2	3	7	2	5	49.1	30.0	66.0
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	1	0	1	1	0	1	7.0	-	13.2
Diseases of heart (390-398, 402, 404-429)	44	25	19	41	23	18	287.7	344.8	237.5
Hypertension (400, 401, 403)	1	0	1	1	0	1	7.0	-	13.2
Cerebrovascular disease (430-438)	28	11	17	26	10	16	182.4	150.0	211.1
Arteriosclerosis (440)	1	0	1	1	0	1	7.0	-	13.2
Influenza and pneumonia (470-486)	5	1	4	9	4	5	63.2	60.0	66.0
Bronchitis, emphysema, asthma (490-493)	2	2	0	2	2	0	14.0	30.0	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	0	1	7.0	-	13.2
Cirrhosis of liver (571)	1	1	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	1	1	0	1	1	0	7.0	15.0	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	1	1	0	7.0	15.0	-
Certain causes of mortality in early infancy (760-778)	4	1	3	8	2	6	30.8	21.3	36.1
Accidents (800-949)	8	5	3	13	8	5	91.2	119.9	66.0
Motor vehicle accidents (810-823)	4	2	2	6	3	3	42.1	45.0	39.6
Suicide (950-959)	0	0	0	2	1	1	14.0	15.0	13.2
Homicide (960-978)	2	0	2	1	0	1	7.0	-	13.2
Ill-defined and unknown (780-796)	6	0	6	7	0	7	49.1	-	92.3

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
JONES COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,860	801	1,059	954	650	304	16.2	14.6	21.2
Fetal deaths	78	15	63	33	8	25	34.6	12.3	82.2
Total deaths	750	466	284	600	419	181	10.2	9.4	12.6
Deaths under 1 year	95	18	77	38	15	23	39.8	23.1	75.6
Deaths under 28 days	77	17	60	25	13	12	26.2	20.0	39.5
Bacillary dysentery and amebiasis (004, 006)	1	1	0	1	1	0	1.7	2.2	-
Enteritis and other diarrheal diseases (008, 009)	4	1	3	3	1	2	5.1	2.2	14.0
Tuberculosis (010-019)	2	0	2	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	95	72	23	81	63	18	137.4	141.2	125.8
Benign and unspecified neoplasms (210-239)	1	0	1	1	0	1	1.7	-	7.0
Diabetes mellitus (250)	20	10	10	11	7	4	18.7	15.7	28.0
Avitaminoses and other nutritional deficiencies (260-269)	1	1	0	0	0	0	-	-	-
Anemias (280-285)	2	1	1	2	1	1	3.4	2.2	7.0
Diseases of heart (390-398, 402, 404-429)	196	145	51	196	150	46	332.6	336.2	321.4
Hypertension (400, 401, 403)	7	3	4	6	3	3	10.2	6.7	21.0
Cerebrovascular disease (430-438)	76	52	24	69	46	23	117.1	103.1	160.7
Arteriosclerosis (440)	11	8	3	7	6	1	11.9	13.4	7.0
Influenza and pneumonia (470-486)	24	12	12	16	11	5	27.2	24.6	34.9
Bronchitis, emphysema, asthma (490-493)	12	10	2	10	10	0	17.0	22.4	-
Peptic ulcer (531-533)	1	0	1	1	0	1	1.7	-	7.0
Hernia and intestinal obstruction (550-553, 560)	8	5	3	2	2	0	3.4	4.5	-
Cirrhosis of liver (571)	6	5	1	5	5	0	8.5	11.2	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	0	1	1	0	1	1.7	-	7.0
Nephritis and nephrosis (580-584)	14	5	9	8	4	4	13.6	9.0	28.0
Infections of kidney (590)	3	3	0	6	5	1	10.2	11.2	7.0
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	4	1	3	1	0	1	10.5	-	32.9
Congenital anomalies (740-759)	6	1	5	3	1	2	5.1	2.2	14.0
Certain causes of mortality in early infancy (760-778)	55	12	43	17	9	8	17.8	13.8	26.3
Accidents (800-949)	68	43	25	44	29	15	74.7	65.0	104.8
Motor vehicle accidents (810-823)	38	27	11	22	16	6	37.3	35.8	41.9
Suicide (950-959)	9	9	0	6	6	0	10.2	13.4	-
Homicide (960-978)	5	1	4	5	1	4	8.5	2.2	28.0
Ill-defined and unknown (780-796)	68	32	36	60	31	29	101.8	69.5	202.6

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
KEMPER COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	88	10	78	207	46	161	17.9	8.7	25.6
Fetal deaths	2	0	2	6	2	4	29.0	43.5	24.8
Total deaths	80	32	48	116	54	62	10.0	10.2	9.9
Deaths under 1 year	3	0	3	6	0	6	29.0	-	37.3
Deaths under 28 days	1	0	1	3	0	3	14.5	-	18.6
Bacillary dysentery and amebiasis (004, 006)	0	0	0	1	1	0	8.6	18.8	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	1	0	1	8.6	-	15.9
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	6	4	2	15	8	7	129.4	150.6	111.5
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	3	0	3	3	0	3	25.9	-	47.8
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	25	15	10	33	19	14	284.7	357.8	222.9
Hypertension (400, 401, 403)	1	0	1	1	0	1	8.6	-	15.9
Cerebrovascular disease (430-438)	7	1	6	15	7	8	129.4	131.8	127.4
Arteriosclerosis (440)	0	0	0	1	1	0	8.6	18.8	-
Influenza and pneumonia (470-486)	10	3	7	10	3	7	86.3	56.5	111.5
Bronchitis, emphysema, asthma (490-493)	0	0	0	1	1	0	8.6	18.8	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	1	1	0	1	1	0	8.6	18.8	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	1	1	0	4	4	0	34.5	75.3	-
Hyperplasia of prostate (600)	1	0	1	1	0	1	8.6	-	15.9
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	0	1	1	0	1	8.6	-	15.9
Certain causes of mortality in early infancy (760-778)	0	0	0	2	0	2	9.7	-	12.4
Accidents (800-949)	4	1	3	5	2	3	43.1	37.7	47.8
Motor vehicle accidents (810-823)	2	0	2	4	1	3	34.5	18.8	47.8
Suicide (950-959)	2	1	1	1	1	0	8.6	18.8	-
Homicide (960-978)	5	2	3	5	2	3	43.1	37.7	47.8
Ill-defined and unknown (780-796)	9	1	8	8	1	7	69.0	18.8	111.5

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
LAFAYETTE COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	496	310	186	435	251	184	16.6	13.0	26.2
Fetal deaths	19	7	12	22	8	14	50.6	31.9	76.1
Total deaths	207	135	72	220	142	78	8.4	7.4	11.1
Deaths under 1 year	9	4	5	9	6	3	20.7	23.9	16.3
Deaths under 28 days	6	2	4	5	3	2	11.5	12.0	10.9
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal disease (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	1	0	1	2	0	2	7.6	-	28.5
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	24	19	5	32	24	8	122.0	124.9	114.1
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	8	5	3	8	5	3	30.5	26.0	42.8
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	1	0	1	1	0	1	3.8	-	14.3
Diseases of heart (390-398, 402, 404-429)	78	58	20	78	55	23	297.4	286.2	328.1
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	20	8	12	23	10	13	87.7	52.0	185.4
Arteriosclerosis (440)	5	5	0	4	4	0	15.2	20.8	-
Influenza and pneumonia (470-486)	4	2	2	5	2	3	19.1	10.4	42.8
Bronchitis, emphysema, asthma (490-493)	4	4	0	2	2	0	7.6	10.4	-
Peptic ulcer (531-533)	3	1	2	4	2	2	15.2	10.4	28.5
Hernia and intestinal obstruction (550-553, 560)	1	1	0	1	0	1	3.8	-	14.3
Cirrhosis of liver (571)	1	1	0	3	3	0	11.4	15.6	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	4	3	1	3	2	1	11.4	10.4	14.3
Infections of kidney (590)	1	1	0	1	1	0	3.8	5.2	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	1	1	0	3.8	5.2	-
Certain causes of mortality in early infancy (760-778)	4	1	3	3	2	1	6.9	8.0	5.4
Accidents (800-949)	14	11	3	15	12	3	57.2	62.4	42.8
Motor vehicle accidents (810-823)	6	3	3	7	4	3	26.7	20.8	42.8
Suicide (950-959)	3	3	0	4	4	0	15.2	20.8	-
Homicide (960-978)	2	1	1	2	1	1	7.6	5.2	14.3
Ill-defined and unknown (780-796)	18	5	13	17	5	12	64.8	26.0	171.2

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
LAMAR COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	86	58	28	281	222	59	17.8	17.0	21.6
Fetal deaths	1	1	0	5	2	3	17.8	9.0	50.8
Total deaths	85	70	15	150	126	24	9.5	9.6	8.8
Deaths under 1 year	1	1	0	7	4	3	24.9	18.0	50.8
Deaths under 28 days	1	1	0	6	3	3	21.4	13.5	50.8
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	7	5	2	21	18	3	132.7	137.5	109.9
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	0	0	0	2	2	0	12.6	15.3	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	30	26	4	46	41	5	290.8	313.2	183.2
Hypertension (400, 401, 403)	0	0	0	2	1	1	12.6	7.6	36.6
Cerebrovascular disease (430-438)	19	16	3	26	23	3	164.3	175.7	109.9
Arteriosclerosis (440)	5	5	0	5	5	0	31.6	38.2	-
Influenza and pneumonia (470-486)	2	1	1	5	3	2	31.6	22.9	73.3
Bronchitis, emphysema, asthma (490-493)	4	2	2	4	2	2	25.3	15.3	73.3
Peptic ulcer (531-533)	1	1	0	1	1	0	6.3	7.6	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	1	0	6.3	7.6	-
Cirrhosis of liver (571)	0	0	0	2	2	0	12.6	15.3	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	1	1	0	6.3	7.6	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	1	1	0	6	3	3	21.4	13.5	50.8
Accidents (800-949)	5	4	1	11	9	2	69.5	68.8	73.3
Motor vehicle accidents (810-823)	1	1	0	5	4	1	31.6	30.6	36.6
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	0	0	0	1	1	0	6.3	7.6	-
Ill-defined and unknown (780-796)	0	0	0	3	3	0	19.0	22.9	-

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
LAUDERDALE COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,966	975	991	1,412	724	688	18.4	13.8	28.3
Fetal deaths	100	68	32	66	46	20	46.7	63.5	29.1
Total deaths	1,109	740	369	821	520	301	10.7	9.9	12.4
Deaths under 1 year	66	31	35	41	17	24	29.0	23.5	34.9
Deaths under 28 days	49	26	23	30	15	15	21.2	20.7	21.8
Bacillary dysentery and amebiasis (004, 006)	1	1	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	11	6	5	3	1	2	3.9	1.9	8.2
Tuberculosis (010-019)	5	1	4	3	1	2	3.9	1.9	8.2
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	143	99	44	107	69	38	139.2	131.2	156.5
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	15	10	5	12	7	5	15.6	13.3	20.6
Avitaminoses and other nutritional deficiencies (260-269)	8	7	1	2	2	0	2.6	3.8	-
Anemias (280-285)	3	1	2	1	0	1	1.3	-	4.1
Diseases of heart (390-398, 402, 404-429)	331	253	78	255	190	65	331.7	361.2	267.7
Hypertension (400, 401, 403)	8	1	7	3	2	1	3.9	3.8	4.1
Cerebrovascular disease (430-438)	161	105	56	125	78	47	162.6	148.3	193.6
Arteriosclerosis (440)	3	3	0	4	3	1	5.2	5.7	4.1
Influenza and pneumonia (470-486)	52	34	18	32	18	14	41.6	34.2	57.7
Bronchitis, emphysema, asthma (490-493)	17	15	2	13	11	2	16.9	20.9	8.2
Peptic ulcer (531-533)	2	2	0	1	1	0	1.3	1.9	-
Hernia and intestinal obstruction (550-553, 560)	3	2	1	2	1	1	2.6	1.9	4.1
Cirrhosis of liver (571)	14	12	2	11	9	2	14.3	17.1	8.2
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	1	1	0	1.3	1.9	-
Nephritis and nephrosis (580-584)	5	2	3	3	2	1	3.9	3.8	4.1
Infections of kidney (590)	25	21	4	10	8	2	13.0	15.2	8.2
Hyperplasia of prostate (600)	2	1	1	3	1	2	3.9	1.9	8.2
Maternal causes (630-678)	2	0	2	1	0	1	7.1	-	14.5
Congenital anomalies (740-759)	10	7	3	6	4	2	7.8	7.6	8.2
Certain causes of mortality in early infancy (760-778)	34	19	15	19	11	8	13.4	15.2	11.6
Accidents (800-949)	71	46	25	50	32	18	65.0	60.8	74.1
Motor vehicle accidents (810-823)	35	25	10	26	19	7	33.8	36.1	28.8
Suicide (950-959)	8	8	0	8	7	1	10.4	13.3	4.1
Homicide (960-978)	10	1	9	9	0	9	11.7	-	37.1
Ill-defined and unknown (780-796)	98	39	59	96	35	61	124.9	66.5	251.2

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
LAWRENCE COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	124	69	55	212	128	84	18.9	16.6	23.8
Fetal deaths	4	0	4	7	1	6	33.0	7.8	71.4
Total deaths	103	46	57	147	80	67	13.1	10.4	19.0
Deaths under 1 year	7	1	6	7	1	6	33.0	7.8	71.4
Deaths under 28 days	5	0	5	5	0	5	23.6	-	59.5
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	1	1	0	8.9	13.0	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	12	5	7	20	11	9	177.9	142.7	255.0
Benign and unspecified neoplasms (210-239)	1	1	0	1	1	0	8.9	13.0	-
Diabetes mellitus (250)	1	1	0	1	1	0	8.9	13.0	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	36	20	16	46	31	15	409.2	402.1	424.9
Hypertension (400, 401, 403)	0	0	0	1	1	0	8.9	13.0	-
Cerebrovascular disease (430-438)	19	6	13	26	9	17	231.3	116.7	481.6
Arteriosclerosis (440)	3	0	3	6	2	4	53.4	25.9	113.3
Influenza and pneumonia (470-486)	3	0	3	7	3	4	62.3	38.9	113.3
Bronchitis, emphysema, asthma (490-493)	0	0	0	0	0	0	-	-	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	1	1	0	8.9	13.0	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	2	2	0	17.8	25.9	-
Nephritis and nephrosis (580-584)	1	1	0	1	1	0	8.9	13.0	-
Infections of kidney (590)	1	1	0	1	1	0	8.9	13.0	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	4	0	4	4	0	4	18.9	-	47.6
Accidents (800-949)	9	4	5	11	6	5	97.9	77.8	141.6
Motor vehicle accidents (810-823)	6	4	2	8	6	2	71.2	77.8	56.6
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	7	4	3	7	4	3	62.3	51.9	85.0

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
LEAKE COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	178	92	86	301	135	166	15.4	11.4	21.6
Fetal deaths	2	1	1	8	4	4	26.6	29.6	24.1
Total deaths	154	90	64	222	133	89	11.3	11.2	11.6
Deaths under 1 year	6	0	6	13	1	12	43.2	7.4	72.3
Deaths under 28 days	1	0	1	5	1	4	16.6	7.4	24.1
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	1	0	1	5.1	-	13.0
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	18	16	2	34	28	6	173.8	235.7	78.1
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	4	3	1	9	8	1	46.0	67.3	13.0
Avitaminoses and other nutritional deficiencies (260-269)	2	0	2	2	0	2	10.2	-	26.0
Anemias (280-285)	0	0	0	1	0	1	5.1	-	13.0
Diseases of heart (390-398, 402, 404-429)	64	37	27	88	50	38	449.9	420.9	494.8
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	27	13	14	29	14	15	148.3	117.8	195.3
Arteriosclerosis (440)	0	0	0	1	1	0	5.1	8.4	-
Influenza and pneumonia (470-486)	4	2	2	8	4	4	40.9	33.7	52.1
Bronchitis, emphysema, asthma (490-493)	2	2	0	2	2	0	10.2	16.8	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	1	1	0	1	1	0	5.1	8.4	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	0	1	1	0	1	5.1	-	13.0
Infections of kidney (590)	5	4	1	6	5	1	30.7	42.1	13.0
Hyperplasia of prostate (600)	0	0	0	1	1	0	5.1	8.4	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	0	0	0	3	1	2	10.0	7.4	12.0
Accidents (800-949)	15	7	8	21	13	8	107.4	109.4	104.2
Motor vehicle accidents (810-823)	7	3	4	11	7	4	56.2	58.9	52.1
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	3	1	2	3	1	2	15.3	8.4	26.0
Ill-defined and unknown (780-796)	3	1	2	6	1	5	30.7	8.4	65.1

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
LEE COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,293	1,017	276	944	678	266	19.4	17.7	25.8
Fetal deaths	23	14	9	22	12	10	23.3	17.7	37.6
Total deaths	571	409	162	466	325	141	9.6	8.5	13.7
Deaths under 1 year	42	22	20	31	13	18	32.8	19.2	67.7
Deaths under 28 days	31	18	13	23	10	13	24.4	14.7	48.9
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	2	0	2	1	0	1	2.0	-	9.7
Tuberculosis (010-019)	1	0	1	2	0	2	4.1	-	19.4
Meningococcal infection (036)	0	0	0	1	1	0	2.0	2.6	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	94	62	32	66	46	20	135.6	119.9	193.8
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	4	4	0	3	3	0	6.2	7.8	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	218	157	61	196	134	62	402.6	349.3	600.8
Hypertension (400, 401, 403)	1	1	0	1	1	0	2.0	2.6	-
Cerebrovascular disease (430-438)	63	45	18	51	37	14	104.8	96.4	135.6
Arteriosclerosis (440)	8	6	2	4	3	1	8.2	7.8	9.7
Influenza and pneumonia (470-486)	19	13	6	14	10	4	28.8	26.1	38.8
Bronchitis, emphysema, asthma (490-493)	6	5	1	6	5	1	12.3	13.0	9.7
Peptic ulcer (531-533)	8	7	1	4	3	1	8.2	7.8	9.7
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	3	3	0	5	5	0	10.3	13.0	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	2	2	0	1	1	0	2.0	2.6	-
Nephritis and nephrosis (580-584)	1	1	0	1	1	0	2.0	2.6	-
Infections of kidney (590)	1	1	0	1	1	0	2.0	2.6	-
Hyperplasia of prostate (600)	2	1	1	3	1	2	6.2	2.6	19.4
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	8	6	2	8	6	2	16.4	15.6	19.4
Certain causes of mortality in early infancy (760-778)	26	14	12	21	9	12	22.2	13.3	45.1
Accidents (800-949)	50	39	11	38	28	10	78.1	73.0	96.9
Motor vehicle accidents (810-823)	34	29	5	25	20	5	51.4	52.1	48.4
Suicide (950-959)	4	4	0	4	4	0	8.2	10.4	-
Homicide (960-978)	4	1	3	3	1	2	6.2	2.6	19.4
Ill-defined and unknown (780-796)	9	5	4	9	5	4	18.5	13.0	38.8

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
LEFLORE COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,104	346	758	1,017	262	755	21.1	14.4	25.1
Fetal deaths	80	35	45	70	27	43	68.8	103.0	57.0
Total deaths	597	246	351	547	209	338	11.3	11.5	11.2
Deaths under 1 year	55	9	46	50	8	42	49.2	30.5	55.6
Deaths under 28 days	34	4	30	31	5	26	30.5	19.1	34.4
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	4	1	3	3	0	3	6.2	-	10.0
Tuberculosis (010-019)	2	0	2	4	1	3	8.3	5.5	10.0
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	91	52	39	80	41	39	165.8	225.0	129.9
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	14	6	8	13	4	9	26.9	22.0	30.0
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	2	0	2	2	0	2	4.1	-	6.7
Diseases of heart (390-398, 402, 404-429)	180	80	100	170	72	98	352.4	395.2	326.4
Hypertension (400, 401, 403)	9	2	7	4	0	4	8.3	-	13.3
Cerebrovascular disease (430-438)	73	37	36	62	26	36	128.5	142.7	119.9
Arteriosclerosis (440)	2	2	0	2	2	0	4.1	11.0	-
Influenza and pneumonia (470-486)	21	9	12	20	10	10	41.4	54.9	33.3
Bronchitis, emphysema, asthma (490-493)	4	3	1	3	3	0	6.2	16.5	-
Peptic ulcer (531-533)	1	1	0	1	1	0	2.1	5.5	-
Hernia and intestinal obstruction (550-553, 560)	2	0	2	2	0	2	4.1	-	6.7
Cirrhosis of liver (571)	3	2	1	4	3	1	8.3	16.5	3.3
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	0	1	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	3	1	2	2	0	2	4.1	-	6.7
Infections of kidney (590)	8	3	5	4	1	3	8.3	5.5	10.0
Hyperplasia of prostate (600)	1	0	1	1	0	1	2.1	-	3.3
Maternal causes (630-678)	1	0	1	2	0	2	19.7	-	26.5
Congenital anomalies (740-759)	3	3	0	3	2	1	6.2	11.0	3.3
Certain causes of mortality in early infancy (760-778)	24	4	20	21	4	17	20.6	15.3	22.5
Accidents (800-949)	41	12	29	41	16	25	85.0	87.8	83.3
Motor vehicle accidents (810-823)	15	6	9	14	9	5	29.0	49.4	16.6
Suicide (950-959)	3	3	0	3	3	0	6.2	16.5	-
Homicide (960-978)	7	1	6	6	0	6	12.4	-	20.0
Ill-defined and unknown (780-796)	62	7	55	63	8	55	130.6	43.9	183.2

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
LINCOLN COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	522	338	184	457	279	178	17.4	15.4	22.0
Fetal deaths	11	6	5	25	8	17	54.7	28.7	95.5
Total deaths	337	249	88	319	216	103	12.2	11.9	12.7
Deaths under 1 year	14	10	4	18	11	7	39.4	39.4	39.3
Deaths under 28 days	9	8	1	9	7	2	19.7	25.1	11.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	2	0	2	7.6	-	24.7
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	45	32	13	40	27	13	152.6	148.9	160.7
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	5	5	0	4	4	0	15.2	22.1	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	1	0	1	1	0	1	3.8	-	12.4
Diseases of heart (390-398, 402, 404-429)	135	110	25	119	88	31	453.8	485.4	383.2
Hypertension (400, 401, 403)	1	0	1	2	1	1	7.6	5.5	12.4
Cerebrovascular disease (430-438)	43	23	20	42	20	22	160.2	110.3	271.9
Arteriosclerosis (440)	21	18	3	16	14	2	61.0	77.2	24.7
Influenza and pneumonia (470-486)	9	6	3	7	4	3	26.7	22.1	37.1
Bronchitis, emphysema, asthma (490-493)	7	7	0	5	5	0	19.1	27.6	-
Peptic ulcer (531-533)	1	1	0	2	2	0	7.6	11.0	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	0	1	3.8	-	12.4
Cirrhosis of liver (571)	0	0	0	1	1	0	3.8	5.5	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	1	1	0	3.8	5.5	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	3	3	0	5	4	1	19.1	22.1	12.4
Hyperplasia of prostate (600)	0	0	0	1	1	0	3.8	5.5	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	3	3	0	5	5	0	19.1	27.6	-
Certain causes of mortality in early infancy (760-778)	7	6	1	6	5	1	13.1	17.9	5.6
Accidents (800-949)	9	7	2	12	7	5	45.8	38.6	61.8
Motor vehicle accidents (810-823)	4	3	1	6	2	4	22.9	11.0	49.4
Suicide (950-959)	3	3	0	4	4	0	15.2	22.1	-
Homocide (960-978)	3	0	3	3	0	3	11.4	-	37.1
Ill-defined and unknown (780-796)	27	14	13	29	14	15	110.6	77.2	185.4

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
LOWNDES COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live Births, Fetal Deaths, Deaths									
Live births	1,131	644	487	992	491	501	17.5	13.1	25.8
Fetal deaths	29	12	17	29	10	19	29.2	20.4	37.9
Total deaths	489	261	228	498	256	242	8.8	6.8	12.4
Deaths under 1 year	30	10	20	31	9	22	31.2	18.3	43.9
Deaths under 28 days	17	9	8	19	9	10	19.2	18.3	20.0
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	4	0	4	4	0	4	7.0	-	20.6
Tuberculosis (010-019)	4	2	2	4	2	2	7.0	5.4	10.3
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	1	0	1	1	0	1	1.8	-	5.1
Malignant neoplasms (140-209)	62	44	18	60	39	21	105.6	104.4	108.0
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	6	5	1	8	6	2	14.1	16.1	10.3
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	155	95	60	155	91	64	272.9	243.6	329.0
Hypertension (400, 401, 403)	4	3	1	6	3	3	10.6	8.0	15.4
Cerebrovascular disease (430-438)	77	33	44	71	29	42	125.0	77.6	215.9
Arteriosclerosis (440)	5	4	1	5	4	1	8.8	10.7	5.1
Influenza and pneumonia (470-486)	20	5	15	21	6	15	37.0	16.1	77.1
Bronchitis, emphysema, asthma (490-493)	5	5	0	5	5	0	8.8	13.4	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	3	1	2	5	2	3	8.8	5.4	15.4
Cirrhosis of liver (571)	4	2	2	4	2	2	7.0	5.4	10.3
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	1	0	1	1.8	-	5.1
Infections of kidney (590)	2	2	0	2	2	0	3.5	5.4	-
Hyperplasia of prostate (600)	2	2	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	3	0	3	4	1	3	7.0	2.7	15.4
Certain causes of mortality in early infancy (760-778)	13	7	6	14	7	7	14.1	14.2	14.0
Accidents (800-949)	35	22	13	33	24	9	58.1	64.2	46.3
Motor vehicle accidents (810-823)	16	10	6	12	8	4	21.1	21.4	20.6
Suicide (950-959)	3	3	0	2	2	0	3.5	5.4	-
Homicide (960-978)	5	2	3	6	2	4	10.6	5.4	20.6
Ill-defined and unknown (780-796)	50	6	44	53	8	45	93.3	21.4	231.4

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
MADISON COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	525	76	449	772	159	613	23.8	14.7	28.3
Fetal deaths	27	5	22	62	21	41	80.3	132.1	66.9
Total deaths	337	125	212	354	118	236	10.9	10.9	10.9
Deaths under 1 year	13	1	12	18	2	16	23.3	12.6	26.1
Deaths under 28 days	7	1	6	11	1	10	14.2	6.3	16.3
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	1	0	1	1	0	1	3.1	-	4.6
Malignant neoplasms (140-209)	32	14	18	41	18	23	126.2	166.2	106.2
Benign and unspecified neoplasms (210-239)	0	0	0	1	1	0	3.1	9.2	-
Diabetes mellitus (250)	10	4	6	6	2	4	18.5	18.5	18.5
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	1	1	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	120	48	72	132	53	79	406.3	489.4	364.7
Hypertension (400, 401, 403)	5	1	4	6	1	5	18.5	9.2	23.1
Cerebrovascular disease (430-438)	35	9	26	32	9	23	98.5	83.1	106.2
Arteriosclerosis (440)	2	0	2	2	0	2	6.2	-	9.2
Influenza and pneumonia (470-486)	6	2	4	8	3	5	24.6	27.7	23.1
Bronchitis, emphysema, asthma (490-493)	5	4	1	8	5	3	24.6	46.2	13.8
Peptic ulcer (531-533)	0	0	0	1	1	0	3.1	9.2	-
Hernia and intestinal obstruction (550-553, 560)	1	1	0	1	1	0	3.1	9.2	-
Cirrhosis of liver (571)	1	1	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	5	2	3	6	2	4	18.5	18.5	18.5
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	3	0	3	3	0	3	9.2	-	13.8
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	0	1	4	1	3	12.3	9.2	13.8
Certain causes of mortality in early infancy (760-778)	3	1	2	6	1	5	7.8	6.3	8.2
Accidents (800-949)	36	21	15	21	7	14	64.6	64.6	64.6
Motor vehicle accidents (810-823)	19	14	5	8	3	5	24.6	27.7	23.1
Suicide (950-959)	3	3	0	2	2	0	6.2	18.5	-
Homicide (960-978)	1	1	0	3	2	1	9.2	18.5	4.6
Ill-defined and unknown (780-796)	53	6	47	50	3	47	153.9	27.7	217.0

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
MARION COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	324	189	135	444	248	196	19.1	16.1	25.2
Fetal deaths	25	18	7	34	18	16	76.6	72.6	81.6
Total deaths	204	144	60	237	162	75	10.2	10.5	9.6
Deaths under 1 year	7	4	3	12	8	4	27.0	32.2	20.4
Deaths under 28 days	6	4	2	10	8	2	22.5	32.2	10.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	1	0	1	4.3	-	12.8
Tuberculosis (010-019)	1	1	0	2	1	1	8.6	6.5	12.8
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	22	17	5	28	24	4	120.7	155.8	51.3
Benign and unspecified neoplasms (210-239)	0	0	0	1	0	1	4.3	-	12.8
Diabetes mellitus (250)	1	1	0	2	1	1	8.6	6.5	12.8
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	53	41	12	59	43	16	254.4	279.2	205.4
Hypertension (400, 401, 403)	2	1	1	3	1	2	12.9	6.5	25.7
Cerebrovascular disease (430-438)	39	31	8	39	28	11	168.2	181.8	141.2
Arteriosclerosis (440)	1	1	0	1	1	0	4.3	6.5	-
Influenza and pneumonia (470-486)	9	4	5	8	3	5	34.5	19.5	64.2
Bronchitis, emphysema, asthma (490-493)	2	2	0	3	3	0	12.9	19.5	-
Peptic ulcer (531-533)	1	1	0	2	2	0	8.6	13.0	-
Hernia and intestinal obstruction (550-553, 560)	1	1	0	1	1	0	4.3	6.5	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	2	1	1	8.6	6.5	12.8
Infections of kidney (590)	2	2	0	2	2	0	8.6	13.0	-
Hyperplasia of prostate (600)	2	1	1	2	1	1	8.6	6.5	12.8
Maternal causes (630-678)	0	0	0	1	1	0	22.5	40.3	-
Congenital anomalies (740-759)	0	0	0	2	1	1	8.6	6.5	12.8
Certain causes of mortality in early infancy (760-778)	4	2	2	6	4	2	13.5	16.1	10.2
Accidents (800-949)	12	7	5	18	12	6	77.6	77.9	77.0
Motor vehicle accidents (810-823)	7	4	3	9	6	3	38.8	39.0	38.5
Suicide (950-959)	1	1	0	1	1	0	4.3	6.5	-
Homicide (960-978)	5	3	2	3	1	2	12.9	6.5	25.7
Ill-defined and unknown (780-796)	31	15	16	32	16	16	138.0	103.9	205.4

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
MARSHALL COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	334	31	303	531	155	376	22.0	17.8	24.4
Fetal deaths	13	0	13	15	0	15	28.2	-	39.9
Total deaths	192	61	131	255	92	163	10.6	10.6	10.6
Deaths under 1 year	13	2	11	21	6	15	39.5	38.7	39.9
Deaths under 28 days	8	0	8	14	4	10	26.4	25.8	26.6
Bacillary dysentery and amebiasis (004, 006)	1	0	1	1	0	1	4.2	-	6.5
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	15	7	8	35	16	19	145.3	184.3	123.4
Benign and unspecified neoplasms (210-239)	0	0	0	2	1	1	8.3	11.5	6.5
Diabetes mellitus (250)	9	3	6	9	2	7	37.4	23.0	45.4
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	1	0	1	4.2	-	6.5
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	58	20	38	79	35	44	328.1	403.2	285.7
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	34	7	27	42	9	33	174.4	103.7	214.3
Arteriosclerosis (440)	5	1	4	5	1	4	20.8	11.5	26.0
Influenza and pneumonia (470-486)	6	0	6	5	0	5	20.8	-	32.5
Bronchitis, emphysema, asthma (490-493)	4	2	2	4	3	1	16.6	34.6	6.5
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	1	0	1	1	0	1	4.2	-	6.5
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	5	2	3	4	1	3	16.6	11.5	19.5
Infections of kidney (590)	1	0	1	1	1	0	4.2	11.5	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	1	1	0	4.2	11.5	-
Certain causes of mortality in early infancy (760-778)	5	0	5	12	4	8	22.6	25.8	21.3
Accidents (800-949)	20	13	7	17	8	9	70.6	92.2	58.4
Motor vehicle accidents (810-823)	11	10	1	6	4	2	24.9	46.1	13.0
Suicide (950-959)	0	0	0	1	1	0	4.2	11.5	-
Homicide (960-978)	2	0	2	4	0	4	16.6	-	26.0
Ill-defined and unknown (780-796)	20	4	16	20	4	16	83.0	46.1	103.9

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
MONROE COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	787	515	272	646	394	252	18.0	15.8	23.0
Fetal deaths	19	9	10	20	8	12	31.0	20.3	47.6
Total deaths	373	226	147	387	242	145	10.8	9.7	13.2
Deaths under 1 year	23	10	13	20	8	12	31.0	20.3	47.6
Deaths under 28 days	17	9	8	15	8	7	23.2	20.3	27.8
Bacillary dysentery and amebiasis (004, 006)	0	0	0	1	0	1	2.8	-	9.1
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	1	1	0	2.8	4.0	-
Syphilis (090-097)	1	1	0	1	1	0	2.8	4.0	-
Malignant neoplasms (140-209)	41	32	9	41	32	9	114.3	128.5	82.0
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	6	2	4	4	1	3	11.1	4.0	27.3
Avitaminoses and other nutritional deficiencies (260-269)	1	0	1	1	0	1	2.8	-	9.1
Anemias (280-285)	2	1	1	2	1	1	5.6	4.0	9.1
Diseases of heart (390-398, 402, 404-429)	146	96	50	151	101	50	420.8	405.4	455.8
Hypertension (400, 401, 403)	3	2	1	4	2	2	11.1	8.0	18.2
Cerebrovascular disease (430-438)	49	29	20	49	28	21	136.6	112.4	191.4
Arteriosclerosis (440)	11	4	7	13	5	8	36.2	20.1	72.9
Influenza and pneumonia (470-486)	6	2	4	9	4	5	25.1	16.0	45.6
Bronchitis, emphysema, asthma (490-493)	1	1	0	1	1	0	2.8	4.0	-
Peptic ulcer (531-533)	0	0	0	1	1	0	2.8	4.0	-
Hernia and intestinal obstruction (550-553, 560)	4	2	2	3	2	1	8.4	8.0	9.1
Cirrhosis of liver (571)	2	1	1	2	1	1	5.6	4.0	9.1
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	9	3	6	7	3	4	19.5	12.0	36.5
Infections of kidney (590)	3	2	1	4	3	1	11.1	12.0	9.1
Hyperplasia of prostate (600)	2	1	1	2	1	1	5.6	4.0	9.1
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	4	3	1	3	2	1	8.4	8.0	9.1
Certain causes of mortality in early infancy (760-778)	12	7	5	10	6	4	15.5	15.2	15.9
Accidents (800-949)	25	12	13	28	16	12	78.0	64.2	109.4
Motor vehicle accidents (810-823)	12	6	6	13	8	5	36.2	32.1	45.6
Suicide (950-959)	4	3	1	4	3	1	11.1	12.0	9.1
Homicide (960-978)	5	1	4	6	1	5	16.7	4.0	45.6
Ill-defined and unknown (780-796)	12	3	9	13	4	9	36.2	16.0	82.0

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
MONTGOMERY COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	321	153	168	254	111	143	17.4	13.0	23.6
Fetal deaths	4	3	1	2	0	2	7.9	-	14.0
Total deaths	194	127	67	170	108	62	11.6	12.6	10.2
Deaths under 1 year	8	4	4	8	3	5	31.5	27.0	35.0
Deaths under 28 days	5	2	3	6	2	4	23.6	18.0	28.0
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	26	20	6	22	19	3	150.8	222.5	49.6
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	4	0	4	4	0	4	27.4	-	66.1
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	1	1	0	6.8	11.7	-
Diseases of heart (390-398, 402, 404-429)	85	55	30	60	37	23	411.2	433.2	380.2
Hypertension (400, 401, 403)	6	4	2	6	2	4	41.1	23.4	66.1
Cerebrovascular disease (430-438)	29	21	8	22	14	8	150.8	163.9	132.2
Arteriosclerosis (440)	1	1	0	1	1	0	6.8	11.7	-
Influenza and pneumonia (470-486)	6	4	2	5	4	1	34.3	46.8	16.5
Bronchitis, emphysema, asthma (490-493)	0	0	0	1	1	0	6.8	11.7	-
Peptic ulcer (531-533)	1	1	0	1	1	0	6.8	11.7	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	1	0	6.8	11.7	-
Cirrhosis of liver (571)	2	2	0	2	2	0	13.7	23.4	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	1	0	2	1	1	13.7	11.7	16.5
Infections of kidney (590)	1	1	0	1	1	0	6.8	11.7	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	1	1	0	6.8	11.7	-
Certain causes of mortality in early infancy (760-778)	3	1	2	4	1	3	15.7	9.0	21.0
Accidents (800-949)	11	8	3	12	11	1	82.2	128.8	16.5
Motor vehicle accidents (810-823)	7	6	1	7	6	1	48.0	70.2	16.5
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	1	1	0	1	1	0	6.8	11.7	-
Ill-defined and unknown (780-796)	7	0	7	10	0	10	68.5	-	165.3

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
NESHOBIA COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths				Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	351	238	113	376	233	143	18.8	15.7	27.2
Fetal deaths	7	2	5	15	6	9	39.9	25.8	62.9
Total deaths	213	144	69	233	167	66	11.6	11.3	12.6
Deaths under 1 year	14	5	9	15	7	8	39.9	30.0	55.9
Deaths under 28 days	6	4	2	11	6	5	29.2	25.8	35.0
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	2	1	1	3	2	1	15.0	13.5	19.0
Tuberculosis (010-019)	1	1	0	4	2	2	20.0	13.5	38.1
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	41	31	10	43	35	8	214.5	236.5	152.4
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	5	4	1	5	4	1	24.9	27.0	19.0
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	1	0	1	5.0	-	19.0
Anemias (280-285)	1	0	1	1	0	1	5.0	-	19.0
Diseases of heart (390-398, 402, 404-429)	64	48	16	61	49	12	304.2	331.1	228.6
Hypertension (400, 401, 403)	1	1	0	3	2	1	15.0	13.5	19.0
Cerebrovascular disease (430-438)	28	22	6	28	23	5	139.6	155.4	95.2
Arteriosclerosis (440)	3	3	0	1	1	0	5.0	6.8	-
Influenza and pneumonia (470-486)	6	3	3	7	4	3	34.9	27.0	57.1
Bronchitis, emphysema, asthma (490-493)	0	0	0	0	0	0	-	-	-
Peptic ulcer (531-533)	1	1	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	1	1	0	1	1	0	5.0	6.8	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	1	0	5.0	6.8	-
Nephritis and nephrosis (580-584)	3	1	2	2	1	1	10.0	6.8	19.0
Infections of kidney (590)	2	1	1	2	2	0	10.0	13.5	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	1	0	1	26.6	-	69.9
Congenital anomalies (740-759)	1	0	1	1	1	0	5.0	6.8	-
Certain causes of mortality in early infancy (760-778)	6	4	2	10	5	5	26.6	21.4	35.0
Accidents (800-949)	9	5	4	17	13	4	84.8	87.8	76.2
Motor vehicle accidents (810-823)	3	3	0	6	6	0	29.9	40.5	-
Suicide (950-959)	3	3	0	3	3	0	15.0	20.3	-
Homicide (960-978)	4	1	3	6	2	4	29.9	13.5	76.2
Ill-defined and unknown (780-796)	20	7	13	21	8	13	104.7	54.0	247.6

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
NEWTON COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	304	181	123	349	181	168	16.9	12.4	27.8
Fetal deaths	29	19	10	28	18	10	80.2	99.4	59.5
Total deaths	223	147	76	270	188	82	13.1	12.9	13.6
Deaths under 1 year	10	2	8	16	6	10	45.8	33.1	59.5
Deaths under 28 days	5	1	4	9	5	4	25.8	27.6	23.8
Bacillary dysentery and amebiasis (004, 006)	2	1	1	2	1	1	9.7	6.9	16.6
Enteritis and other diarrheal diseases (008, 009)	2	0	2	3	0	3	14.6	-	49.7
Tuberculosis (010-019)	1	1	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	30	22	8	33	24	9	160.2	164.8	149.0
Benign and unspecified neoplasms (210-239)	1	1	0	1	1	0	4.8	6.9	-
Diabetes mellitus (250)	7	4	3	7	4	3	34.0	27.5	49.7
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	1	0	1	4.8	-	16.6
Anemias (280-285)	1	0	1	1	0	1	4.8	-	16.6
Diseases of heart (390-398, 402, 404-429)	70	55	15	94	75	19	456.3	515.1	314.6
Hypertension (400, 401, 403)	4	2	2	3	1	2	14.6	6.9	33.1
Cerebrovascular disease (430-438)	30	18	12	35	22	13	169.9	151.1	215.2
Arteriosclerosis (440)	4	4	0	5	5	0	24.3	34.3	-
Influenza and pneumonia (470-486)	5	3	2	7	5	2	34.0	34.3	33.1
Bronchitis, emphysema, asthma (490-493)	3	3	0	3	3	0	14.6	20.6	-
Peptic ulcer (531-533)	2	2	0	1	1	0	4.8	6.9	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	1	0	4.8	6.9	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	0	1	1	0	1	4.8	-	16.6
Nephritis and nephrosis (580-584)	3	2	1	4	2	2	19.4	13.7	33.1
Infections of kidney (590)	3	1	2	5	2	3	24.3	13.7	49.7
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	1	0	1	0	0	0	-	-	-
Congenital anomalies (740-759)	2	1	1	5	3	2	24.3	20.6	33.1
Certain causes of mortality in early infancy (760-778)	4	0	4	6	2	4	17.2	11.0	23.8
Accidents (800-949)	15	8	7	15	12	3	72.8	82.4	49.7
Motor vehicle accidents (810-823)	11	6	5	12	9	3	58.2	61.8	49.7
Suicide (950-959)	0	0	0	2	2	0	9.7	13.7	-
Homicide (960-978)	3	0	3	3	0	3	14.6	-	49.7
Ill-defined and unknown (780-796)	17	10	7	16	9	7	77.7	61.8	115.9

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
NOXUBEE COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	318	67	251	306	61	245	20.1	13.3	23.1
Fetal deaths	11	0	11	13	1	12	42.5	16.4	49.0
Total deaths	167	42	125	210	61	149	13.8	13.3	14.0
Deaths under 1 year	13	1	12	11	1	10	35.9	16.4	40.8
Deaths under 28 days	9	1	8	7	1	6	22.9	16.4	24.5
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	2	0	2	13.1	-	18.8
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	16	5	11	25	10	15	164.4	217.9	141.2
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	5	1	4	5	1	4	32.9	21.8	37.7
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	41	12	29	55	21	34	361.6	457.5	320.2
Hypertension (400, 401, 403)	2	0	2	3	0	3	19.7	-	28.2
Cerebrovascular disease (430-438)	17	7	10	27	11	16	177.5	239.6	150.6
Arteriosclerosis (440)	1	0	1	1	0	1	6.6	-	9.4
Influenza and pneumonia (470-486)	3	0	3	7	2	5	46.0	43.6	47.1
Bronchitis, emphysema, asthma (490-493)	0	0	0	0	0	0	-	-	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	3	0	3	3	0	3	19.7	-	28.2
Infections of kidney (590)	1	0	1	1	0	1	6.6	-	9.4
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	0	1	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	6	1	5	5	1	4	16.3	16.4	16.3
Accidents (800-949)	10	6	4	12	4	8	78.9	87.1	75.3
Motor vehicle accidents (810-823)	2	1	1	5	2	3	32.9	43.6	28.2
Suicide (950-959)	1	1	0	1	1	0	6.6	21.8	-
Homicide (960-978)	2	0	2	2	0	2	13.1	-	18.8
Ill-defined and unknown (780-796)	52	4	48	51	4	47	335.3	87.1	442.6

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
OKTIBBEHA COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	606	272	334	543	285	258	17.5	14.4	23.0
Fetal deaths	17	7	10	19	8	11	35.0	28.1	42.6
Total deaths	233	109	124	249	119	130	8.0	6.0	11.6
Deaths under 1 year	23	6	17	27	8	19	49.7	28.1	73.6
Deaths under 28 days	16	6	10	17	8	9	31.3	28.1	34.9
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	3	0	3	3	0	3	9.7	-	26.7
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	28	14	14	33	18	15	106.4	90.9	133.7
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	5	3	2	4	2	2	12.9	10.1	17.8
Avitaminoses and other nutritional deficiencies (260-269)	1	0	1	1	0	1	3.2	-	8.9
Anemias (280-285)	1	0	1	1	0	1	3.2	-	8.9
Diseases of heart (390-398, 402, 404-429)	98	52	46	103	55	48	332.0	277.8	427.8
Hypertension (400, 401, 403)	9	0	9	4	0	4	12.9	-	35.6
Cerebrovascular disease (430-438)	26	7	19	29	11	18	93.5	55.6	160.4
Arteriosclerosis (440)	3	2	1	3	2	1	9.7	10.1	8.9
Influenza and pneumonia (470-486)	7	3	4	8	4	4	25.8	20.2	35.6
Bronchitis, emphysema, asthma (490-493)	3	2	1	3	2	1	9.7	10.1	8.9
Peptic ulcer (531-533)	1	1	0	2	2	0	6.4	10.1	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	1	0	1	3.2	-	8.9
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	1	1	0	3.2	5.0	-
Nephritis and nephrosis (580-584)	0	0	0	1	0	1	3.2	-	8.9
Infections of kidney (590)	1	1	0	2	2	0	6.4	10.1	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	4	3	1	5	4	1	16.1	20.2	8.9
Certain causes of mortality in early infancy (760-778)	9	3	6	11	4	7	20.2	14.0	27.1
Accidents (800-949)	14	9	5	13	4	9	41.9	20.2	80.2
Motor vehicle accidents (810-823)	12	9	3	7	3	4	22.6	15.2	35.6
Suicide (950-959)	1	1	0	1	1	0	3.2	5.0	-
Homicide (960-978)	1	1	0	2	1	1	6.4	5.0	8.9
Ill-defined and unknown (780-796)	9	2	7	7	1	6	22.6	5.0	53.5

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
PANOLA COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	634	192	442	610	209	401	20.8	16.1	24.5
Fetal deaths	26	4	22	25	4	21	41.0	19.1	52.4
Total deaths	260	109	151	329	140	189	11.2	10.8	11.5
Deaths under 1 year	27	3	24	27	4	23	44.3	19.1	57.4
Deaths under 28 days	15	3	12	12	3	9	19.7	14.4	22.4
Bacillary dysentery and amebiasis (004, 006)	1	0	1	1	0	1	3.4	-	6.1
Enteritis and other diarrheal diseases (008, 009)	6	0	6	7	1	6	23.9	7.7	36.6
Tuberculosis (010-019)	2	2	0	1	1	0	3.4	7.7	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	41	27	14	55	33	22	187.5	254.6	134.4
Benign and unspecified neoplasms (210-239)	1	1	0	1	1	0	3.4	7.7	-
Diabetes mellitus (250)	4	1	3	3	1	2	10.2	7.7	12.2
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	1	1	0	3.4	7.7	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	92	40	52	108	48	60	368.2	370.4	366.5
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	37	12	25	46	13	33	156.8	100.3	201.6
Arteriosclerosis (440)	5	4	1	5	4	1	17.0	30.9	6.1
Influenza and pneumonia (470-486)	5	2	3	10	5	5	34.1	38.6	30.5
Bronchitis, emphysema, asthma (490-493)	1	1	0	3	3	0	10.2	23.1	-
Peptic ulcer (531-533)	1	0	1	1	0	1	3.4	-	6.1
Hernia and intestinal obstruction (550-553, 560)	0	0	0	2	2	0	6.8	15.4	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	0	1	3	0	3	10.2	-	18.3
Infections of kidney (590)	0	0	0	3	2	1	10.2	15.4	6.1
Hyperplasia of prostate (600)	0	0	0	1	1	0	3.4	7.7	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	0	1	2	0	2	6.8	-	12.2
Certain causes of mortality in early infancy (760-778)	11	3	8	7	3	4	11.5	14.4	10.0
Accidents (800-949)	22	8	14	23	7	16	78.4	54.0	97.7
Motor vehicle accidents (810-823)	7	6	1	9	7	2	30.7	54.0	12.2
Suicide (950-959)	3	2	1	3	2	1	10.2	15.4	6.1
Homicide (960-978)	4	1	3	9	1	8	30.7	7.7	48.9
Ill-defined and unknown (780-796)	11	1	10	12	1	11	40.9	7.7	67.2

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
PEARL RIVER COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	597	476	121	555	387	168	19.1	16.1	32.9
Fetal deaths	8	5	3	11	5	6	19.8	12.9	35.7
Total deaths	218	165	53	278	210	68	9.6	8.8	13.3
Deaths under 1 year	17	9	8	21	8	13	37.8	20.7	77.4
Deaths under 28 days	14	8	6	19	8	11	34.2	20.7	65.5
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	2	2	0	6.9	8.3	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	30	27	3	40	35	5	137.6	146.0	97.8
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	4	1	3	5	1	4	17.2	4.2	78.3
Avitaminoses and other nutritional deficiencies (260-269)	2	1	1	2	1	1	6.9	4.2	19.6
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	94	76	18	103	84	19	354.2	350.4	371.8
Hypertension (400, 401, 403)	0	0	0	1	0	1	3.4	-	19.6
Cerebrovascular disease (430-438)	24	15	9	37	24	13	127.2	100.1	254.4
Arteriosclerosis (440)	6	4	2	6	4	2	20.6	16.7	39.1
Influenza and pneumonia (470-486)	7	5	2	8	6	2	27.5	25.0	39.1
Bronchitis, emphysema, asthma (490-493)	0	0	0	0	0	0	-	-	-
Peptic ulcer (531-533)	0	0	0	1	1	0	3.4	4.2	-
Hernia and intestinal obstruction (550-553, 560)	1	0	1	1	0	1	3.4	-	19.6
Cirrhosis of liver (571)	1	1	0	2	1	1	6.9	4.2	19.6
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	1	1	0	3.4	4.2	-
Infections of kidney (590)	4	3	1	5	4	1	17.2	16.7	19.6
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	1	0	1	18.0	-	59.5
Congenital anomalies (740-759)	4	3	1	2	1	1	6.9	4.2	19.6
Certain causes of mortality in early infancy (760-778)	11	7	4	16	7	9	28.8	18.1	53.6
Accidents (800-949)	13	10	3	19	17	2	65.3	70.9	39.1
Motor vehicle accidents (810-823)	7	4	3	10	8	2	34.4	33.4	39.1
Suicide (950-959)	1	1	0	1	1	0	3.4	4.2	-
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	9	4	5	11	6	5	37.8	25.0	97.8

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
PERRY COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	104	75	29	191	125	66	20.9	19.5	24.3
Fetal deaths	2	2	0	5	4	1	26.2	32.0	15.2
Total deaths	73	54	19	103	70	33	11.3	10.9	12.1
Deaths under 1 year	5	3	2	9	4	5	47.1	32.0	75.8
Deaths under 28 days	3	3	0	7	4	3	36.6	32.0	45.4
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	1	1	0	11.0	15.6	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	6	6	0	10	10	0	109.6	156.2	-
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	0	0	0	2	1	1	21.9	15.6	36.8
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	24	20	4	33	26	7	361.8	406.2	257.4
Hypertension (400, 401, 403)	2	1	1	3	1	2	32.9	15.6	73.5
Cerebrovascular disease (430-438)	16	13	3	20	16	4	219.3	250.0	147.0
Arteriosclerosis (440)	0	0	0	1	1	0	11.0	15.6	-
Influenza and pneumonia (470-486)	5	3	2	5	3	2	54.8	46.9	73.5
Bronchitis, emphysema, asthma (490-493)	1	1	0	1	1	0	11.0	15.6	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	1	0	1	1	0	1	11.0	-	36.8
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	0	1	1	0	1	11.0	-	36.8
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	1	0	1	52.4	-	151.5
Congenital anomalies (740-759)	0	0	0	1	1	0	11.0	15.6	-
Certain causes of mortality in early infancy (760-778)	3	3	0	6	3	3	31.4	24.0	45.4
Accidents (800-949)	3	1	2	6	1	5	65.8	15.6	183.8
Motor vehicle accidents (810-823)	3	1	2	4	1	3	43.8	15.6	110.3
Suicide (950-959)	1	1	0	0	0	0	-	-	-
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	5	2	3	5	2	3	54.8	31.2	110.3

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
PIKE COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	624	325	299	583	261	322	18.0	14.2	23.0
Fetal deaths	16	6	10	18	4	14	30.9	15.3	43.5
Total deaths	414	234	180	454	251	203	14.0	13.6	14.5
Deaths under 1 year	22	4	18	22	2	20	37.7	7.7	62.1
Deaths under 28 days	19	4	15	18	2	16	30.9	7.7	49.7
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	2	1	1	2	1	1	6.2	5.4	7.1
Tuberculosis (010-019)	0	0	0	1	0	1	3.1	-	7.1
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	45	30	15	64	45	19	197.5	244.3	135.8
Benign and unspecified neoplasms (210-239)	0	0	0	1	1	0	3.1	5.4	-
Diabetes mellitus (250)	7	2	5	9	2	7	27.8	10.8	50.0
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	2	1	1	2	1	1	6.2	5.4	7.1
Diseases of heart (390-398, 402, 404-429)	146	92	54	159	97	62	490.6	526.6	443.2
Hypertension (400, 401, 403)	3	1	2	3	1	2	9.2	5.4	14.3
Cerebrovascular disease (430-438)	60	40	20	61	38	23	188.2	206.3	164.4
Arteriosclerosis (440)	6	5	1	7	6	1	21.6	32.6	7.1
Influenza and pneumonia (470-486)	20	16	4	15	10	5	46.3	54.3	35.7
Bronchitis, emphysema, asthma (490-493)	2	2	0	5	4	1	15.4	21.7	7.1
Peptic ulcer (531-533)	3	2	1	2	1	1	6.2	5.4	7.1
Hernia and intestinal obstruction (550-553, 560)	1	0	1	1	0	1	3.1	-	7.1
Cirrhosis of liver (571)	1	1	0	2	2	0	6.2	10.8	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	2	1	1	2	1	1	6.2	5.4	7.1
Nephritis and nephrosis (580-584)	1	0	1	1	0	1	3.1	-	7.1
Infections of kidney (590)	1	1	0	1	1	0	3.1	5.4	-
Hyperplasia of prostate (600)	1	0	1	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	3	1	2	2	0	2	6.2	-	14.3
Certain causes of mortality in early infancy (760-778)	15	3	12	15	2	13	25.7	7.7	40.4
Accidents (800-949)	18	9	9	25	11	14	77.1	59.7	100.1
Motor vehicle accidents (810-823)	7	7	0	8	7	1	24.7	38.0	7.1
Suicide (950-959)	2	1	1	2	1	1	6.2	5.4	7.1
Homicide (960-978)	4	1	3	4	1	3	12.3	5.4	21.4
Ill-defined and unknown (780-796)	48	11	37	46	12	34	141.9	65.1	243.0

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
PONTOTOC COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	294	214	80	276	211	65	15.2	14.4	18.7
Fetal deaths	8	8	0	9	8	1	32.6	37.9	15.4
Total deaths	203	171	32	209	173	36	11.5	11.8	10.4
Deaths under 1 year	13	5	8	12	6	6	43.5	28.4	92.3
Deaths under 28 days	9	5	4	8	5	3	29.0	23.7	46.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	2	1	1	11.0	6.8	28.8
Tuberculosis (010-019)	0	0	0	1	0	1	5.5	-	28.8
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	16	14	2	22	17	5	121.3	115.9	144.1
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	2	1	1	2	1	1	11.0	6.8	28.8
Avitaminoses and other nutritional deficiencies (260-269)	1	0	1	1	0	1	5.5	-	28.8
Anemias (280-285)	1	0	1	1	0	1	5.5	-	28.8
Diseases of heart (390-398, 402, 404-429)	96	85	11	88	78	10	485.1	531.7	288.2
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	44	40	4	42	37	5	231.5	252.2	144.1
Arteriosclerosis (440)	0	0	0	0	0	0	-	-	-
Influenza and pneumonia (470-486)	5	4	1	5	4	1	27.6	27.3	28.8
Bronchitis, emphysema, asthma (490-493)	1	1	0	3	3	0	16.5	20.4	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	1	0	5.5	6.8	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	2	2	0	1	1	0	5.5	6.8	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	2	2	0	11.0	13.6	-
Certain causes of mortality in early infancy (760-778)	7	4	3	6	4	2	21.7	19.0	30.8
Accidents (800-949)	9	8	1	11	8	3	60.6	54.5	86.4
Motor vehicle accidents (810-823)	4	4	0	4	4	0	22.0	27.3	-
Suicide (950-959)	1	1	0	1	1	0	5.5	6.8	-
Homicide (960-978)	3	2	1	3	2	1	16.5	13.6	28.8
Ill-defined and unknown (780-796)	6	2	4	9	5	4	49.6	34.1	115.3

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
PRENTISS COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	406	324	82	352	277	75	16.4	14.5	31.8
Fetal deaths	5	5	0	6	6	0	17.0	21.7	-
Total deaths	240	212	28	245	219	26	11.4	11.4	11.0
Deaths under 1 year	8	4	4	9	5	4	25.6	18.0	53.3
Deaths under 28 days	5	2	3	5	2	3	14.2	7.2	40.0
Bacillary dysentery and amebiasis (004, 006)	0	0	0	1	1	0	4.6	5.2	-
Enteritis and other diarrheal diseases (008, 009)	1	1	0	1	1	0	4.6	5.2	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	30	26	4	35	29	6	163.0	151.8	254.2
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	5	3	2	4	3	1	18.6	15.7	42.4
Avitaminoses and other nutritional deficiencies (260-269)	1	1	0	1	1	0	4.6	5.2	-
Anemias (280-285)	1	1	0	1	1	0	4.6	5.2	-
Diseases of heart (390-398, 402, 404-429)	97	90	7	95	89	6	442.5	465.7	254.2
Hypertension (400, 401, 403)	6	5	1	6	5	1	27.9	26.2	42.4
Cerebrovascular disease (430-438)	33	29	4	30	26	4	139.7	136.0	169.5
Arteriosclerosis (440)	6	6	0	3	3	0	14.0	15.7	-
Influenza and pneumonia (470-486)	11	9	2	13	11	2	60.5	57.6	84.7
Bronchitis, emphysema, asthma (490-493)	5	5	0	3	3	0	14.0	15.7	-
Peptic ulcer (531-533)	0	0	0	2	2	0	9.3	10.5	-
Hernia and intestinal obstruction (550-553, 560)	2	0	2	1	0	1	4.6	-	42.4
Cirrhosis of liver (571)	0	0	0	1	1	0	4.6	5.2	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	1	0	4.6	5.2	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	1	1	0	1	1	0	4.6	5.2	-
Hyperplasia of prostate (600)	2	2	0	2	2	0	9.3	10.5	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	2	2	0	9.3	10.5	-
Certain causes of mortality in early infancy (760-778)	3	2	1	1	0	1	2.8	-	13.3
Accidents (800-949)	12	11	1	16	16	0	74.5	83.7	-
Motor vehicle accidents (810-823)	6	5	1	10	10	0	46.6	52.3	-
Suicide (950-959)	2	2	0	2	2	0	9.3	10.5	-
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	10	7	3	9	6	3	41.9	31.4	127.1

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
QUITMAN COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	459	126	333	385	102	283	21.3	14.1	26.1
Fetal deaths	18	2	16	22	3	19	57.1	29.4	67.1
Total deaths	131	32	99	223	80	143	12.3	11.0	13.2
Deaths under 1 year	15	1	14	18	2	16	46.8	19.6	56.5
Deaths under 28 days	10	1	9	11	2	9	28.6	19.6	31.8
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	2	0	2	11.1	-	18.5
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	11	5	6	20	8	12	110.7	110.5	110.8
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	3	1	2	9	5	4	49.8	69.1	36.9
Avitaminoses and other nutritional deficiencies (260-269)	1	1	0	1	1	0	5.5	13.8	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	38	11	27	68	26	42	376.3	359.1	387.8
Hypertension (400, 401, 403)	1	0	1	1	0	1	5.5	-	9.2
Cerebrovascular disease (430-438)	12	4	8	29	13	16	160.5	179.6	147.7
Arteriosclerosis (440)	2	0	2	6	2	4	33.2	27.6	36.9
Influenza and pneumonia (470-486)	8	0	8	11	3	8	60.9	41.4	73.9
Bronchitis, emphysema, asthma (490-493)	1	1	0	3	3	0	16.6	41.4	-
Peptic ulcer (531-533)	0	0	0	1	0	1	5.5	-	9.2
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	1	1	0	1	1	0	5.5	13.8	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	0	1	5.5	-	9.2
Nephritis and nephrosis (580-584)	1	0	1	2	0	2	11.1	-	18.5
Infections of kidney (590)	0	0	0	3	1	2	16.6	13.8	18.5
Hyperplasia of prostate (600)	1	0	1	2	0	2	11.1	-	18.5
Maternal causes (630-678)	1	0	1	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	1	1	0	5.5	13.8	-
Certain causes of mortality in early infancy (760-778)	8	0	8	9	1	8	23.4	9.8	28.3
Accidents (800-949)	17	5	12	24	12	12	132.8	165.7	110.8
Motor vehicle accidents (810-823)	14	5	9	16	8	8	88.5	110.5	73.9
Suicide (950-959)	1	1	0	1	1	0	5.5	13.8	-
Homicide (960-978)	3	0	3	1	0	1	5.5	-	9.2
Ill-defined and unknown (780-796)	14	0	14	16	0	16	88.5	-	147.7

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
RANKIN COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live Births, Fetal Deaths, Deaths									
Live births	78	35	43	755	496	259	17.7	16.6	20.2
Fetal deaths	3	1	2	50	42	8	66.2	84.7	30.9
Total deaths	439	259	180	268	183	85	6.3	6.1	6.6
Deaths under 1 year	6	2	4	19	10	9	25.2	20.2	34.7
Deaths under 28 days	1	1	0	13	8	5	17.2	16.1	19.3
Bacillary dysentery and amebiasis (004, 006)	4	1	3	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	1	0	1	2.3	-	7.8
Tuberculosis (010-019)	6	3	3	1	1	0	2.3	3.3	-
Meningococcal infection (036)	3	3	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	45	21	24	40	28	12	93.7	93.7	93.8
Benign and unspecified neoplasms (210-239)	1	1	0	0	0	0	-	-	-
Diabetes mellitus (250)	13	6	7	10	5	5	23.4	16.7	39.1
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	1	0	1	2.3	-	7.8
Anemias (280-285)	1	0	1	1	1	0	2.3	3.3	-
Diseases of heart (390-398, 402, 404-429)	108	71	37	78	54	24	182.7	180.7	187.5
Hypertension (400, 401, 403)	1	1	0	1	1	0	2.3	3.3	-
Cerebrovascular disease (430-438)	44	20	24	27	20	7	63.2	66.9	54.7
Arteriosclerosis (440)	10	5	5	5	4	1	11.7	13.4	7.8
Influenza and pneumonia (470-486)	51	38	13	17	12	5	39.8	40.1	39.1
Bronchitis, emphysema, asthma (490-493)	9	6	3	4	2	2	9.4	6.7	15.6
Peptic ulcer (531-533)	1	1	0	2	1	1	4.7	3.3	7.8
Hernia and intestinal obstruction (550-553, 560)	7	4	3	1	1	0	2.3	3.3	-
Cirrhosis of liver (571)	3	3	0	3	2	1	7.0	6.7	7.8
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	0	1	0	0	0	-	-	-
Infections of kidney (590)	2	2	0	1	1	0	2.3	3.3	-
Hyperplasia of prostate (600)	1	1	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	1	1	0	2.3	3.3	-
Certain causes of mortality in early infancy (760-778)	1	1	0	7	4	3	9.3	8.1	11.6
Accidents (800-949)	39	25	14	34	26	8	79.6	87.0	62.5
Motor vehicle accidents (810-823)	3	3	0	18	15	3	42.2	50.2	23.4
Suicide (950-959)	4	4	0	2	2	0	4.7	6.7	-
Homicide (960-978)	2	0	2	0	0	0	-	-	-
Ill-defined and unknown (780-796)	33	10	23	18	8	10	42.2	26.8	78.1

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
SCOTT COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	411	285	126	437	256	181	18.2	16.0	22.7
Fetal deaths	11	7	4	14	8	6	32.0	31.2	33.1
Total deaths	223	144	79	252	163	89	10.5	10.2	11.2
Deaths under 1 year	17	5	12	17	6	11	38.9	23.4	60.8
Deaths under 28 days	10	4	6	10	5	5	22.9	19.5	27.6
Bacillary dysentery and amebiasis (004, 006)	1	0	1	1	0	1	4.2	-	12.5
Enteritis and other diarrheal diseases (008, 009)	3	2	1	4	2	2	16.6	12.5	25.1
Tuberculosis (010-019)	0	0	0	1	1	0	4.2	6.2	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	22	16	6	29	20	9	120.8	124.8	112.8
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	0	0	0	1	0	1	4.2	-	12.5
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	118	79	39	117	78	39	487.3	486.6	488.7
Hypertension (400, 401, 403)	2	1	1	2	1	1	8.3	6.2	12.5
Cerebrovascular disease (430-438)	18	11	7	21	13	8	87.5	81.1	100.2
Arteriosclerosis (440)	4	1	3	5	2	3	20.8	12.5	37.6
Influenza and pneumonia (470-486)	4	3	1	5	5	0	20.8	31.2	-
Bronchitis, emphysema, asthma (490-493)	3	3	0	4	4	0	16.6	25.0	-
Peptic ulcer (531-533)	0	0	0	1	1	0	4.2	6.2	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	1	0	4.2	6.2	-
Cirrhosis of liver (571)	1	0	1	1	0	1	4.2	-	12.5
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	1	0	4.2	6.2	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	1	0	1	2	1	1	8.3	6.2	12.5
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	8	4	4	9	5	4	20.6	19.5	22.1
Accidents (800-949)	15	11	4	20	12	8	83.3	74.8	100.2
Motor vehicle accidents (810-823)	9	6	3	9	6	3	37.5	37.4	37.6
Suicide (950-959)	1	1	0	1	1	0	4.2	6.2	-
Homicide (960-978)	1	1	0	1	1	0	4.2	6.2	-
Ill-defined and unknown (780-796)	14	7	7	14	7	7	58.3	43.7	87.7

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
SHARKEY COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	88	11	77	224	42	182	23.5	12.6	29.4
Fetal deaths	2	0	2	13	2	11	58.0	47.6	60.4
Total deaths	88	20	68	108	27	81	11.3	8.1	13.1
Deaths under 1 year	6	2	4	10	2	8	44.6	47.6	44.0
Deaths under 28 days	3	1	2	6	2	4	26.8	47.6	22.0
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	6	2	4	6	1	5	63.0	30.1	80.6
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	2	0	2	2	0	2	21.0	-	32.2
Avitaminoses and other nutritional deficiencies (260-269)	1	0	1	1	0	1	10.5	-	16.1
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	29	9	20	33	10	23	346.6	301.2	371.0
Hypertension (400, 401, 403)	0	0	0	1	1	0	10.5	30.1	-
Cerebrovascular disease (430-438)	17	1	16	16	3	13	168.1	90.4	209.7
Arteriosclerosis (440)	0	0	0	0	0	0	-	-	-
Influenza and pneumonia (470-486)	11	1	10	9	1	8	94.5	30.1	129.0
Bronchitis, emphysema, asthma (490-493)	1	0	1	1	0	1	10.5	-	16.1
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	1	0	1	2	0	2	21.0	-	32.2
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	1	0	1	10.5	-	16.1
Certain causes of mortality in early infancy (760-778)	3	1	2	6	2	4	26.8	47.6	22.0
Accidents (800-949)	7	3	4	8	3	5	84.0	90.4	80.6
Motor vehicle accidents (810-823)	2	1	1	3	1	2	31.5	30.1	32.2
Suicide (950-959)	1	0	1	1	0	1	10.5	-	16.1
Homicide (960-978)	2	1	1	3	1	2	31.5	30.1	32.2
Ill-defined and unknown (780-796)	4	0	4	8	0	8	84.0	-	129.0

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
SIMPSON COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	277	179	98	396	210	186	19.6	14.2	34.2
Fetal deaths	8	4	4	18	10	8	45.4	47.6	43.0
Total deaths	226	152	74	238	169	69	11.8	11.4	12.7
Deaths under 1 year	8	1	7	14	3	11	35.4	14.3	59.1
Deaths under 28 days	5	1	4	10	3	7	25.2	14.3	37.6
Bacillary dysentery and amebiasis (004, 006)	1	0	1	1	0	1	4.9	-	18.4
Enteritis and other diarrheal diseases (008, 009)	0	0	0	1	0	1	4.9	-	18.4
Tuberculosis (010-019)	19	10	9	3	2	1	14.8	13.5	18.4
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	27	23	4	29	24	5	143.4	162.2	92.1
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	2	2	0	1	1	0	4.9	6.8	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	1	0	1	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	81	61	20	88	68	20	435.0	459.4	368.3
Hypertension (400, 401, 403)	2	2	0	2	2	0	9.9	13.5	-
Cerebrovascular disease (430-438)	23	11	12	26	16	10	128.5	108.1	184.2
Arteriosclerosis (440)	3	2	1	6	5	1	29.6	33.8	18.4
Influenza and pneumonia (470-486)	5	3	2	7	5	2	34.6	33.8	36.8
Bronchitis, emphysema, asthma (490-493)	4	4	0	0	0	0	-	-	-
Peptic ulcer (531-533)	2	2	0	3	3	0	14.8	20.3	-
Hernia and intestinal obstruction (550-553, 560)	1	1	0	3	2	1	14.8	13.5	18.4
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	0	1	2	1	1	9.9	6.8	18.4
Infections of kidney (590)	3	1	2	3	2	1	14.8	13.5	18.4
Hyperplasia of prostate (600)	1	1	0	1	1	0	4.9	6.8	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	1	1	0	4.9	6.8	-
Certain causes of mortality in early infancy (760-778)	3	0	3	6	1	5	15.2	4.8	26.9
Accidents (800-949)	17	13	4	20	16	4	98.9	108.1	73.7
Motor vehicle accidents (810-823)	5	4	1	7	6	1	34.6	40.5	18.4
Suicide (950-959)	2	2	0	3	3	0	14.8	20.3	-
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	12	5	7	14	6	8	69.2	40.5	147.3

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
SMITH COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
Live Births, Fetal Deaths, Deaths				Total	White	Non-white	Total	White	Non-white
Live births	13	0	13	271	151	120	20.8	15.7	35.5
Fetal deaths	1	1	0	6	3	3	22.1	19.9	25.0
Total deaths	68	52	16	155	120	35	11.9	12.5	10.4
Deaths under 1 year	5	0	5	16	4	12	59.0	26.5	100.0
Deaths under 28 days	2	0	2	11	3	8	40.6	19.9	66.7
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	5	3	2	21	16	5	161.5	166.3	147.9
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	2	0	2	5	3	2	38.5	31.2	59.2
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	28	26	2	55	53	2	423.1	550.9	59.2
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	7	6	1	18	13	5	138.5	135.1	147.9
Arteriosclerosis (440)	4	3	1	7	6	1	53.8	62.4	29.6
Influenza and pneumonia (470-486)	3	0	3	5	1	4	38.5	10.4	118.3
Bronchitis, emphysema, asthma (490-493)	2	2	0	4	3	1	30.8	31.2	29.6
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	2	2	0	15.4	20.8	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	1	0	7.7	10.4	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	1	1	0	1	1	0	7.7	10.4	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	1	0	1	9	3	6	33.2	19.9	50.0
Accidents (800-949)	8	5	3	13	8	5	100.0	83.2	147.9
Motor vehicle accidents (810-823)	5	4	1	8	5	3	61.5	52.0	88.8
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	5	4	1	6	4	2	46.2	41.6	59.2

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
STONE COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	138	97	41	154	102	52	19.0	16.4	27.4
Fetal deaths	1	1	0	4	3	1	26.0	29.4	19.2
Total deaths	67	49	18	83	56	27	10.2	9.0	14.2
Deaths under 1 year	1	0	1	7	2	5	45.4	19.6	96.2
Deaths under 28 days	0	0	0	4	2	2	26.0	19.6	38.5
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	1	0	2	1	1	24.7	16.1	52.6
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	4	4	0	10	9	1	123.3	144.9	52.6
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	1	1	0	3	3	0	37.0	48.3	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	1	1	0	12.3	16.1	-
Diseases of heart (390-398, 402, 404-429)	28	24	4	27	22	5	332.9	354.3	263.2
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	10	6	4	8	4	4	98.6	64.4	210.5
Arteriosclerosis (440)	0	0	0	1	1	0	12.3	16.1	-
Influenza and pneumonia (470-486)	0	0	0	1	0	1	12.3	-	52.6
Bronchitis, emphysema, asthma (490-493)	2	1	1	2	1	1	24.7	16.1	52.6
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	1	0	12.3	16.1	-
Cirrhosis of liver (571)	0	0	0	1	1	0	12.3	16.1	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	1	0	1	12.3	-	52.6
Certain causes of mortality in early infancy (760-778)	0	0	0	2	2	0	13.0	19.6	-
Accidents (800-949)	9	6	3	9	5	4	111.0	80.5	210.5
Motor vehicle accidents (810-823)	6	4	2	4	1	3	49.3	16.1	157.9
Suicide (950-959)	1	0	1	1	0	1	12.3	-	52.6
Homicide (960-978)	1	1	0	1	1	0	12.3	16.1	-
Ill-defined and unknown (780-796)	4	1	3	4	1	3	49.3	16.1	157.9

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
SUNFLOWER COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	837	136	701	994	192	802	23.8	14.5	28.1
Fetal deaths	24	2	22	43	9	34	43.2	46.9	42.4
Total deaths	375	111	264	455	157	298	10.9	11.8	10.4
Deaths under 1 year	32	4	28	38	6	32	38.2	31.2	39.9
Deaths under 28 days	24	4	20	27	5	22	27.2	26.0	27.4
Bacillary dysentery and amebiasis (004, 006)	1	0	1	1	0	1	2.4	-	3.5
Enteritis and other diarrheal diseases (008, 009)	1	0	1	1	0	1	2.4	-	3.5
Tuberculosis (010-019)	2	0	2	3	0	3	7.2	-	10.5
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	1	0	1	1	0	1	2.4	-	3.5
Malignant neoplasms (140-209)	44	12	32	57	22	35	136.5	165.9	122.8
Benign and unspecified neoplasms (210-239)	3	1	2	3	1	2	7.2	7.5	7.0
Diabetes mellitus (250)	7	4	3	9	6	3	21.6	45.2	10.5
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	1	0	1	2.4	-	3.5
Diseases of heart (390-398, 402, 404-429)	131	51	80	155	69	86	371.2	520.4	301.8
Hypertension (400, 401, 403)	0	0	0	2	0	2	4.8	-	7.0
Cerebrovascular disease (430-438)	56	12	44	62	16	46	148.5	120.7	161.4
Arteriosclerosis (440)	2	2	0	4	3	1	9.6	22.6	3.5
Influenza and pneumonia (470-486)	7	1	6	10	4	6	23.9	30.2	21.0
Bronchitis, emphysema, asthma (490-493)	3	1	2	4	2	2	9.6	15.1	7.0
Peptic ulcer (531-533)	2	1	1	2	1	1	4.8	7.5	3.5
Hernia and intestinal obstruction (550-553, 560)	1	0	1	3	0	3	7.2	-	10.5
Cirrhosis of liver (571)	4	3	1	3	2	1	7.2	15.1	3.5
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	2	0	2	4.8	-	7.0
Nephritis and nephrosis (580-584)	4	1	3	6	1	5	14.4	7.5	17.5
Infections of kidney (590)	5	3	2	6	3	3	14.4	22.6	10.5
Hyperplasia of prostate (600)	1	0	1	3	0	3	7.2	-	10.5
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	3	1	2	7	3	4	16.8	22.6	14.0
Certain causes of mortality in early infancy (760-778)	17	3	14	18	3	15	18.1	15.6	18.7
Accidents (800-949)	29	7	22	30	10	20	71.8	75.4	70.2
Motor vehicle accidents (810-823)	13	6	7	13	6	7	31.1	45.2	24.6
Suicide (950-959)	0	0	0	1	1	0	2.4	7.5	-
Homicide (960-978)	7	1	6	9	1	8	21.6	7.5	28.1
Ill-defined and unknown (780-796)	32	3	29	36	4	32	86.2	30.2	112.3

See page 59 for footnotes.

LIVE BIRTHS, FETAL DSATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
TALLAHATCHIE COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
Live Births, Fetal Deaths, Deaths	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	441	74	367	457	75	382	21.3	9.9	27.5
Fetal deaths	15	2	13	20	5	15	43.8	66.7	39.3
Total deaths	209	69	140	240	88	152	11.2	11.6	10.9
Deaths under 1 year	19	1	18	22	3	19	48.1	40.0	49.7
Deaths under 28 days	9	1	8	13	3	10	28.4	40.0	26.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	3	0	3	3	0	3	14.0	-	21.6
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	20	7	13	26	9	17	121.2	118.9	122.4
Benign and unspecified neoplasms (210-239)	1	0	1	2	1	1	9.3	13.2	7.2
Diabetes mellitus (250)	2	1	1	2	1	1	9.3	13.2	7.2
Avitaminoses and other nutritional deficiencies (260-269)	1	0	1	1	0	1	4.6	-	7.2
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	92	38	54	95	43	52	442.7	568.0	374.4
Hypertension (400, 401, 403)	2	0	2	2	0	2	9.3	-	14.4
Cerebrovascular disease (430-438)	16	6	10	22	10	12	102.5	132.1	86.4
Arteriosclerosis (440)	2	1	1	4	2	2	18.6	26.4	14.4
Influenza and pneumonia (470-486)	9	1	8	10	1	9	46.6	13.2	64.8
Bronchitis, emphysema, asthma (490-493)	3	1	2	5	3	2	23.3	39.6	14.4
Peptic ulcer (531-533)	1	1	0	1	1	0	4.6	13.2	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	1	1	0	4.6	13.2	-
Nephritis and nephrosis (580-584)	0	0	0	1	0	1	4.6	-	7.2
Infections of kidney (590)	3	0	3	2	0	2	9.3	-	14.4
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	7	1	6	10	3	7	21.9	40.0	18.3
Accidents (800-949)	13	2	11	13	3	10	60.6	39.6	72.0
Motor vehicle accidents (810-823)	7	2	5	6	2	4	28.0	26.4	28.8
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	5	3	2	6	3	3	28.0	39.6	21.6
Ill-defined and unknown (780-796)	20	4	16	23	4	19	107.2	52.8	136.8

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
TATE COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	146	1	145	367	119	248	17.8	12.3	22.7
Fetal deaths	5	0	5	12	2	10	32.7	16.8	40.3
Total deaths	112	50	62	189	90	99	9.2	9.3	9.0
Deaths under 1 year	7	0	7	15	0	15	40.9	-	60.5
Deaths under 28 days	5	0	5	11	0	11	30.0	-	44.4
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	2	0	2	9.7	-	18.3
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	15	7	8	21	12	9	101.9	124.1	82.3
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	1	1	0	2	2	0	9.7	20.7	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	53	25	28	78	42	36	378.4	434.3	329.1
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	18	6	12	36	16	20	174.7	165.5	182.8
Arteriosclerosis (440)	0	0	0	2	2	0	9.7	20.7	-
Influenza and pneumonia (470-486)	3	2	1	7	3	4	34.0	31.0	36.6
Bronchitis, emphysema, asthma (490-493)	1	1	0	1	1	0	4.8	10.3	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	1	0	1	4.8	-	9.1
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	0	1	4.8	-	9.1
Nephritis and nephrosis (580-584)	1	1	0	3	3	0	14.6	31.0	-
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	0	1	1	0	1	4.8	-	9.1
Certain causes of mortality in early infancy (760-778)	3	0	3	7	0	7	19.1	-	28.2
Accidents (800-949)	9	4	5	14	5	9	67.9	51.7	82.3
Motor vehicle accidents (810-823)	3	1	2	6	3	3	29.1	31.0	27.4
Suicide (950-959)	1	1	0	1	1	0	4.8	10.3	-
Homicide (960-978)	1	1	0	1	0	1	4.8	-	9.1
Ill-defined and unknown (780-796)	3	0	3	5	0	5	24.3	-	45.7

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
TIPPAH COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
Live Births, Fetal Deaths, Deaths				Total	White	Non-white	Total	White	Non-white
Live births	342	268	74	309	236	73	17.5	16.4	22.4
Fetal deaths	27	18	9	19	11	8	61.5	46.6	109.6
Total deaths	170	138	32	181	149	32	10.2	10.4	9.8
Deaths under 1 year	8	7	1	10	8	2	32.4	33.9	27.4
Deaths under 28 days	6	5	1	7	6	1	22.6	25.4	13.7
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	31	27	4	38	34	4	215.3	236.3	122.7
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	6	6	0	2	2	0	11.3	13.9	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	62	53	9	58	49	9	328.6	340.5	276.1
Hypertension (400, 401, 403)	1	0	1	1	0	1	5.7	-	30.7
Cerebrovascular disease (430-438)	38	23	15	37	24	13	209.6	166.8	398.8
Arteriosclerosis (440)	1	1	0	0	0	0	-	-	-
Influenza and pneumonia (470-486)	3	3	0	4	4	0	22.7	27.8	-
Bronchitis, emphysema, asthma (490-493)	0	0	0	1	1	0	5.7	6.9	-
Peptic ulcer (531-533)	1	1	0	2	2	0	11.3	13.9	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	1	1	0	5.7	6.9	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	1	0	1	1	0	5.7	6.9	-
Infections of kidney (590)	0	0	0	2	2	0	11.3	13.9	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	3	2	1	4	3	1	22.7	20.8	30.7
Certain causes of mortality in early infancy (760-778)	6	5	1	6	5	1	19.4	21.2	13.7
Accidents (800-949)	9	9	0	13	12	1	73.6	83.4	30.7
Motor vehicle accidents (810-823)	5	5	0	8	8	0	45.3	55.6	-
Suicide (950-959)	4	4	0	4	4	0	22.7	27.8	-
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	0	0	0	1	0	1	5.7	-	30.7

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
TISHOMINGO COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live Births, Fetal Deaths, Deaths									
Live births	149	143	6	195	188	7	11.7	11.7	11.7
Fetal deaths	7	5	2	6	4	2	30.8	21.3	285.7
Total deaths	119	112	7	164	156	8	9.8	9.7	13.3
Deaths under 1 year	2	1	1	4	3	1	20.5	16.0	142.8
Deaths under 28 days	2	1	1	4	3	1	20.5	16.0	142.8
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	1	0	1	1	0	6.0	6.2	-
Tuberculosis (010-019)	1	1	0	1	1	0	6.0	6.2	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	21	21	0	28	28	0	167.8	174.0	-
Benign and unspecified neoplasms (210-239)	0	0	0	2	2	0	12.0	12.4	-
Diabetes mellitus (250)	5	4	1	4	3	1	24.0	18.6	166.7
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	29	27	2	46	44	2	275.6	273.5	333.3
Hypertension (400, 401, 403)	0	0	0	1	1	0	6.0	6.2	-
Cerebrovascular disease (430-438)	25	24	1	31	30	1	185.7	186.4	166.7
Arteriosclerosis (440)	0	0	0	2	2	0	12.0	12.4	-
Influenza and pneumonia (470-486)	5	5	0	5	5	0	30.0	31.1	-
Bronchitis, emphysema, asthma (490-493)	1	1	0	1	1	0	6.0	6.2	-
Peptic ulcer (531-533)	1	1	0	2	2	0	12.0	12.4	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	1	1	0	6.0	6.2	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	1	0	2	2	0	12.0	12.4	-
Infections of kidney (590)	4	3	1	4	3	1	24.0	18.6	166.7
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	1	1	0	6.0	6.2	-
Certain causes of mortality in early infancy (760-778)	1	1	0	3	3	0	15.4	16.0	-
Accidents (800-949)	13	12	1	14	12	2	83.9	74.6	333.3
Motor vehicle accidents (810-823)	9	8	1	8	6	2	47.9	37.3	333.3
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	8	7	1	6	5	1	35.9	31.1	166.7

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
TUNICA COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	324	31	293	344	54	290	22.8	12.6	26.8
Fetal deaths	15	1	14	16	2	14	46.5	37.0	48.3
Total deaths	124	20	104	144	22	122	9.5	5.1	11.3
Deaths under 1 year	12	1	11	12	0	12	34.9	-	41.4
Deaths under 28 days	4	0	4	5	0	5	14.5	-	17.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	1	0	1	6.6	-	9.2
Tuberculosis (010-019)	0	0	0	1	0	1	6.6	-	9.2
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	10	3	7	20	7	13	132.4	162.8	120.2
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	1	0	1	0	0	0	-	-	-
Avitaminoses and other nutritional deficiencies (260-269)	2	0	2	1	0	1	6.6	-	9.2
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	50	10	40	54	9	45	357.4	209.3	416.3
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	18	4	14	19	3	16	125.7	69.8	148.0
Arteriosclerosis (440)	0	0	0	1	1	0	6.6	23.2	-
Influenza and pneumonia (470-486)	3	1	2	3	0	3	19.8	-	27.8
Bronchitis, emphysema, asthma (490-493)	1	1	0	0	0	0	-	-	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	0	1	6.6	-	9.2
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	3	0	3	3	0	3	19.8	-	27.8
Infections of kidney (590)	1	0	1	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	1	0	1	2	0	2	5.8	-	6.9
Accidents (800-949)	9	1	8	9	1	8	59.6	23.2	74.0
Motor vehicle accidents (810-823)	6	1	5	5	1	4	33.1	23.2	37.0
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	1	0	1	2	0	2	13.2	-	18.5
Ill-defined and unknown (780-796)	22	0	22	21	0	21	139.0	-	194.3

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
UNION COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	281	197	84	324	241	83	16.3	14.4	26.3
Fetal deaths	9	5	4	11	7	4	34.0	29.0	48.2
Total deaths	193	155	38	212	170	42	10.6	10.1	13.3
Deaths under 1 year	4	1	3	6	2	4	18.5	8.3	48.2
Deaths under 28 days	2	1	1	4	2	2	12.3	8.3	24.1
Bacillary dysentery and amebiasis (004, 006)	1	1	0	1	1	0	5.0	6.0	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	28	23	5	29	21	8	145.6	125.3	253.2
Benign and unspecified neoplasms (210-239)	1	1	0	1	1	0	5.0	6.0	-
Diabetes mellitus (250)	2	1	1	3	2	1	15.1	11.9	31.6
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	77	69	8	83	74	9	416.7	441.5	284.8
Hypertension (400, 401, 403)	2	1	1	2	1	1	10.0	6.0	31.6
Cerebrovascular disease (430-438)	20	16	4	25	20	5	125.5	119.3	158.2
Arteriosclerosis (440)	2	2	0	3	3	0	15.1	17.9	-
Influenza and pneumonia (470-486)	10	6	4	11	7	4	55.2	41.8	126.6
Bronchitis, emphysema, asthma (490-493)	5	5	0	5	5	0	25.1	29.8	-
Peptic ulcer (531-533)	1	0	1	2	1	1	10.0	6.0	31.6
Hernia and intestinal obstruction (550-553, 560)	1	1	0	1	1	0	5.0	6.0	-
Cirrhosis of liver (571)	1	1	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	1	1	0	5.0	6.0	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	3	2	1	2	1	1	10.0	6.0	31.6
Hyperplasia of prostate (600)	1	0	1	1	1	0	5.0	6.0	-
Maternal causes (630-678)	1	0	1	1	0	1	30.9	-	120.5
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	2	1	1	4	2	2	12.3	8.3	24.1
Accidents (800-949)	7	7	0	7	7	0	35.1	41.8	-
Motor vehicle accidents (810-823)	3	3	0	3	3	0	15.1	17.9	-
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	1	0	1	1	0	1	5.0	-	31.6
Ill-defined and unknown (780-796)	16	12	4	17	13	4	85.3	77.6	126.6

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
WALTHALL COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	325	178	147	250	100	150	17.0	12.3	22.9
Fetal deaths	16	8	8	11	6	5	44.0	60.0	33.3
Total deaths	117	72	45	145	90	55	9.9	11.1	8.4
Deaths under 1 year	7	4	3	9	4	5	36.0	40.0	33.3
Deaths under 28 days	4	4	0	6	4	2	24.0	40.0	13.3
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	1	1	0	1	1	0	6.8	12.3	-
Meningococcal infection (036)	1	1	0	1	1	0	6.8	12.3	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	8	3	5	14	6	8	95.4	73.8	122.3
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	2	1	1	2	1	1	13.6	12.3	15.3
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	50	36	14	59	39	20	402.2	479.7	305.8
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	12	8	4	17	14	3	115.9	172.2	45.9
Arteriosclerosis (440)	3	3	0	2	2	0	13.6	24.6	-
Influenza and pneumonia (470-486)	6	3	3	6	3	3	40.9	36.9	45.9
Bronchitis, emphysema, asthma (490-493)	1	1	0	2	2	0	13.6	24.6	-
Peptic ulcer (531-533)	1	1	0	1	1	0	6.8	12.3	-
Hernia and intestinal obstruction (550-553, 560)	2	1	1	3	1	2	20.4	12.3	30.6
Cirrhosis of liver (571)	0	0	0	1	1	0	6.8	12.3	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	1	1	0	2	2	0	13.6	24.6	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	2	2	0	2	2	0	13.6	24.6	-
Certain causes of mortality in early infancy (760-778)	2	2	0	3	2	1	12.0	20.0	6.7
Accidents (800-949)	3	0	3	5	3	2	34.1	36.9	30.6
Motor vehicle accidents (810-823)	1	0	1	3	3	0	20.4	36.9	-
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	2	0	2	3	1	2	20.4	12.3	30.6
Ill-defined and unknown (780-796)	14	4	10	12	2	10	81.8	24.6	152.9

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
WARREN COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,305	534	771	944	472	472	20.6	18.0	24.1
Fetal deaths	142	65	77	94	51	43	99.6	108.0	91.1
Total deaths	710	305	405	560	234	326	12.2	8.9	16.6
Deaths under 1 year	56	9	47	38	7	31	40.2	14.8	65.7
Deaths under 28 days	45	9	36	32	6	26	33.9	12.7	55.1
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	0	0	0	-	-	-
Tuberculosis (010-019)	2	0	2	2	0	2	4.4	-	10.2
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	1	0	1	1	0	1	2.2	-	5.1
Malignant neoplasms (140-209)	102	54	48	79	39	40	172.5	148.8	204.2
Benign and unspecified neoplasms (210-239)	2	0	2	2	0	2	4.4	-	10.2
Diabetes mellitus (250)	14	5	9	10	5	5	21.8	19.1	25.5
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	1	0	1	2	0	2	4.4	-	10.2
Diseases of heart (390-398, 402, 404-429)	207	109	98	166	86	80	362.4	328.1	408.4
Hypertension (400, 401, 403)	8	4	4	4	2	2	8.7	7.6	10.2
Cerebrovascular disease (430-438)	78	31	47	68	24	44	148.5	91.6	224.6
Arteriosclerosis (440)	9	4	5	5	3	2	10.9	11.4	10.2
Influenza and pneumonia (470-486)	22	8	14	21	11	10	45.8	42.0	51.0
Bronchitis, emphysema, asthma (490-493)	13	9	4	10	6	4	21.8	22.9	20.4
Peptic ulcer (531-533)	2	2	0	2	2	0	4.4	7.6	-
Hernia and intestinal obstruction (550-553, 560)	3	2	1	0	0	0	-	-	-
Cirrhosis of liver (571)	2	2	0	2	2	0	4.4	7.6	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	2	0	2	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	10	3	7	7	2	5	15.3	7.6	25.5
Infections of kidney (590)	2	2	0	2	2	0	4.4	7.6	-
Hyperplasia of prostate (600)	1	0	1	1	0	1	2.2	-	5.1
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	6	2	4	3	3	0	6.6	11.4	-
Certain causes of mortality in early infancy (760-778)	42	7	35	29	4	25	30.7	8.5	53.0
Accidents (800-949)	47	23	24	34	17	17	74.2	64.9	86.8
Motor vehicle accidents (810-823)	24	13	11	13	8	5	28.4	30.5	25.5
Suicide (950-959)	4	3	1	4	3	1	8.7	11.4	5.1
Homicide (960-978)	4	0	4	1	0	1	2.2	-	5.1
Ill-defined and unknown (780-796)	76	11	65	73	10	63	159.4	38.2	321.6

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
WASHINGTON COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,850	665	1,185	1,726	571	1,155	22.5	16.4	27.7
Fetal deaths	117	58	59	116	50	66	67.2	87.6	57.1
Total deaths	860	303	557	897	294	603	11.7	8.4	14.5
Deaths under 1 year	72	20	52	76	18	58	44.0	31.5	50.2
Deaths under 28 days	48	18	30	51	16	35	29.5	28.0	30.3
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	6	0	6	6	0	6	7.8	-	14.4
Tuberculosis (010-019)	3	1	2	6	2	4	7.8	5.7	9.6
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	117	53	64	126	51	75	164.6	146.3	179.8
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	23	8	15	23	6	17	30.0	17.2	40.8
Avitaminoses and other nutritional deficiencies (260-269)	2	0	2	2	0	2	2.6	-	4.8
Anemias (280-285)	4	1	3	4	1	3	5.2	2.9	7.2
Diseases of heart (390-398, 402, 404-429)	257	99	158	252	90	162	329.2	258.2	388.5
Hypertension (400, 401, 403)	13	4	9	13	3	10	17.0	8.6	24.0
Cerebrovascular disease (430-438)	89	31	58	97	32	65	126.7	91.8	155.9
Arteriosclerosis (440)	6	5	1	4	3	1	5.2	8.6	2.4
Influenza and pneumonia (470-486)	31	8	23	35	7	28	45.7	20.1	67.1
Bronchitis, emphysema, asthma (490-493)	11	5	6	12	6	6	15.7	17.2	14.4
Peptic ulcer (531-533)	4	3	1	3	2	1	3.9	5.7	2.4
Hernia and intestinal obstruction (550-553, 560)	5	2	3	5	1	4	6.5	2.9	9.6
Cirrhosis of liver (571)	2	1	1	2	1	1	2.6	2.9	2.4
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	3	2	1	2	2	0	2.6	5.7	-
Nephritis and nephrosis (580-584)	3	1	2	4	1	3	5.2	2.9	7.2
Infections of kidney (590)	5	1	4	5	1	4	6.5	2.9	9.6
Hyperplasia of prostate (600)	2	1	1	2	0	2	2.6	-	4.8
Maternal causes (630-678)	3	0	3	3	0	3	17.4	-	26.0
Congenital anomalies (740-759)	5	3	2	7	5	2	9.1	14.3	4.8
Certain causes of mortality in early infancy (760-778)	29	13	16	31	11	20	18.0	19.3	17.3
Accidents (800-949)	46	17	29	54	20	34	70.5	57.4	81.5
Motor vehicle accidents (810-823)	22	10	12	26	12	14	34.0	34.4	33.6
Suicide (950-959)	9	6	3	13	10	3	17.0	28.7	7.2
Homicide (960-978)	13	1	12	13	2	11	17.0	5.7	26.4
Ill-defined and unknown (780-796)	113	20	93	110	19	91	143.7	54.5	218.2

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
WAYNE COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	325	222	103	311	158	153	18.1	14.6	23.9
Fetal deaths	22	14	8	21	12	9	67.5	75.9	58.8
Total deaths	169	101	68	183	107	76	10.6	9.9	11.9
Deaths under 1 year	14	4	10	16	3	13	51.4	19.0	85.0
Deaths under 28 days	10	4	6	13	3	10	41.8	19.0	65.4
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	1	0	1	5.8	-	15.6
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	15	9	6	20	12	8	116.1	110.9	125.0
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	2	2	0	1	1	0	5.8	9.2	-
Avitaminoses and other nutritional deficiencies (260-269)	1	1	0	1	1	0	5.8	9.2	-
Anemias (280-285)	1	0	1	1	0	1	5.8	-	15.6
Diseases of heart (390-398, 402, 404-429)	70	48	22	66	44	22	383.3	406.6	343.8
Hypertension (400, 401, 403)	6	4	2	6	4	2	34.8	37.0	31.2
Cerebrovascular disease (430-438)	23	15	8	25	17	8	145.2	157.1	125.0
Arteriosclerosis (440)	1	0	1	2	0	2	11.6	-	31.2
Influenza and pneumonia (470-486)	9	3	6	11	5	6	63.9	46.2	93.8
Bronchitis, emphysema, asthma (490-493)	2	1	1	2	1	1	11.6	9.2	15.6
Peptic ulcer (531-533)	1	1	0	1	1	0	5.8	9.2	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	0	1	5.8	-	15.6
Cirrhosis of liver (571)	1	1	0	1	1	0	5.8	9.2	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	2	1	1	3	1	2	17.4	9.2	31.2
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	1	0	1	1	0	1	32.2	-	65.4
Congenital anomalies (740-759)	3	1	2	3	2	1	17.4	18.5	15.6
Certain causes of mortality in early infancy (760-778)	5	2	3	9	2	7	28.9	12.6	45.8
Accidents (800-949)	11	5	6	13	7	6	75.5	64.7	93.8
Motor vehicle accidents (810-823)	6	2	4	5	1	4	29.0	9.2	62.5
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	4	1	3	3	1	2	17.4	9.2	31.2
Ill-defined and unknown (780-796)	3	1	2	2	0	2	11.6	-	31.2

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
WEBSTER COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	91	63	28	167	108	59	15.0	12.7	22.6
Fetal deaths	2	0	2	7	3	4	41.9	27.8	67.8
Total deaths	108	94	14	134	116	18	12.0	13.6	6.9
Deaths under 1 year	2	2	0	4	2	2	24.0	18.5	33.9
Deaths under 28 days	2	2	0	2	2	0	12.0	18.5	-
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	2	0	2	18.0	-	76.6
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	11	10	1	21	20	1	188.5	234.5	38.3
Benign and unspecified neoplasms (210-239)	1	0	1	1	0	1	9.0	-	38.3
Diabetes mellitus (250)	0	0	0	3	1	2	26.9	11.7	76.6
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	34	30	4	45	41	4	403.9	480.6	153.2
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	35	29	6	30	25	5	269.3	293.1	191.6
Arteriosclerosis (440)	0	0	0	0	0	0	-	-	-
Influenza and pneumonia (470-486)	3	3	0	5	5	0	44.9	58.6	-
Bronchitis, emphysema, asthma (490-493)	1	1	0	1	1	0	9.0	11.7	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	1	1	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	1	1	0	9.0	11.7	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	1	0	9.0	11.7	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	1	1	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	1	1	0	1	1	0	9.0	11.7	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	2	2	0	2	2	0	12.0	18.5	-
Accidents (800-949)	9	9	0	6	6	0	53.8	70.3	-
Motor vehicle accidents (810-823)	2	2	0	1	1	0	9.0	11.7	-
Suicide (950-959)	1	1	0	2	2	0	18.0	23.4	-
Homicide (960-978)	0	0	0	1	1	0	9.0	11.7	-
Ill-defined and unknown (780-796)	3	2	1	5	4	1	44.9	46.9	38.3

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
WILKINSON COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	221	96	125	220	50	170	19.8	15.0	21.9
Fetal deaths	6	1	5	10	2	8	45.4	40.0	47.0
Total deaths	111	56	55	119	52	67	10.7	15.6	8.6
Deaths under 1 year	6	2	4	8	1	7	36.4	20.0	41.2
Deaths under 28 days	5	1	4	7	1	6	31.8	20.0	35.3
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	1	0	0	0	0	-	-	-
Tuberculosis (010-019)	1	0	1	2	0	2	18.0	-	25.8
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	18	10	8	23	14	9	207.2	419.2	116.0
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	0	0	0	0	0	0	-	-	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	1	0	1	1	0	1	9.0	-	12.9
Diseases of heart (390-398, 402, 404-429)	31	16	15	38	19	19	342.3	568.9	244.8
Hypertension (400, 401, 403)	6	2	4	6	2	4	54.0	59.9	51.5
Cerebrovascular disease (430-438)	10	1	9	6	0	6	54.0	-	77.3
Arteriosclerosis (440)	2	2	0	1	1	0	9.0	29.9	-
Influenza and pneumonia (470-486)	5	4	1	4	0	4	36.0	-	51.5
Bronchitis, emphysema, asthma (490-493)	3	3	0	1	1	0	9.0	29.9	-
Peptic ulcer (531-533)	1	1	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	2	1	1	0	0	0	-	-	-
Cirrhosis of liver (571)	1	1	0	1	1	0	9.0	29.9	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	1	0	1	9.0	-	12.9
Certain causes of mortality in early infancy (760-778)	3	0	3	5	1	4	22.7	20.0	23.5
Accidents (800-949)	10	7	3	14	8	6	126.1	239.5	77.3
Motor vehicle accidents (810-823)	4	4	0	9	6	3	81.1	179.6	38.6
Suicide (950-959)	1	0	1	1	0	1	9.0	-	12.9
Homicide (960-978)	1	0	1	1	0	1	9.0	-	12.9
Ill-defined and unknown (780-796)	4	0	4	5	1	4	45.0	29.9	51.5

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
WINSTON COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	291	136	155	334	145	189	17.9	15.0	21.0
Fetal deaths	13	3	10	15	4	11	44.9	27.6	58.2
Total deaths	176	110	66	226	138	88	12.1	14.3	9.8
Deaths under 1 year	6	1	5	9	2	7	26.9	13.8	37.0
Deaths under 28 days	4	1	3	4	1	3	12.0	6.9	15.9
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	1	0	2	1	1	10.7	10.4	11.1
Tuberculosis (010-019)	1	0	1	1	0	1	5.4	-	11.1
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	29	19	10	34	20	14	182.1	207.2	155.2
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	1	1	0	1	1	0	5.4	10.4	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	1	0	1	5.4	-	11.1
Diseases of heart (390-398, 402, 404-429)	64	44	20	85	58	27	455.3	601.0	299.3
Hypertension (400, 401, 403)	3	2	1	3	2	1	16.1	20.7	11.1
Cerebrovascular disease (430-438)	26	18	8	24	17	7	128.5	177.2	77.6
Arteriosclerosis (440)	2	1	1	5	2	3	26.8	20.7	33.2
Influenza and pneumonia (470-486)	4	2	2	8	5	3	42.8	51.8	33.2
Bronchitis, emphysema, asthma (490-493)	3	1	2	6	4	2	32.1	41.4	22.2
Peptic ulcer (531-533)	0	0	0	1	1	0	5.4	10.4	-
Hernia and intestinal obstruction (550-553, 560)	1	1	0	2	1	1	10.7	10.4	11.1
Cirrhosis of liver (571)	1	1	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	1	0	1	1	0	5.4	10.4	-
Infections of kidney (590)	2	2	0	3	2	1	16.1	20.7	11.1
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	3	1	2	3	1	2	9.0	6.9	10.6
Accidents (800-949)	7	5	2	10	8	2	53.6	82.9	22.2
Motor vehicle accidents (810-823)	1	1	0	4	4	0	21.4	41.4	-
Suicide (950-959)	2	1	1	2	1	1	10.7	10.4	11.1
Homicide (960-978)	2	0	2	1	0	1	5.4	-	11.1
Ill-defined and unknown (780-796)	14	1	13	19	4	15	101.8	41.4	166.3

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
YALOBUSHA COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	168	64	104	210	89	121	16.8	12.4	22.8
Fetal deaths	5	1	4	5	1	4	23.8	11.2	33.0
Total deaths	120	67	53	160	97	63	12.8	13.5	11.9
Deaths under 1 year	3	1	2	5	1	4	23.8	11.2	33.0
Deaths under 28 days	1	1	0	3	1	2	14.3	11.2	16.5
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	11	8	3	22	15	7	176.4	209.2	132.1
Benign and unspecified neoplasms (210-239)	1	1	0	1	1	0	8.0	13.9	-
Diabetes mellitus (250)	1	0	1	2	1	1	16.0	13.9	18.9
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	1	1	0	8.0	13.9	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	41	28	13	49	34	15	392.9	474.2	283.0
Hypertension (400, 401, 403)	1	0	1	1	0	1	8.0	-	18.9
Cerebrovascular disease (430-438)	33	12	21	32	13	19	256.6	181.3	358.5
Arteriosclerosis (440)	0	0	0	1	1	0	8.0	13.9	-
Influenza and pneumonia (470-486)	3	0	3	6	3	3	48.1	41.8	56.6
Bronchitis, emphysema, asthma (490-493)	4	3	1	6	5	1	48.1	69.7	18.9
Peptic ulcer (531-533)	1	1	0	1	1	0	8.0	13.9	-
Hernia and intestinal obstruction (550-553, 560)	1	1	0	1	1	0	8.0	13.9	-
Cirrhosis of liver (571)	1	1	0	1	1	0	8.0	13.9	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	0	1	1	0	1	8.0	-	18.9
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	2	1	1	16.0	13.9	18.9
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	1	1	0	3	1	2	14.3	11.2	16.5
Accidents (800-949)	10	6	4	14	9	5	112.3	125.5	94.3
Motor vehicle accidents (810-823)	6	3	3	8	5	3	64.2	69.7	56.6
Suicide (950-959)	1	1	0	2	2	0	16.0	27.9	-
Homicide (960-978)	0	0	0	1	1	0	8.0	13.9	-
Ill-defined and unknown (780-796)	3	1	2	4	1	3	32.1	13.9	56.6

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
YAZOO COUNTY, MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live Births, Fetal Deaths, Deaths									
Live births	526	203	323	596	215	381	19.0	16.0	21.3
Fetal deaths	23	5	18	36	9	27	60.4	41.9	70.9
Total deaths	302	106	196	388	148	240	12.4	11.0	13.4
Deaths under 1 year	22	9	13	31	11	20	52.0	51.2	52.5
Deaths under 28 days	15	7	8	21	8	13	35.2	37.2	34.1
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	1	0	1	3.2	-	5.6
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	33	16	17	52	26	26	166.2	193.6	145.6
Benign and unspecified neoplasms (210-239)	1	1	0	1	1	0	3.2	7.4	-
Diabetes mellitus (250)	5	0	5	6	2	4	19.2	14.9	22.4
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	69	30	39	85	39	46	271.7	290.4	257.7
Hypertension (400, 401, 403)	1	0	1	4	0	4	12.8	-	22.4
Cerebrovascular disease (430-438)	38	13	25	52	20	32	166.2	148.9	179.3
Arteriosclerosis (440)	16	9	7	16	7	9	51.2	52.1	50.4
Influenza and pneumonia (470-486)	10	4	6	14	5	9	44.8	37.2	50.4
Bronchitis, emphysema, asthma (490-493)	2	1	1	4	2	2	12.8	14.9	11.2
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	1	1	0	1	1	0	3.2	7.4	-
Cirrhosis of liver (571)	1	1	0	1	1	0	3.2	7.4	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	6	2	4	7	3	4	22.4	22.3	22.4
Infections of kidney (590)	0	0	0	2	1	1	6.4	7.4	5.6
Hyperplasia of prostate (600)	1	0	1	1	0	1	3.2	-	5.6
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	1	1	0	3.2	7.4	-
Certain causes of mortality in early infancy (760-778)	8	7	1	11	7	4	18.4	32.6	10.5
Accidents (800-949)	30	10	20	38	13	25	121.5	96.8	140.0
Motor vehicle accidents (810-823)	17	6	11	17	5	12	54.3	37.2	67.2
Suicide (950-959)	2	2	0	1	1	0	3.2	7.4	-
Homicide (960-978)	3	0	3	4	0	4	12.8	-	22.4
Ill-defined and unknown (780-796)	66	6	60	64	5	59	204.6	37.2	330.5

See page 59 for footnotes.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
STATE OF MISSISSIPPI, 1968

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	45706	22896	22810	45719	22807	22912	19.4	15.7	25.4
Fetal deaths	2373	1173	1200	2322	1119	1203	50.8	49.1	52.5
Total deaths	23647	13261	10386	23986	13468	10518	10.2	9.3	11.7
Deaths under 1 year	1588	504	1084	1613	517	1096	35.3	22.7	47.8
Deaths under 28 days	1096	411	685	1116	424	692	24.4	18.6	30.2
Bacillary dysentery and amebiasis (004, 006)	14	5	9	14	5	9	0.6	0.3	1.0
Enteritis and other diarrheal diseases (008, 009)	111	30	81	113	29	84	4.8	2.0	9.3
Tuberculosis (010-019)	92	40	52	89	37	52	3.8	2.6	5.8
Meningococcal infection (036)	5	5	0	5	5	0	0.2	0.3	-
Syphilis (090-097)	14	3	11	12	2	10	0.5	0.1	1.1
Malignant neoplasms (140-209)	3158	2022	1136	3267	2101	1166	139.0	144.9	129.5
Benign and unspecified neoplasms (210-239)	42	22	20	49	28	21	2.1	1.9	2.3
Diabetes mellitus (250)	430	219	211	436	221	215	18.5	15.2	23.9
Avitaminoses and other nutritional deficiencies (260-269)	39	15	24	39	15	24	1.6	1.0	2.7
Anemias (280-285)	53	12	41	56	15	41	2.4	1.0	4.6
Diseases of heart (390-398, 402, 404-429)	7917	5002	2915	7969	5024	2945	339.0	346.4	327.1
Hypertension (400, 401, 403)	209	79	130	209	78	131	8.9	5.4	14.6
Cerebrovascular disease (430-438)	3041	1595	1446	3115	1631	1484	132.5	112.4	164.8
Arteriosclerosis (440)	309	208	101	316	211	105	13.4	14.5	11.7
Influenza and pneumonia (470-486)	830	425	405	825	414	411	35.1	28.5	45.6
Bronchitis, emphysema, asthma (490-493)	304	242	62	303	242	61	12.9	16.7	6.8
Peptic ulcer (531-533)	80	57	23	82	59	23	3.5	4.1	2.6
Hernia and intestinal obstruction (550-553, 560)	91	47	44	88	43	45	3.7	3.0	5.0
Cirrhosis of liver (571)	141	107	34	146	113	33	6.2	7.8	3.7
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	43	30	13	42	29	13	1.8	2.0	1.4
Nephritis and nephrosis (580-584)	181	55	126	181	58	123	7.7	4.0	13.7
Infections of kidney (590)	194	127	67	199	133	66	8.5	9.2	7.3
Hyperplasia of prostate (600)	48	20	28	49	21	28	2.1	1.4	3.1
Maternal causes (630-678)	22	2	20	21	3	18	4.6	1.3	7.8
Congenital anomalies (740-759)	188	117	71	205	135	70	8.7	9.3	7.8
Certain causes of mortality in early infancy (760-778)	773	308	465	791	320	471	17.3	14.0	20.6
Accidents (800-949)	1704	977	727	1723	999	724	73.3	68.9	80.4
Motor vehicle accidents (810-823)	824	542	282	821	542	279	34.9	37.4	31.0
Suicide (950-959)	171	148	23	171	148	23	7.3	10.2	2.6
Homicide (960-978)	240	54	186	256	60	196	10.9	4.1	21.8
Ill-defined and unknown (780-796)	1872	497	1375	1858	487	1371	79.0	33.6	152.3

See page 59 for footnotes.