


VITAL STATISTICS
MISSISSIPPI

1969



Mississippi State Board of Health
Jackson, Mississippi

VITAL STATISTICS

MISSISSIPPI

1969

STATISTICAL SERVICES UNIT
MISSISSIPPI STATE BOARD OF HEALTH
P. O. BOX 1700
JACKSON, MISSISSIPPI 39205

TABLE OF CONTENTS

	Page
SYNOPSIS OF VITAL STATISTICS, MISSISSIPPI, 1969	
Introduction	I
Description of Bulletin	I
Definitions of Live Birth and Fetal Death	I
Occurrence and Residence Allocation	I
Urban-Rural Classification	II
Population	III
Rates	III
Classification of Cause of Death	III
Live Births	IV
Trend	IV
Race	IV
County	V
Natural Increase	V
Attendant	V
Month	VI
Fertility	VII
Age of Mother	VII
Education of Parents	VIII
Legitimacy	IX
Total-Birth Order	X
Last Prior Delivery	XI
Prenatal Care	XII
Complications during Pregnancy and Labor	XIV
Length of Gestation and Birth Weight	XVI
Sex	XVII
Plurality	XVII
Birth Injuries and Congenital Malformations	XVII
Fetal Deaths	XIX
Reporting and Reliability	XIX
Trend and Race	XIX
County	XIX
Attendant	XIX
Age of Mother	XX
Education of Parents	XX
Legitimacy	XX
Total-Birth Order	XXI
Last Prior Delivery	XXII
Prenatal Care	XXII
Length of Gestation and Birth Weight	XXIII
Sex	XXIV
Plurality	XXV
Cause of Fetal Death, Complications during Pregnancy and Labor, Birth Injuries, and Congenital Malformations	XXV
Deaths	XXV
Trend and Race	XXV
County	XXV
Month	XXVI
Age at Death	XXVI
Leading Causes of Death	XXVII
Leading Causes by Age	XXVIII
Ill-defined and Unknown Causes	XXVIII

Deaths, continued	
Specific Causes	XXX
Heart Disease	XXX
Malignant Neoplasms	XXX
Cerebrovascular Disease	XXXI
Chronic Respiratory Diseases	XXXI
Influenza and Pneumonia	XXXI
Accidents	XXXI
Suicides and Homicides	XXXII
Injury Undetermined Whether Accidentally or Purposely Inflicted	XXXIII
Maternal Deaths	XXXIII
Infant Mortality	XXXIV
Marriages	XXXVI
Trend and Race	XXXVI
Month	XXXVII
Officiant	XXXVII
Age of Bride and Groom	XXXVII
Marriage Order and Previous Marital Status	XXXIX
Nonresident Brides and Grooms	XXXIX
Out-of-State Use of Mississippi Licenses	XL
Divorces	XL
Trend and Race	XL
Cause and Complainant	XL
Minor Children	XL
Duration of Marriage	XLI
Additional Information Needed	XLI

POPULATION

Table 1. By County, by Race, Mississippi, Census of April 1, 1970	1
-------------------------------------------------------------------------	---

LIVE BIRTHS

Table 2. By Place of Occurrence, and by Place of Residence of Mother, by Race, with Rates, Mississippi, 1969	3
Table 3. By Place of Residence of Mother, by Race and Attendant, and by Sex, Mississippi, 1969	6

FETAL DEATHS

Table 4. By Place of Occurrence; by Place of Residence of Mother, by Race and Attendant, and by Sex, Mississippi, 1969	10
------------------------------------------------------------------------------------------------------------------------------	----

DEATHS

Table 5. By Cause of Death, by Race, Residence Data, Mississippi, 1969	14
Table 6. By Place of Occurrence, and by Place of Residence, by Race, with Rates, Mississippi, 1969	29
Table 7. By Cause Groups and Certain Selected Causes, by Race, Sex, and Age, with Total Rates, Residence Data, Mississippi, 1969	32

DEATHS UNDER 1 YEAR

Table 8. By Selected Causes, by Race and Age, with Total Rates, Residence Data, Mississippi, 1969 40

MARRIAGES AND DIVORCES

Table 9. Marriages and Divorces, Mississippi, 1969, Marriages by Place of Licensure, by Place of Occurrence, and by Place of Residence of Bride, by Race of Bride; Divorces by Place of Occurrence, by Race 44

Table 10. Divorces, Mississippi, 1969, by Cause of Divorce, by Race; by Number of Minor Children, by Race; by Number of Years Married, by Race 47

TIME SERIES

Table 11. Live Births, Mississippi, 1913-1969, by Year, by Race, with Rates, Occurrence Data 1913-1943, Residence Data 1944-1969 48

Table 12. Deaths, Mississippi, 1913-1969, by Year, by Race, with Rates, Occurrence Data 1913-1943, Residence Data 1944-1969 50

Table 13. Maternal Deaths, Mississippi, 1913-1969, by Year, by Race, with Rates, Occurrence Data 1913-1943, Residence Data 1944-1969 52

Table 14. Deaths Under 1 Year, Mississippi, 1917-1969, by Year, by Race, with Rates, Occurrence Data 1917-1943, Residence Data 1944-1969 .. 54

Table 15. Deaths Under 1 Month, Mississippi, 1924-1969, by Year, by Race, with Rates, Occurrence Data 1924-1944, Residence Data 1945-1969 .. 56

Table 16. Deaths from Heart Diseases, Malignant Neoplasms, Cerebrovascular Disease, Accidents, and Influenza and Pneumonia, Mississippi, 1960-1969, by Year, by Race, with Rates, Residence Data 58

Table 17. Marriages and Divorces, Mississippi, 1926-1969, by Year, by Race, Occurrence Data 59

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS, AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH, by Place of Occurrence, and by Place of Residence, by Race, with Rates, 1969

The Counties (in alphabetical order) 63

State of Mississippi 145

SYNOPSIS OF VITAL STATISTICS
MISSISSIPPI, 1969

INTRODUCTION

Description of Bulletin

The 1969 Vital Statistics Bulletin is the twenty-seventh annual report prepared by Statistical Services from information contained on records of live births, fetal deaths, deaths, marriages, and divorces collected by the office of Vital Records Registration. This synopsis provides certain explanations and definitions necessary for understanding the data presented and also summarizes the most noteworthy facts regarding the vital events of the year. Following the synopsis are ten tables (Numbers 1-10) containing statistics for 1969 and seven tables (Numbers 11-17) showing comparable data over a span of years, in most cases the entire period for which the information is available. The last section of the book contains a one-page summary table for each county and also for the state.

Definitions of Live Birth and Fetal Death

The definitions adopted by the Mississippi State Board of Health for use in Mississippi are those formulated by the World Health Organization and recommended by the National Center for Health Statistics. They are as follows:

Live birth - the complete expulsion or extraction from its mother of a product of human conception, irrespective of the duration of pregnancy, which, after such expulsion or extraction, breathes or shows any other evidence of life such as beating of the heart, pulsation of the umbilical cord, or definite movement of voluntary muscles, whether or not the umbilical cord has been cut or the placenta is attached.

Fetal death - death prior to the complete expulsion or extraction from its mother of a product of human conception, irrespective of the duration of pregnancy; the death is indicated by the fact that after such expulsion or extraction the fetus does not breathe or show any other evidence of life such as beating of the heart, pulsation of the umbilical cord, or definite movement of voluntary muscles.

Occurrence and Residence Allocation

Live births, fetal deaths, and deaths were classified by place of occurrence and by place of residence. Some of the tables contain both occurrence and residence figures, but most of the tables are based on residence data only, as these data are considered more suitable for studying public health problems. All figures given in this synopsis are residence counts unless otherwise stated.

Occurrence data were obtained from certificates filed in Mississippi at the place of the event. Residence data include transcripts of records of events that occurred in other states to Mississippi residents and exclude records filed in this state for non-residents. Residence assignment for live births and fetal deaths was determined by the usual residence of the mother. For deaths it was determined by the usual residence of the decedent, except that deaths of inmates of institutions were reallocated to place of previous residence regardless of length of stay in the institution.

Vital Statistics
Mississippi - 1969

A total of 2,995 transcripts of events that occurred in other places to Mississippi residents were received, and Mississippi filed 2,610 records for events that occurred in this state to nonresidents. Table A lists the states with whom the most records were exchanged in order of the total number of records sent and received. With only one exception, the states listed had more certificates filed in Mississippi for their residents than they filed for Mississippi residents. The reverse was true for Tennessee, the state with whom by far the largest number of records was exchanged. This situation resulted from the fact that many persons who live in the northern part of the state seek health care in Memphis, which is a major medical center and located just across the state line in Tennessee. Mississippi received transcripts from a total of 36 other states and the District of Columbia and filed certificates for residents of 38 other states and also four for foreigners. However, the neighboring states of Alabama, Arkansas, Louisiana, and Tennessee accounted for 89 percent of the transcripts and 83 percent of the Mississippi certificates filed for nonresidents.

Table A. Transcripts of certificates filed in other places for Mississippi residents, by place of occurrence, and certificates filed in Mississippi for nonresidents, by place of residence, by type of certificate

Place	Total		Live Births		Fetal Deaths		Deaths	
	Transcripts	Certificates, Nonresidents						
Total	2,995	2,610	1,680	1,698	42	85	1,273	827
Tennessee	1,800	245	1,085	151	31	6	684	88
Louisiana	439	905	219	661	0	39	220	205
Alabama	376	889	223	598	6	19	147	272
Arkansas	63	126	16	76	1	6	46	44
Florida	51	80	23	35	0	0	28	45
Texas	43	56	11	20	1	2	31	34
Illinois	29	67	12	34	1	2	16	31
Other places	194	242	91	123	2	11	101	108

Marriages were classified according to place of issuance of marriage license, place of occurrence of marriage, and place of residence of bride. However, the figures on brides who were Mississippi residents are incomplete since there is no exchange among the states of transcripts of marriage records for nonresidents and the out-of-state marriages of Mississippi brides could not be included. Divorces were classified only according to place where the decree was granted, as there is no information regarding place of residence on the record.

Urban-Rural Classification

For live births, fetal deaths, and deaths, the specific town or city of occurrence and of residence of the parents or of the deceased was tabulated if it was incorporated and if a population of 2,500 or more had been recorded in the census of 1960 (this was the only population information available when the certificates for 1969 were processed).

Vital Statistics
Mississippi - 1969

All other areas were classified as rural. However, since data from the census of 1970 were released before compilation of vital statistics for 1969 was completed, it was possible to include in Tables 2 and 6 all 24 of the cities that had reached a population of 10,000 or more by 1970. (Only 18 cities were in this category in 1960.) The urban-rural classification was not used for marriages and divorces as information on the marriage records is inadequate for proper allocation and divorce records do not contain an item relating to specific place.

Population

Figures from the 1970 census were used in this bulletin even though the vital statistics relate to the year 1969. Both the final counts from the census and the estimates for 1969 from the series prepared by Mississippi State University were available at the time the material for this report was compiled; but, when the estimates were compared with census figures for 1960 and 1970, it was determined that rates based on the census figures of 1970 would be more reliable than calculations based on the estimated population. There was considerable over-estimation of the population all during the 1960's.

Rates

Rates and ratios were computed from residence data on live births, fetal deaths, and deaths. The base used for rates for live births, fertility, total deaths, and deaths from specific causes (except maternal and certain causes of mortality in early infancy) was population; but the over-estimation of the population on which rates for most of the 1960's were based should be taken into account in comparing the rates for 1969 with those for prior years. Figures on live births were used as the base for calculating fetal death ratios and rates for maternal deaths, infant and neonatal deaths, and deaths from certain causes of mortality in early infancy because they approximate the groups at risk more closely than do general population figures.

The exact base for each type of rate used in this bulletin is given on page 61.

Classification of Cause of Death

Development of a system for classifying diseases began in Europe in the 1700's. By the end of the nineteenth century a standard classification of causes of death was being used in several countries and had been adopted for use in the United States. This classification has been revised approximately every ten years since 1900 in order to keep pace with new knowledge in the field of medicine. The latest version, Eighth Revision International Classification of Diseases, Adapted for Use in the United States, went into effect in Mississippi and a majority of the other states in 1968.

A list of the international classification numbers and group titles for this revision is presented here to show the organization of the system.

Vital Statistics
Mississippi - 1969

000-136	Infective and parasitic diseases
140-239	Neoplasms
240-279	Endocrine, nutritional, and metabolic diseases
280-289	Diseases of the blood and blood-forming organs
290-315	Mental disorders
320-389	Diseases of the nervous system and sense organs
390-458	Diseases of the circulatory system
460-519	Diseases of the respiratory system
520-577	Diseases of the digestive system
580-629	Diseases of the genitourinary system
630-678	Complications of pregnancy, childbirth, and the puerperium
680-709	Diseases of the skin and subcutaneous tissue
710-738	Diseases of the musculoskeletal system and connective tissue
740-759	Congenital anomalies
760-779	Certain causes of mortality in early infancy
780-796	Symptoms and ill-defined conditions
800-999	Accidents, poisonings, and violence

On certificates with more than one cause entered, only that indicated as the underlying cause was classified, but the system can be used for studying all conditions listed on death certificates.

Although standard classification enables comparison of statistics from one country or state with those of another country or state regardless of language or terminology differences, the periodic revisions which have taken place in the past have resulted in an unavoidable lack of comparability over the years for some causes of death. The greatest change was in the Sixth Revision in 1949, and only minor changes were incorporated in the Seventh Revision which went into use in 1958. Differences between the Seventh and Eighth Revisions have not been fully analyzed yet; but it appears that, although they are of more consequence than those between the Sixth and Seventh Revisions, there was a relatively small number of causes with major changes. Because of these revisions, comparisons of data over a period of years should be made with caution. However, the National Center for Health Statistics has prepared and published comparability ratios which show the effect revisions have had on each cause and which may be used to adjust figures for the purpose of analyzing trends.

LIVE BIRTHS (See Tables 2, 3, 11)

Trend

In 1969, for the first time in eight years, the number of live births to Mississippians increased. The total of 47,120 was 1,401 or 3.1 percent more than that reported in 1968 and also more than the count for 1967. Although final figures are not yet available for the United States, provisional data show that Mississippi's seven-year period of decline followed by a reversal of the trend in 1969 paralleled the experience of the country as a whole. The birth rate of 21.2 per 1,000 population for Mississippi in 1969 was higher than the rate for 1968, but part of the increase was due to the change from estimated to census population as a base (see section on population in the Introduction).

Race

The year 1969 was the first one since 1924 in which the number of births to

Vital Statistics
Mississippi - 1969

whites (24,025) exceeded those to nonwhites (23,095), and the difference amounted to 930. In the previous year the excess of nonwhites was only 105. The disparity between the race-specific rates was substantial - the rate of 28.0 for nonwhites was 63 percent higher than the 17.2 computed for whites.

County

A considerable amount of variation in birth rates by county was noted. For whites the range was from 9.1 to 22.4 in Jefferson and Harrison Counties respectively, and it was of note that even the highest rate for the white population was less than the overall rate for nonwhites in the state. The range for nonwhites was even wider - from 16.4 in Hancock County to 39.9 in Stone County. Variation among whites was found to be correlated with the age distribution of the female population but in a negative way; that is, birth rates were low in counties where the median age was high and vice-versa. For example, in Jefferson County, where the rate was lowest, the median age of the white female population was 41.3 years, whereas in Harrison County, which had the highest rate, the median was 25.6 years. This type of relationship was also found to some extent among nonwhites, but it was less consistent.

Natural Increase (Excess of Births over Deaths)

The natural increase registered for Mississippi residents in 1969 was 23,255 persons, and the excess of the birth rate over the death rate was 10.4 per 1,000 population. These figures were higher than comparable ones for 1968 because of a rise in the birth count and slight decline in the death toll, but the natural increase in persons was the second smallest since 1933. The rate for nonwhites was more than double that for whites, 15.9 in comparison with only 7.2. A breakdown by county and race showed that there were eleven counties in which there was no natural increase among the white population but, in fact, an actual loss due to the occurrence of more deaths than births. These counties were: Amite, Choctaw, Claiborne, Copiah, Franklin, Holmes, Jefferson, Kemper, Montgomery, Wilkinson, and Yalobusha. Jackson County again had a higher rate of natural increase among its white population (15.7) than any other county, and Smith County again had the highest rate recorded for the nonwhite population (29.8).

Attendant

A large majority of the children born to Mississippi residents in 1969 were delivered in a hospital or clinic providing in-patient care; as Table B1 shows, they amounted to 87 percent of the total. A small number - less than 1 percent - were delivered by physicians in places other than hospitals or clinics, but for the remaining 12 percent no medical attendant was present at delivery. The total count of deliveries without medical attention was 5,646 and most were to nonwhite mothers. Those to nonwhites made up 24 percent of the total for that race group in contrast to less than 1 percent for whites.

Vital Statistics
Mississippi - 1969

Table B1. Live births and percentage distribution, by attendant and race

Attendant	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>47,120</u>	<u>24,025</u>	<u>23,095</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
Hospital or clinic	41,044	23,911	17,133	87.1	99.5	74.2
Physician, other place	430	25	405	0.9	0.1	1.8
Midwife	5,571	64	5,507	11.8	0.3	23.8
Other	75	25	50	0.2	0.1	0.2

Continued improvement in securing hospitalization for delivery is recorded in Table B2, which shows that the percentage in hospitals or clinics almost doubled in the last twenty years. If the rate of gain registered for nonwhites continues to be as rapid as it has been in recent years, a level of almost 100 percent hospitalization could be reached in the next decade with delivery by a "granny" midwife a rare event. In 1969, however, the proportion of nonwhite children born in hospitals or clinics (74 percent) was far below the latest national figure available for nonwhites (94 percent in 1968).

Table B2. Percentage of live births in hospitals or clinics, by year and race, 1949, 1959, 1965-1969

Year	Total	White	Non-white
1949	44.3	80.1	16.6
1959	72.1	98.9	49.1
1965	77.3	99.4	58.8
1966	79.3	99.4	61.4
1967	82.1	99.5	65.5
1968	84.3	99.5	69.0
1969	87.1	99.5	74.2

Month

The distribution of births during the year was uneven. Although the last six months of the year had only 3 more days than the first six months, the number of births in the last half exceeded the count for the first half by nearly 3,500 and the daily average was 137 in comparison to only 120 for the first half. Table B3 shows that September was the month with the highest daily average and that, as usual, spring was the period of lowest daily incidence.

Vital Statistics
Mississippi - 1969

Table B3. Live births and average per day, by month

Month	Births	Average per Day	Month	Births	Average per Day
<u>Total</u>	<u>47,120</u>	<u>129</u>			
January	3,996	129	July	4,327	140
February	3,629	130	August	4,355	140
March	3,606	116	September	4,410	147
April	3,451	115	October	4,222	136
May	3,535	114	November	3,798	127
June	3,605	120	December	4,186	135

Fertility

For the first time since 1961, fertility rates were calculated because census data by age and sex were again available (see Table B4 in section on age of mother), and it was found that the rates were substantially lower than in 1961. The overall rate for 1969 was 105.1 live births per 1,000 females 15-44 years of age in comparison with 143.2 in 1961. Between the two years the rate for the white population dropped by 24 percent and that for nonwhites by 25 percent. Even though the decline was about the same for both groups, the most recent computations showed that the rate of 147.0 for nonwhites was 78 percent higher than the rate of 82.5 for whites. Latest available figures for the United States (white - 81.5 and nonwhite - 114.9 in 1968) also showed that the rate for nonwhites in Mississippi was excessive.

Age of Mother

Figures in Table B4 show that there were more mothers 20-24 years of age than in any other age group; 37 percent of the total were within this five-year span. There were also large numbers of mothers in the groups 15-19 and 25-29 years of age, and the three groups included in the age range 15-29 years comprised 81 percent of the total. Although there was a difference of only one year between the median ages of 23.6 and 22.6 respectively for white and nonwhite mothers, the percentage distributions by age for the two groups revealed greater variation. Sixty-six percent of the white mothers were concentrated in the period 20-29 years of age, but only 49 percent of the nonwhite mothers were in this range; in addition, excesses of nonwhites over whites occurred among both younger and older mothers.

According to the age-specific fertility rates in Table B4, fertility increased with age reaching a peak among mothers age 20-24 and then declined with age. Even though rates were higher for nonwhites than for whites in every age group, the differences were relatively greater at the lower and upper ends of the childbearing period.

Vital Statistics
Mississippi - 1969

Table B4. Live births, percentage distribution, and fertility rates,
by age of mother and race

Age of Mother	Number			Percent			Fertility Rate*		
	Total	White	Non- white	Total	White	Non- white	Total	White	Non- white
Total	47,120	24,025	23,095	100.0	100.0	100.0	105.1	82.5	147.0
10-14	326	43	283	0.7	0.2	1.2	2.6	0.6	5.0
15-19	11,144	4,375	6,769	23.6	18.2	29.3	96.9	67.9	133.8
20-24	17,230	10,024	7,206	36.6	41.7	31.2	195.0	172.8	237.5
25-29	9,986	5,967	4,019	21.2	24.8	17.4	150.4	129.9	196.5
30-34	4,949	2,404	2,545	10.5	10.0	11.0	82.6	58.5	135.5
35-39	2,545	951	1,594	5.4	4.0	6.9	44.2	23.9	89.4
40-44	835	233	602	1.8	1.0	2.6	13.6	5.5	31.4
45-49	73	20	53	0.2	0.1	0.2	1.2	0.5	2.9
Unknown	32	8	24	0.1	**	0.1			

* For total group - live births per 1,000 females 15-44 years of age; for specific age groups - live births per 1,000 females in specified group.

** Less than 0.06 percent.

Education of Parents

The year 1969 is the second in which information on the highest grade of school completed by the parents was collected. Comparison of the figures in Table B5 with those for the previous year showed that, for both parents, reporting was slightly more complete, educational levels were slightly higher, and the gap between the races was smaller. However, the picture presented by the new data was still an unfavorable one as 45 percent of the fathers and 52 percent of the mothers had not finished high school and 29 percent of each had not gone beyond the ninth grade. The median years of school completed by nonwhite parents was less than the median for whites; for fathers the difference was 2.5 years and for mothers it was 1.9 years. Also, 70 percent of the nonwhite parents had not graduated from high school, whereas, for white fathers and mothers, the percentages were only 30 and 35 respectively. Part of this disparity in educational levels was no doubt due to differences in age distribution since relatively more of the nonwhite than of the white mothers were below the normal age for finishing high school. Additional study which related education and legitimacy revealed that only 53 percent of the white and 12 percent of the nonwhite babies were legitimate and also born to parents both of whom had at least a high school education.

Vital Statistics
Mississippi - 1969

Table B5. Live births and percentage distribution,
by years of school completed by father and mother and race

Years of School Completed by Father and Mother	Number			Percent*			Cumulative Percent*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
FATHER									
Median years completed*	12.2	12.6	10.1						
Total legitimate births	38,379	23,285	15,094	100.0	100.0	100.0			
Less than 1 year	149	45	104	0.4	0.2	0.7	0.4	0.2	0.7
1-3 years	766	161	605	2.1	0.7	4.2	2.5	0.9	5.0
4-6 years	2,912	703	2,209	7.8	3.1	15.5	10.3	4.0	20.5
7-9 years	7,003	2,917	4,086	18.8	12.7	28.7	29.2	16.7	49.2
10, 11 years	6,045	3,109	2,936	16.3	13.6	20.6	45.4	30.3	69.8
12 years	11,078	8,155	2,923	29.8	35.6	20.5	75.3	65.9	90.3
1-3 years college	5,032	4,251	781	13.6	18.6	5.5	88.8	84.5	95.8
4 or more years college	4,149	3,547	602	11.2	15.5	4.2	100.0	100.0	100.0
Unknown	1,245	397	848						
MOTHER									
Median years completed*	11.8	12.4	10.5						
Total births	47,120	24,025	23,095	100.0	100.0	100.0			
Less than 1 year	66	16	50	0.1	0.1	0.2	0.1	0.1	0.2
1-3 years	383	54	329	0.8	0.2	1.4	1.0	0.3	1.7
4-6 years	2,464	371	2,093	5.3	1.6	9.2	6.3	1.9	10.9
7-9 years	10,455	3,315	7,140	22.5	14.0	31.4	28.8	15.8	42.3
10, 11 years	10,793	4,525	6,268	23.2	19.1	27.6	52.0	35.0	69.9
12 years	14,452	9,551	4,901	31.1	40.3	21.6	83.2	75.3	91.4
1-3 years college	5,076	3,846	1,230	10.9	16.2	5.4	94.1	91.5	96.8
4 or more years college	2,726	2,008	718	5.9	8.5	3.2	100.0	100.0	100.0
Unknown	705	339	366						

* Calculations exclude births to parents with education unknown; calculations for fathers also exclude illegitimate births.

Legitimacy

According to the statements on birth certificates, Mississippi mothers had 8,741 illegitimate babies in 1969. This count was 18.6 percent of the total, and there was a marked difference in the reported occurrence by race - 3.1 percent among white infants and 34.6 percent among nonwhites. Although the total percentage was slightly lower than in the previous year, both race-specific figures were higher, continuing well established upward trends. Whites have experienced a threefold increase since 1957, and percentages for nonwhites have climbed steadily since 1949, more than doubling in that time. In 1969, 40 percent of the children born to mothers under the age of 20 were illegitimate with ratios for those 20 and over ranging from 9 percent to 14 percent (refer to Table B6). The high percentage for mothers under 20 was, of course, correlated with the fact that more of the female population in this age group than of older females were unmarried and, therefore, "at risk" of having an illegitimate child. Another finding regarding mothers of illegitimate children was that many had had previous deliveries (20 percent of the whites and 51 percent of the nonwhites); however, information was not available as to whether the earlier ones were also illegitimate.

Vital Statistics
Mississippi - 1969

Table B6. Illegitimate live births and percent of total live births, by age of mother and race

Age of Mother	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>8,741</u>	<u>740</u>	<u>8,001</u>	<u>18.6</u>	<u>3.1</u>	<u>34.6</u>
10-14	274	17	257	84.0	39.5	90.8
15-19	4,276	376	3,900	38.4	8.6	57.6
20-24	2,441	255	2,186	14.2	2.5	30.3
25-29	858	53	805	8.6	0.9	20.0
30-34	492	24	468	9.9	1.0	18.4
35-39	270	9	261	10.6	0.9	16.4
40-44	109	5	104	13.0	2.1	17.3
45-49	9	0	9	12.3	-	17.0
Unknown	12	1	11	37.5	12.5	45.8

Total-Birth Order

Information on number of previous deliveries to the mother has been tabulated in Mississippi only in 1953, 1954, 1968, and 1969. Concern about the validity of the data was the reason such tabulations were seldom done in the past, and there is still a question as to how reliable such reports are.

Total-birth order, as listed in Table B7, is based on live births plus all fetal deaths regardless of length of gestation. These figures for 1969 show that the number of mothers experiencing their first pregnancy and delivery was 15,894, which was one-third of the total. The number of births declined as birth order increased, of course; but there were 9,645 (20 percent of the total) of the fifth or higher orders including 2,084 of the tenth or higher orders. Twenty or more deliveries each were reported for 18 mothers, all nonwhite, with 25 the largest total. Proportionately more of the nonwhites than of the whites had five or more deliveries, 32 percent in contrast to only 10 percent; and the disparity was especially large for births of the tenth or higher orders. However, comparison of figures for 1968 and 1969 revealed that, despite a rise in the total birth count for 1969, the number to both nonwhite and white mothers in each of the orders sixth or higher went down. Some of the decreases were small, but the fact that there was a decline in each of these birth orders for both races is of significance.

The median age of mothers who had their first delivery in 1969 was 20.0 (refer again to Table B7), and there was a difference of 2.1 years between the medians of 20.9 and 18.8 recorded for white and nonwhite mothers respectively. For all other birth orders medians for nonwhites were also less than those for whites with the differences varying from 0.5 to 3.7 years. The fact that there were 7.0 years between the median ages of white mothers having their first child and those having their fourth delivery but only 5.4 years for nonwhite mothers indicates closer spacing of children by the nonwhites.

Vital Statistics
Mississippi - 1969

Table B7. Live births, percentage distribution, and median age of mother, by total-birth order and race

Total-Birth Order*	Number			Percent			Median Age of Mother		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>47,120</u>	<u>24,025</u>	<u>23,095</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>23.2</u>	<u>23.6</u>	<u>22.6</u>
First	15,894	9,273	6,621	33.7	38.6	28.7	20.0	20.9	18.8
Second	10,377	6,273	4,104	22.0	26.1	17.8	22.3	23.5	20.5
Third	6,588	3,831	2,757	14.0	15.9	11.9	24.5	26.0	22.4
Fourth	4,126	2,148	1,978	8.8	8.9	8.6	26.1	27.9	24.2
Fifth	2,637	1,098	1,539	5.6	4.6	6.7	27.7	29.6	26.3
Sixth	1,817	548	1,269	3.8	2.3	5.5	28.9	31.2	28.0
Seventh	1,347	322	1,025	2.8	1.3	4.4	29.8	32.0	29.4
Eighth	993	180	813	2.1	0.7	3.5	31.7	32.8	31.5
Ninth	767	81	686	1.6	0.3	3.0	32.9	35.4	32.6
Tenth & over	2,084	139	1,945	4.4	0.6	8.4	36.9	37.4	36.9
Unknown	490	132	358	1.0	0.5	1.6			

* Includes all live births plus all fetal deaths regardless of length of gestation.

Last Prior Delivery

Outcome of and time since the last delivery were determined from the dates of last live birth and last fetal death, items which were first included on Mississippi birth certificates in 1968. In 1969, as was true in 1968, the numerous obvious errors and inconsistencies noted in processing the records gave cause for concern about the accuracy of the entries.

Study of outcome of the last prior delivery to mothers who had had previous pregnancies showed that 10 percent were fetal deaths and that 12 percent of the last deliveries to whites but only 7 percent of those to nonwhites terminated in this manner. The median interval between the last delivery and the one in 1969 was 26.6 months (see Table B8), but more births occurred during the second year after the previous delivery than in any other twelve-month period. This concentration occurred among both race groups and following both live births and fetal deaths, but median intervals showed some variation. The median length of time between a fetal death and the live birth in 1969 was close to the same for whites and nonwhites, 18.6 and 17.3 months respectively. However, between a live birth and the birth in 1969 the time was somewhat longer for nonwhites (22.8 months), and the time between two live births to white mothers was substantially longer (34.9 months). This marked difference in spacing by the latter group (white mothers with two live births in succession) was also revealed in the percentage distributions of intervals since last delivery.

Vital Statistics
Mississippi - 1969

Table B8. Percentage distribution of live births to mothers who had had previous pregnancies, by months since last delivery, race, and outcome of last delivery; also median number of months since last delivery, by race and outcome of last delivery

Months since Last Delivery	Total			White			Nonwhite		
	Total*	Outcome of Last Delivery		Total*	Outcome of Last Delivery		Total*	Outcome of Last Delivery	
		Live Birth	Fetal Death		Live Birth	Fetal Death		Live Birth	Fetal Death
<u>Total</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
<u>Under 24</u>	<u>41.1</u>	<u>39.2</u>	<u>58.5</u>	<u>30.8</u>	<u>27.0</u>	<u>57.8</u>	<u>50.7</u>	<u>50.0</u>	<u>59.5</u>
<u>Under 12</u>	<u>4.4</u>	<u>3.6</u>	<u>12.1</u>	<u>2.9</u>	<u>1.7</u>	<u>11.7</u>	<u>5.8</u>	<u>5.2</u>	<u>12.8</u>
12-23	36.7	35.6	46.4	27.9	25.3	46.2	44.9	44.7	46.6
24-59	41.6	43.5	25.0	46.7	49.6	26.2	37.0	38.1	23.0
60 and over	13.4	14.2	6.0	19.2	21.1	6.3	7.9	8.1	5.5
Unknown	3.8	3.1	10.5	3.2	2.3	9.6	4.4	3.8	12.0
<u>Median months**</u>	<u>26.6</u>	<u>27.7</u>	<u>18.0</u>	<u>32.8</u>	<u>34.9</u>	<u>18.6</u>	<u>22.5</u>	<u>22.8</u>	<u>17.3</u>

* Excludes those with outcome of last delivery unknown.

** Calculated from ungrouped data.

Prenatal Care

Data on prenatal care were collected in Mississippi for the first time in 1968 when items on month of pregnancy care began and number of visits were added to the birth certificate. Although the proportion of records lacking this information was smaller in 1969 than in 1968, reporting on this subject was still thought to be less reliable than that in most other sections of the certificate. Table B9 shows that only 958 mothers (2 percent of the total) had no medical care during pregnancy, and both this number and the percentage were lower than in 1968. Relatively more nonwhites than whites lacked care, but even the figure of 3.5 percent for nonwhites was not of great consequence. For 1,676 other mothers it was unknown whether there was any care, leaving over 94 percent of the total that were known to have had at least one prenatal visit to a physician or clinic.

Care was sought during the first three months of pregnancy by 58 percent of the mothers, 26 percent had care beginning in the second trimester, and 9 percent did not have any care until the last trimester. It was in the timing and amount of prenatal care rather than whether any was received that there were important differences between the races. For 76 percent of the white mothers, but only 40 percent of the nonwhites, care began during the first trimester; in addition, the proportion delaying care until the last trimester was 4 percent for whites but 14 percent for nonwhites. Study of the relationship between the amount of education the mother had received and the stage of pregnancy in which prenatal care began revealed a strong correlation between the two. As the number of years of schooling increased the percentage of mothers whose care began in the first trimester also increased, and this was true for both whites and nonwhites (see Table B10). However, it was also found, when mothers of the same educational level were compared, that relatively fewer of the nonwhites than of the whites began care in the first trimester.

Vital Statistics
Mississippi - 1969

Table B9. Live births and percentage distribution, by whether mother received prenatal care, time during pregnancy care began, number of visits to physician or clinic, and race

Prenatal Care, Time Care Began, and Number of Visits	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>47,120</u>	<u>24,025</u>	<u>23,095</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
No prenatal care	958	147	811	2.0	0.6	3.5
<u>Some prenatal care</u>	<u>44,486</u>	<u>23,212</u>	<u>21,274</u>	<u>94.4</u>	<u>96.6</u>	<u>92.1</u>
First trimester	27,579	18,247	9,332	58.5	76.0	40.4
Second trimester	12,204	3,958	8,246	25.9	16.5	35.7
Third trimester	4,085	887	3,198	8.7	3.7	13.8
Unknown month	618	120	498	1.3	0.5	2.2
Average number of visits	8.7	10.9	6.2			
1-5 visits	12,564	2,106	10,458	26.7	8.8	45.3
6-10 visits	16,268	8,427	7,841	34.5	35.1	34.0
11-15 visits	12,498	10,223	2,275	26.5	42.6	9.8
16 visits or more	2,464	2,121	343	5.2	8.8	1.5
Unknown number	692	335	357	1.5	1.4	1.5
Unknown whether any prenatal care	1,676	666	1,010	3.6	2.8	4.4

Table B10. Percentage distribution of live births, by race, years of school completed by mother, and prenatal care

Race and Years of School Completed by Mother	Total	No Prenatal Care	Trimester Prenatal Care Began			Prenatal Care Unknown
			First	Second	Third	
ALL RACES						
0-6 years	100.0	4.4	36.0	37.9	14.1	7.6
7-9 years	100.0	3.4	44.0	33.8	13.4	5.3
10, 11 years	100.0	2.4	52.3	31.0	10.6	3.7
12 years	100.0	1.1	68.6	21.8	6.3	2.2
Some college	100.0	0.7	81.3	13.4	3.1	1.5
WHITE						
0-6 years	100.0	2.3	51.9	30.6	10.9	4.3
7-9 years	100.0	1.5	63.4	25.1	7.1	2.8
10, 11 years	100.0	0.7	70.4	21.1	5.1	2.6
12 years	100.0	0.4	79.9	15.1	2.7	1.8
Some college	100.0	0.3	86.5	9.9	2.0	1.3
NONWHITE						
0-6 years	100.0	4.7	33.2	39.2	14.7	8.2
7-9 years	100.0	4.3	35.0	37.9	16.4	6.4
10, 11 years	100.0	3.7	39.2	38.1	14.5	4.5
12 years	100.0	2.4	46.5	35.0	13.2	2.9
Some college	100.0	1.8	65.8	23.7	6.5	2.2

Vital Statistics
Mississippi - 1969

Again referring to Table B9, it can be seen that white mothers not only began prenatal care earlier than nonwhites did but that they also received a greater amount of care, as would be expected if care began earlier. In fact, 51 percent of the whites, in contrast with only 11 percent of the nonwhites, were known to have had more than 10 visits each. The average number of visits per mother for whom the count was reported was 10.9 for whites and only 6.2 for nonwhites.

Complications during Pregnancy and Labor

Medical information regarding complications of the mother during pregnancy and labor were collected in 1968 and 1969. However, in both of the years since items on this subject were added to Mississippi's birth certificates, the data compiled were of doubtful value because of the quality of reporting and the difficulties in classifying material for which there are no defined limits. Although some improvement over the first year was noted for both races, one out of every twelve records for 1969 did not indicate whether there were any complications, and on a number of other records which did show at least one complication the information was missing from one or two of the other major categories. Many complications which were listed were incompletely stated and would have been difficult to classify even if a satisfactory classification system had been developed.

In processing the records it was noted that those for Hinds County appeared to have more information on this subject than those for most other counties. This county, which has almost ten percent of the state's population, is the location of the University of Mississippi Medical Center and several large hospitals and, therefore, could be expected to provide better data because of the concentration of medical facilities and specialists. Information on complications among Hinds County residents was tabulated and results are shown in Table B11 along with data for the entire state.

Table B11. Percentage distribution of live births by whether mother had complications during pregnancy and labor, category of complication, and race, Mississippi and Hinds County residents

Complications	Mississippi			Hinds County		
	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
Without complications	80.8	79.0	82.7	75.3	78.6	72.2
With complications	10.7	12.2	9.1	23.2	19.6	26.4
Related to pregnancy	(4.5)	(4.7)	(4.4)	(10.2)	(7.2)	(12.9)
During but not related to pregnancy	(2.3)	(2.6)	(2.1)	(4.1)	(3.5)	(4.6)
Of labor	(5.5)	(6.6)	(4.3)	(13.0)	(12.2)	(13.8)
Unknown whether any complications	8.5	8.8	8.1	1.6	1.8	1.4

Comparison of the two sets of figures revealed that the reported incidence of complications was significantly higher for Hinds County mothers; 23 percent had complications in contrast with only 11 percent on a statewide basis. Fewer of the Hinds County residents were in the "unknown" category, but this difference could only partially account for the disparity in incidence. Another contrast noted was in the occurrence by race; the state figures showed a lower incidence among nonwhites than among whites but, according to the

Vital Statistics
Mississippi - 1969

Hinds County figures, the reverse was true. Since it can be assumed that the occurrence of complications among mothers in Hinds County is actually no different from that among other mothers in Mississippi, it follows that the reporting for this county was more complete than that for the state as a whole and that the Hinds County figures are, therefore, more reliable.

Table B12. Incidence rates* for complications during pregnancy and labor reported on live birth certificates for residents of Hinds County, by type of complication and race

Type of Complication and International Classification Number**	Total	White	Non-white
<u>Total complications reported</u>	<u>291.7</u>	<u>241.1</u>	<u>337.7</u>
<u>Total related to pregnancy (630-639)</u>	<u>107.1</u>	<u>74.1</u>	<u>137.1</u>
Toxemias (636-639)	53.6	27.7	77.1
Hemorrhage (632)	12.5	8.4	16.2
Premature rupture of membrane without delivery (634.4)	12.5	12.2	12.8
Anemia (633)	9.8	3.3	15.8
Urinary infections (635)	3.1	2.8	3.4
All others	15.6	19.7	11.9
<u>Total during but not related to pregnancy (760, 761)</u>	<u>43.3</u>	<u>36.6</u>	<u>49.4</u>
Chronic (and unspecified) circulatory diseases (760.0-760.3)	6.2	4.7	7.7
Diabetes mellitus (761.1)	6.0	3.3	8.5
Chronic (and unspecified) genitourinary diseases (760.4, 760.5)	4.9	4.7	5.1
All others	26.1	23.9	28.1
<u>Total of labor (651-662)</u>	<u>141.3</u>	<u>130.4</u>	<u>151.2</u>
Fetopelvic disproportion (655)	33.5	41.3	26.4
Malpresentation of fetus (656)	29.2	33.8	25.1
Premature rupture of membranes (661.0)	16.5	12.7	20.0
Obstetrical trauma (658-660)	12.7	3.3	21.3
Hemorrhage (651-653)	9.6	7.5	11.5
Previous cesarean section (661.2)	8.0	8.9	7.2
Umbilical cord complication (661.3)	5.1	4.2	6.0
Abnormality of bony pelvis (654)	4.0	4.7	3.4
All others	22.5	14.1	30.2

* Complications per 1,000 live births.

** From the Eighth Revision International Classification of Diseases, Adapted for Use in the United States.

Complications were classified within the three major categories by using sections of the Eighth Revision International Classification of Diseases, Adapted for Use in the United States. Since the index was not available in 1968 until records for eleven months had already been processed, allocation was undoubtedly more accurate in 1969, but results were still not considered satisfactory. Listed in Table B12 are the incidence rates for the major categories and for the specific classifications most frequently reported for Hinds County mothers. Among white mothers more complications occurred at the time of delivery than during pregnancy, but among nonwhites more occurred during pregnancy. However, nonwhites experienced more of each major category of complications than did the whites.

Vital Statistics
Mississippi - 1969

The highest specific rate on the table was that for toxemias of pregnancy among nonwhite women; it was nearly three times the next highest rate for this group and not far from being twice as high as the highest rate (that for fetopelvic disproportion) for white mothers. Additional complications listed among the top five were malpresentation of the fetus at time of delivery, premature rupture of membranes either during pregnancy or at delivery, and hemorrhages during pregnancy or labor.

Length of Gestation and Birth Weight

Birth weight has been reported and tabulated in the same manner since it was first added to the Mississippi birth certificate in 1949. However, information about length of gestation has been obtained in various ways since 1929, with none completely satisfactory. Beginning with the 1968 revision of the birth certificate, the attendant was asked to provide the date the last normal menstrual period (LMP) prior to delivery began rather than state the length of gestation. Although calculation of the number of completed weeks of gestation from the LMP to the date of delivery has produced more valid results than were obtained by other methods of securing this type of data, certain problems have not been resolved. Inaccuracies in the LMP are probably more numerous than those in any other item on the certificate since it is a date which is often unnoted at the time but must be recalled later, sometimes a considerable length of time later. Findings were essentially the same in 1968 and 1969, with small decreases in the proportions under 40 weeks and of unknown length of gestation and a corresponding rise in the proportion with gestational period 40 or more weeks.

Table B13. Live births and percentage distribution, by maturity and race

Maturity*	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>47,120</u>	<u>24,025</u>	<u>23,095</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
Immature only**	2,030	797	1,233	4.3	3.3	5.3
Premature only**	4,986	1,480	3,506	10.6	6.2	15.2
Both immature and premature	2,422	831	1,591	5.1	3.4	6.9
Total immature	4,452	1,628	2,824	9.4	6.8	12.2
Total premature	7,408	2,311	5,097	15.7	9.6	22.1
Total immature and/or premature	9,438	3,108	6,330	20.0	12.9	27.4
Neither immature nor premature**	37,682	20,917	16,765	80.0	87.1	72.6

* Immature - birth weight 5 1/2 pounds (2,500 grams) or less; premature - gestation less than 37 weeks.

** Births classified as immature only or as premature only include some for which the other measurement was unknown. Those classified as neither immature nor premature include some for which one or both measurements were unknown.

Length of gestation and birth weight were used to compile the information relating to maturity in Table B13. Lack of maturity as measured by birth weight is indicated by the

Vital Statistics
Mississippi - 1969

term "immature" (5 1/2 pounds or less or 2,500 grams or less) and the term "premature" denotes a gestation period of less than 37 weeks. The two measurements are closely associated but, as the figures show, a low birth weight and a shortened gestation period do not necessarily occur simultaneously. Children affected by both prematurity and immaturity numbered 2,422 and made up 5 percent of the total. However, the 2,030 others also classified as immature but not premature raised the total immaturity count and rate to 4,452 and 9.4 percent. Corresponding totals for prematurity were 7,408 and 15.7 percent. Altogether, exactly 20 percent of the babies (9,438) showed a lack of maturity by one or both criteria; and the percentage for nonwhites was more than double that for whites, 27.4 in comparison with 12.9. It was noted, however, that the racial disparity in weight was less than the difference in length of gestation.

Sex

As usual, males outnumbered females. Totals were 24,026 and 23,094 respectively, with the sex ratio calculated from these counts 1,040 males per 1,000 females. The sex ratio is a figure that has varied erratically with no trend or pattern; during the last 26 years it has ranged from 1,024 to 1,052. In this entire period the ratio for whites was always higher than that for nonwhites, and the figures for 1969 were 1,051 for white babies and 1,029 for nonwhites.

Plurality (See Plurality in Fetal Death section also)

Twins and triplets accounted for 2.3 percent of the live births with the total number from plural deliveries 1,096. The percentage for nonwhites was 3.0 but that for whites was only 1.7. It was also noted that the proportion of plural births rose steadily from 1.5 percent among mothers 15-19 years old to a peak of 4.2 percent at ages 35-39 and then declined and that at all ages except 10-14 years the occurrence of multiple births was more frequent among nonwhites than among whites. Study of plurality in relation to birth weight and length of gestation revealed that plural births were far more likely to be of low birth weight or delivered before term than single births. The incidence of "immaturity" for single and plural births was 9 and 59 percent respectively; comparable figures on "prematurity" were 16 and 45 percent. The disparity between single and plural births in lack of maturity (by either measurement) was greater for white babies than for the nonwhites.

Birth Injuries and Congenital Malformations

In 1968 sections on birth injuries and congenital malformations were added to Mississippi's birth certificates. Many difficulties were experienced in obtaining these facts, but data collected during the second year of reporting showed improvement in that the proportion of records with no information or with unusable information declined to about 8 percent for each item. However, the reported incidence of babies with one or both problems remained the same - 1.1 percent. The true incidence was undoubtedly higher as some malformations are not apparent at birth but are only diagnosed later when symptoms appear and others can be recognized by physicians but not by the midwives and other nonmedical persons who attended 12 percent of the deliveries in 1969. Also, it is likely that some malformations were not listed because of the fear of stigmatizing a child and that some birth injuries were not recorded because of possible blame or even court action against the attendant. As Table B14 shows, 523 babies were affected by malformations and/or injuries; 3 of these children had both. Since a larger proportion of the white than

Vital Statistics
Mississippi - 1969

of the nonwhite children were born in hospitals, better diagnosis and reporting may account for the higher incidence noted for whites.

Table B14. Live births and percentage distribution, by whether child had any birth injuries or congenital malformations, whether injury or malformation, and race

Birth Injuries and Congenital Malformations	Number			Percent		
	Total	White	Non- white	Total	White	Non- white
<u>Total</u>	<u>47,120</u>	<u>24,025</u>	<u>23,095</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
Without injury or mal- formation	42,488	21,693	20,795	90.2	90.3	90.0
With injury and/or mal- formation	523	333	190	1.1	1.4	0.8
Injury	(104)	(66)	(38)	(0.2)	(0.3)	(0.2)
Malformation	(422)	(270)	(152)	(0.9)	(1.1)	(0.6)
Other*	4,109	1,999	2,110	8.7	8.3	9.1

* Both categories unknown or one category none and other unknown.

Only 104 babies were reported to have suffered a birth injury; 3 children had two injuries each and 1 child had three injuries making a total of 109 injuries reported. There were 53 injuries to bone or nerve tissue recorded, 19 cases of brain injury, none to the spinal cord, and 37 in other categories.

A total of 503 congenital malformations were reported among 422 children; 353 had only one malformation each, 57 had two each, and 12 had three or more each. According to the numbers in Table B15 which shows the principal types, there were more malformations of the musculoskeletal system than of any other system or site. However, this finding was not surprising as defects in this system are usually quite conspicuous, especially polydactyly (extra fingers or toes) which accounted for 66 of the 179 reported. Ranking second was the digestive system, and it also included a sizeable number of an easily recognized type - cleft palate and cleft lip with 58 of the 102 listed in this category.

Table B15. Congenital malformations reported on live birth certificates, by system or organs affected and race

System or Organs Affected and International Classification Number*	Total	White	Non- white
<u>Total malformations reported</u>	<u>503</u>	<u>334</u>	<u>169</u>
Musculoskeletal system (754-756)	179	90	89
Digestive system (749-751)	102	76	26
Nervous system (740-743)	62	48	14
Genital organs (752)	42	34	8
All others	118	86	32

* From the Eighth Revision International Classification of Diseases, Adapted for Use in the United States.

FETAL DEATHS (See Table 4)

Reporting and Reliability

Regulations of the Mississippi State Board of Health have required the registration of all fetal deaths, regardless of the length of gestation, since 1952. Prior to that year only stillbirths of 20 or more weeks gestation had to be reported. Despite the fact that 1969 was the eighteenth year of all-inclusive registration, there were indications that underreporting of early fetal deaths as well as those that occur in advanced stages of pregnancy was still a serious problem. Because of this, the data tabulated on fetal deaths are incomplete and may be biased.

Trend and Race

Fetal deaths registered in the state in 1969 numbered 2,514, and 2,471 records for Mississippi residents were received. The fetal death ratio of 52.4 per 1,000 live births was the highest recorded since reporting without regard to length of gestation began, but it is not known whether this represents a true increase or merely more complete reporting. Race-specific ratios were 49.0 for whites and 56.0 for nonwhites.

County

Study of county ratios revealed differences far in excess of what could be accounted for by variations in age distribution of the population, fertility, medical care, etc. Among the white population ratios ranged from zero to 135.1 and for nonwhites the range was from zero to 136.4. Undoubtedly much of this extreme variation can be attributed to differences in completeness of reporting. If, as some researchers have estimated, the true incidence of fetal death is one out of every five fetuses or worse, complete reporting would produce ratios of 250 or more.

Attendant

As the figures in Table C1 show, 89 percent of the fetal deaths reported were delivered in hospitals or clinics. Three percent were delivered by physicians in homes or other places, and only midwives or other nonmedical persons were present for 8 percent or 193 deliveries. There has been an upward, but irregular, trend in hospitalization when a fetal death occurs even though the percentage hospitalized dropped slightly in 1969. More of the whites (97 percent) than of the nonwhites (82 percent) were hospitalized, but the discrepancy of 15 percentage points was less than the difference between the two race groups (25 percentage points) in hospitalization for a live birth. In fact, a greater proportion of nonwhites were hospitalized for a fetal death than for a live birth, probably because fetal deaths frequently occur in conjunction with complications and/or premature delivery.

Table C1. Fetal deaths and percentage distribution, by attendant and race

Attendant	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>2,471</u>	<u>1,177</u>	<u>1,294</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
Hospital or clinic	2,199	1,140	1,059	89.0	96.8	81.8
Physician, other place	79	34	45	3.2	2.9	3.5
Midwife	175	0	175	7.1	-	13.5
Other	18	3	15	0.7	0.2	1.2

Vital Statistics
Mississippi - 1969

Age of Mother

Fetal loss and age of mother appear to be correlated, as evidenced by the figures in Table C2. The lowest fetal death ratios were those for mothers 15-24 years of age with increases noted as age increased. Mothers under 15 years of age also experienced a higher than average fetal death ratio. Although the total ratio for nonwhites was greater than that for whites, it was only among mothers in their twenties that excessive ratios for nonwhites occurred (the age group 10-14 years was excluded from this comparison because of the small number of white births). In other age groups the race-specific ratios were about the same or that for the white mothers was the higher.

Table C2. Fetal deaths and fetal death ratios, by age of mother and race

Age of Mother	Number			Ratio*		
	Total	White	Non-white	Total	White	Non-white
<u>All ages</u>	<u>2,471</u>	<u>1,177</u>	<u>1,294</u>	<u>52.4</u>	<u>49.0</u>	<u>56.0</u>
10-14	25	2	23	76.7	#	81.3
15-19	443	176	267	39.8	40.2	39.4
20-24	749	400	349	43.5	39.9	48.4
25-29	527	277	250	52.8	46.4	62.2
30-34	354	171	183	71.5	71.1	71.9
35-39	246	100	146	96.7	105.2	91.6
40-44	98	38	60	117.4	163.1	99.7
45-49	9	4	5	#	#	#
Unknown	20	9	11			

* Fetal deaths per 1,000 live births to mothers in specified group.

Ratio not calculated on less than 100 live births.

Education of Parents

The year 1969 was the second in which data on highest grade of school completed by the father and mother was collected. Although there was some improvement over the previous year in completeness of reporting, counts for 1969 showed an appallingly large proportion of records with no data in these sections - 19 percent (based only on those reported as legitimate) had no information on the father's education and on 17 percent the mother's education was not given. For several of the medical and socio-economic items which are tabulated for both live births and fetal deaths, the data revealed that circumstances or conditions were more unfavorable in cases which ended in fetal death, but tabulations on education of parents showed no such disparity. Figures relating to fetal deaths can be summarized as follows: 44 percent of the fathers and 51 percent of the mothers had not completed high school and about 30 percent of each had not gone beyond the ninth grade; also, there were important differences between educational levels of white and nonwhite parents with the median years of school completed by nonwhite fathers and mothers 2.9 and 2.0 years less respectively than medians for white fathers and mothers.

Legitimacy

Fetal deaths reported as illegitimate made up 21 percent of the total with the figure for nonwhites far higher than that for whites. Among nonwhites the proportion of fetal deaths that were illegitimate was the same as the proportion of live-born children

Vital Statistics
Mississippi - 1969

that were illegitimate - 35 percent. However, 5 percent of the white fetal deaths but only 3 percent of the white live-born children were in this category.

Total-Birth Order

Classification of fetal deaths by number of previous deliveries to the mother began in Mississippi in 1968. The records were so poorly filled out, however, that for 11 percent of the 1969 events the total number of previous deliveries could not be determined and the accuracy of some of the information that was provided was doubtful. Total-birth order, which includes live births plus fetal deaths of all gestational periods, was computed whenever possible; and it was found that the percentage that were first deliveries was exactly the same as in 1968 - 24.5 percent. Although 70 percent of the live births in 1969 were included among mothers' first three deliveries, the first, second, and third total-birth orders accounted for only 53 percent of the fetal deaths. Twenty-three was the largest total number of deliveries reported, but this was less than the high of 25 which was the largest number reported for a mother who had a live birth in 1969. Figures by race showed relatively more whites than nonwhites of the first four orders and a higher proportion of nonwhites than of whites in the higher orders with an especially large excess of nonwhite fetal deaths of tenth or higher orders.

Fetal death ratios given in Table C3 for each total-birth order revealed a pattern of steady increase from a low rate of 38.1 fetal deaths per 1,000 live births among first deliveries to a high of 82.4 among those of the seventh order. Ratios in orders above the seventh varied irregularly, probably at least in part because of the small numbers involved. Even though the overall fetal death ratio for nonwhites was higher than that for whites, only the ratios for nonwhites of the first three orders were higher than those for whites.

Table C3. Fetal death ratios*, by total-birth order and race

Total-Birth Order**	Total	White	Non-white
First	38.1	35.8	41.2
Second	40.6	39.7	41.9
Third	44.6	41.0	49.7
Fourth	52.1	54.9	49.0
Fifth	58.0	64.7	53.3
Sixth	61.6	74.8	55.9
Seventh	82.4	111.8	73.2
Eighth	75.5	105.6	68.9
Ninth	78.2	#	65.6
Tenth and over	77.7	93.5	76.6

* Fetal deaths per 1,000 live births in specified group.

** Includes all live births plus all fetal deaths regardless of length of gestation.

Ratio not calculated on less than 100 live births.

Vital Statistics
Mississippi - 1969

Last Prior Delivery

Dates of the last live birth and last fetal death were new items added to Mississippi's fetal death certificates in the revision of 1968. Incomplete reporting and obviously incorrect entries were serious problems in both 1968 and 1969; and, although the situation improved slightly in the second year, information on 14 percent of the records was inadequate to establish whether the last delivery was a live birth or a fetal death and for 20 percent the length of time since the last delivery could not be determined.

Study of the relationship between the outcome of the last prior delivery to mothers who had had previous pregnancies and the outcome of the deliveries in 1969 revealed that a fetal death was more likely to occur if the previous delivery was also a fetal death. The fetal death ratio among mothers whose last delivery produced a living child was 44.5 fetal deaths per 1,000 live births, but the ratio among mothers whose last delivery resulted in fetal death was more than twice as high - 94.2. In addition, it was noted that, although only 10 percent of the live births in 1969 that were from second or later pregnancies had been preceded by a fetal death, the proportion of fetal deaths preceded by a fetal death was 19 percent.

For mothers who had had previous pregnancies, table C4 provides information about spacing patterns in relation to outcome of delivery. According to these figures the median length of time (25.4 months) between a fetal death in 1969 and the last prior delivery was just one month less than the median interval (26.6 months) between a live birth in 1969 and the last prior delivery. However, there were variations by race and outcome of the last delivery. The longest median intervals were those of about three years for white mothers whose last delivery was a live birth; for nonwhite mothers whose last delivery was a live birth, the median intervals were only about two years. The shortest median intervals were those for mothers whose last delivery was a fetal death, but race was not an important factor here as all these medians were within the range 15-19 months.

Table C4. Median number of months since last delivery to mothers who had had previous pregnancies, by race, outcome of last delivery, and outcome of delivery in 1969

Race and Outcome of Last Delivery	Outcome of Delivery in 1969	
	Live Birth	Fetal Death
<u>Total*</u>	<u>26.6</u>	<u>25.4</u>
White: live birth	34.9	36.1
fetal death	18.6	16.7
Nonwhite: live birth	22.8	23.9
fetal death	17.3	14.9

* Excludes those with outcome of last delivery unknown.

Prenatal Care

Collection of data on month of pregnancy in which prenatal care began and the number of prenatal visits to a physician or clinic was initiated in Mississippi in 1968. Considerable difficulty was experienced in obtaining these facts, and in 1969 there was still

Vital Statistics
Mississippi - 1969

a substantial proportion (19 percent) of fetal death records from which it was not possible to determine whether the mother had any prenatal care. In addition, inaccuracies on and lack of completeness of the records which did indicate that the mother had prenatal care further reduced the validity of this category of information.

It was reported that 16 percent of the mothers had no prenatal care (see Table C5), but since nearly one-fifth of the records had no definite information on this subject it is likely that the true proportion without care was over 20 percent and perhaps over 30 percent. However, even the reported figure of 16 percent was considerably larger than the figure of 2 percent for mothers of children born alive. Sixty-five percent of the mothers who experienced a fetal death were known to have had some prenatal care, 50 percent had care during the first trimester of pregnancy, and only 3 percent had no care until the third trimester. Differences between the percentages of white and non-white mothers with and without prenatal care were small, but there was a disparity of note in the figures on stage of pregnancy in which care began. For white mothers the proportion seeking care in the first trimester was 61 percent in contrast to only 41 percent of nonwhite mothers.

Table C5. Percentage distribution of live births and fetal deaths, by whether mother received prenatal care and race, and average number of visits per mother who received care, by race

Prenatal Care and Visits	Live Births			Fetal Deaths		
	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
No care	2.0	0.6	3.5	16.1	15.9	16.4
Some care	94.4	96.6	92.1	65.2	67.9	62.8
Unknown whether any care	3.6	2.8	4.4	18.6	16.2	20.9
<u>Average number of visits</u>	<u>8.7</u>	<u>10.9</u>	<u>6.2</u>	<u>4.5</u>	<u>4.6</u>	<u>4.4</u>

From data on number of prenatal visits it was determined that the average number was 4.5 per mother who received care. This was only about half as many as the average of 8.7 for mothers of live-born children, but many fetal deaths occurred in the early stages of pregnancy and, therefore, there was not so long a period of time in which the mothers could have received care and the two numbers are not truly comparable. Although race-specific averages were almost identical, this fact actually indicates that non-white mothers had relatively less care because far more of their reported fetal deaths than of those among white mothers occurred during the second and third trimesters thereby increasing the period of time during which prenatal care would have been needed and the number of visits they could have been expected to have.

Length of Gestation and Birth Weight

Determination of the length of gestation by calculating the number of completed weeks between the date of the last normal menstrual period (LMP) and the date of delivery was undertaken for the first time in Mississippi in 1968 when the date of the LMP was added to the certificate. Many problems were encountered, with the result that the

Vital Statistics
Mississippi - 1969

number of weeks gestation could not be calculated for 14 percent of the fetal deaths in 1969 and the accuracy of the intervals that were computed probably was more questionable than that of any other item tabulated.

The distribution of fetal deaths by length of gestation is given in Table C6 which shows that 45 percent of the total occurred in the first 20 weeks of pregnancy with concentration in the period 10-19 weeks and that only 15 percent occurred at or near term (at 37 or more weeks gestation). There was a pronounced contrast in the distributions by race; 59 percent of the whites but only 32 percent of the nonwhites were under 20 weeks of gestation and, conversely, there were proportionately more nonwhites of 20 or more weeks gestation. Calculation of fetal death ratios using only the fetal deaths of 20 or more weeks gestation showed that the ratio for nonwhites was nearly double that for whites. Underreporting is considered to be less of a problem among fetal deaths of 20 or more weeks than among early fetal deaths; and, therefore, the true incidence of fetal deaths of 20 or more weeks gestation probably was higher among nonwhites than among whites as the tabulated data showed. It is not known, however, whether the observed excess of whites among fetal deaths of less than 20 weeks gestation represents the actual relationship or resulted from considerable underreporting of early fetal deaths among nonwhites.

Table C6. Percentage distribution of fetal deaths, by completed weeks of gestation and race

Weeks of Gestation	Total	White	Non-white
<u>Total</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
<u>Under 20</u>	<u>45.2</u>	<u>59.0</u>	<u>32.5</u>
Under 10	11.3	15.2	7.8
10-19	33.8	43.8	24.7
<u>20-36</u>	<u>26.4</u>	<u>19.3</u>	<u>32.9</u>
20-27	13.5	11.3	15.4
28-36	13.0	8.0	17.5
<u>37 and over</u>	<u>14.6</u>	<u>10.5</u>	<u>18.4</u>
37-39	7.3	5.8	8.6
40	2.3	1.4	3.0
Over 40	5.1	3.3	6.7
Unknown	13.8	11.1	16.2

Since a large proportion of fetal deaths occurred when the fetus was quite small, it is not surprising that no birth weight was given for 58 percent of the total delivered. Because of this no analysis of weight is presented.

Sex

The sex ratio calculated from records with sex stated was 1,355 males per 1,000 females and, as usual, was much higher than the sex ratio for live births which was 1,040 in 1969. Due to the fact that a large percentage of the fetal deaths registered occurred in the early weeks of pregnancy, the sex was missing from 1,143 or 46 percent of the records.

Vital Statistics
Mississippi - 1969

It is not known whether this large number of undetermined sex obscured a more even distribution of males and females than the sex ratio indicated or whether a greatly excessive number of males was conceived and then suffered high mortality rates during pregnancy and labor.

Plurality

Fetuses from plural deliveries made up 4.0 percent of the total fetal deaths in contrast with a figure of only 2.3 percent among live births. More of the nonwhite fetal deaths (5.0 percent) than of the white fetal deaths (2.9 percent) were twins and triplets. It is likely that the true occurrence of plurality among fetal deaths was greater than the reported figures show because of failure, particularly by midwives, to recognize the condition among fetal deaths that occur early in pregnancy.

By combining plurality data from live birth and fetal death records it was found that the outcome of 1 out of every 82 pregnancies was a multiple birth and that the frequency was much greater among nonwhites - 1 out of every 64 - than among whites - 1 out of every 115. Altogether, 587 sets of twins and 7 sets of triplets were delivered in 1969. All members of the set were born alive in 89 percent of these cases, in 6 percent there was a combination of live and dead members, and in 5 percent all were born dead. The fetal death ratio for multiple births was 90.3 per 1,000 live births but was only 51.5 for single births.

Cause of Fetal Death, Complications during Pregnancy and Labor, Birth Injuries, and Congenital Malformations

Fetal death records contain sections relating to each of the items listed in the above title. However, the information received on these topics was inadequate for tabulation and analysis.

DEATHS (See Tables 5-8, 12-16)

Trend and Race

A slight drop in the death toll for Mississippians brought the count for 1969 to 23,865. However, this total was the second highest since 1937 and the figure of 13,873 for the white population was the largest ever recorded for that group. In contrast, the number of deaths among nonwhites was 9,992 which was an unusually low total for this group. Race-specific death rates were 10.0 per 1,000 white population and 12.1 for nonwhites. Both of these rates as well as the 10.8 calculated for the total population were higher than those recorded for the previous year; but, as discussed in the section on population, the change from estimated to census population as a base for computing the rates affected their comparability.

County

There was a substantial amount of variation in death rates by county. The highest rate for white persons was 17.4 in Yalobusha County and in two counties, DeSoto and Jackson, a low rate of 5.8 was recorded. The lowest rate for nonwhites was 5.4 in Perry County with

Vital Statistics
Mississippi - 1969

16.6 for Simpson County at the upper limit of the range. Strangely enough, the range of rates for the nonwhites (5.4 to 16.6) was slightly lower than the range for whites (5.8 to 17.4) even though the state figure for nonwhites was higher than that for whites.

Month

According to the monthly distribution of deaths given in Table D1, mortality was, as usual, highest in the coldest part of the year. January was the month with the largest number of deaths and the highest daily average with December ranking second.

Table D1. Deaths and average per day, by month

Month	Deaths	Average per Day	Month	Deaths	Average per Day
<u>Total</u>	<u>23,865</u>	<u>65</u>			
January	2,344	76	July	1,983	64
February	1,915	68	August	2,014	65
March	2,079	67	September	1,779	59
April	1,853	62	October	1,960	63
May	1,894	61	November	1,980	66
June	1,901	63	December	2,163	70

Age at Death

The median age at death for the total population was 68.4; for white persons it was 70.2 and for nonwhites, 65.8. All three of these figures were a fraction of a year less than the comparable medians for 1968, but were still at relatively high levels with no indication of a reversal of the generally upward trends of recent years. The difference between the race-specific medians was 4.4 years, a slightly larger figure than in the previous year but also the second smallest difference recorded in a number of years.

Data on deaths in specific age groups are presented in Table D2. These figures show that, after the high-risk period of infancy, the number of deaths and the death rate dropped sharply to a low level in the groups 5-9 and 10-14 years of age and then rose steadily with the largest numbers of deaths at ages 65-74 and 75-84 and a peak rate in the group 85 and over. At all ages except the oldest group Mississippi rates were higher than corresponding rates for the United States as a whole. It was also noted that the rates for nonwhites were higher than those for whites except in the oldest group and that in some cases the rate for the nonwhite population was more than double that for whites of the same age. This partially explains Mississippi's high rates for the total population, but further comparison revealed that, even when the age-specific rates were also made race-specific, most of those for Mississippi were higher than the corresponding ones for the U.S. The age distribution also indicated how excessive the mortality among nonwhite infants was in that 10 percent of the nonwhites who died were children under one year of age in contrast with a figure of only 4 percent for whites.

Vital Statistics
Mississippi - 1969

Table D2. Deaths, percentage distribution, and rates, by age and race

Age	Number			Percent			Rate**		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>23,865</u>	<u>13,873</u>	<u>9,992</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>	<u>10.8</u>	<u>10.0</u>	<u>12.1</u>
Under 1	1,513	530	983	6.3	3.8	9.8	34.3	22.4	48.3
1-4	237	71	166	1.0	0.5	1.7	1.4	0.8	2.1
5-9	144	68	76	0.6	0.5	0.8	0.6	0.5	0.7
10-14	162	62	100	0.7	0.4	1.0	0.6	0.4	0.9
15-24	665	369	296	2.8	2.6	3.0	1.6	1.5	1.9
25-34	564	271	293	2.4	2.0	2.9	2.3	1.6	4.2
35-44	964	493	471	4.0	3.6	4.7	4.3	3.1	7.4
45-54	2,053	1,179	874	8.6	8.5	8.7	9.4	7.6	13.5
55-64	3,818	2,282	1,536	16.0	16.4	15.4	18.8	16.7	23.3
65-74	5,319	3,095	2,224	22.3	22.3	22.2	37.6	34.4	43.2
75-84	5,373	3,550	1,823	22.5	25.6	18.2	84.5	84.1	85.1
85 and over	3,041	1,900	1,141	12.7	13.7	11.4	175.6	180.9	167.4
Unknown	12	3	9	*	*	0.1			

* Less than 0.06 percent.

** Deaths per 1,000 population in specified group.

Leading Causes of Death

Eighty percent of the deaths resulted from the ten leading causes listed in Table D3. These are the same causes which were listed in 1968, and the order was the same except that certain causes of mortality in early infancy and influenza and pneumonia exchanged places, ranking fifth and sixth respectively in 1969. The first nine of these causes were also the same as those listed for the nation (with a reversal in the order of those ranking fifth and sixth), but homicide was replaced by cirrhosis of the liver in tenth place for the U.S. Relatively more of the deaths among whites than among nonwhites were due to these ten leading causes, chiefly because of a significantly larger proportion of deaths due to heart diseases in the white population. Other differences by race were variations in order, the fact that suicide rather than homicide appeared in tenth place on the list for whites, and the listing of hypertension and nephritis and nephrosis for nonwhites instead of arteriosclerosis and the bronchitis-emphysema-asthma group.

Vital Statistics
Mississippi - 1969

Table D3. Deaths and percent of total deaths from ten leading causes, by cause and race

Cause and International Classification Number	Number			Percent		
	Total	White	Non- white	Total	White	Non- white
<u>Total of ten leading causes</u>	<u>19,107</u>	<u>11,632</u>	<u>7,475</u>	<u>80.1</u>	<u>83.8</u>	<u>74.8</u>
1. Diseases of heart (390-398, 402, 404-429)	7,886	5,142	2,744	33.0	37.1	27.5
2. Malignant neoplasms (140-209)	3,295	2,146	1,149	13.8	15.5	11.5
3. Cerebrovascular disease (430-438)	3,058	1,681	1,377	12.8	12.1	13.8
4. Accidents (800-949)	2,007	1,179	828	8.4	8.5	8.3
5. Certain causes of mortality in early infancy (760-778)	766	329	437	3.2	2.4	4.4
6. Influenza and pneumonia (470-486)	726	385	341	3.0	2.8	3.4
7. Diabetes mellitus (250)	477	223	254	2.0	1.6	2.5
8. Arteriosclerosis (440)	331	232	99	1.4	1.7	1.0
9. Chronic and unqualified bronchitis, emphysema, asthma (490-493)	296	239	57	1.2	1.7	0.6
10. Homicide (960-978)	265	76	189	1.1	0.5	1.9

Leading Causes by Age

Table D4 shows the five leading causes of death for each age group except infants (see infant mortality section). Comparison of these lists revealed that, although a number of the same causes appeared repeatedly, no two lists were alike and the relative significance of the causes that were on more than one list varied considerably by age. For example, accidents appeared on every list and, although they caused more than half of the deaths among persons 5-14 and 15-24 years of age, the death rates in these younger groups were less than half the rate for persons 65 and over; in fact the rate for children 5-14 years old was less than one-fifth the rate for the elderly. Also on every list was malignant neoplasms, and heart diseases was on all but that for the group 1-4 years old; however, the death rates among children and young adults were only a tiny fraction of the rates for older persons. The list for persons 15-24 years of age was unusual in that three of the five leading causes were external causes - accidents, homicide, and suicide - and these three accounted for 75 percent of the total in that group.

Ill-defined and Unknown Causes

The listing of leading causes of death in Tables D3 and D4 was influenced to some degree by the fact that a sizeable number of the death certificates filed in Mississippi had no satisfactory statement of cause. If the cause could have been obtained for the 1,716 deaths in this category (7.3 percent of the total), the ranking and even the causes listed might have been different. The problem was concentrated in the nonwhite population, where the percentage was 12.2 in contrast to only 3.8 for deaths among white persons, and in certain counties. Twenty percent or more of the records filed in three counties - Amite with a high of 34 percent, Benton, and Yazoo - either were without a cause or had only a vague symptom or other ill-defined condition shown. The percentage was in the range 15-19 in six other counties; these were Carroll, DeSoto, Hancock, Madison, Marion, and Noxubee. Another category of deaths which might also be considered as lacking a satisfactory statement of cause is the infant deaths with no reason mentioned except immaturity; if the 207 deaths in this classification were added to the total previously stated, the grand total with ill-defined and unknown causes would be 1,923 and the percentage 8.2.

Vital Statistics
Mississippi - 1969

Table D4. Deaths, percent of total deaths, and death rates for five leading causes, by age and cause

Age	Cause and International Classification Number	Number	Percent	Rate*
1-4	<u>All causes</u>	<u>237</u>	<u>100.0</u>	<u>1.4</u>
	1. Accidents (800-949)	101	42.6	61.0
	2. Influenza and pneumonia (470-486)	27	11.4	16.3
	3. Congenital anomalies (740-759)	20	8.4	12.1
	4. Malignant neoplasms (140-209)	13	5.5	7.8
	5. Meningitis (320)	9	3.8	5.4
5-14	<u>All causes</u>	<u>306</u>	<u>100.0</u>	<u>0.6</u>
	1. Accidents (800-949)	178	58.2	36.3
	2. Malignant neoplasms (140-209)	24	7.8	4.9
	3. Influenza and pneumonia (470-486)	14	4.6	2.8
	4. Congenital anomalies (740-759)	10	3.3	2.0
	5. Diseases of heart (390-398, 402, 404-429)	7	2.3	1.4
15-24	<u>All causes</u>	<u>665</u>	<u>100.0</u>	<u>1.6</u>
	1. Accidents (800-949)	413	62.1	101.6
	2. Homicide (960-978)	57	8.6	14.0
	3. Malignant neoplasms (140-209)	30	4.5	7.4
	4. Suicide (950-959)	27	4.1	6.6
	5. Diseases of heart (390-398, 402, 404-429)	16	2.4	3.9
25-44	<u>All causes</u>	<u>1,528</u>	<u>100.0</u>	<u>3.3</u>
	1. Accidents (800-949)	411	26.9	88.2
	2. Diseases of heart (390-398, 402, 404-429)	224	14.6	48.0
	3. Malignant neoplasms (140-209)	200	13.1	42.9
	4. Homicide (960-978)	122	8.0	26.2
	5. Cerebrovascular disease (430-438)	85	5.6	18.2
45-64	<u>All causes</u>	<u>5,871</u>	<u>100.0</u>	<u>13.9</u>
	1. Diseases of heart (390-398, 402, 404-429)	2,010	34.2	476.0
	2. Malignant neoplasms (140-209)	1,221	20.8	289.2
	3. Cerebrovascular disease (430-438)	552	9.4	130.7
	4. Accidents (800-949)	390	6.6	92.4
	5. Diabetes mellitus (250)	152	2.6	36.0
65 and over	<u>All causes</u>	<u>13,733</u>	<u>100.0</u>	<u>61.8</u>
	1. Diseases of heart (390-398, 402, 404-429)	5,607	40.8	2,522.0
	2. Cerebrovascular disease (430-438)	2,408	17.5	1,083.1
	3. Malignant neoplasms (140-209)	1,805	13.1	811.9
	4. Accidents (800-949)	454	3.3	204.2
	5. Influenza and pneumonia (470-486)	402	2.9	180.8

* Deaths from all causes per 1,000 population in specified group and deaths from specific causes per 100,000 population in specified group.

Vital Statistics
Mississippi - 1969

Specific Causes

Heart disease, ranking first in the state and nation, has been at the top of the list in Mississippi in every year from 1925 to the present except 1929. Almost exactly one-third of the deaths in the state in 1969 were due to this cause, but the count of 7,886 was a little lower than the record high set in 1968. The death rate of 369.0 per 100,000 population for whites was higher than the 333.2 calculated for nonwhites; however, the disparity between the two was not extreme. Ischemic heart disease was responsible for 86 percent of all heart disease deaths and, within this category, acute myocardial infarction alone took 4,068 lives.

Malignant neoplasms caused a record number of deaths in 1969 although the total of 3,295 was only slightly higher than the figure for the previous year. Cancer was the second leading cause, but its death toll was less than half that of heart disease. A higher cancer death rate for whites (154.0 per 100,000 population) than for nonwhites (139.5) was reported; but, as was the case for heart disease deaths, the difference between these race-specific rates was not of great import. It was also noted that death rates for males of both races were larger than those for females. As reported in the section on leading causes of death, cancer ranked among the top five in every age group except infants; however, the deaths among persons under 45 years of age amounted to only 8 percent of the total from this cause. The three most common primary sites of fatal cancers were the digestive organs and peritoneum (866 deaths), the respiratory system (669), and the genital organs (509); together they accounted for 62 percent of the total. As may be seen in Table D5, the highest of the death rates specific for sex and race was that for white males with cancer of the respiratory system even though the digestive organs and peritoneum ranked first in the other groups. Second-ranking sites for each group were: digestive organs and peritoneum for white males, respiratory system for nonwhite males, breast for white females, and genital organs for nonwhite females.

Table D5. Deaths and death rates for malignant neoplasms, by primary site, sex, and race

Primary Site (ranked according to number of deaths among total population) and International Classification Number	Number				Rate*			
	Male		Female		Male		Female	
	White	Non-white	White	Non-white	White	Non-white	White	Non-white
Total	1,214	622	932	527	177.6	159.3	131.4	121.7
1. Digestive organs and peritoneum (150-159)	274	189	257	146	40.1	48.4	36.2	33.7
2. Respiratory system (160-163)	414	139	78	38	60.5	35.6	11.0	8.8
3. Genital organs (180-187)	117	136	132	124	17.1	34.8	18.6	28.6
4. Lymphatic and hematopoietic tissue (200-209)	152	52	101	33	22.2	13.3	14.2	7.6
5. Breast (174)	1	0	174	77	0.1	-	24.5	17.8
6. Urinary organs (188, 189)	52	21	32	15	7.6	5.4	4.5	3.5
7. Skin (172, 173)	34	4	32	6	5.0	1.0	4.5	1.4
8. Brain (191)	30	6	30	2	4.4	1.5	4.2	0.5
9. Buccal cavity and pharynx (140-149)	39	8	11	6	5.7	2.0	1.6	1.4
10. Bone and connective tissue (170, 171)	15	5	13	3	2.2	1.3	1.8	0.7
- Other and unspecified sites (190, 192-199)	86	62	72	77	12.6	15.9	10.1	17.8

* Deaths per 100,000 population in specified group.

Vital Statistics
Mississippi - 1969

Cerebrovascular disease followed relatively closely after malignant neoplasms as the third leading cause with a count of 3,058. This was a somewhat smaller number than was reported for 1968, but study of the figures for all of the 1960's showed no indication of a trend in either direction. Excessive mortality from this cause among nonwhites has been demonstrated over many years, and in 1969 the death rate of 167.2 per 100,000 population for nonwhites was 39 percent higher than that of 120.6 for whites. Among nonwhites, and also among females, cerebrovascular disease was still the second leading cause of death.

Chronic respiratory diseases are significant causes of disability and death in this country. Mississippi mortality figures for the most important of these diseases are listed in Table D6 to show the trend of the past twenty years. Respiratory tuberculosis, which in earlier times had caused over 2,600 deaths in a single year and 552 in 1949, caused only 47 deaths in 1969. On the other hand, the death toll from malignant neoplasms of the respiratory system was over four times greater in 1969 than in 1949 and for a number of years has been responsible for more deaths than all of the other diseases in Table D6 together. The most remarkable increase, however, occurred in emphysema deaths which rose from 6 to 206 in the twenty-year period and in 1969 ranked second among the chronic respiratory diseases. Despite the fact that two revisions of the system of classifying causes of death occurred during the last twenty years, asthma was the only one of these conditions for which the changes were important enough to affect the comparability of the numbers. For this reason, figures adjusted to compensate for the changes in this category are also presented in Table D6. However, comparison of the adjusted numbers did not indicate any trend, either up or down, since 1949. The bronchitis death total for 1969 was higher than in any other year in the period under consideration, but, for this disease also, there was no established trend.

Table D6. Deaths from major chronic respiratory diseases, 1949, 1954, 1959, 1964, 1969

Cause and International Classification Number	1949	1954	1959	1964	1969
Tuberculosis of respiratory system (010-012)	552	210	142	96	47
Malignant neoplasms of respiratory system (160-163)	160	227	339	447	669
Malignant neoplasms of trachea, bronchus, and lung (162)	140	206	308	418	637
Chronic and unqualified bronchitis (490, 491)	22	27	34	32	62
Emphysema (492)	6	26	79	123	206
Asthma (493)	55	47	52	51	28
(Adjusted figures*)	(31)	(26)	(36)	(35)	

* Adjusted according to comparability ratios prepared by the National Center for Health Statistics in order to compensate for breaks in comparability caused by revision of the cause-of-death classification system in 1958 and 1968.

Influenza and pneumonia together caused 726 deaths and ranked sixth on the list of leading causes. The total count was 99 less than the figure for 1968 but larger than that for 1967. These diseases, as usual, were relatively more serious among nonwhites than among whites; death rates were 41.4 per 100,000 population and 27.6 respectively. The majority of the deaths - 642 - were due to pneumonia alone, but 56 were caused by influenza with pneumonia and 28 by influenza without mention of pneumonia.

Accidents, fourth among the major causes of death and first among persons 1-44

Vital Statistics
Mississippi - 1969

years of age, took an unprecedented number of lives in 1969. For deaths occurring in the state the count was 2,010, for deaths to residents it was 2,007, and the death rate was 90.5 per 100,000 population. Each of these figures was far larger than the previous records of 1,861, 1,890, and 80.0 set two years earlier. The year 1969 was marked by two natural disasters and another upswing in deaths due to motor vehicle accidents. Hurricane Camille struck the Gulf Coast in August causing over a hundred deaths and a tornado went through the Hazlehurst area in January with 32 deaths reported. A total of 132 death certificates were filed for hurricane victims including 12 for nonresidents and 4 for unidentified persons.

Information regarding the types of accidents which cause the most fatalities is presented in Table D7. These five types were responsible for 77 percent of the total, but the list for 1969 was different from those for previous years in that it included cataclysm; Hurricane Camille caused this category to appear on the list, thereby excluding firearm deaths which usually is among the top five. Motor vehicle accidents ranked first with more deaths than the other four types together. Fatalities from accidents involving motor vehicles increased substantially to reach totals second only to those reported in 1967; the counts for 1969 were 920 deaths occurring in Mississippi and 903 to residents of the state. Although the death rate of 40.7 per 100,000 population was the highest ever recorded, the rate based on estimated motor vehicle miles traveled was not unusually high.

Table D7. Deaths and percent of total accidental deaths from five leading causes of accidental death, by cause and race

Type of Accident and International Classification Number	Number			Percent		
	Total	White	Non- white	Total	White	Non- white
<u>Total of five leading causes</u>	<u>1,539</u>	<u>911</u>	<u>628</u>	<u>76.7</u>	<u>77.3</u>	<u>75.8</u>
1. Motor vehicle (810-823)	903	586	317	45.0	49.7	38.3
2. Fires and flames (890-899)	202	41	161	10.1	3.5	19.4
3. Falls (880-887)	169	133	36	8.4	11.3	4.3
4. Cataclysm (908)	152	107	45	7.6	9.1	5.4
5. Drowning, except in water transport (910)	113	44	69	5.6	3.7	8.3

Further analysis of accidental deaths that occurred in Mississippi revealed that, for all age groups except children under 5 and persons 65 and over, the chief cause was motor vehicles. Among the elderly, falls ranked first with motor vehicles a close second, and fires and flames lead the list for children 1-4 years old with motor vehicles in second place for this group also. The leading cause for infants under one year of age was mechanical suffocation followed by fires and flames.

Suicides and homicides numbered 174 and 265 respectively among Mississippians in 1969. For the total population, self-inflicted injuries ranked twelfth among the leading causes of death and homicide was in tenth place. However, they were even more important for certain segments of the population. Suicide was the fourth ranking cause of death for young persons 15-24 years of age and eighth among white males. Homicide was second on the list for the population 15-24 years of age, fourth among those 25-44 years old, and seventh among nonwhite males. Table D8 gives comparative data on the incidence of these two events for each specific age-sex-race group. From these figures it can be easily seen that the

Vital Statistics
Mississippi - 1969

likelihood of suicide was greater for white males in the age groups 35 and over than for any other group and that the rates were especially high for white males 55 and over. In addition, the overall suicide rate of 16.4 per 100,000 population for white males was triple the highest rate among the other three sex-race groups. By contrast, nonwhite males suffered the greatest incidence of homicides and their rate of 38.7 was more than double the high suicide rate for white males. For nonwhite males 25-34 and 35-44 years of age the homicide rates were well over 100 and far higher than any other age-sex-race-specific homicide or suicide rates. Firearms and explosives were used to commit 82 percent of the suicides and 72 percent of the homicides.

Table D8. Suicide and homicide rates* by age, sex, and race

Age	Suicide					Homicide				
	Total	Male		Female		Total	Male		Female	
		White	Non-white	White	Non-white		White	Non-white	White	Non-white
<u>Total</u>	<u>7.8</u>	<u>16.4</u>	<u>5.4</u>	<u>4.4</u>	<u>2.3</u>	<u>12.0</u>	<u>8.3</u>	<u>38.7</u>	<u>2.7</u>	<u>8.8</u>
Under 15	0.1	-	-	-	0.6	0.4	0.5	-	0.5	0.6
15-24	6.6	11.0	7.9	3.3	3.7	14.0	6.3	46.1	4.9	9.9
25-34	8.2	14.0	16.2	3.4	-	28.4	12.9	129.5	4.6	35.7
35-44	15.2	28.2	3.7	11.0	5.4	23.7	20.5	107.6	2.4	16.2
45-54	13.7	28.1	14.0	5.0	2.7	17.8	13.4	73.7	3.8	13.7
55-64	16.3	35.4	9.8	8.3	2.8	11.8	7.7	42.7	4.2	8.5
65 and over	13.0	33.4	5.5	6.0	4.6	9.0	10.0	36.0	-	2.3

* Deaths per 100,000 population in specified group.

Injury undetermined whether accidentally or purposely inflicted was the classification for 41 deaths that were not due to natural causes but to external events about which information was inadequate for determining whether they were accidents, suicides, or homicides. This category was first used in 1968 when the reported total was 51. In 1969, 23 of these deaths resulted from injuries caused by firearms and explosives.

Maternal deaths are those stated to have resulted from deliveries and complications of pregnancy, childbirth, and the puerperium. Only 23 such deaths were recorded for Mississippi mothers in 1969 and the maternal death rate was 4.9 per 10,000 live births. Both of these figures were slightly higher than those for 1968 but were the second lowest on record for this state. The rate of 1.2 for white mothers was the smallest ever recorded for that race and considerably less than the rate of 8.6 for nonwhite mothers. Table D9, which contains rates specific for age as well as race, shows that no deaths occurred among the very young mothers or those 45 or over and that the rates increased with age in each 10-year group from 15 through 44 years of age. Two of the maternal deaths in 1969 resulted from abortions and one from an ectopic pregnancy; the death certificates did not indicate whether the abortions were spontaneous or induced. Toxemias, hemorrhage, and sepsis or infection, which for many years were the chief causes of maternal death, took 11, 2, and 4 lives respectively.

Vital Statistics
Mississippi - 1969

Table D9. Maternal deaths and maternal death rates, by age and race

Age	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>23</u>	<u>3</u>	<u>20</u>	<u>4.9</u>	<u>1.2</u>	<u>8.6</u>
10-14	0	0	0	-	-	-
15-24	11	1	10	3.9	0.7	7.2
25-34	8	2	6	5.4	2.4	9.1
35-44	4	0	4	11.8	-	18.2
45-54	0	0	0	-	-	-

* Deaths per 10,000 live births to mothers in specified group.

Infant Mortality

Mississippi's infant death count dropped from a record low in 1968 to another record low of 1,513 in 1969. The same type of decline took place in the mortality rate which stood at 32.1 per 1,000 live births in 1969. This rate following the rate of 35.3 for both 1967 and 1968 indicated an end to the long period from 1945 through 1966 when the rates varied between 36.4 and 41.5 with no downward trend. The curves on the accompanying graph show that, during this time, the death rate for white babies continued to decline slowly but that for nonwhite infants actually increased until the last few years. In 1969 the rate for whites fell to a record low (22.1), and the rate of 42.6 for nonwhites was among the lowest ever recorded. However, provisional data for the United States revealed that Mississippi's rates were still excessive. The rate for this state remained the highest in the country and was 55 percent higher than the national rate. For whites, Mississippi's rate was 20 percent higher than the U.S. rate and for nonwhites it was 35 percent higher.

The number of infants who died in the first day of life was nearly as large as the number who died in the next 27 days, as Table D10 shows, and the total for the first 28 days (neonatal period) was more than double the total for the next 11 months. At each of the ages listed in the table the mortality rate for nonwhite children was higher than that for white infants, but the difference was greatest in the postneonatal period where the rate for nonwhites was over four times higher. Although the unusually large postneonatal death rate for nonwhites has, through the years, been an important component of Mississippi's excessive infant death rates, latest available national information (1968) shows that this state ranked considerably higher than any other in neonatal mortality. It is also of note that this state's high death rates for babies result from excessive mortality not just among nonwhites but also among whites. In 1968 Mississippi had the second highest white infant death rate in the country and the second highest white neonatal death rate.

Vital Statistics
Mississippi - 1969

Infant Mortality Rates by Race
1920-1969




Table D10. Infant deaths and infant death rates, by age and race

Age	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
<u>Total</u>	<u>1,513</u>	<u>530</u>	<u>983</u>	<u>32.1</u>	<u>22.1</u>	<u>42.6</u>
Under one day	531	226	305	11.3	9.4	13.2
One to 28 days	535	215	320	11.4	8.9	13.8
28 days to one year	447	89	358	9.5	3.7	15.5

* Deaths per 1,000 live births.

In Table D11 the most important cause-of-death categories for infants are listed but not ranked as no standards for grouping and ranking infant causes has yet been developed for the new classification system which went into effect in 1968. Among the categories listed, the one with the largest number of deaths reported was anoxic and hypoxic conditions not elsewhere classifiable with a total of 326; hyaline membrane disease (97 deaths), respiratory distress syndrome (87 deaths), and unspecified asphyxia of the

Vital Statistics
Mississippi - 1969

newborn (127 deaths) accounted for all but a few of the deaths in this category. Complications of the mother during pregnancy and childbirth was another group of causes responsible for a sizeable amount of mortality; among these 150 deaths were 54 due to complications of a multiple pregnancy and 33 that resulted from premature separation of the placenta. Congenital anomalies also caused 150 deaths and influenza and pneumonia, 147. Study of the types of accidents responsible for deaths among infants showed that the major type was mechanical suffocation which caused 19 out of the 54 fatalities.

Table D11. Infant deaths and infant death rates
for significant causes of infant death, by cause and race

Cause of Death and International Classification Number	Number			Rate*		
	Total	White	Non- white	Total	White	Non- white
<u>All causes</u>	<u>1,513</u>	<u>530</u>	<u>983</u>	<u>32.1</u>	<u>22.1</u>	<u>42.6</u>
Diarrheal disease (009)	71	9	62	1.5	0.4	2.7
Influenza and pneumonia (470-486)	147	35	112	3.1	1.4	4.8
Congenital anomalies (740-759)	150	85	65	3.2	3.5	2.8
Complications of pregnancy and child- birth, including conditions of placenta and umbilical cord (760-771)	150	64	86	3.2	2.7	3.7
Anoxic and hypoxic conditions not elsewhere classifiable (776)	326	162	164	6.9	6.7	7.1
Immaturity, unqualified (777)	212	71	141	4.5	3.0	6.1
Accidents (800-949)	54	13	41	1.1	0.5	1.8

* Deaths per 1,000 live births.

For all of the causes of death listed in Table D11 except congenital anomalies, the mortality rate for nonwhite infants was higher than that for whites and, in fact, was several times higher for diarrheal disease, influenza and pneumonia, and accidents. It was also noted that 70 percent or more of the deaths from each of these three causes occurred in the postneonatal period. In contrast, 70 percent of the deaths caused by congenital anomalies occurred in the neonatal period and nearly all of the deaths resulting from complications of pregnancy and childbirth and from the anoxic and hypoxic conditions occurred among babies less than a week old.

Immaturity is a vague term as a cause of death and is not assigned unless it is the only condition on a death certificate. However, the number of such certificates in 1969 was 212 which was the second largest total for the categories listed in Table D11. Other certificates lacking a satisfactory statement of cause numbered 191, making a total of 403 (27 percent) for which the cause of death was ill-defined or unknown. Analysis of mortality among nonwhite infants or of deaths in the first day of life is greatly hampered by the fact that the cause was not ascertained for nearly a third of the deaths in each of these categories.

MARRIAGES (See Tables 9, 17)

Trend and Race

In 1969, for the tenth consecutive year, the number of marriages increased. The

Vital Statistics
Mississippi - 1969

upward climb in recent years is related to the tremendous growth of the population of marriageable age. Between the censuses of 1960 and 1970 the total state population increased by only 1.8 percent, whereas the population 16-25 years old (the ten-year span in which the most marriages occurred in 1969) went up 22.3 percent. During most of the 1960's, however, the year-to-year increase in marriages was relatively small and this was again true in 1969. The final figure was 25,956 which was only 601 more than in the previous year. Upward, though slightly irregular, trends have been experienced by both whites and nonwhites with counts at the end of the decade 17,582 and 8,374 respectively.

Month

The popularity of the month of June for weddings can easily be seen in Table E1 which gives the number of marriages and average per day for each month. These figures show that June ranked first, August second, and December third. According to the National Center for Health Statistics, Saturday is the most popular day of the week for weddings; and, therefore, monthly variation depends to some extent on whether there are four or five Saturdays in a month. Although June always ranks first, the order in which August and December ranked during the 1960's appeared to be correlated (with some exceptions) with the number of Saturdays in each.

Table E1. Marriages and average per day, by month

Month	Marriages	Average per Day	Month	Marriages	Average per Day
<u>Total</u>	<u>25,956</u>	<u>71</u>			
January	1,730	56	July	2,259	73
February	1,711	61	August	3,035	98
March	1,868	60	September	1,806	60
April	1,691	56	October	1,756	57
May	2,297	74	November	1,986	66
June	3,282	109	December	2,535	82

Officiant

Ministers officiated at 72 percent of the weddings, and most of the others were conducted by justices of the peace. From 1961 through 1969 the proportion of religious ceremonies remained almost constant with the percentage either 71 or 72 each year.

Age of Bride and Groom

Brides in their teens or younger numbered 10,863 which was 42 percent of the total married in the state. (See Table E2) Corresponding figures on grooms were 4,906 and 19 percent. The youngest bride was only 12 and the youngest groom, 15. In addition, there were 15 brides 13 years of age and 110 who were age 14. Marriage partners of advanced ages were also listed; a total of 186 were reported to be 75 or more years old. Brides in this category numbered 36 with the oldest age 87, and the oldest groom among the 150 in this group was 88. Study of detailed counts by age revealed

Vital Statistics
Mississippi - 1969

that there were more brides 18 years old than of any other single year of age and more grooms who were 21 than any other age. This was true for both whites and nonwhites. The percentage distributions given in Table E2 show only minor differences between the races for any age group.

Table E2. Brides and grooms and percentage distribution, by age and race

Age	Number						Percent					
	Bride			Groom			Bride			Groom		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Total	25,956	17,582	8,374	25,956	17,583	8,373	100.0	100.0	100.0	100.0	100.0	100.0
Under 20	10,863	7,460	3,403	4,906	3,390	1,516	41.8	42.4	40.6	18.9	19.3	18.1
12	1	1	0	0	0	0	*	*	-	-	-	-
13	15	8	7	0	0	0	*	*	0.1	-	-	-
14	110	56	54	0	0	0	0.4	0.3	0.6	-	-	-
15	827	566	261	13	11	2	3.2	3.2	3.1	*	0.1	*
16	1,475	980	495	76	59	17	5.7	5.6	5.9	0.3	0.3	0.2
17	2,308	1,587	721	736	560	176	8.9	9.0	8.6	2.8	3.2	2.1
18	3,252	2,271	981	1,693	1,172	521	12.5	12.9	11.7	6.5	6.7	6.2
19	2,875	1,991	884	2,388	1,588	800	11.1	11.3	10.6	9.2	9.0	9.6
20-24	8,055	5,519	2,536	11,407	7,856	3,551	31.0	31.4	30.3	43.9	44.7	42.4
25-34	3,314	2,158	1,156	4,886	3,295	1,591	12.8	12.3	13.8	18.8	18.7	19.0
35-44	1,719	1,169	550	1,938	1,348	590	6.6	6.6	6.6	7.5	7.7	7.0
45-54	1,064	695	369	1,275	838	437	4.1	4.0	4.4	4.9	4.8	5.2
55-64	658	415	243	907	518	389	2.5	2.4	2.9	3.5	2.9	4.6
65-74	240	146	94	475	248	227	0.9	0.8	1.1	1.8	1.4	2.7
75 & over	36	17	19	150	86	64	0.1	0.1	0.2	0.6	0.5	0.8
Unknown	7	3	4	12	4	8	*	*	*	*	*	0.1

* Less than 0.06 percent.

Median ages of all brides and grooms and of those marrying for the first time are presented in Table E3 for each race group. The medians for grooms were higher than those for brides and those for nonwhites were higher than corresponding figures for whites of the same sex. These relationships held for first marriages as well as for all marriages. Although a cross tabulation of ages of brides and grooms indicated that many persons chose mates whose age was not greatly different from their own, there were some extreme differences, the largest of which was 52 years for a bride of 21 and a groom of 73.

Table E3. Median age of bride and groom, for all marriages and first marriages, by race

All and First Marriages	Bride		Groom	
	White	Non-white	White	Non-white
All marriages	20.8	21.0	22.7	23.2
First marriages	19.5	20.0	22.0	22.5

Vital Statistics
Mississippi - 1969

Marriage Order and Previous Marital Status

Both the bride and the groom married for the first time in 66 percent of the weddings in 1969; but, as Table E4 shows, the total proportions of each sex not previously married were 73 percent for brides and the same for grooms. Among those who had previously been married, the great majority were entering their second marriage. However, persons embarking on a marriage of fifth or higher order numbered 98, and two of the grooms stated that they were being married for the eighth time. Divorced persons far outnumbered the widowed among those being remarried.

Table E4. Brides and grooms and percentage distribution,
by marriage order and previous marital status

Marriage Order and Previous Marital Status	Number		Percent	
	Bride	Groom	Bride	Groom
<u>Total</u>	<u>25,956</u>	<u>25,956</u>	<u>100.0</u>	<u>100.0</u>
First marriage	19,059	19,042	73.4	73.4
<u>Remarriage</u>	<u>6,852</u>	<u>6,875</u>	<u>26.4</u>	<u>26.5</u>
Second	5,213	5,201	20.1	20.0
<u>Third or higher</u>	<u>1,639</u>	<u>1,674</u>	<u>6.3</u>	<u>6.4</u>
Third	1,347	1,316	5.2	5.1
Fourth	246	306	0.9	1.2
Fifth	38	37	0.1	0.1
Sixth	8	10	**	**
Seventh	0	3	-	**
Eighth	0	2	-	**
Widowed	1,627	1,357	6.3	5.2
Divorced*	5,135	5,424	19.8	20.9
Unknown whether widowed or divorced	90	94	0.3	0.4
Unknown whether any previous marriage	45	39	0.2	0.2

* Includes those whose last marriage was annulled.

** Less than 0.06 percent.

Nonresident Brides and Grooms

A total of 3,921 brides or 15 percent of those married in this state were residents of other places, but the number of Mississippi women who went outside the state to get married in 1969 is not known since there is no interchange of marriage records among the states. The neighboring states of Alabama, Arkansas, Louisiana, and Tennessee accounted for 79 percent of the nonresident brides, but they also came from all over the nation - from 45 states and the District of Columbia - and 11 were from foreign countries. It was noted that, as usual, most of the out-of-state brides obtained their marriage licenses in counties along the state boundary lines; a total of 75 percent of the licenses were issued in these border counties with two counties predominating. DeSoto County

Vital Statistics
Mississippi - 1969

(adjacent to Memphis, Tennessee, and long a place to which many out-of-state brides come) and Wilkinson County (on the Louisiana line) issued 1,213 or 31 percent of the total to brides from other places. In addition to the nonresident brides reported, there were 5,146 grooms from other places, and in one out of every eight weddings (12 percent) neither the bride nor the groom was a Mississippi resident.

Out-of-State Use of Mississippi Licenses

Marriage licenses issued in Mississippi are sometimes used for ceremonies performed in other states. Because these weddings did not occur in the state and their legality is questionable, they are not included in the official statistics on Mississippi marriages. However, special tallies of the data from these records are prepared each year. In 1969, 85 reports of such marriages were filed and this was the third lowest count in the thirteen years this information has been collected. The fact that one or both partners lived out of state apparently was the reason many of the weddings were held in other places; 72 percent occurred in the state where either the bride or the groom or both resided. In 1969 all the cases which did not involve persons from the state where the marriage took place involved brides and/or grooms from Mississippi. Louisiana was the site of over half of the out-of-state weddings, and all but 7 occurred in the four states which border on Mississippi. The licenses for 64 percent were issued by Mississippi counties just over the line from the state where the marriage took place.

DIVORCES (See Tables 9, 10, 17)

Trend and Race

The divorce count in Mississippi rose again in 1969. It began going up in the latter part of the previous decade and continued through the 1960's with the total for 1969 reaching 7,506, the second highest figure ever recorded (1946 was the peak year). The upward trend has been more pronounced among the white than among the total population, and in the last two years alone the increase among the whites amounted to 19 percent. Indications are that the nonwhites, after a number of years of irregular variation, have recently entered a period of steady increase also.

Cause and Complainant

Cruel and inhuman treatment was the cause of divorce listed most frequently, accounting for 64 percent of the total. Desertion ranked second with 27 percent, and all other reasons added up to only 9 percent. Although cruel and inhuman treatment was the cause to which 74 percent of the divorces among white couples were attributed, it was listed for only one-third of those among nonwhites, and desertion accounted for 59 percent in this group. Data on complainant showed that, in cases involving whites, 72 percent were filed by the wife, whereas, she was the complainant in only 52 percent among nonwhites.

Minor Children

Forty-three percent of the couples granted divorces reported that they had no minor children. However, the couples who did have children in this category listed a total of 7,506 or an average of 2.2 per family. Three families had 13 children each and the number with five or more children was 279. A higher proportion of the white than of

Vital Statistics
Mississippi - 1969

the nonwhite couples had minor children, but the nonwhites had more children per family.

Duration of Marriage

Couples who were granted divorces in 1969 had had marriages with durations from less than one year up to 63 years. One out of every twelve or 8 percent had not been married an entire year and 43 percent did not remain together as long as five years. In contrast, 15 percent had been married twenty or more years and 19 of these had been married fifty or more years. Calculation of median durations by race revealed a striking contrast. The median of 11.9 years for nonwhites was more than double the median of 5.1 years for white couples.

Additional Information Needed

Mississippi's statistical record of divorce is a form which provides only limited information because of the omission of a number of pertinent items which many other states collect and also because of the lack of clarity regarding some of the items which are on the form. No data at all are available concerning place of marriage, place of residence of the couple at time of separation, their ages at marriage and at divorce, previous marriages, and socio-economic factors. Information which is presently tabulated on minor children, cause of divorce, and complainant probably is erroneous because of lack of definitions and instructions regarding how such facts should be entered on the form and also because of the poor arrangement of the form. These deficiencies seriously hamper meaningful study of the divorce problem on a state level as well as on a national level. Since Mississippi's record does not contain even what is considered minimum information, this state is not eligible to become part of the National Divorce Registration Area. Revision of the statistical record of divorce would be of great help in providing material for analysis of this serious social problem.

TABLE 1
 CENSUS POPULATION
 MISSISSIPPI
 April 1, 1970

By County, by Race

County	Total	White	Nonwhite
<u>STATE</u>	<u>2,216,912</u>	<u>1,393,283</u>	<u>823,629</u>
Adams	37,293	19,366	17,927
Alcorn	27,179	23,951	3,228
Amite	13,763	6,814	6,949
Attala	19,570	11,632	7,938
Benton	7,505	4,350	3,155
Bolivar	49,409	18,750	30,659
Calhoun	14,623	10,807	3,816
Carroll	9,397	4,615	4,782
Chickasaw	16,805	10,819	5,986
Choctaw	8,440	6,069	2,371
Claiborne	10,086	2,536	7,550
Clarke	15,049	9,631	5,418
Clay	18,840	9,517	9,323
Coahoma	40,447	14,232	26,215
Copiah	24,749	12,298	12,451
Covington	14,002	9,426	4,576
DeSoto	35,885	23,235	12,650
Forrest	57,849	43,524	14,325
Franklin	8,011	4,897	3,114
George	12,459	11,007	1,452
Greene	8,545	6,662	1,883
Grenada	19,854	11,154	8,700
Hancock	17,387	14,894	2,493
Harrison	134,582	111,061	23,521
Hinds	214,973	130,592	84,381
Holmes	23,120	7,345	15,775
Humphreys	14,601	5,089	9,512
Issaquena	2,737	1,033	1,704
Itawamba	16,847	15,882	965
Jackson	87,975	73,547	14,428
Jasper	15,994	8,564	7,430
Jefferson	9,295	2,296	6,999
Jeff Davis	12,936	6,435	6,501
Jones	56,357	42,403	13,954
Kemper	10,233	4,488	5,745
Lafayette	24,181	17,313	6,868

Table 1 - Page 2
 Census Population - Mississippi - 1970

County	Total	White	Nonwhite
Lamar	15,209	13,187	2,022
Lauderdale	67,087	46,186	20,901
Lawrence	11,137	7,553	3,584
Leake	17,085	10,452	6,633
Lee	46,148	36,571	9,577
Leflore	42,111	17,550	24,561
Lincoln	26,198	18,138	8,060
Lowndes	49,700	33,336	16,364
Madison	29,737	11,148	18,589
Marion	22,871	15,739	7,132
Marshall	24,027	9,101	14,926
Monroe	34,043	23,650	10,393
Montgomery	12,918	7,121	5,797
Neshoba	20,802	15,091	5,711
Newton	18,983	13,352	5,631
Noxubee	14,288	4,844	9,444
Oktibbeha	28,752	18,569	10,183
Panola	26,829	13,061	13,768
Pearl River	27,802	22,688	5,114
Perry	9,065	6,674	2,391
Pike	31,756	17,903	13,853
Pontotoc	17,363	14,254	3,109
Prentiss	20,133	17,776	2,357
Quitman	15,888	6,687	9,201
Rankin	43,933	31,529	12,404
Scott	21,369	14,254	7,115
Sharkey	8,937	3,125	5,812
Simpson	19,947	13,678	6,269
Smith	13,561	10,609	2,952
Stone	8,101	6,220	1,881
Sunflower	37,047	13,619	23,428
Tallahatchie	19,338	7,657	11,681
Tate	18,544	9,777	8,767
Tippah	15,852	13,260	2,592
Tishomingo	14,940	14,268	672
Tunica	11,854	3,225	8,629
Union	19,096	16,145	2,951
Walthall	12,500	7,403	5,097
Warren	44,981	26,474	18,507
Washington	70,581	31,803	38,778
Wayne	16,650	11,169	5,481
Webster	10,047	7,789	2,258
Wilkinson	11,099	3,588	7,511
Winston	18,406	11,065	7,341
Yalobusha	11,915	7,071	4,844
Yazoo	27,304	12,690	14,614

TABLE 2
LIVE BIRTHS
MISSISSIPPI
1969

By Place of Occurrence, and by Place of Residence of Mother, by Race, with Rates

County	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
<u>STATE</u>	<u>47,138</u>	<u>24,181</u>	<u>22,957</u>	<u>47,120</u>	<u>24,025</u>	<u>23,095</u>	<u>21.2</u>	<u>17.2</u>	<u>28.0</u>
Adams	1,565	753	812	890	393	497	23.9	20.3	27.7
Alcorn	329	265	64	431	365	66	15.8	15.2	20.4
Amite	80	0	80	275	87	188	20.0	12.8	27.0
Attala	366	183	183	394	178	216	20.1	15.3	27.2
Benton	42	1	41	132	64	68	17.6	14.7	21.6
Bolivar	1,741	368	1,373	1,370	339	1,031	27.7	18.1	33.6
Calhoun	179	69	110	276	170	106	18.9	15.7	27.8
Carroll	42	2	40	154	61	93	16.4	13.2	19.4
Chickasaw	286	111	175	358	171	187	21.3	15.8	31.2
Choctaw	172	60	112	157	75	82	18.6	12.4	34.6
Claiborne	83	13	70	202	37	165	20.0	14.6	21.8
Clarke	38	18	20	284	137	147	18.9	14.2	27.1
Clay	237	89	148	316	130	186	16.8	13.6	20.0
Coahoma	855	338	517	943	243	700	23.3	17.1	26.7
Copiah	581	173	408	478	148	330	19.3	12.0	26.5
Covington	163	123	40	276	160	116	19.7	17.0	25.3
DeSoto	209	14	195	781	429	352	21.8	18.5	27.8
Forrest	1,461	1,120	341	1,169	778	391	20.2	17.9	27.3
Franklin	15	2	13	123	56	67	15.4	11.4	21.5
George	293	254	39	279	238	41	22.4	21.6	28.2
Greene	75	36	39	186	129	57	21.8	19.4	30.3
Grenada	426	163	263	446	187	259	22.5	16.8	29.8
Hancock	77	53	24	268	227	41	15.4	15.2	16.4
Harrison	3,548	2,837	711	3,171	2,484	687	23.6	22.4	29.2
Hinds	5,904	2,934	2,970	4,480	2,132	2,348	20.8	16.3	27.8
Holmes	484	80	404	512	104	408	22.1	14.2	25.9
Humphreys	374	46	328	371	83	288	25.4	16.3	30.3
Issaquena	12	0	12	56	18	38	20.5	17.4	22.3
Itawamba	33	25	8	244	224	20	14.5	14.1	20.7
Jackson	1,743	1,287	456	2,060	1,582	478	23.4	21.5	33.1
Jasper	189	131	58	330	124	206	20.6	14.5	27.7
Jefferson	56	7	49	160	21	139	17.2	9.1	19.8
Jeff Davis	166	86	80	292	112	180	22.6	17.4	27.7
Jones	2,003	850	1,153	990	659	331	17.6	15.5	23.7
Kemper	68	10	58	221	57	164	21.6	12.7	28.5
Lafayette	545	340	205	466	285	181	19.3	16.5	26.4

Table 2 - Page 2
Live Births - Mississippi - 1969

County	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Lamar	86	59	27	297	246	51	19.5	18.6	25.2
Lauderdale	2,133	1,138	995	1,525	859	666	22.7	18.6	31.9
Lawrence	94	51	43	200	114	86	18.0	15.1	24.0
Leake	185	112	73	304	154	150	17.8	14.7	22.6
Lee	1,355	1,088	267	897	658	239	19.4	18.0	25.0
Leflore	1,107	398	709	1,002	303	699	23.8	17.3	28.4
Lincoln	579	354	225	532	327	205	20.3	18.0	25.4
Lowndes	1,165	708	457	1,008	502	506	20.3	15.0	30.9
Madison	491	106	385	768	169	599	25.8	15.2	32.2
Marion	377	253	124	485	283	202	21.2	18.0	28.3
Marshall	292	22	270	555	142	413	23.1	15.6	27.7
Monroe	932	626	306	725	444	281	21.3	18.8	27.0
Montgomery	289	127	162	230	96	134	17.8	13.5	23.1
Neshoba	397	238	159	396	229	167	19.0	15.2	29.2
Newton	320	201	119	363	201	162	19.1	15.0	28.8
Noxubee	377	73	304	345	68	277	24.1	14.0	29.3
Oktibbeha	649	314	335	587	318	269	20.4	17.1	26.4
Panola	692	217	475	585	179	406	21.8	13.7	29.5
Pearl River	592	509	83	556	439	117	20.0	19.3	22.9
Perry	122	86	36	192	130	62	21.2	19.5	25.9
Pike	630	325	305	580	256	324	18.3	14.3	23.4
Pontotoc	287	204	83	290	215	75	16.7	15.1	24.1
Prentiss	348	300	48	329	279	50	16.3	15.7	21.2
Quitman	483	115	368	442	130	312	27.8	19.4	33.9
Rankin	99	45	54	802	564	238	18.2	17.9	19.2
Scott	408	272	136	461	244	217	21.6	17.1	30.5
Sharkey	78	9	69	229	43	186	25.6	13.8	32.0
Simpson	303	183	120	428	224	204	21.4	16.4	32.5
Smith	8	0	8	257	152	105	19.0	14.3	35.6
Stone	137	88	49	165	90	75	20.4	14.5	39.9
Sunflower	735	135	600	954	208	746	25.8	15.3	31.8
Tallahatchie	454	78	376	503	99	404	26.0	12.9	34.6
Tate	137	0	137	405	157	248	21.8	16.0	28.3
Tippah	352	271	81	296	230	66	18.7	17.3	25.5
Tishomingo	169	154	15	239	225	14	16.0	15.8	20.8
Tunica	340	18	322	370	53	317	31.2	16.4	36.7
Union	301	220	81	336	264	72	17.6	16.4	24.4
Walthall	299	149	150	248	98	150	19.8	13.2	29.4
Warren	1,378	605	773	917	498	419	20.4	18.8	22.6
Washington	1,918	703	1,215	1,819	613	1,206	25.8	19.3	31.1
Wayne	337	246	91	345	191	154	20.7	17.1	28.1
Webster	109	68	41	174	114	60	17.3	14.6	26.6
Wilkinson	175	71	104	197	40	157	17.7	11.1	20.9
Winston	321	155	166	384	170	214	20.9	15.4	29.2
Yalobusha	166	57	109	220	87	133	18.5	12.3	27.4
Yazoo	492	189	303	637	233	404	23.3	18.4	27.6
Out of state	1,680	1,259	421	1,698	1,415	283			

Table 2 - Page 3
Live Births - Mississippi - 1969

City**	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Biloxi	2,072	1,809	263	1,294	1,072	222	26.7	25.8	31.8
Brookhaven	577	352	225	225	112	113	21.0	16.8	27.9
Canton	319	101	218	307	62	245	29.2	13.6	41.2
Clarksdale	666	338	328	508	166	342	23.4	16.9	28.9
Cleveland	563	293	270	366	168	198	27.5	20.3	39.3
Columbus	874	512	362	598	273	325	23.2	17.0	33.2
Corinth	20	2	18	163	117	46	14.1	12.5	20.5
Greenville	1,220	535	685	1,044	382	662	26.3	20.3	31.8
Greenwood	806	397	409	460	184	276	20.5	16.5	24.5
Gulfport	1,449	1,026	423	949	679	270	23.3	20.5	35.3
Hattiesburg	1,447	1,119	328	768	462	306	20.1	17.0	27.4
Jackson	5,258	2,464	2,794	3,297	1,522	1,775	21.4	16.4	28.9
Laurel	1,924	778	1,146	482	254	228	20.0	16.7	25.5
McComb	492	300	192	217	121	96	18.1	15.2	24.0
Meridian	2,112	1,134	978	1,050	581	469	23.3	19.4	30.8
Moss Point	63	1	62	530	228	302	27.4	20.9	36.0
Natchez	1,563	753	810	570	210	360	28.9	21.6	36.1
Oxford	478	340	138	205	156	49	14.8	13.8	19.4
Pascagoula	1,565	1,193	372	684	576	108	25.1	24.3	30.0
Picayune	486	417	69	210	127	83	20.1	18.3	23.5
Starkville	544	274	270	289	192	97	25.4	22.6	33.6
Tupelo	1,229	1,006	223	354	296	58	17.3	17.5	16.1
Vicksburg	1,374	604	770	550	253	297	21.6	19.7	23.5
Yazoo City	323	189	134	287	111	176	26.6	25.7	27.1

* Births per 1,000 population.

** Cities having 10,000 population and over according to 1970 census.

TABLE 3
LIVE BIRTHS
MISSISSIPPI
1969

By Place of Residence of Mother, by Race and Attendant, and by Sex

County	Total	White				Nonwhite				Sex	
		Total	Hosp. or Clinic	Phys., Other Place	Other	Total	Hosp. or Clinic	Phys., Other Place	Other	Male	Female
<u>STATE</u>	<u>47,120</u>	<u>24,025</u>	<u>23,911</u>	<u>25</u>	<u>89</u>	<u>23,095</u>	<u>17,133</u>	<u>405</u>	<u>5,557</u>	<u>24,026</u>	<u>23,094</u>
Adams	890	393	393	0	0	497	492	1	4	440	450
Alcorn	431	365	361	1	3	66	40	2	24	212	219
Amite	275	87	87	0	0	188	109	0	79	143	132
Attala	394	178	177	0	1	216	107	1	108	190	204
Benton	132	64	62	1	1	68	28	1	39	75	57
Bolivar	1,370	339	339	0	0	1,031	882	1	148	680	690
Calhoun	276	170	170	0	0	106	84	2	20	146	130
Carroll	154	61	60	0	1	93	54	0	39	76	78
Chickasaw	358	171	171	0	0	187	160	1	26	201	157
Choctaw	157	75	75	0	0	82	53	1	28	83	74
Claiborne	202	37	37	0	0	165	114	1	50	100	102
Clarke	284	137	137	0	0	147	139	1	7	136	148
Clay	316	130	130	0	0	186	142	5	39	154	162
Coahoma	943	243	243	0	0	700	461	5	234	492	451
Copiah	478	148	146	2	0	330	282	12	36	231	247
Covington	276	160	159	0	1	116	110	1	5	150	126
DeSoto	781	429	429	0	0	352	273	1	78	383	398
Forrest	1,169	778	777	0	1	391	344	1	46	608	561
Franklin	123	56	56	0	0	67	49	6	12	71	52
George	279	238	237	0	1	41	33	0	8	157	122
Greene	186	129	128	1	0	57	49	0	8	91	95
Grenada	446	187	187	0	0	259	113	4	142	215	231
Hancock	268	227	219	0	8	41	21	0	20	136	132
Harrison	3,171	2,484	2,467	3	14	687	498	8	181	1,588	1,583

Table 3 - Page 2

Live Births - Mississippi - 1969

County	Total	White				Nonwhite				Sex	
		Total	Hosp. or Clinic	Phys., Other Place	Other	Total	Hosp. or Clinic	Phys., Other Place	Other	Male	Female
Hinds	4,480	2,132	2,129	2	1	2,348	2,264	4	80	2,311	2,169
Holmes	512	104	104	0	0	408	139	0	269	275	237
Humphreys	371	83	83	0	0	288	125	2	161	206	165
Issaquena	56	18	18	0	0	38	27	0	11	32	24
Itawamba	244	224	223	0	1	20	19	1	0	124	120
Jackson	2,060	1,582	1,575	0	7	478	380	27	71	1,056	1,004
Jasper	330	124	124	0	0	206	200	2	4	156	174
Jefferson	160	21	21	0	0	139	115	0	24	83	77
Jeff Davis	292	112	112	0	0	180	160	0	20	137	155
Jones	990	659	654	0	5	331	330	1	0	490	500
Kemper	221	57	57	0	0	164	131	1	32	125	96
Lafayette	466	285	285	0	0	181	109	0	72	240	226
Lamar	297	246	246	0	0	51	50	1	0	147	150
Lauderdale	1,525	859	858	0	1	666	647	7	12	762	763
Lawrence	200	114	114	0	0	86	62	5	19	99	101
Leake	304	154	154	0	0	150	121	2	27	150	154
Lee	897	658	653	2	3	239	210	4	25	433	464
Leflore	1,002	303	302	0	1	699	303	0	396	550	452
Lincoln	532	327	325	0	2	205	95	110	0	271	261
Lowndes	1,008	502	496	0	6	506	291	1	214	516	492
Madison	768	169	168	0	1	599	374	9	216	402	366
Marion	485	283	282	0	1	202	148	4	50	250	235
Marshall	555	142	142	0	0	413	288	2	123	291	264
Monroe	725	444	441	2	1	281	240	2	39	357	368
Montgomery	230	96	96	0	0	134	86	0	48	114	116
Neshoba	396	229	228	0	1	167	159	2	6	217	179
Newton	363	201	201	0	0	162	124	38	0	196	167
Noxubee	345	68	68	0	0	277	168	1	108	158	187
Oktibbeha	587	318	317	1	0	269	213	1	55	308	279
Panola	585	179	177	1	1	406	324	23	59	315	270

Table 3 - Page 3
Live Births - Mississippi - 1969

County	Total	White				Nonwhite				Sex	
		Total	Hosp. or Clinic	Phys., Other Place	Other	Total	Hosp. or Clinic	Phys., Other Place	Other	Male	Female
Pearl River	556	439	438	1	0	117	114	1	2	292	264
Perry	192	130	130	0	0	62	62	0	0	109	83
Pike	580	256	256	0	0	324	187	38	99	303	277
Pontotoc	290	215	215	0	0	75	69	1	5	153	137
Prentiss	329	279	279	0	0	50	37	0	13	172	157
Quitman	442	130	130	0	0	312	274	1	37	245	197
Rankin	802	564	560	0	4	238	220	2	16	389	413
Scott	461	244	244	0	0	217	168	11	38	226	235
Sharkey	229	43	42	1	0	186	142	0	44	121	108
Simpson	428	224	224	0	0	204	188	1	15	231	197
Smith	257	152	151	0	1	105	97	3	5	127	130
Stone	165	90	90	0	0	75	50	0	25	89	76
Sunflower	954	208	205	0	3	746	340	4	402	486	468
Tallahatchie	503	99	99	0	0	404	134	2	268	273	230
Tate	405	157	157	0	0	248	112	10	126	191	214
Tippah	296	230	229	1	0	66	60	0	6	146	150
Tishomingo	239	225	225	0	0	14	14	0	0	116	123
Tunica	370	53	53	0	0	317	198	1	118	185	185
Union	336	264	261	0	3	72	39	0	33	161	175
Walthall	248	98	89	0	9	150	78	5	67	145	103
Warren	917	498	497	0	1	419	403	0	16	458	459
Washington	1,819	613	605	5	3	1,206	944	13	249	918	901
Wayne	345	191	190	0	1	154	141	3	10	173	172
Webster	174	114	114	0	0	60	47	2	11	87	87
Wilkinson	197	40	40	0	0	157	103	0	54	108	89
Winston	384	170	170	0	0	214	130	0	84	184	200
Yalobusha	220	87	86	1	0	133	100	2	31	112	108
Yazoo	637	233	232	0	1	404	143	0	261	297	340

Table 3 - Page 4

Live Births - Mississippi - 1969

City*	Total	White				Nonwhite				Sex	
		Total	Hosp. or Clinic	Phys., Other Place	Other	Total	Hosp. or Clinic	Phys., Other Place	Other	Male	Female
Biloxi	1,294	1,072	1,067	1	4	222	196	0	26	656	638
Brookhaven	225	112	112	0	0	113	45	68	0	109	116
Canton	307	62	62	0	0	245	174	5	66	160	147
Clarksdale	508	166	166	0	0	342	293	3	46	262	246
Cleveland	366	168	168	0	0	198	167	0	31	183	183
Columbus	598	273	269	0	4	325	184	0	141	306	292
Corinth	163	117	115	0	2	46	28	2	16	82	81
Greenville	1,044	382	376	4	2	662	559	3	100	522	522
Greenwood	460	184	184	0	0	276	183	0	93	255	205
Gulfport	949	679	674	2	3	270	202	7	61	494	455
Hattiesburg	768	462	462	0	0	306	271	1	34	401	367
Jackson	3,297	1,522	1,520	1	1	1,775	1,749	2	24	1,694	1,603
Laurel	482	254	253	0	1	228	227	1	0	242	240
McComb	217	121	121	0	0	96	58	14	24	113	104
Meridian	1,050	581	580	0	1	469	464	5	0	517	533
Moss Point	530	228	227	0	1	302	243	16	43	284	246
Natchez	570	210	210	0	0	360	357	1	2	281	289
Oxford	205	156	156	0	0	49	43	0	6	108	97
Pascagoula	684	576	572	0	4	108	78	10	20	346	338
Picayune	210	127	126	1	0	83	81	1	1	102	108
Starkville	289	192	192	0	0	97	80	0	17	150	139
Tupelo	354	296	295	1	0	58	54	2	2	178	176
Vicksburg	550	253	253	0	0	297	289	0	8	265	285
Yazoo City	287	111	111	0	0	176	79	0	97	122	165

* Cities having 10,000 population and over according to 1970 census.

TABLE 4
FETAL DEATHS
MISSISSIPPI
1969

By Place of Occurrence; by Place of Residence of Mother, by Race and Attendant, and by Sex

County	Total by Occurrence	Total by Residence	White				Nonwhite				Sex		
			Total	Hosp. or Clinic	Phys., Other Place	Other	Total	Hosp. or Clinic	Phys., Other Place	Other	Male	Female	Unde-terminated
<u>STATE</u>	<u>2,514</u>	<u>2,471</u>	<u>1,177</u>	<u>1,140</u>	<u>34</u>	<u>3</u>	<u>1,294</u>	<u>1,059</u>	<u>45</u>	<u>190</u>	<u>764</u>	<u>564</u>	<u>1,143</u>
Adams	66	38	15	14	1	0	23	23	0	0	14	9	15
Alcorn	7	16	11	10	0	1	5	4	1	0	4	5	7
Amite	1	9	1	1	0	0	8	7	0	1	6	2	1
Attala	16	18	8	8	0	0	10	7	0	3	8	6	4
Benton	1	7	3	3	0	0	4	3	0	1	2	3	2
Bolivar	64	54	9	9	0	0	45	38	1	6	22	20	12
Calhoun	8	12	5	5	0	0	7	6	1	0	4	2	6
Carroll	1	5	1	1	0	0	4	4	0	0	1	2	2
Chickasaw	25	17	8	7	1	0	9	7	0	2	5	3	9
Choctaw	2	3	2	2	0	0	1	1	0	0	1	1	1
Claiborne	4	16	5	5	0	0	11	8	0	3	3	7	6
Clarke	1	11	4	3	1	0	7	5	1	1	3	3	5
Clay	9	11	4	4	0	0	7	5	2	0	4	4	3
Coahoma	64	55	16	16	0	0	39	30	3	6	25	16	14
Copiah	14	19	4	4	0	0	15	15	0	0	5	5	9
Covington	2	6	4	4	0	0	2	2	0	0	4	1	1
DeSoto	2	12	4	4	0	0	8	7	0	1	6	6	0
Forrest	40	36	13	13	0	0	23	19	0	4	11	14	11
Franklin	0	1	0	0	0	0	1	1	0	0	0	1	0
George	6	8	5	5	0	0	3	3	0	0	4	4	0
Greene	2	3	1	1	0	0	2	1	0	1	3	0	0
Grenada	19	22	7	7	0	0	15	10	0	5	12	8	2
Hancock	0	8	8	8	0	0	0	0	0	0	4	1	3
Harrison	300	258	205	200	3	2	53	44	4	5	50	30	178

Table 4 - Page 2
Fetal Deaths - Mississippi - 1969

County	Total by Occurrence	Total by Residence	White				Nonwhite				Sex		
			Total	Hosp. or Clinic	Phys., Other Place	Other	Total	Hosp. or Clinic	Phys., Other Place	Other	Male	Female	Unde-terminated
Hinds	644	464	232	228	4	0	232	225	2	5	69	56	339
Holmes	33	36	3	3	0	0	33	12	0	21	14	17	5
Humphreys	14	22	5	5	0	0	17	10	0	7	10	5	7
Issaquena	1	2	1	1	0	0	1	0	0	1	0	1	1
Itawamba	2	6	6	5	1	0	0	0	0	0	6	0	0
Jackson	37	72	55	54	1	0	17	16	1	0	30	8	34
Jasper	3	6	1	1	0	0	5	4	1	0	3	2	1
Jefferson	0	5	0	0	0	0	5	5	0	0	4	1	0
Jeff Davis	6	12	2	2	0	0	10	8	0	2	4	5	3
Jones	73	39	21	19	2	0	18	18	0	0	20	10	9
Kemper	2	10	5	5	0	0	5	4	0	1	1	5	4
Lafayette	7	6	2	2	0	0	4	3	0	1	3	2	1
Lamar	1	6	5	5	0	0	1	1	0	0	2	3	1
Lauderdale	116	84	60	59	1	0	24	24	0	0	17	13	54
Lawrence	5	10	4	4	0	0	6	6	0	0	5	2	3
Leake	6	12	2	2	0	0	10	5	0	5	5	3	4
Lee	29	23	16	16	0	0	7	7	0	0	14	6	3
Leflore	81	72	27	18	9	0	45	30	7	8	18	16	38
Lincoln	7	17	5	5	0	0	12	10	2	0	9	4	4
Lowndes	42	38	12	12	0	0	26	18	0	8	21	17	0
Madison	27	49	21	21	0	0	28	21	0	7	19	10	20
Marion	44	45	32	31	1	0	13	11	2	0	11	9	25
Marshall	23	31	9	9	0	0	22	15	0	7	16	10	5
Monroe	11	9	5	5	0	0	4	4	0	0	4	5	0
Montgomery	12	11	3	2	1	0	8	7	0	1	3	5	3
Neshoba	11	9	4	4	0	0	5	4	0	1	4	4	1
Newton	18	16	5	5	0	0	11	8	3	0	5	1	10
Noxubee	12	14	4	4	0	0	10	5	0	5	6	4	4
Oktibbeha	11	12	2	2	0	0	10	8	1	1	8	3	1
Panola	25	21	6	6	0	0	15	14	0	1	8	11	2

Table 4 - Page 3
Fetal Deaths - Mississippi - 1969

County	Total by Occurrence	Total by Residence	White				Nonwhite				Sex		
			Total	Hosp. or Clinic	Phys., Other Place	Other	Total	Hosp. or Clinic	Phys., Other Place	Other	Male	Female	Unde-terminated
Pearl River	22	13	9	8	1	0	4	4	0	0	7	2	4
Perry	6	8	5	5	0	0	3	3	0	0	5	3	0
Pike	16	14	4	4	0	0	10	4	0	6	6	6	2
Pontotoc	6	5	4	4	0	0	1	0	0	1	2	3	0
Prentiss	17	13	11	11	0	0	2	2	0	0	9	2	2
Quitman	17	26	7	6	1	0	19	17	1	1	11	12	3
Rankin	3	71	51	51	0	0	20	19	0	1	14	5	52
Scott	10	15	9	9	0	0	6	4	0	2	5	4	6
Sharkey	2	17	5	4	1	0	12	11	0	1	8	3	6
Simpson	5	21	10	10	0	0	11	11	0	0	4	6	11
Smith	0	12	7	7	0	0	5	5	0	0	4	3	5
Stone	3	7	3	3	0	0	4	3	0	1	1	5	1
Sunflower	36	49	6	5	1	0	43	29	2	12	28	17	4
Tallahatchie	15	17	3	3	0	0	14	4	4	6	9	6	2
Tate	4	13	2	2	0	0	11	7	0	4	8	5	0
Tippah	25	25	16	15	1	0	9	8	1	0	4	6	15
Tishomingo	2	3	3	3	0	0	0	0	0	0	1	2	0
Tunica	25	25	1	0	1	0	24	15	0	9	17	5	3
Union	26	21	14	14	0	0	7	6	0	1	6	4	11
Walthall	10	10	3	3	0	0	7	4	0	3	1	6	3
Warren	135	89	48	47	1	0	41	41	0	0	12	9	68
Washington	128	120	38	37	1	0	82	72	2	8	30	34	56
Wayne	9	12	7	7	0	0	5	5	0	0	4	4	4
Webster	7	11	3	3	0	0	8	4	1	3	6	3	2
Wilkinson	7	9	1	1	0	0	8	3	0	5	5	4	0
Winston	6	11	3	3	0	0	8	7	0	1	5	5	1
Yalobusha	1	6	4	4	0	0	2	2	0	0	4	2	0
Yazoo	14	24	2	2	0	0	22	16	2	4	8	7	9

Table 4 - Page 4
Fetal Deaths - Mississippi - 1969

City*	Total by Occurrence	Total by Residence	White				Nonwhite				Sex		
			Total	Hosp. or Clinic	Phys., Other Place	Other	Total	Hosp. or Clinic	Phys., Other Place	Other	Male	Female	Unde-terminated
Biloxi	227	123	101	101	0	0	22	21	0	1	19	18	86
Brookhaven	7	8	1	1	0	0	7	6	1	0	3	3	2
Canton	22	29	12	12	0	0	17	15	0	2	11	5	13
Clarksdale	58	37	15	15	0	0	22	18	2	2	15	11	11
Cleveland	18	10	3	3	0	0	7	6	0	1	4	4	2
Columbus	40	25	7	7	0	0	18	13	0	5	13	12	0
Corinth	0	6	2	2	0	0	4	3	1	0	2	2	2
Greenville	104	73	27	26	1	0	46	42	2	2	13	21	39
Greenwood	66	40	15	10	5	0	25	21	3	1	7	9	24
Gulfport	72	78	53	50	3	0	25	20	3	2	16	7	55
Hattiesburg	38	26	9	9	0	0	17	14	0	3	9	9	8
Jackson	532	368	179	176	3	0	189	188	0	1	49	39	280
Laurel	71	18	7	5	2	0	11	11	0	0	11	4	3
McComb	10	8	4	4	0	0	4	2	0	2	3	3	2
Meridian	115	52	34	34	0	0	18	18	0	0	11	8	33
Moss Point	2	16	8	7	1	0	8	7	1	0	9	4	3
Natchez	66	26	10	9	1	0	16	16	0	0	10	4	12
Oxford	6	4	2	2	0	0	2	2	0	0	2	1	1
Pascagoula	30	13	9	9	0	0	4	4	0	0	9	2	2
Picayune	19	5	3	3	0	0	2	2	0	0	2	0	3
Starkville	10	4	1	1	0	0	3	2	1	0	2	1	1
Tupelo	27	12	8	8	0	0	4	4	0	0	8	3	1
Vicksburg	133	63	34	34	0	0	29	29	0	0	9	5	49
Yazoo City	8	9	0	0	0	0	9	6	1	2	4	3	2

* Cities having 10,000 population and over according to 1970 census.

TABLE 5
DEATHS
MISSISSIPPI
1969

By Cause of Death, by Race, Residence Data

International Classification Number*	Cause of Death**	Total	White	Non-white
000-999	<u>ALL CAUSES</u>	<u>23,865</u>	<u>13,873</u>	<u>9,992</u>
000-136	<u>INFECTIVE AND PARASITIC DISEASES</u>	<u>319</u>	<u>113</u>	<u>206</u>
000-009	<u>Intestinal infectious diseases</u>	<u>126</u>	<u>30</u>	<u>96</u>
002,003	<u>Paratyphoid fever and other Salmonella infections</u>	<u>2</u>	<u>1</u>	<u>1</u>
003	Other Salmonella infections	2	1	1
004,006	<u>Bacillary dysentery and amebiasis</u>	<u>11</u>	<u>2</u>	<u>9</u>
004	Bacillary dysentery	4	0	4
006	Amebiasis	7	2	5
008,009	<u>Enteritis and other diarrheal diseases</u>	<u>113</u>	<u>27</u>	<u>86</u>
008	Enteritis due to other specified organism	7	2	5
009	Diarrheal disease	106	25	81
010-019	<u>Tuberculosis, all forms</u>	<u>54</u>	<u>24</u>	<u>30</u>
010-012	<u>Respiratory</u>	<u>47</u>	<u>21</u>	<u>26</u>
011	Pulmonary	46	20	26
012	Other respiratory	1	1	0
013-018	<u>Other forms</u>	<u>3</u>	<u>0</u>	<u>3</u>
013	Meninges and central nervous system	1	0	1
014	Intestines, peritoneum, and mesenteric glands	1	0	1
017	Other organs	1	0	1
019	Late effects of tuberculosis	4	3	1
020-027	<u>Zoonotic bacterial diseases</u>	<u>0</u>	<u>0</u>	<u>0</u>
030-039	<u>Other bacterial diseases</u>	<u>81</u>	<u>27</u>	<u>54</u>
033	Whooping cough	1	0	1
034	Streptococcal sore throat and scarlet fever	2	2	0
036	Meningococcal infection	7	4	3
037	Tetanus	7	0	7
038	Septicemia	56	18	38
039	Other bacterial diseases	8	3	5
040-046	<u>Poliomyelitis and other enterovirus diseases of central nervous</u>	<u>4</u>	<u>2</u>	<u>2</u>
045	Aseptic meningitis due to enterovirus	1	1	0
046	Other enterovirus diseases of central nervous system	3	1	2
050-057	<u>Viral diseases accompanied by exanthem</u>	<u>2</u>	<u>1</u>	<u>1</u>
055	Measles	1	0	1
056	Rubella	1	1	0
060-068	<u>Arthropod-borne viral diseases</u>	<u>6</u>	<u>5</u>	<u>1</u>
062-065	<u>Viral encephalitis</u>	<u>6</u>	<u>5</u>	<u>1</u>
065	Viral encephalitis, unspecified	6	5	1
070-079	<u>Other viral diseases</u>	<u>21</u>	<u>13</u>	<u>8</u>
070	Infectious hepatitis	16	8	8
072	Mumps	1	1	0

Table 5 - Page 2
Deaths - Mississippi - 1969

International Classification Number*	Cause of Death**	Total	White	Non-white
075	Infectious mononucleosis	1	1	0
079	Other viral diseases	3	3	0
<u>080-089</u>	<u>Rickettsioses and other arthropod-borne diseases</u>	<u>1</u>	<u>1</u>	<u>0</u>
080-083	<u>Typhus and other rickettsioses</u>	<u>1</u>	<u>1</u>	<u>0</u>
082	Tick-borne rickettsioses	1	1	0
<u>090-099</u>	<u>Syphilis and other venereal diseases</u>	<u>12</u>	<u>1</u>	<u>11</u>
090-097	<u>Syphilis and its sequelae</u>	<u>12</u>	<u>1</u>	<u>11</u>
093	Cardiovascular	8	1	7
094	Central nervous system	4	0	4
<u>100-104</u>	<u>Other spirochetal diseases</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>110-117</u>	<u>Mycoses</u>	<u>6</u>	<u>5</u>	<u>1</u>
112	Moniliasis	1	0	1
115	Histoplasmosis	2	2	0
116	Blastomycosis	1	1	0
117	Other systemic mycosis	2	2	0
<u>120-129</u>	<u>Helminthiases</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>130-136</u>	<u>Other infective and parasitic diseases</u>	<u>6</u>	<u>4</u>	<u>2</u>
135	Sarcoidosis	4	3	1
136	Other and unspecified infective and parasitic diseases	2	1	1
<u>140-239</u>	<u>NEOPLASMS</u>	<u>3,332</u>	<u>2,166</u>	<u>1,166</u>
<u>140-209</u>	<u>Malignant neoplasms</u>	<u>3,295</u>	<u>2,146</u>	<u>1,149</u>
<u>140-149</u>	<u>Buccal cavity and pharynx</u>	<u>64</u>	<u>50</u>	<u>14</u>
140	Lip	1	1	0
141	Tongue	9	8	1
142	Salivary gland	12	7	5
143	Gum	3	2	1
144	Floor of mouth	4	4	0
145	Other and unspecified parts of mouth	8	8	0
146	Oropharynx	5	2	3
147	Nasopharynx	7	5	2
148	Hypopharynx	1	1	0
149	Pharynx, unspecified	14	12	2
<u>150-159</u>	<u>Digestive organs and peritoneum</u>	<u>866</u>	<u>531</u>	<u>335</u>
150	Esophagus	41	27	14
151	Stomach	200	89	111
152	Small intestine, including duodenum	9	7	2
153	Large intestine, except rectum	278	184	94
154	Rectum and rectosigmoid junction	66	45	21
155	Liver and intrahepatic bile ducts, specified as primary	20	14	6
156	Gallbladder and bile ducts	15	10	5
157	Pancreas	219	143	76
158	Peritoneum and retroperitoneal tissue	8	8	0
159	Unspecified digestive organs	10	4	6
<u>160-163</u>	<u>Respiratory system</u>	<u>669</u>	<u>492</u>	<u>177</u>
160	Nose, nasal cavities, middle ear, and accessory sinuses	7	3	4
161	Larynx	23	17	6

Table 5 - Page 3
Deaths - Mississippi - 1969

International Classification Number*	Cause of Death**	Total	White	Non-white
162	Trachea, bronchus, and lung	637	470	167
163	Other and unspecified respiratory organs	2	2	0
<u>170-174</u>	<u>Bone, connective tissue, skin, and breast</u>	<u>364</u>	<u>269</u>	<u>95</u>
170	Bone	18	15	3
171	Connective and other soft tissue	18	13	5
172	Melanoma of skin	33	29	4
173	Skin, other	43	37	6
174	Breast	252	175	77
<u>180-189</u>	<u>Genitourinary organs</u>	<u>629</u>	<u>333</u>	<u>296</u>
<u>(180-187)</u>	<u>(Genital organs)</u>	<u>509</u>	<u>249</u>	<u>260</u>
180	Cervix uteri	103	49	54
181	Chorionepithelioma	1	1	0
182	Uterus, other	64	25	39
183	Ovary, fallopian tube, and broad ligament	79	51	28
184	Other and unspecified female genital organs	9	6	3
185	Prostate	245	111	134
186	Testis	6	6	0
187	Other and unspecified male genital organs	2	0	2
<u>(188,189)</u>	<u>(Urinary organs)</u>	<u>120</u>	<u>84</u>	<u>36</u>
188	Bladder	71	45	26
189	Other and unspecified urinary organs	49	39	10
<u>190-199</u>	<u>Other and unspecified sites</u>	<u>365</u>	<u>218</u>	<u>147</u>
190	Eye	4	3	1
191	Brain	68	60	8
192	Other parts of nervous system	19	10	9
193	Thyroid gland	10	7	3
194	Other endocrine glands	4	3	1
195	Ill-defined sites	36	17	19
196	Lymph nodes, secondary and unspecified	4	4	0
197	Respiratory and digestive systems, secondary	84	51	33
198	Other secondary	9	6	3
199	Site unspecified	127	57	70
<u>200-209</u>	<u>Lymphatic and hematopoietic tissue</u>	<u>338</u>	<u>253</u>	<u>85</u>
200	Lymphosarcoma and reticulum-cell sarcoma	66	52	14
201	Hodgkin's disease	34	28	6
202	Lymphoid tissue, other	36	31	5
203	Multiple myeloma	52	33	19
208	Polycythemia vera	4	3	1
209	Myelofibrosis	7	5	2
<u>(204-207)</u>	<u>(Leukemia)</u>	<u>139</u>	<u>101</u>	<u>38</u>
204	Lymphatic	51	43	8
205	Myeloid	55	36	19
206	Monocytic	3	3	0
207	Other and unspecified	30	19	11
<u>210-228</u>	<u>Benign neoplasms</u>	<u>22</u>	<u>10</u>	<u>12</u>
210	Benign neoplasm of buccal cavity and pharynx	1	0	1
211	Benign neoplasm of other parts of digestive system	4	3	1
215	Benign neoplasm of muscular and connective tissue, other	1	1	0

Table 5 - Page 4
Deaths - Mississippi - 1969

International Classification Number*	Cause of Death**	Total	White	Non-white
218	Uterine fibroma	5	0	5
225	Benign neoplasm of brain and other parts of nervous system	7	4	3
226	Benign neoplasm of endocrine glands	3	2	1
228	Benign neoplasm of other and unspecified organs and tissues	1	0	1
<u>230-239</u>	<u>Neoplasms of unspecified nature</u>	<u>15</u>	<u>10</u>	<u>5</u>
231	Respiratory organs	1	1	0
237	Other genitourinary organs	1	1	0
238	Eye, brain, and other parts of nervous system	10	7	3
239	Other and unspecified organs	3	1	2
<u>240-279</u>	<u>ENDOCRINE, NUTRITIONAL, AND METABOLIC DISEASES</u>	<u>550</u>	<u>258</u>	<u>292</u>
<u>240-246</u>	<u>Diseases of thyroid gland</u>	<u>9</u>	<u>4</u>	<u>5</u>
<u>240, 241</u>	<u>Nontoxic goiter</u>	<u>2</u>	<u>0</u>	<u>2</u>
240	Simple	1	0	1
241	Nontoxic nodular	1	0	1
242	Thyrotoxicosis with or without goiter	5	2	3
244	Myxedema	2	2	0
<u>250-258</u>	<u>Diseases of other endocrine glands</u>	<u>490</u>	<u>231</u>	<u>259</u>
250	Diabetes mellitus	477	223	254
251	Disorders of pancreatic internal secretion other than diabetes mellitus	3	2	1
252	Diseases of parathyroid gland	1	1	0
253	Diseases of pituitary gland	3	1	2
255	Diseases of adrenal glands	4	2	2
258	Polyglandular dysfunction and other diseases of endocrine glands	2	2	0
<u>260-269</u>	<u>Avitaminoses and other nutritional deficiency</u>	<u>22</u>	<u>7</u>	<u>15</u>
262	Niacin deficiency	1	1	0
268	Nutritional marasmus	3	1	2
269	Other nutritional deficiency	18	5	13
<u>270-279</u>	<u>Other metabolic diseases</u>	<u>29</u>	<u>16</u>	<u>13</u>
272	Congenital disorders of lipid metabolism	2	2	0
273	Other and unspecified congenital disorders of metabolism	8	6	2
275	Plasma protein abnormalities	1	0	1
277	Obesity not specified as of endocrine origin	15	6	9
278	Other hyperalimentation	1	1	0
279	Other and unspecified metabolic diseases	2	1	1
<u>280-289</u>	<u>DISEASES OF THE BLOOD AND BLOOD-FORMING ORGANS</u>	<u>71</u>	<u>37</u>	<u>34</u>
<u>280-285</u>	<u>Anemias</u>	<u>50</u>	<u>18</u>	<u>32</u>
280	Iron deficiency	1	0	1
282	Hereditary hemolytic	17	1	16
283	Acquired hemolytic	2	1	1
284	Aplastic	12	9	3
285	Other and unspecified	18	7	11
286	Coagulation defects	3	3	0

Table 5 - Page 5
Deaths - Mississippi - 1969

International Classification Number*	Cause of Death**	Total	White	Non-white
287	Purpura and other hemorrhagic conditions	9	9	0
288	Agranulocytosis	6	6	0
289	Other diseases of blood and blood-forming organs	3	1	2
<u>290-315</u>	<u>MENTAL DISORDERS</u>	<u>76</u>	<u>40</u>	<u>36</u>
<u>290-299</u>	<u>Psychoses</u>	<u>16</u>	<u>11</u>	<u>5</u>
290	Senile and presenile dementia	3	2	1
291	Alcoholic psychosis	9	6	3
294	Psychosis associated with other physical conditions	1	1	0
295	Schizophrenia	1	1	0
299	Unspecified psychosis	2	1	1
<u>300-309</u>	<u>Neuroses, personality disorders, and other non-psychotic mental disorders</u>	<u>58</u>	<u>28</u>	<u>30</u>
300	Neuroses	1	1	0
303	Alcoholism	44	16	28
304	Drug dependence	3	3	0
306	Special symptoms not elsewhere classified	2	1	1
309	Mental disorders not specified as psychotic associated with physical conditions	8	7	1
<u>310-315</u>	<u>Mental retardation</u>	<u>2</u>	<u>1</u>	<u>1</u>
315	Unspecified	2	1	1
<u>320-389</u>	<u>DISEASES OF THE NERVOUS SYSTEM AND SENSE ORGANS</u>	<u>169</u>	<u>90</u>	<u>79</u>
<u>320-324</u>	<u>Inflammatory diseases of central nervous system</u>	<u>40</u>	<u>11</u>	<u>29</u>
320	Meningitis	32	8	24
322	Intracranial and intraspinal abscess	1	0	1
323	Encephalitis, myelitis, and encephalomyelitis	4	2	2
324	Late effects of intracranial abscess or pyogenic infection	3	1	2
<u>330-333</u>	<u>Hereditary and familial diseases of nervous system</u>	<u>12</u>	<u>9</u>	<u>3</u>
330	Hereditary neuromuscular disorders	6	3	3
331	Hereditary diseases of the striatopallidal system	4	4	0
333	Other hereditary and familial diseases of nervous system	2	2	0
<u>340-349</u>	<u>Other diseases of central nervous system</u>	<u>108</u>	<u>65</u>	<u>43</u>
340	Multiple sclerosis	5	3	2
342	Paralysis agitans	18	14	4
343	Cerebral spastic infantile paralysis	8	5	3
344	Other cerebral paralysis	9	2	7
345	Epilepsy	34	14	20
347	Other diseases of brain	19	15	4
348	Motor neurone disease	13	10	3
349	Other diseases of spinal cord	2	2	0
<u>350-358</u>	<u>Diseases of nerves and peripheral ganglia</u>	<u>4</u>	<u>1</u>	<u>3</u>
354	Polyneuritis and polyradiculitis	3	1	2
357	Other diseases of peripheral nerves except autonomic	1	0	1

Table 5 - Page 6
Deaths - Mississippi - 1969

International Classification Number*	Cause of Death**	Total	White	Non-white
<u>360-369</u>	<u>Inflammatory diseases of the eye</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>370-379</u>	<u>Other diseases and conditions of eye</u>	<u>1</u>	<u>1</u>	<u>0</u>
374	Cataract	1	1	0
<u>380-389</u>	<u>Diseases of the ear and mastoid process</u>	<u>4</u>	<u>3</u>	<u>1</u>
<u>381-383</u>	<u>Otitis media and mastoiditis</u>	<u>3</u>	<u>3</u>	<u>0</u>
381	Otitis media without mention of mastoiditis	2	2	0
383	Mastoiditis without mention of otitis media	1	1	0
387	Other diseases of ear and mastoid process	1	0	1
<u>390-458</u>	<u>DISEASES OF THE CIRCULATORY SYSTEM</u>	<u>11,864</u>	<u>7,390</u>	<u>4,474</u>
<u>390-392</u>	<u>Active rheumatic fever</u>	<u>5</u>	<u>1</u>	<u>4</u>
390	Rheumatic fever without mention of heart involvement	3	0	3
391	Rheumatic fever with heart involvement	2	1	1
<u>393-398</u>	<u>Chronic rheumatic heart disease</u>	<u>92</u>	<u>54</u>	<u>38</u>
394	Diseases of mitral valve	21	10	11
395	Diseases of aortic valve	28	18	10
396	Diseases of mitral and aortic valves	5	3	2
397	Diseases of other endocardial structures	2	2	0
398	Other heart disease, specified as rheumatic	36	21	15
<u>400-404</u>	<u>Hypertensive disease</u>	<u>405</u>	<u>133</u>	<u>272</u>
<u>400,401,403</u>	<u>Hypertension</u>	<u>161</u>	<u>52</u>	<u>109</u>
400	Malignant hypertension	44	6	38
401	Essential benign hypertension	37	13	24
403	Hypertensive renal disease	80	33	47
<u>402,404</u>	<u>Hypertensive heart disease with or without renal disease</u>	<u>244</u>	<u>81</u>	<u>163</u>
402	Hypertensive heart disease	156	37	119
404	Hypertensive heart and renal disease	88	44	44
<u>410-414</u>	<u>Ischemic heart disease</u>	<u>6,799</u>	<u>4,603</u>	<u>2,196</u>
410	Acute myocardial infarction	4,068	3,041	1,027
411	Other acute and subacute forms of ischemic heart disease	70	40	30
412	Chronic ischemic heart disease	2,657	1,522	1,135
413	Angina pectoris	4	0	4
<u>420-429</u>	<u>Other forms of heart disease</u>	<u>746</u>	<u>403</u>	<u>343</u>
421	Acute and subacute endocarditis	5	3	2
422	Acute myocarditis	8	5	3
423	Chronic disease of pericardium, nonrheumatic	3	1	2
424	Chronic disease of endocardium	20	14	6
425	Cardiomyopathy	12	5	7
426	Pulmonary heart disease	3	2	1
427	Symptomatic heart disease	285	171	114
428	Other myocardial insufficiency	126	71	55
429	Ill-defined heart disease	284	131	153
<u>430-438</u>	<u>Cerebrovascular disease</u>	<u>3,058</u>	<u>1,681</u>	<u>1,377</u>
430	Subarachnoid hemorrhage	91	56	35
431	Cerebral hemorrhage	700	338	362

Table 5 - Page 7
Deaths - Mississippi - 1969

International Classification Number*	Cause of Death**	Total	White	Non-white
432	Occlusion of precerebral arteries	22	13	9
433	Cerebral thrombosis	668	415	253
434	Cerebral embolism	15	6	9
436	Acute but ill-defined cerebrovascular disease	1,230	633	597
437	Generalized ischemic cerebrovascular disease	312	209	103
438	Other and ill-defined cerebrovascular disease	20	11	9
<u>440-448</u>	<u>Diseases of arteries, arterioles, and capillaries</u>	<u>605</u>	<u>423</u>	<u>182</u>
440	Arteriosclerosis	331	232	99
<u>441-448</u>	<u>Other diseases of arteries, arterioles, and capillaries</u>	<u>274</u>	<u>191</u>	<u>83</u>
441	Aortic aneurysm (nonsyphilitic)	124	88	36
442	Other aneurysm	12	10	2
443	Other peripheral vascular disease	8	5	3
444	Arterial embolism and thrombosis	77	64	13
445	Gangrene	42	16	26
446	Polyarteritis nodosa and allied conditions	7	4	3
447	Other diseases of arteries and arterioles	4	4	0
<u>450-458</u>	<u>Diseases of veins and lymphatics, and other diseases of circulatory system</u>	<u>154</u>	<u>92</u>	<u>62</u>
<u>450-453</u>	<u>Venous thrombosis and embolism</u>	<u>138</u>	<u>85</u>	<u>53</u>
450	Pulmonary embolism and infarction	93	56	37
451	Phlebitis and thrombophlebitis	29	21	8
452	Portal vein thrombosis	2	1	1
453	Other venous embolism and thrombosis	14	7	7
<u>454-458</u>	<u>Other diseases of the circulatory system</u>	<u>16</u>	<u>7</u>	<u>9</u>
454	Varicose veins of lower extremities	2	1	1
455	Hemorrhoids	1	1	0
456	Varicose veins of other sites	1	1	0
457	Noninfective disease of lymphatic channels	1	0	1
458	Other diseases of circulatory system	11	4	7
<u>460-519</u>	<u>DISEASES OF THE RESPIRATORY SYSTEM</u>	<u>1,183</u>	<u>711</u>	<u>472</u>
<u>460-466</u>	<u>Acute respiratory infections, except influenza</u>	<u>27</u>	<u>6</u>	<u>21</u>
460	Acute nasopharyngitis (common cold)	7	0	7
463	Acute tonsillitis	1	0	1
464	Acute laryngitis and tracheitis	2	2	0
465	Acute upper respiratory infection of multiple or unspecified sites	4	1	3
466	Acute bronchitis and bronchiolitis	13	3	10
<u>470-474</u>	<u>Influenza</u>	<u>84</u>	<u>42</u>	<u>42</u>
470	Unqualified	25	11	14
471	With pneumonia	56	30	26
472	With other respiratory manifestations	1	0	1
473	With digestive manifestations	1	0	1
474	With nervous manifestations	1	1	0
<u>480-486</u>	<u>Pneumonia</u>	<u>642</u>	<u>343</u>	<u>299</u>
480	Viral	7	3	4
<u>481,482</u>	<u>Bacterial</u>	<u>72</u>	<u>42</u>	<u>30</u>
481	Pneumococcal	52	29	23
482	Other bacterial pneumonia	20	13	7

Table 5 - Page 8
Deaths - Mississippi - 1969

International Classification Number*	Cause of Death**	Total	White	Non-white
483	Pneumonia due to other specified organism	1	0	1
484	Acute interstitial, organism unspecified	12	8	4
485	Broncho, organism unspecified	214	117	97
486	Unspecified	336	173	163
<u>490-493</u>	<u>Bronchitis, emphysema, and asthma</u>	<u>296</u>	<u>239</u>	<u>57</u>
<u>490,491</u>	<u>Chronic and unqualified bronchitis</u>	<u>62</u>	<u>49</u>	<u>13</u>
490	Unqualified	12	5	7
491	Chronic	50	44	6
492	Emphysema	206	179	27
493	Asthma	28	11	17
<u>500-508</u>	<u>Other diseases of upper respiratory tract</u>	<u>7</u>	<u>5</u>	<u>2</u>
500	Hypertrophy of tonsils and adenoids	1	0	1
502	Chronic pharyngitis and nasopharyngitis	1	0	1
503	Chronic sinusitis	1	1	0
508	Other diseases of upper respiratory tract	4	4	0
<u>510-519</u>	<u>Other diseases of respiratory system</u>	<u>127</u>	<u>76</u>	<u>51</u>
510	Empyema	3	2	1
511	Pleurisy	2	1	1
513	Abscess of lung	7	2	5
514	Pulmonary congestion and hypostasis	25	3	22
515	Pneumoconiosis due to silica and silicates	2	1	1
517	Other chronic interstitial pneumonia	26	25	1
518	Bronchiectasis	6	5	1
519	Other diseases of respiratory system	56	37	19
<u>520-577</u>	<u>DISEASES OF THE DIGESTIVE SYSTEM</u>	<u>548</u>	<u>380</u>	<u>168</u>
<u>520-529</u>	<u>Diseases of oral cavity, salivary glands, and jaws</u>	<u>3</u>	<u>2</u>	<u>1</u>
<u>520-525</u>	<u>Diseases of teeth and supporting structures</u>	<u>2</u>	<u>2</u>	<u>0</u>
522	Diseases of pulp and periapical tissues	1	1	0
525	Other diseases and conditions of the teeth and supporting structures	1	1	0
528	Diseases of the oral soft tissues, excluding gingiva and tongue	1	0	1
<u>530-537</u>	<u>Diseases of esophagus, stomach, and duodenum</u>	<u>109</u>	<u>71</u>	<u>38</u>
530	Diseases of esophagus	5	4	1
<u>531-533</u>	<u>Peptic ulcer</u>	<u>87</u>	<u>59</u>	<u>28</u>
531	Ulcer of stomach	22	15	7
532	Ulcer of duodenum	40	26	14
533	Peptic ulcer, site unspecified	25	18	7
534	Gastrojejunal ulcer	5	4	1
535	Gastritis and duodenitis	5	1	4
536	Disorders of function of stomach	5	2	3
537	Other diseases of stomach and duodenum	2	1	1
<u>540-543</u>	<u>Appendicitis</u>	<u>24</u>	<u>8</u>	<u>16</u>
540	Acute appendicitis	23	8	15
541	Appendicitis, unqualified	1	0	1
<u>550-553</u>	<u>Hernia of abdominal cavity</u>	<u>31</u>	<u>14</u>	<u>17</u>
550	Inguinal, without mention of obstruction	3	1	2
551	Other, of abdominal cavity without mention of obstruction	6	4	2

Table 5 - Page 9
Deaths - Mississippi - 1969

International Classification Number*	Cause of Death**	Total	White	Non-white
552	Inguinal, with obstruction	7	2	5
553	Other, of abdominal cavity with obstruction	15	7	8
<u>560-569</u>	<u>Other diseases of intestine and peritoneum</u>	<u>132</u>	<u>89</u>	<u>43</u>
560	Intestinal obstruction without mention of hernia	37	21	16
561	Gastroenteritis and colitis, except ulcerative, of noninfectious origin	1	0	1
562	Diverticula of intestine	34	31	3
563	Chronic enteritis and ulcerative colitis	6	5	1
564	Functional disorders of intestines	1	1	0
565	Anal fissure and fistula	1	0	1
566	Abscess of anal and rectal regions	1	0	1
567	Peritonitis	12	7	5
568	Peritoneal adhesions	1	0	1
569	Other diseases of intestines and peritoneum	38	24	14
<u>570-577</u>	<u>Diseases of liver, gallbladder, and pancreas</u>	<u>249</u>	<u>196</u>	<u>53</u>
570	Acute and subacute necrosis of liver	6	3	3
571	Cirrhosis of liver	140	113	27
572	Suppurative hepatitis and liver abscess	4	3	1
573	Other diseases of liver	18	12	6
<u>574,575</u>	<u>Cholelithiasis, cholecystitis, and cholangitis</u>	<u>46</u>	<u>37</u>	<u>9</u>
574	Cholelithiasis	26	22	4
575	Cholecystitis and cholangitis, without mention of calculus	20	15	5
576	Other diseases of gallbladder and biliary ducts	13	10	3
577	Diseases of pancreas	22	18	4
<u>580-629</u>	<u>DISEASES OF THE GENITOURINARY SYSTEM</u>	<u>466</u>	<u>246</u>	<u>220</u>
<u>580-584</u>	<u>Nephritis and nephrosis</u>	<u>167</u>	<u>65</u>	<u>102</u>
<u>580,581</u>	<u>Acute nephritis and nephrotic syndrome</u>	<u>23</u>	<u>8</u>	<u>15</u>
580	Acute nephritis	12	6	6
581	Nephrotic syndrome	11	2	9
<u>582-584</u>	<u>Chronic and unqualified nephritis and renal sclerosis</u>	<u>144</u>	<u>57</u>	<u>87</u>
582	Chronic nephritis	120	52	68
583	Nephritis, unqualified	21	5	16
584	Renal sclerosis, unqualified	3	0	3
<u>590-599</u>	<u>Other diseases of urinary system</u>	<u>243</u>	<u>150</u>	<u>93</u>
590	Infections of kidney	171	106	65
591	Hydronephrosis	3	1	2
<u>592,594</u>	<u>Calculus of urinary system</u>	<u>8</u>	<u>7</u>	<u>1</u>
592	Of kidney and ureter	8	7	1
593	Other diseases of kidney and ureter	32	18	14
595	Cystitis	1	1	0
596	Other diseases of bladder	3	2	1
598	Stricture of urethra	4	1	3
599	Other diseases of urinary tract	21	14	7
<u>600-607</u>	<u>Diseases of male genital organs</u>	<u>41</u>	<u>21</u>	<u>20</u>
600	Hyperplasia of prostate	36	19	17
601	Prostatitis	5	2	3

Table 5 - Page 10
Deaths - Mississippi - 1969

International Classification Number*	Cause of Death**	Total	White	Non-white
<u>610-616</u>	<u>Diseases of breast, ovary, fallopian tube, and parametrium</u>	<u>3</u>	<u>0</u>	<u>3</u>
614	Salpingitis and oophoritis, unqualified	1	0	1
616	Diseases of parametrium and pelvic peritoneum (female)	2	0	2
<u>620-629</u>	<u>Diseases of uterus and other female genital organs</u>	<u>12</u>	<u>10</u>	<u>2</u>
620	Infective diseases of cervix uteri	1	0	1
623	Uterovaginal prolapse	2	2	0
624	Malposition of uterus	1	0	1
625	Other diseases of uterus	4	4	0
626	Disorders of menstruation	2	2	0
629	Other diseases of female genital organs	2	2	0
<u>630-678</u>	<u>COMPLICATIONS OF PREGNANCY, CHILDBIRTH, AND THE PUERPERIUM</u>	<u>23</u>	<u>3</u>	<u>20</u>
<u>630-634</u>	<u>Complications of pregnancy</u>	<u>3</u>	<u>1</u>	<u>2</u>
630	Infections of genital tract during pregnancy	1	0	1
631	Ectopic pregnancy	1	0	1
632	Hemorrhage of pregnancy	1	1	0
<u>635-639</u>	<u>Urinary infections and toxemias of pregnancy and the puerperium</u>	<u>11</u>	<u>1</u>	<u>10</u>
<u>636-639</u>	<u>Toxemias of pregnancy and the puerperium, except abortion with toxemia</u>	<u>11</u>	<u>1</u>	<u>10</u>
637	Pre-eclampsia, eclampsia, and toxemia, unspecified	11	1	10
<u>640-645</u>	<u>Abortion</u>	<u>2</u>	<u>0</u>	<u>2</u>
644	Not specified as induced or spontaneous	2	0	2
<u>650-662</u>	<u>Delivery</u>	<u>3</u>	<u>1</u>	<u>2</u>
653	Delivery complicated by other postpartum hemorrhage	1	0	1
657	Delivery complicated by prolonged labor of other origin	1	0	1
661	Delivery with other complications	1	1	0
<u>670-678</u>	<u>Complications of the puerperium</u>	<u>4</u>	<u>0</u>	<u>4</u>
<u>670,671,673</u>	<u>Sepsis of childbirth and the puerperium</u>	<u>3</u>	<u>0</u>	<u>3</u>
670	Sepsis of childbirth and the puerperium	1	0	1
671	Puerperal phlebitis and thrombosis	1	0	1
673	Puerperal pulmonary embolism	1	0	1
677	Other and unspecified complications of the puerperium	1	0	1
<u>680-709</u>	<u>DISEASES OF THE SKIN AND SUBCUTANEOUS TISSUE</u>	<u>34</u>	<u>17</u>	<u>17</u>
<u>680-686</u>	<u>Infections of skin and subcutaneous tissue</u>	<u>12</u>	<u>5</u>	<u>7</u>
680	Boil and carbuncle	1	0	1
682	Other cellulitis and abscess	7	2	5
686	Other local infections of skin and subcutaneous tissue	4	3	1

Table 5 - Page 11
Deaths - Mississippi - 1969

International Classification Number*	Cause of Death**	Total	White	Non-white
<u>690-698</u>	<u>Other inflammatory conditions of skin and subcutaneous tissue</u>	<u>7</u>	<u>3</u>	<u>4</u>
695	Erythematous conditions	6	2	4
698	Pruritus and related conditions	1	1	0
<u>700-709</u>	<u>Other diseases of skin and subcutaneous tissue</u>	<u>15</u>	<u>9</u>	<u>6</u>
707	Chronic ulcer of skin	14	8	6
708	Urticaria	1	1	0
<u>710-738</u>	<u>DISEASES OF THE MUSCULOSKELETAL SYSTEM AND CONNECTIVE TISSUE</u>	<u>59</u>	<u>41</u>	<u>18</u>
<u>710-718</u>	<u>Arthritis and rheumatism, except rheumatic fever</u>	<u>34</u>	<u>23</u>	<u>11</u>
<u>710-715</u>	<u>Arthritis and spondylitis</u>	<u>32</u>	<u>22</u>	<u>10</u>
712	Rheumatoid arthritis and allied conditions	18	14	4
713	Osteoarthritis and allied conditions	5	3	2
714	Other specified forms of arthritis	1	1	0
715	Arthritis, unspecified	8	4	4
<u>716-718</u>	<u>Nonarticular rheumatism and rheumatism unspecified</u>	<u>2</u>	<u>1</u>	<u>1</u>
716	Polymyositis and dermatomyositis	2	1	1
<u>720-729</u>	<u>Osteomyelitis and other diseases of bone and joint</u>	<u>5</u>	<u>3</u>	<u>2</u>
720	Osteomyelitis and periostitis	2	1	1
723	Other diseases of bone	1	1	0
725	Displacement of intervertebral disc	2	1	1
<u>730-738</u>	<u>Other diseases of musculoskeletal system</u>	<u>20</u>	<u>15</u>	<u>5</u>
732	Infective myositis and other inflammatory diseases of tendon and fascia	2	2	0
733	Other diseases of muscle, tendon, and fascia	5	4	1
734	Diffuse diseases of connective tissue	13	9	4
<u>740-759</u>	<u>CONGENITAL ANOMALIES</u>	<u>218</u>	<u>127</u>	<u>91</u>
<u>740-743</u>	<u>Nervous system</u>	<u>52</u>	<u>34</u>	<u>18</u>
740	Anencephalus	11	7	4
741	Spina bifida	19	16	3
742	Congenital hydrocephalus	14	7	7
743	Other congenital anomalies of nervous system	8	4	4
745	Ear, face, and neck	1	1	0
<u>746,747</u>	<u>Circulatory system</u>	<u>92</u>	<u>52</u>	<u>40</u>
746	Heart	80	42	38
747	Other of circulatory system	12	10	2
748	Respiratory system	12	9	3
<u>749-751</u>	<u>Digestive system</u>	<u>16</u>	<u>8</u>	<u>8</u>
750	Other congenital anomalies of upper alimentary tract	1	1	0
751	Other congenital anomalies of digestive system	15	7	8
<u>752,753</u>	<u>Genitourinary system</u>	<u>12</u>	<u>8</u>	<u>4</u>
752	Genital organs	1	1	0
753	Urinary system	11	7	4

Table 5 - Page 12
Deaths - Mississippi - 1969

International Classification Number*	Cause of Death**	Total	White	Non-white
<u>754-756</u>	<u>Musculoskeletal system</u>	<u>9</u>	<u>4</u>	<u>5</u>
756	Other congenital anomalies of musculoskeletal system	9	4	5
758	Other and unspecified	6	3	3
759	Congenital syndromes affecting multiple systems	18	8	10
<u>760-778</u>	<u>CERTAIN CAUSES OF MORTALITY IN EARLY INFANCY</u>	<u>766</u>	<u>329</u>	<u>437</u>
761	Other maternal conditions unrelated to pregnancy	7	2	5
762	Toxemia of pregnancy	5	1	4
763	Maternal ante- and intrapartum infection	7	0	7
<u>764-768</u>	<u>Difficult labor</u>	<u>19</u>	<u>11</u>	<u>8</u>
764	With abnormality of bones, organs, or tissues	1	0	1
766	With malposition of fetus	13	6	7
768	With other and unspecified complications	5	5	0
769	Other complications of pregnancy and childbirth	63	28	35
<u>770,771</u>	<u>Conditions of placenta and cord</u>	<u>49</u>	<u>22</u>	<u>27</u>
770	Of placenta	44	21	23
771	Of umbilical cord	5	1	4
772	Birth injury without mention of cause	19	6	13
<u>774,775</u>	<u>Hemolytic disease of newborn</u>	<u>16</u>	<u>14</u>	<u>2</u>
774	With kernicterus	1	1	0
775	Without mention of kernicterus	15	13	2
776	Anoxic and hypoxic conditions not elsewhere classifiable	326	162	164
777	Immaturity, unqualified	212	71	141
778	Other conditions of fetus or newborn	43	12	31
<u>780-796</u>	<u>SYMPTOMS AND ILL-DEFINED CONDITIONS</u>	<u>1,699</u>	<u>502</u>	<u>1,197</u>
<u>780-789</u>	<u>Symptoms referable to systems or organs</u>	<u>80</u>	<u>29</u>	<u>51</u>
780	Certain symptoms referable to nervous system and special senses	6	2	4
782	Symptoms referable to cardiovascular and lymphatic system	41	16	25
783	Symptoms referable to respiratory system	11	2	9
784	Symptoms referable to upper gastrointestinal tract	8	4	4
785	Symptoms referable to abdomen and lower gastrointestinal tract	3	0	3
786	Symptoms referable to genitourinary system	2	1	1
787	Symptoms referable to limbs and joints	1	1	0
788	Other general symptoms	8	3	5
<u>790-796</u>	<u>Senility and ill-defined diseases</u>	<u>1,619</u>	<u>473</u>	<u>1,146</u>
790	Nervousness and debility	1	0	1
792	Uremia	12	4	8
794	Senility without mention of psychosis	50	6	44
795	Sudden death (cause unknown)	21	11	10
796	Other ill-defined and unknown causes	1,535	452	1,083

Table 5 - Page 13
Deaths - Mississippi - 1969

International Classification Number*	Cause of Death**	Total	White	Non-white
<u>800-999</u>	<u>ACCIDENTS, POISONINGS, AND VIOLENCE (EXTERNAL CAUSE)</u>	<u>2,488</u>	<u>1,423</u>	<u>1,065</u>
<u>800-949</u>	<u>Accidents</u>	<u>2,007</u>	<u>1,179</u>	<u>828</u>
<u>800-807</u>	<u>Railway</u>	<u>19</u>	<u>9</u>	<u>10</u>
800	Involving collision with rolling stock	1	1	0
802	Involving derailment without antecedent collision	2	0	2
804	Fall in, on, or from train	1	0	1
805	Hit by rolling stock	14	7	7
806	Other specified railway accident	1	1	0
<u>810-823</u>	<u>Motor vehicle</u>	<u>903</u>	<u>586</u>	<u>317</u>
<u>(810-819)</u>	<u>(Traffic)</u>	<u>889</u>	<u>577</u>	<u>312</u>
810	Involving collision with train	27	16	11
812	Involving collision with another motor vehicle	362	256	106
813	Involving collision with other vehicle	8	6	2
814	Involving collision with pedestrian	99	49	50
815	Other involving collision	109	75	34
816	Noncollision due to loss of control	171	107	64
817	Noncollision while boarding or alighting	1	1	0
818	Other noncollision	14	9	5
819	Of unspecified nature	98	58	40
<u>(820-823)</u>	<u>(Nontraffic)</u>	<u>14</u>	<u>9</u>	<u>5</u>
820	Involving collision with moving object	11	7	4
823	Of other and unspecified nature	3	2	1
<u>825-827</u>	<u>Other road vehicle</u>	<u>4</u>	<u>3</u>	<u>1</u>
826	Pedal cycle	1	0	1
827	Other nonmotor road vehicle	3	3	0
<u>830-838</u>	<u>Water transport</u>	<u>31</u>	<u>26</u>	<u>5</u>
830	Accident to watercraft causing submersion	17	15	2
832	Other submersion or drowning in water transport	13	10	3
838	Other and unspecified water transport accident	1	1	0
<u>840-845</u>	<u>Air and space transport</u>	<u>21</u>	<u>21</u>	<u>0</u>
840	Accident to powered aircraft at take-off or landing	1	1	0
841	Accident to powered aircraft, other and unspecified	19	19	0
844	Other specified air transport accidents	1	1	0
<u>850-877</u>	<u>Poisoning</u>	<u>24</u>	<u>16</u>	<u>8</u>
<u>(850-859)</u>	<u>(By drugs and medicaments)</u>	<u>6</u>	<u>5</u>	<u>1</u>
852	Primarily systemic and hematologic agents	1	1	0
853	Analgesics and antipyretics	1	0	1
854	Other sedatives and hypnotics	2	2	0
859	Other and unspecified drugs and medicaments	2	2	0
<u>(860-869)</u>	<u>(By other solid and liquid substances)</u>	<u>9</u>	<u>4</u>	<u>5</u>
860	Alcohol	2	1	1
864	Petroleum products and other solvents	3	2	1
865	Pesticides, fertilizers, or plant food	2	0	2
869	Other and unspecified solid and liquid substances	2	1	1
<u>(870-877)</u>	<u>(By gases and vapors)</u>	<u>9</u>	<u>7</u>	<u>2</u>

Table 5 - Page 14
Deaths - Mississippi - 1969

International Classification Number*	Cause of Death**	Total	White	Non-white
870	Gas distributed by pipeline	1	1	0
873	Motor vehicle exhaust gas	6	5	1
874	Carbon monoxide from incomplete combustion of domestic fuels	1	1	0
876	Other gases and vapors	1	0	1
<u>880-887</u>	<u>Falls</u>	<u>169</u>	<u>133</u>	<u>36</u>
880	On or from stairs and steps	3	2	1
881	On or from ladders or scaffolding	3	1	2
882	From or out of building or other structure	13	10	3
883	Into hole or other opening in surface	2	2	0
884	Other from one level to another	23	18	5
885	On same level from slipping, stumbling, or tripping	2	2	0
887	Other and unspecified fall	123	98	25
<u>890-899</u>	<u>Caused by fires and flames</u>	<u>202</u>	<u>41</u>	<u>161</u>
890	Conflagration in private dwelling	138	20	118
891	Conflagration in other building or structure	2	2	0
892	Conflagration not in building or structure	1	1	0
893	Ignition of clothing	20	7	13
894	Ignition of highly inflammable material	8	2	6
895	Controlled fire in private dwelling	15	2	13
898	Other specified fires or flames	7	5	2
899	Unspecified fire	11	2	9
<u>900-909</u>	<u>Due to natural and environmental factors</u>	<u>189</u>	<u>120</u>	<u>69</u>
900	Excessive heat	14	3	11
901	Excessive cold	4	1	3
904	Hunger, thirst, exposure, and neglect	9	3	6
905	Bites and stings of venomous animals and insects	2	1	1
906	Other accidents caused by animals	4	2	2
907	Lightning	4	3	1
908	Cataclysm	152	107	45
<u>910-929</u>	<u>Other accidents</u>	<u>409</u>	<u>202</u>	<u>207</u>
910	Drowning and submersion	113	44	69
911	Inhalation and ingestion of food causing obstruction or suffocation	23	11	12
912	Inhalation and ingestion of other object causing obstruction or suffocation	3	1	2
913	Mechanical suffocation	32	14	18
916	Struck by falling object	35	21	14
917	Struck against or struck by objects	8	3	5
918	Caught in or between objects	13	5	8
919	Overexertion and strenuous movements	1	1	0
920	Caused by cutting or piercing instruments	8	3	5
921	Caused by explosion of pressure vessel	2	2	0
922	Caused by firearm missiles	75	40	35
923	Caused by explosive material	10	5	5
924	Caused by hot substance, corrosive liquid, and steam	6	1	5
925	Caused by electric current	31	18	13
928	Machinery accidents not elsewhere classifiable	35	23	12

Table 5 - Page 15
Deaths - Mississippi - 1969

International Classification Number*	Cause of Death**	Total	White	Non-white
929	Other and unspecified accidents	14	10	4
<u>930-936</u>	<u>Surgical and medical complications and misadventure</u>	<u>16</u>	<u>11</u>	<u>5</u>
930	In operative therapeutic procedures	6	3	3
931	In other and unspecified therapeutic procedures	8	6	2
932	In diagnostic procedures	2	2	0
<u>940-949</u>	<u>Late effects of accidental injury</u>	<u>20</u>	<u>11</u>	<u>9</u>
940	Of motor vehicle accident	3	2	1
941	Of other transport accident	1	0	1
943	Of accidental fall	7	6	1
944	Of accident caused by fire	2	1	1
946	Of other accident	7	2	5
<u>950-959</u>	<u>Suicide and self-inflicted injury</u>	<u>174</u>	<u>143</u>	<u>31</u>
<u>950-952</u>	<u>By poisoning</u>	<u>13</u>	<u>12</u>	<u>1</u>
950	By solid or liquid substances	7	6	1
951	By gases in domestic use	1	1	0
952	By other gases	5	5	0
953	By hanging, strangulation, and suffocation	9	7	2
954	By submersion (drowning)	3	2	1
955	By firearms and explosives	143	117	26
956	By cutting and piercing instruments	4	3	1
958	By other and unspecified means	2	2	0
<u>960-978</u>	<u>Homicide and injury purposely inflicted by other persons; legal intervention</u>	<u>265</u>	<u>76</u>	<u>189</u>
<u>960-969</u>	<u>Homicide and injury purposely inflicted by other persons</u>	<u>258</u>	<u>73</u>	<u>185</u>
960	Fight, brawl, rape	6	5	1
963	Assault by hanging and strangulation	6	2	4
965	Assault by firearms and explosives	184	55	129
966	Assault by cutting and piercing instruments	47	7	40
968	Assault by other and unspecified means	15	4	11
<u>970-978</u>	<u>Legal intervention</u>	<u>7</u>	<u>3</u>	<u>4</u>
970	Injury by firearms	7	3	4
<u>980-989</u>	<u>Injury undetermined whether accidentally or purposely inflicted</u>	<u>41</u>	<u>25</u>	<u>16</u>
980	Poisoning by solid or liquid substances	2	2	0
982	Poisoning by other gases	1	0	1
983	Hanging, strangulation, or suffocation	1	0	1
985	Injury by firearms and explosives	23	15	8
986	Injury by cutting and piercing instruments	4	1	3
988	Injury by other and unspecified means	10	7	3
<u>990-999</u>	<u>Injury resulting from operations of war</u>	<u>1</u>	<u>0</u>	<u>1</u>
991	Injury due to war operations by bullets and fragments	1	0	1

* International Classification numbers are those of the Eighth Revision International Classification of Diseases, Adapted for Use in the United States.

** When cause of death and International Classification number are omitted, no deaths were reported.

TABLE 6
DEATHS
MISSISSIPPI
1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

County	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
<u>STATE</u>	<u>23,419</u>	<u>13,573</u>	<u>9,846</u>	<u>23,865</u>	<u>13,873</u>	<u>9,992</u>	<u>10.8</u>	<u>10.0</u>	<u>12.1</u>
Adams	464	204	260	383	164	219	10.3	8.5	12.2
Alcorn	326	278	48	359	308	51	13.2	12.8	15.8
Amite	67	27	40	170	100	70	12.4	14.7	10.1
Attala	249	152	97	275	167	108	14.0	14.4	13.6
Benton	22	12	10	67	44	23	8.9	10.1	7.3
Bolivar	570	126	444	551	152	399	11.2	8.1	13.0
Calhoun	132	103	29	160	133	27	10.9	12.3	7.1
Carroll	48	16	32	107	60	47	11.4	13.0	9.8
Chickasaw	214	125	89	224	134	90	13.3	12.4	15.0
Choctaw	91	56	35	113	84	29	13.4	13.8	12.2
Claiborne	95	25	70	128	41	87	12.7	16.2	11.5
Clarke	116	74	42	184	118	66	12.2	12.2	12.2
Clay	174	82	92	209	107	102	11.1	11.2	10.9
Coahoma	514	181	333	510	149	361	12.6	10.5	13.8
Copiah	280	142	138	347	177	170	14.0	14.4	13.6
Covington	87	57	30	157	101	56	11.2	10.7	12.2
DeSoto	113	35	78	266	136	130	7.4	5.8	10.3
Forrest	700	493	207	590	380	210	10.2	8.7	14.6
Franklin	80	46	34	106	58	48	13.2	11.8	15.4
George	109	93	16	116	105	11	9.3	9.5	7.6
Greene	57	41	16	101	73	28	11.8	11.0	14.9
Grenada	271	170	101	219	116	103	11.0	10.4	11.8
Hancock	166	147	19	202	179	23	11.6	12.0	9.2
Harrison	1,349	1,072	277	1,210	951	259	9.0	8.6	11.0
Hinds	2,791	1,608	1,183	1,843	1,047	796	8.6	8.0	9.4
Holmes	268	78	190	310	108	202	13.4	14.7	12.8
Humphreys	161	45	116	199	67	132	13.6	13.2	13.9
Issaquena	19	6	13	23	8	15	8.4	7.7	8.8
Itawamba	133	126	7	169	159	10	10.0	10.0	10.4
Jackson	506	379	127	568	430	138	6.4	5.8	9.6
Jasper	124	63	61	192	105	87	12.0	12.3	11.7
Jefferson	84	21	63	122	34	88	13.1	14.8	12.6
Jeff Davis	146	65	81	161	65	96	12.4	10.1	14.8
Jones	771	493	278	596	417	179	10.6	9.8	12.8
Kemper	82	41	41	117	60	57	11.4	13.4	9.9
Lafayette	238	164	74	214	134	80	8.8	7.7	11.6

Table 6 - Page 2

Deaths - Mississippi - 1969

County	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Lamar	103	78	25	179	146	33	11.8	11.1	16.3
Lauderdale	1,093	704	389	814	510	304	12.1	11.0	14.5
Lawrence	87	51	36	128	81	47	11.5	10.7	13.1
Leake	157	109	48	220	153	67	12.9	14.6	10.1
Lee	581	428	153	454	330	124	9.8	9.0	12.9
Leflore	567	228	339	513	180	333	12.2	10.2	13.6
Lincoln	286	192	94	296	187	109	11.3	10.3	13.5
Lowndes	473	271	202	484	252	232	9.7	7.6	14.2
Madison	315	135	180	346	134	212	11.6	12.0	11.4
Marion	210	133	77	249	152	97	10.9	9.6	13.6
Marshall	148	49	99	245	96	149	10.2	10.5	10.0
Monroe	381	243	138	413	267	146	12.1	11.3	14.0
Montgomery	183	111	72	175	103	72	13.5	14.5	12.4
Neshoba	188	143	45	229	173	56	11.0	11.5	9.8
Newton	228	160	68	268	188	80	14.1	14.1	14.2
Noxubee	129	43	86	159	57	102	11.1	11.8	10.8
Oktibbeha	207	100	107	230	118	112	8.0	6.4	11.0
Panola	246	129	117	336	171	165	12.5	13.1	12.0
Pearl River	233	184	49	291	227	64	10.5	10.0	12.5
Perry	67	53	14	86	73	13	9.5	10.9	5.4
Pike	405	215	190	415	220	195	13.1	12.3	14.1
Pontotoc	187	164	23	228	202	26	13.1	14.2	8.4
Prentiss	214	184	30	228	192	36	11.3	10.8	15.3
Quitman	90	25	65	174	74	100	11.0	11.1	10.9
Rankin	486	275	211	343	222	121	7.8	7.0	9.8
Scott	219	149	70	256	166	90	12.0	11.6	12.6
Sharkey	66	20	46	105	34	71	11.7	10.9	12.2
Simpson	258	150	108	279	175	104	14.0	12.8	16.6
Smith	55	47	8	156	139	17	11.5	13.1	5.8
Stone	100	77	23	94	73	21	11.6	11.7	11.2
Sunflower	363	119	244	414	141	273	11.2	10.4	11.6
Tallahatchie	219	70	149	257	90	167	13.3	11.8	14.3
Tate	105	47	58	185	99	86	10.0	10.1	9.8
Tippah	141	120	21	151	127	24	9.5	9.6	9.2
Tishomingo	115	107	8	174	165	9	11.6	11.6	13.4
Tunica	157	32	125	165	39	126	13.9	12.1	14.6
Union	207	178	29	211	183	28	11.0	11.3	9.5
Walthall	105	56	49	127	69	58	10.2	9.3	11.4
Warren	670	318	352	514	252	262	11.4	9.5	14.2
Washington	774	286	488	807	281	526	11.4	8.8	13.6
Wayne	191	130	61	203	135	68	12.2	12.1	12.4
Webster	96	78	18	128	102	26	12.7	13.1	11.5
Wilkinson	160	74	86	145	48	97	13.1	13.4	12.9
Winston	161	104	57	194	119	75	10.5	10.8	10.2
Yalobusha	115	68	47	179	123	56	15.0	17.4	11.6
Yazoo	261	90	171	350	134	216	12.8	10.6	14.8
Out of state	1,273	931	342	827	631	196			

Table 6 - Page 3
Deaths - Mississippi - 1969

City**	Place of Occurrence			Place of Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Biloxi	431	375	56	381	318	63	7.8	7.7	9.0
Brookhaven	245	176	69	118	66	52	11.0	9.9	12.8
Canton	224	111	113	158	69	89	15.0	15.2	15.0
Clarksdale	401	170	231	263	91	172	12.1	9.2	14.5
Cleveland	179	75	104	131	50	81	9.8	6.0	16.1
Columbus	381	219	162	298	140	158	11.6	8.7	16.2
Corinth	68	56	12	179	142	37	15.4	15.2	16.5
Greenville	591	249	342	502	201	301	12.7	10.7	14.5
Greenwood	440	191	249	288	121	167	12.8	10.9	14.8
Gulfport	551	415	136	394	277	117	9.6	8.4	15.3
Hattiesburg	643	450	193	378	208	170	9.9	7.7	15.2
Jackson	2,275	1,272	1,003	1,428	848	580	9.3	9.2	9.4
Laurel	647	391	256	289	152	137	12.0	10.0	15.3
McComb	242	138	104	177	110	67	14.8	13.8	16.8
Meridian	1,013	660	353	588	362	226	13.0	12.1	14.9
Moss Point	33	17	16	130	68	62	6.7	6.2	7.4
Natchez	429	193	236	265	113	152	13.4	11.6	15.2
Oxford	184	132	52	83	61	22	6.0	5.4	8.7
Pascagoula	375	285	90	202	160	42	7.4	6.8	11.7
Picayune	156	116	40	110	64	46	10.5	9.2	13.0
Starkville	128	59	69	87	54	33	7.6	6.4	11.4
Tupelo	477	350	127	184	123	61	9.0	7.3	17.0
Vicksburg	602	287	315	369	180	189	14.5	14.0	14.9
Yazoo City	155	61	94	160	59	101	14.8	13.7	15.6

* Deaths per 1,000 population.

** Cities having 10,000 population and over according to 1970 census.

Table 7 - Page 2
Deaths - Mississippi - 1969

Inter-national Classification Number#	Cause of Death	Total		Race	Total by Race	Sex	Total by Race and Sex	Age												
		Number	Rate*					Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 & Over	Unknown		
070	Infectious hepatitis	16	0.7	W	8	M	4	0	0	0	0	1	0	0	1	1	1	0		
						F	4	0	1	0	0	1	0	0	1	1	0			
				C	8	M	2	0	0	0	1	0	1	0	0	0	0	0		
				F	6	0	0	1	0	0	2	1	0	2	0	0	0			
090-097	Syphilis and its sequelae	12	0.5	W	1	M	1	0	0	0	0	0	0	0	1	0	0	0		
						F	0	0	0	0	0	0	0	0	0	0	0	0		
				C	11	M	6	0	0	0	0	0	0	0	0	1	5	0		
				F	5	0	0	0	0	0	0	0	2	1	2	0	0			
140-209	Malignant neoplasms	3,295	148.6	W	2,146	M	1,214	2	4	6	3	14	17	41	146	327	654	0		
						F	932	0	2	2	1	9	18	47	134	205	514	0		
				C	1,149	M	622	0	3	5	3	2	13	23	66	149	358	0		
						F	527	0	4	1	3	5	5	36	84	110	279	0		
								F		0	0	0	0	0	0	0	0	0	0	0
								F		0	0	0	0	0	0	0	0	0	0	0
210-239	Benign and unspecified neoplasms	37	1.7	W	20	M	7	0	0	0	0	0	1	1	1	4	0			
						F	13	0	0	0	0	1	5	2	1	4	0			
				C	17	M	7	1	0	0	0	1	1	1	2	1	0			
				F	10	0	0	0	0	0	1	5	4	0	0					
240-279	Endocrine, nutritional, and metabolic diseases	550	24.8	W	258	M	98	2	0	0	0	0	4	3	16	25	48	0		
						F	160	1	0	1	0	6	5	4	13	26	104	0		
				C	292	M	99	2	1	0	0	4	0	8	6	29	49	0		
				F	193	4	0	1	2	2	4	6	15	41	116	2				
250	Diabetes mellitus	477	21.5	W	223	M	83	0	0	0	0	0	4	2	16	19	42	0		
						F	140	0	0	0	0	4	5	2	12	24	93	0		
				C	254	M	83	0	0	0	0	3	0	8	6	25	41	0		
				F	171	0	0	0	1	2	3	5	12	38	108	2				
260-269	Avitaminoses and other nutritional deficiency	22	1.0	W	7	M	4	0	0	0	0	0	0	0	0	4	0			
						F	3	0	0	0	0	0	0	0	0	3	0			
				C	15	M	7	2	0	0	0	0	0	0	0	5	0			
				F	8	3	0	0	0	0	0	0	0	5	0					

Table 7 - Page 4
Deaths - Mississippi - 1969

International Classification Number#	Cause of Death	Total		Race	Total by Race	Sex	Total by Race and Sex	Age										Unknown
								Number	Rate*	Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	
345	Epilepsy	34	1.5	W	14	M	6	0	0	0	0	1	2	0	1	2	0	0
						F	8	0	0	0	0	2	2	1	0	1	0	0
				C	20	M	12	0	0	3	1	1	1	3	2	0	1	0
						F	8	0	0	0	1	2	2	1	0	1	1	0
390-458	<u>Diseases of the circulatory system</u>	<u>11,864</u>	<u>535.2</u>	W	<u>7,390</u>	M	<u>4,229</u>	<u>0</u>	<u>1</u>	<u>0</u>	<u>1</u>	<u>3</u>	<u>14</u>	<u>111</u>	<u>367</u>	<u>850</u>	<u>2,882</u>	<u>0</u>
						F	<u>3,161</u>	<u>4</u>	<u>3</u>	<u>1</u>	<u>6</u>	<u>3</u>	<u>26</u>	<u>92</u>	<u>288</u>	<u>2,737</u>	<u>0</u>	<u>0</u>
				C	<u>4,474</u>	M	<u>2,284</u>	<u>5</u>	<u>4</u>	<u>2</u>	<u>3</u>	<u>10</u>	<u>20</u>	<u>69</u>	<u>201</u>	<u>403</u>	<u>1,565</u>	<u>2</u>
						F	<u>2,190</u>	<u>3</u>	<u>3</u>	<u>1</u>	<u>3</u>	<u>10</u>	<u>28</u>	<u>74</u>	<u>172</u>	<u>359</u>	<u>1,535</u>	<u>2</u>
390-398, 402, 404-429	Diseases of the heart	7,886	355.7	W	5,142	M	3,099	0	0	0	0	1	9	95	319	698	1,977	0
						F	2,043	3	3	0	0	3	0	14	61	208	1,751	0
				C	2,744	M	1,456	4	3	1	2	7	12	41	125	285	974	2
						F	1,288	3	2	1	3	5	11	42	104	210	905	2
400,401, 403	Hypertension	161	7.3	W	52	M	29	0	0	0	0	1	2	1	2	1	22	0
						F	23	0	0	0	0	0	0	0	3	1	19	0
				C	109	M	61	0	0	0	0	0	2	6	14	6	33	0
						F	48	0	0	0	0	1	2	5	5	12	23	0
430-438	Cerebrovascular disease	3,058	137.9	W	1,681	M	835	0	1	0	1	0	2	13	33	113	672	0
						F	846	1	0	1	0	1	2	9	20	66	746	0
				C	1,377	M	656	1	1	0	1	2	5	21	51	98	476	0
						F	721	0	1	0	0	2	11	22	53	118	514	0
440	Arteriosclerosis	331	14.9	W	232	M	104	0	0	0	0	0	0	0	0	6	98	0
						F	128	0	0	0	0	0	0	0	0	2	126	0
				C	99	M	44	0	0	0	0	0	0	0	3	4	37	0
						F	55	0	0	0	0	0	0	0	0	6	49	0
460-519	<u>Diseases of the respiratory system</u>	<u>1,183</u>	<u>53.4</u>	W	<u>711</u>	M	<u>440</u>	<u>26</u>	<u>5</u>	<u>1</u>	<u>0</u>	<u>2</u>	<u>2</u>	<u>5</u>	<u>27</u>	<u>85</u>	<u>287</u>	<u>0</u>
						F	<u>271</u>	<u>15</u>	<u>5</u>	<u>0</u>	<u>3</u>	<u>2</u>	<u>1</u>	<u>7</u>	<u>19</u>	<u>31</u>	<u>188</u>	<u>0</u>
				C	<u>472</u>	M	<u>288</u>	<u>68</u>	<u>17</u>	<u>5</u>	<u>5</u>	<u>4</u>	<u>3</u>	<u>6</u>	<u>24</u>	<u>34</u>	<u>122</u>	<u>0</u>
						F	<u>184</u>	<u>65</u>	<u>9</u>	<u>3</u>	<u>0</u>	<u>3</u>	<u>2</u>	<u>4</u>	<u>11</u>	<u>17</u>	<u>70</u>	<u>0</u>

Table 7 - Page 5
Deaths - Mississippi - 1969

Inter-national Classification Number#	Cause of Death	Total		Race	Total by Race	Sex	Total by Race and Sex	Age										Unknown
								Number	Rate*	Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	
470-474	Influenza	84	3.8	W	M	24	0	0	0	0	0	0	0	0	2	4	18	0
					F	18	0	0	0	0	1	0	0	2	0	15	0	
				C	M	42	3	1	1	2	0	0	0	1	6	10	0	
					F	18	3	0	0	0	1	0	0	3	0	11	0	
480-486	Pneumonia	642	29.0	W	M	181	22	4	1	0	2	2	3	12	23	112	0	
					F	162	13	4	0	3	1	1	3	6	17	114	0	
				C	M	299	59	12	3	2	1	0	0	9	17	77	0	
					F	119	47	6	2	0	1	1	3	5	9	45	0	
490,491	Chronic and unqualified bronchitis	62	2.8	W	M	32	0	1	0	0	0	0	0	2	9	20	0	
					F	17	0	0	0	0	0	1	4	1	11	0		
				C	M	13	0	0	0	0	0	0	1	1	6	0		
					F	5	1	2	0	0	0	0	0	2	0	0		
492	Emphysema	206	9.3	W	M	150	0	0	0	0	0	0	1	9	34	106	0	
					F	29	0	0	0	0	0	1	3	6	19	0		
				C	M	27	0	0	0	0	0	1	7	2	16	0		
					F	1	0	0	0	0	0	0	0	0	0	1	0	
493	Asthma	28	1.3	W	M	4	0	0	0	0	0	0	0	0	1	3	0	
					F	7	0	0	0	0	0	2	3	2	0	0		
				C	M	17	0	0	0	0	0	1	1	3	3	0		
					F	9	0	0	0	0	1	1	0	1	5	1	0	
<u>520-577</u>	<u>Diseases of the digestive system</u>	<u>548</u>	<u>24.7</u>	W	M	<u>212</u>	<u>2</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>2</u>	<u>6</u>	<u>16</u>	<u>33</u>	<u>49</u>	<u>103</u>	<u>0</u>	
					F	<u>168</u>	<u>4</u>	<u>1</u>	<u>0</u>	<u>0</u>	<u>3</u>	<u>1</u>	<u>8</u>	<u>18</u>	<u>33</u>	<u>100</u>	<u>0</u>	
				C	M	<u>168</u>	<u>6</u>	<u>1</u>	<u>0</u>	<u>3</u>	<u>4</u>	<u>4</u>	<u>10</u>	<u>18</u>	<u>18</u>	<u>47</u>	<u>0</u>	
					F	<u>72</u>	<u>1</u>	<u>0</u>	<u>1</u>	<u>0</u>	<u>3</u>	<u>3</u>	<u>7</u>	<u>12</u>	<u>9</u>	<u>36</u>	<u>0</u>	
531-533	Peptic ulcer	87	3.9	W	M	39	0	0	0	0	0	1	3	3	13	19	0	
					F	20	0	0	0	0	0	2	2	2	14	0		
				C	M	28	0	0	0	0	0	0	1	3	10	0		
					F	14	0	0	1	0	0	2	1	2	8	0		

Table 7 - Page 6
Deaths - Mississippi - 1969

International Classification Number#	Cause of Death	Total		Race	Total by Race	Sex	Total by Race and Sex	Age										Unknown		
		Number	Rate*					Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 & Over			
550-553, 560	Hernia and intestinal obstruction	68	3.1	W	35	M	8	1	0	0	0	1	0	0	1	0	5	0		
						F	27	2	0	0	0	1	0	0	3	4	17	0		
					C	33	M	24	5	1	0	0	0	0	0	0	3	3	12	0
							F	9	0	0	0	0	1	0	0	0	2	6	6	0
571	Cirrhosis of liver	140	6.3	W	113	M	80	0	0	0	0	0	2	9	22	21	26	0		
						F	33	0	0	0	0	0	2	8	9	14	0			
					C	27	M	14	0	0	0	1	2	1	1	7	1	0		
							F	13	0	0	0	0	0	1	2	4	3	3	0	
<u>580-629</u>	<u>Diseases of the genitourinary system</u>	<u>466</u>	<u>21.0</u>	W	<u>246</u>	M	<u>131</u>	0	0	0	1	1	2	2	<u>13</u>	<u>13</u>	<u>99</u>	0		
						F	<u>115</u>	0	0	0	0	1	1	9	7	<u>11</u>	<u>84</u>	0		
					C	<u>220</u>	M	<u>126</u>	0	0	1	0	5	8	7	<u>19</u>	<u>85</u>	0		
							F	<u>94</u>	0	0	1	1	4	4	9	<u>19</u>	<u>51</u>	0		
580-584	Nephritis and nephrosis	167	7.5	W	65	M	41	0	0	0	1	1	1	2	7	5	24	0		
						F	24	0	0	1	0	1	0	1	4	3	14	0		
					C	102	M	55	0	1	1	0	2	4	3	10	34	0		
							F	47	0	0	1	1	0	0	6	13	21	0		
590	Infections of kidney	171	7.7	W	106	M	47	0	0	0	0	0	1	0	2	5	39	0		
						F	59	0	0	0	0	0	2	1	5	51	0			
					C	65	M	33	0	0	0	0	0	2	2	4	23	0		
							F	32	0	0	0	0	1	3	1	4	23	0		
<u>630-678</u>	<u>Complications of pregnancy, childbirth, and the puerperium</u>	<u>23</u>	<u>4.9</u>	W	<u>3</u>	M	<u>0</u>	0	0	0	0	0	0	0	0	0	0	0		
						F	<u>3</u>	0	0	0	0	1	2	0	0	0	0	0		
					C	<u>20</u>	M	<u>0</u>	0	0	0	0	0	0	0	0	0	0	0	
							F	<u>20</u>	0	0	0	0	10	6	4	0	0	0	0	
<u>680-709</u>	<u>Diseases of the skin and subcutaneous tissue</u>	<u>34</u>	<u>1.5</u>	W	<u>17</u>	M	<u>7</u>	1	0	0	0	0	0	1	0	2	3	0		
						F	<u>10</u>	1	0	0	0	0	0	0	1	0	8	0		
					C	<u>17</u>	M	<u>5</u>	1	0	0	0	0	2	2	0	0	2	3	0
							F	<u>12</u>	1	0	1	0	0	1	0	4	1	3	0	

Table 7 - Page 7
Deaths - Mississippi - 1969

International Classification Number#	Cause of Death	Total		Race	Total by Race	Sex	Total by Race and Sex	Age										
		Number	Rate*					Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 & Over	Unknown
710-738	<u>Diseases of the musculoskeletal system and connective tissue</u>	59	2.7	W	41	M	17	0	1	0	0	0	0	0	5	7	4	0
						F	24	0	0	0	2	0	1	3	16	0		
				C	18	M	7	0	0	0	0	0	0	3	4	0		
						F	11	0	0	0	2	1	2	1	1	4	0	
740-759	<u>Congenital anomalies</u>	218	9.8	W	127	M	67	41	5	6	2	4	5	2	2	0	0	0
						F	60	44	1	0	4	1	1	3	5	1	0	
				C	91	M	54	40	8	1	1	1	2	0	0	0	0	
						F	37	25	6	0	1	2	0	2	1	0	0	
760-778	<u>Certain causes of mortality in early infancy</u>	766	16.2	W	329	M	191	191	0	0	0	0	0	0	0	0	0	0
						F	138	138	0	0	0	0	0	0	0	0	0	
				C	437	M	252	252	0	0	0	0	0	0	0	0	0	
						F	185	185	0	0	0	0	0	0	0	0	0	
780-796	<u>Symptoms and ill-defined conditions</u>	1,699	76.6	W	502	M	326	12	1	0	0	1	5	16	52	88	151	0
						F	176	12	0	0	0	2	1	2	18	24	117	0
				C	1,197	M	647	90	9	1	4	4	13	32	60	123	311	0
						F	550	77	6	1	1	4	8	26	52	74	301	0
800-999	<u>Accidents, poisonings, and violence</u>	2,488	112.2	W	1,423	M	1,012	7	21	24	29	232	138	125	136	124	176	0
						F	411	6	13	16	14	58	33	42	37	39	150	3
				C	1,065	M	750	20	36	29	38	168	115	97	77	68	100	2
						F	315	22	33	11	21	44	33	26	25	20	79	1
810-823	<u>Motor vehicle accidents</u>	903	40.7	W	586	M	421	2	7	20	11	136	72	32	50	38	53	0
						F	165	4	8	7	7	35	18	17	14	22	33	0
				C	317	M	238	1	11	13	8	58	32	35	28	25	26	1
						F	79	3	5	1	8	23	11	8	5	4	11	0
800-807, 825-949	<u>All other accidents</u>	1,104	49.8	W	593	M	401	5	12	4	17	71	40	50	50	56	96	0
						F	192	2	5	9	6	12	8	13	15	8	111	3
				C	511	M	329	19	25	16	30	68	32	29	23	27	59	1
						F	182	18	28	10	11	10	7	7	14	11	65	1

Table 7 - Page 8
Deaths - Mississippi - 1969

Inter- national Classi- fication Number#	Cause of Death	Total		Race	Total by Race	Sex	Total by Race and Sex	Age										65 and Over	Unknown
		Number	Rate*					Under 1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64			
950-959	Suicide	174	7.8	W	143	M	112	0	0	0	0	14	12	22	21	23	20	0	
						F	31	0	0	0	0	4	3	9	4	6	5	0	
				C	31	M	21	0	0	0	0	6	5	1	4	3	2	0	
						F	10	0	0	0	1	3	0	2	1	1	2	0	
960-978	Homicide	265	12.0	W	76	M	57	0	1	0	0	8	11	16	10	5	6	0	
						F	19	0	0	0	1	6	4	2	3	3	0	0	
				C	189	M	151	0	0	0	0	35	40	29	21	13	13	0	
						F	38	0	0	0	1	8	14	6	5	3	1	0	

Numbers are those of the Eighth Revision International Classification of Diseases, Adapted for Use in the United States which went into use January 1, 1968.

* Total deaths per 1,000 population. Deaths from specific causes per 100,000 population, except deaths from maternal causes (classification numbers 630-678) are per 10,000 live births and deaths from certain causes of mortality in early infancy (classification numbers 760-778) are 1,000 live births.

Table 8 - Page 2
Deaths Under 1 Year - Mississippi - 1969

Cause of Death and International Classification Number#	Total		Race	Total by Race	Days of Age					Months of Age											
	Number	Rate*			Under 1 Day	1-6	7-13	14-20	21-27	Under 28 Days	1	2	3	4	5	6	7	8	9	10	11
Bronchitis and bronchiolitis (466, 490, 491)	11	0.2	W	3	0	0	0	0	0	0	2	0	1	0	0	0	0	0	0	0	0
			C	8	0	0	0	0	0	0	0	2	2	3	0	1	0	0	0	0	0
Influenza and pneumonia (470-486)	147	3.1	W	35	2	9	2	0	0	13	2	5	6	4	0	0	1	2	1	1	0
			C	112	4	8	3	6	3	24	22	15	14	7	10	4	4	6	2	1	3
Other diseases of the respiratory system (492-519)	8	0.2	W	2	1	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0
			C	6	0	0	0	0	1	1	0	0	2	1	1	0	0	1	0	0	0
Hernia and intestinal obstruction (550-553, 560)	8	0.2	W	3	2	0	0	0	1	3	0	0	0	0	0	0	0	0	0	0	0
			C	5	0	0	1	0	1	2	1	1	0	0	0	0	0	0	0	0	1
Congenital anomalies (740-759)	150	3.2	W	85	31	22	7	5	3	68	3	5	4	2	0	0	2	0	0	0	1
			C	65	16	14	4	0	3	37	10	2	5	1	4	1	0	2	2	1	0
Maternal conditions unrelated to pregnancy (760, 761)	7	0.1	W	2	2	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0
			C	5	4	1	0	0	0	5	0	0	0	0	0	0	0	0	0	0	0
Toxemia of pregnancy (762)	5	0.1	W	1	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0
			C	4	1	3	0	0	0	4	0	0	0	0	0	0	0	0	0	0	0
Maternal antepartum and intrapartum infection (763)	7	0.1	W	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
			C	7	6	1	0	0	0	7	0	0	0	0	0	0	0	0	0	0	0
Difficult labor (764-768)	19	0.4	W	11	7	4	0	0	0	11	0	0	0	0	0	0	0	0	0	0	0
			C	8	5	3	0	0	0	8	0	0	0	0	0	0	0	0	0	0	0
Other complications of pregnancy and childbirth (769)	63	1.3	W	28	25	2	1	0	0	28	0	0	0	0	0	0	0	0	0	0	0
			C	35	28	6	1	0	0	35	0	0	0	0	0	0	0	0	0	0	0
Conditions of placenta (770)	44	0.9	W	21	13	8	0	0	0	21	0	0	0	0	0	0	0	0	0	0	0
			C	23	15	8	0	0	0	23	0	0	0	0	0	0	0	0	0	0	0

Table 8 - Page 3
Deaths Under 1 Year - Mississippi - 1969

Cause of Death and International Classification Number#	Total		Race	Total by Race	Days of Age					Months of Age											
	Number	Rate*			Under 1 Day	1-6	7-13	14-20	21-27	Under 28 Days	1	2	3	4	5	6	7	8	9	10	11
Conditions of umbilical cord (771)	5	0.1	W	1	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0
			C	4	4	0	0	0	0	4	0	0	0	0	0	0	0	0	0	0	0
Birth injury without mention of cause (772)	19	0.4	W	6	1	4	1	0	0	6	0	0	0	0	0	0	0	0	0	0	0
			C	13	4	6	1	0	2	13	0	0	0	0	0	0	0	0	0	0	0
Hemolytic disease of newborn (774, 775)	16	0.3	W	14	7	6	0	0	1	14	0	0	0	0	0	0	0	0	0	0	0
			C	2	1	0	1	0	0	2	0	0	0	0	0	0	0	0	0	0	0
Hyaline membrane disease (776.1)	97	2.0	W	62	16	43	3	0	0	62	0	0	0	0	0	0	0	0	0	0	0
			C	35	11	23	1	0	0	35	0	0	0	0	0	0	0	0	0	0	0
Respiratory distress syndrome (776.2)	87	1.8	W	35	17	15	2	0	1	35	0	0	0	0	0	0	0	0	0	0	0
			C	52	22	26	1	1	2	52	0	0	0	0	0	0	0	0	0	0	0
Asphyxia of newborn, unspecified (776.9)	127	2.7	W	61	40	21	0	0	0	61	0	0	0	0	0	0	0	0	0	0	0
			C	66	40	21	3	2	0	66	0	0	0	0	0	0	0	0	0	0	0
Other anoxic and hypoxic conditions not elsewhere classifiable (776.0, 776.3, 776.4)	15	0.3	W	4	3	1	0	0	0	4	0	0	0	0	0	0	0	0	0	0	0
			C	11	3	7	1	0	0	11	0	0	0	0	0	0	0	0	0	0	0
Immaturity, unqualified (777)	212	4.5	W	71	45	21	3	1	1	71	0	0	0	0	0	0	0	0	0	0	0
			C	141	93	39	4	3	0	139	2	0	0	0	0	0	0	0	0	0	0
Hemorrhagic disease of newborn (778.2)	9	0.2	W	2	1	1	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0
			C	7	0	6	1	0	0	7	0	0	0	0	0	0	0	0	0	0	0
Other conditions of newborn (778.0, 778.1, 778.3, 778.9)	34	0.7	W	10	8	1	1	0	0	10	0	0	0	0	0	0	0	0	0	0	0
			C	24	13	8	2	0	1	24	0	0	0	0	0	0	0	0	0	0	0
Symptoms and ill-defined conditions (780-796)	191	4.0	W	24	2	4	0	1	0	7	2	4	4	1	2	2	0	0	1	1	0
			C	167	31	18	10	4	5	68	33	21	19	8	5	6	0	2	2	1	2

Table 8 - Page 4
 Deaths Under 1 Year - Mississippi - 1969

Cause of Death and International Classi- fication Number#	Total		Race	Total by Race	Days of Age					Months of Age											
	Number	Rate*			Under 1 Day	1- 6	7- 13	14- 20	21- 27	Under 28 Days	1	2	3	4	5	6	7	8	9	10	11
Accidents (800-949)	54	1.1	W	13	0	0	1	0	3	4	0	1	0	1	1	1	1	0	3	1	0
			C	41	1	0	1	4	0	6	6	5	8	3	4	2	1	1	2	1	2
All other causes	32	0.7	W	11	0	2	0	1	0	3	3	0	0	1	0	0	1	0	1	1	1
			C	21	2	3	0	2	1	8	7	1	0	1	0	3	0	0	1	0	0

Numbers are those of the Eighth Revision International Classification of Diseases, Adapted for Use in the United States which went into use January 1, 1968.

* Deaths per 1,000 live births.

** Rate less than 0.06.

TABLE 9
MARRIAGES AND DIVORCES
MISSISSIPPI
1969

Marriages by Place of Licensure, by Place of Occurrence, and by Place of Residence of Bride, by Race of Bride;
Divorces by Place of Occurrence, by Race

County	Marriages									Divorces		
	Place of Licensure			Place of Occurrence			Residence of Bride			Total	White	Non-white
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white			
<u>STATE</u>	<u>25,956</u>	<u>17,582</u>	<u>8,374</u>	<u>25,956</u>	<u>17,582</u>	<u>8,374</u>	<u>22,035</u>	<u>14,367</u>	<u>7,668</u>	<u>7,506</u>	<u>5,616</u>	<u>1,890</u>
Adams	434	267	167	458	299	159	369	213	156	167	106	61
Alcorn	515	462	53	535	477	58	287	255	32	120	111	9
Amite	145	87	58	141	87	54	145	90	55	15	8	7
Attala	178	124	54	169	111	58	177	111	66	48	40	8
Benton	71	42	29	86	61	25	68	41	27	10	4	6
Bolivar	495	252	243	475	234	241	474	228	246	132	74	58
Calhoun	157	123	34	194	155	39	167	129	38	34	27	7
Carroll	61	38	23	71	50	21	66	41	25	11	7	4
Chickasaw	262	170	92	202	134	68	197	133	64	48	40	8
Choctaw	91	68	23	81	65	16	52	38	14	22	18	4
Claiborne	84	24	60	87	29	58	75	25	50	24	6	18
Clarke	158	88	70	148	88	60	156	92	64	41	28	13
Clay	204	128	76	195	125	70	175	108	67	58	37	21
Coahoma	359	158	201	359	163	196	343	149	194	140	64	76
Copiah	210	98	112	238	120	118	229	115	114	62	36	26
Covington	116	81	35	129	96	33	127	88	39	26	21	5
DeSoto	988	594	394	979	579	400	225	149	76	144	123	21
Forrest	685	538	147	657	516	141	614	466	148	288	220	68
Franklin	73	54	19	73	55	18	75	53	22	22	13	9
George	214	200	14	199	185	14	115	107	8	37	31	6
Greene	67	49	18	91	65	26	76	51	25	37	32	5
Grenada	276	163	113	258	145	113	242	138	104	80	54	26
Hancock	326	292	34	315	281	34	135	123	12	72	64	8
Harrison	1,983	1,696	287	2,019	1,720	299	1,665	1,374	291	766	680	86

Table 9 - Page 2
 Marriages and Divorces - Mississippi - 1969

County	Marriages									Divorces		
	Place of Licensure			Place of Occurrence			Residence of Bride			Total	White	Non-white
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white			
Hinds	2,511	1,576	935	2,276	1,380	896	2,340	1,488	852	736	547	189
Holmes	224	71	153	236	73	163	251	79	172	60	20	40
Humphreys	158	64	94	162	67	95	149	52	97	25	13	12
Issaquena	11	9	2	8	1	7	7	1	6	8	2	6
Itawamba	167	154	13	177	166	11	109	102	7	53	45	8
Jackson	983	816	167	985	820	165	823	693	130	362	314	48
Jasper	131	69	62	156	82	74	158	76	82	25	13	12
Jefferson	86	40	46	59	17	42	62	17	45	14	1	13
Jeff Davis	127	70	57	109	59	50	101	56	45	25	16	9
Jones	621	502	119	604	489	115	575	469	106	234	210	24
Kemper	85	39	46	87	38	49	70	30	40	26	13	13
Lafayette	179	125	54	183	125	58	147	96	51	54	43	11
Lamar	149	121	28	179	150	29	156	126	30	52	48	4
Lauderdale	679	464	215	656	448	208	624	413	211	261	204	57
Lawrence	83	49	34	94	56	38	130	80	50	24	16	8
Leake	164	105	59	156	96	60	167	104	63	55	39	16
Lee	498	405	93	501	403	98	579	458	121	197	173	24
Leflore	456	204	252	457	198	259	448	206	242	116	83	33
Lincoln	256	183	73	270	193	77	219	146	73	62	44	18
Lowndes	468	308	160	471	307	164	431	266	165	195	152	43
Madison	303	123	180	317	140	177	296	122	174	101	40	61
Marion	261	166	95	246	151	95	239	151	88	95	76	19
Marshall	392	178	214	377	164	213	230	94	136	69	44	25
Monroe	330	239	91	347	246	101	325	231	94	121	97	24
Montgomery	132	72	60	138	84	54	132	74	58	37	25	12
Neshoba	260	172	88	277	188	89	221	144	77	101	77	24
Newton	151	103	48	179	130	49	197	142	55	31	24	7
Noxubee	111	48	63	120	52	68	118	51	67	21	11	10
Oktibbeha	287	171	116	263	156	107	258	153	105	55	39	16
Panola	254	134	120	268	148	120	240	116	124	63	37	26

Table 9 - Page 3
 Marriages and Divorces - Mississippi - 1969

County	Marriages									Divorces		
	Place of Licensure			Place of Occurrence			Residence of Bride					
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Pearl River	408	360	48	417	369	48	301	261	40	158	143	15
Perry	108	86	22	103	84	19	104	82	22	15	13	2
Pike	344	216	128	362	233	129	320	197	123	96	62	34
Pontotoc	177	150	27	176	152	24	161	139	22	56	47	9
Prentiss	249	224	25	243	224	19	196	185	11	72	70	2
Quitman	209	103	106	202	94	108	184	86	98	45	31	14
Rankin	335	256	79	418	329	89	354	268	86	92	78	14
Scott	222	147	75	212	133	79	210	137	73	47	29	18
Sharkey	108	48	60	104	51	53	94	47	47	31	15	16
Simpson	276	184	92	286	198	88	238	153	85	59	41	18
Smith	89	65	24	127	97	30	93	64	29	39	29	10
Stone	145	109	36	121	90	31	128	98	30	39	30	9
Sunflower	312	126	186	351	141	210	355	142	213	61	34	27
Tallahatchie	207	88	119	206	92	114	180	71	109	44	26	18
Tate	244	169	75	221	152	69	166	94	72	46	30	16
Tippah	154	136	18	153	133	20	161	141	20	30	28	2
Tishomingo	395	387	8	378	372	6	156	153	3	51	49	2
Tunica	143	65	78	134	60	74	108	41	67	24	11	13
Union	204	181	23	212	189	23	170	149	21	40	35	5
Walthall	134	70	64	141	74	67	104	45	59	26	17	9
Warren	536	329	207	532	324	208	484	291	193	184	131	53
Washington	862	488	374	877	496	381	793	428	365	260	157	103
Wayne	188	119	69	182	116	66	124	83	41	78	58	20
Webster	98	79	19	108	88	20	102	79	23	22	16	6
Wilkinson	578	491	87	586	483	103	104	32	72	38	20	18
Winston	164	99	65	165	99	66	170	101	69	61	40	21
Yalobusha	129	89	40	132	86	46	151	102	49	28	19	9
Yazoo	269	142	127	278	154	124	268	145	123	82	47	35
Not reported				12	2	10						
Out of state							3,921	3,215	706			

TABLE 10
DIVORCES
MISSISSIPPI
1969

By Cause of Divorce, by Race

Cause	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>ALL CAUSES</u>	<u>7,506</u>	<u>5,616</u>	<u>1,890</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
Cruel and inhuman treatment	4,789	4,164	625	63.8	74.1	33.1
Desertion	2,041	923	1,118	27.2	16.4	59.2
Adultery	332	247	85	4.4	4.4	4.5
Drunkenness	216	179	37	2.9	3.2	2.0
Penitentiary	34	28	6	0.4	0.5	0.3
Insanity	7	6	1	0.1	0.1	*
Annulment	41	35	6	0.5	0.6	0.3
Bigamy	16	10	6	0.2	0.2	0.3
Miscellaneous and other	15	13	2	0.2	0.2	0.1
Not stated	15	11	4	0.2	0.2	0.2

* Less than 0.06 percent.

By Number of Minor Children, by Race

Number of Minor Children	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>TOTAL DIVORCES</u>	<u>7,506</u>	<u>5,616</u>	<u>1,890</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
None	3,252	2,369	883	43.3	42.2	46.7
One	1,731	1,362	369	23.1	24.2	19.5
Two	1,195	930	265	15.9	16.6	14.0
Three or more	1,241	897	344	16.5	16.0	18.2
Not stated	87	58	29	1.2	1.0	1.5

By Number of Years Married, by Race

Number of Years Married	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>TOTAL DIVORCES</u>	<u>7,506</u>	<u>5,616</u>	<u>1,890</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
<u>Under five</u>	<u>3,225</u>	<u>2,754</u>	<u>471</u>	<u>43.0</u>	<u>49.0</u>	<u>24.9</u>
Under one	625	575	50	8.3	10.2	2.6
One	808	720	88	10.8	12.8	4.6
Two	728	605	123	9.7	10.8	6.5
Three	583	487	96	7.8	8.7	5.1
Four	481	367	114	6.4	6.5	6.0
Five - nine	1,526	1,167	359	20.3	20.8	19.0
Ten - nineteen	1,568	1,067	501	20.9	19.0	26.5
Twenty and over	1,134	594	540	15.1	10.6	28.6
Not stated	53	34	19	0.7	0.6	1.0

TABLE 11
LIVE BIRTHS
MISSISSIPPI
1913-1969

By Year, by Race, with Rates, Occurrence Data 1913-1943,
Residence Data 1944-1969

Year	Number			Rate*		
	Total	White	Non- white	Total	White	Non- white
1913	41,103	21,874	19,229	21.9	26.6	18.2
1914	40,623	21,360	19,263	21.4	25.7	18.0
1915	39,484	20,999	18,485	20.5	23.8	17.1
1916	39,033	20,130	18,903	20.0	23.6	17.2
1917	37,141	19,056	18,085	18.8	22.0	16.3
1918	39,391	20,005	19,386	19.7	22.8	17.2
1919	42,491	22,028	20,463	21.0	24.9	18.0
1920	43,743	22,438	21,305	24.4	26.3	22.7
1921	46,482	23,703	22,779	25.6	27.3	24.1
1922	43,723	21,286	22,437	23.8	24.1	23.6
1923	43,266	21,140	22,126	23.3	23.6	23.1
1924	43,960	22,180	21,780	23.4	24.3	22.5
1925	45,563	22,265	23,298	24.0	24.1	23.9
1926	50,449	24,600	25,849	26.2	26.2	26.3
1927	49,204	24,163	25,041	25.3	25.3	25.3
1928	48,126	23,642	24,484	24.5	24.4	24.5
1929	45,570	22,105	23,465	22.9	22.5	23.3
1930	47,982	23,192	24,790	23.8	23.2	24.0
1931	45,123	21,637	23,486	22.2	21.4	22.9
1932	46,481	21,883	24,598	22.6	21.4	23.9
1933	44,184	20,407	23,777	21.3	19.7	22.9
1934	47,905	22,342	25,563	23.0	21.4	24.6
1935	48,291	21,972	26,319	23.0	20.8	25.1
1936	49,261	22,194	27,067	23.3	20.8	25.7
1937	52,170	23,285	28,885	24.4	21.6	27.3
1938	53,729	24,046	29,683	25.0	22.1	27.8
1939	52,288	23,505	28,783	24.1	21.4	26.8
1940	52,526	23,084	29,442	24.0	20.9	27.3
1941	54,561	24,167	30,394	24.7	21.6	28.0
1942	56,521	25,306	31,215	25.4	22.4	28.6
1943	59,408	27,611	31,797	28.9	26.4	31.6
1944#	56,586	25,647	30,939	28.3	25.2	31.6
1945#	53,540	23,426	30,114	26.9	23.1	30.9
1946#	61,430	29,715	31,715	26.8	25.4	28.2
1947#	66,291	32,423	33,868	28.6	27.5	29.9
1948#	65,291	29,666	35,625	28.0	24.9	31.2
1949#	66,128	28,786	37,342	28.1	24.0	32.5

Table 11 - Page 2
Live Births - Mississippi - 1913-1969

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1950#	64,427	27,188	37,239	29.6	22.9	37.6
1951#	65,512	27,798	37,714	30.1	23.4	38.1
1952#	64,509	28,073	36,436	29.6	23.6	36.8
1953#	64,061	28,312	35,749	29.4	23.8	36.1
1954#	64,748	28,781	35,967	29.7	24.2	36.3
1955#	63,244	27,546	35,698	29.0	23.2	36.0
1956#	62,178	27,278	34,900	29.5	23.8	36.2
1957#	60,712	26,804	33,908	28.8	23.4	35.2
1958#	59,189	26,812	32,377	27.1	20.9	35.9
1959#	60,562	27,979	32,583	27.2	20.9	36.6
1960#	59,210	27,198	32,012	27.2	21.6	34.8
1961#	59,523	27,435	32,088	27.3	21.8	34.8
1962#	58,921	27,512	31,409	26.1	20.8	33.5
1963#	57,316	26,823	30,493	25.0	19.9	32.4
1964#	56,650	25,942	30,708	24.5	18.9	32.5
1965#	51,171	23,369	27,802	22.0	17.4	28.2
1966#	48,230	22,736	25,494	20.4	16.5	25.9
1967#	46,594	22,675	23,919	19.7	15.9	25.6
1968#	45,719	22,807	22,912	19.4	15.7	25.4
1969#	47,120	24,025	23,095	21.2	17.2	28.0

* Births per 1,000 population.

1913-1920, population estimates based on U.S. Censuses of 1910 and 1920.

1921-1930, estimates based on U.S. Censuses of 1920 and 1930.

1931-1940, estimates based on U.S. Censuses of 1930 and 1940.

1941 and 1942, projected estimates based on U.S. Censuses of 1930 and 1940.

1943 and 1944, estimates of civilian population obtained from Ration Book Registration.

1945, estimate of civilian population, July 1, 1945, obtained from Bureau of Census Bulletin, Series P-46, No. 3, dated February 12, 1946.

1946-1949, projected estimates as of July 1, each year based on U.S. Censuses of 1930 and 1940.

1950-1955, U.S. Census population of April 1, 1950.

1956 and 1957, estimate as of July 1, 1956, prepared by Dr. Harald A. Pedersen, Sociology Department, Mississippi State University, with adjustments by Statistical Service, Mississippi State Board of Health.

1958, estimate as of January 1, prepared by Dr. George L. Wilber and Mr. Carl T. Robbins, Sociology Department, Mississippi State University.

1959, estimate as of January 1, prepared by Dr. George L. Wilber and Ellen Bryant, Sociology Department, Mississippi State University.

1960 and 1961, U.S. Census population of April 1, 1960.

1962-1968, estimates as of July 1, prepared by the Division of Sociology and Anthropology, Mississippi State University.

1969, U.S. Census population of April 1, 1970.

On basis of place of residence.

TABLE 12
DEATHS
MISSISSIPPI
1913-1969

By Year, by Race, with Rates, Occurrence Data 1913-1943,
Residence Data 1944-1969

Year	Number			Rate*		
	Total	White	Non- white	Total	White	Non- white
1913	21,305	7,643	13,662	11.4	9.3	13.0
1914	21,485	7,442	14,043	11.3	8.9	13.1
1915	24,001	8,088	15,913	12.5	9.6	14.7
1916	21,794	7,588	14,206	11.2	8.9	13.0
1917	23,579	8,048	15,531	11.9	9.3	14.0
1918	30,437	10,177	20,260	15.2	11.6	18.0
1919	22,667	8,142	14,525	12.7	9.5	15.5
1920	22,072	7,916	14,156	12.3	9.3	15.1
1921	19,991	7,308	12,683	11.0	8.4	13.4
1922	19,278	7,312	11,966	10.5	8.3	12.6
1923	20,363	7,688	12,675	11.0	8.6	13.2
1924	21,155	7,809	13,346	11.3	8.6	13.8
1925	22,229	8,143	14,086	11.7	8.8	14.4
1926	23,929	9,133	14,796	12.5	9.7	15.1
1927	23,257	8,765	14,492	12.0	9.2	14.6
1928	25,965	10,111	15,854	13.2	10.4	15.9
1929	25,868	10,048	15,820	13.0	10.2	15.7
1930	24,098	9,072	15,026	11.9	9.0	14.8
1931	22,007	8,641	13,366	10.8	8.5	13.0
1932	20,322	8,295	12,027	9.9	8.1	11.6
1933	21,617	8,898	12,719	10.4	8.6	12.2
1934	21,786	9,350	12,436	10.5	9.0	12.0
1935	21,285	9,186	12,099	10.1	8.7	11.6
1936	24,127	10,367	13,760	11.4	9.7	13.1
1937	24,005	10,055	13,950	11.2	9.3	13.2
1938	22,814	9,471	13,343	10.6	8.7	12.5
1939	22,701	9,464	13,237	10.5	8.6	12.3
1940	23,185	9,565	13,620	10.6	8.6	12.6
1941	22,573	9,442	13,131	10.2	8.4	12.1
1942	20,693	8,750	11,943	9.3	7.7	10.9
1943	20,850	9,139	11,711	10.2	8.7	11.6
1944#	20,199	8,830	11,369	10.1	8.7	11.6
1945#	19,024	8,452	10,572	9.6	8.3	10.8
1946#	19,321	8,995	10,326	8.4	7.7	9.2
1947#	20,007	9,238	10,769	8.6	7.8	9.5
1948#	20,183	9,216	10,967	8.6	7.7	9.6
1949#	20,489	9,233	11,256	8.7	7.7	9.8

Table 12 - Page 2
Deaths - Mississippi - 1913-1969

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1950#	20,567	9,466	11,101	9.4	8.0	11.2
1951#	21,007	9,593	11,414	9.6	8.1	11.5
1952#	20,657	9,387	11,270	9.5	7.9	11.4
1953#	20,492	9,495	10,997	9.4	8.0	11.1
1954#	19,353	9,453	9,900	8.9	8.0	10.0
1955#	19,403	9,501	9,902	8.9	8.0	10.0
1956#	19,780	9,969	9,811	9.4	8.7	10.2
1957#	20,476	10,339	10,137	9.7	9.0	10.5
1958#	21,541	10,811	10,730	9.9	8.4	11.9
1959#	20,890	10,742	10,148	9.4	8.0	11.4
1960#	21,713	11,204	10,509	10.0	8.9	11.4
1961#	21,232	11,063	10,169	9.7	8.8	11.0
1962#	22,713	11,868	10,845	10.0	9.0	11.6
1963#	23,339	12,118	11,221	10.2	9.0	11.9
1964#	22,559	12,039	10,520	9.7	8.8	11.1
1965#	22,652	12,376	10,276	9.7	9.2	10.4
1966#	23,350	12,976	10,374	9.9	9.4	10.5
1967#	22,609	12,816	9,793	9.6	9.0	10.5
1968#	23,986	13,468	10,518	10.2	9.3	11.7
1969#	23,865	13,873	9,992	10.8	10.0	12.1

* Deaths per 1,000 population. See footnote on Table 11 for source of population estimates.

On basis of place of residence.

TABLE 13
MATERNAL DEATHS
MISSISSIPPI
1913-1969

By Year, by Race, with Rates, Occurrence Data 1913-1943,
Residence Data 1944-1969

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1913	447	162	285	108.8	74.1	148.2
1914	413	124	289	101.7	58.0	150.0
1915	289	88	201	73.2	41.9	108.7
1916	326	102	224	83.5	50.7	118.5
1917	297	105	192	80.0	55.1	106.2
1918	302	82	220	76.7	41.0	113.5
1919	407	173	234	95.8	78.5	114.4
1920	393	127	266	89.8	56.6	124.8
1921	391	145	246	84.1	61.2	108.0
1922	313	110	203	71.6	51.7	90.5
1923	321	108	213	74.2	51.1	96.3
1924	413	137	276	93.9	61.8	126.7
1925	435	143	292	95.5	64.2	125.3
1926	388	153	235	76.9	62.2	90.9
1927	409	134	275	83.1	55.4	109.8
1928	421	161	260	87.5	68.1	106.2
1929	410	152	258	90.0	68.8	110.0
1930	489	171	318	101.9	73.7	128.3
1931	352	131	221	78.0	60.5	94.1
1932	279	103	176	60.0	47.1	71.6
1933	315	116	199	71.3	56.8	83.7
1934	306	116	190	63.9	51.9	74.3
1935	334	137	197	69.2	62.4	74.8
1936	358	134	224	72.7	60.4	82.8
1937	370	150	220	70.9	64.4	76.2
1938	333	116	217	62.0	48.2	73.1
1939	311	97	214	59.5	41.3	74.3
1940	327	106	221	62.2	45.9	75.1
1941	293	93	200	53.7	38.5	65.8
1942	248	77	171	43.9	30.4	54.8
1943	233	66	167	39.2	23.9	52.5
1944#	225	72	153	39.8	28.1	49.4
1945#	216	64	152	40.3	27.3	50.5
1946#	199	67	132	32.4	22.5	41.6
1947#	175	50	125	26.4	15.4	36.9
1948#	177	38	139	27.1	12.8	39.0
1949#	147	32	115	22.2	11.1	30.8

Table 13 - Page 2
 Maternal Deaths - Mississippi - 1913-1969

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1950#	165	30	135	25.6	11.0	36.2
1951#	143	28	115	21.8	10.1	30.5
1952#	119	15	104	18.4	5.3	28.5
1953#	119	24	95	18.6	8.5	26.6
1954#	104	11	93	16.1	3.8	25.8
1955#	76	12	64	12.0	4.4	17.9
1956#	74	12	62	11.9	4.4	17.8
1957#	75	14	61	12.4	5.2	18.0
1958#	53	10	43	9.0	3.7	13.3
1959#	55	9	46	9.1	3.2	14.1
1960#	59	8	51	10.0	2.9	15.9
1961#	49	9	40	8.2	3.3	12.5
1962#	55	7	48	9.3	2.5	15.3
1963#	44	9	35	7.7	3.4	11.5
1964#	57	9	48	10.1	3.5	15.6
1965#	40	3	37	7.8	1.3	13.3
1966#	27	6	21	5.6	2.6	8.2
1967#	36	10	26	7.7	4.4	10.9
1968#	21	3	18	4.6	1.3	7.8
1969#	23	3	20	4.9	1.2	8.6

* Deaths per 10,000 live births.

On basis of place of residence.

TABLE 14
DEATHS UNDER 1 YEAR
MISSISSIPPI
1917-1969

By Year, by Race, with Rates, Occurrence Data 1917-1943,
Residence Data 1944-1969

Year	Number			Rate*		
	Total	White	Non- white	Total	White	Non- white
1917	3,689	1,296	2,393	99.3	68.0	132.3
1918	**	**	**	**	**	**
1919	3,554	**	**	83.6	**	**
1920	3,499	1,344	2,155	80.0	59.9	101.1
1921	3,157	1,243	1,914	67.9	52.4	84.0
1922	2,826	1,067	1,759	64.6	50.1	78.4
1923	2,908	1,104	1,804	67.2	52.2	81.5
1924	3,137	1,188	1,949	71.4	53.6	89.5
1925	3,176	1,179	1,997	69.7	53.0	85.7
1926	3,578	1,447	2,131	70.9	58.8	82.4
1927	3,291	1,303	1,988	66.9	54.0	79.4
1928	3,567	1,416	2,151	74.1	59.9	87.9
1929	3,275	1,268	2,007	71.9	57.4	85.5
1930	3,276	1,182	2,094	68.3	51.0	84.5
1931	2,547	961	1,586	56.4	44.4	67.5
1932	2,482	965	1,517	53.4	44.1	61.7
1933	2,800	1,035	1,765	63.4	50.7	74.2
1934	3,070	1,199	1,871	64.1	53.7	73.2
1935	2,587	1,044	1,543	53.6	47.5	58.6
1936	2,883	1,105	1,778	58.5	49.8	65.7
1937	3,098	1,174	1,924	59.4	50.4	66.6
1938	3,051	1,205	1,846	56.8	50.1	62.2
1939	2,908	1,135	1,773	55.6	48.3	61.6
1940	2,864	1,070	1,794	54.5	46.4	60.9
1941	2,969	1,113	1,856	54.4	46.1	61.1
1942	2,654	957	1,697	47.0	37.8	54.4
1943	2,774	1,137	1,637	46.7	41.2	51.5
1944#	2,489	974	1,515	44.0	38.0	49.0
1945#	2,192	846	1,346	40.9	36.1	44.7
1946#	2,298	1,005	1,293	37.4	33.8	40.8
1947#	2,428	1,009	1,419	36.6	31.1	41.9
1948#	2,470	988	1,482	37.8	33.3	41.6
1949#	2,612	900	1,712	39.5	31.3	45.8

Table 14 - Page 2
Deaths Under 1 Year - Mississippi - 1917-1969

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1950#	2,348	763	1,585	36.4	28.1	42.6
1951#	2,589	814	1,775	39.5	29.3	47.1
1952#	2,592	760	1,832	40.2	27.1	50.3
1953#	2,534	807	1,727	39.6	28.5	48.3
1954#	2,360	744	1,616	36.4	25.8	44.9
1955#	2,338	680	1,658	37.0	24.7	46.4
1956#	2,353	734	1,619	37.8	26.9	46.4
1957#	2,257	694	1,563	37.2	25.9	46.1
1958#	2,424	706	1,718	41.0	26.3	53.1
1959#	2,355	677	1,678	38.9	24.2	51.5
1960#	2,457	714	1,743	41.5	26.2	54.4
1961#	2,250	645	1,605	37.8	23.5	50.0
1962#	2,353	705	1,648	39.9	25.6	52.5
1963#	2,372	610	1,762	41.4	22.7	57.8
1964#	2,230	599	1,631	39.4	23.1	53.1
1965#	2,126	593	1,533	41.5	25.4	55.1
1966#	1,856	520	1,336	38.5	22.9	52.4
1967#	1,645	512	1,133	35.3	22.6	47.4
1968#	1,613	517	1,096	35.3	22.7	47.8
1969#	1,513	530	983	32.1	22.1	42.6

* Deaths per 1,000 live births.

** Not available.

On basis of place of residence.

TABLE 15
DEATHS UNDER 1 MONTH*
MISSISSIPPI
1924-1969

By Year, by Race, with Rates, Occurrence Data 1924-1944,
Residence Data 1945-1969

Year	Number			Rate**		
	Total	White	Non- white	Total	White	Non- white
1924	1,598	730	868	36.4	32.9	39.8
1925	1,639	701	938	36.0	31.5	40.3
1926	1,731	823	908	34.3	33.4	35.1
1927	1,644	786	858	33.4	32.5	34.3
1928	1,788	804	984	37.2	34.0	40.2
1929	1,662	773	889	36.5	35.0	37.9
1930	1,692	715	977	35.3	30.8	39.4
1931	1,339	590	749	29.7	27.3	31.9
1932	1,361	572	789	29.3	26.1	32.1
1933	1,540	636	904	34.8	31.2	38.0
1934	1,505	644	861	31.4	28.8	33.7
1935	1,446	642	804	29.9	29.2	30.5
1936	1,494	676	818	30.3	30.4	30.2
1937	1,678	755	923	32.2	32.4	32.0
1938	1,688	747	941	31.4	31.1	31.7
1939	1,637	735	902	31.3	31.3	31.3
1940	1,613	680	933	30.7	29.4	31.7
1941	1,663	734	929	30.5	30.4	30.6
1942	1,544	638	906	27.3	25.2	29.0
1943	1,652	791	861	27.8	28.6	27.1
1944	1,436	661	775	25.2	25.4	25.1
1945#	1,351	624	727	25.2	26.6	24.1
1946#	1,561	794	767	25.4	26.7	24.2
1947#	1,595	750	845	24.1	23.1	24.9
1948#	1,674	773	901	25.6	26.0	25.3
1949#	1,594	661	933	24.1	23.0	25.0
1950#	1,446	560	886	22.4	20.6	23.8
1951#	1,679	626	1,053	25.6	22.5	27.9
1952#	1,658	581	1,077	25.7	20.7	29.6
1953#	1,630	635	995	25.4	22.4	27.8
1954#	1,521	601	920	23.5	20.9	25.6
1955#	1,440	527	913	22.8	19.1	25.6
1956#	1,451	581	870	23.3	21.3	24.9
1957#	1,425	540	885	23.5	20.1	26.1
1958#	1,475	540	935	24.9	20.1	28.9
1959#	1,450	515	935	23.9	18.4	28.7

Table 15 - Page 2
 Deaths Under 1 Month - Mississippi - 1924-1969

Year	Number			Rate**		
	Total	White	Non-white	Total	White	Non-white
1960#	1,497	550	947	25.3	20.2	29.6
1961#	1,401	509	892	23.5	18.6	27.8
1962#	1,456	550	906	24.7	20.0	28.8
1963#	1,449	470	979	25.3	17.5	32.1
1964#	1,372	460	912	24.2	17.7	29.7
1965#	1,283	470	813	25.1	20.1	29.2
1966#	1,123	386	737	23.3	17.0	28.9
1967#	1,053	388	665	22.6	17.1	27.8
1968#	1,116	424	692	24.4	18.6	30.2
1969#	1,066	441	625	22.6	18.4	27.1

* Figures for 1949-1969 include only the deaths under 28 days.

** Deaths per 1,000 live births.

On basis of place of residence.

TABLE 16
DEATHS FROM HEART DISEASES, MALIGNANT NEOPLASMS,
CEREBROVASCULAR DISEASE, ACCIDENTS, AND INFLUENZA AND PNEUMONIA
(WITH INTERNATIONAL CLASSIFICATION NUMBERS, EIGHTH REVISION)
MISSISSIPPI
1960-1969

By Year, by Race, with Rates, Residence Data

Year	Heart Diseases (390-398, 402, 404-429)			Malignant Neoplasms (140-209)			Cerebrovascular Disease (430-438)			Accidents (800-949)			Influenza & Pneumonia (470-486)		
	Total	White	Non- white	Total	White	Non- white	Total	White	Non- white	Total	White	Non- white	Total	White	Non- white
1960	6,321	3,882	2,439	2,635	1,644	991	2,960	1,513	1,447	1,504	827	677	958	389	569
Rate*	290.2	308.7	264.9	121.0	130.7	107.6	135.9	120.3	157.2	69.0	65.8	73.5	44.0	30.9	61.8
1961	6,561	4,022	2,539	2,748	1,732	1,016	2,891	1,464	1,427	1,427	760	667	721	307	414
Rate*	301.2	319.8	275.8	126.2	137.7	110.4	132.7	116.4	155.0	65.5	60.4	72.4	33.1	24.4	45.0
1962	7,077	4,206	2,871	2,827	1,793	1,034	3,112	1,552	1,560	1,557	846	711	879	362	517
Rate*	313.4	318.4	306.4	125.2	135.7	110.4	137.8	117.5	166.5	69.0	64.0	75.9	38.9	27.4	55.2
1963	7,300	4,396	2,904	2,945	1,859	1,086	3,154	1,650	1,504	1,571	851	720	1,113	420	693
Rate*	318.8	325.9	308.5	128.6	137.8	115.4	137.7	122.3	159.8	68.6	63.1	76.5	48.6	31.1	73.6
1964	6,981	4,313	2,668	2,968	1,861	1,107	3,077	1,588	1,489	1,612	898	714	791	308	483
Rate*	301.4	314.7	282.2	128.2	135.8	117.1	132.9	115.9	157.5	69.6	65.5	75.5	34.2	22.5	51.1
1965	7,038	4,429	2,609	2,931	1,867	1,064	3,064	1,639	1,425	1,610	881	729	764	350	414
Rate*	302.1	329.4	264.8	125.8	138.9	108.0	131.5	121.9	144.6	69.1	65.5	74.0	32.8	26.0	42.0
1966	7,468	4,729	2,739	3,098	1,990	1,108	3,132	1,681	1,451	1,872	1,072	800	773	341	432
Rate*	316.5	343.8	278.2	131.3	144.7	112.6	132.7	122.2	147.4	79.3	77.9	81.3	32.8	24.8	43.9
1967	7,300	4,640	2,660	3,136	2,061	1,075	2,981	1,632	1,349	1,890	1,166	724	657	310	347
Rate*	309.2	325.2	284.7	132.8	144.4	115.1	126.2	114.4	144.4	80.0	81.7	77.5	27.8	21.7	37.1
1968	7,969	5,024	2,945	3,267	2,101	1,166	3,115	1,631	1,484	1,723	999	724	825	414	411
Rate*	339.0	346.4	327.1	139.0	144.9	129.5	132.5	112.4	164.8	73.3	68.9	80.4	35.1	28.5	45.6
1969	7,886	5,142	2,744	3,295	2,146	1,149	3,058	1,681	1,377	2,007	1,179	828	726	385	341
Rate*	355.7	369.0	333.2	148.6	154.0	139.5	137.9	120.6	167.2	90.5	84.6	100.5	32.7	27.6	41.4

* Deaths per 100,000 population.

TABLE 17
MARRIAGES AND DIVORCES
MISSISSIPPI
1926-1969

By Year, by Race, Occurrence Data

Year	Marriages			Divorces		
	Total	White	Non-white	Total	White	Non-white
1926*	28,712			2,766		
1927*	29,048#	11,402	17,383	2,895		
1928	30,231#	11,662	18,257	3,009	1,435	1,574
1929	31,495#	12,864	18,388	3,172	1,598	1,574
1930	25,728#	10,815	14,751	2,743	1,487	1,256
1931	21,589#	9,952	11,629	2,015	1,308	707
1932	22,688#	10,147	12,453	1,918	1,308	610
1933	27,567#	12,201	15,300	1,955	1,396	559
1934	30,149#	13,497	16,541	2,446	1,704	742
1935	28,722#	12,955	15,635	2,425	1,681	744
1936	32,480#	14,120	18,196	2,520	1,690	830
1937	32,055#	14,802	17,246	2,977	1,984	993
1938	31,495	14,224	17,271	2,916	2,069	847
1939	31,956	15,370	16,586	3,034	2,166	868
1940	34,088	16,834	17,254	3,251	2,295	956
1941	42,142	21,611	20,531	3,831	2,444	1,387
1942	49,565	27,227	22,338	4,326	2,606	1,720
1943	46,030	23,445	22,585	5,851	3,062	2,789
1944	41,559	21,173	20,386	6,742	3,425	3,317
1945	44,871	24,738	20,133	6,727	3,919	2,808
1946	64,204	37,099	27,105	9,495	5,867	3,628
1947	53,584	30,514	23,070	7,004	4,323	2,681
1948	53,645	31,504	22,141	6,697	3,864	2,833
1949	52,765	32,191	20,574	6,285	3,621	2,664
1950	56,738	37,871	18,867	6,065	3,711	2,354
1951	57,321	37,857	19,464	5,464	3,601	1,863
1952	58,116	39,162	18,954	5,419	3,607	1,812
1953	61,751	42,961	18,790	5,204	3,515	1,689
1954	61,381	43,268	18,113	5,001	3,424	1,577
1955	66,423	47,618	18,805	4,845	3,372	1,473
1956	66,223	48,140	18,083	5,027	3,452	1,575
1957	62,495	46,868	15,627	4,974	3,350	1,624
1958	36,402	26,998	9,404	5,016	3,485	1,531
1959	20,447	13,547	6,900	5,108	3,557	1,551
1960	21,233	14,029	7,204	5,260	3,553	1,707
1961	21,610	14,438	7,172	5,213	3,545	1,668
1962	22,007	14,432	7,575	5,467	3,827	1,640
1963	22,140	14,508	7,632	5,452	3,930	1,522
1964	22,784	15,159	7,625	5,631	4,077	1,554
1965	23,343	15,613	7,730	5,730	4,220	1,510
1966	23,476	15,860	7,616	6,306	4,650	1,656
1967	23,535	15,752	7,783	6,455	4,704	1,751
1968	25,355	17,142	8,213	7,020	5,202	1,818
1969	25,956	17,582	8,374	7,506	5,616	1,890

* Total number of marriages for 1926 and total number of divorces for 1926 and 1927 furnished by U.S. Bureau of Census; distribution by race not available.

Includes some records on which the race of the couple was not stated.

BIRTHS, FETAL DEATHS, DEATHS
THE COUNTIES AND THE STATE OF MISSISSIPPI, 1969

Footnote for Tables on Pages 63-145

- * Total births and deaths per 1,000 population. Fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from certain causes of mortality in early infancy per 1,000 live births. Maternal deaths per 10,000 live births. Deaths from other specific causes per 100,000 population.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
ADAMS COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,565	753	812	890	393	497	23.9	20.3	27.7
Fetal deaths	66	29	37	38	15	23	42.7	38.2	46.3
Total deaths	464	204	260	383	164	219	10.3	8.5	12.2
Deaths under 1 year	41	16	25	29	13	16	32.6	33.1	32.2
Deaths under 28 days	35	14	21	25	12	13	28.1	30.5	26.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	1	0	1	1	0	1	2.7	-	5.6
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	1	0	1	1	0	1	2.7	-	5.6
Malignant neoplasms (140-209)	66	32	34	51	20	31	136.8	103.3	172.9
Benign and unspecified neoplasms (210-239)	2	1	1	1	0	1	2.7	-	5.6
Diabetes mellitus (250)	15	3	12	10	3	7	26.8	15.5	39.0
Avitaminoses and other nutritional deficiencies (260-269)	1	0	1	0	0	0	-	-	-
Anemias (280-285)	2	0	2	1	0	1	2.7	-	5.6
Diseases of heart (390-398, 402, 404-429)	167	81	86	144	65	79	386.1	335.6	440.7
Hypertension (400, 401, 403)	2	1	1	1	0	1	2.7	-	5.6
Cerebrovascular disease (430-438)	49	18	31	43	13	30	115.3	67.1	167.3
Arteriosclerosis (440)	12	3	9	8	2	6	21.4	10.3	33.5
Influenza and pneumonia (470-486)	13	7	6	10	6	4	26.8	31.0	22.3
Bronchitis, emphysema, asthma (490-493)	5	3	2	3	3	0	8.0	15.5	-
Peptic ulcer (531-533)	3	1	2	2	0	2	5.4	-	11.2
Hernia and intestinal obstruction (550-553, 560)	2	2	0	1	1	0	2.7	5.2	-
Cirrhosis of liver (571)	7	5	2	6	5	1	16.1	25.8	5.6
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	4	1	3	2	1	1	5.4	5.2	5.6
Infections of kidney (590)	2	1	1	1	0	1	2.7	-	5.6
Hyperplasia of prostate (600)	2	0	2	1	0	1	2.7	-	5.6
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	4	2	2	3	3	0	8.0	15.5	-
Certain causes of mortality in early infancy (760-778)	25	8	17	17	6	11	19.1	15.3	22.1
Accidents (800-949)	31	16	15	30	18	12	80.4	92.9	66.9
Motor vehicle accidents (810-823)	13	9	4	12	9	3	32.2	46.5	16.7
Suicide (950-959)	0	0	0	1	0	1	2.7	-	5.6
Homicide (960-978)	9	1	8	10	2	8	26.8	10.3	44.6
Ill-defined and unknown (780-796)	8	2	6	9	3	6	24.1	15.5	33.5

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
ALCORN COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	329	265	64	431	365	66	15.8	15.2	20.4
Fetal deaths	7	2	5	16	11	5	37.1	30.1	75.8
Total deaths	326	278	48	359	308	51	13.2	12.8	15.8
Deaths under 1 year	14	8	6	22	16	6	51.0	43.8	90.9
Deaths under 28 days	12	7	5	20	15	5	46.4	41.1	75.8
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	1	0	1	1	0	3.7	4.2	-
Tuberculosis (010-019)	1	1	0	3	2	1	11.0	8.4	31.0
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	43	37	6	47	41	6	172.9	171.2	185.9
Benign and unspecified neoplasms (210-239)	1	1	0	1	1	0	3.7	4.2	-
Diabetes mellitus (250)	2	1	1	3	2	1	11.0	8.4	31.0
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	154	134	20	149	129	20	548.2	538.6	619.6
Hypertension (400, 401, 403)	4	2	2	5	3	2	18.4	12.5	62.0
Cerebrovascular disease (430-438)	42	37	5	44	37	7	161.9	154.5	216.8
Arteriosclerosis (440)	4	4	0	5	5	0	18.4	20.9	-
Influenza and pneumonia (470-486)	8	6	2	10	8	2	36.8	33.4	62.0
Bronchitis, emphysema, asthma (490-493)	3	2	1	4	3	1	14.7	12.5	31.0
Peptic ulcer (531-533)	1	1	0	1	1	0	3.7	4.2	-
Hernia and intestinal obstruction (550-553, 560)	1	0	1	1	0	1	3.7	-	31.0
Cirrhosis of liver (571)	2	2	0	4	4	0	14.7	16.7	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	1	0	2	2	0	7.4	8.4	-
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	1	1	0	1	1	0	3.7	4.2	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	2	2	0	5	5	0	18.4	20.9	-
Certain causes of mortality in early infancy (760-778)	11	6	5	17	12	5	39.4	32.9	75.8
Accidents (800-949)	15	14	1	15	15	0	55.2	62.6	-
Motor vehicle accidents (810-823)	7	7	0	7	7	0	25.8	29.2	-
Suicide (950-959)	2	2	0	4	4	0	14.7	16.7	-
Homicide (960-978)	6	4	2	5	3	2	18.4	12.5	62.0
Ill-defined and unknown (780-796)	11	10	1	17	16	1	62.5	66.8	31.0

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
AMITE COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	80	0	80	275	87	188	20.0	12.8	27.0
Fetal deaths	1	0	1	9	1	8	32.7	11.5	42.6
Total deaths	67	27	40	170	100	70	12.4	14.7	10.1
Deaths under 1 year	1	0	1	5	2	3	18.2	23.0	16.0
Deaths under 28 days	0	0	0	3	1	2	10.9	11.5	10.6
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	7	2	5	24	18	6	174.4	264.2	86.3
Benign and unspecified neoplasms (210-239)	0	0	0	1	1	0	7.3	14.7	-
Diabetes mellitus (250)	0	0	0	4	2	2	29.1	29.4	28.8
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	2	1	1	14.5	14.7	14.4
Diseases of heart (390-398, 402, 404-429)	22	14	8	53	32	21	385.1	469.6	302.2
Hypertension (400, 401, 403)	0	0	0	2	0	2	14.5	-	28.8
Cerebrovascular disease (430-438)	5	1	4	19	11	8	138.0	161.4	115.1
Arteriosclerosis (440)	1	0	1	4	3	1	29.1	44.0	14.4
Influenza and pneumonia (470-486)	0	0	0	4	4	0	29.1	58.7	-
Bronchitis, emphysema, asthma (490-493)	2	1	1	6	5	1	43.6	73.4	14.4
Peptic ulcer (531-533)	0	0	0	1	1	0	7.3	14.7	-
Hernia and intestinal obastruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	1	1	0	7.3	14.7	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	1	0	1	7.3	-	14.4
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	0	0	0	3	1	2	10.9	11.5	10.6
Accidents (800-949)	4	1	3	6	4	2	43.6	58.7	28.8
Motor vehicle accidents (810-823)	1	1	0	1	1	0	7.3	14.7	-
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	0	0	0	1	0	1	7.3	-	14.4
Ill-defined and unknown (780-796)	23	6	17	24	7	17	174.4	102.7	244.6

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
ATTALA COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	366	183	183	394	178	216	20.1	15.3	27.2
Fetal deaths	16	8	8	18	8	10	45.7	44.9	46.3
Total deaths	249	152	97	275	167	108	14.0	14.4	13.6
Deaths under 1 year	11	4	7	15	5	10	38.1	28.1	46.3
Deaths under 28 days	8	3	5	10	3	7	25.4	16.8	32.4
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	2	0	2	1	0	1	5.1	-	12.6
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	31	20	11	31	19	12	158.4	163.3	151.2
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	8	4	4	8	4	4	40.9	34.4	50.4
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	110	76	34	122	82	40	623.4	705.0	503.9
Hypertension (400, 401, 403)	1	1	0	1	1	0	5.1	8.6	-
Cerebrovascular disease (430-438)	32	19	13	39	27	12	199.3	232.1	151.2
Arteriosclerosis (440)	4	2	2	4	2	2	20.4	17.2	25.2
Influenza and pneumonia (470-486)	6	4	2	7	5	2	35.8	43.0	25.2
Bronchitis, emphysema, asthma (490-493)	2	2	0	1	1	0	5.1	8.6	-
Peptic ulcer (531-533)	2	1	1	3	2	1	15.3	17.2	12.6
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	2	2	0	10.2	17.2	-
Nephritis and nephrosis (580-584)	0	0	0	3	1	2	15.3	8.6	25.2
Infections of kidney (590)	1	0	1	2	1	1	10.2	8.6	12.6
Hyperplasia of prostate (600)	1	0	1	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	2	1	1	4	2	2	20.4	17.2	25.2
Certain causes of mortality in early infancy (760-778)	7	3	4	7	2	5	17.8	11.2	23.1
Accidents (800-949)	15	9	6	15	6	9	76.6	51.6	113.4
Motor vehicle accidents (810-823)	9	5	4	7	3	4	35.8	25.8	50.4
Suicide (950-959)	3	2	1	2	1	1	10.2	8.6	12.6
Homicide (960-978)	1	0	1	1	0	1	5.1	-	12.6
Ill-defined and unknown (780-796)	11	4	7	9	4	5	46.0	34.4	63.0

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
BENTON COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	42	1	41	132	64	68	17.6	14.7	21.6
Fetal deaths	1	0	1	7	3	4	53.0	46.9	58.8
Total deaths	22	12	10	67	44	23	8.9	10.1	7.3
Deaths under 1 year	1	0	1	4	2	2	30.3	31.2	29.4
Deaths under 28 days	0	0	0	3	2	1	22.7	31.2	14.7
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	1	0	1	13.3	-	31.7
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	0	0	0	6	5	1	79.9	114.9	31.7
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	0	0	0	1	1	0	13.3	23.0	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	11	6	5	22	14	8	293.1	321.8	253.6
Hypertension (400, 401, 403)	0	0	0	1	0	1	13.3	-	31.7
Cerebrovascular disease (430-438)	4	3	1	12	8	4	159.9	183.9	126.8
Arteriosclerosis (440)	0	0	0	0	0	0	-	-	-
Influenza and pneumonia (470-486)	0	0	0	2	1	1	26.6	23.0	31.7
Bronchitis, emphysema, asthma (490-493)	0	0	0	1	1	0	13.3	23.0	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	1	1	0	13.3	23.0	-
Infections of kidney (590)	0	0	0	1	1	0	13.3	23.0	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	1	1	0	13.3	23.0	-
Certain causes of mortality in early infancy (760-778)	0	0	0	1	1	0	7.6	15.6	-
Accidents (800-949)	0	0	0	3	2	1	40.0	46.0	31.7
Motor vehicle accidents (810-823)	0	0	0	2	1	1	26.6	23.0	31.7
Suicide (950-959)	1	0	1	2	1	1	26.6	23.0	31.7
Homicide (960-978)	0	0	0	1	0	1	13.3	-	31.7
Ill-defined and unknown (780-796)	5	3	2	5	3	2	66.6	69.0	63.4

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
BOLIVAR COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,741	368	1,373	1,370	339	1,031	27.7	18.1	33.6
Fetal deaths	64	3	61	54	9	45	39.4	26.5	43.6
Total deaths	570	126	444	551	152	399	11.2	8.1	13.0
Deaths under 1 year	52	5	47	49	7	42	35.8	20.6	40.7
Deaths under 28 days	34	4	30	34	6	28	24.8	17.7	27.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	7	0	7	8	0	8	16.2	-	26.1
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	60	9	51	66	20	46	133.6	106.7	150.0
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	6	2	4	4	2	2	8.1	10.7	6.5
Avitaminoses and other nutritional deficiencies (260-269)	1	0	1	1	0	1	2.0	-	3.3
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	176	54	122	180	62	118	364.3	330.7	384.9
Hypertension (400, 401, 403)	4	2	2	5	2	3	10.1	10.7	9.8
Cerebrovascular disease (430-438)	111	17	94	91	19	72	184.2	101.3	234.8
Arteriosclerosis (440)	15	6	9	12	5	7	24.3	26.7	22.8
Influenza and pneumonia (470-486)	19	3	16	18	3	15	36.4	16.0	48.9
Bronchitis, emphysema, asthma (490-493)	3	2	1	3	2	1	6.1	10.7	3.3
Peptic ulcer (531-533)	1	1	0	2	2	0	4.0	10.7	-
Hernia and intestinal obstruction (550-553, 560)	2	0	2	0	0	0	-	-	-
Cirrhosis of liver (571)	5	1	4	5	1	4	10.1	5.3	13.0
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	1	1	0	2.0	5.3	-
Nephritis and nephrosis (580-584)	9	0	9	6	1	5	12.1	5.3	16.3
Infections of kidney (590)	1	0	1	1	0	1	2.0	-	3.3
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	4	1	3	4	1	3	8.1	5.3	9.8
Certain causes of mortality in early infancy (760-778)	21	2	19	20	4	16	14.6	11.8	15.5
Accidents (800-949)	47	13	34	49	15	34	99.2	80.0	110.9
Motor vehicle accidents (810-823)	15	4	11	17	6	11	34.4	32.0	35.9
Suicide (950-959)	3	3	0	2	2	0	4.0	10.7	-
Homicide (960-978)	5	1	4	6	1	5	12.1	5.3	16.3
Ill-defined and unknown (780-796)	41	4	37	40	4	36	81.0	21.3	117.4

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
CALHOUN COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	179	69	110	276	170	106	18.9	15.7	27.8
Fetal deaths	8	1	7	12	5	7	43.5	29.4	66.0
Total deaths	132	103	29	160	133	27	10.9	12.3	7.1
Deaths under 1 year	8	2	6	8	2	6	29.0	11.8	56.6
Deaths under 28 days	4	1	3	5	1	4	18.1	5.9	37.7
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	2	0	2	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	20	17	3	27	22	5	184.6	203.6	131.0
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	4	2	2	5	3	2	34.2	27.8	52.4
Avitaminoses and other nutritional deficiencies (260-269)	1	1	0	1	1	0	6.8	9.2	-
Anemias (280-285)	0	0	0	1	1	0	6.8	9.2	-
Diseases of heart (390-398, 402, 404-429)	59	51	8	72	65	7	492.4	601.5	183.4
Hypertension (400, 401, 403)	1	1	0	2	2	0	13.7	18.5	-
Cerebrovascular disease (430-438)	7	6	1	11	10	1	75.2	92.5	26.2
Arteriosclerosis (440)	0	0	0	0	0	0	-	-	-
Influenza and pneumonia (470-486)	3	2	1	4	2	2	27.4	18.5	52.4
Bronchitis, emphysema, asthma (490-493)	3	3	0	4	4	0	27.4	37.0	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	1	1	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	1	1	0	6.8	9.2	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	1	1	0	6.8	9.2	-
Hyperplasia of prostate (600)	1	1	0	1	1	0	6.8	9.2	-
Maternal causes (630-678)	3	1	2	0	0	0	-	-	-
Congenital anomalies (740-759)	1	0	1	1	0	1	6.8	-	26.2
Certain causes of mortality in early infancy (760-778)	3	1	2	3	1	2	10.9	5.9	18.9
Accidents (800-949)	11	8	3	13	11	2	88.9	101.8	52.4
Motor vehicle accidents (810-823)	7	6	1	7	7	0	47.9	64.8	-
Suicide (950-959)	3	3	0	2	2	0	13.7	18.5	-
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	3	2	1	3	2	1	20.5	18.5	26.2

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
CARROLL COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non- white	Total	White	Non- white	Total	White	Non- white
Live births	42	2	40	154	61	93	16.4	13.2	19.4
Fetal deaths	1	0	1	5	1	4	32.5	16.4	43.0
Total deaths	48	16	32	107	60	47	11.4	13.0	9.8
Deaths under 1 year	1	1	0	5	2	3	32.5	32.8	32.2
Deaths under 28 days	1	1	0	2	2	0	13.0	32.8	-
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	1	0	1	10.6	-	20.9
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	7	2	5	18	9	9	191.6	195.0	188.2
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	0	0	0	3	2	1	31.9	43.3	20.9
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	15	7	8	33	21	12	351.2	455.0	250.9
Hypertension (400, 401, 403)	4	0	4	5	1	4	53.2	21.7	83.6
Cerebrovascular disease (430-438)	2	1	1	10	6	4	106.4	130.0	83.6
Arteriosclerosis (440)	1	0	1	2	1	1	21.3	21.7	20.9
Influenza and pneumonia (470-486)	0	0	0	2	1	1	21.3	21.7	20.9
Bronchitis, emphysema, asthma (490-493)	0	0	0	1	1	0	10.6	21.7	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	2	2	0	21.3	43.3	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	1	1	0	10.6	21.7	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	1	1	0	10.6	21.7	-
Certain causes of mortality in early infancy (760-778)	0	0	0	1	1	0	6.5	16.4	-
Accidents (800-949)	6	0	6	8	3	5	85.1	65.0	104.6
Motor vehicle accidents (810-823)	2	0	2	3	2	1	31.9	43.3	20.9
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	2	0	2	2	0	2	21.3	-	41.8
Ill-defined and unknown (780-796)	9	5	4	9	4	5	95.8	86.7	104.6

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
CHICKASAW COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	286	111	175	358	171	187	21.3	15.8	31.2
Fetal deaths	25	14	11	17	8	9	47.5	46.8	48.1
Total deaths	214	125	89	224	134	90	13.3	12.4	15.0
Deaths under 1 year	11	3	8	12	3	9	33.5	17.5	48.1
Deaths under 28 days	4	1	3	4	1	3	11.2	5.8	16.0
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	3	0	3	17.8	-	50.1
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	27	18	9	30	23	7	178.5	212.6	116.9
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	3	1	2	3	1	2	17.8	9.2	33.4
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	86	58	28	79	56	23	470.1	517.6	384.2
Hypertension (400, 401, 403)	4	0	4	3	0	3	17.8	-	50.1
Cerebrovascular disease (430-438)	33	17	16	36	16	20	214.2	147.9	334.1
Arteriosclerosis (440)	0	0	0	1	1	0	6.0	9.2	-
Influenza and pneumonia (470-486)	9	4	5	7	4	3	41.6	37.0	50.1
Bronchitis, emphysema, asthma (490-493)	2	2	0	2	2	0	11.9	18.5	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	1	1	0	1	1	0	6.0	9.2	-
Cirrhosis of liver (571)	2	2	0	2	2	0	11.9	18.5	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	1	0	3	1	2	17.8	9.2	33.4
Infections of kidney (590)	2	2	0	1	1	0	6.0	9.2	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	3	1	2	83.8	58.5	107.0
Congenital anomalies (740-759)	3	1	2	3	1	2	17.8	9.2	33.4
Certain causes of mortality in early infancy (760-778)	0	0	0	1	0	1	2.8	-	5.3
Accidents (800-949)	16	5	11	18	7	11	107.1	64.7	183.8
Motor vehicle accidents (810-823)	7	0	7	7	2	5	41.6	18.5	83.5
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	3	1	2	4	1	3	23.8	9.2	50.1
Ill-defined and unknown (780-796)	12	6	6	11	6	5	65.4	55.4	83.5

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
CHOCTAW COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	172	60	112	157	75	82	18.6	12.4	34.6
Fetal deaths	2	1	1	3	2	1	19.1	26.7	12.2
Total deaths	91	56	35	113	84	29	13.4	13.8	12.2
Deaths under 1 year	4	2	2	2	1	1	12.7	13.3	12.2
Deaths under 28 days	1	1	0	0	0	0	-	-	-
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	1	0	1	1	0	11.8	16.5	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	8	5	3	10	8	2	118.5	131.8	84.4
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	1	0	1	1	0	1	11.8	-	42.2
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	37	25	12	39	31	8	462.1	510.8	337.4
Hypertension (400, 401, 403)	0	0	0	1	1	0	11.8	16.5	-
Cerebrovascular disease (430-438)	12	6	6	17	13	4	201.4	214.2	168.7
Arteriosclerosis (440)	1	1	0	1	1	0	11.8	16.5	-
Influenza and pneumonia (470-486)	3	3	0	4	4	0	47.4	65.9	-
Bronchitis, emphysema, asthma (490-493)	4	3	1	4	3	1	47.4	49.4	42.2
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	2	0	2	2	0	2	23.7	-	84.4
Cirrhosis of liver (571)	0	0	0	1	1	0	11.8	16.5	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	1	1	0	11.8	16.5	-
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	2	0	2	23.7	-	84.4
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	0	1	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	1	1	0	0	0	0	-	-	-
Accidents (800-949)	6	3	3	9	7	2	106.6	115.3	84.4
Motor vehicle accidents (810-823)	0	0	0	5	4	1	59.2	65.9	42.2
Suicide (950-959)	1	0	1	1	0	1	11.8	-	42.2
Homicide (960-978)	2	2	0	4	3	1	47.4	49.4	42.2
Ill-defined and unknown (780-796)	3	1	2	3	1	2	35.5	16.5	84.4

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
CLAIBORNE COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	83	13	70	202	37	165	20.0	14.6	21.8
Fetal deaths	4	1	3	16	5	11	79.2	135.1	66.7
Total deaths	95	25	70	128	41	87	12.7	16.2	11.5
Deaths under 1 year	8	1	7	11	2	9	54.4	54.0	54.5
Deaths under 28 days	5	1	4	8	2	6	39.6	54.0	36.4
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	11	2	9	22	8	14	218.1	315.4	185.4
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	1	0	1	1	0	1	9.9	-	13.2
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	52	14	38	60	19	41	594.9	749.2	543.0
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	12	1	11	14	2	12	138.8	78.9	158.9
Arteriosclerosis (440)	0	0	0	2	1	1	19.8	39.4	13.2
Influenza and pneumonia (470-486)	5	1	4	5	1	4	49.6	39.4	53.0
Bronchitis, emphysema, asthma (490-493)	1	1	0	1	1	0	9.9	39.4	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	2	2	0	2	2	0	19.8	78.9	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	0	1	9.9	-	13.2
Nephritis and nephrosis (580-584)	0	0	0	2	1	1	19.8	39.4	13.2
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	4	1	3	7	2	5	34.6	54.0	30.3
Accidents (800-949)	0	0	0	1	1	0	9.9	39.4	-
Motor vehicle accidents (810-823)	0	0	0	1	1	0	9.9	39.4	-
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	1	0	1	1	0	1	9.9	-	13.2
Ill-defined and unknown (780-796)	3	0	3	2	0	2	19.8	-	26.5

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
CLARKE COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non- white	Total	White	Non- white	Total	White	Non- white
Live births	38	18	20	284	137	147	18.9	14.2	27.1
Fetal deaths	1	0	1	11	4	7	38.7	29.2	47.6
Total deaths	116	74	42	184	118	66	12.2	12.2	12.2
Deaths under 1 year	1	0	1	5	1	4	17.6	7.3	27.2
Deaths under 28 days	0	0	0	3	1	2	10.6	7.3	13.6
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	1	0	2	1	1	13.3	10.4	18.4
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	11	6	5	23	13	10	152.8	135.0	184.6
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	2	1	1	7	3	4	46.5	31.1	73.8
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	46	34	12	64	50	14	425.3	519.2	258.4
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	24	14	10	32	18	14	212.6	186.9	258.4
Arteriosclerosis (440)	3	3	0	5	5	0	33.2	51.9	-
Influenza and pneumonia (470-486)	2	1	1	5	4	1	33.2	41.5	18.4
Bronchitis, emphysema, asthma (490-493)	0	0	0	1	1	0	6.6	10.4	-
Peptic ulcer (531-533)	1	0	1	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	1	1	0	1	1	0	6.6	10.4	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	1	0	1	1	0	6.6	10.4	-
Infections of kidney (590)	1	1	0	2	2	0	13.3	20.8	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	0	0	0	3	1	2	10.6	7.3	13.6
Accidents (800-949)	7	4	3	16	9	7	106.3	93.4	129.2
Motor vehicle accidents (810-823)	3	1	2	7	3	4	46.5	31.1	73.8
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	1	0	1	2	0	2	13.3	-	36.9
Ill-defined and unknown (780-796)	14	6	8	15	7	8	99.7	72.7	147.6

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
CLAY COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	237	89	148	316	130	186	16.8	13.6	20.0
Fetal deaths	9	3	6	11	4	7	34.8	30.8	37.6
Total deaths	174	82	92	209	107	102	11.1	11.2	10.9
Deaths under 1 year	11	2	9	13	3	10	41.1	23.1	53.8
Deaths under 28 days	5	1	4	7	2	5	22.2	15.4	26.9
Bacillary dysentery and amebiasis (004, 006)	1	0	1	1	0	1	5.3	-	10.7
Enteritis and other diarrheal diseases (008, 009)	1	0	1	0	0	0	-	-	-
Tuberculosis (010-019)	1	0	1	1	0	1	5.3	-	10.7
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	1	0	1	5.3	-	10.7
Malignant neoplasms (140-209)	26	17	9	33	23	10	175.2	241.7	107.3
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	2	2	0	5	5	0	26.5	52.5	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	68	29	39	73	33	40	387.5	346.7	429.0
Hypertension (400, 401, 403)	0	0	0	1	0	1	5.3	-	10.7
Cerebrovascular disease (430-438)	33	13	20	34	15	19	180.5	157.6	203.8
Arteriosclerosis (440)	1	1	0	1	1	0	5.3	10.5	-
Influenza and pneumonia (470-486)	1	0	1	3	2	1	15.9	21.0	10.7
Bronchitis, emphysema, asthma (490-493)	2	2	0	3	3	0	15.9	31.5	-
Peptic ulcer (531-533)	2	0	2	1	0	1	5.3	-	10.7
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	1	0	5.3	10.5	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	1	1	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	1	0	1	1	0	1	5.3	-	10.7
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	2	1	1	10.6	10.5	10.7
Certain causes of mortality in early infancy (760-778)	4	1	3	5	1	4	15.8	7.7	21.5
Accidents (800-949)	13	8	5	23	12	11	122.1	126.1	118.0
Motor vehicle accidents (810-823)	7	6	1	12	6	6	63.7	63.0	64.4
Suicide (950-959)	1	1	0	1	1	0	5.3	10.5	-
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	9	3	6	10	3	7	53.1	31.5	75.1

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
COAHOMA COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	855	338	517	943	243	700	23.3	17.1	26.7
Fetal deaths	64	25	39	55	16	39	58.3	65.8	55.7
Total deaths	514	181	333	510	149	361	12.6	10.5	13.8
Deaths under 1 year	45	9	36	46	7	39	48.8	28.8	55.7
Deaths under 28 days	30	8	22	29	6	23	30.8	24.7	32.8
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	5	0	5	4	0	4	9.9	-	15.2
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	62	33	29	61	26	35	150.8	182.7	133.5
Benign and unspecified neoplasms (210-239)	1	1	0	1	1	0	2.5	7.0	-
Diabetes mellitus (250)	14	7	7	12	4	8	29.7	28.1	30.5
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	167	62	105	165	51	114	407.9	358.3	434.9
Hypertension (400, 401, 403)	4	1	3	2	1	1	4.9	7.0	3.8
Cerebrovascular disease (430-438)	80	30	50	75	23	52	185.4	161.6	198.4
Arteriosclerosis (440)	3	1	2	3	1	2	7.4	7.0	7.6
Influenza and pneumonia (470-486)	14	3	11	16	2	14	39.6	14.0	53.4
Bronchitis, emphysema, asthma (490-493)	5	3	2	4	2	2	9.9	14.0	7.6
Peptic ulcer (531-533)	2	0	2	2	0	2	4.9	-	7.6
Hernia and intestinal obstruction (550-553, 560)	1	0	1	1	0	1	2.5	-	3.8
Cirrhosis of liver (571)	3	3	0	4	4	0	9.9	28.1	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	2	1	1	3	1	2	7.4	7.0	7.6
Nephritis and nephrosis (580-584)	3	0	3	3	0	3	7.4	-	11.4
Infections of kidney (590)	6	1	5	6	2	4	14.8	14.0	15.2
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	4	1	3	9	5	4	22.2	35.1	15.2
Certain causes of mortality in early infancy (760-778)	19	7	12	16	3	13	17.0	12.3	18.6
Accidents (800-949)	41	15	26	41	14	27	101.4	98.4	103.0
Motor vehicle accidents (810-823)	16	10	6	13	7	6	32.1	49.2	22.9
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	5	0	5	6	0	6	14.8	-	22.9
Ill-defined and unknown (780-796)	48	1	47	50	1	49	123.6	7.0	186.9

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
COPIAH COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	581	173	408	478	148	330	19.3	12.0	26.5
Fetal deaths	14	0	14	19	4	15	39.7	27.0	45.4
Total deaths	280	142	138	347	177	170	14.0	14.4	13.6
Deaths under 1 year	12	3	9	24	6	18	50.2	40.5	54.5
Deaths under 28 days	8	3	5	19	6	13	39.7	40.5	39.4
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	3	0	3	12.1	-	24.1
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	35	18	17	46	27	19	185.9	219.5	152.6
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	3	0	3	2	0	2	8.1	-	16.1
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	1	0	1	2	1	1	8.1	8.1	8.0
Diseases of heart (390-398, 402, 404-429)	100	67	33	116	73	43	468.7	593.6	345.4
Hypertension (400, 401, 403)	1	0	1	1	0	1	4.0	-	8.0
Cerebrovascular disease (430-438)	44	24	20	49	27	22	198.0	219.5	176.7
Arteriosclerosis (440)	5	3	2	4	2	2	16.2	16.3	16.1
Influenza and pneumonia (470-486)	8	3	5	7	2	5	28.3	16.3	40.2
Bronchitis, emphysema, asthma (490-493)	2	1	1	4	3	1	16.2	24.4	8.0
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	1	0	4.0	8.1	-
Cirrhosis of liver (571)	2	1	1	2	1	1	8.1	8.1	8.0
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	1	0	4.0	8.1	-
Nephritis and nephrosis (580-584)	4	1	3	4	1	3	16.2	8.1	24.1
Infections of kidney (590)	4	2	2	4	2	2	16.2	16.3	16.1
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	1	0	1	1	0	1	20.9	-	30.3
Congenital anomalies (740-759)	1	0	1	2	1	1	8.1	8.1	8.0
Certain causes of mortality in early infancy (760-778)	7	3	4	16	5	11	33.5	33.8	33.3
Accidents (800-949)	27	9	18	33	13	20	133.3	105.7	160.6
Motor vehicle accidents (810-823)	7	3	4	13	8	5	52.5	65.0	40.2
Suicide (950-959)	2	1	1	4	2	2	16.2	16.3	16.1
Homicide (960-978)	4	0	4	5	0	5	20.2	-	40.2
Ill-defined and unknown (780-796)	14	2	12	14	2	12	56.6	16.3	96.4

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
COVINGTON COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	163	123	40	276	160	116	19.7	17.0	25.3
Fetal deaths	2	2	0	6	4	2	21.7	25.0	17.2
Total deaths	87	57	30	157	101	56	11.2	10.7	12.2
Deaths under 1 year	3	3	0	5	3	2	18.1	18.8	17.2
Deaths under 28 days	2	2	0	3	2	1	10.9	12.5	8.6
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	1	0	1	7.1	-	21.8
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	9	5	4	17	10	7	121.4	106.1	153.0
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	3	2	1	5	4	1	35.7	42.4	21.8
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	27	18	9	53	36	17	378.5	381.9	371.5
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	15	9	6	27	15	12	192.8	159.1	262.2
Arteriosclerosis (440)	0	0	0	0	0	0	-	-	-
Influenza and pneumonia (470-486)	5	3	2	11	7	4	78.6	74.3	87.4
Bronchitis, emphysema, asthma (490-493)	1	1	0	2	2	0	14.3	21.2	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	1	1	0	7.1	10.6	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	1	1	0	7.1	10.6	-
Certain causes of mortality in early infancy (760-778)	1	1	0	2	1	1	7.2	6.2	8.6
Accidents (800-949)	8	4	4	14	8	6	100.0	84.9	131.1
Motor vehicle accidents (810-823)	3	0	3	4	2	2	28.6	21.2	43.7
Suicide (950-959)	1	1	0	1	1	0	7.1	10.6	-
Homicide (960-978)	0	0	0	2	0	2	14.3	-	43.7
Ill-defined and unknown (780-796)	12	10	2	12	9	3	85.7	95.5	65.6

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
DESOTO COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	209	14	195	781	429	352	21.8	18.5	27.8
Fetal deaths	2	0	2	12	4	8	15.4	9.3	22.7
Total deaths	113	35	78	266	136	130	7.4	5.8	10.3
Deaths under 1 year	12	0	12	30	11	19	38.4	25.6	54.0
Deaths under 28 days	5	0	5	18	9	9	23.0	21.0	25.6
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	1	0	1	2.8	-	7.9
Tuberculosis (010-019)	1	0	1	1	0	1	2.8	-	7.9
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	11	2	9	36	22	14	100.3	94.7	110.7
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	1	0	1	5	1	4	13.9	4.3	31.6
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	1	0	1	2.8	-	7.9
Anemias (280-285)	0	0	0	1	0	1	2.8	-	7.9
Diseases of heart (390-398, 402, 404-429)	36	16	20	77	42	35	214.6	180.8	276.7
Hypertension (400, 401, 403)	0	0	0	2	2	0	5.6	8.6	-
Cerebrovascular disease (430-438)	10	3	7	30	13	17	83.6	56.0	134.4
Arteriosclerosis (440)	1	0	1	2	1	1	5.6	4.3	7.9
Influenza and pneumonia (470-486)	2	1	1	7	4	3	19.5	17.2	23.7
Bronchitis, emphysema, asthma (490-493)	0	0	0	6	6	0	16.7	25.8	-
Peptic ulcer (531-533)	0	0	0	1	1	0	2.8	4.3	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	0	1	2.8	-	7.9
Cirrhosis of liver (571)	0	0	0	1	1	0	2.8	4.3	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	0	1	2.8	-	7.9
Nephritis and nephrosis (580-584)	0	0	0	2	0	2	5.6	-	15.8
Infections of kidney (590)	0	0	0	1	0	1	2.8	-	7.9
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	0	1	4	1	3	11.1	4.3	23.7
Certain causes of mortality in early infancy (760-778)	4	0	4	14	8	6	17.9	18.6	17.0
Accidents (800-949)	16	5	11	25	13	12	69.7	56.0	94.9
Motor vehicle accidents (810-823)	7	3	4	11	8	3	30.6	34.4	23.7
Suicide (950-959)	2	2	0	3	2	1	8.4	8.6	7.9
Homicide (960-978)	1	0	1	2	1	1	5.6	4.3	7.9
Ill-defined and unknown (780-796)	21	5	16	22	7	15	61.3	30.1	118.6

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
FORREST COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,461	1,120	341	1,169	778	391	20.2	17.9	27.3
Fetal deaths	40	20	20	36	13	23	30.8	16.7	58.8
Total deaths	700	493	207	590	380	210	10.2	8.7	14.6
Deaths under 1 year	17	11	6	14	5	9	12.0	6.4	23.0
Deaths under 28 days	13	9	4	11	3	8	9.4	3.8	20.5
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	4	1	3	3	1	2	5.2	2.3	14.0
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	100	76	24	81	54	27	140.0	124.1	188.5
Benign and unspecified neoplasms (210-239)	0	0	0	1	1	0	1.7	2.3	-
Diabetes mellitus (250)	19	10	9	12	4	8	20.7	9.2	55.8
Avitaminoses and other nutritional deficiencies (260-269)	1	1	0	0	0	0	-	-	-
Anemias (280-285)	1	1	0	1	1	0	1.7	2.3	-
Diseases of heart (390-398, 402, 404-429)	271	198	73	221	155	66	382.0	356.1	460.7
Hypertension (400, 401, 403)	8	1	7	8	1	7	13.8	2.3	48.9
Cerebrovascular disease (430-438)	82	67	15	63	51	12	108.9	117.2	83.8
Arteriosclerosis (440)	6	5	1	4	2	2	6.9	4.6	14.0
Influenza and pneumonia (470-486)	28	22	6	23	17	6	39.8	39.0	41.9
Bronchitis, emphysema, asthma (490-493)	15	14	1	12	11	1	20.7	25.3	7.0
Peptic ulcer (531-533)	4	4	0	3	3	0	5.2	6.9	-
Hernia and intestinal obstruction (550-553, 560)	3	2	1	2	1	1	3.4	2.3	7.0
Cirrhosis of liver (571)	5	5	0	4	3	1	6.9	6.9	7.0
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	8	2	6	10	1	9	17.3	2.3	62.8
Infections of kidney (590)	4	2	2	4	2	2	6.9	4.6	14.0
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	2	2	0	1	1	0	1.7	2.3	-
Certain causes of mortality in early infancy (760-778)	8	6	2	9	2	7	7.7	2.6	17.9
Accidents (800-949)	48	35	13	52	33	19	89.9	75.8	132.6
Motor vehicle accidents (810-823)	24	18	6	25	17	8	43.2	39.0	55.8
Suicide (950-959)	8	7	1	8	7	1	13.8	16.1	7.0
Homicide (960-978)	8	1	7	6	1	5	10.4	2.3	34.9
Ill-defined and unknown (780-796)	33	10	23	33	10	23	57.0	23.0	160.6

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
FRANKLIN COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	15	2	13	123	56	67	15.4	11.4	21.5
Fetal deaths	0	0	0	1	0	1	8.1	-	14.9
Total deaths	80	46	34	106	58	48	13.2	11.8	15.4
Deaths under 1 year	1	0	1	2	0	2	16.3	-	29.8
Deaths under 28 days	0	0	0	1	0	1	8.1	-	14.9
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	1	0	1	12.5	-	32.1
Tuberculosis (010-019)	0	0	0	1	1	0	12.5	20.4	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	10	8	2	18	9	9	224.7	183.8	289.0
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	4	1	3	2	0	2	25.0	-	64.2
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	27	13	14	33	19	14	411.9	388.0	449.6
Hypertension (400, 401, 403)	2	0	2	2	1	1	25.0	20.4	32.1
Cerebrovascular disease (430-438)	12	8	4	13	10	3	162.3	204.2	96.3
Arteriosclerosis (440)	5	3	2	9	5	4	112.3	102.1	128.4
Influenza and pneumonia (470-486)	0	0	0	0	0	0	-	-	-
Bronchitis, emphysema, asthma (490-493)	0	0	0	0	0	0	-	-	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	1	1	0	1	1	0	12.5	20.4	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	0	0	0	1	0	1	8.1	-	14.9
Accidents (800-949)	10	7	3	13	6	7	162.3	122.5	224.8
Motor vehicle accidents (810-823)	2	2	0	2	1	1	25.0	20.4	32.1
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	1	0	1	2	0	2	25.0	-	64.2
Ill-defined and unknown (780-796)	1	0	1	2	1	1	25.0	20.4	32.1

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
GEORGE COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence						
	Number			Number			Rate*			
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	
Live Births, Fetal Deaths, Deaths										
Live births	293	254	39	279	238	41	22.4	21.6	28.2	
Fetal deaths	6	4	2	8	5	3	28.7	21.0	73.2	
Total deaths	109	93	16	116	105	11	9.3	9.5	7.6	
Deaths under 1 year	5	4	1	9	8	1	32.2	33.6	24.4	
Deaths under 28 days	3	2	1	6	6	0	21.5	25.2	-	
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-	
Enteritis and other diarrheal diseases (008, 009)	0	0	0	1	0	1	8.0	-	68.9	
Tuberculosis (010-019)	1	1	0	1	1	0	8.0	9.1	-	
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-	
Syphilis (090-097)	0	0	0	0	0	0	-	-	-	
Malignant neoplasms (140-209)	18	15	3	19	17	2	152.5	154.4	137.7	
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-	
Diabetes mellitus (250)	4	4	0	5	5	0	40.1	45.4	-	
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-	
Anemias (280-285)	0	0	0	0	0	0	-	-	-	
Diseases of heart (390-398, 402, 404-429)	38	34	4	38	35	3	305.0	318.0	206.6	
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-	
Cerebrovascular disease (430-438)	7	7	0	6	6	0	48.2	54.5	-	
Arteriosclerosis (440)	2	1	1	1	0	1	8.0	-	68.9	
Influenza and pneumonia (470-486)	1	0	1	1	1	0	8.0	9.1	-	
Bronchitis, emphysema, asthma (490-493)	2	2	0	1	1	0	8.0	9.1	-	
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-	
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-	
Cirrhosis of liver (571)	1	1	0	1	1	0	8.0	9.1	-	
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-	
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-	
Infections of kidney (590)	0	0	0	0	0	0	-	-	-	
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-	
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-	
Congenital anomalies (740-759)	6	5	1	8	7	1	64.2	63.6	68.9	
Certain causes of mortality in early infancy (760-778)	3	2	1	4	4	0	14.3	16.8	-	
Accidents (800-949)	23	19	4	22	21	1	176.6	190.8	68.9	
Motor vehicle accidents (810-823)	17	14	3	16	16	0	128.4	145.4	-	
Suicide (950-959)	0	0	0	0	0	0	-	-	-	
Homicide (960-978)	1	0	1	1	0	1	8.0	-	68.9	
Ill-defined and unknown (780-796)	1	1	0	1	1	0	8.0	9.1	-	

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
GREENE COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	75	36	39	186	129	57	21.8	19.4	30.3
Fetal deaths	2	0	2	3	1	2	16.1	7.8	35.1
Total deaths	57	41	16	101	73	28	11.8	11.0	14.9
Deaths under 1 year	5	1	4	10	3	7	53.8	23.2	122.8
Deaths under 28 days	3	1	2	7	3	4	37.6	23.2	70.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	1	1	0	11.7	15.0	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	3	3	0	11	11	0	128.7	165.1	-
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	0	0	0	0	0	0	-	-	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	27	19	8	40	29	11	468.1	435.3	584.2
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	4	4	0	7	6	1	81.9	90.1	53.1
Arteriosclerosis (440)	1	1	0	2	2	0	23.4	30.0	-
Influenza and pneumonia (470-486)	3	1	2	5	3	2	58.5	45.0	106.2
Bronchitis, emphysema, asthma (490-493)	0	0	0	0	0	0	-	-	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	2	2	0	2	2	0	23.4	30.0	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	1	1	0	1	1	0	11.7	15.0	-
Maternal causes (630-678)	1	0	1	1	0	1	53.8	-	175.4
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	3	1	2	7	3	4	37.6	23.2	70.2
Accidents (800-949)	4	4	0	11	5	6	128.7	75.0	318.6
Motor vehicle accidents (810-823)	4	4	0	8	4	4	93.6	60.0	212.4
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	1	0	1	1	0	1	11.7	-	53.1

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
GRENADA COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	426	163	263	446	187	259	22.5	16.8	29.8
Fetal deaths	19	6	13	22	7	15	49.3	37.4	57.9
Total deaths	271	170	101	219	116	103	11.0	10.4	11.8
Deaths under 1 year	15	5	10	15	5	10	33.6	26.7	38.6
Deaths under 28 days	10	4	6	10	4	6	22.4	21.4	23.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	3	1	2	2	0	2	10.1	-	23.0
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	30	21	9	30	21	9	151.1	188.3	103.4
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	2	0	2	2	0	2	10.1	-	23.0
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	2	2	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	84	60	24	59	37	22	297.2	331.7	252.9
Hypertension (400, 401, 403)	1	0	1	1	0	1	5.0	-	11.5
Cerebrovascular disease (430-438)	65	37	28	49	23	26	246.8	206.2	298.8
Arteriosclerosis (440)	4	3	1	1	0	1	5.0	-	11.5
Influenza and pneumonia (470-486)	10	7	3	10	7	3	50.4	62.8	34.5
Bronchitis, emphysema, asthma (490-493)	2	2	0	0	0	0	-	-	-
Peptic ulcer (531-533)	1	1	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	1	1	0	0	0	0	-	-	-
Cirrhosis of liver (571)	2	2	0	4	3	1	20.1	26.9	11.5
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	1	0	0	0	0	-	-	-
Infections of kidney (590)	1	1	0	3	1	2	15.1	9.0	23.0
Hyperplasia of prostate (600)	0	0	0	1	0	1	5.0	-	11.5
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	0	1	2	1	1	10.1	9.0	11.5
Certain causes of mortality in early infancy (760-778)	6	4	2	4	2	2	9.0	10.7	7.7
Accidents (800-949)	19	7	12	15	5	10	75.6	44.8	114.9
Motor vehicle accidents (810-823)	7	3	4	4	2	2	20.1	17.9	23.0
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	1	1	0	1	1	0	5.0	9.0	-
Ill-defined and unknown (780-796)	21	9	12	21	8	13	105.8	71.7	149.4

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
HANCOCK COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	77	53	24	268	227	41	15.4	15.2	16.4
Fetal deaths	0	0	0	8	8	0	29.8	35.2	-
Total deaths	166	147	19	202	179	23	11.6	12.0	9.2
Deaths under 1 year	3	3	0	6	6	0	22.4	26.4	-
Deaths under 28 days	1	1	0	4	4	0	14.9	17.6	-
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	1	1	0	5.8	6.7	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	19	18	1	33	32	1	189.8	214.8	40.1
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	6	6	0	6	6	0	34.5	40.3	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	49	42	7	57	51	6	327.8	342.4	240.7
Hypertension (400, 401, 403)	0	0	0	1	0	1	5.8	-	40.1
Cerebrovascular disease (430-438)	20	18	2	26	21	5	149.5	141.0	200.6
Arteriosclerosis (440)	2	2	0	4	4	0	23.0	26.8	-
Influenza and pneumonia (470-486)	4	3	1	6	5	1	34.5	33.6	40.1
Bronchitis, emphysema, asthma (490-493)	0	0	0	2	2	0	11.5	13.4	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	3	3	0	4	4	0	23.0	26.8	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	1	0	1	1	0	5.8	6.7	-
Infections of kidney (590)	2	2	0	3	3	0	17.2	20.1	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	1	1	0	3	3	0	11.2	13.2	-
Accidents (800-949)	25	22	3	22	18	4	126.5	120.8	160.4
Motor vehicle accidents (810-823)	14	14	0	10	9	1	57.5	60.4	40.1
Suicide (950-959)	1	1	0	1	1	0	5.8	6.7	-
Homicide (960-978)	1	0	1	1	0	1	5.8	-	40.1
Ill-defined and unknown (780-796)	26	22	4	23	19	4	132.3	127.6	160.4

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
HARRISON COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	3,548	2,837	711	3,171	2,484	687	23.6	22.4	29.2
Fetal deaths	300	243	57	258	205	53	81.4	82.5	77.1
Total deaths	1,349	1,072	277	1,210	951	259	9.0	8.6	11.0
Deaths under 1 year	80	59	21	74	52	22	23.3	20.9	32.0
Deaths under 28 days	61	47	14	56	43	13	17.7	17.3	18.9
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	6	4	2	7	5	2	5.2	4.5	8.5
Tuberculosis (010-019)	2	2	0	4	3	1	3.0	2.7	4.2
Meningococcal infection (036)	1	1	0	2	2	0	1.5	1.8	-
Syphilis (090-097)	1	1	0	1	1	0	0.7	0.9	-
Malignant neoplasms (140-209)	194	169	25	177	155	22	131.5	139.6	93.5
Benign and unspecified neoplasms (210-239)	3	2	1	1	1	0	0.7	0.9	-
Diabetes mellitus (250)	17	12	5	18	12	6	13.4	10.8	25.5
Avitaminoses and other nutritional deficiencies (260-269)	1	1	0	1	1	0	0.7	0.9	-
Anemias (280-285)	5	2	3	4	1	3	3.0	0.9	12.8
Diseases of heart (390-398, 402, 404-429)	434	372	62	364	311	53	270.5	280.0	225.3
Hypertension (400, 401, 403)	6	2	4	4	1	3	3.0	0.9	12.8
Cerebrovascular disease (430-438)	112	82	30	111	83	28	82.5	74.7	119.0
Arteriosclerosis (440)	4	4	0	5	5	0	3.7	4.5	-
Influenza and pneumonia (470-486)	40	31	9	37	26	11	27.5	23.4	46.8
Bronchitis, emphysema, asthma (490-493)	36	31	5	21	17	4	15.6	15.3	17.0
Peptic ulcer (531-533)	9	8	1	8	7	1	5.9	6.3	4.2
Hernia and intestinal obstruction (550-553, 560)	4	3	1	4	3	1	3.0	2.7	4.2
Cirrhosis of liver (571)	13	12	1	13	12	1	9.6	10.8	4.2
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	3	3	0	3	3	0	2.2	2.7	-
Nephritis and nephrosis (580-584)	8	4	4	6	2	4	4.4	1.8	17.0
Infections of kidney (590)	6	3	3	6	3	3	4.4	2.7	12.8
Hyperplasia of prostate (600)	1	1	0	1	1	0	0.7	0.9	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	17	16	1	14	13	1	10.4	11.7	4.2
Certain causes of mortality in early infancy (760-778)	44	30	14	40	27	13	12.6	10.9	18.9
Accidents (800-949)	234	181	53	201	155	46	149.4	139.6	195.6
Motor vehicle accidents (810-823)	55	48	7	44	35	9	32.7	31.5	38.3
Suicide (950-959)	14	12	2	11	9	2	8.2	8.1	8.5
Homicide (960-978)	15	9	6	15	10	5	11.1	9.0	21.2
Ill-defined and unknown (780-796)	45	20	25	50	24	26	37.2	21.6	110.5

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
HINDS COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	5,904	2,934	2,970	4,480	2,132	2,348	20.8	16.3	27.8
Fetal deaths	644	339	305	464	232	232	103.6	108.8	98.8
Total deaths	2,791	1,608	1,183	1,843	1,047	796	8.6	8.0	9.4
Deaths under 1 year	214	75	139	134	47	87	29.9	22.0	37.0
Deaths under 28 days	159	60	99	108	42	66	24.1	19.7	28.1
Bacillary dysentery and amebiasis (004, 006)	1	0	1	1	1	0	0.5	0.8	-
Enteritis and other diarrheal diseases (008, 009)	8	4	4	4	4	0	1.9	3.1	-
Tuberculosis (010-019)	6	4	2	5	3	2	2.3	2.3	2.4
Meningococcal infection (036)	3	1	2	2	1	1	0.9	0.8	1.2
Syphilis (090-097)	1	0	1	1	0	1	0.5	-	1.2
Malignant neoplasms (140-209)	522	344	178	297	198	99	138.2	151.6	117.3
Benign and unspecified neoplasms (210-239)	8	4	4	4	2	2	1.9	1.5	2.4
Diabetes mellitus (250)	47	21	26	39	16	23	18.1	12.2	27.2
Avitaminoses and other nutritional deficiencies (260-269)	2	0	2	2	0	2	0.9	-	2.4
Anemias (280-285)	8	5	3	3	0	3	1.4	-	3.6
Diseases of heart (390-398, 402, 404-429)	724	501	223	521	370	151	242.4	283.3	179.0
Hypertension (400, 401, 403)	14	4	10	6	1	5	2.8	0.8	5.9
Cerebrovascular disease (430-438)	246	138	108	168	95	73	78.1	72.7	86.5
Arteriosclerosis (440)	22	18	4	18	15	3	8.4	11.5	3.6
Influenza and pneumonia (470-486)	62	37	25	48	27	21	22.3	20.7	24.9
Bronchitis, emphysema, asthma (490-493)	36	23	13	21	12	9	9.8	9.2	10.7
Peptic ulcer (531-533)	18	15	3	6	5	1	2.8	3.8	1.2
Hernia and intestinal obstruction (550-553, 560)	9	3	6	4	2	2	1.9	1.5	2.4
Cirrhosis of liver (571)	28	18	10	12	10	2	5.6	7.6	2.4
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	7	7	0	1	1	0	0.5	0.8	-
Nephritis and nephrosis (580-584)	23	12	11	8	1	7	3.7	0.8	8.3
Infections of kidney (590)	18	12	6	14	7	7	6.5	5.4	8.3
Hyperplasia of prostate (600)	6	4	2	3	3	0	1.4	2.3	-
Maternal causes (630-678)	4	0	4	2	0	2	4.5	-	8.5
Congenital anomalies (740-759)	48	32	16	20	14	6	9.3	10.7	7.1
Certain causes of mortality in early infancy (760-778)	110	41	69	82	31	51	18.3	14.5	21.7
Accidents (800-949)	245	128	117	139	67	72	64.6	51.3	85.3
Motor vehicle accidents (810-823)	125	71	54	73	36	37	34.0	27.6	43.8
Suicide (950-959)	22	21	1	18	18	0	8.4	13.8	-
Homicide (960-978)	41	11	30	31	11	20	14.4	8.4	23.7
Ill-defined and unknown (780-796)	221	58	163	206	54	152	95.8	41.4	180.1

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
HOLMES COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	484	80	404	512	104	408	22.1	14.2	25.9
Fetal deaths	33	3	30	36	3	33	70.3	28.8	80.9
Total deaths	268	78	190	310	108	202	13.4	14.7	12.8
Deaths under 1 year	15	2	13	20	3	17	39.1	28.8	41.7
Deaths under 28 days	10	2	8	12	3	9	23.4	28.8	22.0
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	2	0	2	4	0	4	17.3	-	25.4
Tuberculosis (010-019)	0	0	0	1	0	1	4.3	-	6.3
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	33	12	21	39	20	19	168.7	272.3	120.4
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	6	2	4	9	3	6	38.9	40.8	38.0
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	2	0	2	1	0	1	4.3	-	6.3
Diseases of heart (390-398, 402, 404-429)	95	36	59	112	48	64	484.4	653.5	405.7
Hypertension (400, 401, 403)	6	1	5	6	1	5	26.0	13.6	31.7
Cerebrovascular disease (430-438)	47	8	39	44	9	35	190.3	122.5	221.9
Arteriosclerosis (440)	0	0	0	0	0	0	-	-	-
Influenza and pneumonia (470-486)	2	0	2	3	0	3	13.0	-	19.0
Bronchitis, emphysema, asthma (490-493)	3	2	1	5	2	3	21.6	27.2	19.0
Peptic ulcer (531-533)	0	0	0	1	0	1	4.3	-	6.3
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	1	1	0	4.3	13.6	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	1	0	4.3	13.6	-
Nephritis and nephrosis (580-584)	5	0	5	4	0	4	17.3	-	25.4
Infections of kidney (590)	2	0	2	1	0	1	4.3	-	6.3
Hyperplasia of prostate (600)	1	0	1	1	0	1	4.3	-	6.3
Maternal causes (630-678)	3	0	3	3	0	3	58.6	-	73.5
Congenital anomalies (740-759)	1	1	0	1	1	0	4.3	13.6	-
Certain causes of mortality in early infancy (760-778)	6	1	5	7	1	6	13.7	9.6	14.7
Accidents (800-949)	9	2	7	15	3	12	64.9	40.8	76.1
Motor vehicle accidents (810-823)	2	0	2	5	0	5	21.6	-	31.7
Suicide (950-959)	4	4	0	4	4	0	17.3	54.4	-
Homicide (960-978)	3	0	3	3	0	3	13.0	-	19.0
Ill-defined and unknown (780-796)	28	3	25	24	3	21	103.8	40.8	133.1

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
HUMPHREYS COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	374	46	328	371	83	288	25.4	16.3	30.3
Fetal deaths	14	2	12	22	5	17	59.3	60.2	59.0
Total deaths	161	45	116	199	67	132	13.6	13.2	13.9
Deaths under 1 year	16	1	15	21	3	18	56.6	36.1	62.5
Deaths under 28 days	11	0	11	15	2	13	40.4	24.1	45.1
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diases (008, 009)	2	0	2	2	0	2	13.7	-	21.0
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	21	10	11	30	14	16	205.5	275.1	168.2
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	2	2	0	3	2	1	20.5	39.3	10.5
Avitaminoses and other nutritional deficiencies (260-269)	1	0	1	1	0	1	6.8	-	10.5
Anemias (280-285)	0	0	0	1	1	0	6.8	19.6	-
Diseases of heart (390-398, 402, 404-429)	66	18	48	73	24	49	500.0	471.6	515.1
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	20	3	17	25	7	18	171.2	137.6	189.2
Arteriosclerosis (440)	1	0	1	0	0	0	-	-	-
Influenza and pneumonia (470-486)	1	0	1	2	1	1	13.7	19.6	10.5
Bronchitis, emphysema, asthma (490-493)	1	1	0	2	1	1	13.7	19.6	10.5
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	1	0	1	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	1	0	1	6.8	-	10.5
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	0	1	1	0	1	6.8	-	10.5
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	1	0	1	6.8	-	10.5
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	2	1	1	2	1	1	13.7	19.6	10.5
Certain causes of mortality in early infancy (760-778)	9	0	9	12	2	10	32.3	24.1	34.7
Accidents (800-949)	12	6	6	15	9	6	102.7	176.8	63.1
Motor vehicle accidents (810-823)	7	3	4	6	3	3	41.1	59.0	31.5
Suicide (950-959)	2	2	0	2	2	0	13.7	39.3	-
Homicide (960-978)	4	1	3	4	1	3	27.4	19.6	31.5
Ill-defined and unknown (780-796)	12	1	11	14	1	13	95.9	19.6	136.7

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
ISSAQUENA COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non- white	Total	White	Non- white	Total	White	Non- white
Live births	12	0	12	56	18	38	20.5	17.4	22.3
Fetal deaths	1	0	1	2	1	1	35.7	55.6	26.3
Total deaths	19	6	13	23	8	15	8.4	7.7	8.8
Deaths under 1 year	2	0	2	3	0	3	53.6	-	78.9
Deaths under 28 days	1	0	1	1	0	1	17.8	-	26.3
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	0	0	0	2	0	2	73.1	-	117.4
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	0	0	0	0	0	0	-	-	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	6	3	3	7	4	3	255.8	387.2	176.0
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	2	0	2	4	1	3	146.1	96.8	176.0
Arteriosclerosis (440)	0	0	0	0	0	0	-	-	-
Influenza and pneumonia (470-486)	0	0	0	0	0	0	-	-	-
Bronchitis, emphysema, asthma (490-493)	0	0	0	0	0	0	-	-	-
Peptic ulcer (531-533)	0	0	0	1	0	1	36.5	-	58.7
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	1	0	1	1	0	1	36.5	-	58.7
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	0	1	1	0	1	36.5	-	58.7
Certain causes of mortality in early infancy (760-778)	0	0	0	0	0	0	-	-	-
Accidents (800-949)	4	2	2	2	2	0	73.1	193.6	-
Motor vehicle accidents (810-823)	3	1	2	1	1	0	36.5	96.8	-
Suicide (950-959)	1	0	1	1	0	1	36.5	-	58.7
Homicide (960-978)	1	0	1	1	0	1	36.5	-	58.7
Ill-defined and unknown (780-796)	1	0	1	2	0	2	73.1	-	117.4

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
ITAWAMBA COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	33	25	8	244	224	20	14.5	14.1	20.7
Fetal deaths	2	2	0	6	6	0	24.6	26.8	-
Total deaths	133	126	7	169	159	10	10.0	10.0	10.4
Deaths under 1 year	3	3	0	3	3	0	12.3	13.4	-
Deaths under 28 days	2	2	0	1	1	0	4.1	4.5	-
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	18	18	0	24	24	0	142.4	151.1	-
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	1	1	0	1	1	0	5.9	6.3	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	59	56	3	68	63	5	403.6	396.7	518.1
Hypertension (400, 401, 403)	1	1	0	1	1	0	5.9	6.3	-
Cerebrovascular disease (430-438)	25	23	2	29	27	2	172.1	170.0	207.2
Arteriosclerosis (440)	4	4	0	4	4	0	23.7	25.2	-
Influenza and pneumonia (470-486)	1	1	0	6	5	1	35.6	31.5	103.6
Bronchitis, emphysema, asthma (490-493)	2	2	0	2	2	0	11.9	12.6	-
Peptic ulcer (531-533)	1	1	0	1	1	0	5.9	6.3	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	2	2	0	2	2	0	11.9	12.6	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	2	2	0	11.9	12.6	-
Nephritis and nephrosis (580-584)	1	1	0	1	1	0	5.9	6.3	-
Infections of kidney (590)	0	0	0	2	2	0	11.9	12.6	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	2	2	0	1	1	0	4.1	4.5	-
Accidents (800-949)	8	6	2	13	11	2	77.2	69.3	207.2
Motor vehicle accidents (810-823)	4	2	2	6	4	2	35.6	25.2	207.2
Suicide (950-959)	1	1	0	1	1	0	5.9	6.3	-
Homicide (960-978)	1	1	0	1	1	0	5.9	6.3	-
Ill-defined and unknown (780-796)	2	2	0	2	2	0	11.9	12.6	-

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
JACKSON COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,743	1,287	456	2,060	1,582	478	23.4	21.5	33.1
Fetal deaths	37	22	15	72	55	17	35.0	34.8	35.6
Total deaths	506	379	127	568	430	138	6.4	5.8	9.6
Deaths under 1 year	45	30	15	54	37	17	26.2	23.4	35.6
Deaths under 28 days	36	25	11	43	30	13	20.9	19.0	27.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	1	1	0	2	2	0	2.3	2.7	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	68	53	15	84	64	20	95.5	87.0	138.6
Benign and unspecified neoplasms (210-239)	1	1	0	2	2	0	2.3	2.7	-
Diabetes mellitus (250)	12	10	2	13	11	2	14.8	15.0	13.9
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	1	0	1	1	0	1	1.1	-	6.9
Diseases of heart (390-398, 402, 404-429)	156	118	38	178	140	38	202.3	190.4	263.4
Hypertension (400, 401, 403)	7	2	5	8	3	5	9.1	4.1	34.6
Cerebrovascular disease (430-438)	47	35	12	56	39	17	63.6	53.0	117.8
Arteriosclerosis (440)	7	5	2	7	5	2	8.0	6.8	13.9
Influenza and pneumonia (470-486)	17	10	7	19	12	7	21.6	16.3	48.5
Bronchitis, emphysema, asthma (490-493)	7	7	0	10	10	0	11.4	13.6	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	2	1	1	2	1	1	2.3	1.4	6.9
Cirrhosis of liver (571)	1	0	1	1	0	1	1.1	-	6.9
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	1	1	0	1.1	1.4	-
Nephritis and nephrosis (580-584)	7	6	1	7	5	2	8.0	6.8	13.9
Infections of kidney (590)	3	3	0	2	2	0	2.3	2.7	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	5	4	1	7	6	1	8.0	8.2	6.9
Certain causes of mortality in early infancy (760-778)	29	20	9	34	24	10	16.5	15.2	20.9
Accidents (800-949)	55	42	13	55	44	11	62.5	59.8	76.2
Motor vehicle accidents (810-823)	21	17	4	22	19	3	25.0	25.8	20.8
Suicide (950-959)	2	2	0	2	2	0	2.3	2.7	-
Homicide (960-978)	2	1	1	4	2	2	4.5	2.7	13.9
Ill-defined and unknown (780-796)	39	28	11	33	23	10	37.5	31.3	69.3

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
JASPER COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	189	131	58	330	124	206	20.6	14.5	27.7
Fetal deaths	3	2	1	6	1	5	18.2	8.1	24.3
Total deaths	124	63	61	192	105	87	12.0	12.3	11.7
Deaths under 1 year	3	2	1	7	3	4	21.2	24.2	19.4
Deaths under 28 days	3	2	1	6	3	3	18.2	24.2	14.6
Bacillary dysentery and amebiasis (004, 006)	1	0	1	1	0	1	6.2	-	13.4
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	15	8	7	31	22	9	193.8	256.9	121.1
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	1	1	0	4	1	3	25.0	11.7	40.4
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	53	30	23	57	33	24	356.4	385.3	323.0
Hypertension (400, 401, 403)	0	0	0	1	1	0	6.2	11.7	-
Cerebrovascular disease (430-438)	21	10	11	26	15	11	162.6	175.2	148.0
Arteriosclerosis (440)	0	0	0	4	3	1	25.0	35.0	13.4
Influenza and pneumonia (470-486)	1	1	0	3	1	2	18.8	11.7	26.9
Bronchitis, emphysema, asthma (490-493)	5	4	1	4	3	1	25.0	35.0	13.4
Peptic ulcer (531-533)	0	0	0	2	1	1	12.5	11.7	13.4
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	1	0	6.2	11.7	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	1	0	4	2	2	25.0	23.4	26.9
Infections of kidney (590)	1	0	1	2	1	1	12.5	11.7	13.4
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	1	1	0	6.2	11.7	-
Certain causes of mortality in early infancy (760-778)	2	1	1	5	2	3	15.2	16.1	14.6
Accidents (800-949)	4	1	3	20	8	12	125.0	93.4	161.5
Motor vehicle accidents (810-823)	1	0	1	12	3	9	75.0	35.0	121.1
Suicide (950-959)	1	1	0	1	1	0	6.2	11.7	-
Homicide (960-978)	1	0	1	3	0	3	18.8	-	40.4
Ill-defined and unknown (780-796)	9	2	7	11	3	8	68.8	35.0	107.7

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
JEFFERSON COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live Births, Fetal Deaths, Deaths									
Live births	56	7	49	160	21	139	17.2	9.1	19.8
Fetal deaths	0	0	0	5	0	5	31.2	-	36.0
Total deaths	84	21	63	122	34	88	13.1	14.8	12.6
Deaths under 1 year	2	0	2	6	0	6	37.5	-	43.2
Deaths under 28 days	0	0	0	3	0	3	18.8	-	21.6
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	11	3	8	12	4	8	129.1	174.2	114.3
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	1	0	1	2	0	2	21.5	-	28.6
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	1	0	1	10.8	-	14.3
Diseases of heart (390-398, 402, 404-429)	40	13	27	50	17	33	537.9	740.4	471.5
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	5	0	5	13	3	10	139.9	130.7	142.9
Arteriosclerosis (440)	0	0	0	0	0	0	-	-	-
Influenza and pneumonia (470-486)	1	0	1	5	1	4	53.8	43.6	57.2
Bronchitis, emphysema, asthma (490-493)	1	1	0	1	1	0	10.8	43.6	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	2	1	1	21.5	43.6	14.3
Cirrhosis of liver (571)	1	1	0	2	1	1	21.5	43.6	14.3
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	0	1	2	0	2	21.5	-	28.6
Infections of kidney (590)	0	0	0	1	1	0	10.8	43.6	-
Hyperplasia of prostate (600)	0	0	0	1	0	1	10.8	-	14.3
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	0	1	2	0	2	21.5	-	28.6
Certain causes of mortality in early infancy (760-778)	0	0	0	2	0	2	12.5	-	14.4
Accidents (800-949)	10	2	8	10	2	8	107.6	87.1	114.3
Motor vehicle accidents (810-823)	8	2	6	6	2	4	64.6	87.1	57.2
Suicide (950-959)	1	0	1	1	0	1	10.8	-	14.3
Homicide (960-978)	2	0	2	1	0	1	10.8	-	14.3
Ill-defined and unknown (780-796)	8	0	8	11	1	10	118.3	43.6	142.9

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
JEFF DAVIS COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	166	86	80	292	112	180	22.6	17.4	27.7
Fetal deaths	6	1	5	12	2	10	41.1	17.8	55.6
Total deaths	146	65	81	161	65	96	12.4	10.1	14.8
Deaths under 1 year	7	2	5	11	3	8	37.7	26.8	44.4
Deaths under 28 days	3	2	1	9	3	6	30.8	26.8	33.3
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	1	0	1	7.7	-	15.4
Tuberculosis (010-019)	1	0	1	1	0	1	7.7	-	15.4
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	17	8	9	19	8	11	146.9	124.3	169.2
Benign and unspecified neoplasms (210-239)	1	0	1	1	0	1	7.7	-	15.4
Diabetes mellitus (250)	6	4	2	4	2	2	30.9	31.1	30.8
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	52	26	26	51	21	30	394.2	326.3	461.5
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	33	12	21	34	13	21	262.8	202.0	323.0
Arteriosclerosis (440)	0	0	0	1	1	0	7.7	15.5	-
Influenza and pneumonia (470-486)	3	1	2	4	3	1	30.9	46.6	15.4
Bronchitis, emphysema, asthma (490-493)	0	0	0	2	1	1	15.5	15.5	15.4
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	2	1	1	2	1	1	15.5	15.5	15.4
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	1	0	2	1	1	15.5	15.5	15.4
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	1	0	1	1	0	1	7.7	-	15.4
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	2	1	1	15.5	15.5	15.4
Certain causes of mortality in early infancy (760-778)	3	2	1	8	3	5	27.4	26.8	27.8
Accidents (800-949)	9	3	6	15	7	8	116.0	108.8	123.0
Motor vehicle accidents (810-823)	2	1	1	7	4	3	54.1	62.2	46.1
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	1	0	1	1	0	1	7.7	-	15.4
Ill-defined and unknown (780-796)	8	2	6	7	1	6	54.1	15.5	92.3

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
JONES COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	2,003	850	1,153	990	659	331	17.6	15.5	23.7
Fetal deaths	73	26	47	39	21	18	39.4	31.9	54.4
Total deaths	771	493	278	596	417	179	10.6	9.8	12.8
Deaths under 1 year	73	22	51	36	16	20	36.4	24.3	60.4
Deaths under 28 days	60	21	39	29	15	14	29.3	22.8	42.3
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	9	1	8	5	1	4	8.9	2.4	28.7
Tuberculosis (010-019)	1	1	0	1	1	0	1.8	2.4	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	106	81	25	90	70	20	159.7	165.1	143.3
Benign and unspecified neoplasms (210-239)	2	2	0	1	1	0	1.8	2.4	-
Diabetes mellitus (250)	25	11	14	15	10	5	26.6	23.6	35.8
Avitaminoses and other nutritional deficiencies (260-269)	2	1	1	3	2	1	5.3	4.7	7.2
Anemias (280-285)	1	0	1	1	0	1	1.8	-	7.2
Diseases of heart (390-398, 402, 404-429)	196	142	54	185	138	47	328.3	325.4	336.8
Hypertension (400, 401, 403)	7	2	5	4	2	2	7.1	4.7	14.3
Cerebrovascular disease (430-438)	93	58	35	71	46	25	126.0	108.5	179.2
Arteriosclerosis (440)	22	17	5	14	12	2	24.8	28.3	14.3
Influenza and pneumonia (470-486)	21	10	11	13	6	7	23.1	14.1	50.2
Bronchitis, emphysema, asthma (490-493)	8	8	0	7	6	1	12.4	14.1	7.2
Peptic ulcer (531-533)	4	2	2	2	1	1	3.5	2.4	7.2
Hernia and intestinal obstruction (550-553, 560)	2	1	1	1	0	1	1.8	-	7.2
Cirrhosis of liver (571)	5	4	1	5	5	0	8.9	11.8	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	3	3	0	2	2	0	3.5	4.7	-
Nephritis and nephrosis (580-584)	20	6	14	10	5	5	17.7	11.8	35.8
Infections of kidney (590)	2	1	1	1	1	0	1.8	2.4	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	1	0	1	0	0	0	-	-	-
Congenital anomalies (740-759)	2	1	1	4	3	1	7.1	7.1	7.2
Certain causes of mortality in early infancy (760-778)	52	18	34	23	11	12	23.2	16.7	36.2
Accidents (800-949)	66	50	16	38	27	11	67.4	63.7	78.8
Motor vehicle accidents (810-823)	37	30	7	18	14	4	31.9	33.0	28.7
Suicide (950-959)	6	5	1	5	5	0	8.9	11.8	-
Homicide (960-978)	13	5	8	8	5	3	14.2	11.8	21.5
Ill-defined and unknown (780-796)	63	38	25	54	32	22	95.8	75.5	157.7

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
KEMPER COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	68	10	58	221	57	164	21.6	12.7	28.5
Fetal deaths	2	0	2	10	5	5	45.2	87.7	30.5
Total deaths	82	41	41	117	60	57	11.4	13.4	9.9
Deaths under 1 year	3	0	3	9	3	6	40.7	52.6	36.6
Deaths under 28 days	1	0	1	5	2	3	22.6	35.1	18.3
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	1	0	1	9.8	-	17.4
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	10	5	5	16	10	6	156.4	222.8	104.4
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	2	1	1	3	1	2	29.3	22.3	34.8
Avitaminoses and other nutritional deficiencies (260-269)	1	0	1	1	0	1	9.8	-	17.4
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	47	27	20	60	36	24	586.3	802.1	417.8
Hypertension (400, 401, 403)	0	0	0	2	2	0	19.5	44.6	-
Cerebrovascular disease (430-438)	5	2	3	10	4	6	97.7	89.1	104.4
Arteriosclerosis (440)	0	0	0	0	0	0	-	-	-
Influenza and pneumonia (470-486)	1	0	1	3	1	2	29.3	22.3	34.8
Bronchitis, emphysema, asthma (490-493)	0	0	0	0	0	0	-	-	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	0	0	0	4	2	2	18.1	35.1	12.2
Accidents (800-949)	7	5	2	4	0	4	39.1	-	69.6
Motor vehicle accidents (810-823)	3	3	0	2	0	2	19.5	-	34.8
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	2	0	2	1	0	1	9.8	-	17.4
Ill-defined and unknown (780-796)	5	0	5	6	0	6	58.6	-	104.4

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
LAFAYETTE COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	545	340	205	466	285	181	19.3	16.5	26.4
Fetal deaths	7	1	6	6	2	4	12.9	7.0	22.1
Total deaths	238	164	74	214	134	80	8.8	7.7	11.6
Deaths under 1 year	10	3	7	9	3	6	19.3	10.5	33.1
Deaths under 28 days	7	3	4	5	2	3	10.7	7.0	16.6
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	1	0	1	1	0	4.1	5.8	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	29	21	8	29	19	10	119.9	109.7	145.6
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	7	6	1	5	4	1	20.7	23.1	14.6
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	89	73	16	69	52	17	285.3	300.4	247.5
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	34	18	16	33	15	18	136.5	86.6	262.1
Arteriosclerosis (440)	1	1	0	1	1	0	4.1	5.8	-
Influenza and pneumonia (470-486)	5	1	4	5	1	4	20.7	5.8	58.2
Bronchitis, emphysema, asthma (490-493)	6	5	1	5	4	1	20.7	23.1	14.6
Peptic ulcer (531-533)	1	0	1	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	1	0	1	1	0	1	4.1	-	14.6
Cirrhosis of liver (571)	1	1	0	3	3	0	12.4	17.3	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	5	3	2	3	2	1	12.4	11.6	14.6
Infections of kidney (590)	4	3	1	3	2	1	12.4	11.6	14.6
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	2	0	2	3	1	2	12.4	5.8	29.1
Certain causes of mortality in early infancy (760-778)	6	3	3	4	2	2	8.6	7.0	11.0
Accidents (800-949)	17	13	4	20	14	6	82.7	80.9	87.4
Motor vehicle accidents (810-823)	13	9	4	13	7	6	53.8	40.4	87.4
Suicide (950-959)	1	1	0	2	1	1	8.3	5.8	14.6
Homicide (960-978)	1	0	1	1	0	1	4.1	-	14.6
Ill-defined and unknown (780-796)	16	5	11	17	6	11	70.3	34.6	160.2

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
LAMAR COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	86	59	27	297	246	51	19.5	18.6	25.2
Fetal deaths	1	1	0	6	5	1	20.2	20.3	19.6
Total deaths	103	78	25	179	146	33	11.8	11.1	16.3
Deaths under 1 year	5	2	3	10	6	4	33.7	24.4	78.4
Deaths under 28 days	1	0	1	6	4	2	20.2	16.3	39.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	11	10	1	28	24	4	184.1	182.0	197.8
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	1	0	1	7	5	2	46.0	37.9	98.9
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	38	28	10	60	47	13	394.5	356.4	642.9
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	23	16	7	29	22	7	190.7	166.8	346.2
Arteriosclerosis (440)	1	1	0	2	2	0	13.2	15.2	-
Influenza and pneumonia (470-486)	6	4	2	8	6	2	52.6	45.5	98.9
Bronchitis, emphysema, asthma (490-493)	3	3	0	5	5	0	32.9	37.9	-
Peptic ulcer (531-533)	1	1	0	3	2	1	19.7	15.2	49.4
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	2	2	0	13.2	15.2	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	1	1	0	6.6	7.6	-
Infections of kidney (590)	2	1	1	3	2	1	19.7	15.2	49.4
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	0	1	3	2	1	19.7	15.2	49.4
Certain causes of mortality in early infancy (760-778)	0	0	0	4	3	1	13.5	12.2	19.6
Accidents (800-949)	4	3	1	12	12	0	78.9	91.0	-
Motor vehicle accidents (810-823)	0	0	0	7	7	0	46.0	53.1	-
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	2	2	0	2	2	0	13.2	15.2	-
Ill-defined and unknown (780-796)	5	5	0	4	4	0	26.3	30.3	-

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
LAUDERDALE COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence						
	Number			Number			Rate*			
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	
Live Births, Fetal Deaths, deaths										
Live births	2,133	1,138	995	1,525	859	666	22.7	18.6	31.9	
Fetal deaths	116	81	35	84	60	24	55.1	69.8	36.0	
Total deaths	1,093	704	389	814	510	304	12.1	11.0	14.5	
Deaths under 1 year	67	27	40	44	16	28	28.8	18.6	42.0	
Deaths under 28 days	48	21	27	27	10	17	17.7	11.6	25.5	
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-	
Enteritis and other diarrheal diseases (008, 009)	4	3	1	1	1	0	1.5	2.2	-	
Tuberculosis (010-019)	1	1	0	1	1	0	1.5	2.2	-	
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-	
Syphilis (090-097)	1	0	1	1	0	1	1.5	-	4.8	
Malignant neoplasms (140-209)	134	92	42	119	81	38	177.4	175.4	181.8	
Benign and unspecified neoplasms (210-239)	2	2	0	2	2	0	3.0	4.3	-	
Diabetes mellitus (250)	22	12	10	11	5	6	16.4	10.8	28.7	
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-	
Anemias (280-285)	2	2	0	1	1	0	1.5	2.2	-	
Diseases of heart (390-398, 402, 404-429)	317	241	76	236	175	61	351.8	378.9	291.8	
Hypertension (400, 401, 403)	9	4	5	7	1	6	10.4	2.2	28.7	
Cerebrovascular disease (430-438)	157	108	49	112	79	33	166.9	171.0	157.9	
Arteriosclerosis (440)	20	13	7	9	4	5	13.4	8.7	23.9	
Influenza and pneumonia (470-486)	54	31	23	39	20	19	58.1	43.3	90.9	
Bronchitis, emphysema, asthma (490-493)	10	8	2	8	6	2	11.9	13.0	9.6	
Peptic ulcer (531-533)	3	2	1	2	1	1	3.0	2.2	4.8	
Hernia and intestinal obstruction (550-553, 560)	5	3	2	1	0	1	1.5	-	4.8	
Cirrhosis of liver (571)	7	5	2	3	2	1	4.5	4.3	4.8	
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	2	2	0	1	1	0	1.5	2.2	-	
Nephritis and nephrosis (580-584)	4	3	1	3	2	1	4.5	4.3	4.8	
Infections of kidney (590)	10	5	5	4	2	2	6.0	4.3	9.6	
Hyperplasia of prostate (600)	5	4	1	3	3	0	4.5	6.5	-	
Maternal causes (630-678)	1	0	1	1	0	1	6.6	-	15.0	
Congenital anomalies (740-759)	6	3	3	6	4	2	8.9	8.7	9.6	
Certain causes of mortality in early infancy (760-778)	38	18	20	21	8	13	13.8	9.3	19.5	
Accidents (800-949)	78	44	34	62	38	24	92.4	82.3	114.8	
Motor vehicle accidents (810-823)	41	25	16	27	20	7	40.2	43.3	33.5	
Suicide (950-959)	5	3	2	5	3	2	7.4	6.5	9.6	
Homicide (960-978)	10	1	9	6	0	6	8.9	-	28.7	
Ill-defined and unknown (780-796)	119	47	72	107	42	65	159.5	90.9	311.0	

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
LAWRENCE COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	94	51	43	200	114	86	18.0	15.1	24.0
Fetal deaths	5	2	3	10	4	6	50.0	35.1	69.8
Total deaths	87	51	36	128	81	47	11.5	10.7	13.1
Deaths under 1 year	2	1	1	3	2	1	15.0	17.5	11.6
Deaths under 28 days	1	0	1	1	1	0	5.0	8.8	-
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	13	8	5	21	13	8	188.6	172.1	223.2
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	2	0	2	6	3	3	53.9	39.7	83.7
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	33	25	8	47	37	10	422.0	489.9	279.0
Hypertension (400, 401, 403)	1	1	0	1	1	0	9.0	13.2	-
Cerebrovascular disease (430-438)	13	4	9	18	8	10	161.6	105.9	279.0
Arteriosclerosis (440)	4	4	0	5	5	0	44.9	66.2	-
Influenza and pneumonia (470-486)	2	1	1	4	2	2	35.9	26.5	55.8
Bronchitis, emphysema, asthma (490-493)	1	1	0	1	1	0	9.0	13.2	-
Peptic ulcer (531-533)	0	0	0	1	1	0	9.0	13.2	-
Hernia and intestinal obstruction (550-553, 560)	1	0	1	2	1	1	18.0	13.2	27.9
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	0	1	1	0	1	9.0	-	27.9
Infections of kidney (590)	1	1	0	1	1	0	9.0	13.2	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	1	1	0	9.0	13.2	-
Certain causes of mortality in early infancy (760-778)	1	0	1	0	0	0	-	-	-
Accidents (800-949)	8	4	4	9	5	4	80.8	66.2	111.6
Motor vehicle accidents (810-823)	6	3	3	7	4	3	62.8	53.0	83.7
Suicide (950-959)	1	1	0	1	1	0	9.0	13.2	-
Homicide (960-978)	0	0	0	1	0	1	9.0	-	27.9
Ill-defined and unknown (780-796)	0	0	0	3	1	2	26.9	13.2	55.8

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
LEAKE COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non- white	Total	White	Non- white	Total	White	Non- white
Live births	185	112	73	304	154	150	17.8	14.7	22.6
Fetal deaths	6	1	5	12	2	10	39.5	13.0	66.7
Total deaths	157	109	48	220	153	67	12.9	14.6	10.1
Deaths under 1 year	6	4	2	8	5	3	26.3	32.5	20.0
Deaths under 28 days	4	4	0	5	5	0	16.4	32.5	-
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	1	0	1	1	0	5.8	9.6	-
Tuberculosis (010-019)	0	0	0	1	0	1	5.8	-	15.1
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	25	17	8	39	25	14	228.3	239.2	211.1
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	8	7	1	9	7	2	52.7	67.0	30.2
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	66	47	19	84	61	23	491.6	583.6	346.8
Hypertension (400, 401, 403)	1	1	0	1	1	0	5.8	9.6	-
Cerebrovascular disease (430-438)	19	13	6	24	15	9	140.5	143.5	135.7
Arteriosclerosis (440)	1	1	0	2	2	0	11.7	19.1	-
Influenza and pneumonia (470-486)	2	0	2	5	3	2	29.3	28.7	30.2
Bronchitis, emphysema, asthma (490-493)	1	1	0	1	1	0	5.8	9.6	-
Peptic ulcer (531-533)	0	0	0	1	0	1	5.8	-	15.1
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	0	1	5.8	-	15.1
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	1	1	0	5.8	9.6	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	3	2	1	17.6	19.1	15.1
Certain causes of mortality in early infancy (760-778)	3	3	0	3	3	0	9.9	19.5	-
Accidents (800-949)	13	5	8	22	15	7	128.8	143.5	105.5
Motor vehicle accidents (810-823)	11	3	8	17	11	6	99.5	105.2	90.4
Suicide (950-959)	1	1	0	2	2	0	11.7	19.1	-
Homicide (960-978)	4	2	2	3	1	2	17.6	9.6	30.2
Ill-defined and unknown (780-796)	4	2	2	3	1	2	17.6	9.6	30.2

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
LEE COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,355	1,088	267	897	658	239	19.4	18.0	25.0
Fetal deaths	29	21	8	23	16	7	25.6	24.3	29.3
Total deaths	581	428	153	454	330	124	9.8	9.0	12.9
Deaths under 1 year	42	28	14	29	17	12	32.3	25.8	50.2
Deaths under 28 days	34	25	9	20	13	7	22.3	19.8	29.3
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	2	0	2	2	0	2	4.3	-	20.9
Tuberculosis (010-019)	0	0	0	1	0	1	2.2	-	10.4
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	89	62	27	81	55	26	175.5	150.4	271.5
Benign and unspecified neoplasms (210-239)	2	1	1	1	0	1	2.2	-	10.4
Diabetes mellitus (250)	5	4	1	5	4	1	10.8	10.9	10.4
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	3	1	2	2	0	2	4.3	-	20.9
Diseases of heart (390-398, 402, 404-429)	208	163	45	179	143	36	387.9	391.0	375.9
Hypertension (400, 401, 403)	0	0	0	1	0	1	2.2	-	10.4
Cerebrovascular disease (430-438)	71	59	12	48	39	9	104.0	106.6	94.0
Arteriosclerosis (440)	2	2	0	2	2	0	4.3	5.5	-
Influenza and pneumonia (470-486)	18	15	3	14	11	3	30.3	30.1	31.3
Bronchitis, emphysema, asthma (490-493)	8	8	0	7	7	0	15.2	19.1	-
Peptic ulcer (531-533)	3	3	0	1	1	0	2.2	2.7	-
Hernia and intestinal obstruction (550-553, 560)	1	0	1	1	0	1	2.2	-	10.4
Cirrhosis of liver (571)	3	3	0	5	5	0	10.8	13.7	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	3	3	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	5	1	4	2	1	1	4.3	2.7	10.4
Infections of kidney (590)	9	6	3	8	5	3	17.3	13.7	31.3
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	7	5	2	5	3	2	10.8	8.2	20.9
Certain causes of mortality in early infancy (760-778)	27	19	8	16	10	6	17.8	15.2	25.1
Accidents (800-949)	57	36	21	21	10	11	45.5	27.3	114.8
Motor vehicle accidents (810-823)	34	27	7	7	6	1	15.2	16.4	10.4
Suicide (950-959)	4	4	0	4	4	0	8.7	10.9	-
Homicide (960-978)	7	3	4	7	2	5	15.2	5.5	52.2
Ill-defined and unknown (780-796)	10	6	4	11	7	4	23.8	19.1	41.8

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
LEFLORE COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,107	398	709	1,002	303	699	23.8	17.3	28.4
Fetal deaths	81	31	50	72	27	45	71.8	89.1	64.4
Total deaths	567	228	339	513	180	333	12.2	10.2	13.6
Deaths under 1 year	47	8	39	40	6	34	39.9	19.8	48.6
Deaths under 28 days	31	8	23	28	6	22	27.9	19.8	31.5
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	4	0	4	3	1	2	7.1	5.7	8.1
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	72	36	36	62	27	35	147.2	153.8	142.5
Benign and unspecified neoplasms (210-239)	1	0	1	1	0	1	2.4	-	4.1
Diabetes mellitus (250)	8	1	7	8	1	7	19.0	5.7	28.5
Avitaminoses and other nutritional deficiencies (260-269)	1	0	1	0	0	0	-	-	-
Anemias (280-285)	2	1	1	4	2	2	9.5	11.4	8.1
Diseases of heart (390-398, 402, 404-429)	180	96	84	158	74	84	375.2	421.6	342.0
Hypertension (400, 401, 403)	6	2	4	4	0	4	9.5	-	16.3
Cerebrovascular (430-438)	78	28	50	75	26	49	178.1	148.1	199.5
Arteriosclerosis (440)	2	2	0	3	1	2	7.1	5.7	8.1
Influenza and pneumonia (470-486)	19	9	10	17	9	8	40.4	51.3	32.6
Bronchitis, emphysema, asthma (490-493)	6	1	5	5	0	5	11.9	-	20.4
Peptic ulcer (531-533)	2	0	2	3	1	2	7.1	5.7	8.1
Hernia and intestinal obstruction (550-553, 560)	1	0	1	2	0	2	4.7	-	8.1
Cirrhosis of liver (571)	2	2	0	1	1	0	2.4	5.7	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	1	1	0	2.4	5.7	-
Nephritis and nephrosis (580-584)	4	1	3	4	0	4	9.5	-	16.3
Infections of kidney (590)	2	2	0	1	1	0	2.4	5.7	-
Hyperplasia of prostate (600)	2	1	1	2	1	1	4.7	5.7	4.1
Maternal causes (630-678)	1	0	1	1	0	1	10.0	-	14.3
Congenital anomalies (740-759)	8	3	5	7	2	5	16.6	11.4	20.4
Certain causes of mortality in early infancy (760-778)	19	4	15	20	4	16	20.0	13.2	22.9
Accidents (800-949)	40	13	27	31	7	24	73.6	39.9	97.7
Motor vehicle accidents (810-823)	19	10	9	12	5	7	28.5	28.5	28.5
Suicide (950-959)	4	4	0	3	3	0	7.1	17.1	-
Homicide (960-978)	6	0	6	6	0	6	14.2	-	24.4
Ill-defined and unknown (780-796)	62	7	55	61	7	54	144.8	39.9	219.9

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
LINCOLN COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non- white	Total	White	Non- white	Total	White	Non- white
Live births	579	354	225	532	327	205	20.3	18.0	25.4
Fetal deaths	7	2	5	17	5	12	32.0	15.3	58.5
Total deaths	286	192	94	296	187	109	11.3	10.3	13.5
Deaths under 1 year	10	8	2	12	6	6	22.6	18.3	29.3
Deaths under 28 days	10	8	2	12	6	6	22.6	18.3	29.3
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	39	26	13	45	29	16	171.8	159.9	198.5
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	7	3	4	7	1	6	26.7	5.5	74.4
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	1	0	1	3.8	-	12.4
Anemias (280-285)	0	0	0	1	1	0	3.8	5.5	-
Diseases of heart (390-398, 402, 404-429)	117	82	35	109	76	33	416.1	419.0	409.4
Hypertension (400, 401, 403)	0	0	0	1	1	0	3.8	5.5	-
Cerebrovascular disease (430-438)	44	29	15	42	27	15	160.3	148.8	186.1
Arteriosclerosis (440)	10	9	1	6	5	1	22.9	27.6	12.4
Influenza and pneumonia (470-486)	6	6	0	6	4	2	22.9	22.0	24.8
Bronchitis, emphysema, asthma (490-493)	0	0	0	0	0	0	-	-	-
Peptic ulcer (531-533)	2	1	1	2	1	1	7.6	5.5	12.4
Hernia and intestinal obstruction (550-553, 560)	2	2	0	1	1	0	3.8	5.5	-
Cirrhosis of liver (571)	0	0	0	1	0	1	3.8	-	12.4
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	1	1	0	3.8	5.5	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	2	2	0	2	2	0	7.6	11.0	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	3	2	1	11.4	11.0	12.4
Certain causes of mortality in early infancy (760-778)	7	5	2	7	4	3	13.2	12.2	14.6
Accidents (800-949)	14	5	9	20	8	12	76.3	44.1	148.9
Motor vehicle accidents (810-823)	8	5	3	11	6	5	42.0	33.1	62.0
Suicide (950-959)	3	3	0	3	3	0	11.4	16.5	-
Homicide (960-978)	1	0	1	1	0	1	3.8	-	12.4
Ill-defined and unknown (780-796)	19	8	11	20	8	12	76.3	44.1	148.9

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
LOWNDES COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,165	708	457	1,008	502	506	20.3	15.0	30.9
Fetal deaths	42	18	24	38	12	26	37.7	23.9	51.4
Total deaths	473	271	202	484	252	232	9.7	7.6	14.2
Deaths under 1 year	28	8	20	31	6	25	30.8	12.0	49.4
Deaths under 28 days	18	7	11	22	6	16	21.8	12.0	31.6
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	2	1	1	1	0	1	2.0	-	6.1
Tuberculosis (010-019)	1	0	1	3	0	3	6.0	-	18.3
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	71	54	17	71	50	21	142.8	150.0	128.3
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	9	6	3	8	5	3	16.1	15.0	18.3
Avitaminoses and other nutritional deficiencies (260-269)	2	0	2	2	0	2	4.0	-	12.2
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	127	82	45	123	75	48	247.5	225.0	293.3
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	49	27	22	64	33	31	128.8	99.0	189.4
Arteriosclerosis (440)	10	10	0	9	9	0	18.1	27.0	-
Influenza and pneumonia (470-486)	25	8	17	27	9	18	54.3	27.0	110.0
Bronchitis, emphysema, asthma (490-493)	2	2	0	4	4	0	8.0	12.0	-
Peptic ulcer (531-533)	1	1	0	1	1	0	2.0	3.0	-
Hernia and intestinal obstruction (550-553, 560)	2	1	1	0	0	0	-	-	-
Cirrhosis of liver (571)	4	3	1	4	3	1	8.0	9.0	6.1
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	1	1	0	2.0	3.0	-
Nephritis and nephrosis (580-584)	2	2	0	1	1	0	2.0	3.0	-
Infections of kidney (590)	1	0	1	2	1	1	4.0	3.0	6.1
Hyperplasia of prostate (600)	1	1	0	1	1	0	2.0	3.0	-
Maternal causes (630-678)	1	0	1	1	0	1	9.9	-	19.8
Congenital anomalies (740-759)	6	1	5	8	2	6	16.1	6.0	36.7
Certain causes of mortality in early infancy (760-778)	13	6	7	15	4	11	14.9	8.0	21.7
Accidents (800-949)	44	29	15	37	21	16	74.4	63.0	97.8
Motor vehicle accidents (810-823)	23	18	5	17	11	6	34.2	33.0	36.7
Suicide (950-959)	3	2	1	2	1	1	4.0	3.0	6.1
Homicide (960-978)	9	1	8	9	1	8	18.1	3.0	48.9
Ill-defined and unknown (780-796)	54	7	47	54	7	47	108.6	21.0	287.2

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
MADISON COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	491	106	385	768	169	599	25.8	15.2	32.2
Fetal deaths	27	8	19	49	21	28	63.8	124.3	46.7
Total deaths	315	135	180	346	134	212	11.6	12.0	11.4
Deaths under 1 year	19	2	17	24	2	22	31.2	11.8	36.7
Deaths under 28 days	11	2	9	12	2	10	15.6	11.8	16.7
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	2	0	2	6.7	-	10.8
Tuberculosis (010-019)	0	0	0	1	1	0	3.4	9.0	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	38	19	19	50	24	26	168.1	215.3	139.9
Benign and unspecified neoplasms (210-239)	0	0	0	2	1	1	6.7	9.0	5.4
Diabetes mellitus (250)	13	3	10	12	2	10	40.4	17.9	53.8
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	1	0	1	1	0	1	3.4	-	5.4
Diseases of heart (390-398, 402, 404-429)	84	42	42	84	38	46	282.5	340.9	247.4
Hypertension (400, 401, 403)	2	1	1	3	1	2	10.1	9.0	10.8
Cerebrovascular disease (430-438)	31	14	17	33	11	22	111.0	98.7	118.3
Arteriosclerosis (440)	3	1	2	3	1	2	10.1	9.0	10.8
Influenza and pneumonia (470-486)	11	4	7	12	4	8	40.4	35.9	43.0
Bronchitis, emphysema, asthma (490-493)	5	5	0	7	7	0	23.5	62.8	-
Peptic ulcer (531-533)	1	0	1	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	1	0	1	1	0	1	3.4	-	5.4
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	1	0	3.4	9.0	-
Nephritis and nephrosis (580-584)	6	1	5	6	1	5	20.2	9.0	26.9
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	6	2	4	7	2	5	9.1	11.8	8.3
Accidents (800-949)	32	17	15	37	16	21	124.4	143.5	113.0
Motor vehicle accidents (810-823)	12	11	1	16	7	9	53.8	62.8	48.4
Suicide (950-959)	4	3	1	4	3	1	13.4	26.9	5.4
Homicide (960-978)	3	0	3	3	0	3	10.1	-	16.1
Ill-defined and unknown (780-796)	57	12	45	55	9	46	185.0	80.7	247.4

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
MARION COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	377	253	124	485	283	202	21.2	18.0	28.3
Fetal deaths	44	33	11	45	32	13	92.8	113.1	64.4
Total deaths	210	133	77	249	152	97	10.9	9.6	13.6
Deaths under 1 year	10	4	6	12	3	9	24.7	10.6	44.6
Deaths under 28 days	7	4	3	8	3	5	16.5	10.6	24.8
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	1	0	1	4.4	-	14.0
Tuberculosis (010-019)	1	1	0	3	2	1	13.1	12.7	14.0
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	29	17	12	37	21	16	161.8	133.4	224.3
Benign and unspecified neoplasms (210-239)	1	1	0	2	2	0	8.7	12.7	-
Diabetes mellitus (250)	3	0	3	3	0	3	13.1	-	42.1
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	45	35	10	59	44	15	258.0	279.6	210.3
Hypertension (400, 401, 403)	2	1	1	3	1	2	13.1	6.4	28.0
Cerebrovascular disease (430-438)	40	29	11	36	26	10	157.4	165.2	140.2
Arteriosclerosis (440)	3	2	1	2	1	1	8.7	6.4	14.0
Influenza and pneumonia (470-486)	6	6	0	4	4	0	17.5	25.4	-
Bronchitis, emphysema, asthma (490-493)	3	2	1	4	3	1	17.5	19.1	14.0
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	1	1	0	1	1	0	4.4	6.4	-
Cirrhosis of liver (571)	0	0	0	1	0	1	4.4	-	14.0
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	1	1	0	4.4	6.4	-
Nephritis and nephrosis (580-584)	1	1	0	4	4	0	17.5	25.4	-
Infections of kidney (590)	1	1	0	1	1	0	4.4	6.4	-
Hyperplasia of prostate (600)	1	0	1	1	0	1	4.4	-	14.0
Maternal causes (630-678)	0	0	0	1	0	1	20.6	-	49.5
Congenital anomalies (740-759)	2	1	1	4	2	2	17.5	12.7	28.0
Certain causes of mortality in early infancy (760-778)	5	4	1	5	3	2	10.3	10.6	9.9
Accidents (800-949)	16	11	5	20	13	7	87.4	82.6	98.1
Motor vehicle accidents (810-823)	12	9	3	10	7	3	43.7	44.5	42.1
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	2	0	2	2	0	2	8.7	-	28.0
Ill-defined and unknown (780-796)	36	8	28	35	7	28	153.0	44.5	392.6

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
MARSHALL COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	292	22	270	555	142	413	23.1	15.6	27.7
Fetal deaths	23	4	19	31	9	22	55.8	63.4	53.3
Total deaths	148	49	99	245	96	149	10.2	10.5	10.0
Deaths under 1 year	14	0	14	23	0	23	41.4	-	55.7
Deaths under 28 days	9	0	9	16	0	16	28.8	-	38.7
Bacillary dysentery and amebiasis (004, 006)	0	0	0	1	0	1	4.2	-	6.7
Enteritis and other diarrheal diseases (008, 009)	2	0	2	4	1	3	16.6	11.0	20.1
Tuberculosis (010-019)	0	0	0	1	0	1	4.2	-	6.7
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	1	0	1	1	0	1	4.2	-	6.7
Malignant neoplasms (140-209)	15	6	9	36	13	23	149.8	142.8	154.1
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	1	0	1	4	1	3	16.6	11.0	20.1
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	2	0	2	8.3	-	13.4
Diseases of heart (390-398, 402, 404-429)	39	17	22	60	33	27	249.7	362.6	180.9
Hypertension (400, 401, 403)	3	1	2	3	1	2	12.5	11.0	13.4
Cerebrovascular disease (430-438)	31	9	22	44	16	28	183.1	175.8	187.6
Arteriosclerosis (440)	2	1	1	4	3	1	16.6	33.0	6.7
Influenza and pneumonia (470-486)	3	0	3	6	3	3	25.0	33.0	20.1
Bronchitis, emphysema, asthma (490-493)	0	0	0	2	2	0	8.3	22.0	-
Peptic ulcer (531-533)	0	0	0	1	0	1	4.2	-	6.7
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	3	3	0	2	2	0	8.3	22.0	-
Infections of kidney (590)	0	0	0	1	1	0	4.2	11.0	-
Hyperplasia of prostate (600)	1	0	1	1	0	1	4.2	-	6.7
Maternal causes (630-678)	1	1	0	1	1	0	18.0	70.4	-
Congenital anomalies (740-759)	1	0	1	3	0	3	12.5	-	20.1
Certain causes of mortality in early infancy (760-778)	7	0	7	10	0	10	18.0	-	24.2
Accidents (800-949)	16	5	11	23	10	13	95.7	109.9	87.1
Motor vehicle accidents (810-823)	7	4	3	11	8	3	45.8	87.9	20.1
Suicide (950-959)	1	1	0	1	1	0	4.2	11.0	-
Homicide (960-978)	0	0	0	5	0	5	20.8	-	33.5
Ill-defined and unknown (780-796)	16	5	11	18	5	13	74.9	54.9	87.1

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
MONROE COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	932	626	306	725	444	281	21.3	18.8	27.0
Fetal deaths	11	7	4	9	5	4	12.4	11.3	14.2
Total deaths	381	243	138	413	267	146	12.1	11.3	14.0
Deaths under 1 year	23	13	10	18	11	7	24.8	24.8	24.9
Deaths under 28 days	18	11	7	14	10	4	19.3	22.5	14.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	1	0	1	2.9	-	9.6
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	37	25	12	41	27	14	120.4	114.2	134.7
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	11	5	6	11	5	6	32.3	21.1	57.7
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	138	98	40	159	113	46	467.0	477.8	442.6
Hypertension (400, 401, 403)	2	0	2	2	0	2	5.9	-	19.2
Cerebrovascular disease (430-438)	63	33	30	64	37	27	188.0	156.4	259.8
Arteriosclerosis (440)	12	9	3	13	10	3	38.2	42.3	28.9
Influenza and pneumonia (470-486)	10	8	2	8	5	3	23.5	21.1	28.9
Bronchitis, emphysema, asthma (490-493)	5	5	0	4	4	0	11.7	16.9	-
Peptic ulcer (531-533)	3	1	2	3	1	2	8.8	4.2	19.2
Hernia and intestinal obstruction (550-553, 560)	1	1	0	1	1	0	2.9	4.2	-
Cirrhosis of liver (571)	0	0	0	1	0	1	2.9	-	9.6
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	5	1	4	6	1	5	17.6	4.2	48.1
Infections of kidney (590)	4	4	0	6	5	1	17.6	21.1	9.6
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	3	1	2	2	1	1	5.9	4.2	9.6
Certain causes of mortality in early infancy (760-778)	14	9	5	12	9	3	16.6	20.3	10.7
Accidents (800-949)	34	17	17	37	21	16	108.7	88.8	153.9
Motor vehicle accidents (810-823)	19	11	8	19	15	4	55.8	63.4	38.5
Suicide (950-959)	3	3	0	3	3	0	8.8	12.7	-
Homicide (960-978)	1	1	0	1	1	0	2.9	4.2	-
Ill-defined and unknown (780-796)	9	4	5	11	5	6	32.3	21.1	57.7

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
MONTGOMERY COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	289	127	162	230	96	134	17.8	13.5	23.1
Fetal deaths	12	2	10	11	3	8	47.8	31.2	59.7
Total deaths	183	111	72	175	103	72	13.5	14.5	12.4
Deaths under 1 year	12	3	9	10	2	8	43.5	20.8	59.7
Deaths under 28 days	9	3	6	9	2	7	39.1	20.8	52.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	15	12	3	18	14	4	139.3	196.6	69.0
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	4	1	3	2	0	2	15.5	-	34.5
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	1	0	1	1	0	1	7.7	-	17.2
Diseases of heart (390-398, 402, 404-429)	90	55	35	76	43	33	588.3	603.8	569.2
Hypertension (400, 401, 403)	2	1	1	1	0	1	7.7	-	17.2
Cerebrovascular disease (430-438)	36	24	12	36	24	12	278.7	337.0	207.0
Arteriosclerosis (440)	1	1	0	2	2	0	15.5	28.1	-
Influenza and pneumonia (470-486)	2	1	1	0	0	0	-	-	-
Bronchitis, emphysema, asthma (490-493)	1	1	0	2	2	0	15.5	28.1	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	2	2	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	3	1	2	3	1	2	23.2	14.0	34.5
Infections of kidney (590)	1	1	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	1	1	0	7.7	14.0	-
Certain causes of mortality in early infancy (760-778)	7	1	6	7	1	6	30.4	10.4	44.8
Accidents (800-949)	4	2	2	9	5	4	69.7	70.2	69.0
Motor vehicle accidents (810-823)	1	1	0	3	2	1	23.2	28.1	17.2
Suicide (950-959)	1	1	0	1	1	0	7.7	14.0	-
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	5	1	4	3	0	3	23.2	-	51.8

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
NESHOBA COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	397	238	159	396	229	167	19.0	15.2	29.2
Fetal deaths	11	5	6	9	4	5	22.7	17.5	29.9
Total deaths	188	143	45	229	173	56	11.0	11.5	9.8
Deaths under 1 year	9	5	4	17	10	7	42.9	43.7	41.9
Deaths under 28 days	7	5	2	14	9	5	35.4	39.3	29.9
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	1	0	1	4.8	-	17.5
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	25	17	8	27	18	9	129.8	119.3	157.6
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	3	2	1	5	4	1	24.0	26.5	17.5
Avitaminoses and other nutritional deficiencies (260-269)	1	0	1	1	0	1	4.8	-	17.5
Anemias (280-285)	1	1	0	1	0	1	4.8	-	17.5
Diseases of heart (390-398, 402, 404-429)	63	55	8	70	61	9	336.5	404.2	157.6
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	27	22	5	28	23	5	134.6	152.4	87.6
Arteriosclerosis (440)	3	3	0	6	6	0	28.8	39.8	-
Influenza and pneumonia (470-486)	1	1	0	5	2	3	24.0	13.2	52.5
Bronchitis, emphysema, asthma (490-493)	4	4	0	4	4	0	19.2	26.5	-
Peptic ulcer (531-533)	1	1	0	1	1	0	4.8	6.6	-
Hernia and intestinal obstruction (550-553, 560)	1	1	0	1	1	0	4.8	6.6	-
Cirrhosis of liver (571)	1	1	0	3	2	1	14.4	13.2	17.5
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	1	1	0	4.8	6.6	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	1	0	1	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	1	1	0	25.2	43.7	-
Congenital anomalies (740-759)	1	1	0	2	1	1	9.6	6.6	17.5
Certain causes of mortality in early infancy (760-778)	4	4	0	9	8	1	22.7	34.9	6.0
Accidents (800-949)	18	15	3	25	19	6	120.2	125.9	105.1
Motor vehicle accidents (810-823)	2	1	1	5	3	2	24.0	19.9	35.0
Suicide (950-959)	2	1	1	2	1	1	9.6	6.6	17.5
Homicide (960-978)	2	0	2	2	1	1	9.6	6.6	17.5
Ill-defined and unknown (780-796)	15	5	10	19	9	10	91.3	59.6	175.1

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
NEWTON COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	320	201	119	363	201	162	19.1	15.0	28.8
Fetal deaths	18	8	10	16	5	11	44.1	24.9	67.9
Total deaths	228	160	68	268	188	80	14.1	14.1	14.2
Deaths under 1 year	9	3	6	12	3	9	33.0	14.9	55.6
Deaths under 28 days	6	3	3	9	2	7	24.8	10.0	43.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	1	1	0	5.3	7.5	-
Tuberculosis (010-019)	1	0	1	1	0	1	5.3	-	17.8
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	1	0	1	1	0	1	5.3	-	17.8
Malignant neoplasms (140-209)	24	21	3	26	23	3	137.0	172.2	53.3
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	8	3	5	9	4	5	47.4	30.0	88.8
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	1	1	0	5.3	7.5	-
Diseases of heart (390-398, 402, 404-429)	107	84	23	106	80	26	558.4	599.2	461.7
Hypertension (400, 401, 403)	6	4	2	7	5	2	36.9	37.4	35.5
Cerebrovascular disease (430-438)	32	18	14	36	21	15	189.6	157.3	266.4
Arteriosclerosis (440)	6	4	2	7	5	2	36.9	37.4	35.5
Influenza and pneumonia (470-486)	4	2	2	10	8	2	52.7	59.9	35.5
Bronchitis, emphysema, asthma (490-493)	1	1	0	1	1	0	5.3	7.5	-
Peptic ulcer (531-533)	0	0	0	1	0	1	5.3	-	17.8
Hernia and intestinal obstruction (550-553, 560)	1	1	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	1	1	0	5.3	7.5	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	2	2	0	10.5	15.0	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	1	0	1	5.3	-	17.8
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	1	1	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	6	3	3	8	2	6	22.0	10.0	37.0
Accidents (800-949)	12	8	4	17	13	4	89.6	97.4	71.0
Motor vehicle accidents (810-823)	7	5	2	10	8	2	52.7	59.9	35.5
Suicide (950-959)	3	3	0	3	3	0	15.8	22.5	-
Homicide (960-978)	0	0	0	3	1	2	15.8	7.5	35.5
Ill-defined and unknown (780-796)	6	3	3	10	7	3	52.7	52.4	53.3

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
NOXUBEE COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	377	73	304	345	68	277	24.1	14.0	29.3
Fetal Deaths	12	1	11	14	4	10	40.6	58.8	36.1
Total deaths	129	43	86	159	57	102	11.1	11.8	10.8
Deaths under 1 year	14	1	13	12	1	11	34.8	14.7	39.7
Deaths under 28 days	8	0	8	6	0	6	17.4	-	21.7
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	3	0	3	3	0	3	21.0	-	31.8
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	1	0	1	1	0	1	7.0	-	10.6
Malignant neoplasms (140-209)	14	7	7	17	7	10	119.0	144.5	105.9
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	2	1	1	1	0	1	7.0	-	10.6
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	29	15	14	41	24	17	287.0	495.4	180.0
Hypertension (400, 401, 403)	1	0	1	1	0	1	7.0	-	10.6
Cerebrovascular disease (430-438)	21	6	15	24	7	17	168.0	144.5	180.0
Arteriosclerosis (440)	2	0	2	2	0	2	14.0	-	21.2
Influenza and pneumonia (470-486)	3	2	1	4	3	1	28.0	61.9	10.6
Bronchitis, emphysema, asthma (490-493)	0	0	0	0	0	0	-	-	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	1	0	7.0	20.6	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	2	1	1	14.0	20.6	10.6
Infections of kidney (590)	2	1	1	3	1	2	21.0	20.6	21.2
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	5	0	5	3	0	3	8.7	-	10.8
Accidents (800-949)	12	5	7	13	5	8	91.0	103.2	84.7
Motor vehicle accidents (810-823)	6	2	4	6	2	4	42.0	41.3	42.4
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	2	0	2	3	0	3	21.0	-	31.8
Ill-defined and unknown (780-796)	25	3	22	28	4	24	196.0	82.6	254.1

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
OKTIBBEHA COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	649	314	335	587	318	269	20.4	17.1	26.4
Fetal deaths	11	2	9	12	2	10	20.4	6.3	37.2
Total deaths	207	100	107	230	118	112	8.0	6.4	11.0
Deaths under 1 year	9	1	8	11	3	8	18.7	9.4	29.7
Deaths under 28 days	6	1	5	5	2	3	8.5	6.3	11.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	2	0	2	3	0	3	10.4	-	29.5
Tuberculosis (010-019)	1	0	1	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	19	9	10	27	13	14	93.9	70.0	137.5
Benign and unspecified neoplasms (210-239)	2	1	1	2	1	1	7.0	5.4	9.8
Diabetes mellitus (250)	3	2	1	4	2	2	13.9	10.8	19.6
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	2	1	1	2	1	1	7.0	5.4	9.8
Diseases of heart (390-398, 402, 404-429)	81	40	41	94	55	39	326.9	296.2	383.0
Hypertension (400, 401, 403)	6	1	5	4	1	3	13.9	5.4	29.5
Cerebrovascular disease (430-438)	31	12	19	27	9	18	93.9	48.5	176.8
Arteriosclerosis (440)	4	3	1	4	3	1	13.9	16.2	9.8
Influenza and pneumonia (470-486)	1	0	1	3	2	1	10.4	10.8	9.8
Bronchitis, emphysema, asthma (490-493)	4	2	2	4	2	2	13.9	10.8	19.6
Peptic ulcer (531-533)	2	2	0	2	2	0	7.0	10.8	-
Hernia and intestinal obstruction (550-553, 560)	1	0	1	1	0	1	3.5	-	9.8
Cirrhosis of liver (571)	1	1	0	2	1	1	7.0	5.4	9.8
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	2	1	1	1	0	1	3.5	-	9.8
Infections of kidney (590)	1	1	0	2	2	0	7.0	10.8	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	1	0	1	1	0	1	17.0	-	37.2
Congenital anomalies (740-759)	2	1	1	4	1	3	13.9	5.4	29.5
Certain causes of mortality in early infancy (760-778)	5	1	4	4	2	2	6.8	6.3	7.4
Accidents (800-949)	21	13	8	22	12	10	76.5	64.6	98.2
Motor vehicle accidents (810-823)	14	12	2	11	8	3	38.2	43.1	29.5
Suicide (950-959)	3	3	0	4	4	0	13.9	21.5	-
Homicide (960-978)	2	2	0	2	1	1	7.0	5.4	9.8
Ill-defined and unknown (780-796)	4	0	4	3	0	3	10.4	-	29.5

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
PANOLA COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	692	217	475	585	179	406	21.8	13.7	29.5
Fetal deaths	25	7	18	21	6	15	35.9	33.5	36.9
Total deaths	246	129	117	336	171	165	12.5	13.1	12.0
Deaths under 1 year	14	4	10	19	5	14	32.5	27.9	34.5
Deaths under 28 days	8	3	5	10	4	6	17.1	22.3	14.8
Bacillary dysentery and amebiasis (004, 006)	1	0	1	1	0	1	3.7	-	7.3
Enteritis and other diarrheal diseases (008, 009)	3	1	2	3	1	2	11.2	7.6	14.5
Tuberculosis (010-019)	0	0	0	1	0	1	3.7	-	7.3
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	21	10	11	41	23	18	152.8	176.1	130.7
Benign and unspecified neoplasms (210-239)	1	0	1	1	0	1	3.7	-	7.3
Diabetes mellitus (250)	2	1	1	7	6	1	26.1	45.9	7.3
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	110	68	42	122	71	51	454.7	543.6	370.4
Hypertension (400, 401, 403)	1	0	1	1	0	1	3.7	-	7.3
Cerebrovascular disease (430-438)	33	12	21	53	17	36	197.5	130.2	261.5
Arteriosclerosis (440)	3	3	0	4	4	0	14.9	30.6	-
Influenza and pneumonia (470-486)	7	1	6	14	4	10	52.2	30.6	72.6
Bronchitis, emphysema, asthma (490-493)	1	1	0	4	4	0	14.9	30.6	-
Peptic ulcer (531-533)	2	2	0	2	2	0	7.4	15.3	-
Hernia and intestinal obstruction (550-553, 560)	1	0	1	3	0	3	11.2	-	21.8
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	1	0	1	3.7	-	7.3
Infections of kidney (590)	1	0	1	2	0	2	7.4	-	14.5
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	2	1	1	1	1	0	3.7	7.6	-
Certain causes of mortality in early infancy (760-778)	6	2	4	9	3	6	15.4	16.8	14.8
Accidents (800-949)	23	15	8	22	12	10	82.0	91.9	72.6
Motor vehicle accidents (810-823)	10	8	2	11	8	3	41.0	61.2	21.8
Suicide (950-959)	5	4	1	6	4	2	22.4	30.6	14.5
Homicide (960-978)	3	0	3	2	0	2	7.4	-	14.5
Ill-defined and unknown (780-796)	9	2	7	12	3	9	44.7	23.0	65.4

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
PEARL RIVER COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non- white	Total	White	Non- white	Total	White	Non- white
Live births	592	509	83	556	439	117	20.0	19.3	22.9
Fetal deaths	22	18	4	13	9	4	23.4	20.5	34.2
Total deaths	233	184	49	291	227	64	10.5	10.0	12.5
Deaths under 1 year	18	12	6	21	12	9	37.8	27.3	76.9
Deaths under 28 days	16	12	4	19	12	7	34.2	27.3	59.8
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	1	0	0	0	0	-	-	-
Tuberculosis (010-019)	1	0	1	2	1	1	7.2	4.4	19.6
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	34	32	2	43	39	4	154.7	171.9	78.2
Benign and unspecified neoplasms (210-239)	0	0	0	1	1	0	3.6	4.4	-
Diabetes mellitus (250)	4	2	2	5	2	3	18.0	8.8	58.7
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	92	74	18	98	78	20	352.5	343.8	391.1
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	19	14	5	29	22	7	104.3	97.0	136.9
Arteriosclerosis (440)	6	4	2	4	2	2	14.4	8.8	39.1
Influenza and pneumonia (470-486)	9	5	4	10	5	5	36.0	22.0	97.8
Bronchitis, emphysema, asthma (490-493)	2	2	0	4	4	0	14.4	17.6	-
Peptic ulcer (531-533)	1	1	0	2	2	0	7.2	8.8	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	1	0	3.6	4.4	-
Cirrhosis of liver (571)	0	0	0	1	1	0	3.6	4.4	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	1	0	3	3	0	10.8	13.2	-
Nephritis and nephrosis (580-584)	0	0	0	1	1	0	3.6	4.4	-
Infections of kidney (590)	0	0	0	1	1	0	3.6	4.4	-
Hyperplasia of prostate (600)	1	1	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	3	3	0	4	4	0	14.4	17.6	-
Certain causes of mortality in early infancy (760-778)	11	8	3	15	9	6	27.0	20.5	51.3
Accidents (800-949)	25	21	4	32	26	6	115.1	114.6	117.3
Motor vehicle accidents (810-823)	21	17	4	21	18	3	75.5	79.3	58.7
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	1	0	1	4	2	2	14.4	8.8	39.1
Ill-defined and unknown (780-796)	10	4	6	10	4	6	36.0	17.6	117.3

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
PERRY COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence						
	Number			Number			Rate*			
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	
Live Births, Fetal Deaths, Deaths										
Live births	122	86	36	192	130	62	21.2	19.5	25.9	
Fetal deaths	6	4	2	8	5	3	41.7	38.5	48.4	
Total deaths	67	53	14	86	73	13	9.5	10.9	5.4	
Deaths under 1 year	4	4	0	8	8	0	41.7	61.5	-	
Deaths under 28 days	3	3	0	5	5	0	26.0	38.5	-	
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-	
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-	
Tuberculosis (010-019)	1	1	0	1	1	0	11.0	15.0	-	
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-	
Syphilis (090-097)	0	0	0	0	0	0	-	-	-	
Malignant neoplasms (140-209)	11	10	1	17	14	3	187.5	209.8	125.5	
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-	
Diabetes mellitus (250)	1	1	0	1	1	0	11.0	15.0	-	
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-	
Anemias (280-285)	0	0	0	0	0	0	-	-	-	
Diseases of heart (390-398, 402, 404-429)	28	21	7	40	35	5	441.2	524.4	209.1	
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-	
Cerebrovascular disease (430-438)	11	8	3	8	5	3	88.2	74.9	125.5	
Arteriosclerosis (440)	1	1	0	1	1	0	11.0	15.0	-	
Influenza and pneumonia (470-486)	1	1	0	1	1	0	11.0	15.0	-	
Bronchitis, emphysema, asthma (490-493)	0	0	0	1	1	0	11.0	15.0	-	
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-	
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-	
Cirrhosis of liver (571)	1	1	0	1	1	0	11.0	15.0	-	
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	1	0	11.0	15.0	-	
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-	
Infections of kidney (590)	0	0	0	0	0	0	-	-	-	
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-	
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-	
Congenital anomalies (740-759)	1	1	0	3	3	0	33.1	45.0	-	
Certain causes of mortality in early infancy (760-778)	2	2	0	3	3	0	15.6	23.1	-	
Accidents (800-949)	7	5	2	4	4	0	44.1	59.9	-	
Motor vehicle accidents (810-823)	2	2	0	1	1	0	11.0	15.0	-	
Suicide (950-959)	0	0	0	0	0	0	-	-	-	
Homicide (960-978)	0	0	0	0	0	0	-	-	-	
Ill-defined and unknown (780-796)	1	0	1	1	0	1	11.0	-	41.8	

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
PIKE COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non- white	Total	White	Non- white	Total	White	Non- white
Live births	630	325	305	580	256	324	18.3	14.3	23.4
Fetal deaths	16	4	12	14	4	10	24.1	15.6	30.9
Total deaths	405	215	190	415	220	195	13.1	12.3	14.1
Deaths under 1 year	19	3	16	17	3	14	29.3	11.7	43.2
Deaths under 28 days	14	2	12	9	0	9	15.5	-	27.8
Bacillary dysentery and amebiasis (004, 006)	0	0	0	1	0	1	3.1	-	7.2
Enteritis and other diarrheal diseases (008, 009)	3	0	3	3	0	3	9.4	-	21.6
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	40	29	11	47	29	18	148.0	162.0	129.9
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	5	1	4	5	1	4	15.7	5.6	28.9
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	2	1	1	1	0	1	3.1	-	7.2
Diseases of heart (390-398, 402, 404-429)	147	82	65	150	88	62	472.4	491.5	447.6
Hypertension (400, 401, 403)	3	2	1	3	2	1	9.4	11.2	7.2
Cerebrovascular disease (430-438)	69	43	26	66	39	27	207.8	217.8	194.9
Arteriosclerosis (440)	6	5	1	8	7	1	25.2	39.1	7.2
Influenza and pneumonia (470-486)	12	8	4	13	9	4	40.9	50.3	28.9
Bronchitis, emphysema, asthma (490-493)	7	4	3	5	2	3	15.7	11.2	21.6
Peptic ulcer (531-533)	1	1	0	1	1	0	3.1	5.6	-
Hernia and intestinal obstruction (550-553, 560)	1	1	0	0	0	0	-	-	-
Cirrhosis of liver (571)	4	4	0	6	6	0	18.9	33.5	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	2	1	1	1	0	1	3.1	-	7.2
Nephritis and nephrosis (580-584)	4	1	3	3	1	2	9.4	5.6	14.4
Infections of kidney (590)	2	1	1	4	2	2	12.6	11.2	14.4
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	0	1	3	1	2	9.4	5.6	14.4
Certain causes of mortality in early infancy (760-778)	11	1	10	8	0	8	13.8	-	24.7
Accidents (800-949)	20	8	12	21	10	11	66.1	55.8	79.4
Motor vehicle accidents (810-823)	9	3	6	10	5	5	31.5	27.9	36.1
Suicide (950-959)	2	2	0	2	2	0	6.3	11.2	-
Homicide (960-978)	7	2	5	7	2	5	22.0	11.2	36.1
Ill-defined and unknown (780-796)	35	6	29	33	6	27	103.9	33.5	194.9

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
PONTOTOC COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	287	204	83	290	215	75	16.7	15.1	24.1
Fetal deaths	6	4	2	5	4	1	17.2	18.6	13.3
Total deaths	187	164	23	228	202	26	13.1	14.2	8.4
Deaths under 1 year	2	1	1	5	3	2	17.2	14.0	26.7
Deaths under 28 days	1	1	0	2	1	1	6.9	4.6	13.3
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	1	0	1	0	0	0	-	-	-
Malignant neoplasms (140-209)	25	25	0	32	31	1	184.3	217.5	32.2
Benign and unspecified neoplasms (210-239)	1	0	1	1	0	1	5.8	-	32.2
Diabetes mellitus (250)	2	0	2	3	1	2	17.3	7.0	64.3
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	1	1	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	88	81	7	94	85	9	541.4	596.3	289.5
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	28	23	5	29	26	3	167.0	182.4	96.5
Arteriosclerosis (440)	5	5	0	4	4	0	23.0	28.1	-
Influenza and pneumonia (470-486)	3	2	1	4	4	0	23.0	28.1	-
Bronchitis, emphysema, asthma (490-493)	2	2	0	2	2	0	11.5	14.0	-
Peptic ulcer (531-533)	0	0	0	1	1	0	5.8	7.0	-
Hernia and intestinal obstruction (550-553, 560)	1	1	0	2	2	0	11.5	14.0	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	0	1	2	0	2	11.5	-	64.3
Infections of kidney (590)	1	1	0	2	2	0	11.5	14.0	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	1	1	0	2	1	1	6.9	4.6	13.3
Accidents (800-949)	7	7	0	27	25	2	155.5	175.4	64.3
Motor vehicle accidents (810-823)	4	4	0	20	18	2	115.2	126.3	64.3
Suicide (950-959)	2	2	0	2	2	0	11.5	14.0	-
Homicide (960-978)	3	2	1	2	2	0	11.5	14.0	-
Ill-defined and unknown (780-796)	3	1	2	4	2	2	23.0	14.0	64.3

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
PRENTISS COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	348	300	48	329	279	50	16.3	15.7	21.2
Fetal deaths	17	16	1	13	11	2	39.5	39.4	40.0
Total deaths	214	184	30	228	192	36	11.3	10.8	15.3
Deaths under 1 year	8	8	0	12	12	0	36.5	43.0	-
Deaths under 28 days	6	6	0	9	9	0	27.4	32.2	-
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	21	20	1	28	25	3	139.1	140.6	127.3
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	3	3	0	2	2	0	9.9	11.2	-
Avitaminoses and other nutritional deficiencies (260-269)	1	1	0	1	1	0	5.0	5.6	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	91	73	18	95	74	21	471.9	416.3	891.0
Hypertension (400, 401, 403)	2	2	0	2	2	0	9.9	11.2	-
Cerebrovascular disease (430-438)	23	21	2	23	21	2	114.2	118.1	84.8
Arteriosclerosis (440)	8	7	1	7	6	1	34.8	33.8	42.4
Influenza and pneumonia (470-486)	11	10	1	8	7	1	39.7	39.4	42.4
Bronchitis, emphysema, asthma (490-493)	5	5	0	5	5	0	24.8	28.1	-
Peptic ulcer (531-533)	0	0	0	1	1	0	5.0	5.6	-
Hernia and intestinal obstruction (550-553, 560)	1	1	0	1	1	0	5.0	5.6	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	1	0	5.0	5.6	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	1	0	1	1	0	1	5.0	-	42.4
Maternal causes (630-678)	1	0	1	1	0	1	30.4	-	200.0
Congenital anomalies (740-759)	1	1	0	2	2	0	9.9	11.2	-
Certain causes of mortality in early infancy (760-778)	5	5	0	8	8	0	24.3	28.7	-
Accidents (800-949)	11	8	3	14	9	5	69.5	50.6	212.1
Motor vehicle accidents (810-823)	4	2	2	8	4	4	39.7	22.5	169.7
Suicide (950-959)	1	1	0	1	1	0	5.0	5.6	-
Homicide (960-978)	2	1	1	2	1	1	9.9	5.6	42.4
Ill-defined and unknown (780-796)	16	16	0	12	12	0	59.6	67.5	-

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
QUITMAN COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	483	115	368	442	130	312	27.8	19.4	33.9
Fetal deaths	17	4	13	26	7	19	58.8	53.8	60.9
Total deaths	90	25	65	174	74	100	11.0	11.1	10.9
Deaths under 1 year	16	2	14	20	5	15	45.2	38.5	48.1
Deaths under 28 days	11	2	9	13	4	9	29.4	30.8	28.8
Bacillary dysentery and amebiasis (004, 006)	0	0	0	1	0	1	6.3	-	10.9
Enteritis and other diarrheal diseases (008, 009)	1	0	1	1	0	1	6.3	-	10.9
Tuberculosis (010-019)	1	0	1	2	1	1	12.6	15.0	10.9
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	4	1	3	14	6	8	88.1	89.7	86.9
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	3	1	2	5	1	4	31.5	15.0	43.5
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	1	0	1	6.3	-	10.9
Diseases of heart (390-398, 402, 404-429)	31	11	20	57	30	27	358.8	448.6	293.4
Hypertension (400, 401, 403)	1	0	1	3	0	3	18.9	-	32.6
Cerebrovascular disease (430-438)	8	2	6	31	14	17	195.1	209.4	184.8
Arteriosclerosis (440)	0	0	0	0	0	0	-	-	-
Influenza and pneumonia (470-486)	6	0	6	8	2	6	50.4	29.9	65.2
Bronchitis, emphysema, asthma (490-493)	2	2	0	2	2	0	12.6	29.9	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	2	0	2	3	1	2	18.9	15.0	21.7
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	0	1	1	0	1	6.3	-	10.9
Infections of kidney (590)	1	0	1	3	1	2	18.9	15.0	21.7
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	1	1	0	6.3	15.0	-
Certain causes of mortality in early infancy (760-778)	10	2	8	11	3	8	24.9	23.1	25.6
Accidents (800-949)	9	4	5	14	7	7	88.1	104.7	76.1
Motor vehicle accidents (810-823)	4	3	1	6	5	1	37.8	74.8	10.9
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	1	0	1	1	0	1	6.3	-	10.9
Ill-defined and unknown (780-796)	9	2	7	10	2	8	62.9	29.9	86.9

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
RANKIN COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	99	45	54	802	564	238	18.2	17.9	19.2
Fetal deaths	3	1	2	71	51	20	88.5	90.4	84.0
Total deaths	486	275	211	343	222	121	7.8	7.0	9.8
Deaths under 1 year	8	2	6	24	14	10	29.9	24.8	42.0
Deaths under 28 days	4	2	2	20	14	6	24.9	24.8	25.2
Bacillary dysentery and amebiasis (004, 006)	4	2	2	1	0	1	2.3	-	8.1
Enteritis and other diarrheal diseases (008, 009)	1	0	1	0	0	0	-	-	-
Tuberculosis (010-019)	4	2	2	1	0	1	2.3	-	8.1
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	53	29	24	53	34	19	120.6	107.8	153.2
Benign and unspecified neoplasms (210-239)	1	0	1	0	0	0	-	-	-
Diabetes mellitus (250)	10	8	2	6	5	1	13.6	15.8	8.1
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	1	1	0	1	1	0	2.3	3.2	-
Diseases of heart (390-398, 402, 404-429)	135	74	61	124	83	41	282.2	263.2	330.5
Hypertension (400, 401, 403)	1	0	1	0	0	0	-	-	-
Cerebrovascular disease (430-438)	51	30	21	31	19	12	70.6	60.3	96.7
Arteriosclerosis (440)	13	10	3	7	4	3	15.9	12.7	24.2
Influenza and pneumonia (470-486)	41	24	17	7	6	1	15.9	19.0	8.1
Bronchitis, emphysema, asthma (490-493)	11	9	2	6	5	1	13.6	15.8	8.1
Peptic ulcer (531-533)	1	1	0	1	1	0	2.3	3.2	-
Hernia and intestinal obstruction (550-553, 560)	3	3	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	0	1	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	0	1	3	3	0	6.8	9.5	-
Infections of kidney (590)	4	3	1	0	0	0	-	-	-
Hyperplasia of prostate (600)	1	1	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	1	0	1	12.5	-	42.0
Congenital anomalies (740-759)	1	1	0	5	5	0	11.4	15.8	-
Certain causes of mortality in early infancy (760-778)	0	0	0	11	8	3	13.7	14.2	12.6
Accidents (800-949)	42	23	19	32	16	16	72.8	50.7	129.0
Motor vehicle accidents (810-823)	16	10	6	19	13	6	43.2	41.2	48.4
Suicide (950-959)	6	4	2	7	6	1	15.9	19.0	8.1
Homicide (960-978)	6	4	2	4	1	3	9.1	3.2	24.2
Ill-defined and unknown (780-796)	26	10	16	17	7	10	38.7	22.2	80.6

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
SCOTT COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	408	272	136	461	244	217	21.6	17.1	30.5
Fetal deaths	10	6	4	15	9	6	32.5	36.9	27.6
Total deaths	219	149	70	256	166	90	12.0	11.6	12.6
Deaths under 1 year	12	4	8	14	4	10	30.4	16.4	46.1
Deaths under 28 days	9	4	5	8	3	5	17.4	12.3	23.0
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	1	0	1	1	0	1	4.7	-	14.0
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	20	16	4	31	23	8	145.1	161.4	112.4
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	1	0	1	2	1	1	9.4	7.0	14.0
Avitaminoses and other nutritional deficiencies (260-269)	1	0	1	1	0	1	4.7	-	14.0
Anemias (280-285)	0	0	0	1	1	0	4.7	7.0	-
Diseases of heart (390-398, 402, 404-429)	103	71	32	102	66	36	477.3	463.0	506.0
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	19	15	4	30	24	6	140.4	168.4	84.3
Arteriosclerosis (440)	8	5	3	10	6	4	46.8	42.1	56.2
Influenza and pneumonia (470-486)	8	6	2	5	3	2	23.4	21.0	28.1
Bronchitis, emphysema, asthma (490-493)	1	0	1	2	1	1	9.4	7.0	14.0
Peptic ulcer (531-533)	0	0	0	1	1	0	4.7	7.0	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	1	0	1	1	0	1	4.7	-	14.0
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	0	1	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	8	3	5	8	3	5	17.4	12.3	23.0
Accidents (800-949)	26	23	3	30	23	7	140.4	161.4	98.4
Motor vehicle accidents (810-823)	20	18	2	20	16	4	93.6	112.2	56.2
Suicide (950-959)	3	1	2	3	1	2	14.0	7.0	28.1
Homicide (960-978)	4	0	4	4	0	4	18.7	-	56.2
Ill-defined and unknown (780-796)	2	1	1	7	4	3	32.8	28.1	42.2

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
SHARKEY COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non- white	Total	White	Non- white	Total	White	Non- white
Live births	78	9	69	229	43	186	25.6	13.8	32.0
Fetal deaths	2	0	2	17	5	12	74.2	116.3	64.5
Total deaths	66	20	46	105	34	71	11.7	10.9	12.2
Deaths under 1 year	6	0	6	13	2	11	56.8	46.5	59.1
Deaths under 28 days	4	0	4	8	2	6	34.9	46.5	32.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	10	3	7	13	4	9	145.5	128.0	154.8
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	0	0	0	3	1	2	33.6	32.0	34.4
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	1	0	1	11.2	-	17.2
Diseases of heart (390-398, 402, 404-429)	27	10	17	35	13	22	391.6	416.0	378.5
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	13	3	10	15	3	12	167.8	96.0	206.5
Arteriosclerosis (440)	1	0	1	2	1	1	22.4	32.0	17.2
Influenza and pneumonia (470-486)	2	0	2	4	0	4	44.8	-	68.8
Bronchitis, emphysema, asthma (490-493)	1	1	0	2	2	0	22.4	64.0	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	0	1	11.2	-	17.2
Cirrhosis of liver (571)	0	0	0	1	1	0	11.2	32.0	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	0	1	1	0	1	11.2	-	17.2
Infections of kidney (590)	0	0	0	1	1	0	11.2	32.0	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	0	1	1	0	1	11.2	-	17.2
Certain causes of mortality in early infancy (760-778)	2	0	2	7	2	5	30.6	46.5	26.9
Accidents (800-949)	2	1	1	7	3	4	78.3	96.0	68.8
Motor vehicle accidents (810-823)	1	0	1	3	0	3	33.6	-	51.6
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	6	2	4	6	1	5	67.1	32.0	86.0

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
SIMPSON COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live Births, Fetal Deaths, Deaths									
Live births	303	183	120	428	224	204	21.4	16.4	32.5
Fetal deaths	5	4	1	21	10	11	49.1	44.6	53.9
Total deaths	258	150	108	279	175	104	14.0	12.8	16.6
Deaths under 1 year	14	2	12	17	3	14	39.7	13.4	68.6
Deaths under 28 days	9	2	7	10	2	8	23.4	8.9	39.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	17	6	11	3	2	1	15.0	14.6	16.0
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	40	26	14	39	28	11	195.5	204.7	175.5
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	3	3	0	2	2	0	10.0	14.6	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	75	51	24	85	60	25	426.1	438.7	398.8
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	26	19	7	30	24	6	150.4	175.5	95.7
Arteriosclerosis (440)	2	1	1	3	2	1	15.0	14.6	16.0
Influenza and pneumonia (470-486)	6	4	2	12	10	2	60.2	73.1	31.9
Bronchitis, emphysema, asthma (490-493)	2	2	0	1	1	0	5.0	7.3	-
Peptic ulcer (531-533)	1	1	0	1	1	0	5.0	7.3	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	2	1	1	10.0	7.3	16.0
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	14	7	7	12	6	6	60.2	43.9	95.7
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	2	1	1	2	1	1	10.0	7.3	16.0
Certain causes of mortality in early infancy (760-778)	7	2	5	8	2	6	18.7	8.9	29.4
Accidents (800-949)	22	7	15	29	9	20	145.4	65.8	319.0
Motor vehicle accidents (810-823)	8	3	5	13	4	9	65.2	29.2	143.6
Suicide (950-959)	2	2	0	2	2	0	10.0	14.6	-
Homicide (960-978)	1	0	1	1	0	1	5.0	-	16.0
Ill-defined and unknown (780-796)	22	8	14	24	10	14	120.3	73.1	223.3

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
SMITH COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non- white	Total	White	Non- white	Total	White	Non- white
Live births	8	0	8	257	152	105	19.0	14.3	35.6
Fetal deaths	0	0	0	12	7	5	46.7	46.0	47.6
Total deaths	55	47	8	156	139	17	11.5	13.1	5.8
Deaths under 1 year	2	1	1	4	1	3	15.6	6.6	28.6
Deaths under 28 days	0	0	0	2	0	2	7.8	-	19.0
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	1	0	1	7.4	-	33.9
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	8	6	2	21	17	4	154.8	160.2	135.5
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	0	0	0	2	1	1	14.7	9.4	33.9
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	1	1	0	7.4	9.4	-
Diseases of heart (390-398, 402, 404-429)	24	22	2	59	55	4	435.1	518.4	135.5
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	6	5	1	18	17	1	132.7	160.2	33.9
Arteriosclerosis (440)	3	3	0	8	8	0	59.0	75.4	-
Influenza and pneumonia (470-486)	1	1	0	4	4	0	29.5	37.7	-
Bronchitis, emphysema, asthma (490-493)	0	0	0	3	3	0	22.1	28.3	-
Peptic ulcer (531-533)	0	0	0	2	2	0	14.7	18.8	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	2	2	0	4	4	0	29.5	37.7	-
Infections of kidney (590)	0	0	0	2	2	0	14.7	18.8	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	1	1	0	7.4	9.4	-
Certain causes of mortality in early infancy (760-778)	0	0	0	1	0	1	3.9	-	9.5
Accidents (800-949)	7	6	1	17	16	1	125.4	150.8	33.9
Motor vehicle accidents (810-823)	0	0	0	8	8	0	59.0	75.4	-
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	0	0	0	2	1	1	14.7	9.4	33.9
Ill-defined and unknown (780-796)	2	1	1	2	1	1	14.7	9.4	33.9

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
STONE COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	137	88	49	165	90	75	20.4	14.5	39.9
Fetal deaths	3	0	3	7	3	4	42.4	33.3	53.3
Total deaths	100	77	23	94	73	21	11.6	11.7	11.2
Deaths under 1 year	5	3	2	3	1	2	18.2	11.1	26.7
Deaths under 28 days	5	3	2	3	1	2	18.2	11.1	26.7
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	1	1	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	7	7	0	9	9	0	111.1	144.7	-
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	1	0	1	1	0	1	12.3	-	53.2
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	41	39	2	40	36	4	493.8	578.8	212.6
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	16	10	6	12	8	4	148.1	128.6	212.6
Arteriosclerosis (440)	1	1	0	0	0	0	-	-	-
Influenza and pneumonia (470-486)	2	1	1	2	1	1	24.7	16.1	53.2
Bronchitis, emphysema, asthma (490-493)	2	2	0	3	3	0	37.0	48.2	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	1	1	0	12.3	16.1	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	1	0	1	1	0	1	60.6	-	133.3
Congenital anomalies (740-759)	1	1	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	4	2	2	3	1	2	18.2	11.1	26.7
Accidents (800-949)	11	6	5	13	10	3	160.5	160.8	159.5
Motor vehicle accidents (810-823)	7	3	4	8	6	2	98.8	96.5	106.3
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	3	0	3	3	0	3	37.0	-	159.5
Ill-defined and unknown (780-796)	4	2	2	2	0	2	24.7	-	106.3

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
SUNFLOWER COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	735	135	600	954	208	746	25.8	15.3	31.8
Fetal deaths	36	3	33	49	6	43	51.4	28.8	57.6
Total deaths	363	119	244	414	141	273	11.2	10.4	11.6
Deaths under 1 year	32	3	29	34	2	32	35.6	9.6	42.9
Deaths under 28 days	20	3	17	19	2	17	19.9	9.6	22.8
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	3	0	3	3	0	3	8.1	-	12.8
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	1	0	1	1	0	1	2.7	-	4.3
Malignant neoplasms (140-209)	32	16	16	41	21	20	110.7	154.2	85.4
Benign and unspecified neoplasms (210-239)	1	0	1	2	0	2	5.4	-	8.5
Diabetes mellitus (250)	15	5	10	13	5	8	35.1	36.7	34.1
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	2	0	2	1	0	1	2.7	-	4.3
Diseases of heart (390-398, 402, 404-429)	127	47	80	136	51	85	367.1	374.5	362.8
Hypertension (400, 401, 403)	4	0	4	4	0	4	10.8	-	17.1
Cerebrovascular disease (430-438)	57	18	39	59	17	42	159.2	124.8	179.3
Arteriosclerosis (440)	5	2	3	8	4	4	21.6	29.4	17.1
Influenza and pneumonia (470-486)	16	1	15	18	2	16	48.6	14.7	68.3
Bronchitis, emphysema, asthma (490-493)	2	1	1	3	2	1	8.1	14.7	4.3
Peptic ulcer (531-533)	1	1	0	2	2	0	5.4	14.7	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	3	2	1	8.1	14.7	4.3
Cirrhosis of liver (571)	0	0	0	1	1	0	2.7	7.3	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	1	0	1	2	1	1	5.4	7.3	4.3
Nephritis and nephrosis (580-584)	4	1	3	5	1	4	13.5	7.3	17.1
Infections of kidney (590)	4	1	3	5	1	4	13.5	7.3	17.1
Hyperplasia of prostate (600)	1	0	1	2	0	2	5.4	-	8.5
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	5	0	5	6	0	6	16.2	-	25.6
Certain causes of mortality in early infancy (760-778)	13	3	10	11	2	9	11.5	9.6	12.1
Accidents (800-949)	29	15	14	36	18	18	97.2	132.2	76.8
Motor vehicle accidents (810-823)	13	9	4	14	8	6	37.8	58.7	25.6
Suicide (950-959)	2	2	0	2	2	0	5.4	14.7	-
Homicide (960-978)	2	0	2	4	0	4	10.8	-	17.1
Ill-defined and unknown (780-796)	23	1	22	24	2	22	64.8	14.7	93.9

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
TALLAHATCHIE COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non- white	Total	White	Non- white	Total	White	Non- white
Live births	454	78	376	503	99	404	26.0	12.9	34.6
Fetal deaths	15	2	13	17	3	14	33.8	30.3	34.6
Total deaths	219	70	149	257	90	167	13.3	11.8	14.3
Deaths under 1 year	23	1	22	26	2	24	51.7	20.2	59.4
Deaths under 28 days	17	1	16	20	2	18	39.8	20.2	44.6
Bacillary dysentery and amebiasis (004, 006)	2	0	2	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	1	0	1	5.2	-	8.6
Tuberculosis (010-019)	1	0	1	1	0	1	5.2	-	8.6
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	25	14	11	36	19	17	186.2	248.1	145.5
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	1	0	1	3	2	1	15.5	26.1	8.6
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	74	29	45	80	31	49	413.7	404.8	419.5
Hypertension (400, 401, 403)	3	1	2	3	1	2	15.5	13.0	17.1
Cerebrovascular disease (430-438)	18	2	16	27	6	21	139.6	78.4	179.8
Arteriosclerosis (440)	6	2	4	6	2	4	31.0	26.1	34.2
Influenza and pneumonia (470-486)	7	2	5	10	3	7	51.7	39.2	59.9
Bronchitis, emphysema, asthma (490-493)	0	0	0	0	0	0	-	-	-
Peptic ulcer (531-533)	1	0	1	1	0	1	5.2	-	8.6
Hernia and intestinal obstruction (550-553, 560)	0	0	0	2	1	1	10.3	13.0	8.6
Cirrhosis of liver (571)	1	1	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	1	0	1	1	0	1	5.2	-	8.6
Infections of kidney (590)	3	1	2	2	1	1	10.3	13.0	8.6
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	12	1	11	13	2	11	25.8	20.2	27.2
Accidents (800-949)	21	10	11	23	12	11	118.9	156.7	94.2
Motor vehicle accidents (810-823)	13	7	6	13	7	6	67.2	91.4	51.4
Suicide (950-959)	3	1	2	3	1	2	15.5	13.0	17.1
Homicide (960-978)	2	1	1	1	1	0	5.2	13.0	-
Ill-defined and unknown (780-796)	28	3	25	30	3	27	155.1	39.2	231.1

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
TATE COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	137	0	137	405	157	248	21.8	16.0	28.3
Fetal deaths	4	0	4	13	2	11	32.1	12.7	44.4
Total deaths	105	47	58	185	99	86	10.0	10.1	9.8
Deaths under 1 year	6	0	6	13	4	9	32.1	25.5	36.3
Deaths under 28 days	5	0	5	11	4	7	27.2	25.5	28.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	2	1	1	2	1	1	10.8	10.2	11.4
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	1	1	0	5.4	10.2	-
Syphilis (090-097)	0	0	0	1	0	1	5.4	-	11.4
Malignant neoplasms (140-209)	15	7	8	27	17	10	145.6	173.9	114.1
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	3	2	1	4	3	1	21.6	30.7	11.4
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	38	25	13	64	39	25	345.1	398.9	285.2
Hypertension (400, 401, 403)	0	0	0	1	0	1	5.4	-	11.4
Cerebrovascular disease (430-438)	12	1	11	24	12	12	129.4	122.7	136.9
Arteriosclerosis (440)	0	0	0	0	0	0	-	-	-
Influenza and pneumonia (470-486)	2	1	1	7	3	4	37.7	30.7	45.6
Bronchitis, emphysema, asthma (490-493)	1	0	1	4	3	1	21.6	30.7	11.4
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	0	0	0	2	2	0	10.8	20.4	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	3	1	2	4	1	3	21.6	10.2	34.2
Certain causes of mortality in early infancy (760-778)	2	0	2	7	4	3	17.3	25.5	12.1
Accidents (800-949)	9	4	5	12	5	7	64.7	51.1	79.8
Motor vehicle accidents (810-823)	5	3	2	6	2	4	32.4	20.4	45.6
Suicide (950-959)	1	1	0	1	1	0	5.4	10.2	-
Homicide (960-978)	1	1	0	5	2	3	27.0	20.4	34.2
Ill-defined and unknown (780-796)	12	3	9	10	2	8	53.9	20.4	91.2

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
TIPPAH COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non- white	Total	White	Non- white	Total	White	Non- white
Live births	352	271	81	296	230	66	18.7	17.3	25.5
Fetal deaths	25	17	8	25	16	9	84.4	69.6	136.4
Total deaths	141	120	21	151	127	24	9.5	9.6	9.2
Deaths under 1 year	8	5	3	3	1	2	10.1	4.3	30.3
Deaths under 28 days	7	5	2	2	1	1	6.8	4.3	15.2
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	0	0	0	-	-	-
Tuberculosis (010-019)	1	1	0	1	1	0	6.3	7.5	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	26	26	0	29	26	3	182.9	196.1	115.7
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	4	4	0	4	4	0	25.2	30.2	-
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	57	49	8	60	55	5	378.5	414.8	192.9
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	23	19	4	22	18	4	138.8	135.7	154.3
Arteriosclerosis (440)	2	2	0	2	2	0	12.6	15.1	-
Influenza and pneumonia (470-486)	2	1	1	6	2	4	37.8	15.1	154.3
Bronchitis, emphysema, asthma (490-493)	1	1	0	0	0	0	-	-	-
Peptic ulcer (531-533)	1	1	0	1	1	0	6.3	7.5	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	3	3	0	3	3	0	18.9	22.6	-
Hyperplasia of prostate (600)	0	0	0	1	1	0	6.3	7.5	-
Maternal causes (630-678)	0	0	0	1	0	1	33.8	-	151.5
Congenital anomalies (740-759)	1	1	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	5	4	1	2	1	1	6.8	4.3	15.2
Accidents (800-949)	5	3	2	6	4	2	37.8	30.2	77.2
Motor vehicle accidents (810-823)	2	1	1	4	3	1	25.2	22.6	38.6
Suicide (950-959)	2	2	0	2	2	0	12.6	15.1	-
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	5	3	2	4	2	2	25.2	15.1	77.2

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
TISHOMINGO COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence						
	Number			Number			Rate*			
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	
Live Births, Fetal Deaths, Deaths										
Live births	169	154	15	239	225	14	16.0	15.8	20.8	
Fetal deaths	2	2	0	3	3	0	12.6	13.3	-	
Total deaths	115	107	8	174	165	9	11.6	11.6	13.4	
Deaths under 1 year	1	1	0	7	7	0	29.3	31.1	-	
Deaths under 28 days	0	0	0	5	5	0	20.9	22.2	-	
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-	
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-	
Tuberculosis (010-019)	1	1	0	0	0	0	-	-	-	
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-	
Syphilis (090-097)	0	0	0	0	0	0	-	-	-	
Malignant neoplasms (140-209)	21	20	1	26	25	1	174.0	175.2	148.8	
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-	
Diabetes mellitus (250)	1	1	0	4	4	0	26.8	28.0	-	
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-	
Anemias (280-285)	0	0	0	0	0	0	-	-	-	
Diseases of heart (390-398, 402, 404-429)	44	42	2	65	62	3	435.1	434.5	446.4	
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-	
Cerebrovascular disease (430-438)	19	16	3	19	16	3	127.2	112.1	446.4	
Arteriosclerosis (440)	0	0	0	3	3	0	20.1	21.0	-	
Influenza and pneumonia (470-486)	1	1	0	7	7	0	46.8	49.1	-	
Bronchitis, emphysema, asthma (490-493)	2	2	0	4	4	0	26.8	28.0	-	
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-	
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-	
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-	
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-	
Nephritis and nephrosis (580-584)	1	1	0	0	0	0	-	-	-	
Infections of kidney (590)	1	1	0	1	1	0	6.7	7.0	-	
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-	
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-	
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-	
Certain causes of mortality in early infancy (760-778)	0	0	0	5	5	0	20.9	22.2	-	
Accidents (800-949)	12	11	1	22	21	1	147.2	147.2	148.8	
Motor vehicle accidents (810-823)	5	5	0	16	16	0	107.1	112.1	-	
Suicide (950-959)	0	0	0	0	0	0	-	-	-	
Homicide (960-978)	1	1	0	1	1	0	6.7	7.0	-	
Ill-defined and unknown (780-796)	5	5	0	8	8	0	53.5	56.1	-	

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
TUNICA COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate *		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	340	18	322	370	53	317	31.2	16.4	36.7
Fetal deaths	25	0	25	25	1	24	67.6	18.9	75.7
Total deaths	157	32	125	165	39	126	13.9	12.1	14.6
Deaths under 1 year	14	0	14	15	1	14	40.5	18.9	44.2
Deaths under 28 days	7	0	7	8	1	7	21.6	18.9	22.1
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	1	0	1	1	0	1	8.4	-	11.6
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	24	6	18	31	7	24	261.5	217.0	278.1
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	5	2	3	5	2	3	42.2	62.0	34.8
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	1	0	1	1	0	1	8.4	-	11.6
Diseases of heart (390-398, 402, 404-429)	49	11	38	51	11	40	430.2	341.1	463.6
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	19	3	16	20	3	17	168.7	93.0	197.0
Arteriosclerosis (440)	1	0	1	1	0	1	8.4	-	11.6
Influenza and pneumonia (470-486)	7	0	7	9	3	6	75.9	93.0	69.5
Bronchitis, emphysema, asthma (490-493)	2	2	0	2	2	0	16.9	62.0	-
Peptic ulcer (531-533)	0	0	0	2	2	0	16.9	62.0	-
Hernia and intestinal obstruction (550-553, 560)	1	0	1	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	3	1	2	4	1	3	33.7	31.0	34.8
Infections of kidney (590)	2	1	1	1	0	1	8.4	-	11.6
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	5	0	5	6	1	5	16.2	18.9	15.8
Accidents (800-949)	8	5	3	5	4	1	42.2	124.0	11.6
Motor vehicle accidents (810-823)	2	2	0	0	0	0	-	-	-
Suicide (950-959)	1	0	1	1	0	1	8.4	-	11.6
Homicide (960-978)	8	1	7	4	1	3	33.7	31.0	34.8
Ill-defined and unknown (780-796)	18	0	18	18	1	17	151.8	31.0	197.0

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
UNION COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	301	220	81	336	264	72	17.6	16.4	24.4
Fetal deaths	26	18	8	21	14	7	62.5	53.0	97.2
Total deaths	207	178	29	211	183	28	11.0	11.3	9.5
Deaths under 1 year	5	3	2	5	3	2	14.9	11.4	27.8
Deaths under 28 days	3	3	0	3	3	0	8.9	11.4	-
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	25	19	6	29	25	4	151.9	154.8	135.5
Benign and unspecified neoplasms (210-239)	0	0	0	1	1	0	5.2	6.2	-
Diabetes mellitus (250)	8	7	1	7	6	1	36.6	37.2	33.9
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	0	0	0	1	1	0	5.2	6.2	-
Diseases of heart (390-398, 402, 404-429)	66	59	7	68	62	6	356.1	384.0	203.3
Hypertension (400, 401, 403)	0	0	0	0	0	0	-	-	-
Cerebrovascular disease (430-438)	22	19	3	24	19	5	125.7	117.7	169.4
Arteriosclerosis (440)	6	6	0	5	5	0	26.2	31.0	-
Influenza and pneumonia (470-486)	15	14	1	14	14	0	73.3	86.7	-
Bronchitis, emphysema, asthma (490-493)	1	1	0	2	2	0	10.5	12.4	-
Peptic ulcer (531-533)	1	0	1	1	0	1	5.2	-	33.9
Hernia and intestinal obstruction (550-553, 560)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	7	6	1	6	5	1	31.4	31.0	33.9
Hyperplasia of prostate (600)	2	2	0	2	2	0	10.5	12.4	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	3	3	0	2	2	0	6.0	7.6	-
Accidents (800-949)	18	17	1	17	16	1	89.0	99.1	33.9
Motor vehicle accidents (810-823)	10	9	1	10	9	1	52.4	55.7	33.9
Suicide (950-959)	2	2	0	2	2	0	10.5	12.4	-
Homicide (960-978)	0	0	0	1	1	0	5.2	6.2	-
Ill-defined and unknown (780-796)	18	11	7	16	9	7	83.8	55.7	237.2

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
WALTHALL COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	299	149	150	248	98	150	19.8	13.2	29.4
Fetal deaths	10	4	6	10	3	7	40.3	30.6	46.7
Total deaths	105	56	49	127	69	58	10.2	9.3	11.4
Deaths under 1 year	6	2	4	8	3	5	32.2	30.6	33.3
Deaths under 28 days	4	2	2	6	3	3	24.2	30.6	20.0
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	0	0	0	-	-	-
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	16	8	8	17	8	9	136.0	108.1	176.6
Benign and unspecified neoplasms (210-239)	1	0	1	1	0	1	8.0	-	19.6
Diabetes mellitus (250)	4	1	3	5	1	4	40.0	13.5	78.5
Avitaminoses and other nutritional deficiencies (260-269)	1	1	0	1	1	0	8.0	13.5	-
Anemias (280-285)	0	0	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	41	26	15	48	31	17	384.0	418.7	333.5
Hypertension (400, 401, 403)	0	0	0	1	0	1	8.0	-	19.6
Cerebrovascular disease (430-438)	7	5	2	14	8	6	112.0	108.1	117.7
Arteriosclerosis (440)	9	9	0	7	7	0	56.0	94.6	-
Influenza and pneumonia (470-486)	1	0	1	3	1	2	24.0	13.5	39.2
Bronchitis, emphysema, asthma (490-493)	1	1	0	2	1	1	16.0	13.5	19.6
Peptic ulcer (531-533)	1	0	1	1	0	1	8.0	-	19.6
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	1	0	8.0	13.5	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	1	0	8.0	13.5	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	1	1	0	0	0	0	-	-	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	1	0	1	1	0	8.0	13.5	-
Certain causes of mortality in early infancy (760-778)	3	1	2	2	1	1	8.1	10.2	6.7
Accidents (800-949)	6	1	5	8	5	3	64.0	67.5	58.8
Motor vehicle accidents (810-823)	5	1	4	6	4	2	48.0	54.0	39.2
Suicide (950-959)	1	0	1	1	0	1	8.0	-	19.6
Homicide (960-978)	0	0	0	0	0	0	-	-	-
Ill-defined and unknown (780-796)	8	0	8	9	0	9	72.0	-	176.6

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
WARREN COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,378	605	773	917	498	419	20.4	18.8	22.6
Fetal deaths	135	57	78	89	48	41	97.0	96.4	97.8
Total deaths	670	318	352	514	252	262	11.4	9.5	14.2
Deaths under 1 year	45	16	29	34	13	21	37.1	26.1	50.1
Deaths under 28 days	39	15	24	29	12	17	31.6	24.1	40.6
Bacillary dysentery and amebiasis (004, 006)	0	0	0	1	1	0	2.2	3.8	-
Enteritis and other diarrheal diseases (008, 009)	5	2	3	3	1	2	6.7	3.8	10.8
Tuberculosis (010-019)	1	0	1	2	0	2	4.4	-	10.8
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	2	0	2	2	0	2	4.4	-	10.8
Malignant neoplasms (140-209)	111	61	50	78	47	31	173.4	177.5	167.5
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	13	1	12	7	0	7	15.6	-	37.8
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	4	0	4	2	0	2	4.4	-	10.8
Diseases of heart (390-398, 402, 404-429)	178	96	82	152	83	69	337.9	313.5	372.8
Hypertension (400, 401, 403)	3	1	2	3	1	2	6.7	3.8	10.8
Cerebrovascular disease (430-438)	98	45	53	73	33	40	162.3	124.6	216.1
Arteriosclerosis (440)	9	6	3	5	4	1	11.1	15.1	5.4
Influenza and pneumonia (470-486)	11	5	6	10	5	5	22.2	18.9	27.0
Bronchitis, emphysema, asthma (490-493)	4	3	1	3	2	1	6.7	7.6	5.4
Peptic ulcer (531-533)	4	2	2	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	2	1	1	1	1	0	2.2	3.8	-
Cirrhosis of liver (571)	8	7	1	7	6	1	15.6	22.7	5.4
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	2	1	1	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	6	1	5	5	2	3	11.1	7.6	16.2
Infections of kidney (590)	13	9	4	9	6	3	20.0	22.7	16.2
Hyperplasia of prostate (600)	4	1	3	1	0	1	2.2	-	5.4
Maternal causes (630-678)	0	0	0	1	0	1	10.9	-	23.9
Congenital anomalies (740-759)	10	3	7	11	4	7	24.4	15.1	37.8
Certain causes of mortality in early infancy (760-778)	28	11	17	17	8	9	18.5	16.1	21.5
Accidents (800-949)	51	26	25	36	20	16	80.0	75.5	86.4
Motor vehicle accidents (810-823)	17	8	9	10	7	3	22.2	26.4	16.2
Suicide (950-959)	4	3	1	3	2	1	6.7	7.6	5.4
Homicide (960-978)	1	1	0	1	1	0	2.2	3.8	-
Ill-defined and unknown (780-796)	50	6	44	44	3	41	97.8	11.3	221.5

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
WASHINGTON COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	1,918	703	1,215	1,819	613	1,206	25.8	19.3	31.1
Fetal deaths	128	53	75	120	38	82	66.0	62.0	68.0
Total deaths	774	286	488	807	281	526	11.4	8.8	13.6
Deaths under 1 year	62	18	44	58	16	42	31.9	26.1	34.8
Deaths under 28 days	46	17	29	44	15	29	24.2	24.5	24.0
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	5	2	3	5	2	3	7.1	6.3	7.7
Tuberculosis (010-019)	1	0	1	1	0	1	1.4	-	2.6
Meningococcal infection (036)	1	0	1	1	0	1	1.4	-	2.6
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	110	44	66	112	48	64	158.7	150.9	165.0
Benign and unspecified neoplasms (210-239)	2	0	2	2	0	2	2.8	-	5.2
Diabetes mellitus (250)	19	2	17	21	2	19	29.8	6.3	49.0
Avitaminoses and other nutritional deficiencies (260-269)	2	0	2	3	0	3	4.2	-	7.7
Anemias (280-285)	1	0	1	1	0	1	1.4	-	2.6
Diseases of heart (390-398, 402, 404-429)	224	95	129	224	87	137	317.4	273.6	353.3
Hypertension (400, 401, 403)	6	1	5	8	1	7	11.3	3.1	18.0
Cerebrovascular disease (430-438)	99	40	59	112	40	72	158.7	125.8	185.7
Arteriosclerosis (440)	6	4	2	8	4	4	11.3	12.6	10.3
Influenza and pneumonia (470-486)	25	6	19	24	6	18	34.0	18.9	46.4
Bronchitis, emphysema, asthma (490-493)	5	1	4	5	1	4	7.1	3.1	10.3
Peptic ulcer (531-533)	2	1	1	3	2	1	4.2	6.3	2.6
Hernia and intestinal obstruction (550-553, 560)	2	0	2	4	1	3	5.7	3.1	7.7
Cirrhosis of liver (571)	2	2	0	2	2	0	2.8	6.3	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	2	1	1	1	0	1	1.4	-	2.6
Nephritis and nephrosis (580-584)	2	1	1	3	1	2	4.2	3.1	5.2
Infections of kidney (590)	6	4	2	4	2	2	5.7	6.3	5.2
Hyperplasia of prostate (600)	2	1	1	1	1	0	1.4	3.1	-
Maternal causes (630-678)	1	0	1	1	0	1	5.5	-	8.3
Congenital anomalies (740-759)	2	1	1	2	0	2	2.8	-	5.2
Certain causes of mortality in early infancy (760-778)	33	15	18	31	13	18	17.0	21.2	14.9
Accidents (800-949)	74	31	43	77	29	48	109.1	91.2	123.8
Motor vehicle accidents (810-823)	31	18	13	28	13	15	39.7	40.9	38.7
Suicide (950-959)	4	3	1	7	6	1	9.9	18.9	2.6
Homicide (960-978)	13	1	12	12	1	11	17.0	3.1	28.4
Ill-defined and unknown (780-796)	77	10	67	79	11	68	111.9	34.6	175.4

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
WAYNE COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	337	246	91	345	191	154	20.7	17.1	28.1
Fetal deaths	9	5	4	12	7	5	34.8	36.6	32.5
Total deaths	191	130	61	203	135	68	12.2	12.1	12.4
Deaths under 1 year	14	5	9	17	5	12	49.3	26.2	77.9
Deaths under 28 days	13	5	8	16	5	11	46.4	26.2	71.4
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	0	0	0	1	0	1	6.0	-	18.2
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	20	12	8	26	17	9	156.2	152.2	164.2
Benign and unspecified neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (250)	11	6	5	8	3	5	48.0	26.9	91.2
Avitaminoses and other nutritional deficiencies (260-269)	1	1	0	1	1	0	6.0	9.0	-
Anemias (280-285)	1	1	0	0	0	0	-	-	-
Diseases of heart (390-398, 402, 404-429)	75	53	22	69	49	20	414.4	438.7	364.9
Hypertension (400, 401, 403)	6	3	3	5	3	2	30.0	26.9	36.5
Cerebrovascular disease (430-438)	16	10	6	18	10	8	108.1	89.5	146.0
Arteriosclerosis (440)	1	0	1	1	0	1	6.0	-	18.2
Influenza and pneumonia (470-486)	6	3	3	7	5	2	42.0	44.8	36.5
Bronchitis, emphysema, asthma (490-493)	7	6	1	6	5	1	36.0	44.8	18.2
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	0	0	0	1	1	0	6.0	9.0	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	0	0	0	-	-	-
Nephritis and nephrosis (580-584)	0	0	0	2	1	1	12.0	9.0	18.2
Infections of kidney (590)	2	2	0	2	2	0	12.0	17.9	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	1	0	1	2	1	1	12.0	9.0	18.2
Certain causes of mortality in early infancy (760-778)	10	5	5	12	5	7	34.8	26.2	45.4
Accidents (800-949)	17	12	5	21	13	8	126.1	116.4	146.0
Motor vehicle accidents (810-823)	9	7	2	11	7	4	66.1	62.7	73.0
Suicide (950-959)	4	4	0	4	4	0	24.0	35.8	-
Homicide (960-978)	2	1	1	0	0	0	-	-	-
Ill-defined and unknown (780-796)	1	1	0	3	3	0	18.0	26.9	-

See page 61 for footnote.

LIVE BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH
(WITH INTERNATIONAL CLASSIFICATION NUMBERS)
WEBSTER COUNTY, MISSISSIPPI, 1969

By Place of Occurrence, and by Place of Residence, by Race, with Rates

Live Births, Fetal Deaths, Deaths	Occurrence			Residence					
	Number			Number			Rate*		
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Live births	109	68	41	174	114	60	17.3	14.6	26.6
Fetal deaths	7	1	6	11	3	8	63.2	26.3	133.3
Total deaths	96	78	18	128	102	26	12.7	13.1	11.5
Deaths under 1 year	3	1	2	5	0	5	28.7	-	83.3
Deaths under 28 days	2	0	2	4	0	4	23.0	-	66.7
Bacillary dysentery and amebiasis (004, 006)	0	0	0	0	0	0	-	-	-
Enteritis and other diarrheal diseases (008, 009)	1	0	1	2	0	2	19.9	-	88.6
Tuberculosis (010-019)	0	0	0	0	0	0	-	-	-
Meningococcal infection (036)	0	0	0	0	0	0	-	-	-
Syphilis (090-097)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-209)	18	14	4	20	16	4	199.1	205.4	177.1
Benign and unspecified neoplasms (210-239)	1	1	0	1	1	0	10.0	12.8	-
Diabetes mellitus (250)	2	0	2	3	1	2	29.8	12.8	88.6
Avitaminoses and other nutritional deficiencies (260-269)	0	0	0	0	0	0	-	-	-
Anemias (280-285)	1	1	0	1	1	0	10.0	12.8	-
Diseases of heart (390-398, 402, 404-429)	20	18	2	27	23	4	268.7	295.3	177.1
Hypertension (400, 401, 403)	0	0	0	1	0	1	10.0	-	44.3
Cerebrovascular disease (430-438)	19	17	2	28	24	4	278.7	308.1	177.1
Arteriosclerosis (440)	0	0	0	3	3	0	29.8	38.5	-
Influenza and pneumonia (470-486)	2	1	1	2	1	1	19.9	12.8	44.3
Bronchitis, emphysema, asthma (490-493)	2	2	0	4	4	0	39.8	51.4	-
Peptic ulcer (531-533)	0	0	0	0	0	0	-	-	-
Hernia and intestinal obstruction (550-553, 560)	1	1	0	1	1	0	10.0	12.8	-
Cirrhosis of liver (571)	0	0	0	0	0	0	-	-	-
Cholelithiasis, cholecystitis, and cholangitis (574, 575)	0	0	0	1	1	0	10.0	12.8	-
Nephritis and nephrosis (580-584)	0	0	0	0	0	0	-	-	-
Infections of kidney (590)	2	2	0	3	3	0	29.8	38.5	-
Hyperplasia of prostate (600)	0	0	0	0	0	0	-	-	-
Maternal causes (630-678)	0	0	0	0	0	0	-	-	-
Congenital anomalies (740-759)	0	0	0	0	0	0	-	-	-
Certain causes of mortality in early infancy (760-778)	0	0	0	1	0	1	5.7	-	16.7
Accidents (800-949)	7	7	0	8	7	1	79.6	89.9	44.3
Motor vehicle accidents (810-823)	5	5	0	6	5	1	59.7	64.2	44.3
Suicide (950-959)	0	0	0	0	0	0	-	-	-
Homicide (960-978)	4	4	0	4	4	0	39.8	51.4	-
Ill-defined and unknown (780-796)	8	3	5	8	3	5	79.6	38.5	221.4